2	STUDY_PD	INITYR	Initial LBCL Diagnosis Year	~~2016 to 2025~~	year(INIT)	let's take all data for now but period will be defined later	DDE 04dec25	DDE 16dec25
3	STUDY_PD	INDEXYR	Index Year	~~2016 to 2025~~	year(INDEX)	let's take all data for now but period will be defined later	DDE 16dec25	
4	STUDY_PD	DLBCLFL	DLBCL diagnosis flag	'Yes'; 'No'	Assign DLBCLFL='Yes' if assessed_timedelta=0 and histology/histology_description %IN% Malignant lymphoma-large B-cell-diffuse-immunoblastic-NOS; Malignant lymphoma-large B-cell-diffuse-centroblastic-NOS; Histiocyte-rich large B-cell lymphoma; T-cell rich large B-cell lymphoma; ALK positive large B-cell lymphoma; HHV8 positive diffuse large B-cell lymphoma; Diffuse large B-cell lymphoma-NOS; T-cell/histiocyte rich large B-cell lymphoma; Thymic large B-cell lymphoma; EBV positive diffuse large B-cell lymphoma; Malignant lymphoma-large B-cell-diffuse-NOS; Anaplastic large B-cell lymphoma; Mediastinal large B-cell lymphoma; Otherwise assign DLBCL='No';	NOTE: Single value for all cohorts. List of histology_description that qualifies as DLBCL subject to change after review by clinical team	DDE16JAN26 (will probably be updated later on)	
5	STUDY_PD	DLBCLFLN	DLBCL diagnosis flag (N)	1,2	Numeric of DLBCLFL: 1=Yes; 2=No;		DDE 16dec25	
6	PRE_INT	STAGE	Stage at LBCL diagnosis	'I'; 'II'; 'III'; 'IV'; 'Unknown/not documented'	lymphoma_stating/principal_stage	Only one variable available at initial diagnosis, i.e. will be the same for all index dates	DDE 04dec25	DDE16JAN26: period should rather be study_period as it's collected once around diagnosis date (so can some days/weeks after)
7	PRE_INT	STAGEN	Stage at LBCL diagnosis (N)	1,2,3,4,5	Numeric of STAGE: 1=I; 2=II; 3=III; 4=IV; 5=Unknown/not documented;	Only one variable available at initial diagnosis, i.e. will be the same for all index dates	DDE 04dec25	DDE16JAN26: period should rather be study_period as it's collected once around diagnosis date (so can some days/weeks after)
	ASSESS_ECOG	ECOG	ECOG	'0' '1' '2' '3'	ECOG values in performance_status/score for performance_status/method=ECOG; The closest ECOG value to index date within the time window (ASSESS_ECOG) will be used. For patients with		DDE 04dec25	

1.Cover | 2.QC Review | 3.Data Prep | 4.StudyPop | 5A. Demos | 5B.ClinChars | 5C.BioVars | 5D.LabVars | 6.TreatVars | 7.Outcomes

Ready Accessibility: Investigate Display Settings

Insert Draw Page Layout Formulas Data Review View Help

Copy Format Painter Clipboard | Arial 11 Font | Alignment | Merge

'ECOG values in performance_status/score for performance_status/method=ECOG;

	A	B	C	D	E	F	G	H
	Time Period	Variable	Label	Values	Definition	Additional Notes	QC Reviewed (Initials + Date)	Date Modified (Initials + Date)
	ASSESS_ECOG	ECOG	ECOG	'0' '1' '2' '3' '4' 'Unknown' 'Missing'	ECOG values in performance_status/score for performance_status/method=ECOG; The closest ECOG value to index date within the time window (ASSESS_ECOG) will be used. For patients with multiple ECOG values available on the same day, the highest value is selected.		DDE 04dec25	
8	ASSESS_ECOG	ECOGN	ECOG (N)	1,2,3,4,5,6,7	Numeric code for ECOG: 1=0; 2=1; 3=2; 4=3; 5=4; 6=Unknown; 7=Missing;		DDE 04dec25	
9	PRE_INT	WEIGHT	Weight (kg)	Numeric	In height_weight table, select row(s) with value_type=weight and assessment date (INIT + assessed_timedelta) closest and prior to INDEX; if unit=kg, WEIGHT=value; else WEIGHT=value/2 205	Convert lbs to kg by dividing by 2.205	DDE 04dec25	
10	PRE_INT	HEIGHT	Height (cm)	Numeric	In height_weight table, select row(s) with value_type=height and assessment date (INIT + assessed_timedelta) closest and prior to INDEX; if unit=cm, HEIGHT=value; else HEIGHT=value*2 54	Convert in to cm by multiplying by 2 54	DDE 04dec25	
11	PRE_INT	BMI	BMI (kg/m^2)	Numeric	BMI=WEIGHT/(HEIGHT/100)^2; if WEIGHT is missing or HEIGHT is missing then BMI is missing.	Height and weight do not need to be on the same date. Height has much less variation for adults than weight.	DDE 04dec25	
12	PRE_INT	BMIGR1	BMI (kg/m^2) Group 1	'Underweight' 'Normal weight' 'Overweight' 'Obese' 'Missing'	if BMI <18 5 then ='Underweight'; else if 18.5 <= BMI < 25 then ='Normal weight'; else if 25 <= BMI < 30 then ='Overweight'; else if BMI >= 30 then ='Obese' else if BMI is missing then ='Missing';		DDE 04dec25	
13	PRE_INT	BMIGR1N	BMI (kg/m^2) Group 1 (N)	1,2,3,4,5	Numeric of BMGR1: 1='Underweight' 2='Normal weight' 3='Overweight' 4='Obese' 5='Missing';		DDE 04dec25	
14	PRE_INT	BSA	Body Surface Area	Numeric	BSA (m2) = 0 007184 * HEIGHT^0.725 * WEIGHT^0.425 if HEIGHT or WEIGHT are missing, then BSA is also missing. Note: formula assumes HEIGHT in cm and	Height and weight do not need to be on the same date. Height has much less variation for adults than weight. Reference: Body Surface Area - StatPearls - NCBI Bookshelf (https://www.ncbi.nlm.nih.gov/books/NBK559005/#:~:text=Ther	DDE 04dec25	

1.Cover | 2.QC Review | 3.Data Prep | 4.StudyPop | 5A. Demos | 5B.ClinChars | 5C.BioVars | 5D.LabVars | 6.TreatVars | 7.Outcomes

Ready Accessibility: Investigate Display Settings

E15 BSA (m2) = 0.007184 * HEIGHT^0.725 * WEIGHT^0.425

Time Period	Variable	Label	Values	Definition	Additional Notes	QC Reviewed (Initials + Date)	Date Modified (Initials + Date)
PRE_INT	BSA	Body Surface Area	Numeric	BSA (m2) = 0.007184 * HEIGHT^0.725 * WEIGHT^0.425 if HEIGHT or WEIGHT are missing, then BSA is also missing. Note: formula assumes HEIGHT in cm and WEIGHT in kg	Height and weight do not need to be on the same date. Height has much less variation for adults than weight. Reference: Body Surface Area - StatPearls - NCBI Bookshelf (https://www.ncbi.nlm.nih.gov/books/NBK559005/#:~:text=There%20exist%20many%20variations%20in,(Weight(kg)0.425)https://www.omnicalculator.com/health/bsa#the-bsa-formula)	DDE 04dec25	
ASSESS_BSYMPT	BSYMPT	B symptoms	'Yes'; 'No'; 'Missing';	In b_symptoms tables; if present=TRUE then BSYMPT='Yes'; else if present=FALSE then BSYMPT='No'; else if present is missing then BSYMPT='Missing'	Captured at initial LBCL diagnosis (within 30 days of initial diagnosis) and at progression or relapse (within 30 days) Value at diagnosis is carried forward until and if a new value is available.	DDE 04dec25	
ASSESS_BSYMPT	BSYMPTN	B symptoms (N)	1,2,3	Numeric of BSYMPT: 1='Yes'; 2='No'; 3='Missing'		DDE 16dec25	
ASSESS_BULKY	BULKY	Bulky disease	'Yes'; 'No'; 'Missing'	In bulky_disease tables; if is_bulky_disease=TRUE then BULKY='Yes'; else if is_bulky_disease=FALSE then BULKY='No'; else if is_bulky_disease is missing then BULKY='Missing'	Captured once at initial LBCL diagnosis (within 30 days of initial diagnosis)	DDE 04dec25	
ASSESS_BULKY	BULKYN	Bulky disease (N)	1,2,3	Numeric of BULKY: 1='Yes'; 2='No'; 3='Missing'		DDE 16dec25	
PRE_INT	SUBTYPE	DLBCL molecular classification (subtype)	'Activated B-cell-like '; 'Activated B-cell-like and germinal B-cell-like'; 'Germinal center B-cell-like'; 'Unknown'	DLBCL subtype as per subtype_dlbcl/subtype	We need the confirm possible values for subtype_dlbcl/subtype and decide if mapping is required.	DDE 04dec25	
PRE_INT	SUBTYPEN	DLBCL molecular classification (subtype) (N)	1,2,3,4	Numeric of SUBTYPE: 1='Activated B-cell-like '; 2='Activated B-cell-like and germinal B-cell-like'; 3='Germinal center B-cell-like'; 4='Unknown'		DDE 16dec25	
PRE_INT	EXTRANOD	Extranodal involvement	'Unspecified'; 'Absent'; 'Inconclusive'; '1'; '2+'	In extranodal_sites tables: if involvement='unspecified' then EXTRANOD='Unspecified'; else if involvement='absent' then EXTRANOD='Absent'; else if involvement='inconclusive' then EXTRANOD='Inconclusive'; else if involvement='present' then use number_of_sites to classify into '1' or '2+'	We need the confirm possible values for extranodal_sites/involvement and decide if mapping is required	DDE 16dec25	
PRE_INT	EXTRANODN	Extranodal involvement (N)	1,2,3,4,5	Numeric of EXTRANOD: 1='Unspecified'; 2='Absent'; 3='Inconclusive'; 4='1'; 5='2+'		DDE 16dec25	

1.Cover | 2.QC Review | 3.Data Prep | 4.StudyPop | 5A. Demos | 5B.ClinChars | 5C.BioVars | 5D.LabVars | 6.TreatVars | 7.Outcomes

Ready Accessibility: Investigate

Time Period	Variable	Label	Values	Definition	Additional Notes	QC Reviewed (Initials + Date)	Date Modified (Initials + Date)
PRE_INT	CCI	Charlson Comorbidity Index (CCI) score	Numeric	Calculate CCI, including the patient's personal history of cancer. Use COTA R package function calculate_CCI(), with argument personal_history_of_cancer = all tables pers_history_of_cancer and add_age=FALSE (default).	NOTE: Single value for all cohorts, because comorbidities are anchored at the time of initial diagnosis. The timedelta variables are irrelevant for the derivations of CCI, we can safely assume (according to COTA) that all comorbidities recorded in a patient's file were present at initial diagnosis.	DDE16JAN26	
PRE_INT	CCIGR	CCI score group	'No comorbidity'; 'Mild comorbidity'; 'Moderate comorbidity'; 'Severe comorbidity'	if CCI = 0 then ='No comorbidity'; else if 1 <= CCI <= 2 then ='Mild comorbidity'; else if 3 <= CCI <= 4 then ='Moderate comorbidity'; else if CCI >= 5 then ='Severe comorbidity'		DDE16JAN26	
PRE_INT	CCIGRN	CCI score group (N)	1,2,3,4	Numeric of CCIGR: 1='No comorbidity'; 2='Mild comorbidity'; 3='Moderate comorbidity'; 4='Severe comorbidity'		DDE16JAN26	