Dr. Julian Bashir
Room 39, Ward 12, Delta Wing
Bajor General Hospital
59 Federation Street,
JB1 DS9

Mr. Ryan Example
13 Example Street,
EX1 9QB,
Exampleville

Dear Mr. Example,

I am writing to confirm your appointment with Dr. Julian Bashir at **10:15am on 15th September 2026** at Bajor General Hospital. Please save this date in your calendar.

What to do when you arrive

Upon arrival please visit reception and inform us that you have arrived. You will directed to a seating area and offered a drink while you wait. The hospital is fully accessible, and assistance is available if required. If you require assistance, please let us know via telephone at 0112 1234 5607.

What to do if you can no longer attend

If you can no longer attend your appointment, please contact us at **enquiries@bajorhospital.gov.**

We look forward to seeing you,

Dr. Julian Bashir

What is measles?



Measles is a very contagious illness caused by a virus. This means it spreads very easily to others.

How common is measles?

Measles is rare in the U.S. because so many people have been vaccinated against it. However, anyone in the U.S. who is unvaccinated can still get measles. Every year unvaccinated travelers bring measles to the U.S. from other countries. So even if you and your family do not travel outside the U.S., measles could still spread in your community.

How do people get measles?



Measles spreads very easily between people. Here's how it spreads:

1. A person sick with measles coughs or sneezes, sending tiny droplets containing the virus into the air.
2. Those tiny droplets then stay in the air or get onto a surface.
3. Another person then gets infected with measles when they:
   - Breathe in that air, or
   - Touch that surface then touch their eyes, nose, or mouth

Measles can be dangerous, especially for:

- Babies and children under age 5
- Adults older than age 20
- People who are pregnant
- People with weakened immune systems

What are the symptoms of measles?

- **1-2 weeks after being exposed to measles, first symptoms are usually:**



High fever (can be up to 104°F)



Cough



Runny nose



Red, watery eyes

- **2-3 days after symptoms start:**



Tiny white spots may appear in the mouth

- **3-5 days after symptoms start:**



A rash breaks out - it usually begins with flat red spots on the forehead. It then spreads down the face, neck, and rest of the body.

MENINGITIS
KNOW THE SYMPTOMS

WORLD MENINGITIS DAY APRIL 24
24 HOURS
Defeat Meningitis

CHILDREN & ADULTS



STIFF NECK

A stiff, sore neck can be a symptom of meningitis. Check for any of the other symptoms of meningitis.




HEADACHE

If you or your child has a severe headache, check for any of the other symptoms of meningitis.



FEVER

A fever, sometimes accompanied by cold hands and feet, can be a symptom of meningitis.



VOMITING

Vomiting is a common symptom of meningitis in both children and adults.



LIGHT SENSITIVITY

A sensitivity to light, or a dislike of bright lights is an early warning sign of meningitis. If this symptom is present, check for any of the other symptoms of meningitis.



DROWSINESS / CONFUSION

If the child or adult is difficult to wake, or in a drowsy and confused state, they could be exhibiting early symptoms of meningitis.



JOINT PAIN

Aching sore joints are a symptom of meningitis. If the person is experiencing pain in their joints check for any of the other symptoms of meningitis.



FITTING

Experiencing convulsive fits or seizures is characteristic of meningitis. Check for any of the other symptoms of meningitis.

Every Second Counts The person's condition may get worse very quickly so it is important to **trust your instincts** – if you suspect meningitis, seek medical help immediately

CALL 111 if it's a medical emergency
CALL 0800 611 116 to talk to a registered nurse for healthcare advice
CALL 0800 IMMUNE (0800 466 863) for immunisation facts

www.meningitis.org.nz | **info@meningitis.org.nz**

MENINGITIS FOUNDATION
AOTEAROA NEW ZEALAND