

# **Coherent Dijets in Ultra-peripheral Pb+Pb Collisions at**

$$\sqrt{s_{NN}} = 5.02 \text{ TeV}$$

By

**Samuel Boren**

Submitted to the graduate degree program in Department of People who read Abstracts and the Graduate Faculty of the University of Kansas in partial fulfillment of the requirements for the degree of Doctor of Philosophy.

---

MEMBER 1, Chairperson

---

MEMBER 2, Occasional Visitor

Committee members

---

MEMBER 3

---

MEMBER 4, The One Who Never Answers Email

---

The One with an Extra Long Name

---

The Fifth Beatle

Date defended: \_\_\_\_\_ October 02, 2016 \_\_\_\_\_

The Thesis Committee for Samuel Boren certifies  
that this is the approved version of the following thesis :

Coherent Dijets in Ultra-peripheral Pb+Pb Collisions at  $\sqrt{s_{NN}} = 5.02$  TeV

---

\_\_\_\_\_  
MEMBER 1, Chairperson

Date approved: \_\_\_\_\_ October 06, 2016 \_\_\_\_\_

## Abstract

This dissertation describes the study of coherent dijet photoproduction in ultra-peripheral heavy-ion collisions. Ultra-peripheral collisions have impact parameters larger  $2R$  where  $R$  is the radius of the heavy-ion. These collisions are mediated entirely by the exchange of low-virtuality photon according to the Weizacker-Williams approximation. The angular correlation of coherently photoproduced dijets is sensitive to elliptic gluon distribution. The azimuthal anisotropy of the ion recoil momentum  $P_T$  and the dijet transverse momentum  $Q_T$  is presented for two bins of  $Q_T$ :  $0 < Q_T < 5$  and  $5 < Q_T < 10$ .

## **Acknowledgements**

# Contents

<b>1 Ultra-relativistic heavy-ion collisions</b>	<b>1</b>
1.1 The Standard Model . . . . .	1
1.2 Quantum electrodynamics . . . . .	2
1.3 Quantum chromodynamics . . . . .	4
1.4 QCD experiments . . . . .	5
1.4.1 Deep inelastic scattering . . . . .	6
1.4.2 Quark gluon plasma . . . . .	8
1.5 Parton distribution functions . . . . .	9
1.5.1 Wigner distribution . . . . .	10
<b>2 Ultra-peripheral collisions at the Large Hadron Collider</b>	<b>14</b>
2.1 Ultra-peripheral heavy-ion collisions . . . . .	14
2.2 Vector meson photoproduction . . . . .	16
2.3 Photon-photon interactions . . . . .	19
2.4 Dijet photoproduction . . . . .	22
2.4.1 Factorization . . . . .	23
<b>3 The experiment</b>	<b>28</b>
3.1 Large Hadron Collider . . . . .	28
3.1.1 ATLAS . . . . .	28
3.1.2 ALICE . . . . .	29
3.1.3 LHCb . . . . .	29
3.1.4 TOTEM . . . . .	29

3.2	Compact Muon Solenoid . . . . .	30
3.2.1	Tracker . . . . .	31
3.2.1.1	Pixel tracker . . . . .	32
3.2.1.2	Strip tracker . . . . .	33
3.2.2	Electromagnetic calorimeter . . . . .	33
3.2.3	Hadronic calorimeter . . . . .	35
3.2.3.1	Hadronic forward calorimeters . . . . .	37
3.2.4	Muon detector . . . . .	37
3.2.5	Zero degree calorimeter . . . . .	38
3.2.6	Particle flow algorithm . . . . .	39
<b>4</b>	<b>Measuring luminosity</b>	<b>40</b>
4.1	Instantaneous luminosity . . . . .	40
4.2	Luminometers . . . . .	40
4.2.1	Pixel cluster counting . . . . .	41
4.2.2	Pixel luminosity telescope . . . . .	41
4.2.3	Hadronic forward calorimeter . . . . .	42
4.2.4	Fast Beam Conditions Monitor . . . . .	42
4.3	Van de Meer scanning calibration . . . . .	43
4.4	Corrections to VDM scans . . . . .	43
4.5	Author's contributions . . . . .	45
<b>5</b>	<b>Trigger development and performance</b>	<b>49</b>
5.1	Introduction to triggering . . . . .	49
5.2	Triggering at CMS . . . . .	49
5.3	Author's contributions . . . . .	51
5.4	Studies from trigger menus . . . . .	53
<b>6</b>	<b>Data analysis</b>	<b>54</b>

<b>7 Systematic uncertainties</b>	<b>59</b>
<b>8 Results and conclusions</b>	<b>60</b>
<b>A My Appendix, Next to my Spleen</b>	<b>78</b>

## List of Figures

1.1	QED Components of Feynman Diagrams (Plot from WikiCommons).	4
1.2	QCD Coupling Constant vs. $Q^2$ [1]	6
1.3	Collision Cross Section vs Bjorken-x, theory and data [2]	7
1.4	QCD Phase Diagram [3]	9
1.5	Subnuclear tomography (Ask Daniel again)	11
1.6	Interconnectedness of Parton Distributions [4]	12
1.7	Feynman Diagram of Coherent Dijets in Dipole Framework [5]	12
2.1	(a.) electromagnetic field of stationary charge (b.) eletromagnetic field of boosted charge [6].	15
2.2	$\gamma + \text{proton} \rightarrow J/\Psi + \text{proton}$ cross-section [7].	17
2.3	Dimuon invariant mass distributions of opposite-sign muon pairs ( $p_T < 1.0 \text{ GeV}$ , $1.8 <  y  < 2.3$ ) [8]	18
2.4	Dimuon transverse momentum distributions of opposite-sign muon pairs ( $p_T < 1.0 \text{ GeV}$ , $1.8 <  y  < 2.3$ ) [8]	19
2.5	Differential cross section versus rapidity for coherent $J/\Psi$ production in ultra-peripheral PbPb collisions at $\sqrt{s_{NN}} = 2.76 \text{ TeV}$ [8]	20
2.6	$\gamma\gamma$ Diagrams of light-by-light scattering, (a.) channels, (b.) analagous [9]	21
2.7	$\gamma\gamma$ kinematics, (a.) acoplanarity, (b.) invariant mass [9]	21
2.8	Event Display of ATLAS photon-photon scattering [10]	22
2.9	2 Figures side by side	24
2.10	Feynman diagrams for coherent jet photoproduction in (a) direct-photon in ep, (b) resolved-photon in ep [11]	24

2.11	Ratio of H1 data cross-section to NLO-QCD cross-section [11]	26
2.12	Beam envelope vs. distance to vertex in H1 [11]	26
3.1	CMS Detector [12]	30
3.2	CMS radial cross-section [13]	31
3.3	Pseudorapidity acceptance of tracker [14]	32
3.4	ECAL components [15]	34
3.5	Ecal energy resolution [16]	34
3.6	Ecal energy resolution [16]	35
3.7	HCAL energy resolution [16]	36
3.8	HCAL Components [16]	36
3.9	Pseudorapidity acceptance of muon detector [16]	38
3.10	Performance of particle flow algorithim compared to calorimeter readout [17]	39
4.1	PCC VdM Scans.	44
4.2	Systematic Uncertainty During 2015 pp Run.	45
4.3	Luminosity by CMS fill.	46
4.4	CMS Run Registry.	47
4.5	CMS lumi during HI-2015.	47
4.6	CMS Lumi by Era.	48
5.1	Rossi circuit with three Geiger-Muller coincidence circuits [18]	49
5.2	High multiplicity PbPb collision	50
5.3	UPC Upsilon candidate	51
5.4	Example of trigger rate.	52
6.1	Raw $\phi$ distribution.	55
6.2	Mixed $\phi$ distribution.	56
6.3	Subtracted $\phi$ distribution.	57

6.4 Final $\phi$ distribution. . . . .	58
--	----

## **List of Tables**

# **Chapter 1**

## **Ultra-relativistic heavy-ion collisions**

### **1.1 The Standard Model**

The Standard Model describes the fundamental particles of the universe in terms of fermions and bosons. Fermions are particles with half-integer spin, while bosons have integer-spin. This difference in spin has far reaching consequences. Fermions must obey the Pauli Exclusion Principle: only one fermion at a time can occupy a given state. However, multiple bosons can simultaneously occupy a specific state. Subnuclear particles – protons and neutrons – contain large numbers of both fermions and bosons.

Among the fermions are the leptons, neutrinos, and quarks. The leptons consist of the electron, muon, and tau, as well as their anti-particles. The leptons are seemingly fundamental: high energy experiments have yet to observe internal lepton-structure. Neutrinos are weakly interacting particles detected primarily through the precise measuring of missing transverse energy in the products of particle collisions. Quarks are the constituent particles of baryons, which contain three valence quarks, and mesons, which contain two valence quarks. In addition to the valence quarks are the sea quarks, which appear and disappear as quark-antiquark pairs within hadrons. The hadrons are particles made of quarks and gluons. Gluons are particles that mediate the strong nuclear force; likewise, photons mediate the electromagnetic force, and weak-gauge bosons mediate the weak nuclear force. A fourth boson, the graviton, is expected to transmit the gravitational force, but this particles existence has not been verified.

The behavior of fundamental particles is best described within the framework of quantum field theory (QFT). QFT defines a Lagrangian for fundamental particles. This Lagrangian then predicts

the outcome of particle collisions. Different terms in the Lagrangian correspond to the various interactions between particles. The Standard Model Lagrangian,  $\mathcal{L}_{StandardModel}$  can be broken down into three basic terms:

$$\mathcal{L}_{StandardModel} = \mathcal{L}_{QED} + \mathcal{L}_{QCD} + \mathcal{L}_{Higgs} + \dots, \quad (1.1)$$

Where  $\mathcal{L}_{QED}$  is the QED Lagrangian,  $\mathcal{L}_{QCD}$  is the QCD Lagrangian, and  $\mathcal{L}_{Higgs}$  is the Higgs Lagrangian. The QED and QCD Lagrangians will be the most important in what follows [19].

The most accessible approach to quantum field theory is through the use of Feynman diagrams. First, one imagines an interaction between particles. Then, one draws this process into a Feynman diagram, which is essentially a pictorial representation of exchanges between particles. The Lagrangian can be interpreted into Feynman rules. These rules describe how the Feynman diagram translates into a calculation for the quantum mechanical amplitude of the process. The quantum mechanical amplitude, in turn, is proportional to the cross-section of the process. This is important because, since the cross-section is invariant between experiments, one can use it to effectively test for invariant quantities in the Lagrangian [20].

## 1.2 Quantum electrodynamics

In classical electrodynamics, all of electrical and magnetic phenomena can be derived from four fundamental equations: Maxwell's laws. Below these equations are given in their differential form.

Gauss's law relates the divergence of a static electric field,  $\vec{E}$ , to its source charge density,  $\rho$ ,

$$\nabla \cdot \vec{E} = \frac{\rho}{\epsilon_0}, \quad (1.2)$$

where  $\epsilon_0$  is permittivity of free space. Gauss's law for a static magnetic field,  $\vec{B}$ , reveals the stark contrast between electricity and magnetism. The divergence of a static magnetic field is always a net 0. This reflects the seeming non-existence of magnetic monopoles.

$$\nabla \cdot \vec{B} = 0 \quad (1.3)$$

Faraday's law describes the curl of an electric field in terms of the dynamics of a magnetic field. The electric field and the magnetic field are coupled, and a change in one will affect the properties of the other. For a given electric field,  $\vec{E}$ , the curl is equal to the partial derivative of a magnetic field,  $\vec{B}$ , with respect to time,  $t$ ,

$$\nabla \times \vec{E} = -\frac{\partial \vec{B}}{\partial t}. \quad (1.4)$$

Ampere's law addresses the curl of a magnetic field,  $\vec{B}$ , and once more the absence of magnetic monopoles gives rise to distinct properties. In addition to the partial derivative of  $\vec{E}$  with respect to time  $t$ , there is a corrective term for the electric charge current density,  $\vec{J}$ , and the sum is multiplied by  $\mu_o$ , the permeability of free space,

$$\nabla \times \vec{B} = \mu_o \left( \vec{J} + \epsilon_o \frac{\partial \vec{E}}{\partial t} \right). \quad (1.5)$$

The complementarity of the classical electric and magnetic fields gives rise to the photon flux so important for ultra-peripheral collisions, and will be discussed in greater detail in the next chapter.

$$\mathcal{L} = \bar{\psi}(i\gamma^\mu D_\mu - m)\psi - \frac{1}{4}F_{\mu\nu}F^{\mu\nu} \quad (1.6)$$

Quantum electrodynamics (QED) is a theory of electromagnetic interaction in terms of relativistic quantum field theory. QED addresses three specific processes: photon motion, electron motion, and the emission, or absorption, of a photon by an electron. To do this, first a Lagrangian is established based on Maxwell's laws and quantum mechanics. The photon constitutes a spin-1 solution to Maxwell's equations. Likewise, electrons are described, at non-relativistic scales, by the Schrodinger equation, at relativistic scales by the Dirac equation. Figure 1.1 shows the resulting processes allowed by QED: the translation of an electron, the translation of a photon, and the scattering of a photon off an electron [21].

The QED coupling constant,  $\alpha_{QED}$ , is approximately 1/137 at perturbative scales. However, at small scales, i.e. non-perturbative momentum transfers  $Q^2$ , the coupling constant increases:

$$\alpha_{QED}(Q^2) = \frac{\alpha_{em}}{(1 - \frac{\alpha_{em}}{3\pi})\ln(\frac{Q^2}{m^2})}, \quad (1.7)$$




Figure 1.1: QED Components of Feynman Diagrams (Plot from WikiCommons).

where  $\alpha_{QED}(Q^2)$  is the QED coupling constant at high  $Q^2$ ,  $\alpha_{em}$ , and  $m$  is the electron mass. High values of  $Q^2$  are non-perturbative because, at small scales and high momentum, the fluctuation of photons into electron-positron pairs begins to saturate.

### 1.3 Quantum chromodynamics

The quarks are a family of fermions that compose the baryons and the mesons. Baryons consist of three quarks in a color neutral state, while mesons consist two quarks in a color neutral state. "Color" in this context refers to the six kinds of strongly-interacting charge available to quarks: red and anti-red, blue and anti-blue, and green and anti-green. Color charge has no relation to optical phenomena, but provides a useful analogy for the stable combinations of quarks. The net color-charge of a baryon or meson is colorless. By way of analogy, a red quark, green quark, and blue quark can together form a hadron, in the way that conventional red, green, and blue can together form white [22].

Gluons are the QCD analogues of the photons in QED. Gluons are spin-1 and massless, but unlike photons, which do not carry electromagnetic charge, gluons carry strongly-interacting charge: color. Color comes six varieties: red, antired, blue, antiblue, green, and antigreen [23].

Unlike QED, the QCD coupling increases with distance [1]. Figure 1.2 shows the running of the QCD coupling with  $Q^2$ :

$$\alpha_{QCD}(Q^2) = \frac{4\pi}{(11 - \frac{2}{3}n_f)\ln(\frac{Q^2}{\Lambda_{QCD}^2})}, \quad (1.8)$$

where  $n_f$  is the number of quark flavors,  $Q^2$  is the momentum transfer, and  $\Lambda_{QCD}^2$  is the mass scale. This has the practical consequence of the strong-interactions being stronger in high momentum transfer collisions. The direct results of the running QCD coupling are the dual phenomena of asymptotic freedom and color confinement. At large distances, string tension describes the binding force of the quarks. At short distances, however, Coulomb-like interactions dominate. The QCD coupling constant can be measured via the cross-section of inelastic proton-proton collisions, and also the cross section of electron-positrons into triple jets.

Within the nucleus, a proton can be thought of as a bubble in a vacuum. Debrye screening exerts a pressure on the proton. This pressure is responsible for the size of the proton.

$$\mathcal{L}_{QCD} = \bar{\psi}_i (i(\gamma^\mu D_\mu)_{ij} - m \delta_{ij}) \psi_j - \frac{1}{4} G_{\mu\nu}^a G_a^{\mu\nu} \quad (1.9)$$

## 1.4 QCD experiments

Scattering experiments are the basic tool for exploring the nucleus. The Large Hadron Collider (LHC) is capable of reaching heavy-ion collision energies of up to 7 TeV per nucleon-nucleon. The higher the energy, the more experiments can probe the nuclear phase-space diagram. Momentum transferred, expressed as  $Q^2$ , is an important quantity for characterizing QCD measurements. In addition to  $Q^2$ , Bjorken-x, also known as Bjorken-scaling is necessary to describe the nuclear phase space. Bjorken-x represents the momentum fraction of partons [24].




Figure 1.2: QCD Coupling Constant vs.  $Q^2$  [1]

At the turn of the century, Ernst Rutherford probed the gold atom by bombarding a gold sheet with alpha-particles, i.e. helium nuclei. The angular distribution of the scattered alpha-particles demonstrates that the mass of the atom is concentrated in a small volume, i.e, the atom is mostly empty space. Further experiments revealed that the atomic nuclei consisted of separate positively and neutrally charged particles: protons and neutrons.

### 1.4.1 Deep inelastic scattering

Deep inelastic scattering commonly refers to the scattering of a leptons off hadrons. Experiments at HERA focused on electron-proton collisions. In these collisions, the electron was used as a source of photons and neutrinos. When these particles scatter off the proton, the dependence of the collision cross section, on momentum transfer and scattering angle of the source electron, reflects the structure of the proton. These experiments provided the first evidence of two phenomena: the parton model and Bjorken-scaling.

The parton model, first proposed by Richard Feynman, posits that hadrons in general, and nucleons in specific, are made of more fundamental constituent particles which may or may not be the quarks implied by the SU(3) symmetry. In addition to the quarks, the partons also include any field quanta associated with nuclear forces. In time, these field quanta are dubbed "gluons".

"Scaling" is an interpretation of the data from deep inelastic scattering (DIS). First proposed by James Bjorken, scaling is reflected in the incoherence of photon-proton interactions at photon energies above  $1 \text{ GeV}/c$ . Predictions from perturbative QCD are in good agreement with DIS data from HERA, as seen in figure 1.3 [2]. In this graph the Bjorken-x momentum fraction is designated  $x$ , and  $\sigma_r$  represents the  $F_2$  structure function, and  $Q^2$  is the transferred momentum from the electron to the proton. The order of magnitude of  $\sigma_r$  set by the Bjorken-x.




Figure 1.3: Collision Cross Section vs Bjorken-x, theory and data [2]

At the small scales probed by high energy photons, the decreasing QCD coupling causes quarks and gluons to interact weakly. This phenomena is called "asymptotic freedom". Because gluons themselves carry color charge, the gluons about a quark tend to have an anti-screening behavior: the gluon color adds to the quark color, increasing the net color charge of the area. At smaller

distances to a quark, then, there are fewer and fewer gluons augmenting the color interaction.

### 1.4.2 Quark gluon plasma

The modern understanding of subnuclear physics is based on results from three laboratories: Lawrence Berkeley National Laboratory (LBNL), Brookhaven National Laboratory (BNL), and CERN. Fixed target experiments hinted at the existence of a quark gluon plasma, but the collision energies were too low for this state to last. These experiments confirmed the developing model of the QCD phase space; see figure 1.4 [3]. Essentially, quark matter organizes itself differently depending on temperature and baryon density. At low energies, quark matter exists in bound states: the hadrons. However, in the high energy limit, quarks and gluons take the form of a strongly interacting plasma: the quark-gluon plasma (QGP). The QGP represents the extreme case of asymptotic freedom; the QCD coupling constant becomes small enough that quarks and gluons no longer behave as bound states. There are two ways of achieving the high energies necessary to form QGP. High baryon densities cause the quarks of separate hadrons to interact at small distances where asymptotic freedom takes effect. It is not currently possible to achieve these densities in laboratory experiments, though this state is thought to occur in neutron stars. By contrast, it particle collider's like the LHC increase the energy density by colliding heavy-ions at ultra-relativistic velocities. The high temperature environment thus produced manifests QGP. The early universe, mere milliseconds after the Big Bang, is thought to have existed as QGP.

There are a number of experimental signatures of QGP. Of particular interest are charm suppression and strangeness enhancement, elliptic flow, jet quenching. The QGP is thought to suppress the production of  $J\Psi$  mesons in heavy-ion collisions. The predicted viscosity of QGP would cause elliptic flow in the overlap region of heavy-ion collisions. Lastly, hadronic jets would interact strongly with the QGP; therefore, dijets will have significant energy imbalance depending on the multiple interactions of the components jet with the QGP. All of these cases require a good understanding the heavy-ion initial state as a basis for comparison.




Figure 1.4: QCD Phase Diagram [3]

## 1.5 Parton distribution functions

QCD describes the interaction between quarks and gluons, but within a nucleon the calculations are too complicated to solve for the behavior of each individual parton. Theorists compromise using the factorisation theorem to use data from DIS experiments to make predictions. The factorisation theorem is discussed in greater detail later in this theorist.

Parton distribution functions (PDFs) are a method of encoding data from DIS experiments in the form of probability density. PDFs give the probability of finding a species of parton with given momentum fraction,  $x$ , and a given squared energy scale,  $Q^2$ .

In electron-proton deep inelastic scattering, the electron interacts with the proton electromagnetically by the emission of a virtual photon. This virtual photon has a four momentum  $q$  and a virtuality  $Q^2 = -q^2$ . When virtuality is low, the virtual photon is approximately "real"; these quasi-photons are discussed in greater detail in the next chapter. The virtual photon, originating from the e. lectrion, interacts with a parton and changes its initial momentum fraction,  $x$ . The  $Q^2$  of the probing virtual photon and Bjorken-x of interacting parton are determined from the collision products. The whole data sample can be used to fill out a probability density,  $F_i(x, Q^2)$ , of finding a parton species  $i$  at a given momentum fraction  $x$  with a probe of virtuality  $Q^2$ .

### 1.5.1 Wigner distribution

The Wigner distribution was first developed as part of an attempt by Eugene Wigner to map solutions to Schrodinger equation into phase probability distributions, i.e. a statistical mechanics interpretation of quantum mechanics.

$$P(x, p) \stackrel{\text{def}}{=} \frac{1}{\pi\hbar} \int_{-\infty}^{\infty} \psi^*(x+y)\psi(x-y)e^{2ipy/\hbar} dy \quad (1.10)$$

The Wigner distribution  $P(x, p)$  can be used to calculate the expectation value of any given variable by considering the Wigner transformation  $g(x, p)$  of that variable's operator,  $\hat{G}$ ,

$$\langle \hat{G} \rangle = \int dx dp P(x, p) g(x, p) . \quad (1.11)$$

Notice that in order to derive expectation values, the probability distribution must be integrated with respect to a function of position or momentum, which are still non-commuting according to the uncertainty principle.

One can use the Wigner distribution to tomographically image the internal structure of the nucleus [5]. The nucleus manifests different structures at varying momentum fractions; specifically, small momentum fractions exhibit gluon saturation [25]; see figure 1.5 for an illustration of how the nucleus appears at varying momentum fractions.

The quantum field theory lagrangian of the strong interaction is relatively simple, but because of confinement and asymptotic freedom the hadronic bound states are too complex for an analytic solution. Furthermore, collider experiment data requires a quantitative interpretation to be useful. The gap between QCD and heavy-ion data is bridged using the parton model, which considers hadrons as composed of quarks and gluons. Parton density functions (PDFs) model the longitudinal momentum distribution of the partons. PDFs are supplemented by transverse momentum distributions (TMDs) and generalized parton distributions (GPDs). In addition to transverse momentum, GPDs describe the transverse spatial distribution. TMDs and GPDs are derived from the final state particles of a collision. Markus Diehl maps the relationship between various distribution functions in figure 1.6 [4].




Figure 1.5: Subnuclear tomography (Ask Daniel again)

The Wigner distribution is a quantum phase space distribution that describes elliptic gluons [26]. Specifically, by considering the color dipole scattering amplitude, the angular correlation of the nucleon recoil momentum and the dijet transverse momentum can provide a three-dimensional, tomographic image of the gluons within a high energy nucleus. This tomographic image takes the form of a Wigner distribution, which contains all the information of both TMDs and GPDs without violating the uncertainty principle. Specifically, the angular correlation directly measures the Fourier transform of the gluons. This is possible because the dipole amplitudes are functions of the impact parameter, and because collinear factorization holds.

TMDs and GPDs manifest non-perturbative QCD effects. The Wigner distribution, at this scale, reflects the relationship between the position and momentum of partons. Integrating the Wigner function over the transverse distance yields the TMD, while integrating over transverse momentum yields a GPD with spatial information.

Yoshitaka Hatta uses the dipole framework to show that the azimuthal angular correlations of coherent dijets are generated by the underlying Wigner distribution of the small- $x$  gluons. Further-




Figure 1.6: Interconnectedness of Parton Distributions [4]

more, these correlations are consistent with predictions based on standard collinear factorization.

Relevant kinematic variables are mapped in the figure 1.7.




Figure 1.7: Feynman Diagram of Coherent Dijets in Dipole Framework [5]

It is expected that the dominant contribution to the angular correlation is the elliptic, cor-

responding to  $n = 1$  in the Fourier transform. The interior of the proton displays an intricate structure.

# Chapter 2

## Ultra-peripheral collisions at the Large Hadron Collider

In this chapter, I will review recent LHC results on ultra-peripheral collisions (UPCs). First I will discuss the unique features of UPCs, and how they are selected. I then present the results from ALICE and CMS on vector meson photoproduction as an example of how UPCs can probe the nuclear gluon distribution. The ATLAS results for light-by-light scattering are described. The final section explains dijet photoproduction in UPCs, and how the factorisation theorem informs the interpretation of coherent dijet correlations.

### 2.1 Ultra-peripheral heavy-ion collisions

Similar to the Rutherford experiment, in heavy-ion collisions the scattered particles carry information about the internal structure of the nucleus. The Rutherford experiment has the three components that still characterize high-energy nuclear experiments: a probe, a medium, and a signal. Alpha particles probe the medium of the gold atom, and the angular distribution of scattered alpha particles signals the internal structure of the atom.

Ultra-peripheral collisions occur at impact parameters greater than the sum of the heavy-ion radii. In these collisions, hadronic interactions are strongly suppressed while photonuclear activity is enhanced proportional to the square of the nuclear charge. The electromagnetic field of an incoming heavy-ion, from the perspective of a target, is equivalent to a flux of virtual photons; figure 2.1 illustrates the Lorentz contraction of the field of a boosted charge.

UPC models typically address two elements: the photon flux,  $N_{\gamma/Pb}$ , and the photoproduction cross-section,  $\sigma_{\gamma Pb}$ . These quantities are related to the UPC cross-section,  $\sigma_{PbPb}$ , in the following




Figure 2.1: (a.) electromagnetic field of stationary charge (b.) electromagnetic field of boosted charge [6].

equation,

$$\frac{d\sigma_{PbPb}}{dy} = N_{\gamma/Pb}(y, M)\sigma_{\gamma Pb}(y) + N_{\gamma/Pb}(-y, M)\sigma_{\gamma Pb}(-y), \quad (2.1)$$

where there are two terms to reflect that either heavy-ion can be a source of photons.

The Weizsacker-Williams appoximation (WWA) calculates the density of photons, about the nucleus, as a function of energy. WWA is a semi-classical formulation. Maxwell's equations are solved for a stationary point charge boosted to an ultra-relativistic velocity. In the target's frame, the Fourier transform of the source field is taken. The Fourier frequency modes are interpreted through the quantum mechanical equation of photon energy. The photon flux as function of energy is given by

$$N(\omega, b) = \frac{\alpha}{\hbar\omega} \left( \frac{Z}{b\beta\pi} \right)^2 \left( \frac{\omega b}{\gamma v} \right) K_1^2 \left( \frac{\omega b}{\gamma v} \right), \quad (2.2)$$

where  $\alpha$  is the QED coupling constant,  $\omega$  is the photon energy,  $Z$  is the atomic number of the nuclei,  $b$  is the impact parameter,  $\beta$  is ratio of the nuclei speed to the speed of light,  $\gamma$  is the Lorentz boost of the nuclei,  $K_1^2$  is a Bessel function,  $v$  is the photon frequency [6].

There are ways to represent the photon flux as solely a function of photon wavelength:

$$n(k) = \frac{2\alpha Z^2}{\pi} \left[ \xi K_0(\xi) - \frac{\xi^2}{Z} (K_1^2(\xi) - K_0^2(\xi)) \right] \quad (2.3)$$

Gluons are the particle exchanged in strong interactions. However, gluons themselves carry

color charge. By analogy, photons transmit the electromagnetic force, but do not themselves have an electric charge. The gluons are spin-1, meaning that more than one can occupy the same quantum state.

When a quark is scattered from a nucleus, the strong force field lines for a tube between the quark and its counterpart. Given that the strong coupling constant increases with distance the strong interaction gathers potential energy until the threshold for quark production is passed, at which point an anti-quark is generated to screen the ejected quark. These quarks continue to separate from each other and continue the quark generation process. The resulting final state hadron distribution takes the form of a cone centered on the path of the initial quark.

QCD factorisation describes the diffractive-photoproduction dijet cross-section as the convolution of the partonic cross-section with the diffractive parton distributions. However, factorisation only describes H1 data if the resolved-photon contribution is suppressed.

The photoproduction cross-section is proportional to the gluon distribution. At low momentum transfer, photons interact electromagnetically, i.e. directly, with partons. High energy, "resolved" photons possess a hadronic structure; instead of directly interacting with the nuclei, these photons fluctuate into mediating quark-antiquark pairs.

## 2.2 Vector meson photoproduction

The virtual photons present in UPC can fluctuate into a quark-antiquark pair which can take the form of a low mass meson. This meson then interacts with the target nuclei via colorless gluon exchange, emitting a vector meson. If the virtual photon interacts coherently with the target nucleus, this is reflected in the transverse momentum of the vector meson. The vector meson decays into a dilepton pair detectable by CMS.

For a given final state  $X$ , the collision cross section is given as

$$\sigma_X = \int d\omega \frac{n(\omega)}{\omega} \sigma_X^\gamma(\omega), \quad (2.4)$$

where  $n(\omega)$  is the number of photons emitted at an energy  $\omega$ , and  $\sigma_X^\gamma(\omega)$  is the photonuclear cross-

section of the photon-nucleon interaction. The Weizacker-Williams approximation provides  $n(\omega)$ . The integral reflects how a state  $X$  can result from the interaction of a quasi-real photon of varying energy.

Ultra-peripheral coherent  $J/\Psi$  photoproduction was studied in the ALICE 2011 Pb-Pb data. Notice that the cross-section of coherent  $J/\Psi$  photoproduction, for  $\gamma + p \rightarrow J/\Psi + p$ , has a power-law dependence on the photon-proton center-of-mass energy, as seen in figure 2.2 [7]. This data can be used to measure the gluon distribution as a function of Bjorken-x.




Figure 2.2:  $\gamma + \text{proton} \rightarrow J/\Psi + \text{proton}$  cross-section [7].

CMS has studied  $J/\Psi$  and  $\rho$  photoproduction off the proton in proton-Pb collisions. In coherent UPC photoproduction of  $J/\Psi$ , the transverse momentum distribution peaks at about 60 MeV because, for coherent production, the photon energy is inversely proportional to the Pb ion diameter,  $1/2R_{Pb}$ . For this process the photon is interacting with the whole nucleus. The low momentum corresponds to large wavelength of the photon with respect to the nuclear radius.

UPC vector mesons are a clean probe of the nuclear initial state. The high temperature environment of heavy-ion collisions produces a near viscosity free medium that exhibits collective flow. However, collective flow phenomena are also observed in heavy-ion collisions below the QGP phase transition. A good understanding of the heavy-ion initial state, before the phase transition, is necessary to detect authentic QGP phenomena. UPC productions can probe the initial




Figure 2.3: Dimuon invariant mass distributions of opposite-sign muon pairs ( $p_T < 1.0 \text{ GeV}$ ,  $1.8 < |y| < 2.3$ ) [8]

state because the interaction is purely electromagnetic; hadronic activity is strongly suppressed, excluding the possibility of QGP forming.

The UPC  $J/\Psi$  in particular is sensitive to the gluon distribution at low Bjorken-x. The UPC  $J/\Psi$  photoproduction cross-section can be calculated from a Glauber model of the heavy-ion. The Glauber model approach depicts the nucleus as a sum of protons and neutrons by scaling up the photon-nuclear cross section derived from electron-proton collisions.

Studies of UPC  $J/\Psi$  at CMS show that the measured cross section is consistent with models of the nucleus that include moderately strong gluon shadowing, in particular EPS09. Fig 2.5 compares the cross section measured by CMS and ALICE against theoretical models. In the data there is a clear difference from the Glauber model prediction. The cross sections indicate that, at the energy scale of the  $J/\Psi$  mass, the nuclear gluon density is suppressed with respect to that of the proton.




Figure 2.4: Dimuon transverse momentum distributions of opposite-sign muon pairs ( $p_T < 1.0$  GeV,  $1.8 < |y| < 2.3$ ) [8]

### 2.3 Photon-photon interactions

Classical electrodynamics forbids the scattering of a photon off another photon. According to Maxwell's equations, charge and current distributions make linear contributions to the electric and magnetic fields. For example, the electric field generated by a grid of static charges is the sum of the electric fields of the each charge considered by itself. This property, called "the principle of superposition", allows Maxwell's equations to be solved as boundary values problems via the separation of variables.

The cross section of a final state from the photon-photon interaction is given by

$$\sigma_X = \int d\omega_1 d\omega_2 \frac{n(\omega_1)}{\omega_2} \sigma_X^{\gamma\gamma}(\omega_1, \omega_2), \quad (2.5)$$

where the subscripts 1 and 2 designate the quasi-real photons that are colliding and  $\sigma_X^{\gamma\gamma}(\omega_1, \omega_2)$  is the cross-section of the final state from a collision of photons with energies  $\omega_1$  and  $\omega_2$ . Once more, notice that the total final state cross-section is an integral of the all the photon energies in




Figure 2.5: Differential cross section versus rapidity for coherent  $J/\Psi$  production in ultra-peripheral PbPb collisions at  $\sqrt{s_{NN}} = 2.76\text{TeV}$  [8]

the photon flux.

Maxwell's equations, being a classical theory, do not take into account the quantum mechanical effects that manifest at the distance scales of sub-atomic particles. The polarization of the vacuum is one such consequence of quantum mechanics. Around a photon there is a cloud of particle-antiparticle pairs, appearing together and then annihilating each other after a time proportional to their energy. In so far as these particles have an electric charge, portions of the local area may carry a non-zero electric charge. Thus, two photons may scatter off each other in three possible exchanges: Diagrams for Delbrück scattering, photon splitting, and elastic light-by-light scattering.




Figure 2.6:  $\gamma\gamma$  Diagrams of light-by-light scattering, (a.) channels, (b.) analogous [9]

Figure 2.6 illustrates the process of light-by-light scattering. The plots in figure 2.6 (a.) are the Feynman diagrams of the three light-by-light channels: Delbrück scattering, photon splitting, and elastic scattering, respectively. Notice that all three channels are variations on a closed electron-loop. Figure 2.6 (b.) shows how these processes arise in heavy-ion collisions. The photon flux from a relativistic heavy-ion is calculated from the Fourier transform of its electric field. The final state  $X$  is empty except for two photons [9].




Figure 2.7:  $\gamma\gamma$  kinematics, (a.) acoplanarity, (b.) invariant mass [9]

Light-by-light scattering has been observed at ATLAS. The relevant data sample contains

$480 \mu b^{-1}$  of Pb+Pb data taken at  $\sqrt{s_{NN}} = 5.02$  TeV. Light-by-light MC was generated using STARLIGHT and processed through a GEANT4 simulation of ATLAS. The reconstructed MC was used to create kinematic templates for comparison against the data. These templates can be seen in figure 2.7. The most important part of the event selection is the cut on diphoton acoplanarity.




Figure 2.8: Event Display of ATLAS photon-photon scattering [10]

Figure 2.8 is an event display for an ATLAS light-by-light candidate [10]. Notice that the two photons are back-to-back in the plane perpendicular to the beam.

## 2.4 Dijet photoproduction

An important experimental signature of deconfinement is how it affects hard-scattering processes. In a hard scattering process, partons are fragmented off from the nucleus and, due to confinement, hadronize into a cone of correlated final state particles, known as a jet. In theory, a jet should be sensitive to the medium that the mothering parton passes through. However, in order to detect the modification of a jet due to its mother parton passing through QGP, one needs a precise measurement of the jet production in of nuclear PDFs but not QGP. One can then determine QGP effects

by comparison to the nuclear PDF baseline. However, there is a significant lack of data for low-x nuclear PDF. As Bjorken-x decreases, the nuclear PDF passes through stages three stages: EMC effects, anti-shadowing, and shadowing. EMC effects and shadowing are suppressions of the nuclear PDF in comparison to the proton PDF; anti-shadowing, by contrast, is the enhancement of the nuclear PDF in comparison to the proton PDF.

The dijet photoproduction cross-section, like that of vector meson photoproduction, displays diffractive dips in its  $|t|$  dependence. Furthermore, according to the color glass condensate formalism, coherent dijet production is sensitive to gluon saturation effects at small Bjorken-x values. Gluon saturation would affect the color-dipole orientation of nucleons in the transverse plane. This effect should be reflected in an enhanced azimuthal angle correlation of the jets.

ATLAS has published preliminary results on UPC dijet production. It has been shown that these photonuclear dijets manifest distinct differences from dijets produced in more "hadronic" collisions. There are rapidity gaps that are similar to those seen in the diffractive productions of electron-proton collisions at HERA. The forward neutron emission spectrum also matches that expected for UPC data set. The ATLAS results are particularly interesting because Strikman, Vogt, and White have shown that UPC dijet photoproduction is sensitive to the low Bjorken-x nuclear PDFs.

ATLAS used a combination of triggering, forward calorimetry, and rapidity gap requirements to select events of photo-nuclear dijet production in ultra-peripheral heavy-ion collisions.

ATLAS reported the cross-section of photo-nuclear dijet production versus various dijet kinematic variables, and compared these cross-sections to those derived from MC. The MC sample was generated by Pythia and re-weighed to fit the quasi-real photon spectrum of a relativistic Pb ion. The photon spectrum itself is taken from the STARLight MC generator.

#### 2.4.1 Factorization

Diffractive dijet photoproduction is not describable in perturbative QCD. For coherent processes the photon energy is small, and therefore the wavelength is large compared to the size of the nu-




Figure 2.9: 2 Figures side by side

cleus. At these large distances, there isn't a hard scale, and so perturbation calculations cannot be done. Gluon splitting interactions dominate the low Bjorken- $x$  partons. QCD collinear factorization describes these soft interactions via the convolution of parton cross sections, taken from perturbative QCD, and diffractive parton distribution functions, taken from experiment [11].




Figure 2.10: Feynman diagrams for coherent jet photoproduction in (a) direct-photon in ep, (b) resolved-photon in ep [11]

In electron-hadron collisions, diffractive photoproduction is characterized by the presence of a

large rapidity gap in the final state and an intact nucleus. The Feynman diagram of electroproduction in lepton-hadron collisions is similar to that of photoproduction in ultraperipheral collisions, as seen in figure 2.10 [11]. The diffractive dijet cross section is expressed by the convolution of partonic cross sections  $d\hat{\sigma}$  and diffractive PDFs  $f_{i/p}^D$ .

$$d\sigma(ep \rightarrow e + 2\text{jets} + X' + p) = \sum_i \int dt \int dx_{\mathbb{P}} \int dz_{\mathbb{P}} d\hat{\sigma}_{ei \rightarrow 2\text{jets}}(\hat{s}, \mu_R^2, \mu_F^2) \times f_{i/p}^D(z_{\mathbb{P}}, \mu_F^2, x_{\mathbb{P}}, t), \quad (2.6)$$

where  $x_{\mathbb{P}}$  is longitudinal momenetum fraction lost by the incoming proton,  $d\hat{\sigma}_{ei \rightarrow 2\text{jets}}$  is the partonic cross-section for the process,  $z_{\mathbb{P}}$  is the longitudinal momentum fraction of the pomeron entering the hard process,  $\hat{s}$  is the squared invariant energy of the subprocess,  $\mu_R^2$  is the squared renormalization scale,  $\mu_F^2$  is the squared factorisation scale,  $f_{i/p}^D$  is the diffractive parton distribution, and  $t$  is the four-momentum transfer squared at the vertex.

In the proton-vertex factorisation hypothesis, the dependence on  $x_{\mathbb{P}}$  and  $|t|$  is factored out of the dependence on  $\mu_F^2$  and  $z_{\mathbb{P}}$ . Furthermore,  $f_{i/p}^D$  is sum of contributions from the pomeron and Reggeon:

$$f_{i/p}^D(z_{\mathbb{P}}, \mu_F^2, x_{\mathbb{P}}, t) = f_{\mathbb{P}/p}(x_{\mathbb{P}}, t) f_{i/\mathbb{P}}(z_{\mathbb{P}}, \mu_F^2) + n_{\mathbb{R}} f_{\mathbb{R}/p}(x_{\mathbb{P}}, t) f_{i/\mathbb{R}}(z_{\mathbb{P}}, \mu_F^2), \quad (2.7)$$

where  $\mathbb{P}/p$  is the pomeron flux factor,  $f_{\mathbb{R}/p}$  is the Reggeon flux factor,  $n_{\mathbb{R}}$  is the normalization factor of the Reggeon,  $f_{i/\mathbb{P}}$  is the pomeron parton distribution, and  $f_{i/\mathbb{R}}$  is the Reggeon parton distribution.

Lepton-hadron collisions were performed at HERA and measured by the H1 experiment. These experiments reported a value for the total diffractive photoproduction cross section that is double that predicted by QCD collinear factorization; figure 2.11 compares the cross-section of H1 data to that predicted by NLO-QCD [11]. Diffractive events were selected for using rapidity gaps or the presence of leading protons in the very forward proton spectrometer (VFPS).

H1 used the Very Forward Proton Spectrometer (VFPS) to trigger on low  $Q^2$  protons. The VFPS consists of two Roman Pots located 218 m and 222 m from the H1 interaction-point in the forward direction. The VFPS can detect protons scattered at very low transverse momentum, corresponding to  $0.008 < x_P < 0.028$  and  $|t| < 0.6$ . Each of the Roman Pots contains layers




Figure 2.11: Ratio of H1 data cross-section to NLO-QCD cross-section [11]

of scintillating fibers, which are covered by a layer of scintillator tiles. The fibers readout to photomultipliers, and the tiles both shield from radiation and trigger on protons. The track efficiency of VFPS is a remarkable 96%, and the background contamination is kept at 1% , making the detector excellent for studying diffractive events. Figure 2.12 shows the  $|t|$  coverage of the Forward Proton Spectrometer (FPS) and VFPS [11].




Figure 2.12: Beam envelope vs. distance to vertex in H1 [11]

The H1 data was compared to predictions based on NLO-QCD convoluted with diffractive parton distribution functions (DPDFs) from HERA inclusive diffractive deep-inelastic scattering (DDIS) data. For diffractive pp collisions the high transverse momentum jets yield a hard scale for perturbative QCD.

# **Chapter 3**

## **The experiment**

### **3.1 Large Hadron Collider**

The Large Hadron Collider (LHC) has a radius of approximately 27 kilometers. As of this writing, it is the largest machine ever constructed. The initial purpose of the LHC was to discover the Higgs boson, but it is capable of investigating a variety of other physics phenomena, such as dark matter, extra-dimensions, and heavy-ion physics.

The beams are accelerated along a circular path using radio frequency cavities, gaining energy with each revolution. LHC is a hadron collider, meaning it is designed to collide particles made of quarks and gluons. The proton-proton, proton-Pb, and Pb-Pb collision energies are the largest ever probed experimentally. The LHC is a circular collider.

Heavy-ion collisions at LHC produce strongly interacting nuclear matter. The temperature and density of this matter is comparable to the state of the universe only a few milliseconds after the Big Bang.

#### **3.1.1 ATLAS**

"ATLAS" stands for "A Toroidal LHC ApparatuS". ATLAS was designed as a general purpose detector for LHC physics. Three toroidal superconducting magnets provide the field for particle identification, in contrast to the titular compact solenoid magnet of CMS. The looser design constraints of the ATLAS magnets allow for a more efficient, stand-alone measurement of muon momentum.

### **3.1.2 ALICE**

"ALICE" stands for "A Large Ion Collider Experiment". ALICE, unlike ATLAS and CMS, is not a general purpose detector. ALICE was designed to study a specific phenomena: the quark-gluon plasma posited to exist during ultra-relativistic heavy-ion collisions. The primary design consideration was the correct identification of particle species at high multiplicities. ALICE retains excellent PID at approximately  $dN/d\eta = 4000$ , and this over a particle momentum range up to 100 GeV. PID is accomplished through a combination of time-of-flight chambers, muon filters, and the analysis of specific ionization energy.

### **3.1.3 LHCb**

"LHCb" stands for "Large Hadron Collider beauty", reflecting the LHCb's purpose: b-quark ("beauty") studies. In this respect it is similar to ALICE, LHC's other specialized experiment. By examining CP violation in heavy-quark hadrons, LHCb hopes to elucidate the asymmetry of matter and antimatter. LHCb is asymmetric in the z-axis: b-hadron pairs tend to scatter in the same forward rapidity, so the experiment only covers  $1.9 < \eta < 4.9$ . A large dipole magnet projects an field in the vertical plane of LHC. Small asymmetries in the transverse scattering of the b-hadron system are used to test for CP violation.

### **3.1.4 TOTEM**

"TOTEM" is a sub-experiment located at CMS. TOTEM uses Roman Pots (RPS), placed far forward of CMS, to measure the total proton+proton interaction cross-section. The RPS have high acceptance for the flux very close to the beam. There are two sets of RPS, one on each side of CMS, located approximately 400 meters down the z-axis from the interaction point. The differential cross-section, of proton-proton scattering, grows exponentially with the  $|t|$  at low values of  $|t|$ . More precise measurements of this differential cross section can help distinguish between competing models of proton structure. Furthermore, the total interaction cross-section is an important

benchmark for analyzing cosmic ray showers, and can be used to independently calibrate the beam luminosity measured at other LHC experiments. Diffractive studies at TOTEM complement those at CMS.

### 3.2 Compact Muon Solenoid

The Compact Muon Solenoid (CMS) is a general-purpose particle detector located at Point-5 of the LHC. CMS was designed to precisely measure the momentum of muons. The titular superconducting solenoid magnet was designed to generate a 4 Tesla field, but operates at 3.8 T to increase longevity. This field is homogeneous and parallel to the beam line close to the interaction point. The momentum of a muon is measured from how it deflects when moving through the magnetic field. Altogether, CMS weighs approximately 12,500 metric tons, with a diameter of 14.6 m and a length of 21.6 meters. Figure 3.1 displays the detector's various sub-systems [12].




Figure 3.1: CMS Detector [12]

Within the solenoid volume are a silicon pixel and strip tracker, a lead tungsten crystal electromagnetic calorimeter (ECAL), and a brass and scintillator hadron calorimeter (HCAL), each composed of a barrel and two endcap sections. Fig.3.2 radial layering of CMS component systems [13].




Figure 3.2: CMS radial cross-section [13]

### 3.2.1 Tracker

The tracker measures the momentum of charged particles via their trajectory through a homogeneous magnetic field. The tracker consists of two units, the pixel tracker and the strip tracker, both of which are made of silicon. Compared to large-volume gas detectors, which are less expensive, silicon detectors have a faster response time. The number of tracker layers is optimized for the highest penetration by charged particles while minimizing the occurrence of multiple scattering.

Multiple scattering reduces the efficiency and resolution of the tracker because of bremsstrahlung. When a charged lepton passes through an electric field, the changes in acceleration will change the energy of lepton. The lepton will release photons that will contaminate the signals read by the tracker.

The high luminosities reached by LHC typically achieve some 20 to 30 collisions per bunch crossing, and increasing the number of hits in the tracker improves the pattern recognition of the particle flow algorithm. The particle flow algorithm is discussed in more detail later in this chapter.

A charged particle causes an electrical signal when passing through a silicon pixel or silicon microstrip. CMS reconstructs these electrical signals, taken at specific points of position and time, into tracks. These tracks are accurate to 10 micrometers. The tracker is meant to have a particle pass all the way through it, with only minimal effect particle's trajectory.

The tracker system is designed for high granularity and fast readout, such that each trajectory can be associated with its corresponding bunch crossing. The tracker is resilient enough to withstand the high flux of particles accompanying every bunch crossing; at design luminosity of  $10^34\text{cm}^{-2}\text{s}^{-1}$ , some 1000 particles will traverse the tracker every 25 ns. However, the mass of the tracker is minimal enough to suppress multiple scattering, off its material, that would distort particle trajectories. These design constraints – resistance and transparency – are satisfied silicon. The tracker has approximately  $200\text{m}^2$  of silicon surface, making it the largest silicon detector ever constructed. The pseudorapidity coverage of the pixel and strip trackers is shown in figure 3.3 [14].




Figure 3.3: Pseudorapidity acceptance of tracker [14]

In addition to excellent track reconstruction, the tracker provides high precision vertex reconstruction. Vertex reconstruction is a key ingredient in several high-profile LHC studies – Higgs, SUSY, extra-dimensions – because of the use of vertices in distinguishing non-prompt decays of heavy-quarks. The tracker is also instrumental in detecting the leptonic decays of W and Z bosons.

### 3.2.1.1 Pixel tracker

The first layer of the tracker is made of silicon pixel modules. This layer has a time resolution on the scale of 25 ns, meaning that it can take data for individual bunch crossings. Pixel technology

is used because the high flux of particles, present at high beam luminosities, require high spatial resolution for proper pattern recognition. A major design constraint is that the pixel tracker had to be able to separate distinct tracks in high multiplicity environments. Tracks reconstructed by the pixels can be used to cross-check the results from the other parts of the tracker.

Every silicon-pixel has a corresponding readout chip. The readout chips are soldered through the bump-bonding method. The readout chip amplifies signals from the pixel. The pixel tracker is precise enough to distinguish the vertices of tracks originating from short-lived particles, such as bottomonia. The innermost elements of the pixel tracker come within 4.4 cms of the CMS interaction point. The pixel tracker covers a pseudorapidity range of  $|\eta| < 2.5$  with some  $66 \times 10^6$  separate pixels.

### 3.2.1.2 Strip tracker

Outside the pixel tracker are the layers of the strip tracker. They function similar to the components of the pixel tracker, except the strip tracker consists of thin silicon plates. The strip tracker itself can be broken down into four components: the inner barrel layer, the inner endcaps, the outer barrel layer, and the outer endcaps. In total, these layers contain some  $9.3 \times 10^6$  strips.

### 3.2.2 Electromagnetic calorimeter

The Electromagnetic Calorimeter (ECAL) is the dedicated CMS calorimeter for detecting electrons and photons. The calorimeter is comprised of lead tungstate ( $PbWO_4$ ) crystals arranged in cylinder about the beam, including two endcaps, as seen in figure 3.4 [15]. The granularity of these crystals gives the ECAL excellent energy resolution, angular resolution, and spatial resolution; for example, the ECAL has the resolution suitable for the decay of the Higgs boson into two photons. The ECAL is both hermetic and homogenous.

The data readout is fast enough that CMS can trigger off signals in the ECAL. It takes about 25 ns for an ECAL hit to scintillate 80 percent of its lights, putting the calorimeter's rate on the same scale as the bunch crossing. Scintillation in the crystals activates photodetectors that transmit




Figure 3.4: ECAL components [15]

information to the L1 trigger. In the barrel these photodetectors are avalanche photodiodes (APDs). The endcaps use vacuum phototriodes (VPTs). ECAL's energy resolution, as a function of energy, is given by figure 3.5 [16]. The resolution increases with energy because, at large energies, parton showers become more similar as their random fluctuations average-out.




Figure 3.5: Ecal energy resolution [16]

In CMS trigger development, "EG" triggers fire based on energy deposits in the ECAL; fig.3.6 gives the EG trigger energy resolution [16]. This energy comes primarily from electrons and

photons. When these particles strike a tungstate crystal, the particle shower size is approximately that of the crystal. Electrons are distinguished from photons by the correlation of track to ECAL energy. The L1 trigger does not fire on tracks, which are reconstructed at the HLT level. Thus, L1 EG triggers do not distinguish between electrons and photons.




Figure 3.6: Ecal energy resolution [16]

### 3.2.3 Hadronic calorimeter

The Hadronic Calorimeter (HCAL) is the next layer outside the ECAL. The HCAL is a sampling calorimeter, meaning that it absorbs particles and measures their energy and momentum via scintillation. HCAL has such a large acceptance that it can indirectly observe non-interacting particles such as neutrinos. The HCAL is designed to be hermetic, so that imbalances of momentum and energy can be precisely measured. Figure 3.7 plots the HCAL's energy resolution; note that the resolution changes with  $|\eta|$  [16].

There are four sub-sections of the HCAL: the inner barrel (HB) and the outer barrel (HO), two




Figure 3.7: HCAL energy resolution [16]

endcaps (HE), and two forward calorimeters (HF). The HF are the most relevant to this analysis because of their use in triggering on non-hadronic events. Fig.3.8 shows the HCAL energy resolution [16].




Figure 3.8: HCAL Components [16]

### 3.2.3.1 Hadronic forward calorimeters

The Hadronic Forward Calorimeters (HF) absorbs the greatest portion of energy from collisions. As named, it is located in the forward region ( $3.0 < |\eta| < 5.2$ ) of CMS and complements the coverage provided by the barrel and endcap detectors.

HCAL is made of quartz fibers and steel absorbers for maximum radiative resistance. When high energy particles pass through the quartz fibers, these particles are moving faster than the speed of light in the medium, depositing energy that causes particle showers. These showers give off light, in the form of Cherenkov radiation, that the fibers transmit to photomultiplier tubes (PMTs). The PMT signals are converted into a digital signal by a charge integration encoder (QIE) chip. Then HCAL trigger and readout boards (HTR) use these digital signals to calculate luminosity. The HTR information is part of the data used for L1-triggering. Triggering on HF can also be done by vetoing on HF activity.

Hits in the HF are used to measure the instantaneous luminosity of CMS. As shown in fig.3.7, the HF have a finer  $E_T$  resolution than the other parts of the HCAL, making it suitable for high precision measurement of luminosity. Furthermore, because HF noise is comparatively small with respect to Minbias HF threshold, the veto of the HF threshold is a reliable measure of HF "emptiness".

### 3.2.4 Muon detector

The outermost layer of CMS consists of the muon detectors, as seen in figure 3.9 [16]. Muons are particles nearly identical to electrons, except for their mass, which exceeds that of the electron by some two orders of magnitude. High mass particles, like the Higgs boson, often decay into a final state containing muons. The muon detector not only identifies muons, but also measures their momentum. The muon detector has readout fast enough for triggering on muons.

The muon detector consists of three types of component: muon drift tubes (DT), cathode strip chambers (CSC), and resistive plate chambers (RPC). The DTs are gas filled chambers that contain a stretched metal wire. When muons pass through the DT gas, electrons are excited. These elec-




Figure 3.9: Pseudorapidity acceptance of muon detector [16]

trons escape from the gas atoms and are attracted to the metal wire, which triggers a signal. The CSC, located in the endcaps, operate under similar principles, but contain perpendicular arrays of positively charged and negatively charged wires immersed in gas. The RPC do not use electrode wires to detect excited electrons; instead, high-resitivitiy plates are used as alternating cathodes and anodes.

### 3.2.5 Zero degree calorimeter

The zero degree calorimeters are both sides of CMS, approximately 140 meters from the interaction point. Each ZDC consists of two independent systems: an electromagnetic calorimeter, for detecting very forward photons, and a hadronic calorimeter, for detecting neutrons. Because these neutrons result from the dissociation of nuclei, the ZDC can measure the centrality of heavy-ion collisions. Hadrons in the forward region have energy on the TeV scale, so the ZDC's hadronic calorimeter is made of thick tungsten plates. For a UPC process, the photon emitting nucleus is most likely to remain intact; therefore, ZDC data can the photon direction of a process, and by extension it's energy.

### 3.2.6 Particle flow algorithm

Raw data from the sub-detectors is combined, for data analysis, by the particle-flow (PF) algorithm. The PF takes the data about tracks in the tracker and energy deposits in the calorimeter, and uses them to reconstruct physics-related data objects, like jets, and to identify specific particles, such as photons and muons. The PF also identifies missing energy and momentum for use in neutrino studies. These data objects are stored in a format similar to that of conventional MC event generators. CMS gains significant jet reconstruction efficiency via the PF. At low transverse momentum, PF reconstructs jets at nearly twice the resolution of HCAL and ECAL. This increase in efficiency comes from the PF integrating in track data with the calorimeter tower data. Figure 3.10 compares the performance of the PF algorithm to the calorimeter reconstruction [17].




Figure 3.10: Performance of particle flow alorgithm compared to calorimeter readout [17]

# Chapter 4

## Measuring luminosity

### 4.1 Instantaneous luminosity

One of the most important quantities measured by CMS is luminosity. Luminosity is necessary to convert the number of events detected, for a given channel, into a collision cross-section. Collision cross-sections are among the primary observables predicted by theoretical physics, specifically quantum field theory. For particle physics, the collision cross-section of a process is typically measured through the relation:

$$\sigma = \frac{R}{L}, \quad (4.1)$$

where  $\sigma$  is the cross-section,  $R$  is the rate at which the process occurs per collision, and  $L$  is the luminosity.

$$L = \frac{R}{\sigma_o(E)A(t, \mu, n_b, \dots)}, \quad (4.2)$$

### 4.2 Luminometers

There are two kinds of CMS luminometer: online and offline. Online luminometers readout the luminosity per bunch in real time. As of 2015 there are three online luminometers: the pixel luminosity telescope (PLT), the HF, and the beam conditions monitor (BCM1f). There is a high-rate, independent data-acquisition system for each of the online luminometers. Offline luminometers measure the rate of reconstructed objects. The primary offline lumimometer is the pixel tracker. In general, the offline lumimometers have better stability over time. [27]

The online and offline luminometers complement each other for high precision data analysis.

Specifically, the offline data can be used to calibrate out imperfections in the online data.

In addition to these hardware luminometers, CMS can use physics processes as luminosity benchmarks. For example, other experiments have measured the Z-boson cross-section to a high accuracy and high precision. Comparing this cross-section to the Z-boson mass-peak ,in CMS data, provides a cross-check to the delivered luminosity.

### 4.2.1 Pixel cluster counting

One of the primary methods of measuring instantaneous luminosity is through pixel cluster counting (PCC) in the pixel-tracker. The collision rate is linked to instantaneous luminosity  $L$  via the total inelastic cross-section  $\sigma_T$

$$v\mu = L\sigma_T, \quad (4.3)$$

where  $v = 11246$  Hz is the beam revolution frequency and  $\mu$  is the number of collisions per bunch crossing. We define the average number of pixel clusters per event,  $\langle n \rangle$ , as

$$\langle n \rangle = \mu n_1, \quad (4.4)$$

and the visible cross section  $\sigma_{vis}$  as

$$\sigma_{vis} = \sigma_T n_1, \quad (4.5)$$

where  $n_1$  is the average number of pixel clusters per inelastic collision. Leveraging the previous two equations, the instantaneous luminosity reduces to

$$L = \frac{v \langle n \rangle}{\sigma_{vis}}, \quad (4.6)$$

such that  $L$  can be calculated after measuring the actual  $\langle n \rangle$  of the data [28]

### 4.2.2 Pixel luminosity telescope

The pixel luminosity telescope (PLT) is one of the online luminometers of CMS. The PLT consists of 8 pixel telescopes arranged around the beamline on either side of the CMS interaction point.

Each telescope has three layers of silicon pixel detector. Triple coincidences in these telescopes are measured as luminosity on a bunch-by-bunch basis. As an independent luminometer, the PLT data is also useful for reducing the systematic uncertainty of the luminosity delivered to CMS.

The PLT was installed as part of the CMS Run-2 upgrades. As of this writing, it is the only subdetector of CMS dedicated solely to measuring instantaneous luminosity. The hardware of the pixel sensors and the readout chip are similar to those used by the pixel tracker. However, the readout chips (ROC) of the PLT are capable of delivering fast-OR formatted data; the pixel tracker cannot do this. The fast-OR readout is optimized for delivering triple-coincidence data; whereas the standard pixel data readout has a max frequency of 100 kHz, the PLT fast-OR readout maxes out at an orders of magnitude greater frequency, at 40 MHz [29].

### 4.2.3 Hadronic forward calorimeter

The hadronic forward calorimeter (HF) of CMS is capable of measuring luminosity. Because of the high flux of particles through the forward region, the HF-luminosity is less subject to statistical uncertainty than the PCC-luminosity. The HF can measure luminosity per bunch through its own dedicated high-rate acquisition system that is separate from the CMS DAQ. Within the HF, the HTR boards each have an additional HF luminosity transmission circuit (HLX). The HLX are mounted in one of the mezzanine expansion slots of the HTR.

The HF luminosity is subject to two difficulties. Because the PMTs require a calibrated voltage, the accumulation of gain changes over long periods of time will cause calibration drifts. Furthermore, the HF detector response has been shown to be nonlinear with respect to vertex pile-up. Substantial data analysis goes into calculating the corrections to these two sources of error [28].

### 4.2.4 Fast Beam Conditions Monitor

The CMS Beam Conditions and Radiation Monitoring System (BRM) provides a wide variety of information on the accumulated radiation dosage into CMS. This information is essential for protecting CMS from radiation damage. There are many subcomponents to the BRM system, one

of which is the Fast Beam Conditions Monitor (BCM1f). The BCM1f is capable of measuring the beam-halo background and the flux from collision products. Similar to the PLT, BCM1f is located close to the beam pipe; however, BCM1f uses diamonds instead of silicon pixels. The readout from BCM1f is not digital, but an analog signal sent through optical fibres.

There are 24 single crystal diamond sensors in the BCM1f system. The BCM1f is located very close to interaction point, specifically 1.8 m on either side. This location is chosen specifically to take advantage of the 25 ns between bunch crossings. From the BCM1f vantage point there is a 12.5 ns delay between particle fluxes from the machine induced background and those coming from actual collisions. Thus the BCM1f can use timing to distinguish between signal and background [30].

### 4.3 Van de Meer scanning calibration

The luminometers of CMS produce signals proportional to the instantaneous luminosity of the LHC beam. However, these signals need to be properly calibrated with respect to a known visible cross-section for each luminometer. This calibration is accomplished via Van de Meer scanning. The opposing beams of LHC are moved back and forth in the transverse plane. During the scan, the detector response is measured as a function of beam displacement. The beam widths are calculated from Gaussian fits to the detector response. The visible cross-section of the luminometer in question is then derived from the width of the beams, and acts as the calibration of the detector response.

$$\sigma_{vis} = \frac{2\pi\Sigma_x\Sigma_y v \langle n \rangle_{\Delta=0}}{N_1 N_2}, \quad (4.7)$$

### 4.4 Corrections to VDM scans

The results of the VDM scan must be corrected for the following phenomena: the length scale, the ghosts and satellite background, the growth in beam emittance, orbit drift, and beam-beam elec-




Figure 4.1: PCC VdM Scans.

tromagnetic effects. These affect the normalization of the VDM scans, and so the corrections take the form of calibration constants derived from additional scans and beam monitoring subdetectors.

The length scale correction is derived by comparing the effects of a length scan to the response of the tracker; checking the difference between the reported beam displacement and that seen in the tracker.

In addition to the circulating bunches, the LHC beams can be polluted by foreign particles that were not intended for injection by LHC commissioning. The presence of these foreign particles – fleeting "ghosts" and persistent "satellites" – can be detected by comparing the total beam current, measured by the DC beam current transformer, against the per bunch current measured by the FBCT. The FBCT, because it is timed for bunch crossings, is less likely to interact with ghosts and satellites.

	Systematic	correction (%)	uncertainty (%)
Integration	Stability	-	1
	type 1	7 – 9	0.6
	type 2	0 – 4	0.7
	CMS deadtime	-	0.5
	Dynamic Inefficiency	-	0.4
Normalization	XY-Correlations	1.1	1.5
	Beam current calibration	-	0.3
	Ghosts and satellites	-	0.2
	Length scale	-3.2	1.5
	Orbit Drift	-	0.4
	Beam-beam deflection	1.8	0.4
	Dynamic- $\beta$	-	0.5
	Total		2.7

Figure 4.2: Systematic Uncertainty During 2015 pp Run.

## 4.5 Author’s contributions

LHC produces a staggering amount of data. The 2015 heavy-ion run itself recorded some  $1.4 pb^{-1}$  in integrated luminosity. This integrated luminosity is reconstructed into data on the order of pentabytes.

I contributed to CMS luminosity validation, and performed studies on the long-term stability of the BRIL luminometers. On a week by week basis, I would examine the data from online and offline luminometers and certify that it met the quality standards of CMS. If the data exhibited non-linear behavior, I would de-certify the corresponding lumi-sections in the BRIL json files.

All CMS data has to go through a data certification workflow. Data certification insures that only valid data is used in CMS analyses. Decisions on data certification are recorded in two systems: the DQM Run Registry, shown here in figure 4.4, and the normtag JSON files of the BRIL-DPG. There are occasions when the sub-detectors of CMS might record poor quality data. For example, the software reading out data from the PLT could crash and thus compromise the output. In this case the certification workers would record the lumi sections during which the PLT software crashed. These lumi sections would be removed from the normtag JSON file passed on to the BRIL group. If all te luminometers are malfunctioning, a whole run could be invalidated in the run registry.

In additon to data certification, I reviewed the long-term stability of the BRIL luminometers

Fill	Begin Time YYYY.MM.DD HH:MM	Duration HH:MM	PeakInstLumi $\times 10^{39} \text{ cm}^{-2} \text{s}^{-1}$ pp $\times 10^{24} \text{ cm}^{-2} \text{s}^{-1}$ ions	DeliveredLumi $\text{pb}^{-1}$ pp $\mu\text{b}^{-1}$ PbPb	RecordedLumi $\text{pb}^{-1}$ pp $\mu\text{b}^{-1}$ PbPb	EffByLumi %
4658	2015.11.25 09:59	05:40	1	0.0	0.0	84.4
4659	2015.11.25 22:51	02:12	89	0.6	0.5	95.6
4661	2015.11.26 07:04	02:55	97	0.8	0.5	69.6
4664	2015.11.26 16:33	04:27	547	5.3	4.5	85.7
4666	2015.11.27 08:34	05:05	721	7.0	6.2	88.8
4669	2015.11.27 21:31	06:10	1327	16.1	12.8	79.2
4671	2015.11.28 11:13	09:03	1115	16.5	14.3	86.9
4672	2015.11.28 23:53	06:55	1085	14.0	11.9	84.6
4677	2015.11.29 17:42	06:54	1011	14.0	12.5	89.6
4679	2015.11.30 08:27	06:58	1092	14.5	12.9	88.8
4680	2015.11.30 21:17	05:13	1820	19.6	16.7	85.4
4681	2015.12.01 07:40	11:51	1856	28.4	25.1	88.2
4685	2015.12.02 05:39	03:28	1040	8.5	6.0	70.4
4688	2015.12.03 02:44	00:18	1887	1.9	1.0	51.1
4689	2015.12.03 09:03	09:04	1254	10.2	9.9	97.1
4690	2015.12.03 22:10	08:27	2030	22.0	21.7	98.5
4691	2015.12.04 12:51	07:31	2072	26.7	25.8	96.6
4692	2015.12.04 23:23	07:40	2334	30.3	29.7	98.1
4693	2015.12.05 10:22	00:60	2314	7.1	6.9	97.2
4695	2015.12.05 17:29	07:14	2216	27.8	26.2	94.3
4696	2015.12.06 07:03	05:29	1795	19.9	19.4	97.7
4697	2015.12.06 16:34	05:30	1879	20.5	20.1	98.2
4698	2015.12.07 01:41	00:09	2102	1.0	0.9	87.2
4699	2015.12.07 05:30	05:40	1885	21.5	20.4	94.8
4705	2015.12.08 05:37	02:24	183	1.1	1.1	97.8
4706	2015.12.08 11:08	04:32	2433	23.5	22.7	96.5
4709	2015.12.09 08:41	05:47	2161	23.3	22.8	97.8
4710	2015.12.09 17:42	05:52	2967	32.8	32.2	98.3
4711	2015.12.10 02:38	06:37	2886	31.6	31.1	98.4
4714	2015.12.11 01:20	06:39	2500	29.3	28.2	96.2
4715	2015.12.11 11:37	06:31	2559	30.6	29.3	95.7
4717	2015.12.11 23:00	10:21	0	0.0	0.0	0.0
4719	2015.12.12 17:26	04:36	2652	27.2	25.6	94.1
4720	2015.12.13 01:10	10:60	3036	44.9	44.2	98.5
Summary		199:09	3036	578.4	543.1	93.9

Figure 4.3: Luminosity by CMS fill.

during the 2015 and 2016 proton-proton runs. Stability refers to the comparative performance of the different luminometers. Ideally, the ratio of the instantaneous luminosity reported by separate luminometers should be a constant ratio, but in fact there is a tendency for this ratio to drift as a result of radiation damage to the luminometers. This drift can be seen in plotting the average ratio as a function of interaction rate and of the different measures of LHC time: runs, lumi-sections, and bunch-crossings. Instantaneous luminosity, in theory, should be independent of machine conditions like acceptance and efficiency.

**CMS DQM Run Registry (GLOBAL)**

Chris Palmer (null) @null    Workspace    Tools    Support    Logout

Offline Application (3.6.1)

Offline Datasets Show Waiting list Refresh | Configure | Export | Page 1 / 1,584 | Help

Common Offline Shift taker actions:

- The runs in this table have been signed-off by online certification and are waiting for offline certification. The component status fields start out with the results of the online certification by default.
- Determine whether run is ready for offline certification, for example checking whether plots are available in DQM GUI (consult shift instructions). This may also be automated.
- Click a row to select it and enable more buttons:
  - Click *Details* to view meta information.
  - Click *Create Dataset* to bring

Ru...	Run Class ...	Dataset Name	Dataset...	Dataset Created	Last Shifter	Cms	Castor	Csc	Dt	Ecal	Es
305842	Collisions17	/PromptReco/Collisions2017F/DQM	OPEN	Sun 05-11-17 02:52:33	DQMGUI Tr...	GOOD	EXCLUDED	GOOD	GOOD	GOOD	GOOD
305841	Collisions17	/PromptReco/Collisions2017F/DQM	OPEN	Sat 04-11-17 15:26:04	DQMGUI Tr...	GOOD	EXCLUDED	GOOD	GOOD	GOOD	GOOD
305839	Collisions17	/PromptReco/Collisions2017F/DQM	OPEN	Sun 05-11-17 05:13:10	DQMGUI Tr...	GOOD	EXCLUDED	GOOD	GOOD	GOOD	GOOD
305833	Collisions17	/PromptReco/Collisions2017F/DQM	OPEN	Sat 04-11-17 12:21:31	DQMGUI Tr...	GOOD	EXCLUDED	GOOD	GOOD	GOOD	GOOD
305832	Collisions17	/PromptReco/Collisions2017F/DQM	OPEN	Mon 06-11-17 11:07:35	DQMGUI Tr...	GOOD	EXCLUDED	GOOD	GOOD	GOOD	GOOD
305821	Collisions17	/PromptReco/Collisions2017F/DQM	OPEN	Sun 05-11-17 23:46:43	DQMGUI Tr...	GOOD	EXCLUDED	GOOD	GOOD	GOOD	GOOD
305814	Collisions17	/PromptReco/Collisions2017F/DQM	OPEN	Sun 05-11-17 17:47:24	DQMGUI Tr...	GOOD	EXCLUDED	GOOD	GOOD	GOOD	GOOD
305809	Collisions17	/PromptReco/Collisions2017F/DQM	OPEN	Sat 04-11-17 03:52:37	DQMGUI Tr...	GOOD	EXCLUDED	GOOD	GOOD	GOOD	GOOD
305766	Collisions17	/PromptReco/Collisions2017F/DQM	OPEN	Fri 03-11-17 22:59:06	DQMGUI Tr...	GOOD	EXCLUDED	GOOD	GOOD	GOOD	GOOD
305758	Collisions1...	/PromptReco/SpecialRun2017/DQM	OPEN	Thu 02-11-17 02:07:15	DQMGUI Tr...	GOOD	EXCLUDED	GOOD	GOOD	GOOD	GOOD
305757	Collisions1...	/PromptReco/SpecialRun2017/DQM	OPEN	Thu 02-11-17 09:02:13	DQMGUI Tr...	GOOD	EXCLUDED	GOOD	GOOD	GOOD	GOOD
305756	Collisions1...	/PromptReco/SpecialRun2017/DQM	OPEN	Wed 01-11-17 23:56...	DQMGUI Tr...	GOOD	EXCLUDED	GOOD	GOOD	GOOD	GOOD
305755	Collisions1...	/PromptReco/SpecialRun2017/DQM	OPEN	Thu 02-11-17 01:32:47	DQMGUI Tr...	GOOD	EXCLUDED	GOOD	GOOD	GOOD	GOOD

Figure 4.4: CMS Run Registry.

### CMS Integrated Luminosity, PbPb, 2015, $\sqrt{s} = 5.02$ TeV/nucleon




Figure 4.5: CMS lumi during HI-2015.




Figure 4.6: CMS Lumi by Era.

# Chapter 5

## Trigger development and performance

### 5.1 Introduction to triggering

Modern "triggering" methods began with Walther Bothe's development of the coincidence circuit. The coincidence circuit accepts two inputs. If these inputs are received within the same time window, approximating coincidence, the circuit passes an output. Bothe's originally used the coincidence circuit to take data for electron-photon production in Compton scattering. Figure 5.1 is the diagram of a coincidence circuit [18].




Figure 5.1: Rossi circuit with three Geiger-Muller coincidence circuits [18]

### 5.2 Triggering at CMS

At stable beams, the LHC delivers bunch crossings every 25 nanoseconds. Each bunch crossing in turn will have some 20 hadron collisions. The resulting interaction rate –  $10^9$  interactions per second – is orders of magnitude greater than the frequency that events can be written to disk,

$10^2$  events per second. CMS therefore needs a means of filtering out the most interesting  $10^2$  interactions per second while declining the other  $10^6$  interactions per second.

CMS uses a two-tiered triggering system. The first tier, the L1 trigger, is hardware based. The second tier, the high-level trigger (HLT), is software based. The L1 trigger receives raw data from the calorimeters and the muon detectors; this determines when the tracker will readout data. The raw data from the tracker, calorimeters, and muon detectors is then passed on to a computer farm running the HLT menu. The HLT then performs a simplistic reconstruction of the raw data into physics objects useful for analysis: jets, tracks, and identifiable particles. If an event passes the HLT, the raw data is permanently stored in preparation for a more complex reconstruction.

The initial interaction rate is approximately  $3.2 \mu\text{s}$ . The L1 trigger can only pass some 1 in 1000 interactions to the HLT. The L1 trigger menu has an output rate of approximately 100 kHz. L1 achieves this rate by only considering data of reduced granularity and reduced resolution. A buffer is used to store the full event data while the L1 runs. The HLT menu reduces this to about 100 Hz as required by the limit on disk writing.

The 2015 UPC triggers were for low multiplicity events and low transverse momentum events. Typical heavy-ion collisions are high multiplicity events. Hard scattering dominates. Fig.5.2 is an event display of one of the first heavy-ion collisions at CMS in 2010.




Figure 5.2: High multiplicity PbPb collision

Fig.5.3 is the event display of a UPC upsilon candidate. Notice that there are only two reconstructed tracks, and that CMS is otherwise empty in all calorimeters. These events constitute a complement to those passing conventional more heavy-ion triggers. However, for Run-2 the Pb-Pb luminosities are high enough that UPC final states can be studied over a wide range of rapidity and  $p_T$ .




Figure 5.3: UPC Upsilon candidate

For this analysis, the L1 trigger applies two selections. First, the L1 checks that at least one of the HF is empty. This is the most important part of the trigger in so far as it suppressed the hadronic contamination of the dataset. Then, if there is at least 5 GeV of energy deposited in the ECAL, the event passes to the HLT.

Low multiplicity events are difficult to distinguish from background. To compensate, the HLT in turn requires that there be at least once reconstructed track from the pixel tracker, to make sure that there are particles that will be reconstructed by the complete tracker. Only the pixel tracker is used for these HLTs to increase the speed of reconstruction while decreasing needed computer cycles.

### 5.3 Author's contributions

In preparation for the 2015 heavy-ion run and the 2016 p-Pb run, I prepared high-level trigger menus for the CMS Forward-HI group. This trigger menu was optimized for firing on ultra-peripheral collisions. I tested the menu's performance on Monte Carlo generated by STARLIGHT

and reconstructed through a GEANT4 simulation of CMS. During the experiment, I was present at CERN to monitor the trigger rates and deliver daily reports on their performance.

CMSSW includes an emulator for the L1 trigger. This software can re-emulate alternative L1 menus on previously taken CMS data. The performance of a trigger on the 2011 PbPb data, taken at  $\sqrt{s_{NN}} = 2.7\text{TeV}$  is extrapolated to the higher energy,  $\sqrt{s_{NN}} = 5.2\text{TeV}$ , of the 2015 PbPb run.

I tested my HLT menu on both STARLIGHT MC and on data from the 2011 Pb+Pb run. STARLIGHT is a MC generator for ultra-peripheral collisions, in particular for the vector-meson photoproduction channels. I used STARLIGHT to generate MC sets for  $Pb + Pb \rightarrow J/\Psi + Pb + Pb$  and  $Pb + Pb \rightarrow \Upsilon(1s) + Pb + Pb$ . I then used CMSSW to test the performance of the component bits of the HLT paths with respect to these MC sets.




Figure 5.4: Example of trigger rate.

It was my duty to carefully observe the state of the UPC triggers during the heavy-ion run. If the total HLT rate ever drifted above 100 Hz, the HLT menu could crash and CMS lose considerable data. It was important for the trigger contacts to make sure that there HLT paths were behaving stably. I also analyzed express physics data to test that the vector-meson triggers saw appropriate mass resonances.

## 5.4 Studies from trigger menus

The UPC HLTs I designed are useful for a variety of physics studies. In particular, vector meson photoproduction, light-by-light scattering, and UPC particle correlations.

The single and double muon UPC triggers are adequate for studies of vector-meson photoproduction. As of this writing, the Forward-HIN group has ongoing studies of UPC  $J/\Psi$  and UPC  $\Upsilon(1s)$  in both the 2015 PbPb and 2016 pPb data.

My trigger menu has produced a data set well suited for studying the heavy-ion initial state.

# **Chapter 6**

## **Data analysis**

In this chapter our data sample will be analyzed according to the framework provided by Yoshikata Hatta. The angular correlation of jets is taken, and event mixing is used to factor out acceptance effects.




Figure 6.1: Raw  $\phi$  distribution.




Figure 6.2: Mixed  $\phi$  distribution.




Figure 6.3: Subtracted  $\phi$  distribution.




Figure 6.4: Final  $\phi$  distribution.

## **Chapter 7**

### **Systematic uncertainties**

The primary sources of systematic uncertainty in this analysis are the measurements of track vertex, jet momentum, jet pseudorapidity, and HF energy.

# **Chapter 8**

## **Results and conclusions**

In this chapter, the angular anisotropy of the elliptic gluons is compared to theoretical predictions from variations of NLO-QCD.

## References

- [1] S. Bethke, “Experimental tests of asymptotic freedom,” *Prog. Part. Nucl. Phys.* **58** (2007) 351–386, arXiv:hep-ex/0606035 [hep-ex].
- [2] **ZEUS, H1** Collaboration, S. Shimizu, “Low  $Q^{**2}$  Structure Functions including the Longitudinal Structure Function,” in *QCD and high energy interactions. Proceedings, 44th Rencontres de Moriond, La Thuile, Italy, March 14-21, 2009*, pp. 301–304. 2009.  
arXiv:0905.4712 [hep-ex].  
<http://inspirehep.net/record/821571/files/arXiv:0905.4712.pdf>.
- [3] R. S. Bhalerao, “Relativistic heavy-ion collisions.”.  
<https://cds.cern.ch/record/1695331>. Comments: Updated version of the lectures given at the First Asia-Europe-Pacific School of High-Energy Physics, Fukuoka, Japan, 14-27 October 2012. Published as a CERN Yellow Report (CERN-2014-001) and KEK report (KEK-Proceedings-2013-8), K. Kawagoe and M. Mulders (eds.), 2014, p. 219. Total 21 pages.
- [4] M. Diehl, “Generalized parton distributions,” *Phys. Rept.* **388** (2003) 41–277,  
arXiv:hep-ph/0307382 [hep-ph].
- [5] Y. Hatta, B.-W. Xiao, and F. Yuan, “Probing the Small-  $x$  Gluon Tomography in Correlated Hard Diffractive Dijet Production in Deep Inelastic Scattering,” *Phys. Rev. Lett.* **116** no. 20, (2016) 202301, arXiv:1601.01585 [hep-ph].
- [6] J. D. Jackson, *Classical electrodynamics*. Wiley, New York, NY, 3rd ed. ed., 1999.  
<http://cdsweb.cern.ch/record/490457>.

- [7] S. Klein and J. Nystrand, “Ultraperipheral nuclear collisions,” *Phys. Today* **70** no. 10, (2017) 40–47.
- [8] CMS Collaboration, V. Khachatryan *et al.*, “Coherent  $J/\psi$  photoproduction in ultra-peripheral PbPb collisions at  $\sqrt{s_{NN}} = 2.76$  TeV with the CMS experiment,” *Phys. Lett. B* **772** (2017) 489–511, arXiv:1605.06966 [nucl-ex].
- [9] ATLAS Collaboration, M. Aaboud *et al.*, “Evidence for light-by-light scattering in heavy-ion collisions with the ATLAS detector at the LHC,” *Nature Phys.* **13** no. 9, (2017) 852–858, arXiv:1702.01625 [hep-ex].
- [10] A. Collaboration, “ATLAS Event Display: Light-by-Light Scattering.” General Photo, Aug, 2017.
- [11] H1 Collaboration, V. Andreev *et al.*, “Diffractive Dijet Production with a Leading Proton in  $ep$  Collisions at HERA,” *JHEP* **05** (2015) 056, arXiv:1502.01683 [hep-ex].
- [12] CMS Collaboration, D. G. d’Enterria *et al.*, “CMS physics technical design report: Addendum on high density QCD with heavy ions,” *J. Phys. G* **34** (2007) 2307–2455.
- [13] L. R. F. Castillo, “The atlas and cms detectors,” in *The Search and Discovery of the Higgs Boson*, 2053-2571, pp. 4–1 to 4–8. Morgan and Claypool Publishers, 2015.  
<http://dx.doi.org/10.1088/978-1-6817-4078-2ch4>.
- [14] CMS Collaboration, S. Chatrchyan *et al.*, “Alignment of the CMS Silicon Tracker during Commissioning with Cosmic Rays,” *JINST* **5** (2010) T03009, arXiv:0910.2505 [physics.ins-det].
- [15] CMS Collaboration, A. Benaglia, “The CMS ECAL performance with examples,” *JINST* **9** (2014) C02008.
- [16] CMS Collaboration, G. L. Bayatian *et al.*, “CMS Physics.”

- [17] CMS Collaboration, A. M. Sirunyan *et al.*, “Particle-flow reconstruction and global event description with the CMS detector,” *JINST* **12** no. 10, (2017) P10003, arXiv:1706.04965 [physics.ins-det].
- [18] L. Bonolis, “Walther Bothe and Bruno Rossi: the birth and development of coincidence methods in cosmic-ray physics,” *Am. J. Phys.* **79** (2011) 1133, arXiv:1106.1365 [physics.hist-ph].
- [19] F. Halzen and A. D. Martin, *QUARKS AND LEPTONS: AN INTRODUCTORY COURSE IN MODERN PARTICLE PHYSICS*. 1984.
- [20] M. E. Peskin and D. V. Schroeder, *An Introduction to quantum field theory*. Addison-Wesley, Reading, USA, 1995.  
<http://www.slac.stanford.edu/~mpeskin/QFT.html>.
- [21] D. Griffiths, *Introduction to elementary particles*. 2008.
- [22] CTEQ Collaboration, R. Brock *et al.*, “Handbook of perturbative QCD: Version 1.0,” *Rev. Mod. Phys.* **67** (1995) 157–248.
- [23] F. Wilczek, “QCD made simple,” *Phys. Today* **53N8** (2000) 22–28.
- [24] J. D. Bjorken, “Asymptotic Sum Rules at Infinite Momentum,” *Phys. Rev.* **179** (1969) 1547–1553.
- [25] D. Boer *et al.*, “Gluons and the quark sea at high energies: Distributions, polarization, tomography,” arXiv:1108.1713 [nucl-th].
- [26] A. V. Belitsky, X.-d. Ji, and F. Yuan, “Quark imaging in the proton via quantum phase space distributions,” *Phys. Rev.* **D69** (2004) 074014, arXiv:hep-ph/0307383 [hep-ph].
- [27] CMS Collaboration, CMS, “Measurement of CMS Luminosity.”

- [28] CMS Collaboration, C. Collaboration, “CMS Luminosity Based on Pixel Cluster Counting - Summer 2013 Update,”.
- [29] CMS Collaboration, A. Kornmayer, “The CMS pixel luminosity telescope,” *Nucl. Instrum. Meth.* **A824** (2016) 304–306.
- [30] CMS Collaboration, M. Guthoff, “Instrumentation for beam radiation and luminosity measurement in the CMS experiment using novel detector technologies,” *Nucl. Instrum. Meth.* **A845** (2017) 565–569.
- [31] S. D. Ellis and D. E. Soper, “Successive combination jet algorithm for hadron collisions,” *Phys. Rev.* **D48** (1993) 3160–3166, arXiv:hep-ph/9305266 [hep-ph].
- [32] M. Thomson, *Modern particle physics*. Cambridge University Press, New York, 2013.  
<http://www-spires.fnal.gov/spires/find/books/www?cl=QC793.2.T46::2013>.
- [33] H1 Collaboration, F. D. Aaron *et al.*, “Diffractive Dijet Photoproduction in ep Collisions at HERA,” *Eur. Phys. J.* **C70** (2010) 15–37, arXiv:1006.0946 [hep-ex].
- [34] F. D. Aaron *et al.*, “Measurement of the cross section for diffractive deep-inelastic scattering with a leading proton at HERA,” *Eur. Phys. J.* **C71** (2011) 1578, arXiv:1010.1476 [hep-ex].
- [35] ZEUS Collaboration, S. Chekanov *et al.*, “Deep inelastic scattering with leading protons or large rapidity gaps at HERA,” *Nucl. Phys.* **B816** (2009) 1–61, arXiv:0812.2003 [hep-ex].
- [36] J. A. Crittenden, “Exclusive production of neutral vector mesons at the electron - proton collider HERA,” arXiv:hep-ex/9704009 [hep-ex].
- [37] V. Guzey and M. Klasen, “A fresh look at factorization breaking in diffractive photoproduction of dijets at HERA at next-to-leading order QCD,” *Eur. Phys. J.* **C76** no. 8, (2016) 467, arXiv:1606.01350 [hep-ph].

- [38] V. Guzey and M. Klasen, “Diffractive dijet photoproduction in ultraperipheral collisions at the LHC in next-to-leading order QCD,” *JHEP* **04** (2016) 158, arXiv:1603.06055 [hep-ph].
- [39] J. Gaiser, “Charmonium Spectroscopy From Radiative Decays of the  $J/\psi$  and  $\psi'$ ”,.
- [40] S. van der Meer, “Calibration of the Effective Beam Height in the ISR,”.
- [41] T. Bauer, R. Spital, D. Yennie, and F. Pipkin, “The Hadronic Properties of the Photon in High-Energy Interactions,” *Rev.Mod.Phys.* **50** (1978) 261.
- [42] S. Donnachie, H. G. Dosch, O. Nachtmann, and P. Landshoff, “Pomeron physics and QCD,” *Camb.Monogr.Part.Phys.Nucl.Phys.Cosmol.* **19** (2002) 1–347.
- [43] **ALICE Collaboration** Collaboration, B. B. Abelev *et al.*, “Exclusive  $J/\psi$  photoproduction off protons in ultra-peripheral p-Pb collisions at  $\sqrt{s_{NN}}=5.02$  TeV,” arXiv:1406.7819 [nucl-ex].
- [44] **ZEUS Collaboration** Collaboration, S. Chekanov *et al.*, “Exclusive photoproduction of  $J/\psi$  mesons at HERA,” *Eur.Phys.J.* **C24** (2002) 345–360, arXiv:hep-ex/0201043 [hep-ex].
- [45] **H1 Collaboration** Collaboration, A. Aktas *et al.*, “Elastic  $J/\psi$  production at HERA,” *Eur.Phys.J.* **C46** (2006) 585–603, arXiv:hep-ex/0510016 [hep-ex].
- [46] **H1 Collaboration** Collaboration, C. Alexa *et al.*, “Elastic and Proton-Dissociative Photoproduction of  $J/\psi$  Mesons at HERA,” *Eur.Phys.J.* **C73** (2013) 2466, arXiv:1304.5162 [hep-ex].
- [47] L. Frankfurt, M. Strikman, and M. Zhalov, “Elastic and large  $t$  rapidity gap vector meson production in ultraperipheral proton-ion collisions,” *Phys.Lett.* **B640** (2006) 162–169, arXiv:hep-ph/0605160 [hep-ph].

- [48] V. Guzey and M. Zhalov, “Rapidity and momentum transfer distributions of coherent  $J/\psi$  photoproduction in ultraperipheral pPb collisions at the LHC,” *JHEP* **1402** (2014) 046, arXiv:1307.6689 [hep-ph].
- [49] **ALICE Collaboration** Collaboration, B. Abelev *et al.*, “Coherent  $J/\psi$  photoproduction in ultra-peripheral Pb-Pb collisions at  $\sqrt{s_{NN}} = 2.76$  TeV,” *Phys.Lett.* **B718** (2013) 1273–1283, arXiv:1209.3715 [nucl-ex].
- [50] K. Eskola, H. Paukkunen, and C. Salgado, “EPS09: A New Generation of NLO and LO Nuclear Parton Distribution Functions,” *JHEP* **0904** (2009) 065, arXiv:0902.4154 [hep-ph].
- [51] K. J. Eskola, H. Paukkunen, and C. A. Salgado, “An Improved global analysis of nuclear parton distribution functions including RHIC data,” *JHEP* **0807** (2008) 102, arXiv:0802.0139 [hep-ph].
- [52] J. Pumplin, D. Stump, J. Huston, H. Lai, P. M. Nadolsky, *et al.*, “New generation of parton distributions with uncertainties from global QCD analysis,” *JHEP* **0207** (2002) 012, arXiv:hep-ph/0201195 [hep-ph].
- [53] A. Martin, W. Stirling, R. Thorne, and G. Watt, “Parton distributions for the LHC,” *Eur.Phys.J.* **C63** (2009) 189–285, arXiv:0901.0002 [hep-ph].
- [54] M. Goldhaber and E. Teller, “On nuclear dipole vibrations,” *Phys.Rev.* **74** (1948) 1046–1049.
- [55] H. Steinwedel and J. H. D. Jensen, “,” *Zeits. f. Naturforsch* **5a** (1950) 343.
- [56] **ALICE Collaboration** Collaboration, E. Abbas *et al.*, “Charmonium and  $e^+e^-$  pair photoproduction at mid-rapidity in ultra-peripheral Pb-Pb collisions at  $\sqrt{s_{NN}} = 2.76$  TeV,” *Eur.Phys.J.* **C73** (2013) 2617, arXiv:1305.1467 [nucl-ex].

- [57] H. Emling, “Electromagnetic excitation of the two phonon giant dipole resonance,” *Prog.Part.Nucl.Phys.* **33** (1994) 729–786.
- [58] E. de Passos, M. Hussein, L. Canto, and B. Carlson, “The Mean energy, strength and width of triple giant dipole resonances,” *Phys.Rev.* **C65** (2002) 034326, arXiv:nucl-th/0108002 [nucl-th].
- [59] M. Ryskin, “Diffractive J / psi electroproduction in LLA QCD,” *Z.Phys.* **C57** (1993) 89–92.
- [60] V. Guzey, M. Strikman, and M. Zhalov, “Disentangling coherent and incoherent  $J/\psi$  photoproduction on nuclei by neutron tagging in ultraperipheral ion collisions at the LHC,” arXiv:1312.6486 [hep-ph].
- [61] L. Frankfurt, V. Guzey, and M. Strikman, “Leading Twist Nuclear Shadowing Phenomena in Hard Processes with Nuclei,” *Phys.Rept.* **512** (2012) 255–393, arXiv:1106.2091 [hep-ph].
- [62] V. Goncalves and M. Machado, “Vector Meson Production in Coherent Hadronic Interactions: An update on predictions for RHIC and LHC,” *Phys.Rev.* **C84** (2011) 011902, arXiv:1106.3036 [hep-ph].
- [63] A. Cisek, W. Schafer, and A. Szczurek, “Exclusive coherent production of heavy vector mesons in nucleus-nucleus collisions at LHC,” *Phys.Rev.* **C86** (2012) 014905, arXiv:1204.5381 [hep-ph].
- [64] T. Lappi and H. Mantysaari, “ $J/|\psi\rangle$  production in ultraperipheral Pb+Pb and  $p$ +Pb collisions at energies available at the CERN Large Hadron Collider,” *Phys.Rev.* **C87** no. 3, (2013) 032201, arXiv:1301.4095 [hep-ph].
- [65] J. Hufner, B. Kopeliovich, and A. B. Zamolodchikov, “Inelastic J / psi photoproduction off nuclei: Gluon enhancement or double color exchange?,” *Z.Phys.* **A357** (1997) 113–120, arXiv:nucl-th/9607033 [nucl-th].

- [66] S. J. Brodsky, L. Frankfurt, J. Gunion, A. H. Mueller, and M. Strikman, “Diffractive lepto production of vector mesons in QCD,” *Phys.Rev.* **D50** (1994) 3134–3144, arXiv:hep-ph/9402283 [hep-ph].
- [67] M. Strikman, M. Tverskoy, and M. Zhalov, “Neutron tagging of quasielastic J/psi photoproduction off nucleus in ultraperipheral heavy ion collisions at RHIC energies,” *Phys.Lett.* **B626** (2005) 72–79, arXiv:hep-ph/0505023 [hep-ph].
- [68] E. Witten, “Cosmic Separation of Phases,” *Phys.Rev.* **D30** (1984) 272–285.
- [69] R. Vogt, “Quark-Gluon Plasma Signatures.”
- [70] C. von Weizsäcker, “Radiation emitted in collisions of very fast electrons,” *Z.Phys.* **88** (1934) 612–625.
- [71] E. Williams, “Nature of the high-energy particles of penetrating radiation and status of ionization and radiation formulae,” *Phys.Rev.* **45** (1934) 729–730.
- [72] **CMS Collaboration** Collaboration, O. A. Grachov *et al.*, “Status of zero degree calorimeter for CMS experiment,” *AIP Conf.Proc.* **867** (2006) 258–265, arXiv:nucl-ex/0608052 [nucl-ex].
- [73] **CMS HCAL Collaboration** Collaboration, G. Baiatian *et al.*, “Design, Performance, and Calibration of CMS Hadron-Barrel Calorimeter Wedges.”
- [74] C. Albajar, J. de Troconiz, J. Rohlf, and G. Wrochna, “Conceptual design of an improved CMS RPC muon trigger using the hadron outer scintillators,” *Nucl.Instrum.Meth.* **A545** (2005) 97–113.
- [75] **CMS Collaboration** Collaboration, S. Dasu *et al.*, “CMS. The TriDAS project. Technical design report, vol. 1: The trigger systems.”
- [76] **CMS Collaboration** Collaboration, P. Sphicas, “CMS: The TriDAS project. Technical design report, Vol. 2: Data acquisition and high-level trigger.”

- [77] S. Margetis, K. Safarik, and O. Villalobos Baillie, “Strangeness production in heavy-ion collisions,” *Ann.Rev.Nucl.Part.Sci.* **50** (2000) 299–342.
- [78] T. Matsui and H. Satz, “ $J/\psi$  Suppression by Quark-Gluon Plasma Formation,” *Phys.Lett.* **B178** (1986) 416.
- [79] M. Gyulassy and M. Plumer, “Jet Quenching in Dense Matter,” *Phys.Lett.* **B243** (1990) 432–438.
- [80] A. Baltz, G. Baur, D. d’Enterria, L. Frankfurt, F. Gelis, *et al.*, “The Physics of Ultraperipheral Collisions at the LHC,” *Phys.Rept.* **458** (2008) 1–171, arXiv:0706.3356 [nucl-ex].
- [81] C. A. Bertulani, S. R. Klein, and J. Nystrand, “Physics of ultra-peripheral nuclear collisions,” *Ann.Rev.Nucl.Part.Sci.* **55** (2005) 271–310, arXiv:nucl-ex/0502005 [nucl-ex].
- [82] F. Karsch, E. Laermann, and A. Peikert, “Quark mass and flavor dependence of the QCD phase transition,” *Nucl.Phys.* **B605** (2001) 579–599, arXiv:hep-lat/0012023 [hep-lat].
- [83] S. Hands, “The Phase diagram of QCD,” *Contemp.Phys.* **42** (2001) 209–225, arXiv:physics/0105022 [physics.ed-ph].
- [84] L. Evans and P. Bryant, “LHC Machine,” *JINST* **3** (2008) S08001.
- [85] M. L. Miller, K. Reygers, S. J. Sanders, and P. Steinberg, “Glauber modeling in high energy nuclear collisions,” *Ann.Rev.Nucl.Part.Sci.* **57** (2007) 205–243, arXiv:nucl-ex/0701025 [nucl-ex].
- [86] G. Roland, K. Safarik, and P. Steinberg, “Heavy-ion collisions at the LHC,” *Prog.Part.Nucl.Phys.* **77** (2014) 70–127.

- [87] **CMS Collaboration** Collaboration, S. Chatrchyan *et al.*, “Multiplicity and transverse momentum dependence of two- and four-particle correlations in pPb and PbPb collisions,” *Phys.Lett.* **B724** (2013) 213–240, arXiv:1305.0609 [nucl-ex].
- [88] **ALICE Collaboration** Collaboration, B. Abelev *et al.*, “Long-range angular correlations on the near and away side in  $p$ -Pb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV,” *Phys.Lett.* **B719** (2013) 29–41, arXiv:1212.2001 [nucl-ex].
- [89] **ALICE Collaboration** Collaboration, K. Aamodt *et al.*, “The ALICE experiment at the CERN LHC,” *JINST* **3** (2008) S08002.
- [90] **ATLAS Collaboration** Collaboration, G. Aad *et al.*, “The ATLAS Experiment at the CERN Large Hadron Collider,” *JINST* **3** (2008) S08003.
- [91] **LHCb Collaboration** Collaboration, J. Alves, A. Augusto *et al.*, “The LHCb Detector at the LHC,” *JINST* **3** (2008) S08005.
- [92] J. Bjorken, “Highly Relativistic Nucleus-Nucleus Collisions: The Central Rapidity Region,” *Phys.Rev.* **D27** (1983) 140–151.
- [93] **STAR Collaboration** Collaboration, K. Ackermann *et al.*, “Elliptic flow in Au + Au collisions at  $(S(NN))^{**}(1/2) = 130$  GeV,” *Phys.Rev.Lett.* **86** (2001) 402–407, arXiv:nucl-ex/0009011 [nucl-ex].
- [94] J.-Y. Ollitrault, “Anisotropy as a signature of transverse collective flow,” *Phys.Rev.* **D46** (1992) 229–245.
- [95] P. Sorensen, “Elliptic Flow: A Study of Space-Momentum Correlations In Relativistic Nuclear Collisions,” arXiv:0905.0174 [nucl-ex].
- [96] N. Borghini, R. Bhalerao, and J. Ollitrault, “Anisotropic flow from Lee-Yang zeroes: A Practical guide,” *J.Phys.* **G30** (2004) S1213–S1216, arXiv:nucl-th/0402053 [nucl-th].

- [97] R. Bhalerao, N. Borghini, and J. Ollitrault, “Genuine collective flow from Lee-Yang zeroes,” *Phys.Lett.* **B580** (2004) 157–162, arXiv:nucl-th/0307018 [nucl-th].
- [98] N. Borghini, P. M. Dinh, and J.-Y. Ollitrault, “Flow analysis from cumulants: A Practical guide,” arXiv:nucl-ex/0110016 [nucl-ex].
- [99] V. Andreev, X. Aslanoglou, A. Azman, M. Bakirci, S. Basegmez, *et al.*, “Performance studies of a full-length prototype for the CASTOR forward calorimeter at the CMS experiment,” *Eur.Phys.J.* **C67** (2010) 601–615.
- [100] L. McLerran, “The Color Glass Condensate and Glasma,” arXiv:0804.1736 [hep-ph].
- [101] F. Gelis, “Color glass condensate and glasma,” *Nuclear Physics A* **854** no. 1, (2011) 10 – 17. <http://www.sciencedirect.com/science/article/pii/S0375947410006615>.  
Saturation, the Color Glass Condensate and the Glasma: What Have We Learned from RHIC?
- [102] M. M. I. Bloch, E. J. A., E. D. Barge, C. Campagnari, P. Kalavase, V. Krutelyov, D. Kovalskyi, J. Ribnik, and N. Amapane, “Muon identification in cms,” *CMS Note* **2008/098** (2008) . <http://cdsweb.cern.ch/record/962045>.
- [103] L. Frankfurt, M. Strikman, and M. Zhalov, “Large t diffractive  $\rho$ -meson photoproduction with target dissociation in ultraperipheral p a and a a collisions at lhc,” hep-ph/0612072.
- [104] **H1** Collaboration, A. Aktas *et al.*, “Diffractive photoproduction of  $\rho$  mesons with large momentum transfer at hera,” *Phys. Lett.* **B638** (2006) 422–431, hep-ex/0603038.
- [105] **STAR** Collaboration, C. Adler *et al.*, “Coherent  $\rho^0$  production in ultra-peripheral heavy ion collisions,” *Phys. Rev. Lett.* **89** (2002) 272302, nucl-ex/0206004.
- [106] T. Teubner, “Diffractive production of vector mesons and the gluon at small  $x$ ,” *AIP Conf. Proc.* **792** (2005) 416–419.

- [107] M. Klasen, “Hard photoproduction at hera,” hep-ph/0702052.
- [108] D. d’Enterria, “Hard spectra and qcd matter: Experimental review,” *J. Phys. G***30** (2004) S767–S774, nucl-ex/0404018.
- [109] M. Strikman, R. Vogt, and S. White, “Probing small  $x$  parton densities in ultraperipheral a a and p a collisions at the lhc,” *Phys. Rev. Lett.* **96** (2006) 082001, hep-ph/0508296.
- [110] J. Nystrand, “Electromagnetic interactions in nucleus nucleus and proton proton collisions,” *Nucl. Phys.* **A752** (2005) 470–479, hep-ph/0412096.
- [111] S. R. Klein and J. Nystrand, “Photoproduction of quarkonium in proton proton and nucleus nucleus collisions,” *Phys. Rev. Lett.* **92** (2004) 142003, hep-ph/0311164.
- [112] A. J. Baltz, S. R. Klein, and J. Nystrand, “Coherent vector meson photoproduction with nuclear breakup in relativistic heavy ion collisions,” *Phys. Rev. Lett.* **89** (2002) 012301, nucl-th/0205031.
- [113] S. Klein and J. Nystrand, “Exclusive vector meson production in relativistic heavy ion collisions,” *Phys. Rev.* **C60** (1999) 014903, hep-ph/9902259.
- [114] L. Frankfurt, M. Strikman, and C. Weiss, “Small- $x$  physics: From hera to lhc and beyond,” *Ann. Rev. Nucl. Part. Sci.* **55** (2005) 403–465, hep-ph/0507286.
- [115] V. P. Goncalves and M. V. T. Machado, “Heavy quarks and meson production in ultraperipheral heavy ion collisions,” hep-ph/0601131.
- [116] B. L. Berman and S. C. Fultz *Rev. Mod. Phys.* **47** (1975) 713.
- [117] R. Vogt, “Heavy quark production in heavy ion colliders,” *Heavy Ion Phys.* **18** (2003) 11–20, hep-ph/0205330.
- [118] K. Piotrzkowski, “Tagging two-photon production at the lhc,” *Phys. Rev.* **D63** (2001) 071502, hep-ex/0009065.

- [119] M. W. Krasny, J. Chwastowski, and K. Slowikowski, “Luminosity measurement method for lhc: The theoretical precision and the experimental challenges,” hep-ex/0610052.
- [120] **Particle Data Group** Collaboration, W.-M. Yao *et al.*, “Review of particle physics,” *J. Phys. G* **33** (2006) 1–1232.
- [121] V. A. Khoze, A. D. Martin, and M. G. Ryskin, “Double-diffractive processes in high-resolution missing- mass experiments at the tevatron,” *Eur. Phys. J. C* **19** (2001) 477–483, hep-ph/0011393.
- [122] M. Albrow, 2001. private communication.
- [123] I. A. Pshenichnov, J. P. Bondorf, I. N. Mishustin, A. Ventura, and S. Masetti, “Mutual heavy ion dissociation in peripheral collisions at ultrarelativistic energies,” *Phys. Rev. C* **64** (2001) 024903, nucl-th/0101035.
- [124] M. Boonekamp *et al.* Atl-gen-2004-001, March, 2004.
- [125] S. Baffioni, C. Charlot, *et al.*, “Electron reconstruction in cms,” *CMS NOTE 2006/40* (2006) .
- [126] P. Yepes, “Studying light objects in ultra-peripheral heavy ion collisions,” *CMS NOTE 2003/022* (2003) .
- [127] A. Lardeux, “ $\text{J}/\psi$  production in Pb-Pb collisions at  $\sqrt{s_{NN}} = 2.76 \text{ TeV}$  in the ALICE experiment,” *J.Phys.Conf.Ser.* **446** (2013) 012042.
- [128] S. Chatrchyan *et al.*, “The cms experiment at the cern lhc,” *Journal of Instrumentation* **3** no. 08, (2008) S08004. <http://stacks.iop.org/1748-0221/3/i=08/a=S08004>.
- [129] M. B. G. Ducati, M. T. Griep, and M. V. T. Machado, “Exclusive photoproduction of  $\text{j}/\psi$  and  $\psi(2s)$  states in proton-proton collisions at the cern lhc,” Tech. Rep. arXiv:1305.4611, May, 2013. Comments: 04 pages, 2 figures. arXiv admin note: text overlap with arXiv:1305.2407.

- [130] E. Fermi, “On the Theory of Collisions between Atoms and Electrically Charged Particles,” in *Electromagnetic Probes of Fundamental Physics*, W. Marciano and S. White, eds., pp. 243–252. Sept., 2003. hep-th/0205086.
- [131] **PHENIX Collaboration** Collaboration, A. Adare *et al.*, “Cold Nuclear Matter Effects on  $J/\psi$  Yields as a Function of Rapidity and Nuclear Geometry in Deuteron-Gold Collisions at  $\sqrt{s_{NN}} = 200$  GeV,” *Phys.Rev.Lett.* **107** (2011) 142301, arXiv:1010.1246 [nucl-ex].
- [132] C. A. Brau, *Modern Problems in Classical Electrodynamics*. Oxford, New York, NY, 1st ed. ed., 2004.
- [133] S. R. Klein and J. Nystrand, “Exclusive vector meson production in relativistic heavy ion collisions,” *Phys. Rev. C* **60** (Jun, 1999) 014903.  
<http://link.aps.org/doi/10.1103/PhysRevC.60.014903>.
- [134] S. R. Klein and J. Nystrand, “Interference in exclusive vector meson production in heavy-ion collisions,” *Phys. Rev. Lett.* **84** (Mar, 2000) 2330–2333.  
<http://link.aps.org/doi/10.1103/PhysRevLett.84.2330>.
- [135] V. Rebyakova, M. Strikman, and M. Zhalov, “Coherent rho and J/psi photoproduction in ultraperipheral processes with electromagnetic dissociation of heavy ions at RHIC and LHC,” *Phys.Lett.* **B710** (2012) 647–653, arXiv:1109.0737 [hep-ph].
- [136] V. Rebyakova, M. Strikman, and M. Zhalov, “Coherent  $\rho$  and  $J/\Psi$  photoproduction in ultraperipheral processes with electromagnetic dissociation of heavy ions at RHIC and LHC,” *Physics Letters B* **710** no. 4–5, (2012) 647 – 653.  
<http://www.sciencedirect.com/science/article/pii/S0370269312003152>.
- [137] V. Guzey, E. Kryshen, M. Strikman, and M. Zhalov, “Evidence for nuclear gluon shadowing from the ALICE measurements of pbpb ultraperipheral exclusive production,” *Physics Letters B* **726** no. 1–3, (2013) 290 – 295.  
<http://www.sciencedirect.com/science/article/pii/S0370269313006825>.

- [138] A. Adeluyi and C. A. Bertulani, “Gluon distributions in nuclei probed at energies available at the CERN large hadron collider,” *Phys. Rev. C* **84** (Aug, 2011) 024916.  
<http://link.aps.org/doi/10.1103/PhysRevC.84.024916>.
- [139] A. Adeluyi and T. Nguyen, “Coherent photoproduction of  $\psi$  and  $\Upsilon$  mesons in ultraperipheral pPb and PbPb collisions at the CERN Large Hadron Collider at  $\sqrt{S_{NN}} = 5$  TeV and  $\sqrt{S_{NN}} = 2.76$  TeV,” *Phys. Rev. C* **87** (Feb, 2013) 027901, arXiv:1302.4288 [nucl-th]. <http://link.aps.org/doi/10.1103/PhysRevC.87.027901>.
- [140] T. C. collaboration, “Performance of cms muon reconstruction in pp collision events at  $\sqrt{s} = 7$  tev,” *Journal of Instrumentation* **7** no. 10, (2012) P10002.  
<http://stacks.iop.org/1748-0221/7/i=10/a=P10002>.
- [141] **CMS Collaboration** Collaboration, S. Chatrchyan *et al.*, “Measurement of CMS Luminosity.”
- [142] **CMS Collaboration** Collaboration, G. Bayatian *et al.*, “CMS technical design report, volume II: Physics performance,” *J.Phys. G* **34** (2007) 995–1579.
- [143] **CMS Collaboration** Collaboration, *The CMS tracker: addendum to the Technical Design Report*. Technical Design Report CMS. CERN, Geneva, 2000.
- [144] P. Faccioli, C. Lourenco, J. Seixas, and H. K. Wohri, “Towards the experimental clarification of quarkonium polarization,” *Eur.Phys.J. C* **69** (2010) 657–673, arXiv:1006.2738 [hep-ph].
- [145] J. Alwall, A. Ballestrero, P. Bartalini, S. Belov, E. Boos, *et al.*, “A Standard format for Les Houches event files,” *Comput.Phys.Commun.* **176** (2007) 300–304, arXiv:hep-ph/0609017 [hep-ph].

- [146] T. Sjöstrand, S. Mrenna, and P. Skands, “Pythia 6.4 physics and manual,” *Journal of High Energy Physics* **2006** no. 05, (2006) 026.  
<http://stacks.iop.org/1126-6708/2006/i=05/a=026>.
- [147] P. Steinberg, “Relativistic heavy ion physics: Results from AGS to RHIC,” *eConf* **C020620** (2002) FRBT03, arXiv:nucl-ex/0210009 [nucl-ex].
- [148] H. Satz, “The SPS heavy ion programme,” *Phys.Rept.* **403-404** (2004) 33–50, arXiv:hep-ph/0405051 [hep-ph].
- [149] M. Chiu, A. Denisov, E. Garcia, J. Katzy, A. Makeev, M. Murray, and S. White, “Measurement of mutual coulomb dissociation in  $\sqrt{s_{NN}} = 130\text{GeV}$   $au + au$  collisions,” *Phys. Rev. Lett.* **89** (Jun, 2002) 012302.  
<http://link.aps.org/doi/10.1103/PhysRevLett.89.012302>.
- [150] M. Wilde, “Measurement of direct photons in pp and pb–pb collisions with {ALICE},” *Nuclear Physics A* **904–905** no. 0, (2013) 573c – 576c.  
<http://www.sciencedirect.com/science/article/pii/S0375947413001954>. The Quark Matter 2012 Proceedings of the {XXIII} International Conference on Ultrarelativistic Nucleus-Nucleus Collisions.
- [151] J. D. Bjorken, “Highly relativistic nucleus-nucleus collisions: The central rapidity region,” *Phys. Rev. D* **27** (Jan, 1983) 140–151.  
<http://link.aps.org/doi/10.1103/PhysRevD.27.140>.
- [152] P. Stankus, “Direct photon production in relativistic heavy-ion collisions,” *Annual Review of Nuclear and Particle Science* **55** no. 1, (2005) 517–554,  
<http://dx.doi.org/10.1146/annurev.nucl.53.041002.110533>.  
<http://dx.doi.org/10.1146/annurev.nucl.53.041002.110533>.

- [153] H. Song, S. A. Bass, U. Heinz, T. Hirano, and C. Shen, “200  $a$  gev Au + Au collisions serve a nearly perfect quark-gluon liquid,” *Phys. Rev. Lett.* **106** (May, 2011) 192301.  
<http://link.aps.org/doi/10.1103/PhysRevLett.106.192301>.
- [154] B. Alver *et al.*, “Importance of correlations and fluctuations on the initial source eccentricity in high-energy nucleus-nucleus collisions,” *Phys. Rev. C* **77** (Jan, 2008) 014906. <http://link.aps.org/doi/10.1103/PhysRevC.77.014906>.
- [155] A. J. Baltz, C. Chasman, and S. N. White, “Correlated forward–backward dissociation and neutron spectra as a luminosity monitor in heavy-ion colliders,” *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment* **417** no. 1, (1998) 1 – 8.  
<http://www.sciencedirect.com/science/article/pii/S0168900298005750>.
- [156] I. A. Pshenichnov, J. P. Bondorf, I. N. Mishustin, A. Ventura, and S. Masetti, “Mutual heavy ion dissociation in peripheral collisions at ultrarelativistic energies,” *Phys. Rev. C* **64** (Jul, 2001) 024903. <http://link.aps.org/doi/10.1103/PhysRevC.64.024903>.
- [157] I. A. Pshenichnov, I. N. Mishustin, J. P. Bondorf, A. S. Botvina, and A. S. Iljinov, “Particle emission following coulomb excitation in ultrarelativistic heavy-ion collisions,” *Phys. Rev. C* **60** (Sep, 1999) 044901. <http://link.aps.org/doi/10.1103/PhysRevC.60.044901>.
- [158] P. Chomaz, “Collective excitations in nuclei,”  
<http://starlight.hepforge.org>.
- [159] <http://starlight.hepforge.org>.
- [160] S. Chatrchyan *et al.*, “J/ $\psi$  and  $\psi(2s)$  production in pp collisions at  $\sqrt{s} = 7$  tev,” *Journal of High Energy Physics* **2012** no. 2, (2012) .  
<http://dx.doi.org/10.1007/JHEP02%282012%29011>.

## **Appendix A**

### **My Appendix, Next to my Spleen**

There could be lots of stuff here