
Reparameterization invariance in approximate Bayesian inference

Hrittik Roy[†], Marco Miani[†]

Technical University of Denmark
{hroy, mmia}@dtu.dk

Carl Henrik Ek

University of Cambridge,
Karolinska Institutet
che29@cam.ac.uk

Philipp Hennig, Marvin Pfortner, Lukas Tatzel

University of Tübingen, Tübingen AI Center
{philipp.hennig, lukas.tatzel,
marvin.pfoertner}@uni-tuebingen.de

Søren Hauberg

Technical University of Denmark
sohau@dtu.dk

Why linearised Laplace > regular Laplace

October 2024

Section 1: Laplace Approximation in BNNs

- Neural network $f_{\mathbf{w}} : \mathbb{R}^I \rightarrow \mathbb{R}^O$ with likelihood $p(\mathbf{y}|f_{\mathbf{w}}(\mathbf{x}))$ and prior $p(\mathbf{w})$

Section 1: Laplace Approximation in BNNs

- Neural network $f_{\mathbf{w}} : \mathbb{R}^I \rightarrow \mathbb{R}^O$ with likelihood $p(\mathbf{y}|f_{\mathbf{w}}(\mathbf{x}))$ and prior $p(\mathbf{w})$
- Take second-order Taylor expansion of log-posterior around a mode $\hat{\mathbf{w}}$

$$\log p(\mathbf{x}, \mathbf{y}; \mathbf{w}) \approx \log p(\mathbf{x}, \mathbf{y}; \hat{\mathbf{w}}) - \frac{1}{2}(\mathbf{w} - \hat{\mathbf{w}})^T (-\nabla_{\mathbf{w}}^2 \log p(\mathbf{x}, \mathbf{y}; \mathbf{w})|_{\hat{\mathbf{w}}})(\mathbf{w} - \hat{\mathbf{w}})$$

Section 1: Laplace Approximation in BNNs

- Neural network $f_{\mathbf{w}} : \mathbb{R}^I \rightarrow \mathbb{R}^O$ with likelihood $p(\mathbf{y}|f_{\mathbf{w}}(\mathbf{x}))$ and prior $p(\mathbf{w})$
- Take second-order Taylor expansion of log-posterior around a mode $\hat{\mathbf{w}}$

$$\log p(\mathbf{x}, \mathbf{y}; \mathbf{w}) \approx \log p(\mathbf{x}, \mathbf{y}; \hat{\mathbf{w}}) - \frac{1}{2}(\mathbf{w} - \hat{\mathbf{w}})^T (-\nabla_{\mathbf{w}}^2 \log p(\mathbf{x}, \mathbf{y}; \mathbf{w})|_{\hat{\mathbf{w}}})(\mathbf{w} - \hat{\mathbf{w}})$$

- Approximate posterior as $p(\mathbf{w}|\mathbf{x}, \mathbf{y}) \approx \mathcal{N}(\mathbf{w}|\hat{\mathbf{w}}, -\mathbf{H}_{\hat{\mathbf{w}}}^{-1})$
- with Hessian matrix $\mathbf{H}_{\hat{\mathbf{w}}} = \nabla_{\mathbf{w}}^2 \log p(\mathbf{x}, \mathbf{y}; \mathbf{w})|_{\hat{\mathbf{w}}}$

Linearised Laplace BNNs

- Linearise the NN at $\hat{\mathbf{w}}$

$$f_{\mathbf{w}}(\mathbf{x}) \approx f_{\hat{\mathbf{w}}}(\mathbf{x}) + \mathbf{J}_{\hat{\mathbf{w}}}(\mathbf{x})(\mathbf{w} - \hat{\mathbf{w}})$$

- using Jacobian $\mathbf{J}_{\hat{\mathbf{w}}}(\mathbf{x}) = \partial_{\mathbf{w}} f_{\mathbf{w}}(\mathbf{x})|_{\mathbf{w}=\hat{\mathbf{w}}} \in \mathbb{R}^{O \times D}$ $D = \dim(\mathbf{w})$

Linearised Laplace BNNs

- Linearise the NN at $\hat{\mathbf{w}}$

$$f_{\mathbf{w}}(\mathbf{x}) \approx f_{\hat{\mathbf{w}}}(\mathbf{x}) + \mathbf{J}_{\hat{\mathbf{w}}}(\mathbf{x})(\mathbf{w} - \hat{\mathbf{w}})$$

- using Jacobian $\mathbf{J}_{\hat{\mathbf{w}}}(\mathbf{x}) = \partial_{\mathbf{w}} f_{\mathbf{w}}(\mathbf{x})|_{\mathbf{w}=\hat{\mathbf{w}}} \in \mathbb{R}^{O \times D}$ $D = \dim(\mathbf{w})$

- Linearised Laplace approximation with prior $\mathcal{N}(\mathbf{0}, \alpha^{-1}\mathbf{I})$ gives

$$q(\mathbf{w}|\mathcal{D}) = \mathcal{N}(\mathbf{w} \mid \hat{\mathbf{w}}, (\text{GGN}_{\hat{\mathbf{w}}} + \alpha\mathbf{I})^{-1})$$

Linearised Laplace BNNs

- Linearise the NN at $\hat{\mathbf{w}}$

$$f_{\mathbf{w}}(\mathbf{x}) \approx f_{\hat{\mathbf{w}}}(\mathbf{x}) + \mathbf{J}_{\hat{\mathbf{w}}}(\mathbf{x})(\mathbf{w} - \hat{\mathbf{w}})$$

- using Jacobian $\mathbf{J}_{\hat{\mathbf{w}}}(\mathbf{x}) = \partial_{\mathbf{w}} f_{\mathbf{w}}(\mathbf{x})|_{\mathbf{w}=\hat{\mathbf{w}}} \in \mathbb{R}^{O \times D}$ $D = \dim(\mathbf{w})$

- Linearised Laplace approximation with prior $\mathcal{N}(\mathbf{0}, \alpha^{-1}\mathbf{I})$ gives

$$q(\mathbf{w}|\mathcal{D}) = \mathcal{N}(\mathbf{w} \mid \hat{\mathbf{w}}, (\text{GGN}_{\hat{\mathbf{w}}} + \alpha\mathbf{I})^{-1})$$

$$\text{GGN}_{\hat{\mathbf{w}}} = \sum_{n=1}^N \mathbf{J}_{\hat{\mathbf{w}}}(\mathbf{x}_n)^\top \mathbf{H}(\mathbf{x}_n) \mathbf{J}_{\hat{\mathbf{w}}}(\mathbf{x}_n)$$

Linearised Laplace BNNs

- Linearise the NN at $\hat{\mathbf{w}}$

$$f_{\mathbf{w}}(\mathbf{x}) \approx f_{\hat{\mathbf{w}}}(\mathbf{x}) + \mathbf{J}_{\hat{\mathbf{w}}}(\mathbf{x})(\mathbf{w} - \hat{\mathbf{w}})$$

- using Jacobian $\mathbf{J}_{\hat{\mathbf{w}}}(\mathbf{x}) = \partial_{\mathbf{w}} f_{\mathbf{w}}(\mathbf{x})|_{\mathbf{w}=\hat{\mathbf{w}}} \in \mathbb{R}^{O \times D}$ $D = \dim(\mathbf{w})$

- Linearised Laplace approximation with prior $\mathcal{N}(\mathbf{0}, \alpha^{-1}\mathbf{I})$ gives

$$q(\mathbf{w}|\mathcal{D}) = \mathcal{N}(\mathbf{w} \mid \hat{\mathbf{w}}, (\text{GGN}_{\hat{\mathbf{w}}} + \alpha\mathbf{I})^{-1})$$

$$\text{GGN}_{\hat{\mathbf{w}}} = \sum_{n=1}^N \mathbf{J}_{\hat{\mathbf{w}}}(\mathbf{x}_n)^\top \mathbf{H}(\mathbf{x}_n) \mathbf{J}_{\hat{\mathbf{w}}}(\mathbf{x}_n) \quad \mathbf{H}(\mathbf{x}) = -\partial_{f_{\hat{\mathbf{w}}}(\mathbf{x})}^2 \log p(\mathbf{y}|f_{\hat{\mathbf{w}}}(\mathbf{x})) \in \mathbb{R}^{O \times O}$$

Laplace vs Linearised Laplace

- GGN is commonly used to approximate the Hessian $\mathbf{H}_{\hat{\mathbf{w}}}$
- Immer et al. (2021) argue that this choice implicitly linearises the BNN

$$q(\mathbf{w}|\mathcal{D}) \approx \mathcal{N}(\mathbf{w}|\hat{\mathbf{w}}, -\mathbf{H}_{\hat{\mathbf{w}}}^{-1}) \qquad q(\mathbf{w}|\mathcal{D}) = \mathcal{N}(\mathbf{w}|\hat{\mathbf{w}}, (\text{GGN}_{\hat{\mathbf{w}}} + \alpha\mathbf{I})^{-1})$$

Laplace vs Linearised Laplace

- GGN is commonly used to approximate the Hessian $\mathbf{H}_{\hat{\mathbf{w}}}$
- Immer et al. (2021) argue that this choice implicitly linearises the BNN

$$q(\mathbf{w}|\mathcal{D}) \approx \mathcal{N}(\mathbf{w}|\hat{\mathbf{w}}, -\mathbf{H}_{\hat{\mathbf{w}}}^{-1}) \qquad q(\mathbf{w}|\mathcal{D}) = \mathcal{N}(\mathbf{w} | \hat{\mathbf{w}}, (\text{GGN}_{\hat{\mathbf{w}}} + \alpha \mathbf{I})^{-1})$$

$$\mathbf{H}_{\hat{\mathbf{w}}} = \nabla_{\mathbf{w}}^2 \log p(\mathbf{x}, \mathbf{y}; \mathbf{w})|_{\hat{\mathbf{w}}}$$

Laplace vs Linearised Laplace

- GGN is commonly used to approximate the Hessian $\mathbf{H}_{\hat{\mathbf{w}}}$
- Immer et al. (2021) argue that this choice implicitly linearises the BNN

$$q(\mathbf{w}|\mathcal{D}) \approx \mathcal{N}(\mathbf{w}|\hat{\mathbf{w}}, -\mathbf{H}_{\hat{\mathbf{w}}}^{-1})$$

$$q(\mathbf{w}|\mathcal{D}) = \mathcal{N}(\mathbf{w} | \hat{\mathbf{w}}, (\text{GGN}_{\hat{\mathbf{w}}} + \alpha \mathbf{I})^{-1})$$

$$\mathbf{H}_{\hat{\mathbf{w}}} = \nabla_{\mathbf{w}}^2 \log p(\mathbf{x}, \mathbf{y}; \mathbf{w})|_{\hat{\mathbf{w}}}$$

$$\text{GGN}_{\hat{\mathbf{w}}} = \sum_{n=1}^N \mathbf{J}_{\hat{\mathbf{w}}}(\mathbf{x}_n)^{\top} \mathbf{H}(\mathbf{x}_n) \mathbf{J}_{\hat{\mathbf{w}}}(\mathbf{x}_n)$$
$$\mathbf{H}(\mathbf{x}) = -\partial_{f_{\hat{\mathbf{w}}}(\mathbf{x})}^2 \log p(\mathbf{y}|f_{\hat{\mathbf{w}}}(\mathbf{x})) \in \mathbb{R}^{O \times O}$$


(If likelihood is gaussian, then $\mathbf{H}(\mathbf{x}) = \mathbb{I}_O$)

Predictive distributions

$$q(\mathbf{w}|\mathcal{D}) = \mathcal{N}(\mathbf{w} \mid \hat{\mathbf{w}}, (\mathbf{G}\mathbf{G}\mathbf{N}_{\hat{\mathbf{w}}} + \alpha\mathbf{I})^{-1})$$

- Regular Laplace severely underfits

$$p(\mathbf{y}^*|\mathbf{x}^*, \mathcal{D}) = \mathbb{E}_{\mathbf{w} \sim q}[p(\mathbf{y}^*|f(\mathbf{w}, \mathbf{x}^*))] \approx \frac{1}{S} \sum_{i=1}^S p(\mathbf{y}^*|f(\mathbf{w}_i, \mathbf{x}^*)), \quad \mathbf{w}_i \sim q$$




$$q(\mathbf{w}|\mathcal{D}) = \mathcal{N}(\mathbf{w} \mid \hat{\mathbf{w}}, (\text{GGN}_{\hat{\mathbf{w}}} + \alpha \mathbf{I})^{-1})$$

Predictive distributions


- Regular Laplace severely underfits

$$p(\mathbf{y}^*|\mathbf{x}^*, \mathcal{D}) = \mathbb{E}_{\mathbf{w} \sim q}[p(\mathbf{y}^*|f(\mathbf{w}, \mathbf{x}^*))] \approx \frac{1}{S} \sum_{i=1}^S p(\mathbf{y}^*|f(\mathbf{w}_i, \mathbf{x}^*)), \quad \mathbf{w}_i \sim q$$



- Linearised predictions perform much better

$$p(\mathbf{y}^*|\mathbf{x}^*, \mathcal{D}) = \mathbb{E}_{\mathbf{w} \sim q}[p(\mathbf{y}^*|f_{\text{lin}}^{\hat{\mathbf{w}}}(\mathbf{w}, \mathbf{x}^*))] \approx \frac{1}{S} \sum_{i=1}^S p(\mathbf{y}^*|f_{\text{lin}}^{\hat{\mathbf{w}}}(\mathbf{w}_i, \mathbf{x}^*)), \quad \mathbf{w}_i \sim q$$




$$q(\mathbf{w}|\mathcal{D}) = \mathcal{N}(\mathbf{w} \mid \hat{\mathbf{w}}, (\text{GGN}_{\hat{\mathbf{w}}} + \alpha \mathbf{I})^{-1})$$

Predictive distributions


- Regular Laplace severely underfits

$$p(\mathbf{y}^*|\mathbf{x}^*, \mathcal{D}) = \mathbb{E}_{\mathbf{w} \sim q}[p(\mathbf{y}^*|f(\mathbf{w}, \mathbf{x}^*))] \approx \frac{1}{S} \sum_{i=1}^S p(\mathbf{y}^*|f(\mathbf{w}_i, \mathbf{x}^*)), \quad \mathbf{w}_i \sim q$$



- Linearised predictions perform much better

$$p(\mathbf{y}^*|\mathbf{x}^*, \mathcal{D}) = \mathbb{E}_{\mathbf{w} \sim q}[p(\mathbf{y}^*|f_{\text{lin}}^{\hat{\mathbf{w}}}(\mathbf{w}, \mathbf{x}^*))] \approx \frac{1}{S} \sum_{i=1}^S p(\mathbf{y}^*|f_{\text{lin}}^{\hat{\mathbf{w}}}(\mathbf{w}_i, \mathbf{x}^*)), \quad \mathbf{w}_i \sim q$$



- Why does adding another degree of approximation improve performance?

This paper: linearised laplace is better because it's invariant to reparameterisation


- BNNs are massively overparameterised, with many points in weight-space corresponding to identical functions

This paper: linearised laplace is better because it's invariant to reparameterisation

- BNNs are massively overparameterised, with many points in weight-space corresponding to identical functions
- Consider the NN $f(x) = w_1 \text{ReLU}(w_2 x); f : \mathbb{R} \rightarrow \mathbb{R}$
- For any $\alpha > 0$, (w_1, w_2) and $(w_1/\alpha, \alpha w_2)$ are equivalent


This paper: linearised laplace is better because it's invariant to reparameterisation

- BNNs are massively overparameterised, with many points in weight-space corresponding to identical functions
- Consider the NN $f(x) = w_1 \text{ReLU}(w_2 x)$; $f : \mathbb{R} \rightarrow \mathbb{R}$
- For any $\alpha > 0$, (w_1, w_2) and $(w_1/\alpha, \alpha w_2)$ are equivalent




This paper: linearised laplace is better because it's invariant to reparameterisation

- BNNs are massively overparameterised, with many points in weight-space corresponding to identical functions
- Consider the NN $f(x) = w_1 \text{ReLU}(w_2 x); f: \mathbb{R} \rightarrow \mathbb{R}$
- For any $\alpha > 0$, (w_1, w_2) and $(w_1/\alpha, \alpha w_2)$ are equivalent
- The GGN covariance naturally aligns the linearised Laplace approx. post. with the reparameterisation (kernel) direction




This paper: linearised laplace is better because it's invariant to reparameterisation

- The regular Laplace approximation puts a crude pdf over the space of (nonlinear) NN functions
 - Identical functions are given different masses (this also messes up marginal likelihood estimates)



This paper: linearised laplace is better because it's invariant to reparameterisation

- The regular Laplace approximation puts a crude pdf over the space of (nonlinear) NN functions
 - Identical functions are given different masses (this also messes up marginal likelihood estimates)



Section 2: Reparameterisation of linear functions

- Consider a linear function $f(\mathbf{w}) = \mathbf{A}\mathbf{w} + \mathbf{b}$
and a reparameterisation $g : \mathbb{R}^D \rightarrow \mathbb{R}^D$ such that $f(g(\mathbf{w})) = f(\mathbf{w})$

Section 2: Reparameterisation of linear functions

- Consider a linear function $f(\mathbf{w}) = \mathbf{A}\mathbf{w} + \mathbf{b}$
and a reparameterisation $g : \mathbb{R}^D \rightarrow \mathbb{R}^D$ such that $f(g(\mathbf{w})) = f(\mathbf{w})$
- Then $\mathbf{A}(g(\mathbf{w}) - \mathbf{w}) = \mathbf{0}$

Section 2: Reparameterisation of linear functions

- Consider a linear function $f(\mathbf{w}) = \mathbf{A}\mathbf{w} + \mathbf{b}$
and a reparameterisation $g : \mathbb{R}^D \rightarrow \mathbb{R}^D$ such that $f(g(\mathbf{w})) = f(\mathbf{w})$
- Then $\mathbf{A}(g(\mathbf{w}) - \mathbf{w}) = \mathbf{0}$
- So the direction of movement between \mathbf{w} and $g(\mathbf{w})$ lies in the kernel/nullspace of \mathbf{A}

$$g(\mathbf{w}) - \mathbf{w} \in \ker(\mathbf{A})$$

Find the kernel of the linearised NN (and avoid it!)

$$f_{\text{lin}}^{\mathbf{w}'} : \mathbf{w}, \mathbf{x} \mapsto f_{\mathbf{w}'}(\mathbf{x}) + \mathbf{J}_{\mathbf{w}'}(\mathbf{x})(\mathbf{w} - \mathbf{w}')$$

Find the kernel of the linearised NN (and avoid it!)

$$f_{\text{lin}}^{\mathbf{w}'} : \mathbf{w}, \mathbf{x} \mapsto f_{\mathbf{w}'}(\mathbf{x}) + \mathbf{J}_{\mathbf{w}'}(\mathbf{x})(\mathbf{w} - \mathbf{w}')$$

$$\ker(\mathbf{J}_{\mathbf{w}}) = \ker(\text{GGN}_{\mathbf{w}})$$

$$\begin{aligned}\text{GGN}_{\mathbf{w}} &= \mathbf{J}_{\mathbf{w}}^{\top} \mathbf{J}_{\mathbf{w}} \\ \text{NTK}_{\mathbf{w}} &= \mathbf{J}_{\mathbf{w}} \mathbf{J}_{\mathbf{w}}^{\top}\end{aligned}$$

Find the kernel of the linearised NN (and avoid it!)

$$f_{\text{lin}}^{\mathbf{w}'} : \mathbf{w}, \mathbf{x} \mapsto f_{\mathbf{w}'}(\mathbf{x}) + \mathbf{J}_{\mathbf{w}'}(\mathbf{x})(\mathbf{w} - \mathbf{w}')$$

$$\ker(\mathbf{J}_{\mathbf{w}}) = \ker(\text{GGN}_{\mathbf{w}})$$

$$\begin{aligned}\text{GGN}_{\mathbf{w}} &= \mathbf{J}_{\mathbf{w}}^{\top} \mathbf{J}_{\mathbf{w}} \\ \text{NTK}_{\mathbf{w}} &= \mathbf{J}_{\mathbf{w}} \mathbf{J}_{\mathbf{w}}^{\top}\end{aligned}$$

- Because the GGN is a self-adjoint operator (positive semi-definite matrix)

$$\text{im}(\text{GGN}_{\mathbf{w}}) \oplus \ker(\text{GGN}_{\mathbf{w}}) = \mathbb{R}^D$$

Find the kernel of the linearised NN (and avoid it!)


$$f_{\text{lin}}^{\mathbf{w}'} : \mathbf{w}, \mathbf{x} \mapsto f_{\mathbf{w}'}(\mathbf{x}) + \mathbf{J}_{\mathbf{w}'}(\mathbf{x})(\mathbf{w} - \mathbf{w}')$$

$$\ker(\mathbf{J}_{\mathbf{w}}) = \ker(\text{GGN}_{\mathbf{w}})$$

$$\begin{aligned}\text{GGN}_{\mathbf{w}} &= \mathbf{J}_{\mathbf{w}}^{\top} \mathbf{J}_{\mathbf{w}} \\ \text{NTK}_{\mathbf{w}} &= \mathbf{J}_{\mathbf{w}} \mathbf{J}_{\mathbf{w}}^{\top}\end{aligned}$$

- Because the GGN is a self-adjoint operator (positive semi-definite matrix)

$$\text{im}(\text{GGN}_{\mathbf{w}}) \oplus \ker(\text{GGN}_{\mathbf{w}}) = \mathbb{R}^D$$



Find the kernel of the linearised NN (and avoid it!)

$$f_{\text{lin}}^{\mathbf{w}'} : \mathbf{w}, \mathbf{x} \mapsto f_{\mathbf{w}'}(\mathbf{x}) + \mathbf{J}_{\mathbf{w}'}(\mathbf{x})(\mathbf{w} - \mathbf{w}')$$


$$\ker(\mathbf{J}_{\mathbf{w}}) = \ker(\text{GGN}_{\mathbf{w}})$$

$$\begin{aligned}\text{GGN}_{\mathbf{w}} &= \mathbf{J}_{\mathbf{w}}^{\top} \mathbf{J}_{\mathbf{w}} \\ \text{NTK}_{\mathbf{w}} &= \mathbf{J}_{\mathbf{w}} \mathbf{J}_{\mathbf{w}}^{\top}\end{aligned}$$

- Because the GGN is a self-adjoint operator (positive semi-definite matrix)

$$\text{im}(\text{GGN}_{\mathbf{w}}) \oplus \ker(\text{GGN}_{\mathbf{w}}) = \mathbb{R}^D$$

- $\text{im}(\text{GGN}_{\mathbf{w}})$ spans to the *effective parameters*
 - (ideally we'd just explore this)



Find the kernel of the linearised NN (and avoid it!)

$$f_{\text{lin}}^{\mathbf{w}'} : \mathbf{w}, \mathbf{x} \mapsto f_{\mathbf{w}'}(\mathbf{x}) + \mathbf{J}_{\mathbf{w}'}(\mathbf{x})(\mathbf{w} - \mathbf{w}')$$


$$\ker(\mathbf{J}_{\mathbf{w}}) = \ker(\text{GGN}_{\mathbf{w}})$$

$$\begin{aligned}\text{GGN}_{\mathbf{w}} &= \mathbf{J}_{\mathbf{w}}^{\top} \mathbf{J}_{\mathbf{w}} \\ \text{NTK}_{\mathbf{w}} &= \mathbf{J}_{\mathbf{w}} \mathbf{J}_{\mathbf{w}}^{\top}\end{aligned}$$

- Because the GGN is a self-adjoint operator (positive semi-definite matrix)

$$\text{im}(\text{GGN}_{\mathbf{w}}) \oplus \ker(\text{GGN}_{\mathbf{w}}) = \mathbb{R}^D$$

- $\text{im}(\text{GGN}_{\mathbf{w}})$ spans to the *effective parameters*
 - (ideally we'd just explore this)
- $\ker(\text{GGN}_{\mathbf{w}})$ corresponds to the directions of reparameterisation



Laplace samples can be decomposed into image and kernel contributions

- Let $\mathbf{U}^T \mathbf{\Lambda} \mathbf{U}$ be the eigendecomposition of $\mathbf{GGN}_{\hat{\mathbf{w}}}$ with \mathbf{U}_1 and \mathbf{U}_2 corresponding to non-zero and zero eigenvalues respectively

Laplace samples can be decomposed into image and kernel contributions

- Let $\mathbf{U}^T \mathbf{\Lambda} \mathbf{U}$ be the eigendecomposition of $\text{GGN}_{\hat{\mathbf{w}}}$ with \mathbf{U}_1 and \mathbf{U}_2 corresponding to non-zero and zero eigenvalues respectively

$$\Sigma = \left(\begin{bmatrix} \mathbf{U}_1 \\ \mathbf{U}_2 \end{bmatrix}^T \begin{bmatrix} \tilde{\mathbf{\Lambda}} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} \begin{bmatrix} \mathbf{U}_1 \\ \mathbf{U}_2 \end{bmatrix} + \alpha \mathbf{I} \right)^{-1} = \mathbf{U}_1^T (\tilde{\mathbf{\Lambda}} + \alpha \mathbf{I}_k)^{-1} \mathbf{U}_1 + \alpha^{-1} \mathbf{U}_2^T \mathbf{U}_2$$

Laplace samples can be decomposed into image and kernel contributions

- Let $\mathbf{U}^T \mathbf{\Lambda} \mathbf{U}$ be the eigendecomposition of $\text{GGN}_{\hat{\mathbf{W}}}$ with \mathbf{U}_1 and \mathbf{U}_2 corresponding to non-zero and zero eigenvalues respectively

$$\Sigma = \left(\begin{bmatrix} \mathbf{U}_1 \\ \mathbf{U}_2 \end{bmatrix}^T \begin{bmatrix} \tilde{\mathbf{\Lambda}} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} \begin{bmatrix} \mathbf{U}_1 \\ \mathbf{U}_2 \end{bmatrix} + \alpha \mathbf{I} \right)^{-1} = \mathbf{U}_1^T (\tilde{\mathbf{\Lambda}} + \alpha \mathbf{I}_k)^{-1} \mathbf{U}_1 + \alpha^{-1} \mathbf{U}_2^T \mathbf{U}_2$$

So any Laplace sample can be written $\mathbf{W} = \hat{\mathbf{W}} + \mathbf{W}_{\text{im}} + \mathbf{W}_{\text{ker}}$

Laplace samples can be decomposed into image and kernel contributions

- Let $\mathbf{U}^T \mathbf{\Lambda} \mathbf{U}$ be the eigendecomposition of $\text{GGN}_{\hat{\mathbf{W}}}$ with \mathbf{U}_1 and \mathbf{U}_2 corresponding to non-zero and zero eigenvalues respectively

$$\Sigma = \left(\begin{bmatrix} \mathbf{U}_1 \\ \mathbf{U}_2 \end{bmatrix}^T \begin{bmatrix} \tilde{\mathbf{\Lambda}} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} \begin{bmatrix} \mathbf{U}_1 \\ \mathbf{U}_2 \end{bmatrix} + \alpha \mathbf{I} \right)^{-1} = \mathbf{U}_1^T (\tilde{\mathbf{\Lambda}} + \alpha \mathbf{I}_k)^{-1} \mathbf{U}_1 + \alpha^{-1} \mathbf{U}_2^T \mathbf{U}_2$$

So any Laplace sample can be written $\mathbf{W} = \hat{\mathbf{W}} + \mathbf{W}_{\text{im}} + \mathbf{W}_{\text{ker}}$

(Note that all probability mass in the kernel comes from the prior)

Linearised laplace automatically ignores \mathbf{W}_{ker}

$$f_{\text{lin}}^{\hat{\mathbf{W}}}(\hat{\mathbf{W}} + \mathbf{W}_{\text{ker}} + \mathbf{W}_{\text{im}}, \mathbf{X}) = f_{\text{lin}}^{\hat{\mathbf{W}}}(\hat{\mathbf{W}} + \mathbf{W}_{\text{im}}, \mathbf{X})$$

Linearised laplace automatically ignores \mathbf{W}_{ker}

$$f_{\text{lin}}^{\hat{\mathbf{W}}}(\hat{\mathbf{W}} + \mathbf{W}_{\text{ker}} + \mathbf{W}_{\text{im}}, \mathbf{X}) = f_{\text{lin}}^{\hat{\mathbf{W}}}(\hat{\mathbf{W}} + \mathbf{W}_{\text{im}}, \mathbf{X})$$

- It doesn't matter which reparameterisation gets sampled if your model is linear!

Linearised laplace automatically ignores \mathbf{W}_{ker}

$$f_{\text{lin}}^{\hat{\mathbf{W}}}(\hat{\mathbf{W}} + \mathbf{W}_{\text{ker}} + \mathbf{W}_{\text{im}}, \mathbf{X}) = f_{\text{lin}}^{\hat{\mathbf{W}}}(\hat{\mathbf{W}} + \mathbf{W}_{\text{im}}, \mathbf{X})$$

- It doesn't matter which reparameterisation gets sampled if your model is linear!
- Regular Laplace doesn't ignore \mathbf{W}_{ker} because the kernel doesn't correspond to reparameterisations

Linearised laplace automatically ignores \mathbf{W}_{ker}

$$f_{\text{lin}}^{\hat{\mathbf{W}}}(\hat{\mathbf{W}} + \mathbf{W}_{\text{ker}} + \mathbf{W}_{\text{im}}, \mathbf{X}) = f_{\text{lin}}^{\hat{\mathbf{W}}}(\hat{\mathbf{W}} + \mathbf{W}_{\text{im}}, \mathbf{X})$$

- It doesn't matter which reparameterisation gets sampled if your model is linear!
- Regular Laplace doesn't ignore \mathbf{W}_{ker} because the kernel doesn't correspond to reparameterisations
 - Adds 'incorrect' degrees of freedom to the approx. post.

Linearised laplace automatically ignores \mathbf{W}_{ker}

$$f_{\text{lin}}^{\hat{\mathbf{W}}}(\hat{\mathbf{W}} + \mathbf{W}_{\text{ker}} + \mathbf{W}_{\text{im}}, \mathbf{X}) = f_{\text{lin}}^{\hat{\mathbf{W}}}(\hat{\mathbf{W}} + \mathbf{W}_{\text{im}}, \mathbf{X})$$


- It doesn't matter which reparameterisation gets sampled if your model is linear!
- Regular Laplace doesn't ignore \mathbf{W}_{ker} because the kernel doesn't correspond to reparameterisations
 - Adds 'incorrect' degrees of freedom to the approx. post.
- This suggests that if $\text{ker}(\text{GGN}_{\hat{\mathbf{W}}})$ is small (i.e. $\text{GGN}_{\hat{\mathbf{W}}}$ has a high rank) then regular Laplace should perform similarly to linearised Laplace

Linearised laplace automatically ignores \mathbf{W}_{ker}

$$f_{\text{lin}}^{\hat{\mathbf{W}}}(\hat{\mathbf{W}} + \mathbf{W}_{\text{ker}} + \mathbf{W}_{\text{im}}, \mathbf{X}) = f_{\text{lin}}^{\hat{\mathbf{W}}}(\hat{\mathbf{W}} + \mathbf{W}_{\text{im}}, \mathbf{X})$$


- It doesn't matter which reparameterisation gets sampled if your model is linear!
- Regular Laplace doesn't ignore \mathbf{W}_{ker} because the kernel doesn't correspond to reparameterisations
 - Adds 'incorrect' degrees of freedom to the approx. post.
- This suggests that if $\text{ker}(\text{GGN}_{\hat{\mathbf{W}}})$ is small (i.e. $\text{GGN}_{\hat{\mathbf{W}}}$ has a high rank) then regular Laplace should perform similarly to linearised Laplace

Small CNN on MNIST so GGN can be computed exactly – (rank increases with more data if we keep the same number of parameters)




Section 3: Reparameterisation of NNs

- They do the same analysis but for nonlinear models (e.g. regular NNs)





Section 3: Reparameterisation of NNs

- They do the same analysis but for nonlinear models (e.g. regular NNs)
- We can similarly decompose Laplace samples into a combination of two manifolds embedded in \mathbb{R}^D , $(\mathcal{P}_{\mathbf{w}}, \mathfrak{m})$ and $(\mathcal{P}_{\mathbf{w}}^\perp, \mathfrak{m}^\perp)$, which act like $\text{im}(\text{GGN}_{\mathbf{w}})$ and $\text{ker}(\text{GGN}_{\mathbf{w}})$





Section 3: Reparameterisation of NNs

- They do the same analysis but for nonlinear models (e.g. regular NNs)
- We can similarly decompose Laplace samples into a combination of two manifolds embedded in \mathbb{R}^D , $(\mathcal{P}_{\mathbf{w}}, \mathbf{m})$ and $(\mathcal{P}_{\mathbf{w}}^\perp, \mathbf{m}^\perp)$, which act like $\text{im}(\text{GGN}_{\mathbf{w}})$ and $\text{ker}(\text{GGN}_{\mathbf{w}})$



Section 3: Reparameterisation of NNs

- They do the same analysis but for nonlinear models (e.g. regular NNs)
- We can similarly decompose Laplace samples into a combination of two manifolds embedded in \mathbb{R}^D , $(\mathcal{P}_{\mathbf{w}}, \mathbf{m})$ and $(\mathcal{P}_{\mathbf{w}}^\perp, \mathbf{m}^\perp)$, which act like $\text{im}(\text{GGN}_{\mathbf{w}})$ and $\text{ker}(\text{GGN}_{\mathbf{w}})$
- Then we can ignore the reparameterisation directions by only exploring $(\mathcal{P}_{\mathbf{w}}, \mathbf{m})$ (*Laplace diffusion*)



Define Effective Parameter-Space as a Quotient Group

$\mathbf{w}_1 \sim \mathbf{w}_2$ iff there exists a smooth path between \mathbf{w}_1 and \mathbf{w}_2 such that

$$f(\mathbf{w}_1, \mathbf{x}) = f(\mathbf{w}_2, \mathbf{x}) \quad \forall \mathbf{x} \in \mathcal{D}_{\text{train}}$$

Define Effective Parameter-Space as a Quotient Group

$\mathbf{w}_1 \sim \mathbf{w}_2$ iff there exists a smooth path between \mathbf{w}_1 and \mathbf{w}_2 such that

$$f(\mathbf{w}_1, \mathbf{x}) = f(\mathbf{w}_2, \mathbf{x}) \quad \forall \mathbf{x} \in \mathcal{D}_{\text{train}}$$

Then we define the effective parameter space as the quotient group $\mathcal{P} = \mathbb{R}^D / \sim$

(i.e. only consider parameters that give us unique functions)

Define a Reparameterisation Distance and get a (Pseudo-)Riemannian manifold

We'd like to define a distance such that $\text{dist}(\mathbf{w}_1, \mathbf{w}_2) = 0 \iff \mathbf{w}_1 \sim \mathbf{w}_2$

Define a Reparameterisation Distance and get a (Pseudo-)Riemannian manifold

We'd like to define a distance such that $\text{dist}(\mathbf{w}_1, \mathbf{w}_2) = 0 \iff \mathbf{w}_1 \sim \mathbf{w}_2$

$$\text{dist}^2(\mathbf{w}, \mathbf{w} + \boldsymbol{\epsilon}) = \sum_{n=1}^N \|f(\mathbf{w}, \mathbf{x}_n) - f(\mathbf{w} + \boldsymbol{\epsilon}, \mathbf{x}_n)\|^2 = \boldsymbol{\epsilon}^\top \text{GGN}_{\mathbf{w}} \boldsymbol{\epsilon} + \mathcal{O}(\epsilon^3)$$

Define a Reparameterisation Distance and get a (Pseudo-)Riemannian manifold

We'd like to define a distance such that $\text{dist}(\mathbf{w}_1, \mathbf{w}_2) = 0 \iff \mathbf{w}_1 \sim \mathbf{w}_2$

$$\text{dist}^2(\mathbf{w}, \mathbf{w} + \epsilon) = \sum_{n=1}^N \|f(\mathbf{w}, \mathbf{x}_n) - f(\mathbf{w} + \epsilon, \mathbf{x}_n)\|^2 = \epsilon^\top \text{GGN}_{\mathbf{w}} \epsilon + \mathcal{O}(\epsilon^3)$$

So the GGN matrix infinitesimally defines an inner product, a (pseudo-)Riemannian metric!

Define a Reparameterisation Distance and get a (Pseudo-)Riemannian manifold

We'd like to define a distance such that $\text{dist}(\mathbf{w}_1, \mathbf{w}_2) = 0 \iff \mathbf{w}_1 \sim \mathbf{w}_2$

$$\text{dist}^2(\mathbf{w}, \mathbf{w} + \epsilon) = \sum_{n=1}^N \|f(\mathbf{w}, \mathbf{x}_n) - f(\mathbf{w} + \epsilon, \mathbf{x}_n)\|^2 = \epsilon^\top \text{GGN}_{\mathbf{w}} \epsilon + \mathcal{O}(\epsilon^3)$$

So the GGN matrix infinitesimally defines an inner product, a (pseudo-)Riemannian metric!

NOTE:

1. GGN is rank-deficient (hence pseudo-riemannian)

Define a Reparameterisation Distance and get a (Pseudo-)Riemannian manifold

We'd like to define a distance such that $\text{dist}(\mathbf{w}_1, \mathbf{w}_2) = 0 \iff \mathbf{w}_1 \sim \mathbf{w}_2$

$$\text{dist}^2(\mathbf{w}, \mathbf{w} + \epsilon) = \sum_{n=1}^N \|f(\mathbf{w}, \mathbf{x}_n) - f(\mathbf{w} + \epsilon, \mathbf{x}_n)\|^2 = \epsilon^\top \text{GGN}_{\mathbf{w}} \epsilon + \mathcal{O}(\epsilon^3)$$

So the GGN matrix infinitesimally defines an inner product, a (pseudo-)Riemannian metric!

NOTE:

1. GGN is rank-deficient (hence pseudo-riemannian)
2. $\mathbf{w}_1 \sim \mathbf{w}_2$ doesn't then necessarily mean $\text{GGN}_{\mathbf{w}_1} = \text{GGN}_{\mathbf{w}_2}$ (pseudo-metric might be different based on your Laplace centre/mode (... but infinitesimally that doesn't matter too much))

Equivalence of the manifold and the quotient group

Proposition 4.4. *There exists a bijection between $(\mathbb{R}^D, \text{GGN}_{\mathbf{w}})$ and \mathcal{P} .*

Equivalence of the manifold and the quotient group

Proposition 4.4. *There exists a bijection between $(\mathbb{R}^D, \text{GGN}_{\mathbf{w}})$ and \mathcal{P} .*

(Basically, using the tangent space of the GGN gives us a path along the homotopies in the equivalence classes.)

Equivalence of the manifold and the quotient group

Proposition 4.4. *There exists a bijection between $(\mathbb{R}^D, \text{GGN}_{\mathbf{w}})$ and \mathcal{P} .*

(Basically, using the tangent space of the GGN gives us a path along the homotopies in the equivalence classes.)

- Why do we care? Answer: Riemannian diffusion.

Equivalence of the manifold and the quotient group

Proposition 4.4. *There exists a bijection between $(\mathbb{R}^D, \text{GGN}_{\mathbf{w}})$ and \mathcal{P} .*

(Basically, using the tangent space of the GGN gives us a path along the homotopies in the equivalence classes.)

- Why do we care? Answer: Riemannian diffusion.
- We can run a Markov chain to give us samples from the Riemannian manifold (\mathbf{M}, \mathbf{G})

Equivalence of the manifold and the quotient group

Proposition 4.4. *There exists a bijection between $(\mathbb{R}^D, \text{GGN}_{\mathbf{w}})$ and \mathcal{P} .*

(Basically, using the tangent space of the GGN gives us a path along the homotopies in the equivalence classes.)

- Why do we care? Answer: Riemannian diffusion.
- We can run a Markov chain to give us samples from the Riemannian manifold (\mathbf{M}, \mathbf{G})

$$\mathbf{w}_{t+1} = \mathbf{w}_t + \sqrt{2h_t} \mathbf{G}(\mathbf{w}_t)^{-\frac{1}{2}} \epsilon, \text{ where } \epsilon \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$$

Equivalence of the manifold and the quotient group

Proposition 4.4. *There exists a bijection between $(\mathbb{R}^D, \text{GGN}_{\mathbf{w}})$ and \mathcal{P} .*


(Basically, using the tangent space of the GGN gives us a path along the homotopies in the equivalence classes.)

- Why do we care? Answer: Riemannian diffusion.
- We can run a Markov chain to give us samples from the Riemannian manifold (\mathbf{M}, \mathbf{G})

$$\mathbf{w}_{t+1} = \mathbf{w}_t + \sqrt{2h_t} \mathbf{G}(\mathbf{w}_t)^{-\frac{1}{2}} \epsilon, \text{ where } \epsilon \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$$


Problem: $(\mathbb{R}^D, \text{GGN}_{\mathbf{w}})$ is only pseudo-Riemannian (GGN is rank-deficient)

Define two Riemannian manifolds which act as $\text{im}(\text{GGN}_{\mathbf{w}})$ and $\text{ker}(\text{GGN}_{\mathbf{w}})$




Define two Riemannian manifolds which act as $\text{im}(\text{GGN}_{\mathbf{w}})$ and $\text{ker}(\text{GGN}_{\mathbf{w}})$

1. $(\mathcal{P}_{\mathbf{w}}, \mathfrak{m})$ where $\mathfrak{m} = \text{GGN}_{\mathbf{w}}^+$




Define two Riemannian manifolds which act as $\text{im}(\text{GGN}_{\mathbf{w}})$ and $\text{ker}(\text{GGN}_{\mathbf{w}})$

1. $(\mathcal{P}_{\mathbf{w}}, \mathfrak{m})$ where $\mathfrak{m} = \text{GGN}_{\mathbf{w}}^+$
 - Never intersects the same equivalence class more than once (at least locally)



Define two Riemannian manifolds which act as $\text{im}(\text{GGN}_{\mathbf{w}})$ and $\text{ker}(\text{GGN}_{\mathbf{w}})$


1. $(\mathcal{P}_{\mathbf{w}}, \mathfrak{m})$ where $\mathfrak{m} = \text{GGN}_{\mathbf{w}}^+$
 - Never intersects the same equivalence class more than once (at least locally)
 - Like the effective parameter manifold (but actually Riemannian)



Define two Riemannian manifolds which act as $\text{im}(\text{GGN}_{\mathbf{w}})$ and $\text{ker}(\text{GGN}_{\mathbf{w}})$


1. $(\mathcal{P}_{\mathbf{w}}, \mathfrak{m})$ where $\mathfrak{m} = \text{GGN}_{\mathbf{w}}^+$
 - Never intersects the same equivalence class more than once (at least locally)
 - Like the effective parameter manifold (but actually Riemannian)

2. $(\mathcal{P}_{\mathbf{w}}^{\perp}, \mathfrak{m}^{\perp})$ where $\mathfrak{m}^{\perp} = \alpha \mathbf{I}$ for $\alpha > 0$



Define two Riemannian manifolds which act as $\text{im}(\text{GGN}_{\mathbf{w}})$ and $\text{ker}(\text{GGN}_{\mathbf{w}})$

1. $(\mathcal{P}_{\mathbf{w}}, \mathfrak{m})$ where $\mathfrak{m} = \text{GGN}_{\mathbf{w}}^+$
 - Never intersects the same equivalence class more than once (at least locally)
 - Like the effective parameter manifold (but actually Riemannian)
2. $(\mathcal{P}_{\mathbf{w}}^{\perp}, \mathfrak{m}^{\perp})$ where $\mathfrak{m}^{\perp} = \alpha \mathbf{I}$ for $\alpha > 0$
 - Entirely contained in the same equivalence class $\mathcal{P}_{\mathbf{w}}^{\perp} \subseteq [\mathbf{w}]$




Define two Riemannian manifolds which act as $\text{im}(\text{GGN}_{\mathbf{w}})$ and $\text{ker}(\text{GGN}_{\mathbf{w}})$

1. $(\mathcal{P}_{\mathbf{w}}, \mathfrak{m})$ where $\mathfrak{m} = \text{GGN}_{\mathbf{w}}^+$

- Never intersects the same equivalence class more than once (at least locally)
- Like the effective parameter manifold (but actually Riemannian)

2. $(\mathcal{P}_{\mathbf{w}}^{\perp}, \mathfrak{m}^{\perp})$ where $\mathfrak{m}^{\perp} = \alpha \mathbf{I}$ for $\alpha > 0$

- Entirely contained in the same equivalence class $\mathcal{P}_{\mathbf{w}}^{\perp} \subseteq [\mathbf{w}]$
- So contains only functions which are identical on the training set





Section 4: Diffusion on various manifolds

0. Effective-parameter is only pseudo-Riemannian $(\mathbb{R}^D, \bar{G}G\mathbf{N}_{\mathbf{w}})$



Section 4: Diffusion on various manifolds

0. Effective-parameter is only pseudo-Riemannian $(\mathbb{R}^D, \mathbf{G}\mathbf{G}\mathbf{N}_{\mathbf{w}})$
1. Laplace Approximation $(\mathbb{R}^D, \mathbf{G}\mathbf{G}\mathbf{N}_{\mathbf{w}} + \alpha\mathbf{I})$





Section 4: Diffusion on various manifolds

0. Effective-parameter is only pseudo-Riemannian $(\mathbb{R}^D, \mathbf{G}\mathbf{G}\mathbf{N}_{\mathbf{w}})$
1. Laplace Approximation $(\mathbb{R}^D, \mathbf{G}\mathbf{G}\mathbf{N}_{\mathbf{w}} + \alpha\mathbf{I})$





Section 4: Diffusion on various manifolds

0. Effective-parameter is only pseudo-Riemannian $(\mathbb{R}^D, \mathbf{G}\mathbf{G}\mathbf{N}_{\mathbf{w}})$
1. Laplace Approximation $(\mathbb{R}^D, \mathbf{G}\mathbf{G}\mathbf{N}_{\mathbf{w}} + \alpha\mathbf{I})$
2. Kernel manifold (“never underfits”) $(\mathcal{P}_{\mathbf{w}}^{\perp}, \alpha\mathbf{I})$



Section 4: Diffusion on various manifolds

0. Effective-parameter is only pseudo-Riemannian $(\mathbb{R}^D, \mathbf{G}\mathbf{G}\mathbf{N}_{\mathbf{w}})$
1. Laplace Approximation $(\mathbb{R}^D, \mathbf{G}\mathbf{G}\mathbf{N}_{\mathbf{w}} + \alpha\mathbf{I})$
2. Kernel manifold (“never underfits”) $(\mathcal{P}_{\mathbf{w}}^{\perp}, \alpha\mathbf{I})$
3. Non-kernel-parameter manifold, i.e. “Laplace Diffusion” $(\mathcal{P}_{\mathbf{w}}, \mathbf{G}\mathbf{G}\mathbf{N}_{\mathbf{w}}^+)$



Section 4: Diffusion on various manifolds



0. Effective-parameter is only pseudo-Riemannian $(\mathbb{R}^D, \mathbf{G}\mathbf{G}\mathbf{N}_{\mathbf{w}})$

1. Laplace Approximation $(\mathbb{R}^D, \mathbf{G}\mathbf{G}\mathbf{N}_{\mathbf{w}} + \alpha\mathbf{I})$

2. Kernel manifold (“never underfits”) $(\mathcal{P}_{\mathbf{w}}^{\perp}, \alpha\mathbf{I})$

3. Non-kernel-parameter manifold, i.e. “Laplace Diffusion” $(\mathcal{P}_{\mathbf{w}}, \mathbf{G}\mathbf{G}\mathbf{N}_{\mathbf{w}}^+)$

4. Alternate between the two



Results

Table 1: In-distribution performance across methods trained on MNIST, FMNIST and CIFAR-10.

		Conf. (\uparrow)	NLL (\downarrow)	Acc. (\uparrow)	Brier (\downarrow)	ECE (\downarrow)	MCE (\downarrow)
MNIST	Laplace Diffusion (ours)	0.988\pm0.001	0.042\pm0.007	0.987\pm0.002	0.022\pm0.003	0.137\pm0.019	0.775\pm0.043
	Sampled Laplace	0.589 \pm 0.008	3.812 \pm 0.284	0.146 \pm 0.032	1.176 \pm 0.046	0.443 \pm 0.026	0.985 \pm 0.002
	Linearised Laplace	0.968 \pm 0.004	0.306 \pm 0.041	0.926 \pm 0.008	0.117 \pm 0.012	0.251 \pm 0.034	0.855 \pm 0.041
FMNIST	Laplace Diffusion (ours)	0.900\pm0.001	0.001\pm0.000	0.906\pm0.007	0.141\pm0.006	0.108\pm0.015	0.729\pm0.092
	Sampled Laplace	0.618 \pm 0.021	4.507 \pm 0.000	0.098 \pm 0.010	1.295 \pm 0.014	0.518 \pm 0.013	0.986 \pm 0.001
	Linearised Laplace	0.897 \pm 0.003	0.423 \pm 0.000	0.862 \pm 0.005	0.207 \pm 0.006	0.147 \pm 0.017	0.756 \pm 0.048
CIFAR-10	Laplace Diffusion (ours)	0.952\pm0.007	0.345\pm0.062	0.905\pm0.007	0.155\pm0.019	0.259 \pm 0.008	0.870 \pm 0.021
	Sampled Laplace	0.843 \pm 0.004	0.997 \pm 0.222	0.717 \pm 0.049	0.422 \pm 0.081	0.221\pm0.047	0.804 \pm 0.080
	Linearised Laplace	0.951 \pm 0.007	0.614 \pm 0.020	0.863 \pm 0.001	0.222 \pm 0.002	0.337 \pm 0.022	0.789\pm0.035

Results


Table 1: In-distribution performance across methods trained on MNIST, FMNIST and CIFAR-10.

		Conf. (\uparrow)	NLL (\downarrow)	Acc. (\uparrow)	Brier (\downarrow)	ECE (\downarrow)	MCE (\downarrow)
MNIST	Laplace Diffusion (ours)	0.988\pm0.001	0.042\pm0.007	0.987\pm0.002	0.022\pm0.003	0.137\pm0.019	0.775\pm0.043
	Sampled Laplace	0.589 \pm 0.008	3.812 \pm 0.284	0.146 \pm 0.032	1.176 \pm 0.046	0.443 \pm 0.026	0.985 \pm 0.002
	Linearised Laplace	0.968 \pm 0.004	0.306 \pm 0.041	0.926 \pm 0.008	0.117 \pm 0.012	0.251 \pm 0.034	0.855 \pm 0.041
FMNIST	Laplace Diffusion (ours)	0.900\pm0.001	0.001\pm0.000	0.906\pm0.007	0.141\pm0.006	0.108\pm0.015	0.729\pm0.092
	Sampled Laplace	0.618 \pm 0.021	4.507 \pm 0.000	0.098 \pm 0.010	1.295 \pm 0.014	0.518 \pm 0.013	0.986 \pm 0.001
	Linearised Laplace	0.897 \pm 0.003	0.423 \pm 0.000	0.862 \pm 0.005	0.207 \pm 0.006	0.147 \pm 0.017	0.756 \pm 0.048
CIFAR-10	Laplace Diffusion (ours)	0.952\pm0.007	0.345\pm0.062	0.905\pm0.007	0.155\pm0.019	0.259 \pm 0.008	0.870 \pm 0.021
	Sampled Laplace	0.843 \pm 0.004	0.997 \pm 0.222	0.717 \pm 0.049	0.422 \pm 0.081	0.221\pm0.047	0.804 \pm 0.080
	Linearised Laplace	0.951 \pm 0.007	0.614 \pm 0.020	0.863 \pm 0.001	0.222 \pm 0.002	0.337 \pm 0.022	0.789\pm0.035

Table 2: Out-of-distribution AUROC (\uparrow) performance for MNIST, FMNIST and CIFAR-10.

Trained on	MNIST			FMNIST			CIFAR-10	
Tested on	FMNIST	EMNIST	KMNIST	MNIST	EMNIST	KMNIST	CIFAR-100	SVHN
Laplace Diffusion (ours)	0.909\pm0.033	0.625\pm0.018	0.929\pm0.008	0.759\pm0.045	0.741\pm0.010	0.749\pm0.023	0.851\pm0.002	0.862\pm0.010
Sampled Laplace	0.500 \pm 0.026	0.494 \pm 0.006	0.482 \pm 0.013	0.495 \pm 0.037	0.503 \pm 0.036	0.493 \pm 0.033	0.687 \pm 0.033	0.599 \pm 0.038
Linearised Laplace	0.758 \pm 0.070	0.602 \pm 0.027	0.790 \pm 0.018	0.625 \pm 0.050	0.628 \pm 0.013	0.624 \pm 0.020	0.837 \pm 0.006	0.854 \pm 0.024

Rotated MNIST (shift = rotation angle)



Summary

- Linearised Laplace works better than full Laplace because it implicitly ignores the reparameterisation issues of BNNs

Summary

- Linearised Laplace works better than full Laplace because it implicitly ignores the reparameterisation issues of BNNs
- Their new method (*Laplace diffusion*) also tries to remove those reparameterisation issues (but is significantly more complicated)

Summary

- Linearised Laplace works better than full Laplace because it implicitly ignores the reparameterisation issues of BNNs
- Their new method (*Laplace diffusion*) also tries to remove those reparameterisation issues (but is significantly more complicated)
- In practice, we often use an approximation of the GGN (e.g. KFAC), which would break the motivation behind their new method

Links


Paper: <https://arxiv.org/pdf/2406.03334>

Slides from a talk by one of the authors:




https://www2.compute.dtu.dk/~sohau//talks/2024_MTNS/

Immer et al. (2021): “Improving predictions of Bayesian neural nets via local linearization,” Alexander Immer, Maciej Korzepa, Matthias Bauer. Proceedings of The 24th International Conference on Artificial Intelligence and Statistics, PMLR 130:703-711, 2021. <https://proceedings.mlr.press/v130/immer21a.html>


pics





— Reparametrization family ○ Metric



From slides





Section 4: They do diffusion

$$(\mathbf{M}, \mathbf{G})$$

$$d\mathbf{w} = \sqrt{2\tau} \mathbf{G}(\mathbf{w})^{-\frac{1}{2}} dW + \tau \Gamma dt \quad \text{where} \quad \Gamma_i(\mathbf{w}) = \sum_{j=1}^D \frac{\partial}{\partial \mathbf{w}_j} (\mathbf{G}(\mathbf{w})^{-1})_i$$

$$\mathbf{w}_{t+1} = \mathbf{w}_t + \sqrt{2h_t} \mathbf{G}(\mathbf{w}_t)^{-\frac{1}{2}} \boldsymbol{\epsilon}, \quad \text{where} \quad \boldsymbol{\epsilon} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$$