


- Often we wish to make predictions of quantities like  $\underline{C}\beta + Du$  for some estimable  $C\beta$ .
- The BLUP of such a quantity is  $C\hat{\beta}_{\Sigma} + D\hat{u}$ , the BLUE of  $C\beta$  plus  $D$  times the BLUP of  $u$ .

(E) for empirical  
whenever we need  
to estimate our  
variance components first:

$\begin{array}{c} \text{BLUE} \quad \text{(E)BLUP} \\ \underbrace{\hspace{10em}} \\ \text{(E)BLUP of } C\beta + Du \end{array}$

$u_1, \dots, u_n \sim N(0, \underline{\sigma_u^2})$  - variability in students reading ability scores


$e_1, \dots, e_n \sim N(0, \sigma_e^2)$  - variability in the actual test scores



we observe a student's test score:

$$y_{i,true} = \mu + \underbrace{u_i}_{\text{predict}} \quad y_i^*$$

- Suppose reading ability for a population of students are normally distributed with a mean  $\mu$  and variance  $\sigma_u^2$ .
- Suppose a reading ability test was given to an i.i.d. sample of such students. *sample from  $N(\mu, \sigma_u^2)$*
- Suppose that, given the true reading ability of a student at that time, the test score for that student is normally distributed with a mean equal to the student's reading ability and a variance  $\sigma_e^2$  and is independent of the test score of any other student.



- Suppose it is known that  $\sigma_u^2 / \sigma_e^2 = 9$ .

$$\bar{y} = 86$$

- If the sample mean of the students' test scores was 86, what is the best prediction of the reading ability of a student who scored 96 on the test?

the student's true reading ability

$$y_{i, \text{true}} = \mu + u_i$$

we observe

however  $y_i^* = 96$

$u$  &  $e$  are indep.

- Suppose  $\underbrace{u_1, \dots, u_n}_{i.i.d.} \stackrel{i.i.d.}{\sim} \mathcal{N}(0, \sigma_u^2)$  independent of  $\underbrace{e_1, \dots, e_n}_{i.i.d.} \stackrel{i.i.d.}{\sim} \mathcal{N}(0, \sigma_e^2)$ .
- If we let  $\mu + u_i$  denote the reading ability of student  $i$  ( $i = 1, \dots, n$ ), then the reading ability scores of the students are  $\mathcal{N}(\mu, \sigma_u^2)$  as in the statement of the problem.
- If we let  $y_i = \mu + u_i + e_i$  denote the test score of student  $i$  ( $i = 1, \dots, n$ ), then  $(y_i | \mu + u_i) \sim \mathcal{N}(\mu + u_i, \sigma_e^2)$  as in the problem statement.

We have  $\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \mathbf{Z}\mathbf{u} + \mathbf{e}$ , where

$$\underline{\mathbf{X} = \mathbf{1}}, \quad \underline{\boldsymbol{\beta} = \mu}, \quad \underline{\mathbf{Z} = \mathbf{I}}, \quad \underline{\mathbf{G} = \sigma_u^2 \mathbf{I}}, \quad \underline{\mathbf{R} = \sigma_e^2 \mathbf{I}}, \quad \text{and}$$
$$\boldsymbol{\Sigma} = \mathbf{Z}\mathbf{G}\mathbf{Z}^\top + \mathbf{R} = \underline{(\sigma_u^2 + \sigma_e^2)\mathbf{I}}.$$

Thus,

$$\hat{\boldsymbol{\beta}}_{\boldsymbol{\Sigma}} = (\mathbf{X}^\top \boldsymbol{\Sigma}^{-1} \mathbf{X})^{-1} \mathbf{X}^\top \boldsymbol{\Sigma}^{-1} \mathbf{y} = (\mathbf{1}^\top \mathbf{1})^{-1} \mathbf{1}^\top \mathbf{y} = \underline{\bar{y}}.$$

and

$$\mathbf{G}\mathbf{Z}^\top \boldsymbol{\Sigma}^{-1} = \frac{\sigma_u^2}{\sigma_u^2 + \sigma_e^2} \mathbf{I}.$$

in our ex.  
observed  
mean = 86

end lecture  
36  
4-25-25

Thus, the BLUP for  $u$  is

$$\hat{\mathbf{u}} = \underline{\mathbf{GZ}^\top \Sigma^{-1}} (\underline{\mathbf{y} - \mathbf{X}\hat{\boldsymbol{\beta}}_\Sigma}) = \boxed{\frac{\sigma_u^2}{\sigma_u^2 + \sigma_e^2}} (\mathbf{y} - \mathbf{1}\bar{y}).$$

The  $i^{th}$  element of this vector is

$$\hat{u}_i = \frac{\sigma_u^2}{\sigma_u^2 + \sigma_e^2} (y_i - \bar{y}).$$

Thus, the BLUP for  $\mu + u_i$  (the reading ability score of student  $i$ ) is

$$\hat{\mu} + \hat{u}_i = \bar{y}. + \frac{\sigma_u^2}{\sigma_u^2 + \sigma_e^2} (y_i - \bar{y}.) = \frac{\sigma_u^2}{\sigma_u^2 + \sigma_e^2} y_i + \frac{\sigma_e^2}{\sigma_u^2 + \sigma_e^2} \bar{y}.$$

Note that the BLUP is a convex combination of the individual score and the overall mean score.

$$\hat{\mu} + \hat{u}_i = \frac{\sigma_u^2}{\sigma_u^2 + \sigma_e^2} y_i + \frac{\sigma_e^2}{\sigma_u^2 + \sigma_e^2} \bar{y}.$$

empirical BLUP



Because  $\frac{\sigma_u^2}{\sigma_e^2}$  is assumed to be 9, the weights are

$$\frac{\sigma_u^2}{\sigma_u^2 + \sigma_e^2} = \frac{\frac{\sigma_u^2}{\sigma_e^2}}{\frac{\sigma_u^2}{\sigma_e^2} + 1} = \frac{9}{9 + 1} = 0.9$$

and

$$\frac{\sigma_e^2}{\sigma_u^2 + \sigma_e^2} = 0.1.$$

We would predict the IQ of a student who scored 130 on the test to be

$$0.9(96) + 0.1(86) = \underline{\underline{95}}$$

## EBLUPs in R for Seedling Dry Weight Example

2 genotype



```
> library(lme4)
>
> #Fit the linear mixed-effects model
> #with fixed genotype effects and
> #random tray effects.
>
> o = lmer(SeedlingWeight~Genotype+(1|Tray),data=d)
```

## EBLUPs in R for Seedling Dry Weight Example

```
> #uhat is the vector of the EBLUPs of Tray effects.  
>  
> ranef(o)  
$Tray  
  (Intercept)  
1  -4.985886  
2   2.622632  
3  -1.226723  
4   3.589977  
5   1.200572  
6  -1.666317  
7   3.104183  
8  -2.638439  
> uhat = ranef(o)$Tray[[1]]
```

# EBLUPs in R for Seedling Dry Weight Example

```
> #Get EBLUPs of genotype mean + tray effects
>
> betahat = fixef(o)
> betahat
(Intercept)      Genotype2
   15.288837    -3.550201
```

Handwritten annotations:

- $\hat{\mu}_i$  above genotype mean
- $\hat{u}_{ij}$  above tray
- $\hat{\mu}_1$  above Genotype2
- $\hat{\mu}_2 - \hat{\mu}_1 = -3.55$  next to the value -3.550201

```
> estGeno1Mean = as.numeric(betahat[1])
> estGeno2Mean = as.numeric(betahat[1] + betahat[2])
> EBLUPs = c(estGeno1Mean + uhat[1:4],
+            estGeno2Mean + uhat[5:8])
```

# EBLUPs in R for Seedling Dry Weight Example

```
> #Compare EBLUPs with tray averages.  
> trayAverages=tapply(d$SeedlingWeight,d$Tray,FUN = mean)  
> rbind(trayAverages, EBLUPs)
```

	1	2	3	4	5	6	7	8
trayAverages	10.0	18.0	14.0	19.0	13.0	10.0	15.0	9.00
<u>EBLUPs</u>	<u>10.3</u>	17.9	14.1	18.9	12.9	10.1	14.8	9.10

```
> estGenolMean
```

```
[1] 15.28884
```

```
> estGeno2Mean
```

```
[1] 11.73864
```

$$15.28884 - 4.98586$$

$$\approx \underline{\underline{10.3}}$$

end  
lecture 37  
4-28-25