

# Independence

## Examples

(\*\*)

Example 1: sampling with replacement (Bernoulli trials)

Let  $X_1, \dots, X_n$  be iid Bernoulli( $p$ ) r.v.s & let  $Y = \sum_{i=1}^n X_i$

$$M_{X_1+X_2+\dots+X_n}(t) \stackrel{\text{def}}{=} \mathbb{E} \left[ e^{t(X_1+\dots+X_n)} \right]$$

$$= \mathbb{E} \left[ e^{tX_1} e^{tX_2} \dots e^{tX_n} \right]$$

$$\stackrel{\substack{X_1, \dots, X_n \text{ are} \\ \text{ind.}}}{=} \mathbb{E} \left[ e^{tX_1} \right] \dots \mathbb{E} \left[ e^{tX_n} \right]$$

$$\stackrel{\substack{X_1, X_2, \dots, X_n \text{ are} \\ \text{Ber}(p)}}{=} [1-p+pe^t]^n$$

$$\Rightarrow Y = \sum_{i=1}^n X_i \sim \text{Bin}(n, p) \Rightarrow$$

$$\begin{cases} \mathbb{E}(Y) = np \checkmark \\ \text{Var}(Y) = np(1-p) \checkmark \end{cases}$$

$$\mathbb{E}(Y) = \mathbb{E} \left( \sum_{i=1}^n X_i \right) = \sum_{i=1}^n \mathbb{E}(X_i) = \sum_{i=1}^n p = np$$

$$\text{Var}(Y) = \text{Var} \left( \sum_{i=1}^n X_i \right) \stackrel{\substack{X_1, \dots, X_n \\ \text{are ind.}}}{=} \sum_{i=1}^n \text{Var}(X_i) \stackrel{\substack{X_1, \dots, X_n \\ \text{are Ber}(p)}}{=} \sum_{i=1}^n p(1-p) = np(1-p)$$

# Independence

Introduction

definition

Recall: Two events  $A, B$  are independent if  $P(A \cap B) = P(A)P(B)$

We next want to extend the concept of independence to random variables

Definition: Random variables  $X$  and  $Y$  are **independent** if

$$P(X \in A, Y \in B) = P(X \in A)P(Y \in B) \quad \text{for any } A, B \subset \mathbb{R}.$$

*Some Characterizations of Independence*

1. **Theorem:** In general, random variables  $X$  and  $Y$  are **independent** iff

$$P(X \leq x, Y \leq y) = F(x, y) = F_X(x)F_Y(y) \quad \text{for any } x, y \in \mathbb{R}.$$

$P(X \leq x) \cdot P(Y \leq y)$

If and only If

2. **Theorem:** Jointly discrete variables  $X$  and  $Y$  are **independent** iff

$$P(X = x, Y = y) = f(x, y) = f_X(x)f_Y(y) = P(X = x)P(Y = y) \quad \text{for any } x, y \in \mathbb{R}.$$

joint pmf

Product of marginal pmfs

3. **Theorem:** Jointly continuous variables  $X$  and  $Y$  are **independent** iff (there exist versions of the pdfs  $f(x, y)$ ,  $f_X(x)$  and  $f_Y(y)$  such that)

$$f(x, y) = f_X(x)f_Y(y) \quad \text{for any } x, y \in \mathbb{R}.$$


(The “there exist versions” business is needed because one can change densities at a few values (i.e., endpoints) without changing the integrals computed from them.)

# Independence

## Introduction (cont'd)

4. Furthermore, if  $X$  and  $Y$  are independent so that  $f(x, y) = f_X(x)f_Y(y)$  then

$$\begin{aligned}
 &f(x|y) = f_X(x) \quad \text{for any } x \in \mathbb{R}, \text{ any } y \text{ with } \underline{f_Y(y) > 0}; \\
 &f(y|x) = f_Y(y) \quad \text{for any } y \in \mathbb{R}, \text{ any } x \text{ with } \underline{f_X(x) > 0}.
 \end{aligned}$$



$$f(x|y) = \frac{f(x, y)}{f_Y(y)} \stackrel{\text{ind.}}{=} \frac{f_X(x) \cancel{f_Y(y)}}{\cancel{f_Y(y)}} = f_X(x)$$

In other words, under independence, the conditional distribution of  $Y$  given  $X = x$  is the same as the marginal distribution of  $Y$ . This is another way to think of independence: given information about one variable, the distribution of the 2nd variable does not change.

### *Two uses of independence*

1. examine a joint pdf to determine if the r.v.s are independent
2. assume independence and derive joint distribution from marginal distributions

# Independence

Some Examples

Discrete example:

		$x$			
		1	2	3	
$y$	3	1/12	1/12	1/6	1/3
	2	1/12	1/6	1/12	1/3
	1	1/6	1/12	1/12	1/3
		1/3	1/3	1/3	

Are  $X$  and  $Y$  independent?

$$f_{X,Y}(x,y) = f_X(x) f_Y(y)$$

$$\mathbb{P}(X=x, Y=y) = \mathbb{P}(X=x) \cdot \mathbb{P}(Y=y)$$

$$\mathbb{P}(X=2, Y=1) = 1/12 \quad *$$

$$\mathbb{P}(X=2) \cdot \mathbb{P}(Y=1) = 1/3 \cdot 1/3 = 1/9 \quad **$$

\*) \*  $\Rightarrow X$  and  $Y$  are NOT independent.

Continuous example (the familiar one):  $f(x,y) = 1/x, 0 < y < x < 1$ .

Are  $X$  and  $Y$  independent?

$$\rightarrow Y|X \sim \text{Uni}(0,x) \Rightarrow f(y|x) = \frac{1}{x} \quad \forall 0 < y < x$$

$$f_Y(y) = \begin{cases} -\log y & 0 < y < 1 \\ 0 & \text{otherwise} \end{cases} \quad f_Y(y) = -\log y \quad 0 < y < 1$$

$$\Rightarrow f(y|x) \neq f_Y(y) \Rightarrow X \text{ and } Y \text{ are NOT independent.}$$

Another continuous example:  $f(x,y) = 2y, 0 \leq x < 1, 0 \leq y < 1$ .

Are  $X$  and  $Y$  independent?

$$f_X(x) = \int_0^1 f_{X,Y}(x,y) dy = \int_0^1 2y dy = y^2 \Big|_0^1 = 1$$

$$\Rightarrow f_X(x) = \begin{cases} 0 & \text{If } x \notin (0,1) \\ 1 & \text{If } x \in (0,1) \end{cases} \quad \text{I}$$

$$f_Y(y) = \int_0^1 f(x,y) dx = \int_0^1 2y dx = 2y x \Big|_0^1 = 2y \quad \forall 0 < y < 1$$


$$f_Y(y) = \begin{cases} 2y & 0 < y < 1 \\ 0 & \text{otherwise} \end{cases} \quad \text{II}$$

$$2y = f_{X,Y}(x,y) = f_X(x) f_Y(y) = 1 \cdot 2y \quad \forall (x,y) \in (0,1) \times (0,1)$$

$X$  and  $Y$  are independent.

# Independence

Some important results & facts



1. Theorem: If  $X, Y$  are independent then  $g(X)$  and  $h(Y)$  are independent.

Proof:  $\mathbb{P}(g(X) \in A \text{ and } h(Y) \in B)$

$= \mathbb{P}(X \in g^{-1}(A) \text{ and } Y \in h^{-1}(B))$

$\xrightarrow{\text{X and Y are ind.}} \mathbb{P}(X \in g^{-1}(A)) \mathbb{P}(Y \in h^{-1}(B))$

$= \mathbb{P}(g(X) \in A) \cdot \mathbb{P}(h(Y) \in B)$

ex:  $X$  and  $Y$  are independent then  $e^X$  and  $\sin Y$  are independent.

2. Theorem: If  $X, Y$  are independent and the mean of  $g(X)h(Y)$  exists, then

$\mathbb{E}[g(X)h(Y)] = \mathbb{E}[g(X)]\mathbb{E}[h(Y)]$

$\mathbb{E}(XY) = \mathbb{E}(X)\mathbb{E}(Y)$

$\mathbb{E}(Z(X,Y)) \stackrel{\text{def}}{=} \int \int Z(x,y) f_{XY}(x,y) dx dy$

$= \int \int g(x)h(y) f_X(x) f_Y(y) dx dy$

$= \int g(x) f_X(x) dx \int h(y) f_Y(y) dy = \mathbb{E}[g(X)] \mathbb{E}[h(Y)]$

3. Corollary: If  $X, Y$  are independent and  $EXY$  exists, then

Proof:  $\text{Cov}(X,Y) \stackrel{\text{def}}{=} \mathbb{E}[XY] - \mathbb{E}[X]\mathbb{E}[Y]$

$\stackrel{\text{X and Y are ind.}}{=} \mathbb{E}[X]\mathbb{E}[Y] - \mathbb{E}[X]\mathbb{E}[Y] = 0$

So independent  $\Rightarrow$  0 correlation. But, it is easy enough to find examples showing that 0 correlation does NOT imply independence.

# Independence

Some important results & facts

4. **Corollary:** If  $X, Y$  are independent and  $EX^2, EY^2$  exist, then

$$\text{Var}(aX + bY) = \text{Var}(aX) + \text{Var}(bY) = a^2 \text{Var}(X) + b^2 \text{Var}(Y)$$

$$\text{Var}(aX + bY) = \text{Var}(aX) + \text{Var}(bY) + 2ab \underbrace{\text{Cov}(X, Y)}_{=0}$$

In general

$$\text{Var}\left(\sum_{i=1}^n a_i X_i\right) = \sum_{i=1}^n a_i^2 \text{Var}(X_i) \quad (\text{If } X_1, X_2, \dots, X_n \text{ are independent})$$

(\*) 5. **Corollary:** If  $X, Y$  are independent and  $X$  has mgf  $M_X(t)$  and  $Y$  has  $M_Y(t)$  then

$$\rightarrow M_{X+Y}(t) = M_X(t) M_Y(t)$$

holds for any  $t$  where  $M_X(t)$  and  $M_Y(t)$  exist.

(Note  $M_X(t)$  must exist in some  $(-h_1, h_1)$  and  $M_Y(t)$  must exist in some  $(-h_2, h_2)$ , so that both  $M_X(t)$  and  $M_Y(t)$  exist in some neighborhood of zero.)

$$M_{X+Y}(t) \stackrel{\text{def}}{=} \mathbb{E}[e^{t(X+Y)}] = \mathbb{E}[e^{tX} \cdot e^{tY}] \quad \begin{array}{l} X \text{ \& } Y \text{ are} \\ \text{independent} \end{array}$$

$$\mathbb{E}[e^{tX}] \cdot \mathbb{E}[e^{tY}] = M_X(t) M_Y(t).$$

$$M_{X_1+X_2+\dots+X_n}(t) = M_{X_1}(t) \cdot M_{X_2}(t) \cdots M_{X_n}(t) \quad (\text{If } X_1, X_2, \dots, X_n \text{ are independent})$$

6. **Theorem:** Suppose  $X$  has mgf  $M_X(t)$  and  $Y$  has  $M_Y(t)$ . Then,  $X, Y$  are independent iff

$$M_{X,Y}(t_1, t_2) \stackrel{\text{def}}{=} \mathbb{E}e^{t_1 X + t_2 Y} = \mathbb{E}e^{t_1 X} \mathbb{E}e^{t_2 Y} = M_X(t_1) M_Y(t_2) \quad \begin{array}{l} X \text{ and } Y \\ \text{are independent.} \end{array}$$

two arguments  $t_1, t_2$

holds for all  $t_1, t_2 \in (-h, h)$  in some neighborhood of zero.

$$M_{X,Y}(t_1, t_2) = \frac{1}{(1-t_1)(1-t_1-t_2)} \quad \begin{array}{l} t_1 < 1 \\ t_1+t_2 < 1 \end{array}$$

$$M_X(t_1) = M_{X,Y}(t_1, 0) = \frac{1}{(1-t_1)^2}$$

$$M_Y(t_2) = M_{X,Y}(0, t_2) = \frac{1}{(1-t_2)}$$

$$M_{X,Y}(t_1, t_2) = e^{\frac{t_1^2 + t_2^2}{2}} \quad M_X(t_1) = e^{t_1^2/2} \quad M_Y(t_2) = e^{t_2^2/2}$$

$X$  and  $Y$  are NOT independent.

# Independence

## Multivariate case

*Definition:* Random variables  $X_1, \dots, X_n$  are independent if

$$\longrightarrow P(X_1 \in A_1, \dots, X_n \in A_n) = \prod_{i=1}^n P(X_i \in A_i) \quad \text{for any } A_1, \dots, A_n \subset \mathbb{R}$$

### Characterizations

1.  $X_1, \dots, X_n$  are **independent** iff

$$F(x_1, \dots, x_n) = P(X_1 \leq x_1, \dots, X_n \leq x_n) = \prod_{i=1}^n F_{X_i}(x_i) \quad \text{for any } x_1, \dots, x_n \in \mathbb{R}$$

2.  $X_1, \dots, X_n$  are **independent** iff

$$f(x_1, \dots, x_n) = \prod_{i=1}^n f_{X_i}(x_i) \quad \text{for any } x_1, \dots, x_n \in \mathbb{R}$$

~~XX~~

*Important Properties:* If  $X_1, \dots, X_n$  are independent r.v.s, then

1.  $g_1(X_1), \dots, g_n(X_n)$  are independent r.v.s

2. it holds that  $E \left[ \prod_{i=1}^n g_i(X_i) \right] = \prod_{i=1}^n E[g_i(X_i)]$

3.  $\text{Cov}(X_i, X_j) = 0$  for  $i \neq j$

4.  $\text{Var} \left( \sum_{i=1}^n a_i X_i \right) = \sum_{i=1}^n a_i^2 \text{Var}(X_i)$

5. if the mgf  $M_{X_i}(t)$  exists for each  $i = 1, \dots, n$ , then the mgf of  $X_1 + \dots + X_n$  is

$$\underline{M_{X_1 + \dots + X_n}}(t) = \prod_{i=1}^n M_{X_i}(t)$$

for all  $t$  where each  $M_{X_i}(t)$  exists.

6. MGF characterization:  $X_1, \dots, X_n$  are independent iff

$$M_{X_1, \dots, X_n}(t_1, \dots, t_n) = \prod_{i=1}^n M_{X_i}(t_i)$$