

HW3

2024-09-28

HW 3

NAME: SAM OLSON

COLLABORATORS: **The Hatman**

Overview

- 1: Fix up $f(y)$ calculation
- 2: WIP
- 3: Part (c) left TO-DO
- 4: DONE
- 5:
- 6:
- 7:
- 8:

1. 2.23(b)

Question 1

Let X have the pdf

$$f(x) = \frac{1}{2}(1+x)$$


, $-1 < x < 1$

Define the random variable Y by $Y = X^2$

(b): Find $E(Y)$ and $\text{Var}(Y)$.

Answer 1

```
x <- seq(from = -1, to = 1, by = 0.01)
y <- 1/2 * (1+x)
plot(x, y)
```



(b):

(From prior HW) Note from the results of theorem 2.1.8:

$$f_Y(y) = \begin{cases} \sum_{i=1}^k f_X(g_i^{-1}(y)) \left| \frac{d}{dy} g_i^{-1}(y) \right| & y \in \mathbb{Y} \\ 0 & \text{otherwise} \end{cases}$$

Over the following partitions, we have monotonicity,

$A_1 = (-1, 0)$, $A_2 = (0, 1)$, with

$g_1(x) = x^2$ on their respective intervals.

Giving

$$f_Y(y) = \frac{1}{2\sqrt{y}}, \quad 0 < y < 1$$

Calculations:

$$E(Y) = \int_{y \in \mathbb{Y}} y f(y) dy = \int_{y=0}^1 y \left(\frac{1}{2\sqrt{y}} \right) dy$$

$$E(Y) = \int_{y=0}^1 \sqrt{y} \left(\frac{1}{2} \right) dy = \frac{1}{2} \frac{2}{3} y^{3/2} \Big|_{y=0}^{y=1} = \frac{1}{2} \frac{2}{3} (1) - 0 = \frac{1}{3}$$

To calculate $\text{Var}(Y)$, let us consider $E(Y^2)$,

$$E(Y^2) = \int_{y \in \mathbb{Y}} y^2 f(y) dy = \int_{y=0}^1 y^2 \left(\frac{1}{2\sqrt{y}}\right) dy$$

$$E(Y^2) = \int_{y=0}^1 y^{3/2} \frac{1}{2} dy = \frac{2}{5} \left(\frac{1}{2}\right) y^{5/2} \Big|_{y=0}^{y=1} = \frac{2}{5} \left(\frac{1}{2}\right) (1) - 0 = \frac{2}{10} = \frac{1}{5}$$

Taking $\text{Var}(Y) = E(Y^2) - (E(Y))^2$, then,

$$\text{Var}(Y) = \frac{1}{5} - \left(\frac{1}{3}\right)^2 = \frac{1}{5} - \frac{1}{9} = \frac{9}{45} - \frac{5}{45} = \frac{4}{45}$$

2.

Question 2

A family continues to have children until they have one female child. Suppose, for each birth, a single child is born and the child is equally likely to be male or female. The gender outcomes are independent across births.

(a): Let X be a random variable representing the number of children born to this family. Find the distribution of X .

(b): Find the expected value $E(X)$

(c): Let X_m denote the number of male children in this family and let X_f denote the number of female children. Find the expected value of X_m and the expected value of X_f

Answer 2

(a): We can frame X as the number of children until the family has their first (one) female child. So we can think of X as a Geometric distribution with probability $p = 0.5$ since it is equally likely that they have a male/female for each birth.

Notation-wise we write this as:

$$X \sim \text{Geometric}(p = 0.5)$$

(b):

Knowing the distribution of X , we know its pmf (discrete!) is given by:

For X number of children, $k = 1, 2, \dots$, we have:

$$f_X(x) = P(X = x) = p(1 - p)^{x-1}$$

$$E(X) = \sum_{x=1}^{\infty} xP(X = x) = \sum_{k=x}^{\infty} x(p(1 - p)^{x-1}) = p \sum_{x=1}^{\infty} x((1 - p)^{x-1})$$

Note, for the infinite geometric series we have, for $|r| < 1$, k some positive integer, the following holds:

$$\sum_{k=1}^{\infty} r^{k-1} = \frac{1}{1 - r}$$

Note: as $0 < p < 1 \rightarrow 0 < 1 - p < 1$, giving us:

$$E(X) = p \sum_{k=1}^{\infty} k((1 - p)^{k-1})$$

$$E(X) = X$$

(c):

3. 2.30 (a), (b), (c)

Question 3

Find the moment generating function corresponding to:

(a): $f(x) = \frac{1}{c}, 0 < x < c$

(b): $f(x) = \frac{2x}{c^2}, 0 < x < c$

(c): $f(x) = \frac{1}{2\beta} e^{\frac{-|x-\alpha|}{\beta}}, -\infty < x < \infty, -\infty < \alpha < \infty, \beta > 0$

Answer 3

Note, for a continuous random variable X, we may write the moment generating function as:

$$M_X(t) = \int_{-\infty}^{\infty} e^{tx} f_X(x) dx$$

Using this method, we then calculate the following:

(a):

$$M_X(t) = \int_{-\infty}^{\infty} e^{tx} f_X(x) dx = \int_0^c e^{tx} \frac{1}{c} dx = \frac{1}{ct} e^{tx} \Big|_{x=0}^{x=c} = \frac{1}{ct} e^{tc} - \frac{1}{ct} (1)$$

$$M_X(t) = \frac{1}{ct} e^{tc} - \frac{1}{ct} (1) = \frac{1}{ct} (e^{tc} - 1)$$

(b):

$$M_X(t) = \int_{-\infty}^{\infty} e^{tx} f_X(x) dx = \int_0^c e^{tx} \frac{2x}{c^2} dx = \frac{2}{c^2 t^2} e^{tx} (tx - 1) \Big|_{x=0}^{x=c}$$

$$M_X(t) = \frac{2}{c^2 t^2} e^{tc} (tc - 1) - \left(\frac{2}{c^2 t^2} 1(-1) \right) = \frac{2}{c^2 t^2} (tce^{tc} - e^{tc} + 1)$$

(c):

Question 4

Does a distribution exist for which $M_X(t) = \frac{t}{(1-t)}$, $|t| < 1$? If yes, find it. If no, prove it.

Answer 4

Let us suppose that the distribution exists.

Then by the definition of a(n) mfg:

$$M_X(t) = E(e^{tX})$$

We know for $t = 0$ that the relation $|t| = |0| = 0 < 1$ holds.

Thus we know the 0-th moment is defined, as:

$$M_X(0) = E(e^{0X}) = E(e^0) = E(1) = 1$$

However, if we evaluate $M_X(t)$ directly using the mgf as given, for $t = 0$ as given, we have:

$$M_X(t) = \frac{t}{(1-t)} = \frac{0}{1-0} = 0$$

And we arrive at a contradiction. Thus we must conclude that such a distribution does not exist.

5.

Question 5

Suppose that X has the standard normal distribution with pdf:

$$f(x) = \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}}$$

, $-\infty < x < \infty$

Then the random variable Y , $Y = e^X$ has a log-normal distribution.

(a): Find $E(Y^r)$ for any r .

(b): Show the moment generating function of Y does not exist (even though all moments of Y exist).

Answer 5

(a):

(b):

6.

Question 6

Suppose that X has a normal distribution with pdf:

$$f(x) = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$

, $-\infty < x < \infty$

The mean of X is μ . Show that the moment generating function of X satisfies $M_X(t) \geq e^{t\mu}$

Answer 6

7.

Question 7

Suppose that X has pmf $f(x) = p(1-p)^{x-1}$, for $x = 1, 2, 3, \dots$ where $0 < p < 1$. Find the mgf $M_X(t)$ and use this to derive the mean and variance of X .

Answer 7

8.

Question 8

Suppose for one month a company purchases c copies of a software package at a cost of d_1 dollars per copy. The packages are sold to customers for d_2 dollars per copy; any unsold copies are destroyed at the end of the month. Let X represent the demand for this software package in the month. Assume that X is a discrete random variable with pmf $f(x)$ and cdf $F(x)$.

(a): Let $s = \min\{X, c\}$ represent the number of sales during the month. Show that:

$$E(S) = \sum_{x=0}^c xf(x) + c(1 - F(c))$$

(b): Let $Y = S * d_2 - cd_1$ represent the profit for the company, the total income from sales minus the total cost of all copies. Find $E(Y)$

(c): As $Y \equiv Y_c$ depends on integer $c \geq 0$, write the expected profit function as $g(c) \equiv E(Y_c)$ from part (b). The company should choose the value of c which maximizes $g(c)$; that is, choose the smallest c such that $g(c + 1)$ is less than or equal to $g(c)$. Show that such $c \geq 0$ is the smallest integer with $F(c) \geq \frac{d_2 - d_1}{d_2}$

Answer 8

(a):

(b):

(c):