1.

Let A be an $n \times m$ matrix of rank r.

(a)

Show that $\text{rank}(I - AA^-) = n - r$.

Answer

(Preface: I in this problem is I_n; this is done to ensure matrix arithmetic is well-defined. Additionally, I have an inclination that Lemma 2.7 of the course notes *could* be used to prove the desired result (if one were to make a projection matrix argument); however, I am sticking with a more direct argument without note/use of projection matrices.)

To start, note that A^- is an $m \times n$ matrix. Let $P = AA^-$ (P is an $n \times n$ matrix).

The goal then is to determine the rank of $I_n - P$. And as given, we know $\text{rank}(P) = \text{rank}(AA^-) \leq \text{rank}(A) = r$.

The above inequality holds via Lemma 2.2, though this is a general linear algebra result. The goal then is to prove $\text{rank}(P) = r$.

We'll utilize a "squeeze" (or sandwich) argument, showing $r = \text{rank}(A) \leq \text{rank}(P)$ to establish equality. One way to do this is to show that $A = PA$, i.e., by showing that $A = PA \Rightarrow \text{rank}(A) = \text{rank}(PA) \leq \text{rank}(P)$ (again, utilizing Lemma 2.2 instead of directly jumping to Lemma 2.3).

To that end, note that A^- is a generalized inverse of A, meaning that, by definition $AA^-A = A$.

Then consider PA:

$$PA = AA^-A = A.$$

Thus, we've shown our desired outcome, namely $r = \text{rank}(A) = \text{rank}(PA) \leq \text{rank}(P)$, which, combined with our earlier result, shows that $\text{rank}(P) = r$.

Next, we show that P is idempotent (as it makes the rank difference calculation tractable via the Corollary to Lemma 2.5).

By definition, a matrix P is idempotent if $PP = P$. Evaluating this expression directly and simplifying:

$$PP = AA^-(AA^-) = (AA^-A)A^- = AA^- = P.$$

(Noting associativity of matrix multiplication *and* the definition of a generalized inverse for simplification.)

Thus, P is idempotent.

Now, consider $\text{rank}(I_n - P)$. The goal again is to show this is idempotent, as that allows us to calculate its rank via trace and apply the Corollary to Lemma 2.5 as well as Lemma 2.7.

To show that $I_n - P$ is also idempotent, we evaluate and simplify the following expression:

$$(I_n - P)^2 = (I_n - P)(I_n - P) = I_n - P - P + P^2 = I_n - 2P + P = I_n - P.$$

Thus, as desired, $I_n - P$ is idempotent.

Note: Via the corollary to Lemma 2.5, the trace of an idempotent matrix is equal to its rank. Applying this result to $I_n - P$, we obtain $\text{rank}(I_n - P) = \text{tr}(I_n - P)$.

Using the linearity of the trace, specifically additivity, we have

$$\text{tr}(I_n - P) = \text{tr}(I_n) - \text{tr}(P).$$

Since I_n is an $n \times n$ identity matrix, $\text{tr}(I_n) = n$. Furthermore, since P is idempotent, we know that

$$\text{tr}(P) = \text{rank}(P) = r.$$

Combining the two statements together, we conclude:

$$\text{rank}(I_n - P) = \text{tr}(I_n) - \text{tr}(P) = \text{rank}(I_n) - \text{rank}(P) = n - r.$$

Where $P = AA^-$.

(b)

Find the rank of $I - A^-A$.

Answer

(Preface: I in this problem is I_m; this is done to ensure matrix arithmetic is well-defined, similar to part (a). The other prefaced comments from part (a) similarly apply here.)

Let $Q = A^-A$ (Q is an $m \times m$ matrix). To start note that Q is idempotent, as:

$$QQ = A^-AA^-A = A^-(AA^-A) = A^-A = Q.$$

As in part (a), $\text{rank}(Q) = \text{rank}(A^-A) \leq \text{rank}(A) = r$ (via Lemma 2.2). To get the other side of the inequality (to conclude equality), note: $AQ = A(A^-A) = A$, such that $r = \text{rank}(A) = \text{rank}(AQ) \leq \text{rank}(Q)$.

Thus, via a "squeeze" argument we've shown that $\text{rank}(Q) = r$.

Next, we show that $I_m - Q$ is idempotent. We evaluate the expression directly and simplify:

$$(I_m - Q)^2 = I_m - 2Q + Q^2 = I_m - 2Q + Q = I_m - Q.$$

(Noting that Q is idempotent for simplifying the expression.)

By the Corollary to Lemma 2.5, the trace of an idempotent matrix is equal to its rank; and again, invoking known properties of the trace operation (additivity), we have: $\text{rank}(I_m - Q) = \text{tr}(I_m - Q) = \text{tr}(I_m) - \text{tr}(Q)$.

Evaluating this expression, we then conclude:

$$\text{rank}(I_m - Q) = \text{rank}(I_m) - \text{rank}(Q) \ = m - r.$$

Where $Q = A^-A$.

2.

Suppose that $\underline{y}$ is an $n \times 1$ observable random vector that follows the model

$$\underline{y} = X(\underline{\beta} + \underline{b}) + \underline{e},$$

where X is an $n \times p$ known matrix, and $\underline{\beta}$ is a $p \times 1$ vector of unknown parameters. Furthermore, suppose that $\underline{b}$ and $\underline{e}$ are independent random vectors with $\underline{b} \sim N(\underline{0}, D)$ and $\underline{e} \sim N(\underline{0}, \sigma^2 I)$, where D is an unknown non-negative definite matrix, and σ^2 is an unknown positive parameter.

(a)

Show that the simple least squares estimator (SLSE) of any estimable function $\underline{\lambda}'\underline{\beta}$ is the best linear unbiased estimator (BLUE) of $\underline{\lambda}'\underline{\beta}$.

Answer

Let $\underline{\lambda}'\underline{\beta}$ be an estimable function. By Theorem 2.7, $\underline{\lambda}'\underline{\beta}$ is estimable if and only if the conjugate normal equations are consistent. That is to say, we know there exists a solution $\underline{r} \in \mathbb{R}^p$ such that

$$X'X\underline{r} = \underline{\lambda}.$$

The SLSE of $\underline{\lambda}'\underline{\beta}$ is $\underline{\lambda}'\widehat{\underline{\beta}}$, where $\widehat{\underline{\beta}}$ is any solution to the ordinary normal equations $X'X\widehat{\underline{\beta}} = X'\underline{y}$.

Because $X'X\underline{r} = \underline{\lambda}$, noting transpose of matrices, $\underline{r}'X'X = \underline{\lambda}'$.

Taken together, this gives us:

$$\underline{\lambda}'\widehat{\underline{\beta}}_{\text{SLSE}} = \underline{\lambda}'\widehat{\underline{\beta}} = (\underline{r}'X'X)\widehat{\underline{\beta}} = \underline{r}'(X'X\widehat{\underline{\beta}}) = \underline{r}'X'\underline{y} = (X\underline{r})'\underline{y}.$$

The goal then is to establish that the SLSE is BLUE, i.e., that the SLSE is an unbiased estimator with the smallest variance among the class of linear unbiased estimators.

On unbiasedness: Since $\underline{b}$ and $\underline{e}$ are independent random vectors with mean zero, taking expectation of $\underline{y}$:

$$E(\underline{y}) = E(X(\underline{\beta} + \underline{b}) + \underline{e}) = E(X\underline{\beta}) + E(X\underline{b}) + E(\underline{e}) = X\underline{\beta},$$

noting linearity of expectation (and that $\underline{\beta}$ is non-random).

Then, taking expectation of the SLSE,

$$E(\underline{\lambda}'\widehat{\underline{\beta}}_{\text{SLSE}}) = E(\underline{r}'X'\underline{y}) = \underline{r}'X'E(\underline{y}) = \underline{r}'X'X\underline{r} = \underline{r}'\underline{\lambda} = \underline{\lambda}'\underline{\beta}.$$

So the SLSE is unbiased, as desired.

Next, on the variance-covariance matrix: Starting with $\underline{y}$:

$$\text{Var}(\underline{y}) = \text{Var}(X(\underline{\beta} + \underline{b}) + \underline{e}) = \text{Var}(X\underline{b} + \underline{e}) = XDX' + \sigma^2 I_n,$$

noting independence of $\underline{b}$ and $\underline{e}$.

The variance of the SLSE is then

$$\text{Var}(\underline{\lambda}'\widehat{\underline{\beta}}_{\text{SLSE}}) = \text{Var}(\underline{r}'X'\underline{y}) = \underline{r}'X'VX\underline{r} = \underline{r}'X'(XDX' + \sigma^2 I_n)X\underline{r} = \underline{\lambda}'D\underline{\lambda} + \sigma^2\underline{\lambda}'\underline{r}.$$

(Noting that $X'X\underline{r} = \underline{\lambda} \Rightarrow \underline{r}'X'X = \underline{\lambda}'$.)

But can any other unbiased estimator have lower variance compared to the SLSE? To answer that, consider an (other) arbitrary linear unbiased estimator $c'\underline{y}$ of $\underline{\lambda}'\underline{\beta}$. By Lemma 2.1, unbiasedness requires $c'X = \underline{\lambda}'$; and since $X'X\underline{r} = \underline{\lambda} \Rightarrow \underline{r}'X'X = \underline{\lambda}'$.

Let $\underline{d} = c - X\underline{r}$. Then

$$\underline{d}'X = c'X - \underline{r}'X'X = \underline{\lambda}' - \underline{\lambda}' = 0.$$

The difference in variances between these two estimators is then

$$\text{Var}(c'\underline{y}) - \text{Var}(\underline{r}'X'\underline{y}) = (X\underline{r} + \underline{d})'V(X\underline{r} + \underline{d}) - (X\underline{r})'V(X\underline{r}) = 2\underline{d}'VX\underline{r} + \underline{d}'V\underline{d}.$$

We consider the two quantities that make up $\text{Var}(c'\underline{y}) - \text{Var}(\underline{r}'X'\underline{y})$ in turn. To start, note that the term $2\underline{d}'VX\underline{r}$ is zero because

$$VX\underline{r} = (XDX' + \sigma^2 I_n)X\underline{r} = X(DX'X\underline{r} + \sigma^2\underline{r}) = X(D\underline{\lambda} + \sigma^2\underline{r}).$$

We may then write

$$2\underline{d}'VX\underline{r} = 2\underline{d}'\left[X(D\underline{\lambda} + \sigma^2\underline{r})\right] = 2(\underline{d}'X)(D\underline{\lambda} + \sigma^2\underline{r}).$$

and $\underline{d}'X = 0$, making the whole term zero.

The remaining term, $\underline{d}'V\underline{d}$, is non-negative as

$$\underline{d}'V\underline{d} = \underline{d}'XDX'\underline{d} + \sigma^2\underline{d}'\underline{d} = (X'\underline{d})'D(X'\underline{d}) + \sigma^2\underline{d}'\underline{d} = \sigma^2\underline{d}'\underline{d} \geq 0,$$

because $\underline{d}'X = 0 \Rightarrow X'\underline{d} = 0$.

Consequently,

$$\text{Var}(c'\underline{y}) - \text{Var}(\underline{r}'X'\underline{y}) \geq 0.$$

So any other linear unbiased estimator can do "no better" in terms of its variance, and we conclude that the SLSE of any estimable function $\underline{\lambda}'\underline{\beta}$ is the BLUE.

An Aside

I have a hunch that I may have done more work than is necessary. In reviewing the Summary Notes again, we can argue that the model in this problem is of the form of Model VII. Thus, if we can identify a suitable matrix Q such that $VX = XQ$, we have a clear path to the goal.

To that end, consider the variance-covariance matrix of $\underline{y}$ and call it V. We know $V = XDX' + \sigma^2 I_n$; then consider $Q = DX'X + \sigma^2 I_n$.

Then, we can verify that $VX = XQ$ holds, as:

$$VX = (XDX' + \sigma^2 I_n)X = X(DX'X + \sigma^2 I_n) = XQ.$$

Then we can invoke Corollary 1 to Theorem 2.21 and conclude that the SLSE is the BLUE for every estimable function $\underline{\lambda}'\underline{\beta}$ for the model specified in this problem.

(b)

Show that the variance of the BLUE of any estimable function $\underline{\lambda}'\underline{\beta}$ is $\sigma^2\underline{\lambda}'(X'X)^-\underline{\lambda} + \underline{\lambda}'D\underline{\lambda}$.

Answer

Using the conjugate normal equations noted in Theorem 2.7, let $\underline{r}$ be a solution to $X'X\underline{r} = \underline{\lambda}$.

Using the results from part (a), and taking as given that the SLSE is BLUE, we have

$$\begin{aligned}\mathrm{Var}(\underline{\lambda}'\widehat{\underline{\beta}}_{\mathrm{BLUE}}) = \mathrm{Var}(\underline{\lambda}'\widehat{\underline{\beta}}_{\mathrm{SLSE}}) &= \mathrm{Var}(\underline{r}'X'\underline{y}) \\ &= \underline{r}'X'VX\underline{r} \\ &= \underline{r}'X'(XDX' + \sigma^2 I_n)X\underline{r} \\ &= \sigma^2\underline{r}'X'X\underline{r} + \underline{r}'X'XDX'X\underline{r} \\ &= \sigma^2\underline{\lambda}'\underline{r} + \underline{\lambda}'D\underline{\lambda}.\end{aligned}$$

For an estimable $\underline{\lambda}'\underline{\beta}$, take $\underline{r} = (X'X)^-\underline{\lambda}$, where $(X'X)^-$ denotes *a* generalized inverse of $X'X$.

By the corollary to Theorem 2.7, $\underline{\lambda}'\underline{\beta}$ is estimable iff

$$X'X(X'X)^-\underline{\lambda} = \underline{\lambda},$$

which confirms that the conjugate normal equation ($X'X\underline{r} = \underline{\lambda}$) holds.

Because we are working with a generalized inverse, there are potentially (infinitely) many solutions for $\underline{r}$ (i.e., the choice of generalized inverse $(X'X)^-$). To address this (potential) issue, note that because $\underline{\lambda}'\underline{\beta}$ is estimable, Theorem 2.1 guarantees that $\underline{\lambda}' \in \mathcal{R}(X)$. Thus, there exists a vector $\underline{a}$ such that $\underline{\lambda}' = \underline{a}'X$.

Substituting this into the variance expression gives:

$$\sigma^2\underline{\lambda}'(X'X)^-\underline{\lambda} = \sigma^2\underline{a}'X(X'X)^-X'\underline{a} = \sigma^2\underline{a}'P_X\underline{a},$$

where $P_X = X(X'X)^-X'$.

Note: We know P_X is an orthogonal projection matrix with the properties of P_X following *after* Corollary 4 to Lemma 1.5 (i.e., the 5 noted properties, specifically property (3)). Furthermore, by property (1) of P_X, and with note of Corollary 2 to Lemma 1.5, P_X is invariant to the choice of the generalized inverse $(X'X)^-$.

In toto then, the expression for the variance of the BLUE of any estimable function $\underline{\lambda}'\underline{\beta}$ is given by:

$$\mathrm{Var}(\underline{\lambda}'\widehat{\underline{\beta}}_{\mathrm{BLUE}}) = \sigma^2\underline{\lambda}'(X'X)^-\underline{\lambda} + \underline{\lambda}'D\underline{\lambda}.$$

3.

Suppose that $\underline{y}$ is an $n \times 1$ observable random vector that follows the linear model

$$\underline{y} = X\underline{\beta} + \underline{\epsilon},$$

where $\underline{\beta}$ is a $p \times 1$ vector of unknown parameters, $\underline{\epsilon}$ is an unobservable random vector whose distribution is $N(\underline{0}, \sigma^2 I)$, and σ^2 is an unknown positive parameter. Further, let $X\underline{\beta} = X_1\underline{\beta}_1 + X_2\underline{\beta}_2$, where $X = (X_1, X_2)$, $\underline{\beta} = (\underline{\beta}_1', \underline{\beta}_2')'$, and $\underline{\beta}_1$ is a $q \times 1$ $(1 \leq q < p)$ vector. Suppose that the parameters $\underline{\beta}$ satisfy the constraints $\underline{\beta}_2 = \underline{b}$. That is, we now have Model IIa.

(a)

Let $\underline{\lambda}'\underline{\beta} = \underline{\lambda}_1'\underline{\beta}_1 + \underline{\lambda}_2'\underline{\beta}_2$ be a linear function. Show that $\underline{\lambda}'\underline{\beta}$ is estimable if and only if $\underline{\lambda}_1' \in \mathcal{R}(X_1)$, where $\mathcal{R}(X_1)$ denotes the row space of X_1.

Answer

To prove $\iff$, we prove each direction.

$(\Rightarrow)$ Suppose that $\underline{\lambda}'\underline{\beta}$ is estimable.

By the definition of estimability (for Models IIa, as noted in the Definition *under* Lemma 2.9), a linear function $\underline{\lambda}'\underline{\beta}$ is estimable if there exists a linear unbiased estimator $t(\underline{y}) = c + \underline{a}'\underline{y}$, such that $E(c + \underline{a}'\underline{y}) = \underline{\lambda}'\underline{\beta}$ for all $\underline{\beta}$ satisfying $A\underline{\beta} = \underline{b}$.

Note: In the above definition, c is a constant and $\underline{a}$ is an $n \times 1$ vector of constants.

Under the given constraint $\underline{\beta}_2 = \underline{b}$, and taking expectation, we then know

$$E[t(\underline{y})] = E[c + \underline{a}'\underline{y}] = c + E[\underline{a}'\underline{y}] = c + \underline{a}'X\underline{\beta} = c + \underline{a}'X_1\underline{\beta}_1 + \underline{a}'X_2\underline{\beta}_2 = c + \underline{a}'X_1\underline{\beta}_1 + \underline{a}'X_2\underline{b}.$$

Here the constraint may be written $A\underline{\beta} = \underline{b}$ with $A = (0, I_{p-q})$, such that $A\underline{\beta} = \underline{\beta}_2 = \underline{b}$.

By Lemma 2.9, because $t(\underline{y}) = c + \underline{a}'\underline{y}$ is an unbiased estimator of $\underline{\lambda}'\underline{\beta}$ under Model IIa, there exists a vector $\underline{k}$ such that

$$c = \underline{k}'\underline{b} \quad \text{and} \quad \underline{\lambda}' = \underline{a}'X + \underline{k}'A.$$

We may "split" (well, partition) $\underline{\lambda}'$ from the above expression into two components, and noting $X = (X_1, X_2)$ and $A = (0, I_{p-q})$):

$$\underline{\lambda}_1' = \underline{a}'X_1 \quad \text{and} \quad \underline{\lambda}_2' = \underline{a}'X_2 + \underline{k}'.$$

The first equation given above says that $\underline{\lambda}_1'$ is a linear combination of the rows of X_1, i.e., $\underline{\lambda}_1' \in \mathcal{R}(X_1)$, as desired.

$(\Leftarrow)$ Suppose that

$$\underline{\lambda}_1' \in \mathcal{R}(X_1).$$

By the definition of the row space of X_1 (i.e., all $1 \times q$ vectors spanned by the rows of X_1), there exists an $n \times 1$ vector $\underline{a}$ such that $\underline{a}'X_1 = \underline{\lambda}_1'$.

We may then construct a linear unbiased estimator using the above linear combination. Specifically, let

$$\underline{k}' = \underline{\lambda}_2' - \underline{a}' X_2, \quad \text{and let} \quad c = \underline{k}'\underline{b} = \underline{\lambda}_2'\underline{b} - \underline{a}' X_2 \underline{b}.$$

Taken together, the estimator we have constructed, $t(\underline{y}) = c + \underline{a}'\underline{y}$, is a linear estimator of $\underline{\lambda}'\underline{\beta}$. The goal then is to verify that this estimator is unbiased.

To that end, we take expectation

$$E[t(\underline{y})] = E[c + \underline{a}'\underline{y}] = c + E[\underline{a}'\underline{y}] = c + \underline{a}' X_1 \underline{\beta}_1 + \underline{a}' X_2 \underline{\beta}_2 = c + \underline{a}' X_1 \underline{\beta}_1 + \underline{a}' X_2 \underline{b}$$

Continuing simplification

$$E[t(\underline{y})] = \underline{\lambda}_2'\underline{b} - \underline{a}' X_2 \underline{b} + \underline{\lambda}_1'\underline{\beta}_1 + \underline{a}' X_2 \underline{b} = \underline{\lambda}_1'\underline{\beta}_1 + \underline{\lambda}_2'\underline{b} = \underline{\lambda}'\underline{\beta},$$

noting use of the given constraint $\underline{\beta}_2 = \underline{b}$.

As desired, the estimator $t(\underline{y})$ is unbiased, so we in fact constructed a linear unbiased estimator of $\underline{\lambda}'\underline{\beta}$. What's more, this function is estimable (with note of the Definition *under* Lemma 2.9 for Model IIa), as desired.

Then, combining the two directions, we may then conclude

$$\underline{\lambda}'\underline{\beta} \text{ is estimable } \iff \underline{\lambda}_1' \in \mathcal{R}(X_1).$$

(b)

Let $\underline{\lambda}'\underline{\beta} = \underline{\lambda}_1'\underline{\beta}_1 + \underline{\lambda}_2'\underline{\beta}_2$ be an estimable function. Show that the BLUE of $\underline{\lambda}'\underline{\beta}$ is given by

$$\underline{\lambda}_1'(X_1'X_1)^- X_1'(\underline{y} - X_2\underline{b}) + \underline{\lambda}_2'\underline{b}.$$

Answer

(Preface: This problem took the most time for me to figure out; that is why I start by rather explicitly noting where we're heading and why.)

The overall goal here is to show that the given expression is the BLUE of $\underline{\lambda}'\underline{\beta}$. By Theorem 2.18, the BLUE is $\underline{\lambda}'\widehat{\underline{\beta}}$, where $\widehat{\underline{\beta}}$ is the first component of a solution to the constrained normal equations for Model IIa. That is to say, it suffices to solve the constrained normal equations and then evaluate $\underline{\lambda}'\widehat{\underline{\beta}}$ to identify the BLUE.

As noted in part (a) the constraint in this problem may be written as $A\underline{\beta} = \underline{b}$ with $A = (0, I_{p-q})$, such that $A\underline{\beta} = \underline{\beta}_2 = \underline{b}$. The constrained normal equations are then (with note of the Definition *under* Theorem 2.11)

$$\begin{pmatrix} X'X & A' \\ A & 0 \end{pmatrix} \begin{pmatrix} \underline{\beta} \\ \underline{\rho} \end{pmatrix} = \begin{pmatrix} X'\underline{y} \\ \underline{b} \end{pmatrix}.$$

Our main focus then is the top of the above expression, $X'X\underline{\beta} + A'\underline{\rho} = X'\underline{y}$.

Using the given information, i.e., $X = (X_1, X_2)$, $\underline{\beta} = (\underline{\beta}_1', \underline{\beta}_2')'$, and $A = (0, I_{p-q})$, we evaluate each term in this equation.

To that end, note

$$X'X = \begin{pmatrix} X_1' \\ X_2' \end{pmatrix} \begin{pmatrix} X_1 & X_2 \end{pmatrix} = \begin{pmatrix} X_1'X_1 & X_1'X_2 \\ X_2'X_1 & X_2'X_2 \end{pmatrix}.$$

And the term $A'\underline{\rho}$ is equivalent to

$$A'\underline{\rho} = \begin{pmatrix} 0 \\ I_{p-q} \end{pmatrix} \underline{\rho} = \begin{pmatrix} \underline{0} \\ \underline{\rho} \end{pmatrix}.$$

And lastly, the term $X'\underline{y}$ is equivalent to

$$X'\underline{y} = \begin{pmatrix} X_1' \\ X_2' \end{pmatrix} \underline{y} = \begin{pmatrix} X_1'\underline{y} \\ X_2'\underline{y} \end{pmatrix}.$$

Via substitution of the above quantities then, the constrained normal equations from the Definition *under* Theorem 2.11 is equivalent to

$$\begin{pmatrix} X_1'X_1 & X_1'X_2 \\ X_2'X_1 & X_2'X_2 \end{pmatrix} \begin{pmatrix} \underline{\beta}_1 \\ \underline{\beta}_2 \end{pmatrix} + \begin{pmatrix} \underline{0} \\ \underline{\rho} \end{pmatrix} = \begin{pmatrix} X_1'\underline{y} \\ X_2'\underline{y} \end{pmatrix}.$$

Simplifying and evaluating this expression

$$\begin{pmatrix} X_1'X_1\underline{\beta}_1 + X_1'X_2\underline{\beta}_2 + \underline{0} \\ X_2'X_1\underline{\beta}_1 + X_2'X_2\underline{\beta}_2 + \underline{\rho} \end{pmatrix} = \begin{pmatrix} X_1'\underline{y} \\ X_2'\underline{y} \end{pmatrix}.$$

This may then be treated as a system of equations (specifically, the two equations)

$$X_1'X_1\underline{\beta}_1 + X_1'X_2\underline{\beta}_2 = X_1'\underline{y}, \qquad X_2'X_1\underline{\beta}_1 + X_2'X_2\underline{\beta}_2 + \underline{\rho} = X_2'\underline{y}.$$

Using the constraint $\underline{\beta}_2 = \underline{b}$ and gathering like terms (X_1'), the first of these equations may be written

$$X_1'X_1\underline{\beta}_1 = X_1'(\underline{y} - X_2\underline{b}).$$

Let $\underline{y}^* = \underline{y} - X_2\underline{b}$, then the above equation is equivalent to $X_1'X_1\underline{\beta}_1 = X_1'\underline{y}^*$. However, this is exactly the form of the ordinary normal equations for a Model II with design matrix X_1 and response $\underline{y}^*$!

Looking ahead at the desired expression for the BLUE, we need $\widehat{\underline{\beta}}_1 = (X_1'X_1)^- X_1'\underline{y}^*$.

To motivate such a choice, note that by Lemma 1.4,

$$X_1'X_1(X_1'X_1)^- X_1' = X_1',$$

so this $\widehat{\underline{\beta}}_1$ indeed solves $X_1'X_1\underline{\beta}_1 = X_1'\underline{y}^*$. However, we need to ensure that this solution also satisfies the second equation; to that end, set

$$\widehat{\underline{\rho}} = X_2'\underline{y} - X_2'X_1\widehat{\underline{\beta}}_1 - X_2'X_2\underline{b}, \qquad \widehat{\underline{\beta}} = \begin{pmatrix} \widehat{\underline{\beta}}_1 \\ \underline{b} \end{pmatrix}.$$

By construction, the vector $\begin{pmatrix} \widehat{\underline{\beta}} \\ \widehat{\underline{\rho}} \end{pmatrix}$ solves the constrained normal equations. Therefore, by Theorem 2.18, the BLUE of $\underline{\lambda}'\underline{\beta}$ is

$$\underline{\lambda}'\widehat{\underline{\beta}}_{\text{BLUE}} = \underline{\lambda}'\widehat{\underline{\beta}} = \underline{\lambda}_1'\widehat{\underline{\beta}}_1 + \underline{\lambda}_2'\underline{b} = \underline{\lambda}_1'(X_1'X_1)^- X_1'(\underline{y} - X_2\underline{b}) + \underline{\lambda}_2'\underline{b}.$$

While the above expression is our desired goal, there is one (potential) issue left to address, namely that the above expression does not depend on the choice of the generalized inverse of $(X_1'X_1)$. To prove this

statement, note that, with reference to part (a), because $\underline{\lambda}_1' \in \mathcal{R}(X_1)$, there exists a vector $\underline{a}$ such that $\underline{\lambda}_1' = \underline{a}' X_1$. Then

$$\underline{\lambda}_1'(X_1'X_1)^- X_1' = \underline{a}' X_1 (X_1'X_1)^- X_1' = \underline{a}' P_{X_1},$$

where $P_{X_1} = X_1(X_1'X_1)^- X_1'$ is an orthogonal projection matrix (similar the argument made in question 2 part (b)). That is, by Corollary 2 to Lemma 1.5, P_{X_1} is invariant to the choice of generalized inverse $(X_1'X_1)^-$.

4.

Consider the following linear model

$$y_i = \mu + \alpha x_i + \gamma z_i + \epsilon_i, \quad i = 1, \ldots, n, \tag{1}$$

where μ, α, and γ are unknown fixed parameters; the y_i's are observable random variables; the x_i's and z_i's are known constants; the ϵ_i's are unobservable and uncorrelated random variables with mean 0 and variance σ^2, where σ^2 is an unknown positive parameter. Further, let $\underline{1} = (1, \ldots, 1)'$ be an $n \times 1$ vector of 1's, $\underline{y} = (y_1, \ldots, y_n)'$, $\underline{x} = (x_1, \ldots, x_n)'$, and $\underline{z} = (z_1, \ldots, z_n)'$. Suppose also that $\underline{1}'\underline{x} = \underline{1}'\underline{z} = \underline{x}'\underline{z} = 0$ and $\underline{x}'\underline{x} = \underline{z}'\underline{z} = n$.

(a)

Formulate model (1) as a Model II: $\underline{y} = X\underline{\beta} + \underline{\epsilon}$. Carefully write out X, $\underline{\beta}$, and $\underline{\epsilon}$.

Answer

For reference, when we say Model II, we are referring to a Gauss-Markov model:

$$\underline{y} = X\underline{\beta} + \underline{\epsilon}$$

where $E(\underline{\epsilon}) = \underline{0}$, $\text{Var}(\underline{\epsilon}) = \sigma^2 I_n$ with parameter space $\{\underline{\beta}, \sigma^2 : \underline{\beta} \in \mathbb{R}^p, \sigma^2 > 0\}$.

Using this notation for model (1), as formulated in the problem statement, we may write:

$$\underline{y} = X\underline{\beta} + \underline{\epsilon},$$

where

$$E(\underline{\epsilon}) = \underline{0}, \qquad \text{Var}(\underline{\epsilon}) = \sigma^2 I_n.$$

And where

$$X = \begin{pmatrix} 1 & x_1 & z_1 \\ 1 & x_2 & z_2 \\ \vdots & \vdots & \vdots \\ 1 & x_n & z_n \end{pmatrix}, \qquad \underline{\beta} = \begin{pmatrix} \mu \\ \alpha \\ \gamma \end{pmatrix}, \qquad \underline{\epsilon} = \begin{pmatrix} \epsilon_1 \\ \vdots \\ \epsilon_n \end{pmatrix}.$$

Note: X is an $n \times 3$ (design) matrix, $\underline{\beta}$ is a 3×1 vector, and $\underline{\epsilon}$ is an $n \times 1$ vector

(b)

Calculate $X'X$ and use this to argue that $n \geq 3$.

Answer

Note: X is an $n \times 3$ matrix, meaning X' is an $3 \times n$ matrix, such that $X'X$ is a 3×3 matrix.

X as specified in part (a) may equivalently be represented as:

$$X = \begin{pmatrix} 1 & x_1 & z_1 \\ 1 & x_2 & z_2 \\ \vdots & \vdots & \vdots \\ 1 & x_n & z_n \end{pmatrix} = \begin{pmatrix} \underline{1} & \underline{x} & \underline{z} \end{pmatrix}$$

Then, $X'X$ is given by:

$$X'X = \begin{pmatrix} \underline{1} & \underline{x} & \underline{z} \end{pmatrix}' \begin{pmatrix} \underline{1} & \underline{x} & \underline{z} \end{pmatrix}$$

Calculating:

$$X'X = \begin{pmatrix} \underline{1}'\underline{1} & \underline{1}'\underline{x} & \underline{1}'\underline{z} \\ \underline{x}'\underline{1} & \underline{x}'\underline{x} & \underline{x}'\underline{z} \\ \underline{z}'\underline{1} & \underline{z}'\underline{x} & \underline{z}'\underline{z} \end{pmatrix} = \begin{pmatrix} n & 0 & 0 \\ 0 & n & 0 \\ 0 & 0 & n \end{pmatrix} = n \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} = nI_3.$$

Noting $\underline{1}'\underline{x} = \underline{1}'\underline{z} = \underline{x}'\underline{z} = 0$ and $\underline{x}'\underline{x} = \underline{z}'\underline{z} = n$ as given, and $\underline{1}'\underline{1} = \sum_{i=1}^{n} 1 \cdot 1 = n$.

We know then that $X'X = nI_3$ is a nonsingular matrix of dimension 3, such that $\text{rank}(X'X) = 3$ (assuming $n > 0$ to avoid the trivial case).

Noting Lemma 2.3, $\text{rank}(X'X) = \text{rank}(X) = 3$.

As X is an $n \times 3$ matrix, by the definition of the rank of a matrix (i.e., the rank of a matrix is the maximum number of linearly independent row vectors or column vectors in the matrix), we have:

$$3 = \text{rank}(X) \leq n.$$

This forces $n \geq 3$.

Additional Note

While I believe the above argument is correct (and hopefully clear), I wanted to make one part of the argument airtight. Specifically, for $0 < n < 3$, we effectively have fewer than 3 rows (observations). In that case, the individual coefficients cannot all be uniquely identified, i.e., only certain linear combinations of the coefficients may be estimable. While I'd say this intuitively makes sense, such a statement does not get at the *why*, and the why is this: If $0 < n < 3$, then $\text{rank}(X) \leq n < 3$, such that X cannot have full column rank. Consequently for model (1) as specified, the three unknown fixed parameters cannot *all* be estimable, meaning that $n \geq 3$ is a necessary condition for the separate estimation of all three parameters (with a tip of the cap to Theorem 2.2).

(c)

Show that μ, α, and γ are estimable functions.

Answer

From part (b), we know that $X'X = nI_3$ is nonsingular, meaning X has full column rank ($\text{rank}(X) = 3$).

Under the definition of estimability, a linear function $\underline{\lambda}'\underline{\beta}$ is said to be estimable if there exists a linear unbiased estimator of $\underline{\lambda}'\underline{\beta}$.

Invoking Theorem 2.1 (which is appropriate here given our work in specifying model (1) as Model II), a necessary and sufficient condition for estimability is that we can express the coefficient vector $\underline{\lambda}'$ in the form $\underline{\lambda}' = \underline{a}'X$, and that $\underline{\lambda}'$ belong to the rowspace of X.

To that end, note that

$$\mathcal{R}(X') = \mathbb{R}^3.$$

That is to say, a linear function $\underline{\lambda}'\underline{\beta}$ is estimable $\iff \underline{\lambda}' \in \mathcal{R}(X) \iff \underline{\lambda} \in \mathcal{C}(X')$.

As model (1) is a full-rank linear model, with note of Corollary 3, since $\mathcal{C}(X') = \mathbb{R}^3$, every linear function $\underline{\lambda}'\underline{\beta}$ is estimable. Specifically, taking

$$\underline{\lambda}_1 = \begin{pmatrix}1\\0\\0\end{pmatrix}, \quad \underline{\lambda}_2 = \begin{pmatrix}0\\1\\0\end{pmatrix}, \quad \underline{\lambda}_3 = \begin{pmatrix}0\\0\\1\end{pmatrix}, \quad \text{gives the three estimable functions} \quad \underline{\lambda}_1'\underline{\beta} = \mu, \quad \underline{\lambda}_2'\underline{\beta} = \alpha, \quad \underline{\lambda}_3'\underline{\beta} = \gamma.$$

And we conclude that μ, α, and γ are estimable functions.

(d)

Find the best linear unbiased estimators (BLUE's) of μ, α, and γ in non-matrix, explicit expressions.

Answer

A necessary condition (noted in Section 2.3 of the Summary Notes) is that our estimator $\widehat{\underline{\beta}}$ must satisfy the normal equations, which are given by:

$$X'X\widehat{\underline{\beta}} = X'\underline{y}.$$

From our work in the prior problems, we've calculated $X'X = nI_3$ and know it is nonsingular (such that we know its inverse exists, and we can take that inverse rather than calculating/using generalized inverses for computation). Specifically, by multiplying both sides of the normal equations by $(X'X)^{-1}$,

$$\begin{pmatrix}\widehat{\mu}\\\widehat{\alpha}\\\widehat{\gamma}\end{pmatrix} = \widehat{\underline{\beta}} = (X'X)^{-1}X'X\widehat{\underline{\beta}} = (X'X)^{-1}X'\underline{y} = \frac{1}{n}X'\underline{y} = \frac{1}{n}\begin{pmatrix}\underline{1}'\underline{y}\\\underline{x}'\underline{y}\\\underline{z}'\underline{y}\end{pmatrix}.$$

Where $(X'X)^{-1} = \frac{1}{n}I_3$.

Writing these terms component-wise, i.e., as non-matrix, explicit expressions:

$$\widehat{\mu} = \frac{1}{n}\underline{1}'\underline{y} = \frac{1}{n}\sum_{i=1}^{n} y_i, \qquad \widehat{\alpha} = \frac{1}{n}\underline{x}'\underline{y} = \frac{1}{n}\sum_{i=1}^{n} x_i y_i, \qquad \widehat{\gamma} = \frac{1}{n}\underline{z}'\underline{y} = \frac{1}{n}\sum_{i=1}^{n} z_i y_i.$$

And by the Gauss–Markov theorem (Theorem 2.9), these least squares estimators are BLUEs of μ, α, and γ respectively.

5.

Consider model (1) with the constraint $\gamma = \mu + \alpha$. That is, we now have a constrained Model IIa. Let $\underline{\beta} = \begin{pmatrix} \mu \\ \alpha \\ \gamma \end{pmatrix}$ and $\underline{\lambda}' = (\lambda_1, \lambda_2, \lambda_3)$.

(a)

Find all estimable functions $\underline{\lambda}'\underline{\beta}$.

Answer

From question 4, we have the original parameter vector and design matrix given by

$$\underline{\beta} = \begin{pmatrix} \mu \\ \alpha \\ \gamma \end{pmatrix}, \qquad \text{and } X = \begin{pmatrix} 1 & x_1 & z_1 \\ 1 & x_2 & z_2 \\ \vdots & \vdots & \vdots \\ 1 & x_n & z_n \end{pmatrix} = \begin{pmatrix} \underline{1} & \underline{x} & \underline{z} \end{pmatrix}.$$

Note: We can write the constraint $\gamma = \mu + \alpha$ as $A\underline{\beta} = \underline{0}$, where

$$A = \begin{pmatrix} 1 & 1 & -1 \end{pmatrix}, \quad \text{as} \quad \mu + \alpha - \gamma = 0 \Rightarrow \mu + \alpha = \gamma$$

Under this constraint, every linear function $\underline{\lambda}'\underline{\beta}$ can be written as

$$\underline{\lambda}'\underline{\beta} = \lambda_1\mu + \lambda_2\alpha + \lambda_3\gamma = \lambda_1\mu + \lambda_2\alpha + \lambda_3(\mu + \alpha) = (\lambda_1 + \lambda_3)\mu + (\lambda_2 + \lambda_3)\alpha.$$

That is to say, we can write every linear function $\underline{\lambda}'\underline{\beta}$ as a function of μ and α only, effectively reducing $\underline{\lambda}'\underline{\beta}$ to $\tilde{\underline{\lambda}}'\tilde{\underline{\beta}}$ with $\tilde{\underline{\beta}} = (\mu, \alpha)'$.

The question then becomes whether $\tilde{\underline{\lambda}}'\tilde{\underline{\beta}}$ is estimable in the model for $\underline{y}$ with parameter $\tilde{\underline{\beta}}$. Estimability here is determined by the (design) matrix W satisfying $E(\underline{y}) = W\tilde{\underline{\beta}}$. To identify W, we consider the constrained mean model given by

$$E(\underline{y}) = \mu\underline{1} + \alpha\underline{x} + \gamma\underline{z} = \mu\underline{1} + \alpha\underline{x} + (\mu + \alpha)\underline{z} = \mu(\underline{1} + \underline{z}) + \alpha(\underline{x} + \underline{z}).$$

To that end, let

$$\tilde{\beta} = \begin{pmatrix} \mu \\ \alpha \end{pmatrix}, \qquad W = \begin{pmatrix} 1 + z_1 & x_1 + z_1 \\ 1 + z_2 & x_2 + z_2 \\ \vdots & \vdots \\ 1 + z_n & x_n + z_n \end{pmatrix}.$$

And for shorthand convenience, similar to question 4, we may write

$$W = \begin{pmatrix} 1 + z_1 & x_1 + z_1 \\ 1 + z_2 & x_2 + z_2 \\ \vdots & \vdots \\ 1 + z_n & x_n + z_n \end{pmatrix} = \begin{pmatrix} \underline{1} + \underline{z} & \underline{x} + \underline{z} \end{pmatrix}.$$

At any rate, the constrained model can be written as

$$\underline{y} = W\tilde{\beta} + \underline{\epsilon}.$$

Note: W is an $n \times 2$ matrix, W' is a $2 \times n$ matrix, and $W'W$ is a 2×2 matrix. Our goal then is to compute $W'W$ and calculate its rank; the "idea" here being that we can easily calculate the rank of a nonsingular matrix, and know that $\text{rank}(W'W) = \text{rank}(W)$. To that end then, we compute $W'W$:

$$W'W = \begin{pmatrix} \underline{1} + \underline{z} & \underline{x} + \underline{z} \end{pmatrix}' \begin{pmatrix} \underline{1} + \underline{z} & \underline{x} + \underline{z} \end{pmatrix} = \begin{pmatrix} (\underline{1} + \underline{z})'(\underline{1} + \underline{z}) & (\underline{1} + \underline{z})'(\underline{x} + \underline{z}) \\ (\underline{x} + \underline{z})'(\underline{1} + \underline{z}) & (\underline{x} + \underline{z})'(\underline{x} + \underline{z}) \end{pmatrix}$$

Continuing simplification of this expression

$$W'W = \begin{pmatrix} \underline{1}'\underline{1} + \underline{1}'\underline{z} + \underline{z}'\underline{1} + \underline{z}'\underline{z} & \underline{1}'\underline{x} + \underline{1}'\underline{z} + \underline{z}'\underline{x} + \underline{z}'\underline{z} \\ \underline{x}'\underline{1} + \underline{x}'\underline{z} + \underline{z}'\underline{1} + \underline{z}'\underline{z} & \underline{x}'\underline{x} + \underline{x}'\underline{z} + \underline{z}'\underline{x} + \underline{z}'\underline{z} \end{pmatrix} = n \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}.$$

Noting the given information from question 4, namely $\underline{1}'\underline{x} = \underline{1}'\underline{z} = \underline{x}'\underline{z} = 0$, $\underline{x}'\underline{x} = \underline{z}'\underline{z} = n$, and $\underline{1}'\underline{1} = \sum_{i=1}^{n} 1 \cdot 1 = n$.

Note: For a standard 2x2 matrix,

$$kA = k \begin{pmatrix} a & b \\ c & d \end{pmatrix}, \qquad \det(A) = k^2(ad - bc).$$

Applying this to the above matrix, and noting $n > 0$, we have $\det(W'W) = (2n)(2n) - n^2 = 3n^2 > 0$, meaning that, $W'W$ is nonsingular (and consequently W has full column rank).

Then, by Lemma 2.3, $2 = \text{rank}(W'W) = \text{rank}(W)$, and $\text{rank}(W) = 2$.

Since W has full column rank, Corollary 3 to Theorem 2.1 gives that $\tilde{\underline{\lambda}}'\tilde{\underline{\beta}}$ is estimable for every $\tilde{\underline{\lambda}} \in \mathbb{R}^2$. Now for any $\underline{\lambda} \in \mathbb{R}^3$, set

$$\tilde{\underline{\lambda}} = \begin{pmatrix} \lambda_1 + \lambda_3 \\ \lambda_2 + \lambda_3 \end{pmatrix}.$$

Then $\underline{\lambda}'\underline{\beta} = \tilde{\underline{\lambda}}'\tilde{\underline{\beta}}$ on the constrained space, and since $\tilde{\underline{\lambda}}'\tilde{\underline{\beta}}$ is estimable, $\underline{\lambda}'\underline{\beta}$ is also estimable. And because $\underline{\lambda} \in \mathbb{R}^3$ was arbitrary, we may then conclude that all linear functions $\underline{\lambda}'\underline{\beta}$ are estimable under $\gamma = \mu + \alpha$.

(b)

Give the constrained conjugate normal equations, in an explicit matrix form, for an estimable $\underline{\lambda}'\underline{\beta}$.

Answer

In general, with note of Lemma 2.14, the constrained conjugate normal equations are given by

$$\begin{pmatrix} X'X & A' \\ A & 0 \end{pmatrix} \begin{pmatrix} \underline{r}_1 \\ \underline{r}_2 \end{pmatrix} = \begin{pmatrix} \underline{\lambda} \\ \underline{0} \end{pmatrix}.$$

Because we are considering an estimable function $\underline{\lambda}'\underline{\beta}$, Theorem 2.16 guarantees that this system is consistent (i.e., a solution for $\underline{r}_1$ and $\underline{r}_2$ exists).

Note: The congujate normal equations may equivalently be written as

$$X'X\underline{r}_1 + A'\underline{r}_2 = \underline{\lambda} \text{ and } A\underline{r}_1 = \underline{0}$$

From part (a), we may write $A = \begin{pmatrix} 1 & 1 & -1 \end{pmatrix}$ and $A' = \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix}$, and from question 4, $X'X = nI_3$.

We need then to write out the matrix forms of

$$\begin{pmatrix} X'X & A' \\ A & 0 \end{pmatrix}, \qquad \begin{pmatrix} \underline{r}_1 \\ \underline{r}_2 \end{pmatrix}, \qquad \text{and } \begin{pmatrix} \underline{\lambda} \\ \underline{0} \end{pmatrix}.$$

To that end, note that $X'X$ is a 3×3 matrix, A' is a 3×1 matrix, A is a 1×3 matrix, and the lower-right 0 is 1×1. Combining together, we know that

$$\begin{pmatrix} X'X & A' \\ A & 0 \end{pmatrix}$$

is a 4×4 matrix; similarly, $\underline{\lambda}$ is a 3×1 vector, $\underline{r}_1$ is a 3×1 vector, and $\underline{r}_2$ (which we will henceforth refer to as ρ) is 1×1.

Note: While not ideal notationally, I want to call out that r_2 and $\underline{r}_2$ are two different quantities here.

"Plugging in" known quantities, we then have:

$$\begin{pmatrix} n & 0 & 0 & 1 \\ 0 & n & 0 & 1 \\ 0 & 0 & n & -1 \\ 1 & 1 & -1 & 0 \end{pmatrix} \begin{pmatrix} r_1 \\ r_2 \\ r_3 \\ \rho \end{pmatrix} = \begin{pmatrix} \lambda_1 \\ \lambda_2 \\ \lambda_3 \\ 0 \end{pmatrix}.$$

(c)

Give the BLUE of any an estimable function $\underline{\lambda}'\underline{\beta}$ in a non-matrix, explicit expression.

Answer

From part (b), the constrained conjugate normal equations are

$$\begin{pmatrix} n & 0 & 0 & 1 \\ 0 & n & 0 & 1 \\ 0 & 0 & n & -1 \\ 1 & 1 & -1 & 0 \end{pmatrix} \begin{pmatrix} r_1 \\ r_2 \\ r_3 \\ \rho \end{pmatrix} = \begin{pmatrix} \lambda_1 \\ \lambda_2 \\ \lambda_3 \\ 0 \end{pmatrix}.$$

Writing this system out component-wise, we have

$$\begin{aligned} nr_1 + \rho &= \lambda_1, \\ nr_2 + \rho &= \lambda_2, \\ nr_3 - \rho &= \lambda_3, \\ r_1 + r_2 - r_3 &= 0. \end{aligned}$$

As a system of linear equations, we can solve for ρ:

$$\lambda_1 + \lambda_2 - \lambda_3 = (nr_1 + \rho) + (nr_2 + \rho) - (nr_3 - \rho) = 3\rho + n(r_1 + r_2 - r_3) = 3\rho.$$

Thus,

$$\rho = \frac{\lambda_1 + \lambda_2 - \lambda_3}{3}.$$

"Plugging this in" to then solve for r_1, r_2, r_3, we have

$$\begin{aligned} r_1 &= \frac{\lambda_1 - \rho}{n} = \frac{2\lambda_1 - \lambda_2 + \lambda_3}{3n}, \\ r_2 &= \frac{\lambda_2 - \rho}{n} = \frac{-\lambda_1 + 2\lambda_2 + \lambda_3}{3n}, \\ r_3 &= \frac{\lambda_3 + \rho}{n} = \frac{\lambda_1 + \lambda_2 + 2\lambda_3}{3n}. \end{aligned}$$

By Theorem 2.17, the least-squares estimate of an estimable function (under the constrained model) is given by

$$\underline{r}_1' X' \underline{y} + \underline{r}_2' \underline{b},$$

where $\begin{pmatrix} \underline{r}_1 \\ \underline{r}_2 \end{pmatrix}$ is any solution to the constrained conjugate normal equations.

Note: With regards to notation, $\underline{r}_2' = \rho$. Additionally note the original model constraint is $\gamma = \mu + \alpha \Rightarrow \mu + \alpha - \gamma = 0$. Thus, $A\underline{\beta} = \underline{b} \Rightarrow (1, 1, -1) \begin{pmatrix} \mu \\ \alpha \\ \gamma \end{pmatrix} = \mu + \alpha - \gamma = \underline{0}$, meaning $\underline{b} = \underline{0}$, such that $\underline{r}_2' \underline{b}$ is also zero.

Also, by Theorem 2.18, this least-squares estimator is the BLUE under Model IIa. Now we just need to provide an (non-matrix) explicit expression.

From question 4, $X = (\underline{1}, \underline{x}, \underline{z})$, so we need only compute and simplify using known quantities. Specifically

$$\underline{\lambda}' \underline{\widehat{\beta}}_{\text{BLUE}} = \underline{r}_1' X' \underline{y} = r_1 \underline{1}' \underline{y} + r_2 \underline{x}' \underline{y} + r_3 \underline{z}' \underline{y}.$$

Expanding

$$\underline{\lambda}' \underline{\widehat{\beta}}_{\text{BLUE}} = \left(\frac{2\lambda_1 - \lambda_2 + \lambda_3}{3n} \right) \sum_{i=1}^{n} y_i + \left(\frac{-\lambda_1 + 2\lambda_2 + \lambda_3}{3n} \right) \sum_{i=1}^{n} x_i y_i + \left(\frac{\lambda_1 + \lambda_2 + 2\lambda_3}{3n} \right) \sum_{i=1}^{n} z_i y_i.$$

Note: This is the BLUE of *any* estimable function $\underline{\lambda}' \underline{\beta}$, since, by part (a), every $\underline{\lambda}' \underline{\beta}$ is estimable under the constraint.