

1. Solution:

- (i) $\forall \omega \in (0, 1), X_n(\omega) = 0$ for large n since $\frac{1}{n^2} < \omega$ for $n \geq \frac{1}{\sqrt{\omega}}$ if $\omega > 0$. Also we have $X_n(0) = X_n(1) = 0, \forall n \geq 1$. So $X_n(\omega) \rightarrow X(\omega), \forall \omega \in [0, 1]$, i.e. $X_n \rightarrow X$ a.e. (P) .
- (ii) $m(\{\omega : |X_n(\omega) - X(\omega)| > \epsilon\}) = m((0, \frac{1}{n^2})) = \frac{1}{n^2} \rightarrow 0$ as $n \rightarrow \infty, \forall \epsilon > 0$. So $X_n \xrightarrow{p} X$.
- (iii) $(E|X_n - X|^p) = (\int |X_n - X|^p dP) = (\sqrt{n})^p \cdot m((0, \frac{1}{n^2})) = n^{p/2-2} \rightarrow 0$ as $n \rightarrow \infty$ iff $0 < p < 4$. Hence $X_n \rightarrow X$ in L^p for $0 < p < 4$, but this convergence doesn't hold for $4 \leq p < \infty$.

2. Solution:

(a) Note that

$$\begin{aligned} \text{Var}(X + Y) &= E([(X - EX) + (Y - EY)]^2) \\ &= \text{Var}(X) + \text{Var}(Y) + 2E[(X - EX)(Y - EY)] \\ &= \text{Var}(X) + \text{Var}(Y) + 2[E(XY) - EXEY] \end{aligned}$$

Now $E|XY| \leq \sqrt{EX^2EY^2} < \infty$ by Cauchy-Schwarz inequality. So $XY \in L^1(\Omega, \mathcal{F}, P)$. Then, by using Fubini theorem,

$$\begin{aligned} E(XY) &= \int_{\Omega} X(\omega)Y(\omega)dP(\omega) \\ &= \int_{\mathbb{R}^2} xy \, d(P_X \times P_Y)(x, y) \\ &= \int_{\mathbb{R}} x \, dP_X(x) \int_{\mathbb{R}} y \, dP_Y(y) \\ &= \int_{\Omega} X(\omega) \, dP_X(\omega) \int_{\Omega} Y(\omega) \, dP_Y(\omega) \\ &= EXEY \end{aligned}$$

Hence $\text{Var}(X + Y) = \text{Var}(X) + \text{Var}(Y)$.

- (b) Let $\phi(x) = \sqrt{1 + x^2}$, then $\phi(x)$ is convex, so $\sqrt{1 + (EX)^2} = \phi(EX) \leq E\phi(X) = E\sqrt{1 + X^2}$ given that $EX < \infty$. When $EX = \infty$, this is automatically true. As $1 + X^2 \leq (1 + |X|)^2$, so $E\sqrt{1 + X^2} \leq E(1 + |X|) = 1 + E|X|$.
- (c) From $\frac{dP_X}{dm} = f, \frac{dP_Y}{dm} = g$, we have

$$\begin{aligned} P_{(X,Y)}(A_1 \times A_2) &= (P_X \times P_Y)(A_1 \times A_2) \text{ (by definition of product measures)} \\ &= P_X(A_1)P_Y(A_2) \text{ (by definition of RN derivative)} \\ &= \left(\int I_{A_1} f dm \right) \left(\int I_{A_2} g dm \right) \text{ (by Tonelli's theorem since } f, g \geq 0) \\ &= \int I_{A_1}(x)I_{A_2}(y)f(x)g(y) \, dm(x)dm(y) \\ &= \int_{A_1 \times A_2} f(x)g(y) \, dm^2(x, y) \end{aligned}$$

Since on rectangles $A_1 \times A_2$ on $\mathbb{R}^2$,

$$P_{(X,Y)}(A_1 \times A_2) = \int_{A_1 \times A_2} f(x)g(y) \, dm^2(x, y),$$

By uniqueness of Caratheodory extension theorem, we have

$$P_{(X,Y)}(A) = \int_A f g \, dm^2, \quad \forall A \in \mathcal{B}(\mathbb{R}^2)$$

Note: another argument is the following

$$\begin{aligned} (P_X \times P_Y)(A) &= \int P_Y(A_{1x}) \, dP_X(x) \\ &= \int \left[\int I_{A_{1x}} \, dP_Y(y) \right] \, dP_X(x) \\ &= \int \left[\int I_{A_{1x}} g(y) \, dm(y) \right] \, dP_X(x) \\ &= \int \int I_{A_{1x}}(y) g(y) \, dm(y) f(x) dm(x) \\ &= \int \int I_A(x, y) f(x) g(y) \, dm^2(x, y) \end{aligned}$$

[More rigorous proof is: define $\mu(A) = \int_A f g \, dm^2$, $\forall A \in \mathcal{B}(\mathbb{R}^2)$, then $\mu = P_{(X,Y)}$ on $\mathcal{C} = \{A \times B : A, B \in \mathcal{B}(\mathbb{R}^2)\}$, and it is a finite measure on $\mathcal{C}$. Hence by uniqueness theorem, $\mu = P_{(X,Y)}$ on $(\mathbb{R}^2, \mathcal{B}(\mathbb{R}^2))$.]

From this, it is clear that $P_{(X,Y)} \ll m^2$ and $\frac{dP_{(X,Y)}}{dm^2} = f \cdot g$ a.e. (m^2).

- (d) If $P_X \ll m, P_Y \ll m$, then we have $f = \frac{dP_X}{dm}, g = \frac{dP_Y}{dm}$ by the definition of Radon-Nikodym derivatives, the following holds for $A \in \mathcal{B}(\mathbb{R})$ (proving the claim):

$$\begin{aligned} (P_X * P_Y)(A) &= \int \int I_A(x+y) \, dP_X(x) dP_Y(y) \\ &= \int \left(\int I_A(x+y) f(x) dm(x) \right) dP_Y(y) \quad (\text{by change of variable}) \\ &= \int \left(\int I_A(u) f(u-y) dm(u) \right) dP_Y(y) \quad (\text{by Fubini's theorem}) \\ &= \int I_A(u) \left(\int f(u-y) dP_Y(y) \right) dm(u) \\ &= \int I_A(u) \left(\int f(u-y) g(y) dm(y) \right) dm(u) \\ &= \int I_A(u) (f * g)(u) dm(u) \\ &= \int_A (f * g)(u) dm(u) \end{aligned}$$

Hence by the uniqueness of the RN derivatives, $\frac{\lambda}{dm} = f * g$ a.e. (m).

- (e) It says Z has the same law as $X + Y$. Part (d) says if X, Y has densities f, g respectively, then Z will have density $f * g$.

3. Solution:

- (i) Define $\mathcal{C} = \{(a, b], (b, \infty) : -\infty \leq a, b < \infty\}$ and $\mu_\Phi((a, b]) = \Phi(b) - \Phi(a)$, $\mu_\Phi((b, \infty)) = 1 - \Phi(b)$. This defines a measure on the semi-algebra $\mathcal{C}$. Caratheodory extension theorem states that there exists an extension μ^* of μ_Φ to $(\mathbb{R}, \mathcal{M})$, and $\mathcal{C} \subset \mathcal{M}$, where

$$\mu^*(A) = \inf \left\{ \sum_{n=1}^{\infty} \mu_\Phi(A_n) : \{A_n\} \subset \mathcal{C}, A \subset \bigcup_{n=1}^{\infty} A_n \right\}, \quad A \subset \mathbb{R},$$

and $\mathcal{M}$ is the set of μ^* -measurable sets, i.e. $\mathcal{M} = \{A \subset \mathbb{R} : \mu^*(E) = \mu^*(E \cap A) + \mu^*(E \cap A^c), \forall E \subset \mathbb{R}\}$. Since $\mathcal{C} \subset \mathcal{M}$, and we know that $\mathcal{B}(\mathbb{R}) = \sigma(\mathcal{C})$, then we have $\mathcal{B}(\mathbb{R}) \subset \mathcal{M}$. Hence, we get an extension of μ_Φ to $(\mathbb{R}, \mathcal{B}(\mathbb{R}))$ by restricting μ^* to $(\mathbb{R}, \mathcal{B}(\mathbb{R}))$. We will denote this measure as μ on $(\mathbb{R}, \mathcal{B}(\mathbb{R}))$. Define $(\Omega, \mathcal{F}, P) = (\mathbb{R}, \mathcal{B}(\mathbb{R}), \mu)$. Define $X : \Omega \rightarrow \mathbb{R}$ as $X(\omega) = \omega$. Clearly, it is measurable and μ is a probability (and hence X is a random variable). Also, the law of X is $P_X(A) = P(X^{-1}(A)) = \mu(A)$ for all $A \in \mathcal{B}(\mathbb{R})$. $F_X(x) = P(X \leq x) = \mu((-\infty, x]) = \Phi(x)$, for all $x \in \mathbb{R}$.

- (ii) Here we use the following change of variable formula for Lebesgue integrals: If X is a measurable transformation from $(\Omega, \mathcal{F}, P)$ to $(\mathbb{R}, \mathcal{B}(\mathbb{R}))$, then $X \in L^1(\Omega, \mathcal{F}, P)$ iff $I \in L^1(\mathbb{R}, \mathcal{B}(\mathbb{R}), P_X)$, where $I(x) = x$, for $x \in \mathbb{R}$, and in that case,

$$\int X dP = \int_{\mathbb{R}} I(x) dP_X.$$

From the definition of P_X in the problem and relationship between Riemann and Lebesgue integrals, we know that the right side of the above equation is

$$\int_{\mathbb{R}} I(x) dP_X = \int_{\mathbb{R}} I(x) \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}} dm(x) = \int_{-\infty}^{\infty} x \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}} dx,$$

where the last expression is a usual Riemann integral. We know from elementary calculus that this integral is equal to 0.

- (iii) From Kolmogorov's Consistency theorem, only condition we need here is that $\mu^{(n+1)}(A \times \mathbb{R}) = \mu^n(A)$ for all $A \in \mathcal{B}(\mathbb{R}^n)$ for all $n \geq 1$. For the sequence of random variables, define for $\omega = (\omega_1, \omega_2, \dots) \in \mathbb{R}^\infty$, $X_n(\omega) = \omega_n$, $n \geq 1$. Then the joint law of $(X_1, \dots, X_n)$ is given by

$$P_{(X_1, \dots, X_n)}(A) = P((X_1, \dots, X_n) \in A) = P(A \times \mathbb{R}^\infty) = \mu^n(A), \text{ for } \forall A \in \mathcal{B}(\mathbb{R}^n), n \geq 1.$$

- (iv) In part (i), define $\mu^{(1)} = \mu \equiv N(0, 1)$ and $\mu^{(n+1)} = \mu^{(n)} \times \mu$, for $n \geq 1$. That is, for each $n \geq 1$, μ^n is the product measure generated by μ on $(\mathbb{R}^n, \mathcal{B}(\mathbb{R}^n))$. Then for each $n \geq 1$, μ^n is a probability on $(\mathbb{R}^n, \mathcal{B}(\mathbb{R}^n))$ and $\mu^{(n+1)}(A \times \mathbb{R}) = \mu^n(A)$ for $A \in \mathcal{B}(\mathbb{R}^n)$. Hence, as in part (i), we can construct a sequence of random variables $\{X_n : n \geq 1\}$ on some probability space $(\Omega, \mathcal{F}, P)$ such that joint distribution of $(X_1, \dots, X_n)$ is given by μ^n , which is the product measure generated by μ . Hence $\{X_n : n \geq 1\}$ is an i.i.d. sequence.