Minimum Absolute Difference in an Array



The absolute difference is the positive difference between two values a and b, is written |a-b| or |b-a| and they are equal. If a=3 and b=2, |3-2|=|2-3|=1. Given an array of integers, find the minimum absolute difference between any two elements in the array.

Example. arr = [-2,2,4]

There are 3 pairs of numbers: [-2,2], [-2,4] and [2,4]. The absolute differences for these pairs are |(-2)-2|=4, |(-2)-4|=6 and |2-4|=2. The minimum absolute difference is 2.

Function Description

Complete the *minimumAbsoluteDifference* function in the editor below. It should return an integer that represents the minimum absolute difference between any pair of elements.

minimumAbsoluteDifference has the following parameter(s):

• int arr[n]: an array of integers

Returns

• int: the minimum absolute difference found

Input Format

The first line contains a single integer n, the size of arr.

The second line contains n space-separated integers, arr[i].

Constraints

- $2 \le n \le 10^5$
- $-10^9 \le arr[i] \le 10^9$

Sample Input 0

3 3 -7 0

Sample Output 0

3

Explanation 0

The first line of input is the number of array elements. The array, arr = [3, -7, 0] There are three pairs to test: (3, -7), (3, 0), and (-7, 0). The absolute differences are:

•
$$|3--7| \Rightarrow 10$$

•
$$|3-0| \Rightarrow 3$$

•
$$|-7-0| \Rightarrow 7$$

Remember that the order of values in the subtraction does not influence the result. The smallest of these absolute differences is 3.

Sample Input 1

```
10
-59 -36 -13 1 -53 -92 -2 -96 -54 75
```

Sample Output 1

1

Explanation 1

The smallest absolute difference is |-54--53|=1.

Sample Input 2

```
5
1 -3 71 68 17
```

Sample Output 2

3

Explanation 2

The minimum absolute difference is |71-68|=3.