DGCA CPL / ATPL AVIATION METEOROLOGY

COMPLETE THEORY & REVISION HANDBOOK (21 CHAPTERS) — FOR DAIVAM MEHTA

Chapter 01: Atmosphere, ISA & Vertical Structure

> **DGCA Syllabus:** Physical properties of the atmosphere, composition, extent, vertical division (Troposphere, Stratosphere, Mesosphere, Thermosphere), Tropopause characteristics, ICAO Standard Atmosphere (ISA).

> **Primary References:** IC Joshi Chapter 1; Oxford/Nordian Chapter 1; Air Meteorology Notes Page 1-3.

- **Atmosphere:** Gaseous envelope surrounding the Earth, rotating with it, bound by gravity.
- **Permanent Gas Composition (Dry Clean Air up to 80 km):**
- **Nitrogen (N_2): 78.08\%** by volume (majority).
- **Oxygen (O_2): 20.95\%** (**~ 21\%**) by volume (supports combustion and life).
- **Argon (Ar): 0.93\%** by volume (inert gas).
- **Carbon Dioxide (CO_2): 0.04\%** (greenhouse gas, absorbs terrestrial radiation).
- *Memory Code:* **N-O-A-C** (**78\% - 21\% - 1\% - 0.04\%**).
- **Variable Gases:**
- **Water Vapour (H_{2O}): 0\%** to **4\%** by volume. Concentrated strictly in lower troposphere (**<5 km**). Primary driver of weather!
- **Ozone (O_3):** Concentrated in **Stratosphere (20-50 km)**, maximum concentration at **25 km** (**75,000 ft**). Absorbs harmful solar Ultraviolet (UV-B/UV-C) radiation.
- **Atmospheric Mass Distribution:**
- **50\%** of total mass lies below **18,000 ft** (**5.5 km** / **500 hPa**).
- **75\%** of total mass lies below **36,000 ft** (**11 km** / **250 hPa**).
- **99\%** of total mass lies below **32 km** (**100,000 ft** / **10 hPa**).

---

graph TD

EXO["EXOSPHERE (Above 500 km) -> Outer Space"]

Auroras occur
99% water vapour
Tropopause
Stratopause
Mesopause

THERMO --> EXO

Layer	Height Range	Temperature Trend	Key Aviation Significance
Troposphere	Surface to $8-18\text{ km}$	**Decreases** at $2^\circ\text{C} / 1,000\text{ ft}$ ($6.5^\circ\text{C}/\text{km}$)	Contains almost all weather, water vapour, clouds, convection, and turbulence.

Tropopause	Boundary layer	**Isothermal** (constant temp)	Boundary lid on convection; jet stream cores and maximum CAT lie just below it.
Stratosphere	Tropopause to 50 km	**Increases** (temperature inversion due to Ozone)	Excellent cruising conditions, free of convective clouds (except occasional volcanic ash or mother-of-pearl clouds).
Mesosphere	50 km to 85 km	**Decreases** to -90°C (coldest zone)	Noctilucent clouds; meteors burn up.
Thermosphere	85 km to 500 km	**Increases** rapidly ($>1000^\circ\text{C}$)	Ionized gases; reflects HF communications; Auroras.

• **Definition:** The transition boundary between Troposphere and Stratosphere where the lapse rate decreases to **2^°C/km or less** for a depth of at least **2 km**.

• **Variation with Latitude:**

• **Equator:** Higher and colder! Height = **16-18 km (55,000 ft)**; Temp = **-75^°C to -80^°C**.

• **Poles:** Lower and warmer! Height = **8-10 km (26,000-30,000 ft)**; Temp = **-45^°C to -50^°C**.

• **Mid-latitudes:** Height = **11 km (36,000 ft)**; Temp = **-56.5^°C**.

• **Variation with Season:**

• Higher and colder in **Summer** than in **Winter**.

• **Tropopause Breaks:**

• The tropopause is not continuous. It has distinct breaks / steps between Tropical, Subtropical, and Polar air masses.

• **Jet stream cores** reside in these tropopause breaks!

graph LR

MID --> BREAK2["Polar Front Jet Break"]

Because the real atmosphere is constantly fluctuating, ICAO established a baseline reference:

• **Mean Sea Level Pressure (P_0): 1013.25 hPa** (**29.92 inHg = 760 mm Hg = 14.7 psi**).

• **Mean Sea Level Temperature (T_0): +15.0^°C** (**288.15 K**).

• **Mean Sea Level Density (rho $_0$): 1.225 kg/m^3**.

• **Standard Lapse Rate (Troposphere up to 11 km): 1.98^°C / 1,000 ft** (**~ 2^°C/1,000 ft** or **6.5^°C/km**).

• **Standard Tropopause: 11.0 km (36,090 ft)** at **-56.5^°C**.

• **Stratosphere (11 km to 20 km): Isothermal** at **-56.5^°C**.

• **Stratosphere (20 km to 32 km):** Temperature **increases at +1.0^°C/km**.

---

1. ■■ **The Equator Tropopause is COLDER than the Pole Tropopause!**

Trap: Candidates intuitively guess the Equator is warmer. The surface is warmer, causing massive vertical convective expansion that pushes the tropopause up to 18 km, allowing it to cool to **-80^°C**!

2. ■■ **The Stratosphere has a Temperature INVERSION:**

Temperature increases with height due to UV absorption by ozone (**O_3**).

3. ■■ **Water Vapour is NOT permanent:**

Permanent gases maintain a constant percentage ratio up to 80 km. Water vapour varies (**0-4\%**).

• **A:** It remains constant throughout the year.

• **B:** It decreases from south to north.

- **C:** It remains constant from north to south.
- **D:** It increases from south to north.

View Answer

Correct Answer: Option B (It decreases from ~18 km at the equator/south to ~8 km at the pole/north).

- **A:** +3°C
- **B:** 0°C
- **C:** -3°C
- **D:** +7°C

View Answer

Correct Answer: Option D (FL 110 to FL 50 is a descent of 6,000 ft. At 2°C/1000 ft, temperature increases by +12°C. -5 + 12 = +7°C).

- **A:** 78% Nitrogen, 21% Oxygen, 1% other gases.
- **B:** 50% Nitrogen, 50% Oxygen.
- **C:** 21% Nitrogen, 78% Oxygen, 1% Argon.
- **D:** 70% Nitrogen, 20% Oxygen, 10% Carbon Dioxide.

View Answer

Correct Answer: Option A.

Chapter 02: Atmospheric Pressure & Altimetry

> **DGCA Syllabus:** Atmospheric pressure definition, units, isobars, pressure systems (High, Low, Trough, Ridge, Col), pressure lapse rate, altimeter working principle, QNH, QFE, QNE, QFF, altimeter errors (pressure and temperature errors), Transition Altitude (TA), Transition Level (TL).

> **Primary References:** IC Joshi Chapter 2; Oxford/Nordian Chapters 2 & 3; Volare Question Bank.

- **Atmospheric Pressure:** Force exerted by the weight of a column of air over a unit horizontal area.
- At sea level: **~ 1.033 kg/cm^2 = 14.7 psi**.
- **Units:**
- **Hectopascal (hPa):** Standard SI unit. **1 hPa = 100 Pa = 1 millibar (mb)**.
- **Inches of Mercury (inHg):** Used in North America. **1013.25 hPa = 29.92 inHg**.
- **Millimeters of Mercury (mmHg): 1013.25 hPa = 760 mmHg**.
- **Isobar:** Line joining places of equal atmospheric pressure reduced to Mean Sea Level (MSL).
- Drawn at standard intervals of **2 hPa** or **4 hPa**.
- Close isobars **\implies** Steep pressure gradient **\implies** Strong winds!
- Wide isobars **\implies** Weak pressure gradient **\implies** Light winds!

---

graph TD

subgraph Pressure Systems

end

Feature	Cyclone (Low / Depression)	Anticyclone (High)	Trough	Ridge	Col

Pressure Pattern	Enclosed Low	Enclosed High	'V' or 'U' shaped isobar extension from Low	Elongated extension from High	Saddle region between 2 Highs & 2 Lows
Vertical Air Motion	**Ascending (upward)**	**Descending (subsiding)**	Ascending	Descending	Variable / Weak
Associated Weather	Clouds, turbulence, rain, storms	Clear skies, calm winds, radiation fog/smog	Cloud bands, showers, windshear	Fine, settled weather	In summer: thunderstorms; in winter: dense fog

---

Because air is compressible, pressure drops rapidly near the ground and more slowly higher up:

Altitude Range Pressure Lapse Rate

0 to 5,000 ft 1 hPa = 30 ft (Standard DGCA constant)

10,000 ft (700 hPa) 1 hPa = 37 ft

18,000 ft (500 hPa) 1 hPa = 50 ft (Half MSL pressure)

30,000 ft (300 hPa) 1 hPa = 75 ft

39,000 ft (200 hPa) 1 hPa = 100 ft

- **Cold Air Column:** Cold air is dense and contracts downward. Pressure drops **faster** with height.
- **Warm Air Column:** Warm air is light and expands upward. Pressure drops **slower** with height.
- **Key Exam Consequence:** At any high altitude level (e.g., **10,000 ft**), the pressure is **HIGHER over warm air** and **LOWER over cold air**!

graph LR

COLD["COLD AIR COLUMN
Air contracts down
Isobars packed tightly
Low pressure aloft"]

WARM["WARM AIR COLUMN
Air expands upward
Isobars spaced widely
High pressure aloft"]

---

graph TD

QFE["QFE (Field Elevation Pressure)
Reference: Airfield Elevation
On Ground: Reads 0 ft
In Flight: Reads Height AGL"]

QNH["QNH (Altimeter Subscale Setting)
Reference: Mean Sea Level (MSL) via ISA
On Ground: Reads Airfield Elevation
In Flight: Reads Altitude AMSL"]

QNE["QNE (Standard 1013.25 hPa)
Reference: Standard 1013.25 Datum
Above Transition Altitude
Reads Flight Level (FL)"]

QFF["QFF (Synoptic Chart Pressure)
Reference: MSL via ACTUAL Air Temp
Used for surface weather maps"]

$\mathbf{QNH = QFE + (Aerodrome Elevation (ft) / 30)}$

$\mathbf{QFE = QNH - (Aerodrome Elevation (ft) / 30)}$

Condition	QNH vs QFF Relationship
Airfield at Sea Level (Elevation = 0)	$\mathbf{\text{QNH} = \text{QFE} = \text{QFF}}$
Actual Temperature = ISA Temperature	$\mathbf{\text{QNH} = \text{QFF}}$
Air column is COLDER than ISA ($\text{OAT} < \text{ISA}$)	$\mathbf{\text{QFF} > \text{QNH}}$
Air column is WARMER than ISA ($\text{OAT} > \text{ISA}$)	$\mathbf{\text{QFF} < \text{QNH}}$

If a pilot flies from an area of High pressure into Low pressure without resetting the altimeter:

- The subscale remains set to the higher pressure.
- The instrument measures ambient low pressure and assumes the aircraft has climbed.
- It indicates a higher altitude than the aircraft is actually flying!
- **Result:** Aircraft is flying **dangerously LOWER than indicated**!

$\mathbf{True Altitude = Indicated Altitude + [(Current QNH - Subscale Setting) x 30 ft]}$

$\mathbf{Correction (ft) = 4 x <= ft((Indicated Altitude / 1,000) \right) x (OAT - T_{ISA})}$

$$\begin{aligned}

\mathbf{\text{"High to Low, Look Out Below!"}} &\implies \text{Altimeter OVER-READS (You are LOWER than indicated!)} \\

\mathbf{\text{"Low to High, Clear Sky!"}} &\implies \text{Altimeter UNDER-READS (You are HIGHER than indicated!)} \\

\mathbf{\text{"Cold to Warm, Do No Harm!"}} &\implies \text{You are HIGHER than indicated} \\

\mathbf{\text{"Warm to Cold, Danger Foretold!"}} &\implies \text{You are LOWER than indicated (True Alt < Indicated)}

\end{aligned}$$

---

graph TD

FL["FLIGHT LEVELS (Standard 1013.25 hPa)"]

TL["TRANSITION LEVEL (Lowest usable Flight Level by ATC)"]

T_LAYER["TRANSITION LAYER (Buffer zone - do not cruise here!)"]

TA["TRANSITION ALTITUDE (Fixed altitude on departure - Set QNH)"]

ALT["ALTITUDES (QNH Setting - Referenced to MSL)"]

Climbing through TA: Change QNH to 1013.25
Descending through TL: Change 1013.25 to QNH

- **Transition Altitude (TA):** Fixed published altitude for an aerodrome. During climb, at or above TA, altimeter setting is switched from local **QNH to Standard 1013.25 hPa**.
- **Transition Level (TL):** The lowest available flight level above TA, determined dynamically by ATC depending on the local QNH.
- **Transition Layer:** Airspace between TA and TL. Used only for climbing/descending; level cruising is strictly prohibited.
- **Formula:**
- When **QNH >= 1013.25 hPa \implies Transition Layer thickness is maximum**.
- When **QNH < 1013.25 hPa \implies Transition Layer thickness is minimum**.

- **A:** The elevation = 0.

- **B:** T actual = T standard.
- **C:** T actual > T standard.
- **D:** T actual < T standard.

View Answer

Correct Answer: Option A (At sea level, elevation is 0, so QNH = QFE).

- **A:** 3,600 ft.
- **B:** 3,000 ft.
- **C:** 2,400 ft.
- **D:** 3,200 ft.

View Answer

Correct Answer: Option C

Explanation: High to Low, look out below! Pressure drop = **1020 - 1000 = 20 hPa**.

Height difference = **20 x 30 = 600 ft**.

Actual True Altitude = **3000 - 600 = \mathbf{2,400 ft}**.

- **A:** Overcast clouds, strong turbulence, and severe precipitation.
- **B:** Subsiding air, light winds, clear skies, and potential radiation fog.
- **C:** Continuous squalls and towering cumulonimbus clouds.
- **D:** Steep pressure gradient and hurricane-force winds.

View Answer

Correct Answer: Option B.

Chapter 03: Temperature, Heat Transfer & Inversions

> **DGCA Syllabus:** Heating and cooling of the atmosphere, radiation (solar vs terrestrial), conduction, convection, advection, latent heat of phase changes, diurnal variation of temperature, temperature inversions (types, formation, and aviation hazards).

> **Primary References:** IC Joshi Chapter 3; Air Meteorology Notes Page 1-2; Oxford/Nordian Chapter 1; Volare Question Bank.

- **Atmosphere is Heated from BELOW, Not from Above!**
- Solar radiation passes through the atmosphere with very little direct absorption.
- The ground absorbs incoming solar short-wave radiation, heats up, and re-radiates **terrestrial long-wave infrared radiation**, which is absorbed by greenhouse gases (**H_{2O}** vapour and **CO_2**) in the lower atmosphere.
- **Heat Transfer Mechanisms:**
- **Radiation:** Transfer of energy by electromagnetic waves without requiring a material medium.
- *Solar Radiation (Insolation):* Short-wave (**0.15 to 4.0 \mum**), visible and UV.
- *Terrestrial Radiation:* Long-wave (**4.0 to 100 \mum**), infrared. Occurs continuously day and night.
- **Conduction:** Heat transfer by direct molecular contact. Air is a very poor conductor; conduction only affects the thin layer in direct contact with the ground (first few centimeters/inches).
- **Convection:** Heat transfer by vertical buoyant motion of warm, less dense air parcels. Creates thermals, cumulus clouds, and thunderstorms.

• **Advection:** Heat transfer by the **horizontal movement** of an air mass (e.g., warm air blowing over a cold ocean surface, triggering advection fog).

• **Turbulence / Mechanical Mixing:** Eddies caused by wind friction against surface irregularities mixing warm and cold air layers.

---

graph TD

subgraph Heat Released to Environment Warming

V_TO_L["Condensation (Vapour -> Liquid)"]

L_TO_S["Freezing (Liquid -> Solid)"]

V_TO_S["Deposition (Vapour -> Solid Ice)"]

end

subgraph Heat Absorbed from Environment Cooling

S_TO_L["Melting / Thawing (Solid -> Liquid)"]

L_TO_V["Evaporation (Liquid -> Vapour)"]

S_TO_V["Sublimation (Solid -> Vapour)"]

end

• **Latent Heat of Evaporation / Condensation: ~ $\mathbf{2.5 \times 10^6}$ J/kg}** (**597 cal/g**).

• **Latent Heat of Fusion (Melting / Freezing): ~ $\mathbf{3.34 \times 10^5}$ J/kg}** (**80 cal/g**).

• **Sensible Heat:** Heat that causes a change in temperature that can be measured with a thermometer. Contributes **\mathbf{ ~ 23\%}** to total atmospheric heating.

• **Latent Heat Release:** The primary engine powering thunderstorms, tropical revolving storms (cyclones), and vertical updrafts!

---

graph LR

SUNRISE["Sunrise (~05:00 - 06:00)
MINIMUM TEMPERATURE
(Radiation loss exceeded insolation all night)"] --> NOON["Solar Noon (~12:00)
Maximum Solar Insolation"]

NOON --> PEAK["Afternoon (~14:00 - 15:00)
MAXIMUM TEMPERATURE
(Lag effect: Outgoing terrestrial radiation balances incoming insolation)"]

1. **Cloud Cover:**

• Day: Clouds reflect incoming solar radiation **\implies** Lower maximum day temperature.

• Night: Clouds absorb and re-radiate terrestrial heat back to ground **\implies** Higher minimum night temperature (warmer night).

• *Result:* **Clouds drastically reduce the diurnal temperature range.**

2. **Wind Speed:**

• Calm winds: Pronounced cooling at surface, large diurnal range.

• Strong winds: Mixes air mechanically, distributes heat over a deeper layer **\implies** Smaller diurnal range.

3. **Moisture Content (Humidity):**

• Humid tropical climates have a small diurnal range (water vapour traps heat).

• Dry desert climates have extreme diurnal ranges (**>25^°C** difference between day and night).

4. **Surface Nature (Land vs Sea):**

• Land has low specific heat capacity, heats up and cools down rapidly.

• Water has high specific heat capacity, transparent (heat penetrates deep), constantly mixes, and evaporates **\implies Diurnal range of sea surface temperature is <1^°C!**

An inversion occurs when temperature **INCREASES with altitude** (**ELR < 0**). It acts as an impenetrable lid to vertical air motion.

graph TD

INV["TEMPERATURE INVERSIONS"]

INV --> RAD["1. Ground / Radiation Inversion
Clear calm winter nights
Forms from surface up to 500-1,000 ft"]

INV --> SUB["2. Subsidence Inversion
In Anticyclones / Highs
Sinking air warms adiabatically at DALR (3°C/1000ft) aloft"]

INV --> FRON["3. Frontal Inversion
Warm air overrides cold air at front
Sloping layer aloft"]

INV --> TURB["4. Turbulence / Friction Inversion
Wind mixing lowest 1,000-2,000 ft
Cools at top, forms Sc clouds"]

Inversion Type	Formation Mechanism	Typical Level	Associated Cloud/Weather
Ground / Radiation	Rapid terrestrial radiative cooling of ground on clear, calm night	Surface to $1,000\text{ ft}$ AGL	Radiation fog, heavy ground frost, dense smog
Subsidence	Slow sinking (subsidence) of air in High pressure; warms adiabatically at DALR ($3^\circ\text{C}/1000\text{ ft}$)	Upper levels ($3,000\text{ to }10,000\text{ ft}$)	Clear skies, haze layer, persistent dry weather
Frontal	Warm air mass overrunning a denser cold air mass at a frontal zone	Sloping aloft ($2,000\text{ to }15,000\text{ ft}$)	Ns, As, continuous rain, freezing rain hazard
Turbulence / Friction	Strong surface wind mixing the boundary layer; top cools adiabatically	Top of friction layer ($1,500\text{ to }3,000\text{ ft}$)	Stratocumulus (Sc) cloud deck

---

1. **Severe Degradation of Visibility:**

• Inversions trap all smoke, pollutants, moisture, and dust beneath the inversion base.

• Forms a sharp, clearly visible "haze line" from above.

2. **Low-Level Wind Shear (LLWS):**

• The air above the inversion is uncoupled from ground friction and may blow rapidly (Low Level Jet).

• An aircraft on takeoff climb or final approach crossing the inversion layer encounters abrupt changes in airspeed, pitch, and glide path.

3. **Severe Turbulence:**

• Intense wind shear generates violent mechanical turbulence right across the inversion boundary.

4. **Carburettor & Airframe Icing:**

• In frontal inversions with rain falling from warm air into sub-zero cold air below, **freezing rain (FZRA)** forms, causing catastrophic clear ice accretion!

• **A:** Low

- **B:** High
- **C:** Within +/-5°C of ISA
- **D:** Standard

View Answer

Correct Answer: Option A (Low)

Explanation: In standard atmosphere, 700 hPa corresponds to 10,000 ft (FL 100).

T_{ISA} = 15 - (2 x 10) = -5^°C.
Actual temperature is **-15^°C**, which is **10^°C** colder than ISA (Low).

- **A:** Temperature increases with height.
- **B:** Temperature remains constant with height.
- **C:** Temperature decreases faster than the standard lapse rate.
- **D:** Relative humidity decreases to zero.

View Answer

Correct Answer: Option A.

- **A:** Strong winds, overcast cloud cover, and summer afternoon.
- **B:** Calm or light wind, clear skies, and a long winter night.
- **C:** Gale force winds and a passing warm front.
- **D:** High humidity, thick cumulus clouds, and solar noon.

View Answer

Correct Answer: Option B.

Chapter 04: Air Density & Aircraft Performance

> **DGCA Syllabus:** Definition of air density, relationship between pressure, temperature, and density (Ideal Gas Law), influence of humidity on air density, Density Altitude (calculation and significance), impact on engine performance, takeoff/landing distance, and rate of climb.

> **Primary References:** IC Joshi Chapter 4; Oxford/Nordian Chapter 2.4; Volare Question Bank.

- **Air Density (rho):** Mass of air per unit volume (**kg/m^3** or **lb/ft^3**).
- ISA Sea Level Density: **\mathbf{1.225 kg/m^3** (**0.0765 lb/ft^3**).
- **The Equation of State (Ideal Gas Law):**

$\mathbf{P = rho R T \implies rho = (P / R T)}$

- **rho** is **directly proportional** to Pressure (**P**).
- **rho** is **inversely proportional** to Absolute Temperature (**T** in Kelvin).
- **The Humidity Paradox (DGCA Exam Favorite!):**
- **Moist air is LESS DENSE than dry air at the same pressure and temperature!**
- *Why?* Molecular weight of water vapour (**H_{2O}**) is **18 g/mol**, whereas dry air (mostly **N_2** and **O_2**) has an average molecular weight of **28.96 g/mol**. When water vapour enters a parcel of air, it displaces heavier nitrogen and oxygen molecules!

graph TD

DENSITY["AIR DENSITY (ρ)"]

Directly Proportional

Inversely Proportional
Inversely Proportional

• **Definition:** The altitude in the standard atmosphere (ISA) at which the air density is equal to the current air density of the actual atmosphere.

• **In simple terms:** It is the altitude that the aircraft "feels" it is flying at!

• **High Density Altitude = LOW Air Density = Thin Air!**

graph LR

HOT["High OAT
+ Low Pressure (Low QNH)
+ High Humidity"] --> HIGH_DA["HIGH DENSITY ALTITUDE
(Thin Air)"]

$\mathbf{Pressure Altitude (PA) = Elevation + [(1013.25 - QNH) x 30]}$

$\mathbf{Density Altitude (DA) = PA + [120 x (OAT - T_{ISA})]}$

> **Question:** An airfield has an elevation of **4,000 ft**, QNH is **998 hPa**, and the current ambient temperature (OAT) is **+30^°C**. Calculate the Density Altitude.

>

> **Step 1: Calculate Pressure Altitude (PA)**

> **PA = 4,000 + (1013 - 998) x 30 = 4,000 + (15 x 30) = \mathbf{4,450 ft}**

>

> **Step 2: Calculate ISA Temperature at PA**

> **T_{ISA} = 15^°C - (2 x 4.45) = 15 - 8.9 = \mathbf{+6.1^°C}**

>

> **Step 3: Calculate ISA Deviation (Delta ISA)**

> **Delta ISA = OAT - T_{ISA} = 30 - 6.1 = \mathbf{+23.9^°C}** (ISA **+24**)

>

> **Step 4: Calculate Density Altitude (DA)**

> **DA = 4,450 + (120 x 23.9) = 4,450 + 2,868 = \mathbf{7,318 ft}**

> *Result:* Even though the physical airfield is at **4,000 ft**, the aircraft will perform as if it were taking off from a high-altitude runway at **7,318 ft**!

---

1. **Thrust & Power Output:**

• Naturally aspirated piston engines suffer loss of power because less oxygen mass enters the cylinders per intake stroke.

• Jet engines suffer reduced thrust due to reduced mass flow rate through the compressor and turbines.

2. **True Airspeed (TAS) vs Indicated Airspeed (IAS):**

• The pitot tube measures dynamic pressure: **q = (1 / 2) rho V_{TAS}^2**.

• At lower density (**rho**), the aircraft must fly at a **higher True Airspeed (TAS)** to generate the same dynamic pressure and lift (same IAS).

• *Rule of Thumb:* TAS increases by approximately **1.75\% to 2\% per 1,000 ft** of altitude for a constant IAS.

3. **Takeoff & Landing Distances:**

• Because stall speed and liftoff speed occur at a fixed IAS, the aircraft must reach a much higher ground speed to become airborne.

• Combined with reduced engine acceleration, takeoff distance can double or triple on a hot, high, and humid day!

---

1. ■■ **"Hot, High, and Humid" \to Triple Threat!**

All three factors combine to create maximum density altitude and worst performance.

2. ■■ **Altimeter vs Air Density:**

The pressure altimeter DOES NOT sense air density; it only senses static pressure! An altimeter set to 1013 will read Pressure Altitude, not Density Altitude.

• **A:** Pressure is high, temperature is low, and humidity is low.

• **B:** Pressure is low, temperature is high, and humidity is high.

• **C:** Pressure is high, temperature is high, and humidity is high.

• **D:** Altitude is high, temperature is high, and air is dry.

View Answer

Correct Answer: Option A (Cold, dry, high-pressure air is the densest possible air).

• **A:** Decreases takeoff distance because moisture lubricates the wings.

• **B:** Increases takeoff distance because moist air is less dense, reducing engine power and wing lift.

• **C:** Has no effect because air density depends solely on temperature.

• **D:** Increases rate of climb due to latent heat release in the boundary layer.

View Answer

Correct Answer: Option B.

Chapter 05: Humidity & Adiabatic Processes

> **DGCA Syllabus:** Water vapour in the atmosphere, saturation, vapour pressure, Dew Point (T_d), Relative Humidity (RH), mixing ratio, wet-bulb temperature, adiabatic expansion and cooling, Dry Adiabatic Lapse Rate (DALR), Saturated Adiabatic Lapse Rate (SALR), Lifting Condensation Level (LCL).

> **Primary References:** IC Joshi Chapter 5; Oxford/Nordian Chapter 4; Air Meteorology Notes Page 1-2; Volare Question Bank.

• **Water Vapour:** Invisible gas in the atmosphere, ranging from **0\%** (cold dry polar regions) to **4\%** (warm tropical oceans).

• **Vapour Pressure (e):** Partial pressure exerted by water vapour molecules alone (hPa).

• **Saturation Vapour Pressure (e_s):** Maximum vapour pressure possible at a given temperature.

• Increases exponentially with temperature: **Warm air can hold significantly more water vapour than cold air!**

• **Dew Point Temperature (T_d):** The temperature to which moist air must be cooled at constant pressure to become saturated.

• Higher dew point **\implies** Greater moisture content.

• When Air Temperature (**T**) = Dew Point (T_d) **\implies \mathbf{Relative Humidity = 100\%}** (Condensation begins **\to** Fog or Cloud!).

• **Depression of the Dew Point (Spread):**

$\mathbf{Spread = T - T_d}$

• A small spread (**0-2^°C**) indicates high humidity and imminent fog or low cloud.

• **Relative Humidity (RH):**

$\mathbf{RH = (Actual Vapour Pressure (e) / Saturation Vapour Pressure (e_s)) x 100\% ~ (Actual Mixing Ratio (w) / Saturation Mixing Ratio (w_s)) x 100\%}$

graph TD

HUMIDITY["HUMIDITY MEASURES"]

HUMIDITY --> DP["Dew Point (Td)
Direct measure of absolute moisture quantity"]

HUMIDITY --> RH["Relative Humidity (RH)
Ratio of moisture present to moisture capacity"]

HUMIDITY --> SPREAD["Spread (T - Td)
Closeness to saturation
When Spread = 0 => Fog / Clouds form"]

- **Dry-Bulb Thermometer:** Measures actual ambient air temperature (**T**).
- **Wet-Bulb Thermometer:** Bulb enclosed in a wet muslin wick exposed to airflow.
- Evaporation cools the bulb.
- In dry air: Rapid evaporation **\implies** Large wet-bulb depression (**$T - T_w$**).
- In saturated air (**RH = 100\%**): No evaporation **\implies $\mathbf{T = T_w = T_d}$**.

An **adiabatic process** is a thermodynamic change of state taking place **without any heat transfer** into or out of the parcel of air.

- Rising air expands due to lower ambient pressure aloft **\implies** Does work on surrounding air **\implies Cools adiabatically**.
- Sinking air is compressed by higher ambient pressure **\implies** Work done on the parcel **\implies Warms adiabatically**.

graph TD

RISING["RISING AIR PARCEL"]

RISING --> EXP["Expands due to lower pressure aloft"]

EXP --> COOL1["Cools at DALR (3.0°C / 1,000 ft) until saturated"]

COOL1 --> LCL["Reaches Dew Point at LCL (Lifting Condensation Level)"]

LCL --> COND["Condensation occurs: Cloud droplets form!"]

COND --> LATENT["LATENT HEAT RELEASED into parcel"]

LATENT --> COOL2["Further ascent cools at SLOWER rate: SALR (~1.5°C / 1,000 ft)"]

Type	Value in Imperial	Value in Metric	Constancy	Description
DALR (Dry Adiabatic)	**$3.0^\circ\text{C} / 1,000\text{ ft}$**	**$0.98^\circ\text{C} / 100\text{ m} \approx 10 ^\circ\text{C}/\text{km}$**	**Constant**	Applies to unsaturated air ($\text{RH} < 100\%$).
SALR (Saturated Adiabatic)	**$\approx 1.5^\circ\text{C} / 1,000\text{ ft}$**	**$\approx 5^\circ\text{C}\text{ to }6^\circ\text{C} / \text{km}$**	**Variable!**	Applies once air saturates. Slower cooling because latent heat of condensation is released.
ELR (Environmental)	Variable	Variable	Variable	Actual ambient temperature profile measured by radiosonde balloon.

> **Key Rule:** SALR is always **less than** DALR because latent heat released during condensation offsets the expansion cooling!

> In very cold air (e.g., upper troposphere, **-40^°C**), very little water vapour can condense, so SALR approaches DALR.

When an unsaturated surface parcel is lifted convective heating:

- Temperature cools at the DALR: **3.0^°C / 1,000 ft**.
- Dew Point cools at the Dew Point Lapse Rate: **~ 0.5^°C / 1,000 ft**.
- Net rate of convergence (Spread closure): **3.0 - 0.5 = \mathbf{2.5^°C / 1,000 ft}**.

$\mathbf{Cloud Base (ft AGL) = (T$_{surface}$ - T$_{d}$ (surface)} / 2.5) x 1,000 = 400 x (T - T$_{d}$)}$

- **A:** 2°C / 1,000 ft
- **B:** 1.5°C / 1,000 ft
- **C:** 3°C / 1,000 ft
- **D:** 0.65°C / 100 m

View Answer

Correct Answer: Option C (**3.0^°C / 1,000 ft** or **~ 10^°C/km**).

- **A:** Because moist air is denser than dry air.
- **B:** Because of the release of latent heat of condensation during ascent.
- **C:** Because solar radiation heats saturated air parcels.
- **D:** Because the pressure lapse rate decreases inside clouds.

View Answer

Correct Answer: Option B.

Chapter 06: Winds & Circulation Dynamics

> **DGCA Syllabus:** Forces acting on horizontal airflow (PGF, Coriolis, Centrifugal, Friction), Geostrophic wind, Gradient wind, Cyclostrophic wind, Buys Ballot's Law, Veering and Backing, Diurnal variation of surface wind, Local winds (Sea/Land Breeze, Anabatic/Katabatic winds, Föhn/Chinook wind, Mountain winds).

> **Primary References:** IC Joshi Chapter 6; Oxford/Nordian Chapter 6; Volare Question Bank.

- **Wind:** Horizontal motion of air relative to the Earth's surface (vertical motion is termed *current* or *updraft/downdraft*).
- **Wind Direction:** The direction **FROM WHICH** the wind blows (e.g., wind from **270°** is a Westerly wind).
- *Tower / ATC Reports:* Given in **Degrees Magnetic** (aligned with runway heading).
- *METAR / TAF / Met Charts:* Given in **Degrees True**!
- *Memory Rule:* **"If you read it, it's True; if you hear it, it's Magnetic!"**
- **Veering vs Backing:**
- **Veering:** Clockwise change in wind direction (e.g., **090° \to 120°**).
- **Backing:** Counter-clockwise change in wind direction (e.g., **120° \to 090°**).
- **Gale:** Sustained wind speed of **>= 34 kt** (Beaufort Force 8).
- **Squall:** Sudden abrupt increase in wind speed by at least **16 kt**, reaching **>= 22 kt**, and lasting for at least **1 minute**.
- **Gust:** Brief, sharp increase in wind speed lasting only a few seconds.

---

graph TD

CENT["3. Centrifugal Force
Acts radially outwards on curved paths"]

FRIC["4. Friction Force
Acts opposite to motion
Confined to boundary layer (<3,000 ft)"]

PGF --> BALANCE["Equilibrium Wind Types"]

CF --> BALANCE

CENT --> BALANCE

FRIC --> BALANCE

1. **Geostrophic Wind (V_g):**
• Equilibrium between **PGF** and **Coriolis Force**.
• Valid for **straight, parallel isobars** in the **free atmosphere** (above friction layer, **>3,000 ft**).
• Blows strictly parallel to isobars.
• Formula: **$\mathbf{V_g = (1 / 2 \rho \Omega \sin\phi) (\Delta P / \Delta n)}$**.
• Strongest at **low latitudes** (small **\sinphi**) and in **low air density** for a given pressure gradient.
2. **Gradient Wind (V_{grad}):**
• Equilibrium between **PGF, Coriolis Force, and Centrifugal Force** along **curved isobars**.
• **Anticyclone (High):** Centrifugal force aids PGF **\implies $\mathbf{V_{grad} > V_g}$** (**Supergeostrophic**).
• **Cyclone (Low):** Centrifugal force opposes PGF **\implies $\mathbf{V_{grad} < V_g}$** (**Subgeostrophic**).
3. **Cyclostrophic Wind:**
• Balance between **PGF** and **Centrifugal Force** alone (Coriolis force is negligible).
• Occurs in tight, intense vortices near the equator or on small scales: **Tornadoes, Waterspouts, Dust Devils**.

> **In the Northern Hemisphere:** Stand with your back to the wind **\implies Low pressure is on your LEFT**, and High pressure is on your RIGHT!

> *(In the Southern Hemisphere: Low is on your RIGHT).*

• Below **2,000-3,000 ft**, surface friction slows the wind down.
• Slower wind **\implies** Weaker Coriolis force **\implies** PGF pulls wind across isobars toward low pressure.
• As an aircraft **climbs out of the friction layer**:
• Wind speed **increases** (**2 x** over land).
• Wind **VEERS** (turns clockwise by **~ 30°** over land, **~ 10°** over sea).
• As an aircraft **descends into the friction layer**:
• Wind speed **decreases**.
• Wind **BACKS** (turns counter-clockwise) and crosses isobars at **~ 30°** toward Low pressure.
• **By Day:** Solar heating triggers thermal convection, mixing high-momentum air from aloft down to the surface **\implies** Surface wind **veers and increases in speed with gusts**.
• **By Night:** Ground cools, inversion forms, decouples surface from upper winds **\implies** Surface wind **backs and decreases to calm or light**.

---

graph TD

LOCAL["LOCAL WINDS"]

LOCAL --> SEA_LAND["Sea & Land Breezes
Differential heating of land vs water"]

LOCAL --> MOUNTAIN["Anabatic & Katabatic
Mountain slope heating / cooling"]

LOCAL --> FOHN["Föhn / Chinook
Warm dry lee-side wind"]

Feature	Sea Breeze	Land Breeze
Timing	Daytime (maximum $14:00 - 15:00$ local)	Nighttime (around dawn / sunrise)
Flow Direction	**From Sea to Land**	**From Land to Sea**
Mechanism	Land heats faster than sea $\to$ thermal low over land	Land cools faster than sea $\to$ shallow high over land
Speed & Extent	$10-20\text{ kt}$, extends $15-30\text{ NM}$ inland	Lighter ($5-10\text{ kt}$), extends only $5-10\text{ NM}$ offshore

• **Anabatic Wind (Upslope):** Daytime solar heating warms mountain slopes faster than valley air; warm air rises **upslope**. Lighter (**5-10 kt**).

• **Katabatic Wind (Drainage / Downslope):** Nighttime radiative cooling chills dense air on high plateaus/glaciers; gravity drains cold heavy air **down into valleys**. Can be violently fast (**>50 kt** in Greenland/Antarctica, e.g., Mistral, Bora).

• A warm, dry wind that blows down the **leeward side** of a mountain range.

• **Thermodynamic Cycle:**

1. Moist air ascends windward slope **\implies** Cools at **DALR** (**3^°C/1000 ft**) until saturation **\implies** Cools at **SALR** (**~ 1.5^°C/1000 ft**) with heavy rain/snow.

2. Moisture is stripped out at the summit.

3. Air descends leeward slope dry **\implies** Warms at the rapid **DALR** (**3^°C/1000 ft**) the entire way down!

4. **Result:** Leeward air arrives drastically **WARMER and DRIER** than it started!

• **A:** From the left (Port drift).

• **B:** From the right (Starboard drift).

• **C:** Headwind.

• **D:** Tailwind.

View Answer

Correct Answer: Option B (From the right)

Explanation: By Buys Ballot's Law, when heading directly toward the Low, the Low is ahead; wind blows counter-clockwise around Lows in NH, crossing from right to left across track.

• **A:** Backs and decreases in speed.

• **B:** Veers and increases in speed.

• **C:** Backs and increases in speed.

• **D:** Veers and decreases in speed.

View Answer

Correct Answer: Option B (Veers and increases in speed).

• **A:** NE trade winds.

• **B:** SE trade winds.

• **C:** Westerly winds.

• **D:** Polar easterlies.

View Answer

Correct Answer: Option A (NE trade winds).

Chapter 07: Visibility, Fog, Mist & Haze

> **DGCA Syllabus:** Definitions of visibility, Meteorological Optical Range (MOR), Runway Visual Range (RVR), Slant vs Vertical visibility, distinction between Fog, Mist, Haze, and Smoke, types of Fog (Radiation, Advection, Upslope, Steam, Frontal), formation conditions, dissipation, and operational flight hazards.

> **Primary References:** IC Joshi Chapter 7; Oxford/Nordian Chapter 8; Volare Question Bank.

• **Visibility (Meteorological):** The greatest horizontal distance at which a black object of suitable dimensions can be seen and recognized against the horizon sky by normal day vision.

• **Runway Visual Range (RVR):** The range over which the pilot of an aircraft on the centreline of a runway can see the runway surface markings or the lights delineating the runway or identifying its centreline.

• Measured automatically by **Transmissometers** or **Forward-Scatter Meters**.

• Reported in METAR when visibility or RVR is **< 1,500 m**.

• **Slant Visibility vs Vertical Visibility:**

• **Slant Visibility:** Pilot's line-of-sight visibility looking diagonally from the cockpit down toward the approach lights.

Critical Hazard: In shallow radiation fog, looking straight down from overhead, runway lights may be visible (good vertical visibility), but on 3-degree glide slope (slant path), the aircraft enters dense fog and visual references vanish!

• **Vertical Visibility (`VV`):** Maximum vertical distance into an obscuring layer (such as fog or snow) from which ground-based objects are visible.

Phenomenon	METAR Code	Visibility Limit	Relative Humidity (RH)	Particle Nature
Fog	**`FG`**	**$< 1,000\text{ m}$**	**$\approx 100\%$** (usually $\ge 95\%$)	Microscopic liquid water droplets (or ice crystals) suspended in air.
Mist	**`BR`**	**$1,000\text{ m to }5,000\text{ m}$**	**$\ge 95\%$**	Liquid water droplets; feels damp; grayish appearance.
Haze	**`HZ`**	**$\le 5,000\text{ m}$**	**$< 95\%$** (usually $< 80\%$)	Extremely fine dry particles (dust, salt, pollution); opalescent blue against dark ground, yellowish against sun.
Smoke	**`FU`**	Variable	Any	Suspended combustion particles; distinct odor; sun appears red.

---

graph TD

FOG["FOG TYPES"]

itions	Location	Wind Requirement
night, high RH, radiative cooling	**Land ONLY** (Never over open sea due to negligible diurnal SST variation)	**Light breeze: $2\text{ to }8\text{ kt}$** *(Calm $>10\text{ kt}$ gives low stratus)*
g horizontally over a cold surface	**Sea and Coastal Land** (can blow inland)	**Moderate: $8\text{ to }15\text{ kt}$** (Persiste winds!)
up terrain by wind, cooling adiabatically	Windward mountain slopes and elevated plateaus	**Moderate upslope wind: $10-20\text{ kt}$**

over warm water; intense evaporation	Polar waters, ice-free lakes in winter	Light to moderate
air aloft evaporates into cold stable surface air	Ahead of **Warm Front** or along cold front	Variable

---

1. ■■ **Radiation Fog NEVER forms at sea:**

The diurnal variation of sea surface temperature is **<1^°C**, which is insufficient to trigger radiative cooling to dew point.

2. ■■ **The Wind Sweet Spot for Radiation Fog:**

• **Calm (<2 kt) \implies** Only Dew or Frost forms (no vertical mixing).

• **\mathbf{2 to 8 kt} \implies Ideal Radiation Fog** (slight turbulence mixes cold layer through 100-500 ft).

• **>10 kt \implies** Stratus cloud deck forms instead of fog!

3. ■■ **Advection Fog can persist in daylight:**

Unlike radiation fog, advection fog does NOT burn off under the midday sun because the underlying cold surface keeps cooling the air continuously.

• **A:** Strong winds, overcast clouds, and afternoon sunshine.

• **B:** High relative humidity, clear skies, and light breeze of 2 to 8 knots on a long winter night.

• **C:** Gale force winds over a warm ocean current.

• **D:** Rapid passage of a fast-moving cold front.

View Answer

Correct Answer: Option B.

• **A:** Advection fog only forms in mountainous areas.

• **B:** Advection fog requires calm air and occurs exclusively in summer.

• **C:** Advection fog forms when warm moist air moves over a colder surface and can form over both sea and land day or night.

• **D:** Advection fog burns off rapidly 30 minutes after sunrise.

View Answer

Correct Answer: Option C.

Chapter 08: Clouds & Vertical Motion

> **DGCA Syllabus:** Cloud formation mechanisms (convection, orography, frontal lift, turbulence), the 10 basic Cloud Genera, cloud classification by height, special cloud forms (Lenticular, Rotor, Castellanus, Mammatus), Cloud Ceiling vs Cloud Base, reporting coverage in oktas.

> **Primary References:** IC Joshi Chapter 8; Oxford/Nordian Chapter 8; Volare Question Bank.

• **Cloud:** Visible aggregate of minute water droplets, ice crystals, or a mixture of both, suspended in free air.

• **Condensation Nuclei (Aerosols):** Microscopic hygroscopic particles (salt, dust, combustion soot) essential for water vapour to condense upon.

• **Vertical Motion Mechanisms:**

1. **Thermal Convection:** Solar heating creates rising thermals **\implies** Cumuliform clouds (**Cu, Cb**).
2. **Orographic Lifting:** Wind forced upward by mountains **\implies** Cap clouds, Lenticular clouds, banner clouds.
3. **Frontal Uplift:** Warm air sliding up a cold air wedge **\implies** Extensive stratiform decks (**Cs, As, Ns**).
4. **Turbulence Mixing:** Surface friction mixing moist air in the boundary layer **\implies** Stratocumulus (**Sc**).

---

graph TD

	Altitude Level	Physical Composition
	High ($>20,000\text{ ft}$)	Ice crystals
	High ($>20,000\text{ ft}$)	Ice crystals
	High ($>20,000\text{ ft}$)	Ice crystals
	Medium ($6,500-20,000\text{ ft}$)	Supercooled water & ice
	Medium ($6,500-20,000\text{ ft}$)	Supercooled water & ice
	Low / Medium ($1,000-10,000\text{ ft}$)	Water droplets, raindrops, snow
	Low ($1,000-6,500\text{ ft}$)	Liquid water
	Low (Surface to $1,500\text{ ft}$)	Liquid water
	Vertical ($2,000-15,000\text{ ft}$)	Liquid water (ice at top)
	Great Vertical ($1,000\text{ to }>45,000\text{ ft}$)	Water droplets, supercooled drops, hail, ice

---

graph LR

LENT["Altocumulus lenticularis
Smooth lens-shaped
Marker for Mountain Waves & CAT"]

MAMM["Mammatus (Mammatocumulus)
Bag-like pouches hanging under Cb anvil
Extreme turbulence warning!"]

CAST["Altocumulus castellanus
Turreted castle-like towers
Instability aloft -> Afternoon T-storms"]

ROTOR["Rotor Clouds
Turbulent roll cloud under mountain wave crest
Deadly low-level violent eddies"]

- **`FEW`: 1 to 2 oktas** (Trace to **25\%**).
- **`SCT` (Scattered): 3 to 4 oktas** (**37\% to 50\%**).
- **`BKN` (Broken): 5 to 7 oktas** (**62\% to 87\%**).
- **`OVC` (Overcast): 8 oktas** (**100\%** full cover).

> **Cloud Ceiling:** The height above ground or water (AGL) of the base of the lowest layer of cloud that covers **more than half the sky** (i.e., **`BKN` (5-7 oktas)** or **`OVC` (8 oktas)**), or the vertical visibility in an obscured sky.

> *(Note: `FEW` and `SCT` layers DO NOT constitute a ceiling!)*

- **A:** Altostratus
- **B:** Cirrostratus
- **C:** Stratocumulus
- **D:** Nimbostratus

View Answer

Correct Answer: Option B (Cirrostratus)

Explanation: Halos are formed by refraction of light through hexagonal ice crystals found in thin Cirrostratus sheets. (Altostratus produces a watery sun without halos; Altocumulus produces coronas through diffraction).

- **A:** Nimbostratus
- **B:** Cirrus
- **C:** Altocumulus castellanus
- **D:** Cumulus humilis

View Answer

Correct Answer: Option A (Nimbostratus).

- **A:** Severe ground fog.
- **B:** Standing mountain waves and severe turbulence.
- **C:** Rapidly falling barometric pressure at sea.
- **D:** Approaching tropical hurricane.

View Answer

Correct Answer: Option B.

Chapter 09: Atmospheric Stability & Lapse Rates

> **DGCA Syllabus:** Atmospheric equilibrium (stable, unstable, neutral), comparison of Environmental Lapse Rate (ELR) with DALR and SALR, Absolute Stability, Absolute Instability, Conditional Instability, Latent Instability, convective condensation level, flight weather characteristics in stable vs unstable atmospheres.

> **Primary References:** IC Joshi Chapter 9; Oxford/Nordian Chapter 4; Volare Question Bank.

- **Atmospheric Stability:** The tendency of an air parcel displaced vertically to return to its original level (stable), accelerate away from its original level (unstable), or remain at its new displaced level (neutral).
- **The Core Rule of Buoyancy:**
- If a lifted parcel is **COLDER** (denser) than the ambient environment **\implies** It sinks back **\implies STABLE**.
- If a lifted parcel is **WARMER** (lighter) than the ambient environment **\implies** It continues to rise **\implies UNSTABLE**.
- If a lifted parcel is the **SAME TEMPERATURE** as the ambient environment **\implies NEUTRAL**.

graph TD

PARCEL["LIFTED AIR PARCEL"]

Parcel Colder than Environment
Parcel Warmer than Environment
Parcel Same Temp as Environment

The state of stability is determined solely by comparing the **Environmental Lapse Rate (ELR)** with the **Dry Adiabatic Lapse Rate (DALR = 3.0^°C/1000 ft)** and the **Saturated Adiabatic Lapse Rate (SALR ~ 1.5^°C/1000 ft)**.

graph LR

SUB["ELR < 1.5°C/1000 ft
(ABSOLUTE STABILITY)"] --- COND["1.5°C < ELR < 3.0°C
(CONDITIONAL INSTABILITY)"]

COND --- SUPER["ELR > 3.0°C/1000 ft
(ABSOLUTE INSTABILITY)"]

Stability State	Lapse Rate Condition	Behavior if Unsaturated (Dry)	Behavior if Saturated (Wet)	Associated Weather
Absolute Stability	$\mathbf{\text{ELR} < \text{SALR} < \text{DALR}}$ ($\text{ELR} < 1.5^\circ\text{C}/1000\text{ ft}$)	Stable (sinks back)	Stable (sinks back)	Stratus clouds, fog, smooth air, poor visibility, haze/smog
Absolute Instability	$\mathbf{\text{ELR} > \text{DALR} > \text{SALR}}$ ($\text{ELR} > 3.0^\circ\text{C}/1000\text{ ft}$)	**Unstable (rises)**	**Unstable (rises)**	Vigorous convection, Cumulus, Cb, severe turbulence, gusty winds
Conditional Instability	$\mathbf{\text{SALR} < \text{ELR} < \text{DALR}}$ ($1.5^\circ\text{C} < \text{ELR} < 3.0^\circ\text{C}/1000\text{ ft}$)	**STABLE** if unsaturated	**UNSTABLE** if saturated!	Stable until lifted to condensation level (LCL); once clouds form, latent heat triggers rapid instability!
Neutral Equilibrium (Dry)	$\mathbf{\text{ELR} = \text{DALR}}$ ($3.0^\circ\text{C}/1000\text{ ft}$)	Neutral	Unstable	Well-mixed boundary layer
Neutral Equilibrium (Saturated)	$\mathbf{\text{ELR} = \text{SALR}}$ ($\approx 1.5^\circ\text{C}/1000\text{ ft}$)	Stable	Neutral	Overcast nimbostratus
Temperature Inversion	$\mathbf{\text{ELR} < 0}$ (Temp rises with height)	Extremely Stable	Extremely Stable	Traps pollutants, ground fog, severe low-level windshear

---

This is one of the most heavily tested comparative themes in DGCA exams:

Weather Element	Stable Atmosphere	Unstable Atmosphere
Cloud Type	**Stratiform** (St, Sc, As, Ns) or cloudless	**Cumuliform** (Cu, Cb, TCU)
Airflow / Turbulence	**Smooth air**, laminar flow, negligible turbulence	**Turbulent**, choppy, strong thermals, updrafts/downdrafts
Visibility	**Poor** (smoke, haze, mist trapped near surface)	**Good / Excellent** outside showers (active vertical mixing)
Precipitation	**Continuous**, steady rain or drizzle (`RA`, `DZ`)	**Showers** (`SHRA`), intermittent, sudden onset, hail
Surface Winds	Steady, calm to light	Gusty, squally, variable direction
Altimeter Hazards	Temperature inversions create altimeter errors	Severe vertical windshear & microbursts

---

1. ■■ **Conditional Instability is the NORMAL state of the Troposphere:**

Standard atmosphere lapse rate is **1.98^°C/1000 ft**, which lies right between SALR (**1.5^°C**) and DALR (**3.0^°C**)!

2. ■■ **Superadiabatic Lapse Rate:**

When **ELR > 3.0^°C/1000 ft** (Absolute Instability), it is also termed *superadiabatic*. Confined strictly to the lowest few hundred feet over sunbaked desert ground.

- **A:** The ELR is greater than the DALR.
- **B:** The ELR is less than the SALR.
- **C:** The ELR is between the SALR and the DALR.
- **D:** The ELR is negative.

View Answer

Correct Answer: Option C (ELR between SALR and DALR)

Explanation: The parcel is stable when dry (displaced at DALR) but becomes buoyant and unstable once saturated (displaced at SALR).

- **A:** Smooth air, stratiform clouds, and poor visibility in haze.
- **B:** Turbulent air, cumuliform clouds, good visibility outside showers, and gusty surface winds.
- **C:** Continuous light drizzle and calm surface winds.
- **D:** Uniform fog and zero vertical air currents.

View Answer

Correct Answer: Option B.

Chapter 10: Optical & Electrical Phenomena

> **DGCA Syllabus:** Refraction, reflection, and diffraction in the atmosphere, Solar and Lunar Halos, Coronas, Rainbows, Glory (Brocken Spectre), Mirages (Superior vs Inferior), St. Elmo's Fire, static electricity hazards in flight.

> **Primary References:** IC Joshi Chapter 10; Oxford/Nordian Chapter 8; Volare Question Bank.

- **Atmospheric Optics:** Produced by interaction of sunlight or moonlight with atmospheric particles:
- **Refraction:** Bending of light rays passing through media of different densities or ice crystal prisms.
- **Reflection:** Bouncing of light off water droplet or ice crystal surfaces.
- **Diffraction:** Bending of light waves around the edges of minute liquid water droplets.
- **Scattering:** Rayleigh scattering by gas molecules (causes blue sky and red sunsets).

---

graph TD

OPT["OPTICAL PHENOMENA"]

OPT --> HALO["HALO (22° & 46° Ring)
Cloud: Cirrostratus (Cs)
Mechanism: Refraction through Ice Crystals"]

OPT --> CORONA["CORONA (Small concentric rings)
Cloud: Altocumulus (Ac) / Stratus
Mechanism: Diffraction around Water Droplets"]

OPT --> RAINBOW["RAINBOW (42° Arc)
Mechanism: Internal Reflection & Refraction in Raindrops
Requires Sun behind observer!"]

OPT --> GLORY["GLORY (Colored rings around shadow)
Seen around aircraft shadow on cloud tops
Diffraction of light by cloud droplets"]

OPT --> MIRAGE["MIRAGE (Superior & Inferior)
Extreme temperature gradients bending light"]

Phenomenon	Required Particle Type	Associated Cloud	Physical Mechanism	Visual Appearance & Exam Points
Halo	**Hexagonal Ice Crystals**	**Cirrostratus (`Cs`)**	**Refraction**	Large ring of 22° (most common) or 46° around Sun or Moon. Red inside, blue outside. **Strong indicator of approaching warm front!**
Corona	**Minute Water Droplets**	**Altocumulus (`Ac`)** or `Sc`	**Diffraction**	Small series of colored rings closely hugging the Sun or Moon. Blue inside, red outside.
Rainbow	**Liquid Raindrops**	Shower / Rain area	**Refraction + Internal Reflection**	Arc centered on anti-solar point (42° radius). Primary bow has red on outside, violet inside. Observer must have **Sun behind back**!
Glory	**Cloud Water Droplets**	Fog or Stratus tops	**Diffraction**	Colored rings seen directly around the shadow of the aircraft projected onto a cloud layer below.
Inferior Mirage	Hot surface air	None (Clear sunny desert/runway)	**Refraction (Upward bending)**	Looking down at hot ground, light from sky bends upward $\to$ appears like a pool of water on the runway/road.
Superior Mirage	Strong ground temperature inversion	Cold sea / Arctic surface	**Refraction (Downward bending)**	Looking toward cold horizon, light bends downward $\to$ objects appear elevated or floating in the sky above their true position.

- **St. Elmo's Fire:**
- A luminous, silent electrical brush discharge (corona discharge) emanating from sharp pointed extremities of the aircraft (windshield corners, nose cone, wingtips, propeller tips).
- Occurs when flying through highly charged atmospheres: **near thunderstorms (`Cb`), in heavy snow, or charged ice crystal clouds**.
- Emits a ghostly blue-violet or greenish glow, often accompanied by crackling static in aircraft communication radios ("precipitation static" / P-static).
- *Hazard Level:* Does **not** cause structural damage to the aircraft, but serves as an immediate warning of entering a high-voltage electrostatic field with potential impending **lightning strike**!

- **Static Discharge Wicks:**
- Trailing edge conductive braided cords that slowly bleed accumulated static charge back into the air stream to prevent radio blackout and spark ignition.

- **A:** Diffraction of light by water droplets in Altocumulus clouds.
- **B:** Refraction of light through hexagonal ice crystals in Cirrostratus clouds.
- **C:** Internal reflection of light in heavy rain showers.
- **D:** Scattering of light by smoke particles in an inversion.

View Answer

Correct Answer: Option B.

- **A:** Halos are caused by ice crystals in high clouds (refraction), while Coronas are caused by water droplets in medium clouds (diffraction).
- **B:** Halos are small, Coronas are large.
- **C:** Halos only occur at night, Coronas only occur during the day.
- **D:** Halos indicate fair weather, Coronas indicate tornadoes.

View Answer

Correct Answer: Option A.

Chapter 11: Precipitation & Mechanisms

> **DGCA Syllabus:** Physical processes of precipitation formation (Collision-Coalescence and Bergeron-Findeisen processes), Supercooled Liquid Water (SLW), precipitation forms (Rain, Drizzle, Freezing Rain, Snow, Sleet/Ice Pellets, Hail, Graupel), Virga, and associated flight hazards.

> **Primary References:** IC Joshi Chapter 11; Oxford/Nordian Chapter 9; Volare Question Bank.

- **Precipitation:** Any product of the condensation of atmospheric water vapour that falls under gravity from clouds and reaches the ground.
- **The Raindrop Problem:** Typical cloud droplets have diameters of **~ 0.02 mm** (**20 \mum**) and fall at only **0.01 m/s**. They evaporate long before hitting the ground. A raindrop (**1-2 mm**) contains the volume of roughly **one million cloud droplets**!

---

graph TD

GROWTH["PRECIPITATION GROWTH THEORIES"]

GROWTH --> WARM["1. Collision-Coalescence Theory
Warm clouds (Temp > 0°C)
Dominant in tropical & summer maritime clouds"]

GROWTH --> COLD["2. Bergeron-Findeisen Ice Crystal Theory
Cold / Mixed clouds (Temp < 0°C to -40°C)
Dominant in mid-latitudes & deep convective clouds"]

- Operates in clouds where the temperature is entirely above freezing (**>0^°C**).
- Larger hygroscopic droplets have higher terminal fall velocities than smaller droplets.
- As larger droplets fall through the cloud, they collide with and absorb smaller droplets (**coalescence**), sweeping up moisture until they grow heavy enough to fall out as rain.

• In the temperature zone between **0^°C and -40^°C**, clouds consist of a mixture of **ice crystals and Supercooled Liquid Water Droplets (SLWD)**.

• **Fundamental Physical Law: The Saturation Vapour Pressure over water is GREATER than the Saturation Vapour Pressure over ice!** (**e_s (water)} > e_s (ice)}**).

• **The Consequence:**

1. The air is simultaneously **undersaturated** relative to liquid water and **supersaturated** relative to ice.
2. Liquid water droplets continuously evaporate into water vapour.
3. Water vapour rapidly deposits directly onto the ice crystals by sublimation/deposition.
4. The ice crystals grow rapidly into large snowflakes at the expense of evaporating water droplets!
5. As they fall through the freezing level (**0^°C** isotherm), they melt into raindrops.

Precipitation Form	METAR Code	State & Size	Originating Cloud Type	Critical Aviation Flight Hazard
Drizzle	**`DZ`**	Liquid droplets **$< 0.5\text{ mm}$** diameter; falls slowly	**Stratus (`St`)** or Fog	Extremely low cloud ceiling and poor visibility.
Rain	**`RA`**	Liquid drops **$\ge 0.5\text{ mm}$** diameter	**Nimbostratus (`Ns`)** (continuous) or **`As`**	Decreased runway braking friction (aquaplaning / hydroplaning).
Showers	**`SH`**	Intermittent rain, snow, or hail	**Cumulus (`Cu`)** or **Cumulonimbus (`Cb`)**	Rapidly fluctuating visibility, sudden strong wind shifts.
Freezing Rain	**`FZRA`**	Supercooled liquid drops freezing instantly on impact	Warm air aloft over cold sub-zero surface wedge	**EXTREME DANGER!** Forms instantaneous heavy Clear / Glaze ice on airframe; can cause aerodynamic stall in minutes!
Snow	**`SN`**	White ice crystals in hexagonal branched structures	**Nimbostratus (`Ns`)** or `Cb`	Severe reduction in visibility; snow drifts on runway; airframe icing.
Ice Pellets (Sleet)	**`PL`**	Hard, transparent frozen raindrops ($<5\text{ mm}$)	Rain freezing as it falls through deep cold layer aloft	Positive warning of **Freezing Rain aloft**!
Hail	**`GR`**	Hard balls of ice **$\ge 5\text{ mm}$** up to $10+\text{ cm}$	Strictly **Cumulonimbus (`Cb`)**	Severe structural airframe damage, punctured radomes, shattered windshields.
Small Hail / Snow Pellets	**`GS`**	White, opaque, crisp ice grains ($<5\text{ mm}$)	`Cb` or `TCU`	Runway slippery; moderate turbulence.

• **Virga:** Precipitation falling from the base of a cloud that completely evaporates in dry sub-cloud air before reaching the ground.

• **Aviation Trap:** Evaporation is an endothermic process (absorbs latent heat from surrounding air).

• The air column under virga cools rapidly **\implies** Becomes significantly denser than ambient air **\implies** Accelerates violently downward.

• Generates devastating **Dry Microbursts** and low-level windshear with zero visual rain on the runway!

• **A:** The difference between the saturation vapour pressure over water and over ice.

• **B:** Strong thermal updrafts carrying water droplets upward.

• **C:** Friction between falling raindrops and aircraft wings.

- **D:** High air density at the cloud base.

View Answer

Correct Answer: Option A.

- **A:** The aircraft has entered an isothermal stratosphere layer.
- **B:** There is Freezing Rain (`FZRA`) and warmer air directly above the aircraft.
- **C:** The cloud is composed entirely of dry snow.
- **D:** A strong radiation inversion exists on the ground.

View Answer

Correct Answer: Option B (Freezing Rain exists aloft)

Explanation: Ice pellets form when rain from a warm layer aloft falls through a deep freezing layer near the ground and freezes solid before impact.

Chapter 12: Aircraft Icing & Accretion Hazards

> **DGCA Syllabus:** Conditions favorable for structural and induction icing, Supercooled Liquid Water (SLW), Supercooled Large Droplets (SLD), Clear/Glaze ice, Rime ice, Mixed ice, Hoar frost, aerodynamic penalties (lift reduction, drag increase, stall speed increase), Carburettor icing, Anti-icing vs De-icing systems.

> **Primary References:** IC Joshi Chapter 12; Oxford/Nordian Chapter 14; Volare Question Bank.

- **Structural Icing Conditions:** Requires TWO factors simultaneously:

1. **Visible Liquid Moisture** in the atmosphere (clouds, freezing rain, drizzle, or fog).
2. **Airframe Surface Temperature at or below 0^°C** (**0^°C to -40^°C**).

- **The Temperature Danger Zones:**
- **0^°C to -10^°C: MOST DANGEROUS!** High concentration of Supercooled Large Droplets (SLD) **\implies** Forms severe **Clear / Glaze Ice**!
- **-10^°C to -20^°C:** Mixture of supercooled droplets and ice crystals **\implies** Forms **Mixed Ice** and **Rime Ice**.
- **-20^°C to -40^°C:** Mostly small droplets **\implies** Light **Rime Ice**.
- **Below -40^°C:** Spontaneous freezing (homogenous nucleation) occurs **\implies** Clouds consist almost entirely of dry ice crystals **\implies** Ice crystals do not adhere to cold airframes; structural icing risk is negligible!

graph LR

ZONE1["0°C to -10°C
High SLD content
CLEAR / GLAZE ICE
(Severe Danger)"] --> ZONE2["-10°C to -20°C
Mixed SLD & Ice
MIXED / RIME ICE
(Moderate Danger)"]

ZONE2 --> ZONE3["-20°C to -40°C
Small droplets
LIGHT RIME ICE"]

ZONE3 --> ZONE4["Below -40°C
Pure ice crystals
NEGLIGIBLE ICING RISK"]

---

graph TD

ICE["STRUCTURAL ICING TYPES"]

Large Droplets (SLD)
Small Droplets

ICE --> HOAR["HOAR FROST
Sub-zero parked aircraft or rapid descent
Direct deposition of vapour to ice crystals"]

Ice Type	Temperature Range	Droplet Size & Cloud Type	Freezing Mechanism	Physical Appearance & Hazard Level
Clear Ice (Glaze)	**$0^\circ\text{C}\text{ to }-10^\circ\text{C}$**	**Large droplets (SLD)** in **`Cb`, `Cu`, `Ns`**, or Freezing Rain	**Freezes SLOWLY**. Droplet impacts leading edge, only a fraction freezes instantly; the remainder flows rearward over aerofoil before freezing.	Smooth, glassy, transparent, heavy, highly adhesive. **Most dangerous!** Distorts wing camber, destroys laminar airflow, difficult to dislodge with boots.
Rime Ice	**$-10^\circ\text{C}\text{ to }-20^\circ\text{C}$**	**Small droplets** in **`St`, `Sc`, `As`** (stratiform clouds)	**Freezes INSTANTANEOUSLY**. Droplet freezes on initial contact without spreading; traps tiny air bubbles inside.	Milky white, opaque, rough granular surface, brittle. Confined to leading edges; easily shed by de-icing boots.
Mixed Ice	**$-10^\circ\text{C}\text{ to }-15^\circ\text{C}$**	Mixture of small and large droplets in convective or deep stratiform	Simultaneous rapid and slow freezing.	Rough, grey, irregular mushroom-like horns on leading edges. High drag penalty.
Hoar Frost	Surface $<0^\circ\text{C}$	Water vapour (Clear air)	**Direct Deposition (Sublimation)**: Vapour turns directly into ice crystals on cold metal skin.	White crystalline feathery coating. Disables laminar flow; increases stall speed by $10-15\%$! **Must be removed before takeoff!**

---

graph TD

ACCRETION["ICE ACCRETION ON WINGS"]

ACCRETION --> LIFT["Lift: DECREASES by 30% to 50%!"]

ACCRETION --> DRAG["Drag: INCREASES by 100% to 300%!"]

ACCRETION --> WEIGHT["Weight: INCREASES (hundreds of kg)"]

ACCRETION --> STALL["STALL SPEED: INCREASES by 15 - 30 kt!"]

ACCRETION --> TAIL["Tailplane Stall: Ice on horizontal stabilizer causes violent uncommanded nose-down pitch!"]

> **Crucial Rule:** Even a thin layer of frost or ice equivalent to coarse sandpaper on the wing leading edge can reduce wing maximum lift by **30\%** and increase drag by **40\%**, causing aerodynamic stall at normal takeoff rotation speeds!

Carburettor icing can occur **IN CLEAR AIR (No Clouds!)** with outside air temperatures as high as **+25^°C to +30^°C** and high relative humidity (**>50\%**)!

1. **Fuel Vaporization Cooling:** Liquid avgas sprayed from the jet absorbs latent heat of vaporization, cooling the intake air by **~ 15^°C**.
2. **Venturi / Throttle Butterfly Expansion:** Pressure drop in the carburettor venturi and across the partially closed throttle valve causes adiabatic cooling of **~ 5^°C**.

3. **Total Temperature Drop:** Up to **20^°C** across the carburettor!

• If ambient OAT is **+15^°C**, the carburettor throat temperature drops to **-5^°C**.

• Any moisture in the air condenses and freezes solid on the throttle butterfly, choking fuel/air flow to the engine!

• **Indication of Carb Icing:** In fixed-pitch propeller aircraft **\to Drop in Engine RPM**; in constant-speed propeller aircraft **\to Drop in Manifold Pressure**.

• **Action:** Apply full Carburettor Heat. Initial drop in RPM will occur (hot air is less dense), followed by rough running as ice melts through the engine, then RPM recovers!

System Type	Operating Philosophy	Timing of Activation	Common Mechanisms
Anti-Icing	**Prevention**: Prevents ice from adhering to surfaces	Activated **BEFORE** entering icing conditions (based on OAT $\le +10^\circ\text{C}$ and visible moisture)	Hot bleed air (engine cowl & wing leading edges), electrical heating (pitot tubes, static ports, windshields), TKS fluid.
De-Icing	**Removal**: Removes ice after a thin layer has accreted	Activated **AFTER** a measurable accumulation of ice (e.g., $1/4\text{ to }1/2\text{ inch}$)	Pneumatic rubber boots (inflatable rubber bladders that crack and shed brittle ice).

• **A:** Slowly and do not spread out.

• **B:** Slowly and spread out.

• **C:** Rapidly and spread out.

• **D:** Rapidly and do not spread out.

View Answer

Correct Answer: Option D (Rapidly and do not spread out)

Explanation: Rime ice forms at cold temperatures (**-10^°C** to **-20^°C**) with small droplets. The latent heat is dissipated instantly, freezing the droplet immediately upon impact without allowing it to flow or spread out, trapping air within the structure.

• **A:** High humidity (RH > 50%) and ambient air temperatures between -5°C and +25°C.

• **B:** Zero humidity at temperatures below -40°C.

• **C:** Clear skies at FL 350 with OAT of -56°C.

• **D:** Full throttle takeoff roll on a cold dry winter runway.

View Answer

Correct Answer: Option A.

Chapter 13: Thunderstorms, Microbursts & Severe Weather

> **DGCA Syllabus:** Prerequisites for thunderstorm formation, the 3 life-cycle stages (Cumulus, Mature, Dissipating), Supercells, Multicell clusters, Squall lines, Microbursts and Low-Level Wind Shear (LLWS), Gust fronts and Roll clouds, Hail hazards, Lightning, Airborne weather radar detection, and in-flight avoidance techniques.

> **Primary References:** IC Joshi Chapter 13; Oxford/Nordian Chapter 16; Volare Question Bank.

A thunderstorm (`**Cb**`) cannot develop unless **ALL THREE** of the following conditions exist simultaneously:

graph TD

INGREDIENTS["THUNDERSTORM PREREQUISITES"]

INGREDIENTS --> M["1. HIGH MOISTURE CONTENT
High surface dew point (>18°C) in lower atmosphere"]

INGREDIENTS --> I["2. DEEP CONDITIONAL INSTABILITY
Steep ELR through a deep layer (up to tropopause)"]

INGREDIENTS --> L["3. TRIGGER / LIFTING MECHANISM
Forces moist parcel to its Level of Free Convection (LFC)"]

- **Thermal Convection:** Solar heating of dry land in summer (Air Mass / Heat Thunderstorms).
- **Orographic Lift:** Moist wind forced up mountain barriers (e.g., Western Ghats, Himalayas).
- **Frontal Lifting:** Cold dense air wedging beneath warm moist air (Cold Front Thunderstorms).
- **Convergence:** Intertropical Convergence Zone (ITCZ) or monsoon trough convergence.

A single-cell thunderstorm lasts roughly **45 to 60 minutes** and progresses through three distinct stages:

graph LR

CUMULUS["1. CUMULUS STAGE
Updrafts only (up to 3,000 ft/min)
Cloud grows to 20,000 ft
No precipitation yet"] --> MATURE["2. MATURE STAGE
Starts when RAIN HITS SURFACE!
Updrafts (6,000 ft/min) & Downdrafts coexist
Maximum turbulence, lightning, hail, microbursts"]

MATURE --> DISSIPATE["3. DISSIPATING STAGE
Downdrafts dominate entire cloud
Precipitation weakens
Anvil 'orphan anvil' spreads out"]

Stage	Typical Duration	Dominant Vertical Motion	Precipitation	Defining Diagnostic Features
1. Cumulus (Building)	$15\text{ to }20\text{ minutes}$	**Updrafts ONLY** (speeds $1,000-3,000\text{ ft/min}$)	None hitting ground	Rapid vertical growth of towering cumulus (`TCU`); droplets suspended by strong updrafts.
2. Mature	**$20\text{ to }30\text{ minutes}$**	**Updrafts AND Downdrafts coexist** side by side!	**Heavy rain, hail, or snow**	**Starts the precise moment rain reaches the ground!** Anvil top (`incus`) develops. Maximum severe turbulence, lightning, and microbursts.
3. Dissipating	$20\text{ to }30\text{ minutes}$	**Downdrafts ONLY** throughout cell	Rain gradually stops	Downdrafts starve the cell of warm moist inflow. Cloud top frays into fuzzy stratiform anvil.

A **Microburst** is an extremely intense, localized column of sinking air (downdraft) that hits the ground and fans out violently in all directions.

graph TD

DOWNDRAFT["INTENSE DOWNDRAFT (up to 6,000 ft/min)"] --> IMPACT["Hits Ground Surface"]

IMPACT --> OUTFLOW["Violent Radial Outflow (> 45 kt windshear)"]

- **Spatial Dimension:** Core diameter **< 4 km** (**2.2 NM**); outflow extends to **~ 5 km**.

• **Temporal Duration:** Peak intensity lasts **2 to 5 minutes**; total lifespan **< 15 minutes**.

• **Wind Shear Velocity Difference:** Across the core can exceed **45 to 90 knots**!

• **Vertical Downdraft Speed:** Up to **6,000 ft/min** (**30 m/s**).

1. **Entry (Increasing Headwind):** Aircraft encounters strong outflow headwind **\implies** Airspeed jumps **\implies** Aircraft balloons above glide slope. *Pilot reaction: reduces engine power and pushes nose down.*

2. **Core (Violent Downdraft):** Aircraft enters central downdraft **\implies** Severe sink rate.

3. **Exit (Abrupt Severe Tailwind):** Aircraft encounters tailwind outflow **\implies** Indicated airspeed drops catastrophically by **30-50 kt \implies** Loss of lift, aerodynamic stall, and ground impact before engines can spool up!

sequenceDiagram

participant AC as Aircraft on Approach

participant MB as Microburst Cell

AC->>MB: 1. Outflow Headwind: Airspeed Increases (+20 kt), Pitches Up

AC->>MB: 2. Core: Violent Downdraft (-4,000 ft/min)

AC->>MB: 3. Outflow Tailwind: Airspeed Plummets (-40 kt), Wing Stalls -> Crash!

---

1. **Distance Clearances (ICAO / FAA / DGCA Standards):**

• Below FL 200: Clear any severe storm echo by at least **10 to 15 NM**.

• Above FL 200: Clear by at least **20 NM**!

• Clear the **downwind side (under the anvil)** by at least **20 NM** because violent updrafts inside the Cb shoot large hail stones miles out through the anvil into apparently clear air!

2. **Overflying a Cb:**

• If overflying is unavoidable, clear the cloud top by at least **1,000 ft for every 10 knots of wind speed** at that level (minimum clearance **5,000 ft** above the top).

3. **Penetration Rules (If Trapped inside Cb):**

• **Do NOT turn back** (structural stress is doubled in a turn, and you may fly into the core).

• Set **Turbulence Penetration Speed (V_B / V_{RA})**.

• Maintain a **constant pitch attitude**; do NOT fight altitude variations (let the altimeter drift, fighting altitude stresses the wings!).

• Turn cockpit lights to **FULL BRIGHT** to prevent temporary blindness from lightning flash.

• **A:** The cloud first reaches the freezing level.

• **B:** Rain or precipitation first reaches the ground surface.

• **C:** Downdrafts completely replace updrafts.

• **D:** Lightning is first recorded on ADF.

View Answer

Correct Answer: Option B (Precipitation reaches the surface)

Explanation: The onset of precipitation at the surface marks the development of the precipitation-induced downdraft alongside the existing updraft, which defines the Mature stage.

• **A:** Severe carburettor icing.

• **B:** A sudden loss of airspeed and lift caused by transitioning rapidly from a strong headwind into a severe tailwind.

• **C:** Compass deviation caused by static discharge.

• **D:** High barometric pressure causing altimeter under-reading.

View Answer

Correct Answer: Option B.

Chapter 14: Air Masses, Fronts & Frontal Depressions

> **DGCA Syllabus:** Air mass classification (cT, cP, cA, mT, mP), Fronts (Warm front, Cold front, Stationary front, Occlusions), cloud sequences, weather changes across frontal boundaries, Polar front depression life-cycle, Western Disturbances (origin, track, and weather over India).

> **Primary References:** IC Joshi Chapter 14; Oxford/Nordian Chapters 10 & 11; Volare Question Bank.

- **Air Mass:** An immense body of air, horizontally uniform in temperature and moisture characteristics over thousands of kilometers.
- **Source Region:** Large, flat, uniform surface of the Earth (ocean, ice sheet, desert) with light winds where air remains stagnant for days acquiring surface properties.
- **Classification Code:**
- Moisture: **`c`** (Continental = Dry) vs **`m`** (Maritime = Moist).
- Temperature: **`P`** (Polar = Cold), **`T`** (Tropical = Warm), **`A`** (Arctic = Frigid), **`E`** (Equatorial = Very Warm & Humid).

---

graph TD

FRONTS["FRONTAL TYPES"]

Element	Warm Front Passage	Cold Front Passage
Slope	**Gentle: $1:150$** (Warm air slides up smoothly)	**Steep: $1:50\text{ to }1:80$** (Cold air acts like a bulldozer)
Speed	Slower ($10-20\text{ kt}$)	Faster ($20-35\text{ kt}$)
Cloud Sequence Ahead	**$\text{Ci} \to \text{Cs} \to \text{As} \to \text{Ns}$** (Precursor 600 NM ahead!)	Cumulus, Towering Cumulus, **Cumulonimbus (`Cb`)**
Precipitation	**Continuous steady rain or drizzle** over wide band ($200-400\text{ NM}$)	**Intense showers, hail, lightning**, narrow squall band ($20-50\text{ NM}$)
Pressure Trend	Falls steadily ahead $\to$ Levels off in warm sector	Falls ahead $\to$ **Rises abruptly (sharp pressure kick!)** behind
Wind Shift (NH)	**Veers** (e.g., $180^\circ \to 240^\circ$)	**Veers sharply** (e.g., $240^\circ \to 320^\circ$) with violent gusts
Temperature	Rises after passage	**Drops sharply** ($5-15^\circ\text{C}$ drop)
Dew Point	Rises	Drops sharply
Visibility	Poor (mist, fog, steady rain)	Very poor during front $\to$ **Exceptional / crystal clear** behind!

Because cold fronts move significantly faster than warm fronts (**2 x** speed), the cold front eventually catches up with the warm front, lifting the entire warm sector off the ground!

graph TD

OCC["OCCLUDED FRONT"]

OCC --> COLD_OCC["Cold Occlusion
Air behind cold front is COLDER than air ahead of warm front
Behaves like a cold front"]

OCC --> WARM_OCC["Warm Occlusion
Air behind cold front is WARMER than frigid air ahead of warm front
Behaves like a warm front"]

> **Western Disturbances (WD):** Shallow extra-tropical cyclonic depressions that originate over the **Mediterranean Sea, Black Sea, or Caspian Sea**, embedded in the **Subtropical Westerly Jet Stream**.

graph LR

MED["Origin: Mediterranean Sea / Caspian Sea"] --> TRACK["Steered Eastward by Subtropical Jet across Iran & Afghanistan"]

TRACK --> INDIA["Enters NW India: J&K;, Himachal, Punjab, Haryana, Rajasthan, UP"]

1. **Winter Precipitation (December to March):**

• Brings moderate to heavy **snowfall** over Jammu & Kashmir, Ladakh, Himachal Pradesh, and Uttarakhand.

• Brings vital winter rains (**Mawat**) to Punjab, Haryana, Rajasthan, and Western UP, essential for the *Rabi* (wheat) crop.

2. **Dense Radiation Fog Hazard:**

• Rain from a passing WD moistens the soil.

• Clear skies and light winds immediately behind the WD trigger **widespread, severe radiation fog** across Delhi, Punjab, and Uttar Pradesh, causing catastrophic flight disruptions at Delhi (VIDP).

3. **Cold Wave:** Behind the WD, frigid northerly winds from the snow-covered Himalayas cause severe **Cold Wave conditions** across Northern and Central India.

• **A:** Cb followed by St and then Ci.

• **B:** Cirrus, Cirrostratus, Altostratus, and Nimbostratus.

• **C:** Stratocumulus, Cumulus, and Altocumulus.

• **D:** Stratus, Fog, and Clear skies.

View Answer

Correct Answer: Option B (Ci -> Cs -> As -> Ns).

• **A:** Warm sea breezes from the Bay of Bengal.

• **B:** Soil moistening by WD rainfall followed by clear skies and radiative nighttime cooling in the high-pressure ridge behind the system.

• **C:** Katabatic winds blowing directly from the Indian Ocean.

• **D:** Industrial smoke trapping without moisture.

View Answer

Correct Answer: Option B.

Chapter 15: Jet Streams & Clear Air Turbulence (CAT)

> **DGCA Syllabus:** Definition and dimensions of Jet Streams, Polar Front Jet (PFJ), Subtropical Westerly Jet (STJ), Tropical Easterly Jet (TEJ), Somali Low-Level Jet, core location relative to tropopause, Clear Air Turbulence (CAT) causes, location of maximum windshear, and pilot avoidance/escape procedures.

> **Primary References:** IC Joshi Chapter 15; Oxford/Nordian Chapter 12; Volare Question Bank.

• **WMO Definition:** A strong, narrow, quasi-horizontal current of wind, concentrated along a quasi-horizontal axis in the upper troposphere or stratosphere, characterized by strong vertical and lateral wind shears and featuring one or more velocity maxima.

• **Minimum Speed Threshold:** The wind speed along the jet axis must be **at least 60 knots** (**110 km/h ~ 30 m/s**).

• **Standard Dimensions:**

- **Length:** Thousands of kilometers (**1,500 to 5,000 km**).
- **Width:** Hundreds of kilometers (**100 to 500 km**).
- **Depth (Thickness):** Several kilometers (**1 to 5 km**).

graph TD

CORE["JET STREAM CORE (Max Speed > 100 - 200 kt)"]

CORE --> CYCLONIC["Cold Air Side (Cyclonic Shear Side)
MAXIMUM WIND SHEAR & CAT!"]

CORE --> ANTICYCLONIC["Warm Air Side (Anticyclonic Shear Side)
Wider, moderate wind shear"]

CORE --> LOC["Core Location:
Located in the WARM air mass
Just BELOW the warm-air tropopause break"]

Jet Stream	Direction	Typical Flight Level	Seasonal Behavior	Key Aviation & Indian Climatology Significance
Polar Front Jet (PFJ)	**Westerly** (270°)	**FL 300 to FL 350** ($9-10\text{ km} \approx 300\text{ hPa}$)	Strongest in winter ($150-250\text{ kt}$); shifts equatorward	Associated with polar front; severe CAT.
Subtropical Jet (STJ)	**Westerly** (270°)	**FL 400** ($12\text{ km} \approx 200\text{ hPa}$)	Highly persistent in winter over North India ($25^\circ-30^\circ\text{N}$)	Steers Western Disturbances into India. Disappears/shifts north of Himalayas in summer.
Tropical Easterly Jet (TEJ)	**EASTERLY** (090°)	**FL 450 to FL 500** ($14-16\text{ km} \approx 100-150\text{ hPa}$)	**Summer ONLY** (June to September)	Runs from South China Sea across Peninsular India ($12^\circ-15^\circ\text{N}$) to Africa. **Essential driver of Indian Southwest Monsoon!**
Somali Jet (LLJ)	**South-Westerly**	**850 hPa** ($1.5\text{ km} \approx 5,000\text{ ft}$)	**Summer ONLY** (SW Monsoon)	Low-level jet blowing from Somali coast across Arabian Sea; feeds intense moisture into Western Ghats.

Clear Air Turbulence (CAT) is high-altitude turbulence occurring in cloud-free air, without visual meteorological cues.

- **Cold Air / Cyclonic Side of Jet Core:** Horizontal wind shear is much stronger on the polar/cold side of the jet stream core than on the equatorial/warm side.
- **Directly above and below the Jet Core:** Where vertical wind shear is maximum (**>6 kt** per **1,000 ft**).
- **Tropopause Breaks:** Around sharp folds where polar and subtropical air masses meet.
- **Mountain Wave Interaction:** Where jet streams cross mountain ranges perpendicular to the ridge.

1. **Vertical Escape is Fastest:**

Because vertical wind shear (**>6-10 kt/1,000 ft**) is thousands of times sharper than horizontal shear, a climb or descent of **2,000 to 4,000 ft** exits the turbulent shear layer faster than flying horizontally!

2. **Speed Management:**

Immediately reduce airspeed to recommended **Turbulence Penetration Speed (V_B / V_{RA})**.

3. **Temperature Trend Rule:**

- Flying across a jet stream, watch the outside air temperature (OAT) gauge:
- If temperature is **falling rapidly \implies** You are penetrating into the cold (cyclonic) side **\implies Expect severe CAT!**

- If temperature stops falling and levels off **\implies** You have crossed the tropopause boundary.

- **A:** In the warm air mass.
- **B:** In the cold air mass.
- **C:** Just above the warm-air tropopause.
- **D:** Just below the cold-air tropopause.

View Answer

Correct Answer: Option A (In the warm air mass)

Explanation: The jet stream core lies in the warm air mass, in the upper troposphere, just below the tropopause break on the warm side of the polar front.

- **A:** Blows westerly across Central Europe in winter.
- **B:** Blows easterly across Peninsular India at FL 450-500 during the Southwest Monsoon summer months.
- **C:** Operates exclusively at sea level over Antarctica.
- **D:** Dissipates the Indian monsoon rains.

View Answer

Correct Answer: Option B.

Chapter 16: Mountain Waves & Orographic Hazards

> **DGCA Syllabus:** Conditions required for Mountain Wave (Standing Lee Wave) formation, wave anatomy (Cap cloud, Lenticular clouds, Rotor clouds), hazards (severe downdrafts, altimeter errors, turbulence), and mountain flying rules of thumb.

> **Primary References:** IC Joshi Chapter 17; Oxford/Nordian Chapter 18; Volare Question Bank.

---

graph TD

COND["MOUNTAIN WAVE PREREQUISITES"]

COND --> W1["1. Wind Direction Perpendicular to Ridge
Within 30° of normal to mountain barrier"]

COND --> W2["2. Wind Speed ≥ 20 - 25 kt at Ridge Crest
Must increase or remain constant with height"]

COND --> S3["3. Stable Layer Sandwich
Stable layer near ridge height, capped by less stable air aloft"]

---

graph LR

WIND["Windward Flow (≥ 25 kt)"] --> RIDGE["Mountain Ridge"]

RIDGE --> CAP["1. Cap Cloud (Föhn Wall)
Stationary over crest"]

RIDGE --> WAVE["Standing Lee Waves (Wavelength 5 - 20 NM)"]

WAVE --> LENT["2. Lenticular Clouds (Ac lenticularis)
Stationary lens clouds in wave crests aloft"]

WAVE --> ROTOR["3. Rotor Clouds (Roll Clouds)
Violent turbulent vortex below wave crest"]

WAVE --> DOWN["Severe Downdrafts (2,000 - 5,000 ft/min)
On downwind/leeward slope!"]

1. **Cap Cloud (Föhn Wall):** Sits directly over the mountain ridge. Air rises up the windward slope, condenses at the crest, then plunges down the leeward slope, dissipating by adiabatic warming.

2. **Lenticular Clouds (`Ac lenticularis`):** Smooth, lens-shaped clouds formed at the crest of each standing wave downwind. **They appear stationary even in winds >60 kt** because water continuously condenses at the upwind edge and evaporates at the downwind edge!

3. **Rotor Clouds:** Violent cylindrical tumbling roll clouds formed beneath the wave crests near ridge level. The most dangerous turbulence zone in the atmosphere!

---

1. **Catastrophic Downdrafts:**

• Sinking air on the leeward slope can exceed **3,000 to 5,000 ft/min**, far exceeding the maximum climb capability of most transport aircraft!

2. **Severe Altimeter Over-Reading (Venturi Effect):**

• High-speed airflow squeezing over the mountain crest accelerates (Bernoulli's principle), creating a localized **pressure drop**.

• An altimeter senses this low pressure and assumes the aircraft is higher than it really is.

• Combined with cold air column error, the altimeter can **over-read by 1,000 to 2,000 ft \implies** The pilot flies into the mountain in IMC!

3. **Severe Structural Turbulence in Rotors:**

• Extreme vertical gust loads can cause structural airframe failure.

• **Crossing Angle:** Cross ridges at a **45° angle**, never at **90°** (perpendicular). A **45°** approach allows a quick **90°** turn away from the mountain if unmanageable downdrafts are encountered!

• **Terrain Clearance:** Maintain a minimum terrain clearance of at least **3,000 to 5,000 ft** above the highest ridge in mountain wave conditions.

• **Flight Path:** Never fly up the center of a narrow mountain valley; hug the upwind (lift) side of the valley to preserve turnaround room.

• **A:** Calm surface winds, unstable air mass, and dense cumulonimbus clouds.

• **B:** Strong wind (≥ 20 kt) blowing roughly perpendicular to a ridge, wind increasing with height, and a marked stable layer near crest level.

• **C:** Gale force winds blowing parallel to the mountain valley in summer.

• **D:** Strong frontal inversion in flat terrain.

View Answer

Correct Answer: Option B.

• **A:** The altimeter under-reads, indicating lower than true altitude.

• **B:** The altimeter over-reads due to dynamic pressure drop (Bernoulli effect), causing the aircraft to fly dangerously closer to terrain than indicated.

• **C:** The altimeter indicates zero.

• **D:** The error depends solely on humidity.

View Answer

Correct Answer: Option B.

Chapter 17: Tropical Meteorology & Cyclones

> **DGCA Syllabus:** Tropical weather characteristics, Intertropical Convergence Zone (ITCZ), Easterly Waves, Tropical Revolving Storms (TRS / Cyclones), genesis requirements, structural anatomy (Eye, Eyewall, Spiral bands), IMD storm classification in North Indian Ocean, Dangerous vs Navigable semicircle.

> **Primary References:** IC Joshi Chapter 18; Oxford/Nordian Chapter 17; Volare Question Bank.

- **Latitude Band:** Between **30^°N** and **30^°S**.
- **Weak Pressure Gradients:** Horizontal pressure differences are very small; isobaric analysis is generally replaced by **Streamline and Isotach analysis**.
- **Negligible Coriolis Force near the Equator:** Geostrophic balance breaks down within **5°** to **10°** of the Equator. Wind blows almost directly down the pressure gradient.
- **Diurnal Variation Dominates:** Daily cycle of solar heating and nocturnal cooling drives almost all local weather (land/sea breezes, afternoon thunderstorms).
- **Intertropical Convergence Zone (ITCZ):**
- The equatorial trough where the Trade Winds of both hemispheres converge (NE Trades meeting SE Trades).
- Characterized by sluggish, calm winds (**The Doldrums**), towering cumulonimbus lines, heavy convective rain, and severe turbulence.
- Migrates north during the Northern Hemisphere summer (reaching **20°-25^°N** over India as the Monsoon Trough) and south during winter.

A Tropical Revolving Storm is an intense cyclonic vortex originating over warm tropical oceans.

graph TD

GENESIS["TRS / CYCLONE GENESIS REQUIREMENTS"]

GENESIS --> SST["1. Warm Sea Surface Temp (SST) ≥ 26.5°C
To a depth of at least 50 meters (Thermal Engine)"]

GENESIS --> COR["2. Latitude ≥ 5° from Equator (5° to 15°)
Coriolis force is required to initiate spin
(NEVER forms between 0° and 5°!)"]

GENESIS --> SHEAR["3. Low Vertical Wind Shear (< 10-15 kt)
Strong shear tears vertical chimney apart"]

GENESIS --> MOIST["4. High Mid-Tropospheric Humidity
Prevents dry air entrainment"]

GENESIS --> TRIGGER["5. Pre-Existing Low-Level Disturbance
Easterly Wave or monsoon trough vortex"]

Ocean Basin	Local Name
Bay of Bengal & Arabian Sea (North Indian Ocean)	**Cyclonic Storm (Cyclone)**
North Atlantic Ocean & Northeast Pacific	**Hurricane**
Northwest Pacific Ocean (China, Japan, Philippines)	**Typhoon**
South Indian Ocean & Southwest Pacific (Australia)	**Tropical Cyclone (formerly Willy-Willy)**

---

graph LR

SPIRAL1["Outer Spiral Rainbands
Squalls, Cu/Cb, gusty winds"] --> EYEWALL["EYEWALL (Most Violent!)

Wall of towering Cb to 50,000 ft
Max winds (>100 kt), torrential rain, extreme turbulence"]

EYE --> EYEWALL2["Rear Eyewall
Immediate sudden resumption of hurricane winds from opposite direction!"]

1. **The Eye:**

• Central core, diameter typically **20 to 50 km** (**10-30 NM**).

• **Calm or light winds (<10 kt)**, descending/subsiding warm air, clear skies or scattered low stratus, zero precipitation.

• Lowest barometric pressure (frequently drops below **950 hPa**, record lows **<880 hPa**).

2. **The Eyewall:**

• Dense circular ring of towering cumulonimbus clouds surrounding the eye, extending up to **50,000-60,000 ft**.

• **Location of maximum sustained wind speeds**, highest storm surge, torrential rain, and extreme turbulence.

3. **Spiral Feeder Rainbands:**

• Long curved bands of convective showers spiraling inward toward the center across hundreds of kilometers.

When facing in the direction of the cyclone's movement (track):

graph TD

TRACK["CYCLONE MOVING FORWARD (NH)"]

TRACK --> RIGHT["RIGHT-HAND SIDE: DANGEROUS SEMICIRCLE
1. Wind speed + Storm forward speed add up (Maximum Winds!)
2. Wind direction blows aircraft/ships directly toward storm track!"]

TRACK --> LEFT["LEFT-HAND SIDE: NAVIGABLE SEMICIRCLE
1. Storm forward speed subtracts from wind speed
2. Wind blows aircraft/ships away from storm path"]

The India Meteorological Department (IMD) classifies cyclonic disturbances by maximum sustained **3**-minute surface wind speed:

Category	Sustained Wind Speed (Knots)	Wind Speed (km/h)	Pressure Deficit
Low Pressure Area	$< 17\text{ kt}$	$< 31\text{ km/h}$	$< 1\text{ hPa}$
Depression (`D`)	**$17\text{ to }27\text{ kt}$**	$31\text{ to }49\text{ km/h}$	$1-3\text{ hPa}$
Deep Depression (`DD`)	**$28\text{ to }33\text{ kt}$**	$50\text{ to }61\text{ km/h}$	$3-4.5\text{ hPa}$
Cyclonic Storm (`CS`)	**$34\text{ to }47\text{ kt}$**	$62\text{ to }88\text{ km/h}$	$4.5-8.5\text{ hPa}$
Severe Cyclonic Storm (`SCS`)	**$48\text{ to }63\text{ kt}$**	$89\text{ to }117\text{ km/h}$	$8.5-15.5\text{ hPa}$
Very Severe Cyclonic Storm (`VSCS`)	**$64\text{ to }89\text{ kt}$**	$118\text{ to }166\text{ km/h}$	$15.5-31.5\text{ hPa}$
Extremely Severe Cyclonic Storm (`ESCS`)	**$90\text{ to }119\text{ kt}$**	$167\text{ to }221\text{ km/h}$	$31.5-65.5\text{ hPa}$
Super Cyclonic Storm (`SuCS`)	**$\ge 120\text{ kt}$**	**$\ge 222\text{ km/h}$**	$> 65.5\text{ hPa}$

- **A:** The sea surface temperature is too cold.
- **B:** The Coriolis force is zero or negligible near the equator, which prevents cyclonic rotation from initiating.
- **C:** The trade winds blow too fiercely.
- **D:** High vertical wind shear always exists at the equator.

View Answer

Correct Answer: Option B (Coriolis force is negligible).

- **A:** Maximum winds and extreme hail.
- **B:** Light winds, subsiding air, little or no cloud, and absence of precipitation.
- **C:** Violent low-level squalls and dense fog.
- **D:** Continuous towering cumulonimbus with severe icing.

View Answer

Correct Answer: Option B.

Chapter 18: Climatology of India & Monsoons

> **DGCA Syllabus:** Climatological seasons of India (Winter, Pre-Monsoon, SW Monsoon, Post-Monsoon), Southwest Monsoon mechanism, Somali Jet, Tropical Easterly Jet, Tibetan High, Monsoon Trough, Breaks in Monsoon, Northeast Monsoon, Pre-monsoon severe storms (Nor'westers/Kalbaishakhi, Andhi, Loo), Western Disturbances, and Indo-Gangetic winter fog.

> **Primary References:** IC Joshi Chapter 19; Volare Question Bank.

---

1. **Winter (Cold Weather Season):** January to February.
2. **Pre-Monsoon (Hot Weather / Summer Season):** March to May.
3. **Southwest Monsoon (Rainy Season):** June to September.
4. **Post-Monsoon (Retreating Monsoon Season):** October to December.

The Southwest Monsoon accounts for **over 75\%** of India's total annual rainfall.

graph TD

MONSOON["DRIVING ENGINES OF SW MONSOON"]

MONSOON --> TIBET["1. Tibetan Plateau Heating Engine
High-altitude thermal engine -> Upper-level Tibetan High at 200 hPa"]

MONSOON --> TEJ["2. Tropical Easterly Jet (TEJ)
FL 450 - 500 (100 - 150 hPa)
Runs from S. China Sea -> Peninsular India -> Africa"]

MONSOON --> SOMALI["3. Somali Low-Level Jet (LLJ)
850 hPa (1.5 km)
Cross-equatorial flow feeding moisture into Western Ghats"]

MONSOON --> TROUGH["4. Monsoon Trough (Shifted ITCZ)
Extends from Ganganagar (Rajasthan) to Head Bay of Bengal"]

- **Onset Date:** Standard normal onset date over **Kerala is 1st June** (Car Nicobar **~** 15th-20th May).
- Advances northward in two major branches:

1. **Arabian Sea Branch:** Hits the steep barrier of the **Western Ghats**, producing torrential orographic rain on the windward coastal belt (Kerala, Karnataka, Goa, Konkan/Mumbai).

2. **Bay of Bengal Branch:** Moves into Bangladesh and Myanmar, gets deflected by the Arakan mountains and Himalayas, and travels westward along the Gangetic plains.

- Covers the entire Indian subcontinent by **~ 8th July**.

- Form at the head of the **Bay of Bengal** (near **20^°N**).

- Move in a **West-Northwesterly (WNW)** direction along the axis of the Monsoon Trough across Central India.
- Cause widespread, heavy, prolonged rainfall in Odisha, West Bengal, MP, Chhattisgarh, and Gujarat.

graph LR

NORMAL["NORMAL MONSOON:
Monsoon Trough over Central Plains
Widespread rain across India"] --> SHIFT["TROUGH SHIFTS NORTHWARD
Hugs the Himalayan Foothills"]

SHIFT --> BREAK["BREAK IN MONSOON:
1. Heavy rain & flash floods in Himalayan foothills, Assam, Bihar
2. Severe drought & clear dry skies over Plains of India!"]

- Occurs in **July or August**, typically lasting from a few days to 2 weeks.
- The axis of the monsoon trough shifts from its normal position to the **foothills of the Himalayas**.
- **Rainfall pattern reverses completely:**
- Plains of India experience **dry, rainless, hot weather**.
- Himalayan foothills, Assam, and Bihar experience **catastrophic cloudbursts and flooding**.

- As the Asian continent cools, high pressure builds over Siberia and North India.
- The ITCZ retreats southward back toward the equator.
- Wind direction reverses: blows from **Northeast to Southwest**.
- Picks up moisture over the warm waters of the **Bay of Bengal**.
- Brings major rainfall (**50-60\%** of their annual total) to **Tamil Nadu, Coastal Andhra Pradesh, and Rayalaseema**.
- **Peak Season for Destructive Cyclones:** October and November are the most active cyclone months in the Bay of Bengal and Arabian Sea.

During the scorching summer months, intense solar insolation interacting with dry continental and moist maritime air triggers violent local convective storms:

Local Name	Region Affected	Defining Meteorological Characteristics	Aviation Hazard
Nor'westers (Kalbaishakhi)	**West Bengal, Odisha, Assam, Bihar**	Destructive line of severe Cb thunderstorms moving from the **Northwest**. Accompanied by hurricane-force gusts, torrential rain, and giant hail.	Severe low-level windshear, microbursts, destroyed runways, hail damage.
Andhi (Dust Storm)	**Rajasthan, Punjab, Haryana, Delhi, Western UP**	Convective downdrafts from dry thunderstorms kicking up choking walls of sand and dust. Often has **no rain reaching ground**!	Visibility drops to **near ZERO** in seconds; engine ingestion of abrasive dust.
Loo	**Indo-Gangetic Plains (NW & Central India)**	Searing, blisteringly hot and dry westerly wind ($>45^\circ\text{C}$ to 50°C), blowing during afternoons of May and June.	Extreme density altitude; drastic reduction in engine thrust and wing lift.

- **A:** The monsoon winds reverse to northeasterly throughout the entire troposphere.
- **B:** The monsoon trough moves north to the Himalayan foothills, causing heavy rainfall in the foothills and drought-like dry conditions over the central plains.
- **C:** Rainfall ceases completely over the entire Indian subcontinent and Sri Lanka.

- **D:** Tropical cyclones form simultaneously over Delhi and Nagpur.

View Answer

Correct Answer: Option B.

- **A:** Peak winter (January) in Punjab.
- **B:** Pre-monsoon season (March to May) in West Bengal, Odisha, and Assam.
- **C:** Northeast monsoon (November) in Tamil Nadu.
- **D:** Post-monsoon season in Gujarat.

View Answer

Correct Answer: Option B.

Chapter 19: General Planetary Circulation

> **DGCA Syllabus:** Tri-cellular planetary circulation model (Hadley, Ferrel, and Polar cells), global pressure belts (Equatorial Low/ITCZ, Subtropical Highs/Horse Latitudes, Subpolar Lows, Polar Highs), global wind systems (Trade Winds, Prevailing Westerlies, Polar Easterlies), seasonal shifts of wind and pressure belts.

> **Primary References:** IC Joshi Chapter 20; Oxford/Nordian Chapter 13; Volare Question Bank.

Because the Earth rotates on its axis and has Coriolis force, the simple single-cell equator-to-pole circulation breaks down into **three distinct cells** in each hemisphere:

graph TD

subgraph Northern Hemisphere Tri-Cellular Model

NP["North Pole (90°N) - Frigid sinking air -> POLAR HIGH"]

POLAR_CELL["POLAR CELL (60°N to 90°N)
Polar Easterlies blow SW"]

POLAR_FRONT["Subpolar Low (60°N) -> POLAR FRONT (Convergence & Lift)"]

FERREL_CELL["FERREL CELL (30°N to 60°N)
Prevailing Westerlies blow NE"]

HORSE["Horse Latitudes (30°N) -> SUBTROPICAL HIGH (Sinking air)"]

HADLEY_CELL["HADLEY CELL (0° to 30°N)
Northeast Trade Winds blow SW"]

EQ["Equator (0°) -> ITCZ / DOLDRUMS (Convective Lift)"]

NP --> POLAR_CELL --> POLAR_FRONT --> FERREL_CELL --> HORSE --> HADLEY_CELL --> EQ

end

Cell	Latitude Band	Driving Mechanism	Dominant Surface Winds	Characteristic Weather
Hadley Cell	**$0^\circ\text{ to }30^\circ$**	**Thermally Direct** (Warm air rises at Equator, travels poleward aloft, sinks at 30°)	**Trade Winds** (NE in NH, SE in SH)	Convective squalls and thunderstorms at ITCZ; clear skies and deserts at 30°.
Ferrel Cell	**$30^\circ\text{ to }60^\circ$**	**Thermally Indirect** (Forced like a mechanical gear between Hadley and Polar cells)	**Prevailing Westerlies** (SW in NH, NW in SH)	Unsettled, dynamic frontal depressions, mobile lows and highs, variable weather.

Polar Cell	**$60^\circ\text{ to }90^\circ$**	**Thermally Direct** (Frigid dense air sinks at poles, spreads equatorward, rises at polar front)	**Polar Easterlies** (NE in NH, SE in SH)	Frigid, dry, light precipitation, cold blizzard winds.

---

graph LR

Polar Easterlies
Prevailing Westerlies
NE Trade Winds
SE Trade Winds
Roaring Forties Westerlies
Polar Easterlies

1. **The Doldrums (Equatorial Trough / ITCZ):**

• Region between **5^°N** and **5^°S** characterized by low pressure, light and baffling calm winds, high humidity, and squally afternoon cumulonimbus storms.

2. **Horse Latitudes (Subtropical Highs):**

• High pressure belts at **~ 30^°N** and **30^°S**.

• Characterized by calm winds, subsiding dry air, clear skies, and lack of rain. Location of the world's major hot deserts (Sahara, Arabian, Thar, Atacama).

3. **The Trade Winds:**

• Steady, persistent winds blowing from the Subtropical Highs toward the Equatorial Trough.

• **Northeast Trades** in the Northern Hemisphere; **Southeast Trades** in the Southern Hemisphere.

• Most consistent winds on Earth (**15-20 kt**).

4. **Roaring Forties & Furious Fifties:**

• In the Southern Hemisphere (**40^°S to 60^°S**), there is almost no land barrier.

• The Westerlies blow uninterrupted across open ocean with gale force and massive swell.

• As the thermal equator shifts with the Sun's declination:

• In **July (Northern Summer)**: Entire planetary wind system shifts **North** by **5° to 15°**. (Draws SE trade winds across the equator into the Arabian Sea **\implies** deflected by Coriolis force to become the Indian SW Monsoon!).

• In **January (Northern Winter)**: Entire planetary wind system shifts **South** by **5° to 15°**.

• **A:** SE trade winds.

• **B:** NE trade winds.

• **C:** Prevailing westerlies.

• **D:** Polar easterlies.

View Answer

Correct Answer: Option B (NE trade winds).

• **A:** Continuous heavy tropical rain and cyclonic gales.

• **B:** Subtropical high pressure, subsiding air, light winds, and fine settled weather.

• **C:** The lowest barometric pressures on Earth.

• **D:** Violent cold front thunderstorms.

View Answer

Correct Answer: Option B.

Chapter 20: Aviation Met Services, SIGMET & Flight Briefing

> **DGCA Syllabus:** Meteorological organization for international air navigation, World Area Forecast System (WAFS), Meteorological Watch Offices (MWO), Aerodrome Meteorological Offices (AMO), VOLMET, ATIS, Significant Weather (SIGWX) charts, Upper Wind/Temp charts, SIGMET and AIRMET criteria/validity, Aerodrome warnings, Station Model decoding.

> **Primary References:** IC Joshi Chapters 21, 24, 27, 28; Oxford/Nordian Chapter 19; Volare Question Bank.

---

graph TD

WAFC["WAFC (World Area Forecast Centres)
London & Washington
Global Upper Wind/Temp & SIGWX charts"] --> MWO["MWO (Meteorological Watch Offices)
Continuous flight watch over FIR
Issues SIGMET & AIRMET"]

MWO --> AMO["AMO (Aerodrome Meteorological Offices)
At major airports
Issues TAF, warnings, pilot flight folders"]

AMO --> AMS["AMS (Aeronautical Met Stations)
Observations: METAR & SPECI"]

---

graph TD

SIGMET["SIGMET
Severe weather hazards for ALL aircraft
Valid: 4 hours (6 hours for VA & TC)"]

AIRMET["AIRMET
Weather hazards for LOW-LEVEL flights (Below FL 100/150)
Valid: 4 hours"]

Parameter	SIGMET	AIRMET
Target Aircraft	**ALL aircraft**, regardless of size or altitude	**Low-level flights** (below FL 100, or FL 150 in mountainous terrain)
Issuing Authority	**Meteorological Watch Office (MWO)**	**Meteorological Watch Office (MWO)**
Standard Validity	**4 hours**	**4 hours**
Special Validity	**6 hours for Volcanic Ash (`VA`) & Tropical Cyclone (`TC`)**	None
Trigger Phenomena	- **Severe Turbulence** (`SEV TURB`) - **Severe Icing** (`SEV ICE` / `FZRA`) - **Severe Mountain Waves** (`SEV MTW`) - **Heavy Duststorm / Sandstorm** (`HVY DS` / `HVY SS`) - **Volcanic Ash Cloud** (`VA CLD`) - **Tropical Cyclone** (`TC`) - **Thunderstorms:** Obscured (`OBSC TS`), Embedded (`EMBD TS`), Frequent (`FRQ TS`), Squall Line (`SQL TS`) with or without hail (`GR`)	- Moderate Turbulence (`MOD TURB`) - Moderate Icing (`MOD ICE`) - Moderate Mountain Waves (`MOD MTW`) - Widespread surface wind $>30\text{ kt}$ - Surface visibility $<5,000\text{ m}$ - Isolated/Occasional thunderstorms - Mountain obscuration

---

SIGWX charts provide a graphical forecast of en-route weather hazards:

1. **High-Level SIGWX:** Covers **FL 250 to FL 630** (standard for jet transports). Depicts Jet streams (speed, height, core), Tropopause heights, CB clouds, CAT areas, Volcanic ash, and Tropical cyclones.
2. **Medium-Level SIGWX:** Covers **FL 100 to FL 250**. Includes icing, turbulence, fronts, freezing levels.
3. **Low-Level SIGWX:** Covers **Surface to FL 100** (or FL 150). Used for general aviation, depicting cloud bases/tops, visibility, and weather.

• **VOLMET (Meteorological Information for Aircraft in Flight):**

• Continuous broadcast of current **METAR, SPECI, and TAF** for designated groups of major international aerodromes.

• Broadcast via VHF (locally) or HF (long-range over oceanic and desert routes).

• **ATIS (Automatic Terminal Information Service):**

• Continuous VHF voice broadcast (or D-ATIS datalink) for arriving and departing aircraft at a specific busy airport.

• Contains: Identification letter (Alpha, Bravo, Charlie...), observation time, active runway, transition level, surface wind, visibility, weather, clouds, temperature, dew point, QNH, and special NOTAM remarks.

• Updated upon every new METAR/SPECI or immediate operational change.

• **A:** 2 hours

• **B:** 4 hours

• **C:** 6 hours

• **D:** 12 hours

View Answer

Correct Answer: Option C (6 hours)

Explanation: Routine SIGMETs for turbulence, icing, or thunderstorms have a maximum validity of 4 hours. Special SIGMETs for Volcanic Ash (`VA`) and Tropical Cyclones (`TC`) are valid for 6 hours.

• **A:** World Area Forecast Centre (WAFC).

• **B:** Meteorological Watch Office (MWO).

• **C:** Control Tower (TWR).

• **D:** Airline Flight Operations Center (FOC).

View Answer

Correct Answer: Option B (MWO).

Chapter 21: Meteorological Radar, Satellites & Instruments

• **Airborne Weather Radar (AWR):** Primary function is **hazard avoidance** (detecting wet precipitation, turbulence, hail), **NOT** terrain navigation or penetrating severe storms. Operates on **X-band (approx. 3 cm / 9375 MHz)** for optimal balance between antenna size and droplet reflection.

• **Reflectivity:** Radar detects **liquid water droplets** (reflectivity **proportional to D^6**, diameter to 6th power). Dry hail and dry snow reflect very weakly; wet hail reflects strongly.

• **Radar Tilt & Beam Geometry:** Beam width typically **3° to 5°**. Tilt control is essential to distinguish ground returns from weather cells and to scan storm cores at altitude.

• **Dangerous Radar Signatures: Hook echo** (tornado/mesocyclone), **Finger/Pendant echo** (severe updraft/hail), **U-shape / V-notch**, **Steep reflectivity gradient** (color change from green to red in very short distance = severe turbulence).

• **Radar Shadow / Attenuation:** Heavy rainfall attenuates radar pulses; an empty/black area behind an intense red core often hides a deadlier cell (**Radar Shadow / Blind Alley**).

• **Meteorological Satellites:**

• **Geostationary (GEO):** ~36,000 km over the equator, orbit matches Earth rotation, continuous coverage of same region (**60° N to 60° S**). Poor polar resolution (INSAT-3D, METEOSAT).

• **Polar Orbiting (LEO):** 800–1000 km altitude, sun-synchronous, passes over poles, global high-resolution twice daily.

• **Spectral Channels:**

• **Visible (VIS):** Reflects sunlight (daytime only). Thick clouds = Bright white; Water/Land = Dark.

• **Infrared (IR):** Measures thermal radiation / cloud-top temperature (Day & Night). High cold cloud tops (CB/Cirrus) = Bright white; Low warm clouds = Grey.

• **Water Vapour (WV):** 6.7 **\mum** channel; detects moisture in mid-to-upper troposphere (jet streams, upper lows), even without visible clouds.

• **Surface Meteorological Instruments:**

• **Barometer:** Mercury barometer (Kew / Fortin) or Aneroid barometer; calibrated to sea level.

• **Stevenson Screen:** Double-louvred wooden shelter painted white, 1.25 m (4 ft) above ground, doors open North (in Northern Hemisphere) to prevent direct solar heating during reading. Houses Dry-bulb, Wet-bulb, Maximum, and Minimum thermometers.

• **Transmissometer & Forward Scatter Meter:** Used to measure Runway Visual Range (RVR).

---

graph TD

A[Radar Transmitter X-band 3 cm] --> B[Pulse Transmitted into Cloud]

B --> C{Hydrometeor Type}

Wet Hail / Large Raindrops
Moderate Rain
Light Rain / Drizzle
Dry Hail / Ice Crystals / Cirrus

D --> H[High Turbulence Gradient / Severe Hazard Avoid by >= 20 NM]

Radar Display Color	Reflectivity (dBZ)	Rainfall Rate (mm/hr)	Associated Turbulence / Weather	Pilot Action
Black	< 20	< 0.5 (Trace)	None / Light	Safe to fly
Green	$20 - 30$	$0.5 - 2.5$	Light precipitation	Normal transit
Yellow	$30 - 40$	$2.5 - 12.5$	Moderate rain, light-moderate turbulence	Exercise caution
Red	$40 - 50$	$12.5 - 50$	Heavy rain, severe turbulence, hail probable	**AVOID by $\ge 20\text{ NM}$**
Magenta / Purple	> 50	> 50	Extreme precipitation / Doppler wind shear / Hail	**Immediate divergence**

**$Beam Width Offset (ft) = Distance (NM) x \text{Tilt Angle (**^\circ**)} x 100**$

• At 60 NM, **1°** tilt moves beam center by:

**$60 x 1 x 100 ~ 6,000 ft**$

1. **Radar Attenuation (Shadow Effect):** High-density water droplets absorb the microwave energy, preventing it from reaching clouds beyond. A black hole directly behind a bright red return is almost always **another massive cell hidden behind**, not clear sky.

2. **Stealth Hail (Upper Levels):** Above the freezing level, hail is dry. Dry ice reflectivity is **approx. 1/5th (17%)** of equivalent liquid water. A dry hail shaft will paint green or not at all on radar, but causes catastrophic airframe damage.

3. **Ground Clutter:** Mountains and terrain return strong signals. Tilting radar upwards at lower altitudes avoids confusing ground targets with storm cells.

---

graph LR

Sat[Weather Satellites] --> Geo[Geostationary GEO]

Sat --> Pol[Polar Orbiting LEO]

	Daytime Only?	Principle
	Yes (Requires daylight)	Albedo (re
	No (Day and Night)	Thermal r
	No (Day and Night)	Absorptio

---

graph TD

Inst[Met Instruments] --> P[Pressure]

Inst --> T[Temperature & Humidity]

Inst --> W[Wind]

Inst --> V[Visibility & RVR]

P --> P1[Mercury Barometer & Aneroid Barograph]

T --> T1[Stevenson Screen: Dry/Wet Bulb, Max/Min]

W --> W1[Cup Anemometer: Speed / Wind Vane: Dir at 10m AGL]

V --> V1[Transmissometer & Forward Scatter Sensor]

- **Stevenson Screen Standard:**
- **Height:** Thermometer bulbs at **1.25 to 2.0 m (4 to 6.5 ft)** above ground.
- **Orientation:** Door opens **facing North** in Northern Hemisphere (prevents direct sun falling on thermometers while reading).
- **Color & Construction:** Painted double-coated **gloss white** with double louvres for free airflow and thermal reflection.
- **Wind Measurement:**
- Standard anemometer mast height is **10 m (33 ft)** above ground in open, undisturbed terrain.
- Gust reported if peak wind exceeds mean by **>= 10 kt** during the observation period.
- **Runway Visual Range (RVR) Instruments:**
- **Transmissometer:** Light projector and receiver separated by a fixed baseline (e.g., 30–50 m). Attenuation of light beam gives transmissivity **\rightarrow** converted to RVR.
- **Forward-scatter Meter:** Measures light scattered forward at an angle (**30° - 45°**); compact single-unit mast.

---

1. **Airborne Weather Radar Wavelength: X-band (\sim 3 cm / 9375 MHz)** is standard for airborne radar. Ground weather radars use **S-band (10 cm)** or **C-band (5 cm)** to minimize attenuation over long ranges.
2. **Avoidance Distance:** Avoid severe CB / Red echoes by at least:

- **20 NM laterally** at cruise altitudes (**>= FL200**).
- **5,000 ft vertically** over cloud top if clearing above.
- Never fly beneath the anvil of a severe CB (severe turbulence, hail, microburst).

3. **Hail Return vs Rain Return:** Liquid water reflects **approx. 5 times more power** than dry ice of the same diameter; however, hail is typically much larger (**D^6** factor dominates when wet).
4. **Blind Alley Trap:** Flying between two intense radar echoes with a narrow gap. Turbulence can merge echoes, trapping the aircraft. Maintain minimum **40 NM** clearance between cells.

5. **Geostationary Satellite Invisibility:** GEO satellites cannot image polar regions (above **75° - 80°** latitude) due to the extreme curvature angle (oblique horizon). Polar-orbiting satellites are required for polar routes.

• Test your concepts with:

• [`question_bank/06_metar_taf_briefing.md`](../question_bank/06_metar_taf_briefing.md)

• [`question_bank/08_thunderstorms_hazards.md`](../question_bank/08_thunderstorms_hazards.md)

• Run the interactive quiz CLI: `python scripts/interactive_quiz.py`