



INSIDE A CELL


Dr. Manu S Singh

Department of Biotechnology


Bennett University

Table 5.2 A Comparison of Bacterial, Animal, and Plant Cells

Bacterium	Animal	Plant
EXTERIOR STRUCTURES		
Cell wall	Present (protein-polysaccharide)	Absent
Cell membrane	Present	Present
Flagella	May be present (single strand)	May be present
INTERIOR STRUCTURES		
ER	Absent	Usually present
Ribosomes	Present	Present
Microtubules	Absent	Present
Centrioles	Absent	Present
Golgi apparatus	Absent	Present
Nucleus	Absent	Present
Mitochondria	Absent	Present
Chloroplasts	Absent	Absent
Chromosomes	A single circle of DNA	Multiple; DNA-protein complex
Lysosomes	Absent	Usually present
Vacuoles	Absent	Absent or small









Plant Cell Magnification: 14000X




Plant vs Animal Cells

Animal Cell Magnification: 40500X
(White Blood Cells)


ZOOM IN



- Most cells are large enough to be resolved from each other with **light microscopes (LM)**
 - Cells were discovered by Robert Hooke in 1665
 - he saw the remains of cell walls in cork with a LM
 - his microscope had about 30x magnification
 - Modern LMs can reach up to 1000x



SCALE





- How do proteins get outside of a cell?
- How do proteins get into a cell membrane?
- How does a cell digest its food?
- How does a cell commit suicide?
- Why would a cell commit suicide?

PLASMA MEMBRANE



FUNCTIONS OF PLASMA MEMBRANE

- **Protective**:- Forms outermost boundary of the cells.
- **Digestive**: Takes in food and excretes waste products.
- **Selective Permeability**:-Helps in transport across the membrane.
- Contains cell **surface receptors** (e.g: Glycoprotein receptors present on RBCs).
- **Cell Adhesion Molecules** (Cadherins) present on the plasma membrane of certain cells plays an important role in the process of inflammation.
- **Junctions**: Helps in formation of various types of junction (Adherens & Anchoring) along with the help of cytoskeleton elements.





[Video Link](#)

PLASMA MEMBRANE



[Video Link](#)

OSMOSIS VS REVERSE OSMOSIS



BLOOD GROUPS







Figure 1. ABO antigen specificity. The ABO antigens differ by just one sugar at the antigen terminus. Only the carbohydrate portion of the antigen is illustrated.


		Blood Type			
		A	B	AB	O
Red Blood Cell Type					
Antibodies in Plasma				None	
Antigens in Red blood Cell				A and B antigens	None
Blood Types Compatible in an Emergency	A, O	B, O	A, B, AB, O (AB ⁺ is the universal recipient)		O (O is the universal donor)

PLASMA MEMBRANE-TRANSPORT



CYTOPLASM

- Water-like substance fills the space between the plasma membrane and the nuclear membrane.
- Consists of **cytosol** and **cellular organelles** except for the cell nucleus.
- **Cytosol** is made up of water, salts, organic molecules and many enzymes that catalyze reactions.





CYTOSKELETON

- Maintains cell shape.
- Protects the cell.
- Plays important roles in intra-cellular transport (the movement of vesicles and organelles).
- Plays important role in cell division.


RIBOSOME- PROTEIN FACTORIES

Granular bodies which are made up of ribonucleoprotein (RNA+Protein)

- two main subunits- large and small
- perform the enzymatic activity for forming peptide bonds, and serve as the sites of translation of genetic information (mRNA) into protein
- some are free in the cytoplasm while others are associated with the endoplasmic reticulum (ER)




RIBOSOME-ZOOM IN





- Three different types of RNA:
 - mRNA (messenger) used as template to make proteins
 - rRNA (ribosomal) makes up ribosomes
 - tRNA (transfer) matches amino acids to mRNA to help make proteins


RNA Functions



RIBOSOME-ZOOM IN





POLYRIBOSOME IN ACTION



Single mRNA can lead to formation of multiple proteins as per requirement


ENDOPLASMIC RETICULUM (ER)

- Make and package proteins and lipids
- Much like an assembly line
- Found in eukaryotic organisms
- Forms an interconnected network of flattened, membrane-enclosed sacs or tubes known as cisternae.
- Largest organelle in eukaryotic cell.
- It provides separate chemical environment which allows for correct protein folding.
- Two types
 - Rough Endoplasmic Reticulum
 - Smooth Endoplasmic Reticulum




Smooth ER – primary site of lipid synthesis, many detoxification reactions in liver cells

Rough ER – ribosomes that attach & insert proteins into the ER lumen as they are synthesized




ENDOPLASMIC RETICULUM (ER)

- Endoplasmic reticulum (**ER**) – membrane network that winds through the cytoplasm
 - winding nature of the ER provides a lot of surface area
 - many important cell reactions or sorting functions require ER membrane surface
 - **ER lumen** – internal aqueous compartment
- Proteins processed:
 - Soluble- ER Lumen
 - Membrane bound- ER membrane



SMOOTH VS ROUGH ER




Smooth ER



- Synthesizes lipids
- Metabolizes carbohydrates
- Detoxifies drugs and poisons
- Stores calcium ions

Rough ER

- Has bound ribosomes, which secrete **glycoproteins**
- Distributes transport vesicles
- Is a membrane factory for the cell



ROUGH ER TO GOLGI BODIES




[Video Link](#)


GOLGI BODIES-POST OFFICE

- Glycosylation of proteins and lipids
- Processing, sorting and transport of membrane molecules
- Carbohydrate and lipid metabolism
- Synthesis of complex polysaccharides for cell wall
- Produce mucus
- Packages products into vesicles for transport

RER → ER-Golgi vesicles → Golgi Cisternae Network
→ Secretory Vesicles → Cell Exterior





GOLGI BODIES-POST OFFICE/ EXOCYTOSIS





LYSOSOME- DIGESTIVE SYSTEM


- Vary in size and shape
- Acidic pH (5) maintained by H⁺ ATPase pumps
- Hydrolytic enzymes- protease, nuclease, glycosidase, lipase
- Microbial killing via phagocytosis of immune cells
- Heterophagy and autophagy (self-eating)



VACUOLES



- Plant and fungal cells
- Membrane called tonoplast
- Filled with water, ions, acids, sugars, enzymes
- Amoeba and paramecium have contractile vacuole

ANIMAL CELL	PLANT CELL
	
<ul style="list-style-type: none"> • several small vacuoles • used for storage • can contain nutrients, water, or waste 	<ul style="list-style-type: none"> • one large vacuole • used to store water and push against the cell wall • keeps the plant rigid




MITOCHONDRIA

- Powerhouse of cells
- Produce cellular energy in the form of ATP through aerobic respiration
- Double membrane
- Humans have 16.6 kilobase sized DNA in mitochondria (mtDNA)
- Maternal Inheritance
- Divide on their own





MITOCHONDRIA




CHLOROPLAST


- Double membrane bound organelle called plastids
- Differentiate as leucoplast, chromoplast, chloroplast
- Flattened sacs called granum, stacked together called thylakoids
- Comprised mainly of lipids
- Self-replicating organelles



MOVEMENT APPENDAGES




(a) Flagella



(b) Cilia


Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings.

Protists




Gut cells

CILIA



Structure of Eukaryotic Cilium



- Occur throughout the cell
- Beat in coordination
- Found in lining of body tubes
- Nasal passage clean by preventing dust accumulation and secretion of mucus
- Locomotion, feeding, circulation


[Animation Link](#)

NUCLEUS

1. Nuclear Envelope
2. Matrix
3. Nucleolus
4. Chromatin

Nuclear Matrix

- Structural framework for organizing chromatin
- High-salt buffers, nucleases & non-ionic detergents



[Video Link](#)


NUCLEAR ENVELOPE & NUCLEUS

Nuclear Envelop

- Contains pores- nuclear pore complex
- Transportation between cytoplasm & nucleus

Nucleolus

- Non-membrane bound dynamic body
- Disappears during late prophase
- Site of transcription of ribosomal RNA
- Assembly of ribosome



CHROMATIN






Figure: The human cell nucleus.

- Organized structure of DNA and protein that is found in the nucleus
- In G0 stage, one chromatin contains a single dsDNA in coiled and condensed form
- Chromosomes are more condensed than chromatin
- Regions- Euchromatin and Heterochromatin
- Euchromatin- Region active in transcription
- Heterochromatin- transcriptionally silent and stains darker



HUMAN CHROMOSOMES



Karyotyping

NUCLEUS

Centromere-

- Constricted region of linear chromosome
- Site of attachment of sister chromatids

Telomeres

- Caps the ends of chromosomes
- Rich in G

Origin of replication





Table 5.1 Eukaryotic Cell Structures and Their Functions

Structure	Description	Function
Cell wall	Outer layer of cellulose or chitin; or absent	Protection; support
Cytoskeleton	Network of protein filaments	Structural support; cell movement
Flagella (cilia)	Cellular extensions with 9 + 2 arrangement of pairs of microtubules	Motility or moving fluids over surfaces
Plasma membrane	Lipid bilayer with embedded proteins	Regulates what passes into and out of cell; cell-to-cell recognition
Endoplasmic reticulum	Network of internal membranes	Forms compartments and vesicles; participates in protein and lipid synthesis
Nucleus	Structure (usually spherical) surrounded by double membrane that contains chromosomes	Control center of cell; directs protein synthesis and cell reproduction
Golgi apparatus	Stacks of flattened vesicles	Packages proteins for export from cell; forms secretory vesicles
Lysosomes	Vesicles derived from Golgi apparatus that contain hydrolytic digestive enzymes	Digest worn-out organelles and cell debris; play role in cell death
Microbodies	Vesicles formed from incorporation of lipids and proteins containing oxidative and other enzymes	Isolate particular chemical activities from rest of cell
Mitochondria	Bacteria-like elements with double membrane	“Power plants” of the cell; sites of oxidative metabolism
Chloroplasts	Bacteria-like elements with membranes containing chlorophyll, a photosynthetic pigment	Sites of photosynthesis
Chromosomes	Long threads of DNA that form a complex with protein	Contain hereditary information
Nucleolus	Site of genes for rRNA synthesis	Assembles ribosomes
Ribosomes	Small, complex assemblies of protein and RNA, often bound to endoplasmic reticulum	Sites of protein synthesis



Questions?

ANNEXURE

Blood Type	Antigens Present	Antibodies in Serum
<i>AB+</i>	<i>ABH, Rh D</i>	<i>None</i>
AB-	ABH	Rh D
<i>A+</i>	<i>AH, Rh D</i>	<i>B</i>
A-	AH	B, Rh D
<i>B+</i>	<i>BH, Rh D</i>	<i>A</i>
B-	BH	A, Rh D
<i>O+</i>	<i>H, Rh D</i>	<i>AB</i>
O-	H	AB, Rh D
<i>Bombay</i>	<i>None</i>	<i>ABH, Rh D</i>

**BOMBAY
BLOOD GROUP
OR
HH BLOOD
GROUP**