



Networks used in every experiment — identical except for the widths of the data fix;
biases are drawn from the same distribution as the weights, and only the optimizer varies

Vehicle 18-50-4




4 vehicle silhouettes · sigmoid throughout · MSE loss
batch 64 · 950 epochs · Glorot-uniform init

Satimage 36-50-6




6 soil types · sigmoid throughout · MSE loss
batch 64 · 125 epochs · Glorot-uniform init

Dry Bean 16-50-7




7 bean cultivars · sigmoid throughout · MSE loss
batch 64 · 60 epochs · Glorot-uniform init

Segment 19-50-7




7 image segments · sigmoid throughout · MSE loss
batch 64 · 350 epochs · Glorot-uniform init

Optdigits 64-50-10




10 digits · sigmoid throughout · MSE loss
batch 64 · 145 epochs · Glorot-uniform init

Pendigits 16-50-10




10 digits · sigmoid throughout · MSE loss
batch 64 · 75 epochs · Glorot-uniform init

MNIST 784-50-10



10 digits · sigmoid throughout · MSE loss
batch 64 · 10 epochs · Glorot-uniform init

Letter Recognition 16-50-26



26 letters · sigmoid throughout · MSE loss
batch 64 · 60 epochs · Glorot-uniform init