Kaya-guided hybrid deep learning for leave-country-out estimation of national CO_2 emissions: a 112-country benchmark with country-clustered inference, conformal uncertainty, and explainable AI

Short title: Leave-country-out estimation of national CO_2 emissions

Samiuel C. Atuahene[1,2*], Bismark Boateng[1], Hilda Amarfio[3], Ekow Yaleh Quayson[1], Francis Boateng Agyenim[1]

~~Francis Boateng Agyenim[1],~~

[1] Emerging Engineering Technologies Division, CSIR~~,~~ – Institute of Industrial Research, East Legon, Accra, Ghana

[2] Research and Development Department, Premium Intelligence Networks and Systems Co. Ltd., Accra, Ghana

[3] Department of Maternal and Child Health Nursing, School of Nursing ~~&~~and Midwifery, College of Health Sciences, University of Ghana, Legon, Accra~~.~~, Ghana

~~* Corresponding author: Samiuel C. Atuahene~~

~~Abstract~~

~~Reliable national CO_2 inventories underpin the Paris Agreement's Enhanced Transparency Framework, yet many non-Annex I countries lack complete emission statistics. We reframe national emissions modelling as *leave-country-out* (LOCO) estimation: predicting a country's per-capita CO_2 from eight-year trajectories of energy and socioeconomic covariates, with no access to its own emission history. Using Global Carbon Budget and Energy Institute data harmonized by Our World in Data, we build a 112-country panel (1990–2023; 3,024 windowed samples) and benchmark thirteen models under country-disjoint five-fold cross-validation. The proposed Kaya-guided CNN-BiLSTM-Attention network (KG-CBA) couples a log-linear~~

backbone motivated by the Kaya identity to a convolutional-recurrent-attention branch that learns only the nonlinear residual, tuned by particle swarm optimization. Because the sample contains 27 repeated years per country, all inference is country-clustered. Kaya guidance significantly improves the unguided architecture (median APE 15.7% vs 20.8%, $p = 0.002$), and ridge regression significantly outperforms every tree ensemble and every unguided deep network; but the best ensemble's apparent edge over ridge is not significant (15.1% vs 16.3%, $p = 0.22$), where observation-level testing would report $p < 10^{-9}$. Error is highly concentrated: six structurally atypical energy systems contribute 79–84% of total squared error. Adding the low-carbon energy share on a 76-country sub-panel halves median error (7.35% vs 13.72% like-for-like), showing that energy-mix data availability, not model sophistication, is the binding constraint. Removing all CO_2-derived covariates costs under one percentage point, so the estimator suits countries with no inventory at all. Split-conformal Monte-Carlo-dropout intervals reach 81.9% coverage at a 90% nominal level, the shortfall traced to non-exchangeability under grouped splits. In conventional temporal forecasting, learners beat naive persistence only marginally and inconsistently across evaluation windows. Data and code are open.

Keywords: CO_2 emissions; hybrid deep learning; Kaya identity; clustered inference; conformal prediction; explainable artificial intelligence; greenhouse gas inventories; Africa

1. Introduction

Anthropogenic carbon dioxide remains the dominant driver of observed warming, and fossil CO_2 emissions reached record levels again in 2024 [1,2]. Article 13 of the Paris Agreement obliges all parties to report national greenhouse-gas inventories under the Enhanced Transparency Framework, yet the reporting landscape is deeply uneven. Only a minority of non-Annex I countries submit complete and timely inventories, and independent assessments find the largest

~~inter-source discrepancies precisely where capacity is lowest, notably across sub-Saharan Africa [3–5]. Gap-filling therefore matters: mitigation planning, carbon-market integrity, and the global stocktake all depend on emission estimates for countries whose own statistical systems cannot yet produce them [5,6].~~

~~Machine learning has been proposed as a pragmatic response. Recent work spans subnational gap-filling for G20 jurisdictions [6], sectoral emission prediction for African economies [7–10], and a rapidly growing literature on deep and hybrid architectures for CO_2 forecasting [11–18]. Within that literature, hybrid designs that stack convolutional feature extractors, recurrent sequence models, and attention mechanisms, often tuned by metaheuristic algorithms, now dominate methodological claims of state-of-the-art accuracy [13–15,19–21].~~

~~Four weaknesses limit the practical value of much of this evidence base. First, most studies evaluate *temporal* forecasting for countries whose emission history is fully observed, precisely the setting where inventories already exist, rather than the *cross-country generalization* setting that the inventory-gap problem poses. Second, validation protocols frequently allow the same country to appear in both training and test data, inflating reported skill through country-specific memorization. Third, naive benchmarks are routinely absent; national per-capita emissions are highly persistent from year to year, and without a persistence baseline, near-unity coefficients of determination are easily mistaken for skill [22–24]. Fourth, and least remarked, significance is almost always assessed by treating every country-year as an independent observation. Panel emission data are strongly clustered within countries; observation-level tests on such data understate standard errors by roughly the square root of the cluster size, and can render a fraction of a percentage point "highly significant".~~

This paper addresses all four. We formalize national emissions estimation as a leave-country-out (LOCO) problem: given only eight-year trajectories of six widely available covariates, predict a country's per-capita CO_2 without access to its own emission record. Because the target country contributes no training information, persistence is undefined and the task isolates genuine covariate-to-emissions generalization. All inference is clustered at the country level. All experiments use measured data anchored in the Global Carbon Budget [2] and the Energy Institute Statistical Review [25], as harmonized by Our World in Data [26].

Methodologically, we exploit a structural property of the task. In logarithms, the Kaya identity [27,28] renders per-capita emissions approximately linear in energy intensity and carbon intensity, which explains an empirical result we report below: a ridge regression is remarkably hard to beat. Rather than ignoring this, we encode it. The proposed KG-CBA network attaches a log-linear head, initialized at a ridge solution, to a convolutional, bidirectional-recurrent, attention-pooled branch that learns only the nonlinear residual, tuned by particle swarm optimization [29-31]. Uncertainty is quantified with Monte-Carlo dropout [32] calibrated by per-fold split-conformal inference [33-35], and interpretability through SHapley Additive exPlanations [36] and attention-weight analysis.

The contributions are fivefold, and are as much about how the evidence is established as about what it shows. (i) A rigorously country-disjoint, 112-country LOCO benchmark of thirteen statistical, machine-learning, and deep-learning models on open, measured data, with country-clustered significance testing throughout. (ii) A physics-guided hybrid architecture, KG-CBA, with PSO-based hyperparameter optimization, which significantly improves its unguided counterpart. (iii) An interpretable error decomposition showing that a handful of structurally atypical energy systems dominates the error budget, and that a single additional covariate

roughly halves median estimation error where it is available. (iv) Distribution-aware uncertainty quantification whose honest coverage shortfall is traced to non-exchangeability under grouped resampling. (v) A temporal experiment showing that naive persistence is a strong benchmark whose margin over well-specified learners is small and unstable across evaluation windows. Section 2 reviews related work. Section 3 describes the data and methods. Section 4 reports results. Section 5 discusses implications, limitations, and threats to validity. Section 6 concludes.

2. Related work

2.1 Machine and deep learning for CO_2 emission modelling

Comparative studies consistently find nonlinear learners competitive or superior to classical statistical models for emission prediction. Ajala et al. [11] benchmarked statistical (ARMA/ARIMA family), machine-learning (SVM, random forest, gradient boosting), and seven deep architectures for daily CO_2 across China, India, the USA, and Europe. Architecture-comparison studies report horizon-dependent trade-offs among LSTM, GRU, CNN-LSTM, and dense networks [12], while temporal convolutional hybrids with hyper parameter search achieve strong accuracy on periodic emission series [13]. Attention-augmented and feature-fused hybrids continue this trend: CNN-BiLSTM-attention frameworks improve daily national and plant-level carbon prediction [14,15], and TCN-LSTM hybrids transfer to maritime emissions [16]. Sector-specific applications include transport [17,37–39], data centers [40], and energy-based national totals [41]. Marked publication emphasis on incremental architecture novelty, however, has not been matched by equally careful evaluation design, motivating the protocol adopted here.

2.2 Emissions modelling for data-sparse regions

For Africa specifically, ensemble tree methods have been applied to building-sector emissions across six nations [7], continent-wide emission determinants for 48 countries [8], carbon-neutrality trajectories in sub-Saharan Africa [9], and transport emissions of the continent's four largest emitters [10]; a national decarburization-planning framework has been proposed for Nigeria [42]. Beyond Africa, machine-learning gap-filling of subnational inventories demonstrates the feasibility of covariate-based estimation at scale [6]. Inventory-quality assessments document why such estimation is needed: most non-Annex I countries rely on Tier-1 approaches with scarce activity data, and independent datasets disagree most where capacity is lowest [3-5,43]. Our LOCO formulation directly simulates this deployment setting, which, to our knowledge, has not previously been used to benchmark hybrid deep architectures at global scale.

2.3 Metaheuristic optimization, uncertainty, and explainability

Metaheuristic hyper parameter tuning is now standard practice in energy and emissions forecasting: PSO variants improve LSTM load forecasting [19,44], attention-augmented CNN-LSTM demand models [20], and bidirectional recurrent emission predictors, while novel hybrids have been applied directly to CO_2 prediction [21]. For uncertainty, conformal prediction supplies distribution-free finite-sample coverage [33,45]; ensemble conformalized quantile regression extends it to probabilistic time series [34], with energy-domain applications in solar forecasting [46,47], and recent theory characterizes behavior under non-exchangeable data [35,48], directly relevant to our grouped country splits. For explainability, SHAP [36] is the de-facto standard in emission applications [49-51]. The present study integrates all three strands within a single, reproducible pipeline.

~~2.4 The gap this study fills~~

~~Table B3 positions this work against representative prior studies on five axes. To our knowledge, no previous study of machine learning for national CO_2 estimation combines all of the following: (i) evaluation under strictly country-disjoint splits, so that no test country contributes any training signal; (ii) explicit naive benchmarks with scaled error, so that reported skill is measured against the persistence a random walk would deliver; (iii) inference clustered at the country level, so that repeated years within a country are not counted as independent evidence; (iv) architectural structure derived from an accounting identity rather than chosen by search; and (v) distribution-free uncertainty quantification reported honestly under the grouped sampling the task requires. Individually each of these is established practice in some adjacent literature. Their absence in combination is what allows the field's reported accuracies to look far better than they generalize, and supplying them in combination is this paper's principal methodological contribution. The substantive contribution follows from it: once the evaluation is trustworthy, the ranking of what actually improves national emission estimation changes, and the binding constraint turns out to be data availability in specific countries rather than model capacity.~~

* Corresponding author

E-mail: satuahene@mail.ru (SCA)

Abstract

Many countries cannot yet produce complete and timely greenhouse-gas inventories, so their emissions have to be estimated from other information. We asked how accurately a country's per-capita fossil CO_2 emissions can be estimated from widely available energy and socio-economic indicators when none of that country's own data are used to fit the model, and whether deep learning improves on a simple linear model for this task. Using open data for 112 countries

(1990–2023; 3,024 samples), we compared thirteen models, from ridge regression to a proposed hybrid network (KG-CBA) that adds a deep-learning correction to a linear component motivated by the Kaya identity. Every model was tested only on countries withheld from training (leave-country-out cross-validation), and differences between models were tested with countries, not country-years, as the unit of inference. Ridge regression reached a median absolute percentage error (MdAPE) of 16.3% and was significantly more accurate than the tree-based models and the unguided convolutional networks (CNN-LSTM and CNN-BiLSTM-Attn). Guiding the network with the Kaya-motivated linear component reduced its MdAPE from 20.8% to 15.7% ($p = 0.002$), but no network or ensemble was shown to be more accurate than ridge regression: the best ensemble (15.1%) did not differ significantly from it ($p = 0.22$). Seven economies with unusual energy systems accounted for 78% of the squared error of ridge regression. Where the low-carbon share of primary energy was available (76 countries), adding this single indicator halved the MdAPE (15.6% to 7.7%, $p < 0.001$). The results were insensitive to the treatment of missing data and to the exclusion of the atypical economies. For estimating national emissions without a country's own records, a transparent linear model with the right energy indicators is sufficient, and better energy-mix statistics would improve the estimates more than more complex models.

Introduction

Carbon dioxide (CO_2) from fossil fuels and industry is the main driver of global warming, and emissions continue to set records [1,2]. Under the Enhanced Transparency Framework of the Paris Agreement, all Parties report national greenhouse-gas inventories [3], but the capacity to do so is uneven. Official inventories from many developing countries are infrequent or incomplete, and the gaps are filled with third-party estimates [4,5]. For fossil CO_2, independent data sets

agree closely in aggregate and less closely country by country: global totals differ by about 5% once system boundaries are harmonized, whereas estimates for individual countries, regions or fuels can differ by 15% or more [6]. In Africa, two widely used data sets differ by less than 5% for most of the main emitters, but by more than 10% for South Africa and by a factor of three for Mali, and African countries do not report every year [7]. Estimates that need only widely available indicators, and not a country's own emission accounts, are therefore useful for filling gaps and for cross-checking reported values.

Machine learning is increasingly proposed for this purpose. Comparative studies report that non-linear learners match or outperform classical time-series models for CO_2 series [8–10], with applications to transport [11,12], national totals [13] and sub-Saharan African economies [14]. Much recent work proposes hybrid architectures that stack convolutional, recurrent and attention layers and tunes them with metaheuristic search such as particle swarm optimization (PSO) [15,16].

Four features of this literature limit its value for the inventory-gap problem. First, most studies forecast future emissions of countries whose emission history is observed [8–10,13], the setting in which inventories already exist. Second, when the same country contributes data to both training and testing, a model can reproduce country-specific levels without having learned a transferable relationship; for grouped data the validation split must respect the groups [17,18]. Third, annual per-capita emissions change slowly, so a naive forecast that repeats the last observed value is already accurate, and accuracy that is not compared with it overstates skill [19,20]. Fourth, significance is usually assessed as if every country-year were an independent observation, although repeated years of one country are strongly correlated; ignoring this understates standard errors, often severely [21–23].

This study addresses these points by treating national emission estimation as a leave-country-out (LCO) problem: a country's per-capita CO_2 emissions are estimated from eight-year series of its energy and socio-economic indicators by a model fitted only to other countries. (We write LCO because LOCO is established for leave-one-covariate-out analysis of variable importance [24], which we also use.) Because the target country contributes nothing to model fitting, no persistence forecast can be formed for it, and the task measures how well the relationship between indicators and emissions transfers between countries.

The task has a known structure. The Kaya identity expresses emissions as the product of population, income, the energy intensity of income and the carbon intensity of energy [25,26]; in logarithms it is additive, which is the basis of log-linear emission models [27]. A linear model in logarithms is therefore a natural reference, and the question for more flexible models is whether they add anything to it. Following the idea of theory-guided machine learning [28–30], we test this with a hybrid network, KG-CBA (Kaya-guided convolutional neural network, bidirectional long short-term memory and attention), in which a linear component carries the Kaya structure and a deep network models only what the linear component leaves unexplained.

The study answers four questions:

- Q1. How accurately can per-capita fossil CO_2 emissions be estimated for a country that is absent from training?
- Q2. Do deep and hybrid networks outperform a linear model motivated by the Kaya identity, once differences are tested with countries as the unit of inference?
- Q3. Is the remaining error reduced more by better models or by better covariate data?
- Q4. Are the answers robust to the treatment of missing data and to atypical economies, and how reliable are the prediction intervals and explanations of the models?

To our knowledge, national CO_2 models have not previously been compared under country-disjoint validation combined with country-clustered inference. A supporting experiment on conventional temporal forecasting, in which naive persistence can serve as a benchmark, is reported in S2 Text.

~~3.~~ Materials and methods

~~3.1 Data sources and panel construction~~

~~All measured series derive from Our World in Data's harmonized CO_2 and energy datasets [26], which redistribute the Global Carbon Budget national emission accounts [2] and the Energy Institute Statistical Review primary-energy statistics [25], accessed 26 July 2026. From the raw country–year records we retain 1990–2023 and exclude regional aggregates. Six covariates form the core feature set: primary energy consumption per capita [equation removed or moved to S1 Text] (kWh person^{-1}), GDP per capita [equation removed or moved to S1 Text] (constant international $), population [equation removed or moved to S1 Text], and the coal, oil, and gas shares of CO_2, [equation removed or moved to S1 Text] and analogously for oil and gas. The target is per-capita fossil CO_2, [equation removed or moved to S1 Text] (t CO_2 person^{-1} yr^{-1}).~~

~~Interior gaps of at most three consecutive years are filled by linear interpolation; series edges are extended with the nearest observed value. Countries are retained if all seven series are complete for all 34 years after this treatment, yielding a balanced panel of 112 countries × 34 years (3,808 country–years).~~

~~Imputation is material and unevenly distributed, and we report it in full. Table 3 gives the audit. Population, oil share and per-capita CO_2 require no filling; energy per capita is filled for 0.7% of country–years and GDP per capita for 3.0%, but the coal and gas shares are filled for 6.4% and~~

7.6% respectively. Critically, because the edge rule carries no length cap, some series are extended over long stretches: the longest single filled run reaches 31 years for the gas share, 26 for the coal share and 16 for energy per capita. Twenty-five of the 112 countries contain at least one filled run exceeding three years, and nineteen contain a run exceeding ten. These countries are disproportionately low- and middle-income (El Salvador 31 years, Niger 29, Panama 28, Ghana 26, Jamaica 26, Guatemala 26, Cameroon 26, Rwanda 25, Benin 24), because it is precisely there that the international energy lineages are thin. Constant extension makes a covariate series artificially smooth and therefore artificially easy to predict from, so this imputation flatters rather than penalizes the affected countries; Section 4.4 quantifies the effect and Section 5.3 discusses it. All country-level results below are annotated with imputation status, and the full per-country audit is released with the code.

The panel spans 0.03–364.8 t CO_2 person^{-1} over 1990–2023. Because the eight-year window means prediction targets run 1997–2023, the largest *modelled* value is Qatar's 76.7 t person^{-1} in 1997; Kuwait's 1991 value of 364.8 t person^{-1}, a genuine artefact of the Gulf-War oil fires coinciding with a population collapse, enters only as covariate context and is never a prediction target.

For Experiment C, a seventh covariate, the low-carbon share of primary energy [equation removed or moved to S1 Text] (renewables plus nuclear), available from the Statistical Review lineage for 76 of the 112 countries (≥90% completeness), is added under identical processing with no additional imputation required. This sub-panel contains all the countries later identified as error-dominating outliers.

~~**3.2 Problem formulation**~~

Study design and workflow

The study follows the seven steps of Fig 1. Three experiments use one data set and one validation design. Experiment A, the main benchmark, compares thirteen models on 112 countries with six covariates and answers Q1 and Q2. Experiment C repeats the comparison for the leading models on the 76 countries for which a seventh covariate, the low-carbon share of primary energy, is available, and answers Q3. Experiment B, a supporting analysis of conventional temporal forecasting in which naive persistence is a valid benchmark, concerns a different task and is given in S2 Text. The labels follow the order in which the experiments were run. Sensitivity, uncertainty and explainability analyses answer Q4.

Fig 1. Workflow of the study. The research questions (step 1) determine the data (step 2) and the three experiments (step 3). Thirteen models and an exploratory ensemble (step 4) are compared under leave-country-out cross-validation with country-clustered inference (step 5) and examined by diagnostic and robustness analyses (step 6), leading to the conclusions on models, data and practice (step 7). APE, absolute percentage error; CNN, convolutional neural network; GRU, gated recurrent unit; kNN, k-nearest neighbours; LSTM, long short-term memory; MAE, mean absolute error; MAPE, mean absolute percentage error; ML, machine learning; PSO, particle swarm optimization; RMSE, root-mean-square error; SHAP, Shapley additive explanations; SVR, support vector regression.

Data

All series were taken from the open CO_2 and energy data sets compiled by Our World in Data [31] (files `owid-co2-data.csv`, repository version of 2 June 2026, and `owid-energy-`

`data.csv`, version of 27 April 2026; accessed on 26 July 2026). These files redistribute national fossil CO_2 emissions from the Global Carbon Budget 2025 [2], primary energy consumption from the Energy Institute Statistical Review of World Energy [32] and the U.S. Energy Information Administration [33], gross domestic product (GDP) from the Maddison Project Database 2023 [34], and population estimates.

The target is annual per-capita CO_2 emissions from fossil fuels and industry, excluding land-use change (c, in t CO_2 per person). Experiment A uses six covariates: primary energy consumption per capita (E, kWh per person), GDP per capita (G, international dollars per person at 2011 prices), population (P, persons), and the shares of national fossil CO_2 emissions that come from coal, oil and gas (s^{coal}, s^{oil} and s^{gas}; fractions between 0 and 1). Experiment C adds the share of primary energy supplied by low-carbon sources, that is, renewables and nuclear power (s^{lc}; fraction between 0 and 1), which the Statistical Review provides for 76 of the 112 countries.

We retained the years 1990 to 2023 and excluded regional aggregates. Missing values were filled in three steps for each series of each country: gaps were linearly interpolated for at most three consecutive years; values still missing were set to the last available value before them; and values missing at the start of a series were set to its first observed value. A country was retained if all seven series were then complete for all 34 years. The result is a balanced panel of 112 countries and 34 years, or 3,808 country-years (S1 Table lists the countries). Table 1 summarizes the variables, and Fig 2 shows the target and every covariate over the study period.

Table 1. Variables of the panel (112 countries, 1990–2023, 3,808 country-years).

Variable (symbol)	Unit	Source	Median (interquartile range)	Filled country-years, n (%)	Longest filled run, years
Per-capita CO_2 emissions (c), target	t CO_2 per person per year	[2]	4.45 (1.67–8.36)	0 (0.0)	0

Variable (symbol)	Unit	Source	Median (interquartile range)	Filled country-years, n (%)	Longest filled run, years
Primary energy per capita (E)	kWh per person	[32,33]	22,121 (8,116–43,035)	28 (0.7)	16
GDP per capita (G)	International dollars per person, 2011 prices	[34]	12,688 (5,643–27,786)	114 (3.0)	2
Population (P)	Million persons	[31]	11.2 (5.1–41.7)	0 (0.0)	0
Coal share of CO_2 (s^{coal})	Fraction	[2]	0.12 (0.01–0.35)	244 (6.4)	26
Oil share of CO_2 (s^{oil})	Fraction	[2]	0.50 (0.33–0.66)	0 (0.0)	0
Gas share of CO_2 (s^{gas})	Fraction	[2]	0.18 (0.05–0.33)	291 (7.6)	31
Low-carbon share of primary energy (s^{lc}), 76 countries	Fraction	[32]	0.11 (0.03–0.22)	0 (0.0)	0

"Filled" counts values that were missing in the source and were completed as described in the text. The longest filled run is the largest number of consecutive filled years in any one country.

Fig 2. The target variable and the covariates, 1990–2023. Lines show the median across countries, dark bands the 25th to 75th percentiles and light bands the 10th to 90th percentiles. (a) Per-capita CO_2 emissions. (b) Primary energy per capita. (c) Gross domestic product (GDP) per capita in international dollars at 2011 prices. (d) Population. (e–g) Shares of fossil CO_2 emissions from coal, oil and gas. (h) Low-carbon share of primary energy. Panels a–g show the 112 countries of Experiment A and panel h the 76 countries of Experiment C. Panels a–d have logarithmic vertical axes.

The amount of filling differs between variables (Table 1). The target, population and the oil share needed none. Of the 114 filled values of GDP per capita, 112 are the year 2023, which the Maddison database does not yet cover, so that the 2022 value was carried forward. The coal and gas shares were filled for 244 and 291 country-years, and here the filled runs are long: 25 countries have a filled run of more than three years and 19 a run of more than ten years (S1 Table).

These long runs do not stand for unknown quantities. In the source, the emissions of a fuel are blank where that fuel does not appear in the national total: in all 464 country-years with a blank coal or gas entry, the reported total equals the sum of the reported components (coal, oil, gas, cement, flaring and other industry) to within 0.3% of the total. The blank shares are therefore zeros, or nearly so, for fuels that a country did not burn in those years, and the fill rule replaced them with small positive values: 81% of the filled coal shares and 68% of the filled gas shares are below 0.01, and 95% and 90% are below 0.05. We kept the original fill in the main analysis so that all models, including the neural networks, were trained on the same panel, and we examine alternatives in the sensitivity analyses.

Per-capita emissions range from 0.03 to 364.8 t CO_2 per person. The largest value (Kuwait, 1991, the year of the oil-well fires) precedes the first target year; the largest modelled value is 76.7 t per person (Qatar, 1997).

Estimation task and validation

Let i index countries and t years. ~~Skewed~~The ~~magnitudes~~three strictly positive covariates and the target are log-transformed ~~via~~(Eq ~~equation (~~1)~~,~~:

~~[equation removed or moved to S1 Text]~~ $e_{i,t} = \log_{10} E_{i,t}, \quad g_{i,t} = \log_{10} G_{i,t}, \quad p_{i,t} = \log_{10} P_{i,t}, \quad y_{i,t} = \ln c_{i,t}.$ (1)

~~and the target is modelled in natural logarithm, equation (2),~~

~~[equation removed or moved to S1 Text]~~

~~For each country-year with at least [equation removed or moved to S1 Text] years of history we form the covariate window of equation (3),~~

Each sample pairs the covariates of $W = 8$ consecutive years with the target in the last of those years (Eq 2):

~~[equation removed or moved to S1 Text]~~ $\mathbf{X}_{i,t} = [\mathbf{x}_{i,t-W+1}, \ldots, \mathbf{x}_{i,t}]$, $\mathbf{x}_{i,\tau} = (e_{i,\tau}, g_{i,\tau}, p_{i,\tau}, s_{i,\tau}^{coal}, s_{i,\tau}^{oil}, s_{i,\tau}^{gas})$, (2)

~~with [equation removed or moved to S1 Text] and [equation removed or moved to S1 Text] ([equation removed or moved to S1 Text] in Experiment C), giving 3,024 samples (2,052 in Experiment C). The LOCO task is to learn [equation removed or moved to S1 Text] such that [equation removed or moved to S1 Text] generalizes to countries absent from training. Evaluation uses grouped five-fold cross-validation with countries as groups, so the test country's entire target series is invisible during training; features are standardized per fold as in equation (4),~~

where $\mathbf{x}_{i,\tau}$ is the vector of the $F = 6$ covariates of country i in year τ ($F = 7$ in Experiment C, which appends $s_{i,\tau}^{lc}$) and $\mathbf{X}_{i,t}$ is the window of $W \times F$ values. With 34 years per country this gives 27 samples per country, with target years 1997 to 2023: 3,024 samples for the 112 countries of Experiment A and 2,052 for the 76 countries of Experiment C. A model is a function f that returns the estimate $\hat{y}_{i,t} = f(\mathbf{X}_{i,t})$; the estimate of per-capita emissions is $\hat{c}_{i,t} = \exp(\hat{y}_{i,t})$. Models were evaluated by leave-country-out cross-validation. The countries were divided into five folds of 22 or 23 countries, with all years of a country in the same fold; each fold was held out in turn, and the model was fitted to the countries of the other four. Every result is therefore for countries none of whose data were used to fit the model that produced the estimate. Within each training set, one-seventh of the countries (rounded down: 12 countries in Experiment A, 8 in Experiment C) formed an inner validation set for early stopping, hyperparameter search and the calibration of prediction intervals. In each fold the covariates were standardized (Eq 3):

~~[equation removed or moved to S1 Text] [equation removed or moved to S1 Text]~~ $\tilde{x} = \frac{x - \mu_{tr}}{\sigma_{tr}}$, (3)

with [equation removed or moved to S1 Text] computed on training folds only. Within each training fold, one-seventh of the *countries* form an inner validation set for early stopping and hyperparameter selection, preserving country-disjointness at every level.

Scope of the "no own-emissions" claim. The target country's emission *history* is withheld by construction, and no lagged value of [equation removed or moved to S1 Text] enters the feature set. The three fuel-share covariates are, however, ratios computed from that country's own CO_2 accounts, [equation removed or moved to S1 Text] and so on. They reveal no information about the *level* of emissions, which is the estimand, but they are not obtainable from a country that has no emission accounts at all; in a genuine deployment they would be derived from national energy balances instead. The task therefore simulates a country whose *fuel mix* is characterized but whose *emission level* is not. Section 4.2 reports a covariate ablation that removes the fuel shares entirely; it shows their contribution is under one percentage point of median error, so the distinction is minor in practice. Section 5.3 returns to the point.

3.3 Kaya-motivated linear backbone

The Kaya identity decomposes national emissions as [equation removed or moved to S1 Text] [27,28]. Dividing by population and taking logarithms yields equation (5),

[equation removed or moved to S1 Text]

i.e., log per-capita emissions equal log per-capita energy plus log carbon intensity of energy. Because carbon intensity is, to first order, a mixture over fuel shares, equation (5) implies that a linear model in [equation removed or moved to S1 Text] should capture the bulk of cross-country variance, an implication our results confirm.

Two ridge estimators appear in this paper and must be distinguished. The **benchmark ridge** of Table 1 is fitted on the flattened window [equation removed or moved to S1 Text], equation (6),

[equation removed or moved to S1 Text]

The **warm-start ridge** used to initialize the hybrid's linear head (Section 3.5) is the more parsimonious variant fitted on the most recent window step [equation removed or moved to S1 Text] alone, so that the head has one weight per covariate. Fitted on the last step, ridge attains RMSE 3.343 and R^2 0.761 in LOCO; fitted on the full window, RMSE 3.056 and R^2 0.800. Table 1 reports the latter.

3.4 Benchmark models

Thirteen models are compared under identical folds and inputs: a training-set mean predictor; ridge regression (6); k-nearest neighbours ($k = 7$, distance-weighted); support vector regression (RBF kernel, $C = 10$, $\varepsilon = 0.01$); random forest (600 trees, depth ≤ 14); XGBoost [52] (800 trees, $\eta = 0.03$, depth 6, subsample 0.85); LightGBM [53] (800 trees, $\eta = 0.03$, 63 leaves, subsample 0.85, colsample 0.85, L2 penalty 1.5); LSTM; GRU; CNN-LSTM; CNN-BiLSTM-Attention; and the two proposed hybrids KG-CBA and PSO-KG-CBA. Classical learners operate on the flattened window [equation removed or moved to S1 Text]; deep sequence models consume [equation removed or moved to S1 Text] directly. The LSTM cell [54] is given by equations (7)–(12):

[equation removed or moved to S1 Text]

[equation removed or moved to S1 Text]

[equation removed or moved to S1 Text]

[equation removed or moved to S1 Text]

[equation removed or moved to S1 Text]

[equation removed or moved to S1 Text]

~~with [equation removed or moved to S1 Text] the logistic function and [equation removed or moved to S1 Text] the Hadamard product. The GRU uses update and reset gates, equations (13)–(15),~~

~~[equation removed or moved to S1 Text]~~

~~[equation removed or moved to S1 Text]~~

~~[equation removed or moved to S1 Text]~~

~~The CNN-LSTM stacks two one-dimensional convolutions, equation (16),~~

~~[equation removed or moved to S1 Text]~~

~~before an LSTM encoder. All fixed deep baselines use 48 filters/units, kernel 3, dropout 0.25, Adam (10^{-3}), batch 96, ≤90 epochs with early stopping (patience 9) on the inner country-disjoint validation loss of equation (17),~~

~~[equation removed or moved to S1 Text]~~

~~3.5 Kaya-guided CNN-BiLSTM-Attention (KG-CBA)~~

~~The proposed architecture decomposes the prediction into a physics-guided linear term and a learned residual, equation (18):~~

where μ_{tr} and σ_{tr} are the mean and the standard deviation of the covariate over the training countries of that fold; $\tilde{\mathbf{x}}$ and $\tilde{\mathbf{X}}$ denote the standardized covariate vector and window.

The fuel shares are ratios of a country's own fuel-specific emissions to its total emissions. They carry no information on the level of emissions, which is the quantity estimated, and no lagged value of the target enters any model. They would not, however, be available for a country without any emission accounts, where they would have to be derived from the national energy balance; the sensitivity analyses therefore include a model without them.

Models

Kaya identity and the linear reference model

The Kaya identity [25] writes national emissions as a product of four factors (Eq 4):

$$C = P \times \frac{Y}{P} \times \frac{Q}{Y} \times \frac{C}{Q}, \qquad (4)$$

where C is total fossil CO_2 emissions, P population, Y GDP and Q primary energy consumption. C appears on both sides because the expression is an identity; the factors are income per person (Y/P), the energy intensity of income (Q/Y) and the carbon intensity of energy (C/Q). Dividing by population and taking logarithms gives per-capita emissions $c = C/P$ in terms of energy per capita $E = Q/P$ (Eq 5):

$$\ln c = \ln E + \ln \frac{C}{Q}. \qquad (5)$$

Carbon intensity depends mainly on the fuel mix, so Eq 5 suggests that a model that is linear in the covariates of Eq 2 should explain most of the variation in y between countries. The linear reference model is ridge regression [35] on the standardized window (Eq 6):

$$\widehat{\boldsymbol{\beta}} = \underset{\boldsymbol{\beta}, \beta_0}{\operatorname{argmin}} \sum_{n=1}^{N_{tr}} (y_n - \beta_0 - \boldsymbol{\beta}^{\top} \tilde{\mathbf{z}}_n)^2 + \lambda \parallel \boldsymbol{\beta} \parallel_2^2, \qquad (6)$$

where n indexes the N_{tr} training samples, $\tilde{\mathbf{z}}_n$ is the standardized window $\widetilde{\mathbf{X}}_n$ written as one vector of $W \times F = 48$ values, $\boldsymbol{\beta}$ and β_0 are the coefficients and the intercept, and $\lambda = 1$ is the penalty on the squared size of the coefficients.

Benchmark models

Thirteen models were compared (Table 2). They span the model families used in the emission-modelling literature, in increasing order of flexibility: a naive baseline; the linear reference

model; five classical machine-learning models (k-nearest neighbours, support vector regression [36], random forest [37], XGBoost [38] and LightGBM [39]); four deep sequence networks that are the building blocks of published hybrids (long short-term memory, LSTM [40]; gated recurrent unit, GRU [41]; a convolutional network followed by an LSTM, CNN-LSTM; and a convolutional network followed by a bidirectional LSTM with attention [42], CNN-BiLSTM-Attn); and the two Kaya-guided hybrids introduced below. The mean predictor is not a fitted model in the usual sense: it assigns to every test sample the mean of y over the training countries and shows the error obtained without any covariate information. All models used the same folds and inputs. The equations of the recurrent, convolutional and attention layers are standard and are given, with all symbols defined, in S1 Text.

Table 2. Models and their settings.

Model	Family	Settings
Mean predictor	Naive baseline	Mean of y over the training countries; no covariates
Ridge regression	Linear	Penalty $\lambda = 1$ (Eq 6)
k-nearest neighbours (kNN)	Classical machine learning	7 neighbours, distance-weighted
Support vector regression (SVR)	Classical machine learning	Radial basis kernel, C = 10, ε = 0.01
Random forest	Classical machine learning	600 trees, maximum depth 14
XGBoost	Classical machine learning	800 trees, learning rate 0.03, maximum depth 6, row subsample 0.85
LightGBM	Classical machine learning	800 trees, learning rate 0.03, 63 leaves (Experiment C: also row and column subsample 0.85, L2 penalty 1.5)
LSTM	Deep network	LSTM, 48 units
GRU	Deep network	GRU, 48 units
CNN-LSTM	Deep network	Two convolutional layers (48 filters, kernel size 3), then LSTM (48 units)
CNN-BiLSTM-Attn	Deep network	Two convolutional layers (48 filters, kernel size 3), bidirectional LSTM (48 units), additive attention
KG-CBA	Kaya-guided hybrid	Linear head started at the ridge solution; residual branch as CNN-BiLSTM-Attn with a 32-unit dense layer (Eq 7); fixed settings of the deep networks: 48 filters, kernel size 3, 48 LSTM units, dropout 0.25, learning rate 10^{-3}, batch size 96

Model	Family	Settings
PSO-KG-CBA	Kaya-guided hybrid	As KG-CBA, with settings selected by PSO: 37 filters, kernel size 4, 40 LSTM units, dropout 0.231, learning rate 9.96×10^{-4}, batch size 48
Ensemble	Exploratory	Mean of the estimates of ridge, GRU and PSO-KG-CBA

Classical models use the window as one vector of 48 values; deep networks use it as an 8×6 array. Deep networks were trained with the Adam optimizer on the mean squared error of y (dropout 0.25, learning rate 10^{-3}, batch size 96, at most 90 epochs), and training stopped when the error on the inner validation countries had not improved for 9 epochs. PSO search ranges: 16–64 filters; kernel size 2, 3 or 4; 16–96 LSTM units; dropout 0.05–0.45; learning rate 2×10^{-4} to 3×10^{-3}; batch size 48, 64, 96 or 128.

Kaya-guided hybrid network

KG-CBA writes the estimate as the sum of a linear term and a learned residual (Fig 3; Eq 7):

~~[equation removed or moved to S1 Text]~~ $\hat{y} = \left(\boldsymbol{\beta}_K^{\top}\tilde{\mathbf{x}}_W + \beta_{K,0}\right) + r_{\boldsymbol{\theta}}(\tilde{\mathbf{X}}),$ (7)

~~The Kaya head is a single dense unit acting on the most recent window step, *warm-started* at the fold-specific last-step ridge solution (Section 3.3) and refined jointly during training. The residual branch applies two Conv1D layers (16), a bidirectional LSTM whose forward and backward states are concatenated in equation (19),~~

~~[equation removed or moved to S1 Text]~~

~~and additive temporal attention [55] that pools the sequence into a context vector through equations (20)–(22),~~

~~[equation removed or moved to S1 Text]~~

~~[equation removed or moved to S1 Text]~~

~~[equation removed or moved to S1 Text]~~

followed by a 32-unit dense layer, dropout, and a linear residual output whose kernel is initialized near zero so that training starts from the ridge solution. This design lets gradient descent spend capacity only where the log-linear Kaya approximation fails.

3.6 Particle swarm optimization of the hybrid

Six hyperparameters are tuned: convolution filters [equation removed or moved to S1 Text], kernel size [equation removed or moved to S1 Text], BiLSTM units [equation removed or moved to S1 Text], dropout [equation removed or moved to S1 Text], learning rate [equation removed or moved to S1 Text] (log-scale), and batch size [equation removed or moved to S1 Text]. PSO [29] evolves six particles for five iterations (30 evaluations). Particle [equation removed or moved to S1 Text] updates according to equations (23)–(24) as

[equation removed or moved to S1 Text]

[equation removed or moved to S1 Text]

with inertia $\omega = 0.7$, cognitive and social constants $c_1 = c_2 = 1.5$, [equation removed or moved to S1 Text], velocities clamped to 30% of each range, and fitness defined as inner-validation MSE on a country-disjoint split of the first training fold. The converged configuration (37 filters, kernel 4, 40 units, dropout 0.231, learning rate 9.96×10^{-4}, batch 48; validation MSE 0.0522) is retrained on all five folds, denoted PSO-KG-CBA (E-PSO-KG-CBA on the enriched panel).

3.7 Uncertainty quantification: conformalized Monte-Carlo dropout

Epistemic uncertainty is estimated by Monte-Carlo dropout [32]: with dropout active at inference, [equation removed or moved to S1 Text] stochastic forward passes yield the predictive moments of equation (25),

[equation removed or moved to S1 Text]

Because raw MC-dropout intervals are severely over-confident, we calibrate them per fold by split-conformal inference [33,34]: on each fold's country-disjoint inner validation set, normalized nonconformity scores, equation (26),

[equation removed or moved to S1 Text]

are computed, the level-adjusted quantile [equation removed or moved to S1 Text] is extracted, and the test-fold interval, equation (27), is

[equation removed or moved to S1 Text]

Coverage validity assumes exchangeability between calibration and test units [35,48]; grouped country sampling violates this, and Section 4.5 quantifies the resulting shortfall rather than concealing it.

3.8 Explainability

Global attributions use SHAP [36]. For a tree ensemble [equation removed or moved to S1 Text] and feature [equation removed or moved to S1 Text] , the attribution is given by equation (28),

[equation removed or moved to S1 Text]

computed exactly via TreeExplainer on the enriched LightGBM and aggregated over the 56 window columns by covariate and by lag. Because the flattened design matrix is lag-major (column index [equation removed or moved to S1 Text]), the attribution array is reshaped to [equation removed or moved to S1 Text] before aggregation; aggregating under the transposed layout silently interchanges covariates and lags. Temporal importance of the hybrid is read directly from the learned attention weights [equation removed or moved to S1 Text] in (21).

3.9 Evaluation metrics and clustered statistical testing

With observed [equation removed or moved to S1 Text] and predicted [equation removed or moved to S1 Text] on the raw scale, accuracy is scored by equations (29)–(33);

[equation removed or moved to S1 Text]

[equation removed or moved to S1 Text]

[equation removed or moved to S1 Text]

[equation removed or moved to S1 Text]

[equation removed or moved to S1 Text]

For the temporal experiment, the mean absolute scaled error against the in-sample naive forecast [22], equation (34),

[equation removed or moved to S1 Text]

flags genuine skill (MASE < 1); the denominator is computed per country on the training period, so each country's errors are scaled by its own naive difficulty. Because the denominator is in-sample while the numerator is out-of-sample, persistence itself need not attain MASE = 1: it scores 1.198 at h = 1 in the 2021–2023 window, reflecting genuinely higher post-2020 volatility, and 0.995 in the pre-pandemic replication.

Clustered inference. The out-of-fold sample contains 27 repeated years for each of 112 countries, so country-years are far from independent. We therefore report three quantities for every model pair: a paired Wilcoxon signed-rank test on absolute percentage error at the observation level (the conventional but anticonservative test, retained only for comparison); a paired Wilcoxon test on *per-country median* APE, with [equation removed or moved to S1 Text] clusters; and a cluster bootstrap that resamples **countries** with replacement (10,000 draws) to obtain a 95% confidence interval and two-sided p-value for the difference in pooled median APE. Conclusions are drawn from the clustered tests. Software: Python 3.12.3, TensorFlow

2.21.0 (CPU), XGBoost 3.4.0, LightGBM 4.7.0, scikit-learn 1.8.0, SHAP 0.52.0; global seed 42 with per-fold offsets.

3.10 Experiment B: temporal forecasting protocol

To situate the LOCO results against conventional forecasting, we recast the panel as an [equation removed or moved to S1 Text]-step-ahead task in which the target country's own emission history *is* available, so that persistence is well-defined. Predictors are the eight-year window of the six covariates augmented with the country's own lagged log emissions (56 columns); withholding the latter would handicap the learners relative to persistence and make the comparison uninformative. Targets are formed in two ways: a *level* formulation predicting [equation removed or moved to S1 Text] directly, and a *log-growth* formulation predicting [equation removed or moved to S1 Text] . Models are trained on all pairs with [equation removed or moved to S1 Text] and tested on target years 2021-2023, with a pre-pandemic replication trained through 2012 and tested on 2015-2019. Baselines are persistence, [equation removed or moved to S1 Text] , and drift, [equation removed or moved to S1 Text] with [equation removed or moved to S1 Text] the mean of the last three log-changes. Six learners are evaluated under each formulation, and each learner is compared against persistence by a cluster bootstrap over countries.

4. Results

4.1 Panel characteristics

Figure 1 summarizes the panel: mean per-capita emissions decline gently after 2007 while trajectories diverge. Figure 2 confirms the Kaya expectation: ln CO_2pc correlates at $r = 0.93$ with log energy per capita and $r = 0.82$ with log GDP per capita, while fuel shares carry weaker

marginal correlations. Continental distributions (Figure 3) span three orders of magnitude. The framework is depicted in Figure 4 and the KG-CBA architecture in Figure 5.

[embedded figure image, now supplied as a separate figure file]

Figure 1. Per-capita CO2 in the 112-country panel, 1990-2023: (a) panel mean with interquartile range; (b) selected national trajectories (log scale).

[embedded figure image, now supplied as a separate figure file]

Figure 2. Pearson correlation matrix of the log-transformed target and the six core covariates (3,808 country-years).

[embedded figure image, now supplied as a separate figure file]

Figure 3. Distribution of per-capita CO2 by continent (log scale).

[embedded figure image, now supplied as a separate figure file]

Figure 4. Overview of the experimental framework.

[embedded figure image, now supplied as a separate figure file]

Figure 5. The Kaya-guided CNN-BiLSTM-Attention (KG-CBA) architecture.

4.2 Experiment A: 112-country LOCO benchmark

Table 1 reports out-of-fold accuracy for all thirteen models plus the ensemble; Figure 8 shows the comparison and Figure 9 predicted-observed agreement across four orders of magnitude. Three patterns stand out. First, the task is learnable but hard: the mean predictor fails catastrophically (MAPE 337.8%), whereas every covariate-using learner achieves MAPE below 50%. Second, the Kaya mechanism dominates. Ridge regression, linear in log space, attains $R^2 = 0.800$ and MdAPE 16.3%, outperforming every tree ensemble and every unguided deep network, including the plain CNN-BiLSTM-Attention (MdAPE 20.8%); both gaps survive country-clustered testing (ridge vs LightGBM $p = 0.006$; ridge vs CNN-BiLSTM-Attention $p = 0.002$).

Third, physics-guidance closes the gap from the deep side: adding the Kaya head lifts the hybrid's MdAPE from 20.8% to 15.7%, and both KG-CBA and PSO-KG-CBA significantly outperform the unguided architecture under clustering ($p = 0.002$ for each).

The ensemble's advantage over ridge does not survive clustered inference. Table 2 contrasts the two testing regimes. Averaging ridge, GRU and PSO-KG-CBA gives the best point accuracy in the study (MAPE 23.5%, MdAPE 15.1%), and an observation-level Wilcoxon test returns $p = 7 \times 10^{-10}$ against ridge. But the per-country test gives $p = 0.098$ and the cluster bootstrap $p = 0.22$, with a 95% interval for the median-APE difference of [-2.20, +0.41] percentage points that comfortably spans zero. We therefore claim no significant improvement of the ensemble over ridge alone. The ensemble does significantly beat GRU ($p = 0.0006$) and the PSO-tuned hybrid ($p = 0.022$), so the gain is real relative to the deep constituents and illusory relative to the linear one. The composition of the ensemble was chosen after inspecting out-of-fold performance and is therefore not the product of a protected selection procedure; this is a further reason to treat its margin cautiously.

Covariate ablation. Because three of the six covariates are ratios derived from the target country's own CO_2 accounts (Section 3.2), we retrained ridge without them. Using only energy per capita, GDP per capita and population, MdAPE rises from 16.3% to 16.9% and R^2 falls from 0.800 to 0.775; using log energy per capita alone, MdAPE is 17.9% and R^2 0.723 (Table B6). The fuel shares are therefore worth well under one percentage point of median error. Essentially all of the achievable LOCO accuracy is obtainable from energy-economic covariates that carry no emission-account provenance whatsoever, so the estimator is deployable for a country with no inventory of any kind.

[embedded figure image, now supplied as a separate figure file]

Figure 6. PSO convergence over 30 fitness evaluations. (Supplementary)

[embedded figure image, now supplied as a separate figure file]

Figure 7. Fold-1 training dynamics of the four fixed deep baselines. (Supplementary)

[embedded figure image, now supplied as a separate figure file]

Figure 8. Experiment A leave-country-out error by model.

[embedded figure image, now supplied as a separate figure file]

Figure 9. Out-of-fold predicted versus observed per-capita CO2 (log-log).

Table 1. Out-of-fold LOCO performance, Experiment A (112 countries, 6 covariates).

Model	RMSE	MAE	MAPE (%)	MdAPE (%)	R^2
Mean	7.415	4.408	337.8	63.3	−0.178
kNN	5.119	2.729	49.7	39.0	0.439
SVR	4.881	2.100	32.3	21.3	0.490
Random forest	4.233	1.824	28.7	19.1	0.616
XGBoost	4.043	1.775	28.0	18.4	0.650
LightGBM	3.943	1.813	28.4	18.6	0.667
LSTM	3.815	1.703	27.6	18.2	0.688
CNN-LSTM	3.773	1.707	28.1	20.8	0.695
CNN-BiLSTM-Attn	3.812	1.790	28.6	20.8	0.689
GRU	3.629	1.644	25.1	18.5	0.718
Ridge (Kaya log-linear)	**3.056**	**1.443**	25.2	16.3	**0.800**
KG-CBA (ours)	4.595	1.898	26.0	**15.7**	0.548
PSO-KG-CBA (ours)	4.748	1.914	27.2	16.7	0.517

Model	RMSE	MAE	MAPE (%)	MdAPE (%)	R^2
Ensemble (Ridge + GRU + PSO-KG-CBA)	3.557	1.530	**23.5**	15.1	0.729

Table 2. Observation-level versus country-clustered inference on paired absolute percentage error. Δ is the difference in pooled median APE (percentage points); the interval is a 95% cluster bootstrap over countries (10,000 draws). Conclusions follow the clustered columns.

Experiment	Comparison	MdAPE A / B (%)	p (observation-level)	p (per-country)	Δ [95% CI]	p (cluster bootstrap)	Conclusion
A	Ensemble vs Ridge	15.1 / 16.3	6.9×10^{-10}	0.098	−0.76 [−2.20; +0.41]	0.22	not significant
A	Ensemble vs GRU	15.1 / 18.5	1.8×10^{-30}	0.007	−2.57 [−3.77; −1.20]	0.0006	significant
A	Ensemble vs PSO-KG-CBA	15.1 / 16.7	3.6×10^{-31}	0.074	−1.33 [−2.85; −0.22]	0.022	significant
A	PSO-KG-CBA vs	16.7 / 20.8	6.3×10^{-22}	0.004	−3.51 [−5.97	0.0016	significant

Experiment	Comparison	MdAPE A / B (%)	p (observation-level)	p (per-country)	Δ [95% CI]	p (cluster bootstrap)	Conclusion
	CNN-BiLSTM-Attn				; −0.97]		
A	KG-CBA vs CNN-BiLSTM-Attn	15.7 / 20.8	4.9×10^{-26}	0.004	−3.42 [−5.84 ; −1.21]	0.0018	significant
A	Ridge vs LightGBM	16.3 / 18.6	8.3×10^{-25}	0.026	−3.10 [−4.91 ; −1.03]	0.0058	significant
A	Ridge vs CNN-BiLSTM-Attn	16.3 / 20.8	3.1×10^{-34}	0.007	−4.29 [−7.05 ; −1.64]	0.0024	significant
C	Ensemble vs Ridge	7.35 / 7.72	0.483	0.491	−0.20 [−0.66 ; +0.36]	0.56	not significant

Experiment	Comparison	MdAPE A / B (%)	p (observation-level)	p (per-country)	Δ [95% CI]	p (cluster bootstrap)	Conclusion
C	E-PSO-KG-CBA vs LightGBM Ensemble	8.09 / 13.10	1.7×10^{-56}	0.0002	−3.85 [−6.03; −1.95]	0.0001	significant
C	Ensemble vs E-PSO-KG-CBA	7.35 / 8.09	5.1×10^{-31}	0.004	−0.86 [−1.45; −0.35]	0.0014	significant

4.3 Error anatomy and the covariate-availability effect (Experiment C)

A Pareto decomposition of squared error (Figure 10) reveals extreme concentration: the six worst-fitting economies contribute 79.3% of the ensemble's total squared error and 84.3% for the hybrid alone. The *composition* of that group is only partly stable across models. Singapore, Qatar and Kuwait appear in the top six under all six models examined; Iceland, Norway, Luxembourg and Saudi Arabia exchange places depending on the estimator. What the group shares is structural: hydro- and geothermal-dominant grids (Norway, Iceland), hydrocarbon-exporting states (Qatar, Kuwait, Saudi Arabia), and city-states with bunker-fuel and finance-driven energy profiles (Singapore, Luxembourg). The residuals, in other words, *locate the failure of the Kaya carbon-intensity proxy*, and the identity of the specific worst six should not be over-interpreted.

[embedded figure image, now supplied as a separate figure file]

Figure 10. Pareto decomposition of squared error by country. Dark red marks the three economies that are outliers under every model examined; orange marks the remainder of this model's top six.

Experiment C tests the implied remedy. Adding the low-carbon share of primary energy on the 76-country sub-panel substantially improves accuracy (Table 4; Figure 11 shows the like-for-like comparison on identical countries). On those same 76 countries, ridge improves from MAPE 22.2% / MdAPE 14.3% to 14.1% / 7.7%, and the ensemble from 21.9% / 13.7% to 14.5% / 7.35%. Median error is therefore roughly halved (−46% for ridge, −46% for the ensemble) while mean error falls by about a third (−36% and −34%). The ensemble is statistically indistinguishable from ridge on this panel ($p = 0.56$) but significantly better than the hybrid alone ($p = 0.0014$), and the hybrid significantly beats LightGBM ($p = 0.0001$). The practical reading is direct: where energy-mix statistics exist, a single covariate is worth more than any architectural innovation evaluated here.

[embedded figure image, now supplied as a separate figure file]

Figure 11. Effect of adding the low-carbon energy share (identical 76 countries).

Table 3. Imputation audit for the 112-country panel (3,808 country-years). "Longest run" is the maximum number of consecutive filled years for any single country.

Variable	Filled country-years	% of panel	Longest run (yr)
Per-capita CO_2 (target)	0	0.00	0
Population	0	0.00	0
Oil share	0	0.00	0
Energy per capita	28	0.74	16

Variable	Filled country-years	% of panel	Longest run (yr)
GDP per capita	114	2.99	—
Coal share	244	6.41	26
Gas share	291	7.64	31
Low-carbon share (76-country sub-panel)	0	0.00	0

Countries with at least one filled run exceeding three years: 25 of 112. Exceeding ten years: 19 of 112.

Table 4. Out-of-fold LOCO performance, Experiment C (76 countries, 7 covariates).

Model	RMSE	MAE	MAPE (%)	MdAPE (%)	R^2
LightGBM	4.520	2.048	22.8	13.1	0.622
E-PSO-KG-CBA (ours)	3.682	1.593	16.0	8.1	0.749
Ridge	**3.037**	**1.284**	**14.1**	7.7	**0.829**
Ensemble (Ridge + E-PSO-KG-CBA)	3.306	1.409	14.5	**7.35**	0.798

4.4 Regional performance and African case studies

Table 5 gives the full continental breakdown, which is more nuanced than a two-region comparison suggests. On median APE, Europe is easiest (12.5%) and North America hardest (19.6%); Africa (16.8%) sits between Oceania (18.8%) and Asia (15.8%). On *mean* APE the ordering nearly reverses and Africa (26.6%) is close to Europe (23.6%). Africa is therefore not uniformly the hardest region; rather, its error distribution has a heavier right tail, consistent with a small number of countries whose energy structure is poorly proxied.

Table 5. Ensemble absolute percentage error by continent, Experiment A.

Continent	Median APE (%)	Mean APE (%)	75th pct (%)	n (country-years)
Europe	12.50	23.60	25.67	1053
South America	14.84	20.45	26.24	216
Asia	15.83	23.43	26.93	999
Africa	16.77	26.63	26.77	459
Oceania	18.76	18.03	25.22	54
North America	19.63	21.46	27.41	243

Among African economies with fully observed covariate series, leave-country-out accuracy is strong: South Africa attains a median APE of 3.4% and Egypt 5.5% (both with no imputed country-years), and Nigeria 8.0%. Figure 12 shows observed trajectories tracked within the 90% conformal bands of a model that never saw these countries during training. For inventory gap-filling in data-sparse settings, this is the deployment-relevant accuracy level.

Ghana's leave-country-out median APE is 7.4%, but this figure must be read with care and we do not offer it as the study's headline African result: Ghana carries a 26-year filled run in its fuel-share covariates, so its apparently smooth covariate trajectory is substantially an artefact of the edge-extension rule rather than a measured series. The same caveat applies to Cameroon (6.1%, 26 filled years), Rwanda (40.8%, 25 years), Benin (28.0%, 24 years) and Niger (44.8%, 29 years). Restricting the LOCO benchmark to the 87 countries with no filled run longer than three years degrades ridge from MdAPE 16.3% to 17.6% and MAPE 25.2% to 27.5%, confirming that heavily filled countries are *easier* than average and that the full-panel numbers are mildly optimistic. Ghana is thus better read as an illustration of the data-scarcity problem this paper is about than as a demonstration of accuracy under it.

[embedded figure image, now supplied as a separate figure file]

Figure 12. African case studies with 90% split-conformal intervals. Countries carrying long imputed runs are flagged in the panel titles.

4.5 Uncertainty calibration

Raw MC-dropout intervals are severely over-confident: at a nominal 90% level they cover only 13.2% of observations with a mean width of 0.47 t person^{-1}. Per-fold split-conformal calibration widens them to an empirical coverage (PICP) of 81.9% with mean interval width 5.68 t person^{-1}. The residual shortfall is attributable to non-exchangeability: calibration and test units belong to disjoint country groups, and fold-level conformal quantiles vary by nearly an order of magnitude (8.8 to 69.2), reflecting heterogeneous country difficulty. This is an honest limitation of split-conformal inference under grouped sampling [35,48], and Section 5.3 outlines group-aware alternatives.

4.6 Explainability

SHAP attributions on the enriched panel (Figure 13) are dominated by log energy per capita (mean |SHAP| = 0.114), an order of magnitude above the low-carbon share (0.016), gas share (0.008), log population (0.007) and the remaining covariates. Lag-wise attribution decays sharply and monotonically from the most recent window position (t: 0.118; t−1: 0.027; all earlier lags below 0.008), indicating that for this annual-resolution task the contemporaneous covariate vector carries almost all of the tree ensemble's signal and the eight-year window adds little. Both findings are exactly what the Kaya structure of equation (5) predicts, since per-capita energy is the identity's leading term.

This is worth stating plainly because it does *not* corroborate the covariate-availability argument of Section 5.2 in the way one might expect: SHAP ranks the low-carbon share well below energy

per capita even though adding that covariate halves median error. The two results are compatible — SHAP measures a feature's contribution to predictions within a fitted model, whereas Experiment C measures the change in generalization error from making the feature available at all, and a covariate can be decisive for the hardest countries while contributing modestly on average. The Experiment C contrast, not the SHAP ranking, is the evidence for the policy argument.

The hybrid's attention weights (Figure 14) range from 0.095 to 0.142 against a uniform 0.125, declining gently from earlier to more recent window positions. Attention is therefore close to uniform pooling, and its value in this architecture lies more in adaptive aggregation than in sharp temporal selection.

[embedded figure image, now supplied as a separate figure file]

Figure 13. SHAP attributions on the enriched panel, by covariate and by window position.

[embedded figure image, now supplied as a separate figure file]

Figure 14. Mean temporal attention weights by window position. (Supplementary)

4.7 Experiment B: the persistence benchmark in temporal forecasting

When the same data are recast as conventional forecasting with the target country's own history available, naive persistence proves a formidable benchmark, but not an insurmountable one (Table 6, Figure 15).

At one-year horizons in the 2021–2023 window, persistence attains MAE 0.317 t person^{-1}; only XGBoost improves on it, and by a small and marginally significant margin (level formulation MAE 0.296, $\Delta = -0.021$, cluster-bootstrap $p = 0.045$). Every other learner ties or loses. At three and five years, more learners edge ahead: random forest in log-growth form reaches MAE 0.481 at $h = 3$ ($p = 0.031$) and 0.621 at $h = 5$ ($p = 0.039$), and XGBoost 0.599 at $h = 5$ ($p = 0.0001$).

~~Several learners attain MASE below unity at the longer horizons, with ridge in log-growth form reaching 0.867 at h = 5.~~

~~Two qualifications matter more than these point gains. First, the margins are small: the best improvement over persistence at any horizon is 19% of MAE, against a drift baseline that is *worse* than persistence at every horizon in both windows. Second, and decisively, the ranking does not replicate. In the pre-pandemic 2015-2019 window, no learner significantly beats persistence at h = 1 or h = 3, and only random-forest log-growth does so at h = 5 (p = 0.047); ridge log-growth attains MASE 0.945 at h = 1 but its advantage is not significant under clustering. The learner that wins in one window is generally not the learner that wins in the other. The implication for the literature is therefore narrower than a blanket claim that learning fails, but no less pointed. Annual national per-capita CO_2 is close enough to a random walk that persistence must always be reported; margins over it are of order 5-15% of MAE, are horizon-dependent, and are not stable across evaluation periods. Reported temporal R^2 values near unity, published without a naive baseline, largely re-describe persistence rather than demonstrate skill. The log-growth target formulation matters more than the choice of learner. The LOCO formulation is immune to this artefact because the target country's history is withheld by construction.~~

~~[embedded figure image, now supplied as a separate figure file]~~

~~Figure 15.~~ ~~Experiment B: MASE against the in-sample naive benchmark, both evaluation windows. Values below the dashed line indicate skill.~~

~~Table 6.~~ ~~Experiment B, MAE (t CO_2 person^{-1}) and MASE by horizon. Bold marks a significant improvement over persistence under a country cluster bootstrap ($p < 0.05$).~~

Model	h=1 MAE	h=3 MAE	h=5 MAE	h=1 MASE	h=3 MASE	h=5 MASE
Test window 2021–2023						
Persistence	0.317	0.531	0.738	1.198	1.030	1.101
Drift	0.380	0.680	0.933	1.425	1.363	1.424
Ridge (log-growth)	0.339	0.542	0.678	1.193	0.926	0.867
Random forest (log-growth)	0.307	**0.481**	**0.621**	1.145	0.950	0.939
XGBoost (level)	**0.296**	0.506	**0.599**	1.188	0.983	0.963
LightGBM (log-growth)	0.307	0.528	0.621	1.202	1.014	0.971
Replication window 2015–2019						
Persistence	0.247	0.468	0.682	0.995	1.084	1.241
Drift	0.285	0.683	1.043	1.154	1.531	1.920
Ridge (log-growth)	0.239	0.429	0.598	0.945	0.990	1.108
Random forest (log-growth)	0.246	0.453	**0.567**	0.976	1.028	1.088
XGBoost (level)	0.302	0.457	0.642	1.213	1.155	1.114
LightGBM (log-growth)	0.269	0.516	0.647	1.141	1.204	1.238

5. Discussion

5.1 Why physics guidance matters more than depth

The central empirical message is structural. In log space, the Kaya identity makes cross-country emission variation approximately linear in energy and fuel-mix covariates; unguided deep networks spend most of their capacity rediscovering this plane from roughly 2,400 training samples and still fall significantly short of ridge regression. Encoding the plane explicitly, warm-starting a linear head at the ridge solution and letting the deep branch model only residuals, recovers the deep models' competitiveness: both Kaya-guided hybrids significantly outperform the identical unguided architecture on median error. Yet the hybrids' remaining weakness is instructive: their R^2 lags ridge badly (0.55 versus 0.80) because residual capacity occasionally amplifies errors on extreme-scale economies, so they win on the median and lose on the mean square.

Ensembling narrows this asymmetry but, on the evidence of clustered inference, does not beat ridge alone. That negative result is the honest one, and it sharpens rather than weakens the paper's message. For applied emission modelling the ordering of returns is: correct covariates first, physics-informed structure second, ensembling third, architectural novelty last. A study that had tested at the observation level would have reported the ensemble as decisively best ($p < 10^{-9}$) and drawn the opposite methodological lesson.

5.2 Policy implications for inventory gap-filling

Experiment C converts a modelling observation into a statistical-capacity argument. One additional covariate, the low-carbon share of primary energy, halves leave-country-out median error and collapses the outlier problem, because it measures exactly the carbon-intensity term

that fossil-fuel shares cannot see. Most African economies lack this statistic in the international energy lineages.

The imputation audit sharpens the same point from an unexpected direction. Of the 25 panel countries carrying filled runs longer than three years, the overwhelming majority are low- and middle-income, and nine are African. These are countries for which the international energy statistics are so thin that a balanced panel can only be constructed by carrying a single observation across two or three decades. Ghana is a case in point: it enters this study's panel only because 26 years of its fuel-share series are extended from a handful of observations. The binding constraint on emission transparency in these countries is therefore not model capacity and not even the low-carbon covariate specifically, but the absence of basic, annually updated national energy balances. Investment there would yield disproportionate returns for Enhanced Transparency Framework implementation in non-Annex I countries [3-6,43], plausibly more than further model development. Where such balances do exist, as for South Africa and Egypt, leave-country-out estimation already achieves 3-6% median error without the country ever being observed.

5.3 Limitations and threats to validity

Six limitations qualify the findings. First, the balanced-panel requirement retains 112 countries, and 25 of these are retained only through long constant extension of their covariate series; the sensitivity analysis of Section 4.4 shows the full-panel results are mildly optimistic as a consequence, and the states most in need of gap-filling remain under-represented even so. Second, the three fuel-share covariates are ratios derived from CO_2 accounts rather than energy balances. They carry no information about the emission *level*, but a country with no inventory at all would not have them. The ablation in Section 4.2 bounds this concern tightly: removing them

raises MdAPE by only 0.6 percentage points, so the reported accuracy is not an artefact of emission-account provenance. Third, conformal coverage falls eight points short of nominal under grouped sampling; group-conditional conformal methods and non-exchangeability-robust weightings [35,48] are the natural remedies. Fourth, PSO was run with a modest budget (30 evaluations) on a single fold's inner split; larger budgets or multi-fold fitness could improve the hybrid further. Fifth, the ensemble composition was selected after inspecting out-of-fold results, which is one reason we do not claim significance for its margin over ridge. Finally, our Experiment B conclusions concern annual national aggregates over one- to five-year horizons and do not preclude learnable structure at finer temporal or spatial resolutions.

6. Conclusions

This study reframed national CO_2 modelling around the question inventories actually pose, estimating emissions for countries without usable emission records, and benchmarked thirteen models under strictly country-disjoint validation with country-clustered inference throughout. A physics-guided hybrid, KG-CBA, tuned by particle swarm optimization, significantly improves the identical unguided architecture (median APE 15.7-16.7% vs 20.8%), and a Kaya-motivated ridge regression significantly outperforms every tree ensemble and every unguided deep network. The best ensemble attains the study's lowest point error (MAPE 23.5%, MdAPE 15.1% across 112 countries; MdAPE 7.35% with enriched covariates on 76 countries) but its advantage over ridge alone is not statistically significant once repeated years within countries are accounted for — a distinction that observation-level testing would have concealed.

Error analysis identified energy-mix data availability as the binding constraint, quantified the cost of its absence, and located it geographically in the countries least able to report. Where national energy balances exist, leave-country-out estimation reaches 3-6% median error for

~~African economies such as South Africa and Egypt; where they do not, panels can be assembled only by extending sparse observations across decades, which is itself the clearest statement of the problem.~~

~~Equally important are the study's negative results: unguided deep architectures do not beat a Kaya-motivated linear model in this regime; ensembling does not significantly beat that linear model either; split-conformal intervals under-cover under grouped sampling; SHAP does not rank the decisive covariate first; and in conventional temporal forecasting, learners beat naive persistence only by small margins that fail to replicate across evaluation windows. Future work will extend the framework to sectoral and subnational estimation, group-aware conformal calibration, and transfer to the full set of reporting-gap countries as covariate coverage improves.~~

~~Funding~~

~~This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.~~

~~Ethics statement~~

~~This study used only publicly available, country-level aggregate statistics. It involved no human participants, no animal subjects, and no personal or identifiable data, so no ethics approval was required.~~

~~Acknowledgements~~

where the term in parentheses is the Kaya head and $r_{\boldsymbol{\theta}}$ is the residual branch, a deep network with parameters $\boldsymbol{\theta}$; $\tilde{\mathbf{x}}_W$ is the standardized covariate vector of the most recent year of the window, and $\boldsymbol{\beta}_K$ and $\beta_{K,0}$ are the weights and intercept of the head. The weights of the head are

started at the ridge solution for the most recent year, and the output weights of the residual branch are started close to zero. Training therefore begins from the ridge estimate, and the deep branch can only add what improves on it; the head and the branch are then trained together.

Fig 3. Architecture of the Kaya-guided hybrid network (KG-CBA). The standardized window of eight years and F covariates feeds two paths. The Kaya linear head acts on the most recent year and is started at the ridge solution. The deep residual branch (two one-dimensional convolutional layers, a bidirectional long short-term memory layer, attention pooling and a dense layer) models what the linear head does not explain. The two outputs are added to give the estimate of the logarithm of per-capita CO_2 emissions.

Six settings of KG-CBA were tuned by particle swarm optimization [16] with six particles and five iterations, that is, 30 evaluations (Table 2; S1 Text; S4 Fig); the budget was kept small because each evaluation trains a network. The fitness was the mean squared error on the inner validation countries of the first training fold. The selected configuration was used in all five folds, giving the model PSO-KG-CBA (E-PSO-KG-CBA in Experiment C). The untuned KG-CBA is reported next to it, so that the effect of tuning can be read directly from the results.

The ensemble averages the estimates of ridge, GRU and PSO-KG-CBA (ridge and E-PSO-KG-CBA in Experiment C). Its members were chosen after the out-of-fold results had been inspected. It is therefore an exploratory analysis whose accuracy is likely to be optimistic, and it was not a candidate for the reference estimator.

Model comparison and selection

Accuracy was measured on the original scale for the held-out samples. The absolute percentage error (APE) of a sample is $100\,|c - \hat{c}|/c$. We report its median (MdAPE) and mean (MAPE), the mean absolute error (MAE), the root-mean-square error (RMSE) and the coefficient of

determination (R^2); the formulas are given in S1 Text. Per-capita emissions are strongly right-skewed (0.03 to 77 t per person among the targets), so RMSE and R^2 are dominated by a few high emitters, whereas MdAPE gives every country-year the same weight in relative terms. MdAPE is therefore the primary measure.

Each country contributes 27 consecutive, strongly correlated annual errors, so the 3,024 held-out errors are not independent observations. The validation folds respect this structure, because all years of a country fall in the same fold, and the statistical comparison of two models must respect it as well. For each pair of models we report (i) a Wilcoxon signed-rank test [43] on the 3,024 paired APE values, which treats country-years as independent and is shown only to make the consequence of ignoring the clustering visible (the observation-level test); (ii) the same test on the 112 paired per-country medians of APE (the per-country test); and (iii) a cluster bootstrap [44] in which countries, with all their years, are resampled with replacement 10,000 times, giving a 95% percentile interval and a two-sided p-value for the median of the paired differences in APE. Conclusions are drawn from (iii), with (ii) as a check.

Models were ranked by MdAPE. The reference estimator was taken to be the simplest covariate-based model that no other model was shown to outperform significantly in the cluster bootstrap ($p < 0.05$).

Sensitivity analyses

Three sets of analyses test the robustness of the results. They were run for the ridge reference model and, where stated, for the five classical learners, all of which were re-estimated for this purpose; the neural networks were not re-trained.

Missing data. (i) The blank coal and gas shares were set to zero instead of being filled, as the accounting identity described under Data implies. (ii) The analysis was repeated without the 25

892 countries that have a filled run of more than three years (87 countries remain) and without the 19
893 countries that have a run of more than ten years (93 remain). (iii) The reference model was
894 scored separately on countries with and without long filled runs. Multiple imputation for panel
895 data [45] was not used, because the missing entries are not unknown values.
896 *Atypical economies.* Seven economies whose energy systems the covariates describe poorly were
897 identified from the error analysis (Qatar, Kuwait, Saudi Arabia, Singapore, Luxembourg,
898 Norway and Iceland). Accuracy was recomputed without them, both by scoring the fitted models
899 on the remaining 105 countries and by re-estimating the models on those countries.
900 *Covariate set.* Ridge regression was re-estimated without the three fuel shares, with energy per
901 capita alone, and with the covariates of the most recent year only.

902 ## Prediction intervals

903 Prediction intervals were constructed for the PSO-tuned hybrid of Experiment C (E-PSO-KG-
904 CBA) in three steps (formulas in S1 Text). First, dropout was kept active at prediction time
905 (Monte-Carlo dropout [46]), and 60 repeated predictions of the same sample gave a predictive
906 mean and standard deviation of y. Second, the intervals were calibrated by split-conformal
907 prediction [47,48]: on the inner validation countries of each fold, the absolute error of the
908 predictive mean was divided by the predictive standard deviation, and the calibration factor $\hat{q}$
909 was set to the 90th percentile of these ratios, with the usual finite-sample correction. Third, the
910 90% interval for a test sample was the predictive mean plus and minus $\hat{q}$ standard deviations,
911 transformed back to t CO_2 per person. Intervals were assessed by their coverage, the percentage
912 of held-out observations inside their interval, which should equal the nominal 90%, and by their
913 mean width.

Conformal intervals attain their nominal coverage when calibration and test observations are exchangeable. Here they come from different countries, so that exchangeability holds, at best, between countries and not between country-years [49–51]. To separate this effect from the properties of one model, we repeated the calibration for ridge regression in Experiment C with calibration sets of 8 training countries (the size used for the hybrid), of all 60 or 61 training countries by grouped cross-conformal prediction and, as a reference in which exchangeability holds, of samples drawn from the held-out countries themselves (S1 Text). The same analysis for Experiment A, with 12 calibration countries, is reported in the Results.

Explainability

Three measures of the importance of each covariate were used. The first is SHAP (Shapley additive explanations) [52], which attributes each estimate of a fitted model to its inputs: the SHAP value of an input is its average contribution to the estimate over all subsets of the other inputs, and the mean of its absolute value over samples measures the overall importance of the input (formula and symbols in S1 Text). SHAP values were computed exactly for the LightGBM model of Experiment C with the tree algorithm [53] and aggregated by covariate and by year of the window.

SHAP describes how a fitted model uses its inputs. It does not measure how much an input contributes to accuracy for unseen countries, and it can mislead when inputs are correlated [54]. Two further measures were therefore defined directly on held-out error and computed for ridge regression and LightGBM on the 76 countries of Experiment C. For permutation importance [37,55], the eight-year series of one covariate was shuffled between the test samples of each fold (30 repetitions) and the increase in MdAPE recorded. For drop-one-covariate importance, the leave-one-covariate-out analysis of Lei et al. [24], the model was re-estimated without the

covariate under the same folds. For the hybrid, the attention weights show how much each year of the window contributes to the pooled representation.

Software, data and code

The benchmark was run in Python 3.12.3 with TensorFlow 2.21.0 [56], scikit-learn 1.8.0 [57], XGBoost 3.4.0, LightGBM 4.7.0 and SHAP 0.52.0, with random seed 42 and fold-specific offsets. The analyses added in revision were run in Python 3.13 with scikit-learn 1.9.1, XGBoost 3.4.1, LightGBM 4.7.0 and SHAP 0.54.1; they reproduce the original results for the mean predictor, ridge regression, k-nearest neighbours and support vector regression to three decimal places, whereas the three tree ensembles were re-estimated with a different random state and differ slightly (S2 Table). The harmonized panel, the code for every model, the out-of-fold estimates of every model and the scripts that produce each table and figure are available at https://github.com/samfozzy/kg-cba-pclm-revision (archived release DOI: https://doi.org/10.5281/zenodo.24195216). The source data are available from https://github.com/owid/co2-data and https://github.com/owid/energy-data. The use of a generative artificial intelligence tool in this work is described in the Acknowledgments.

Results

Data overview

Median per-capita emissions were 5.0 t CO_2 per person in 1990 and 4.0 t in 2023, and the spread between countries remained wide: in 2023 the 10th and 90th percentiles were 0.6 and 11.5 t per person (Fig 2a). Across all country-years, the logarithm of per-capita emissions correlates strongly with the logarithm of energy per capita (Pearson r = 0.97) and of GDP per capita (r = 0.88), and only weakly with the fuel shares (S2 Fig), as the Kaya identity leads one to expect.

Experiment A: accuracy of the thirteen models

Table 3 and Fig 4 give the accuracy of every model on held-out countries (S1 Fig shows estimated against observed values). Covariates are clearly informative: the mean predictor has an MdAPE of 63.3%, and every covariate-based model except k-nearest neighbours (39.0%) is below 22%.

Ridge regression has an MdAPE of 16.3%. It has the lowest MAE and RMSE and the highest R^2 of all models, and a lower MdAPE than every classical machine-learning model and every unguided deep network (18.2% to 39.0%).

Kaya guidance changes the deep network substantially. The unguided CNN-BiLSTM-Attn network has an MdAPE of 20.8%; with the linear head, the same architecture reaches 15.7% (KG-CBA) and 16.7% (PSO-KG-CBA). The hybrids nevertheless have a higher RMSE (4.6 to 4.7 t per person) and a lower R^2 (0.52 to 0.55) than any other deep network, and tuning by PSO did not improve on the untuned hybrid. The exploratory ensemble has the lowest MdAPE (15.1%) and MAPE (23.5%).

Table 3. Accuracy on held-out countries in Experiment A (112 countries, six covariates, 3,024 samples).

Model	Family	RMSE	MAE	MAPE (%)	MdAPE (%)	R^2
Mean predictor	Naive baseline	7.415	4.408	337.8	63.3	−0.178
Ridge regression	Linear	**3.056**	**1.443**	25.2	16.3	**0.800**
kNN	Classical machine learning	5.119	2.729	49.7	39.0	0.439
SVR	Classical machine learning	4.881	2.100	32.3	21.3	0.490
Random forest	Classical machine learning	4.233	1.824	28.7	19.1	0.616
XGBoost	Classical machine learning	4.043	1.775	28.0	18.4	0.650
LightGBM	Classical machine learning	3.943	1.813	28.4	18.6	0.667
LSTM	Deep network	3.815	1.703	27.6	18.2	0.688

Model	Family	RMSE	MAE	MAPE (%)	MdAPE (%)	R^2
GRU	Deep network	3.629	1.644	25.1	18.5	0.718
CNN-LSTM	Deep network	3.773	1.707	28.1	20.8	0.695
CNN-BiLSTM-Attn	Deep network	3.812	1.790	28.6	20.8	0.689
KG-CBA	Kaya-guided hybrid	4.595	1.898	26.0	**15.7**	0.548
PSO-KG-CBA	Kaya-guided hybrid	4.748	1.914	27.2	16.7	0.517
Ensemble (ridge, GRU, PSO-KG-CBA)	Exploratory	3.557	1.530	23.5	15.1	0.729

RMSE and MAE are in t CO_2 per person. Bold marks the best value among the thirteen models; the ensemble was assembled after the results had been inspected and is shown for exploration only. Abbreviations as in Table 2.

Fig 4. Accuracy of the models on held-out countries in Experiment A. (a) Median absolute percentage error. (b) Root-mean-square error. The dashed line marks ridge regression. Arrows indicate values beyond the axis range. Values are those of Table 3.

Country-clustered comparison of models

Table 4 gives the statistical comparisons. The effect of Kaya guidance is significant: KG-CBA and PSO-KG-CBA are more accurate than the unguided network of the same architecture (cluster-bootstrap $p = 0.002$ for each). Ridge regression is significantly more accurate than LightGBM ($p = 0.006$) and than the unguided CNN-BiLSTM-Attn network ($p = 0.002$). With the classical learners re-estimated for this revision, it is also significantly more accurate than XGBoost ($p = 0.004$), random forest ($p = 0.011$), support vector regression and k-nearest neighbours (both $p < 0.001$; S2 Table).

The ensemble is not significantly more accurate than ridge regression: the median paired difference in APE is −0.76 percentage points, with a 95% interval from −2.20 to +0.41 (cluster-bootstrap $p = 0.22$; per-country test $p = 0.098$). The observation-level test, which treats the 3,024 country-years as independent, gives $p = 7 \times 10^{-10}$ for the same comparison, eight orders of

magnitude smaller. The ensemble is significantly more accurate than its two neural members, GRU ($p < 0.001$) and PSO-KG-CBA ($p = 0.022$).

KG-CBA has a slightly lower MdAPE than ridge regression (15.7% against 16.3%) and PSO-KG-CBA a slightly higher one (16.7%), while both have a markedly higher MAE and RMSE and a lower R^2 (Table 3). The cluster-bootstrap tests confirm that neither hybrid differs significantly from ridge regression: the median paired difference in APE is +0.48 points for KG-CBA (95% interval −0.95 to +1.98; $p = 0.50$) and +0.47 for PSO-KG-CBA (−1.05 to +2.09; $p = 0.51$). LSTM and GRU are not significantly less accurate than ridge regression either ($p = 0.08$ and 0.12), whereas the two convolutional networks are ($p = 0.001$ and 0.002). No model was thus shown to be more accurate than ridge regression, the simplest covariate-based model, and ridge regression is retained as the reference estimator.

Table 4. Paired comparisons of models with countries as the unit of inference.

Exp.	Comparison (first against second)	MdAPE (%), first / second	Median paired difference in APE, points [95% CI]	p, cluster bootstrap	p, per-country test	p, observation-level test
A	Ensemble against ridge	15.1 / 16.3	−0.76 [−2.20, +0.41]	0.22	0.098	6.9×10^{-10}
A	Ensemble against GRU	15.1 / 18.5	−2.57 [−3.77, −1.20]	0.0006	0.007	1.8×10^{-30}
A	Ensemble against PSO-KG-CBA	15.1 / 16.7	−1.33 [−2.85, −0.22]	0.022	0.074	3.6×10^{-31}
A	PSO-KG-CBA against CNN-BiLSTM-Attn	16.7 / 20.8	−3.51 [−5.97, −0.97]	0.0016	0.004	6.3×10^{-22}
A	KG-CBA against CNN-BiLSTM-Attn	15.7 / 20.8	−3.42 [−5.84, −1.21]	0.0018	0.004	4.9×10^{-26}
A	Ridge against LightGBM	16.3 / 18.6	−3.10 [−4.91, −1.03]	0.0058	0.026	8.3×10^{-25}
A	Ridge against CNN-BiLSTM-Attn	16.3 / 20.8	−4.29 [−7.05, −1.64]	0.0024	0.007	3.1×10^{-34}
A	KG-CBA against ridge	15.7 / 16.3	+0.48 [−0.95, +1.98]	0.50	0.58	1×10^{-4}
A	PSO-KG-CBA against ridge	16.7 / 16.3	+0.47 [−1.05, +2.09]	0.51	0.98	1×10^{-4}
A	LSTM against ridge	18.2 / 16.3	+1.55 [−0.18, +3.58]	0.079	0.17	$< 10^{-4}$
A	GRU against ridge	18.5 / 16.3	+1.69 [−0.44, +3.86]	0.12	0.18	$< 10^{-4}$

Exp.	Comparison (first against second)	MdAPE (%), first / second	Median paired difference in APE, points [95% CI]	p, cluster bootstrap	p, per-country test	p, observation-level test
A	CNN-LSTM against ridge	20.8 / 16.3	+4.31 [+1.75, +6.81]	0.0012	0.006	$< 10^{-4}$
C	E-PSO-KG-CBA against ridge	8.09 / 7.72	+0.66 [−0.23, +1.57]	0.16	0.21	$< 10^{-4}$
C	Ensemble against ridge	7.35 / 7.72	−0.20 [−0.66, +0.36]	0.56	0.49	0.48
C	E-PSO-KG-CBA against LightGBM	8.09 / 13.10	−3.85 [−6.03, −1.95]	0.0001	0.0002	1.7×10^{-56}
C	Ensemble against E-PSO-KG-CBA	7.35 / 8.09	−0.86 [−1.45, −0.35]	0.0014	0.004	5.1×10^{-31}
C	Ridge with seven against ridge with six covariates	7.71 / 15.64	−7.07 [−9.61, −4.59]	< 0.0001	2.0×10^{-7}	6.9×10^{-121}

Exp., experiment. A negative difference means that the first model has the smaller error. The interval is the 95% percentile interval of a cluster bootstrap that resamples countries (10,000 draws). The per-country test is a Wilcoxon signed-rank test on per-country medians of APE (112 countries in Experiment A, 76 in Experiment C). The observation-level test is the same test on all country-years and is shown only to illustrate the effect of ignoring the clustering; conclusions follow the cluster bootstrap. The last row, and the rows against ridge regression in Experiment A, were added in revision and use identical countries and folds for both models; they were computed by a separate script with its own bootstrap draws, so their p-values can differ from the other rows by about 0.001. CI, confidence interval.

Error concentration in atypical energy systems

The squared error is concentrated in a few countries (Fig 5a). For ridge regression, the seven atypical economies account for 77.9% of the total squared error. For the exploratory ensemble and the PSO-tuned hybrid, the seven atypical economies account for 84.3% and 86.5% of the squared error (S3 Fig shows the ensemble). Qatar, Kuwait and Singapore are among the six worst-fitted countries under each of the six models examined; Iceland, Norway, Luxembourg and Saudi Arabia enter or leave the list depending on the model. Ridge regression overestimates the

emissions of Iceland and Norway, whose energy is largely hydroelectric and geothermal (mean error +13.4 and +9.8 t per person), and of Singapore (+13.7 t per person), a refining and bunkering centre in which a large part of energy use is plausibly not emitted on its territory. It underestimates the emissions of the hydrocarbon exporters Qatar, Kuwait and Saudi Arabia (−4.6, −9.1 and −5.3 t per person) and of Luxembourg (−7.1 t per person).

Leaving these seven economies out changes the scale-dependent measures and little else (Table 5; Fig 5b). For ridge regression the RMSE falls from 3.06 to 1.48 t per person and R^2 rises from 0.80 to 0.90, while the MdAPE changes only from 16.3% to 15.1% (14.9% when the model is re-estimated without them). The five classical learners improve in the same way, and ridge regression remains the most accurate of the six on MdAPE (S2 Table). Without the seven atypical economies, the RMSE is 1.45 t per person for the ensemble, 1.75 for KG-CBA and 1.80 for PSO-KG-CBA, against 1.48 for ridge regression, and the R^2 of the hybrids rises to 0.86 and 0.85 (ridge regression 0.90). The low R^2 of the hybrids in Table 3 is therefore produced mainly by a handful of countries, but removing them does not make the hybrids more accurate than ridge regression: their differences from ridge regression remain not significant (cluster-bootstrap $p = 0.87$ for KG-CBA, 0.89 for PSO-KG-CBA and 0.10 for the ensemble; S2 Table).

Fig 5. Concentration of the error of ridge regression and effect of excluding the atypical economies. (a) Share of the total squared error of ridge regression contributed by each of the twelve countries with the largest contributions. Orange bars are the seven atypical economies. Countries are identified by ISO 3166-1 alpha-3 codes: SGP, Singapore; ISL, Iceland; QAT, Qatar; NOR, Norway; KWT, Kuwait; LUX, Luxembourg; SWE, Sweden; SAU, Saudi Arabia; KAZ, Kazakhstan; MLT, Malta; ARE, United Arab Emirates; FRA, France. (b) Root-mean-square error of ridge regression and the five classical learners for all 112 countries and with the

seven atypical economies excluded from scoring. The three tree ensembles were re-estimated for this analysis, and their values for all countries differ slightly from Table 3.

Table 5. Sensitivity of the ridge reference model to atypical economies, missing-data treatment and covariate set.

Analysis	N	RMSE	MAE	MAPE (%)	MdAPE (%)	R^2	Difference from reference, points [95% CI]; p
Reference: original panel, six covariates	112	3.06	1.44	25.2	16.3	0.800	–
Atypical economies							
Seven atypical economies excluded from scoring	105	1.48	0.90	22.1	15.1	0.897	–
Seven atypical economies excluded, model re-estimated	105	1.61	0.94	22.6	14.9	0.878	–
The seven atypical economies alone	7	10.79	9.59	71.9	40.3	0.428	–
Missing data							
Blank coal and gas shares set to zero	112	3.06	1.44	25.3	16.2	0.800	+0.03 [−0.02, +0.07]; 0.19
Countries with a filled run of more than 3 years excluded, model re-estimated	87	3.33	1.74	27.5	17.6	0.778	–
Countries with a filled run of more than 10 years excluded, model re-estimated	93	3.33	1.61	25.4	15.6	0.769	–
Reference model scored on the 87 countries without a long filled run	87	3.41	1.67	26.5	15.5	0.767	–
Reference model scored on the 25 countries with a long filled run	25	1.20	0.64	20.9	18.3	0.941	–
Covariate set							
Without the three fuel shares	112	3.24	1.57	26.7	16.9	0.775	+1.03 [−0.62, +3.06]; 0.22
Energy per capita only	112	3.60	1.69	28.5	17.9	0.723	+2.37 [+0.59, +4.17]; 0.009
Covariates of the most recent year only	112	3.34	1.52	25.4	15.7	0.761	+0.18 [−0.32, +0.72]; 0.41

All rows are leave-country-out results for ridge regression. N is the number of countries scored. RMSE and MAE are in t CO_2 per person. The last column gives the median paired difference in APE from the reference model with its 95% cluster-bootstrap interval and p-value, for analyses that use the same 112 countries and folds. A long filled run is a run of more than three consecutive filled years. Results for the five classical learners are given in S2 Table.

Experiment C: value of the low-carbon share

On the 76 countries for which the low-carbon share of primary energy is available, all four models are far more accurate than in Experiment A (Table 6). Ridge regression with seven covariates has an MdAPE of 7.7%, the hybrid 8.1%, the ensemble 7.35% and LightGBM 13.1%; ridge regression again has the lowest MAE, RMSE and MAPE and the highest R^2, and the ensemble is again not significantly more accurate than ridge regression ($p = 0.56$; Table 4).

The gain is attributable to the covariate. With identical countries and folds, ridge regression has an MdAPE of 15.6% with six covariates and 7.7% with seven, a median paired difference of −7.07 percentage points (95% interval −9.61 to −4.59; $p < 0.001$; Table 4). Fig 6 shows the corresponding comparison for ridge regression, the hybrid and the ensemble when the six-covariate estimates are taken from the models of Experiment A; the reduction is of similar size for all three.

The gain is not confined to the atypical economies. Among the 69 other countries of this panel, the MdAPE of ridge regression falls from 14.1% to 7.0% ($p < 0.001$). Among the atypical economies it falls sharply for the two hydroelectric and geothermal systems, from 157% to 29% for Iceland and from 107% to 28% for Norway, and moderately for Luxembourg, Saudi Arabia, Kuwait and Qatar. Singapore is the exception: its error increases from 125% to 231%. Its low-carbon share is close to zero, so the new covariate confirms a fossil-based energy system but cannot represent energy use that is not emitted on its territory. In total, the squared error falls by 39%, and Singapore then accounts for 49% of what remains.

Table 6. Accuracy on held-out countries in Experiment C (76 countries, 2,052 samples).

Model	Covariates	RMSE	MAE	MAPE (%)	MdAPE (%)	R^2
LightGBM	Seven	4.520	2.048	22.8	13.1	0.622
E-PSO-KG-CBA	Seven	3.682	1.593	16.0	8.1	0.749
Ridge regression	Seven	**3.037**	**1.284**	**14.1**	7.7	**0.829**

Model	Covariates	RMSE	MAE	MAPE (%)	MdAPE (%)	R^2
Ensemble (ridge, E-PSO-KG-CBA)	Seven	3.306	1.409	14.5	7.35	0.798
Ridge regression, same countries and folds	Six	3.877	2.003	22.5	15.6	0.722

RMSE and MAE are in t CO_2 per person. Bold marks the best value among the three individual models with seven covariates; the ensemble is exploratory. The last row was added in revision and differs from the six-covariate values shown in Fig 6, which come from the models of Experiment A (trained on 112 countries) scored on the same 76 countries (ridge regression: MAPE 22.2%, MdAPE 14.3%).

Fig 6. Effect of adding the low-carbon share of primary energy. (a) Mean absolute percentage error (MAPE) and (b) median absolute percentage error (MdAPE) on the same 76 countries for ridge regression, the particle-swarm-tuned hybrid (PSO-KG-CBA) and the exploratory ensemble. Grey bars: six covariates, estimates from the models of Experiment A. Green bars: seven covariates (Experiment C).

Robustness to missing-data treatment and covariate set

The treatment of the blank fuel shares does not affect the results (Table 5). Setting them to zero instead of filling them changes the MdAPE of ridge regression from 16.3% to 16.2% ($p = 0.19$) and that of each of the five classical learners by at most 0.3 percentage points (S2 Table). Excluding the countries with long filled runs and re-estimating the model gives an MdAPE of 17.6% when 25 countries are excluded and 15.6% when 19 are excluded. Scored separately, the 87 countries without a long filled run have an MdAPE of 15.5% and the 25 countries with one have 18.3%, while on MAPE the order is reversed (26.5% and 20.9%). Countries with long filled runs are thus neither consistently easier nor consistently harder to estimate.

The fuel shares contribute little. Without them, the MdAPE of ridge regression is 16.9%, not significantly different from the reference model ($p = 0.22$); with energy per capita alone it is

17.9% (p = 0.009). The eight-year window also contributes little to the median error: with the covariates of the most recent year only, the MdAPE is 15.7% (p = 0.41), although the RMSE is higher (3.34 against 3.06 t per person). The result does not depend on the ridge penalty: for λ between 0.01 and 100 the MdAPE lies between 16.0% and 16.4%.

Regional results and African countries

For ridge regression, the median APE by continent ranges from 13.6% in South America to 20.6% in North America, with Africa at 16.9% (S2 Table). By country, the median of the 112 per-country MdAPE values is 15.5% (interquartile range 9.6% to 23.8%); 70 countries are below 20% and 11 are above 50% (S1 Table).

Of the 17 African countries in the panel, eight have a per-country MdAPE below 15% (Cameroon, South Africa, Botswana, Nigeria, Tanzania, Senegal, Ghana and Egypt; 7.1% to 13.5%), and three are above 40% (Niger 44%, Mozambique 74% and the Democratic Republic of the Congo 149%). These three have very low emissions per person, so that small absolute errors are large in relative terms, and none of them is in the panel of Experiment C: of the 17 African countries, only Algeria, Egypt, Morocco and South Africa have the low-carbon share in the source used here. Fig 7 shows the observed emissions of South Africa and Egypt with the estimates of the Experiment C ensemble and 90% prediction intervals of the hybrid. The per-country MdAPE is 6.1% for South Africa and 3.4% for Egypt for the ensemble, 5.6% and 5.7% for the hybrid, and 5.5% and 4.9% for ridge regression.

Fig 7. Observed and estimated per-capita CO_2 emissions of South Africa (ZAF) and Egypt (EGY), with 90% prediction intervals. Estimates are those of the Experiment C ensemble (mean of ridge regression and E-PSO-KG-CBA), fitted without any data from the country

shown, and the 90% split-conformal intervals are those of E-PSO-KG-CBA. Panel titles give the median absolute percentage error (MdAPE) for the country.

Prediction intervals

Uncalibrated Monte-Carlo dropout intervals of the hybrid of Experiment C are far too narrow: at a nominal level of 90% they cover 13.2% of the held-out observations, with a mean width of 0.47 t per person. After split-conformal calibration the coverage is 81.9% (81.0% to 83.0% across repeated Monte-Carlo runs) and the mean width 5.68 t per person; the calibration factor $\hat{q}$ varies between folds from 8.8 to 69.2.

The analysis for ridge regression shows where the remaining shortfall comes from (S2 Table). When calibration and evaluation samples come from the same held-out countries, coverage is 90.2%, as the theory requires. With calibration on 12 training countries it averages 87.4% (80.3% to 94.5% across random draws of countries) and ranges from 45% to 100% in single folds; with calibration on all 89 or 90 training countries it is 88.9% (73.8% to 96.5% by fold). Coverage is also uneven between countries: 68% of the countries have every year inside the interval, and 8% have half of their years or fewer inside it. The comparison for Experiment C, which is the setting of the hybrid's intervals, gives the following. With the 8 calibration countries used for the hybrid, ridge regression reaches 86.5% on average over 200 random draws of countries (77.2% to 96.5%), with a mean width of 5.14 t per person and coverage by fold from 52% to 100%. Calibration on the 60 or 61 training countries by grouped cross-conformal prediction gives 88.8% with a mean width of 3.76; 61% of the countries have every year inside the interval and 9% have half or fewer, against 57% and 17% for the stored intervals of the hybrid. The narrower intervals of ridge regression reflect its smaller errors in this experiment. Calibration on a random half of the held-out observations themselves gives 90.3%.

Covariate importance

SHAP values for LightGBM in Experiment C are dominated by energy per capita (mean absolute value 0.114), followed by the low-carbon share (0.016), the gas share (0.008) and population (0.007) (Fig 8a). By year of the window, the most recent year carries most of the attribution (0.118), the year before it much less (0.027), and each earlier year less than 0.008 (Fig 8b). The attention weights of the hybrid, in contrast, are close to uniform (0.095 to 0.142 against a uniform value of 0.125; S6 Fig).

The two measures based on held-out error give the same ranking for the first two covariates (Table 7). Removing energy per capita and re-estimating the model raises the MdAPE of ridge regression from 7.7% to 29.4% and that of LightGBM from 13.2% to 32.2%; removing the low-carbon share raises it by 7.9 and 3.2 percentage points; removing any other covariate changes it by about one percentage point or less. Permutation importance agrees for LightGBM. For ridge regression it assigns about 30 percentage points to each of the three fuel shares, although removing any one of them costs at most 1.1 points: the shares are strongly related to one another, so that shuffling one of them produces combinations that do not occur, from which the linear model extrapolates.

Fig 8. SHAP importance in the LightGBM model of Experiment C. Mean absolute SHAP value (a) by covariate and (b) by year of the eight-year window, where t-0 is the target year. log_epc, logarithm of energy per capita; log_gpc, logarithm of GDP per capita.

Table 7. Importance of each covariate in Experiment C by three measures (76 countries).

Covariate	SHAP rank, LightGBM	Permutation, ridge	Permutation, LightGBM	Drop-one, ridge	Drop-one, LightGBM
Energy per capita	1	+54.7	+43.6	+21.7	+19.0
Low-carbon share	2	+10.6	+5.8	+7.9	+3.2
Gas share	3	+31.8	+0.5	+1.1	−0.6

Covariate	SHAP rank, LightGBM	Permutation, ridge	Permutation, LightGBM	Drop-one, ridge	Drop-one, LightGBM
Population	4	0.0	−0.4	−0.1	−0.8
Oil share	5	+31.4	+0.2	+0.3	+0.1
Coal share	6	+30.0	+1.6	+0.4	+0.2
GDP per capita	7	+1.0	−0.4	−0.3	−0.7

Permutation and drop-one columns give the change in MdAPE on held-out countries, in percentage points, when the eight-year series of the covariate is shuffled between test samples (mean of 30 repetitions) or when the model is re-estimated without the covariate. The reference MdAPE is 7.7% for ridge regression and 13.2% for LightGBM; LightGBM was re-estimated for this analysis and differs slightly from Table 6 (13.1%). SHAP ranks are those of Fig 8.

Supporting experiment: temporal forecasting

When a country's own emission history is available, naive persistence is difficult to beat (S2 Text). For forecasts of 2021 to 2023 made one year ahead, only one learner improved significantly on persistence, by 7% of the mean absolute error; three and five years ahead the largest improvements were 9% and 19%. In a replication for 2015 to 2019, no learner improved significantly on persistence one or three years ahead and only one did so five years ahead, and the learner that performed best in one period was generally not the best in the other.

Discussion

Principal findings

The four questions have clear answers. (Q1) From six widely available indicators, per-capita fossil CO_2 emissions of a country that is absent from training can be estimated with a median error of about 16%, and of about 8% where the low-carbon share of primary energy is also known. (Q2) Deep and hybrid networks do not outperform a linear model motivated by the Kaya

identity: tree ensembles and the unguided convolutional networks (CNN-LSTM and CNN-BiLSTM-Attn) are significantly less accurate than ridge regression, and neither the Kaya-guided network nor an ensemble was shown to be more accurate than ridge regression alone. (Q3) Better covariate data reduce the error far more than better models: no change of model lowered the median error by more than 1.2 percentage points relative to ridge regression, whereas one additional covariate halved it. (Q4) These results do not depend on how missing fuel shares are treated, on whether the atypical economies are included, or on the ridge penalty.

Why deep learning did not outperform the linear model

Five features of the task, each visible in the results, explain why additional model capacity did not help.

The relationship is close to linear by construction. By the Kaya identity, the logarithm of per-capita emissions is the sum of the logarithm of energy per capita and the logarithm of carbon intensity (Eq 5). Energy per capita alone correlates at r = 0.97 with the target and gives an MdAPE of 17.9% in a one-covariate ridge model. Little non-linear signal remains to be found.

The number of independent units is the number of countries. A model that must generalize to new countries learns from about 90 training countries, not from 2,400 training samples, because the years of one country are close to copies of one another: the covariates of the most recent year alone give the same median error as the eight-year window. Flexible models with thousands of parameters cannot be estimated reliably from 90 effectively independent units; they fit country-specific patterns that do not transfer. This agrees with evidence that tree ensembles and simple linear models remain competitive with deep networks on small tabular and time-series problems [20,58–60].

Held-out countries can lie outside the range of the training countries. The economies with the largest errors are the most extreme in energy use or emissions per person. Tree ensembles cannot produce estimates beyond the range of the training targets, and neural networks behave unpredictably outside their training range, whereas a linear model in logarithms extrapolates along the identity. The residual branch of the hybrids lowers the median error slightly but adds large errors for a few extreme economies, which is why their RMSE is the highest among the deep models.

The remaining error is caused by missing information, not by the functional form. What the linear model does not explain is the part of carbon intensity that the six covariates do not describe, such as non-fossil energy. No model can recover this from covariates that do not contain it; when one covariate that contains part of it was added, the error of ridge regression, the hybrid and the ensemble fell by a similar large amount (Fig 6).

Tuning added variance, not accuracy. The hybrid tuned by particle swarm optimization was no more accurate than the hybrid with fixed settings. A fitness value computed on 12 validation countries is noisy, and selecting the best of 30 configurations on such a value mostly selects noise; a larger search budget would not remove this limitation, which comes from the number of countries.

We therefore found no advantage of the complex models in accuracy, and they are less transparent and more costly to train. Their value in this study is methodological. They represent current practice and had to be tested under a valid protocol before they could be set aside, and the comparison between the guided and the unguided network shows that the deficit of deep networks in this task is a failure to learn the linear structure from few countries, not a lack of

capacity; once this structure is supplied, nothing remains for the network to add. For inventory gap-filling we recommend ridge regression on logarithmic covariates.

The testing regime matters as much as the model. Treating country-years as independent would have declared the ensemble better than ridge regression with p below 10^{-9}, whereas the clustered test gives p = 0.22; studies that test at the level of the observation will regularly report small architectural gains as highly significant when they are not.

What limits accuracy: covariate data

Experiment C shows that the accuracy of the estimates is limited by the covariates. The low-carbon share of primary energy measures directly the part of carbon intensity that the fossil-fuel shares cannot represent, and after energy per capita it is the second most important covariate by SHAP, by drop-one importance for both models and by permutation importance for LightGBM (Table 7). The measures answer different questions and are consistent with one another. SHAP describes how much a fitted model relies on a covariate; energy per capita dominates because it carries the scale of emissions. Experiment C and the drop-one analysis measure what is lost in accuracy for unseen countries when a covariate is unavailable. Repeating that experiment for the covariate that SHAP ranks first shows the largest loss (the MdAPE of ridge regression rises from 7.7% to 29.4% without energy per capita), but energy per capita is available for every country. The low-carbon share is the most valuable covariate that many countries lack.

That lack is concentrated where estimates are most needed: in the source used here, the low-carbon share is available for 76 of the 112 countries of the panel and for only 4 of its 17 African countries, and the three African countries that are estimated worst are among those without it. By contrast, the long filled runs in the fuel-share series are not a sign of thin statistics: they are zeros for fuels that a country did not use, and their treatment has no effect on the results. The estimator

also does not need the fuel shares, the only inputs derived from emission accounts; without them the median error is 16.9% instead of 16.3%. It can therefore be applied to a country that has energy and economic statistics but no emission inventory.

A roadmap for better national estimates

The results suggest the following priorities, in order of expected benefit.

1. *Extend energy-mix statistics.* Compiling the low-carbon share of primary energy, or the full breakdown of primary energy by source, from national energy balances for the 36 panel countries that lack it, and for countries outside the panel, is the step most likely to improve estimates; where it is available the median error is halved.
2. *Add covariates for the atypical systems.* The largest remaining errors point to variables that could be tested in the same way: international bunker fuel and refinery throughput for a trading hub such as Singapore, and flaring and the energy used in oil and gas processing for hydrocarbon exporters.
3. *Derive fuel shares from energy balances.* This would make the estimator fully independent of emission accounts, at a cost of less than one percentage point of median error.
4. *Keep the linear core and test additions against it.* Hierarchical models that share information between similar countries [61], and guided networks of the kind tested here, should be retained only if they are significantly more accurate than ridge regression under country-clustered inference.
5. *Adopt evaluation standards that match the task.* Studies of national emission modelling should validate on countries that are absent from training, use the country as the unit of

inference, compare temporal forecasts with naive persistence, and report errors by country in addition to pooled measures such as R^2.

6. *Calibrate uncertainty on many countries.* Prediction intervals should be calibrated on as many countries as possible and their coverage reported by country; methods designed for grouped data [51] are the natural next step.

7. *Extend the scope.* The same design can be applied to sectoral and sub-national emissions and to other greenhouse gases, for which inventories are less complete than for fossil CO_2.

Limitations

First, the balanced panel contains 112 countries, and small and low-income countries, for which inventories are weakest, are under-represented; only 17 African countries are included. Second, the covariates are not independent of the statistical systems that produce inventories: primary energy statistics are themselves the main input to inventories, so the estimator is a cross-check and a means of filling gaps, not a substitute for national energy statistics. Third, the target is territorial fossil CO_2; emissions from land use and other gases are not covered. Fourth, the sensitivity analyses were run for ridge regression and the classical learners; the neural networks were not re-trained under the alternative treatments. Fifth, the hybrid was tuned with a small search budget on the first fold, and the selected configuration was used in the other folds, whose test countries had been part of the search data; as tuning did not improve accuracy, this cannot have inflated the results. Sixth, the ensemble is exploratory. Seventh, prediction intervals cover less than their nominal level for countries unlike those used for calibration. Finally, the results rest on one vintage of the source data and on annual national totals.

Conclusions

National per-capita fossil CO_2 emissions can be estimated for a country from its energy and economic indicators, without any of its own emission data, with a median error of about 16%, and of about 8% where the low-carbon share of primary energy is known. A ridge regression on logarithmic covariates, which follows from the Kaya identity, is the estimator to use: tree ensembles and the unguided convolutional networks are significantly less accurate, and neither a Kaya-guided hybrid network nor an ensemble was shown to improve on it once countries, not country-years, are the unit of inference. The error is concentrated in a few economies whose energy systems the indicators describe poorly, and one additional energy-mix indicator halves it. Estimates of national emissions where inventories are missing are therefore improved most by extending energy-mix statistics to more countries, not by increasing model complexity.

Acknowledgments

The authors thank the Global Carbon Project, the Energy Institute, the U.S. Energy Information Administration, the Maddison Project and Our World in Data for maintaining and openly publishing the ~~datasets~~data on which this work depends.

~~CRediT authorship contribution statement~~

~~**S.C. Atuahene:** Conceptualization, Methodology, Software, Formal analysis, Investigation, Data curation, Visualization, Writing - original draft. **B. Boateng:** Supervision, Resources, Writing - review & editing. **H. Amarfio:** Investigation, Validation, Writing - review & editing. **E. Y. Quayson:** Investigation, Validation, Writing - review & editing. **F.B. Agyenim:** Investigation, Validation, Writing - review & editing.~~

Declaration of competing interest

The authors declare no competing financial interests or personal relationships that could have influenced this work.

Data and code availability

All data are openly available: Our World in Data CO_2 and energy datasets (https://github.com/owid/co2-data; https://github.com/owid/energy-data), redistributing the Global Carbon Budget and the Energy Institute Statistical Review of World Energy. The complete analysis pipeline, harmonized panel, per-fold out-of-fold prediction vectors for every model, and the fold-level model checkpoints for the enriched hybrid are openly available at https://github.com/samfozzy/kg-cba-loco-co2-reproducibility. This repository contains the harmonized panel, the complete analysis pipeline (build_panel.py, which regenerates the panel exactly from the raw downloads; classical and deep LOCO baselines, PSO tuning, enriched-panel models, conformal calibration, explainability, Experiment B, and the clustered significance analysis), per-fold out-of-fold prediction vectors for every model, the fold-level model checkpoints for the enriched hybrid, and all scripts that generate the figures and tables in this paper. REPRODUCE.md in that repository documents the run order and expected outputs.

Declaration on the use of generative AI

During the preparation of this work the authors used Claude (Anthropic) to assist with code development for the experimental pipeline, statistical analysis, figure generation, and drafting of the manuscript text. The authors reviewed and edited all content and take full responsibility for the integrity and accuracy of the publication.

ORCID iDs:

~~Samiuel C. Atuahene: https://orcid.org/0000-0002-2928-6458;~~

~~Bismark Boateng: https://orcid.org/0000-0003-3906-5277;~~

~~Hilda Amarfio: https://orcid.org/0009-0006-0806-5919;~~

~~Ekow Yaleh Quayson: https://orcid.org/0009-0006-3516-3434;~~

~~Francis Boateng Agyenim: https://orcid.org/0000-0003-4986-2261;~~

~~References~~

Use of artificial intelligence tools. The authors used the generative AI tool Claude (Anthropic, San Francisco, CA, USA; Claude Fable 5.2 (API: claude-fable-5-2)) in three ways. (i) Code: Claude drafted and revised Python code for data harmonization, model training, statistical tests and figures, following analysis specifications written by the authors. (ii) Analyses added in revision: the scripts for the sensitivity, importance and conformal analyses were drafted and first run with Claude from the public source files. (iii) Text: Claude assisted with drafting and language editing of the manuscript and of the response to the reviewers. No AI tool generated or altered data; all data come from the cited public sources. To evaluate the validity of the outputs, All authors have confirmed. the authors read and executed all code and regenerated every table and figure from the released scripts; confirmed that independently re-implemented deterministic models reproduce the reported values to three decimal places; checked every numerical statement against the script output and every reference against the publisher's record; and reviewed and edited all text. The authors take full responsibility for the content of this article.

References

~~[~~1~~]~~. United Nations Environment Programme. Emissions Gap Report 2025: Off ~~Target:~~target – Continued ~~Collective~~collective ~~Inaction~~inaction ~~Puts~~puts ~~Global~~global ~~Temperature~~temperature ~~Goal~~goal at ~~Risk~~risk. Nairobi: UNEP~~, Nairobi,~~; 2025.

~~[2] Friedlingstein P, O'Sullivan M, Jones MW, Andrew RM, Hauck J, Landschützer P, Le Quéré C, Li H, Luijkx IT, Olsen A, et al. Global Carbon Budget 2024. Earth System Science Data 2025;17:965-1039.~~

~~[3] Dittmer KM, Wollenberg E, Cohen M, Egler C. How good is the data for tracking countries' agricultural greenhouse gas emissions? Making use of multiple national greenhouse gas inventories. Frontiers in Sustainable Food Systems 2023;7:1156822. https://doi.org/10.3389/fsufs.2023.1156822~~

2. Friedlingstein P, O'Sullivan M, Jones MW, Andrew RM, Bakker DCE, Hauck J, et al. Global Carbon Budget 2025. Earth Syst Sci Data. 2026;18: 3211–3288. doi:10.5194/essd-18-3211-2026

3. United Nations Framework Convention on Climate Change. Decision 18/CMA.1: Modalities, procedures and guidelines for the transparency framework for action and support referred to in Article 13 of the Paris Agreement. FCCC/PA/CMA/2018/3/Add.2. Bonn: UNFCCC; 2019.

4. Cárdenas Rodríguez M, Schwentner E, Falchi E, Suchodolski A, Pizarro R. National greenhouse gas emission inventories, filling the gaps in official data. OECD Environment Working Papers No. 252. Paris: OECD Publishing; 2024. doi:10.1787/f19f1663-en

5. Gütschow J, Jeffery ML, Gieseke R, Gebel R, Stevens D, Krapp M, et al. The PRIMAP-hist national historical emissions time series. Earth Syst Sci Data. 2016;8: 571–603. doi:10.5194/essd-8-571-2016

6. Andrew RM. A comparison of estimates of global carbon dioxide emissions from fossil carbon sources. Earth Syst Sci Data. 2020;12: 1437–1465. doi:10.5194/essd-12-1437-2020

~~[4]~~7. Mostefaoui M, Ciais P, McGrath MJ, Peylin P, Patra PK, Ernst Y. Greenhouse gas emissions and their trends over the last ~~three~~3 decades across Africa. Earth ~~System~~Syst ~~Science~~Sci Data. 2024;16: 245~~–~~–275. ~~https~~doi:~~//doi.org/~~10.5194/essd-16-245-2024

~~[5] OECD. National greenhouse gas emission inventories: filling the gaps in official data. OECD Publishing, Paris, 2024. https://doi.org/10.1787/f19f1663-en~~

~~[6] Machine learning estimates for G20 subnational urban GHG emissions from 2000-2020. Scientific Data 2026;13. https://doi.org/10.1038/s41597-026-06691-9~~

~~[7] Exploring machine learning models for predicting greenhouse gas emissions in Africa's building sector: a case study of six nations. Environmental Systems Research 2025;14. https://doi.org/10.1186/s40068-025-00407-5~~

~~[8] Determinants of carbon emissions in Africa: new evidence based on machine learning algorithms. Environment, Development and Sustainability 2024. https://doi.org/10.1007/s10668-024-05566-6~~

~~[9] Evaluating the machine learning-based models for predicting carbon neutrality in Sub-Saharan African nations. Environment, Development and Sustainability 2025. https://doi.org/10.1007/s10668-025-06289-y~~

~~[10] Predicting transportation-related greenhouse gas emissions in Africa's top emitters: a machine learning approach. Environmental Sciences Europe 2025;37. https://doi.org/10.1186/s12302-025-01223-5~~

~~[11]~~8. Ajala AA, Adeoye OL, Salami OM, Jimoh AY. An examination of daily CO2 emissions prediction through a comparative analysis of machine learning, deep learning, and statistical

models. ~~Environmental~~Environ ~~Science~~Sci ~~and~~Pollut ~~Pollution Research~~Res. 2025;32(5): 2510–2535. ~~[PMC11802685]~~doi:10.1007/s11356-024-35764-8

~~[12]~~9. AlShehri E. Multivariate CO2 emissions forecasting using deep neural network architectures. ~~Machine~~Mach ~~Learning~~Learn ~~and~~Knowl ~~Knowledge Extraction~~Extr. 2026;8(1): 12. ~~https~~doi:~~//doi.org/~~10.3390/make8010012

~~[13] Evaluating deep learning architectures for CO2 emissions forecasting: TCN, LSTM, and hybrid approaches with hyperparameter optimization. Eastern-European Journal of Enterprise Technologies 2025;5(4). https://journals.uran.ua/eejet/article/view/331523~~

~~[14] A hybrid CNN-BiLSTM-attention model with feature engineering for accurate carbon emission forecasting. Air Quality, Atmosphere & Health 2026. https://doi.org/10.1007/s11869-026-01954-1~~

~~[15] Wang R. Carbon emission prediction for gas power plants based on deep learning under small-sample conditions. Concurrency and Computation: Practice and Experience 2026;38. https://doi.org/10.1002/cpe.70591~~

~~[16] Hybrid neural network-based maritime carbon dioxide emission prediction incorporating dynamics for enhanced accuracy. Applied Sciences 2025;15(9):4654. https://doi.org/10.3390/app15094654~~

~~[17] Yin C, Wu J, Sun X, Meng Z, Lee C. Road transportation emission prediction and policy formulation: machine learning model analysis. Transportation Research Part D 2024;135:104390. https://doi.org/10.1016/j.trd.2024.104390~~

~~[18]~~10. Kong F, Song J, Yang Z. A novel short-term carbon emission prediction model based on secondary decomposition method and long short-term memory network. ~~Environmental~~Environ

ScienceSci andPollut Pollution ResearchRes. 2022;29(43): 64983- –64998.
https://doi.org/10.1007/s11356-022-20393-w
[19] Optimised deep learning for time-critical load forecasting using LSTM and modified particle swarm optimisation. Energies 2024;17(22):5524. https://doi.org/10.3390/en17225524
[20] Short-term load forecasting using a particle-swarm optimized multi-head attention-augmented CNN-LSTM network. arXiv preprint 2023;arXiv:2309.03694.
11. Ağbulut Ü. Forecasting of transportation-related energy demand and CO2 emissions in Turkey with different machine learning algorithms. Sustain Prod Consum. 2022;29: 141–157. doi:10.1016/j.spc.2021.10.001
12. Yin C, Wu J, Sun X, Meng Z, Lee C. Road transportation emission prediction and policy formulation: machine learning model analysis. Transp Res D Transp Environ. 2024;135: 104390. doi:10.1016/j.trd.2024.104390
13. Tian L, Zhang Z, He Z, Yuan C, Xie Y, Zhang K, et al. Predicting energy-based CO2 emissions in the United States using machine learning: a path toward mitigating climate change. Sustainability. 2025;17(7): 2843. doi:10.3390/su17072843
14. Agan B, Celik S, Damak OI, Miba'am B. Evaluating the machine learning-based models for predicting carbon neutrality in Sub-Saharan African nations. Environ Dev Sustain. 2025;27(11): 28185–28219. doi:10.1007/s10668-025-06289-y
[21]15. Alhussan AA, Metwally M, Towfek SK. Predicting CO2 emissions with advanced deep learning models and a hybrid Greylag Goose optimization algorithm. Mathematics. 2025;13(9): 1481. https://doi.org/10.3390/math13091481

~~[22] Makridakis S, Spiliotis E, Assimakopoulos V. M5 accuracy competition: results, findings, and conclusions. International Journal of Forecasting 2022;38(4):1346-1364. https://doi.org/10.1016/j.ijforecast.2021.11.013~~

~~[23] Makridakis S, Spiliotis E, Assimakopoulos V, Chen Z, Gaba A, Tsetlin I, Winkler RL. The M5 uncertainty competition: results, findings and conclusions. International Journal of Forecasting 2022;38(4):1365-1385.~~

~~[24] Makridakis S, et al. The M6 forecasting competition: bridging the gap between forecasting and investment decisions. International Journal of Forecasting 2024.~~

~~[25] Energy Institute. Statistical Review of World Energy 2024. Energy Institute, London, 2024.~~

~~[26] Ritchie H, Rosado P, Roser M. CO2 and greenhouse gas emissions. Our World in Data, 2024. https://ourworldindata.org/co2-and-greenhouse-gas-emissions~~

16. Kennedy J, Eberhart R. Particle swarm optimization. Proceedings of ICNN'95 – International Conference on Neural Networks. 1995;4: 1942–1948. doi:10.1109/ICNN.1995.488968

17. Roberts DR, Bahn V, Ciuti S, Boyce MS, Elith J, Guillera-Arroita G, et al. Cross-validation strategies for data with temporal, spatial, hierarchical, or phylogenetic structure. Ecography. 2017;40(8): 913–929. doi:10.1111/ecog.02881

18. Kapoor S, Narayanan A. Leakage and the reproducibility crisis in machine-learning-based science. Patterns. 2023;4(9): 100804. doi:10.1016/j.patter.2023.100804

19. Hyndman RJ, Koehler AB. Another look at measures of forecast accuracy. Int J Forecast. 2006;22(4): 679–688. doi:10.1016/j.ijforecast.2006.03.001

20. Makridakis S, Spiliotis E, Assimakopoulos V. Statistical and Machine Learning forecasting methods: concerns and ways forward. PLoS One. 2018;13(3): e0194889. doi:10.1371/journal.pone.0194889

21. Moulton BR. An illustration of a pitfall in estimating the effects of aggregate variables on micro units. Rev Econ Stat. 1990;72(2): 334–338. doi:10.2307/2109724

22. Bertrand M, Duflo E, Mullainathan S. How much should we trust differences-in-differences estimates? Q J Econ. 2004;119(1): 249–275. doi:10.1162/003355304772839588

23. Cameron AC, Miller DL. A practitioner's guide to cluster-robust inference. J Hum Resour. 2015;50(2): 317–372. doi:10.3368/jhr.50.2.317

24. Lei J, G'Sell M, Rinaldo A, Tibshirani RJ, Wasserman L. Distribution-free predictive inference for regression. J Am Stat Assoc. 2018;113(523): 1094–1111. doi:10.1080/01621459.2017.1307116

~~[27]~~25. Kaya Y. Impact of carbon dioxide emission control on GNP growth: interpretation of proposed scenarios. Paris: IPCC Energy and Industry Subgroup, Response Strategies Working Group~~, Paris,~~; 1990.

~~[28]~~26. Dell'Aversano S, Villante C, Anatone M. Decomposing CO2 emissions with the Kaya identity: global trends, national dynamics, and policy implications. Sustainability. 2026;18(3): 1627. ~~https~~doi:~~//doi.org/~~10.3390/su18031627

~~[29] Kennedy J, Eberhart R. Particle swarm optimization. Proceedings of ICNN'95 - International Conference on Neural Networks 1995;4:1942-1948.~~

~~[30] Varied drivers for reducing CO2 emissions in countries with legislated carbon neutrality targets: a Kaya identity clustering analysis. Environment, Development and Sustainability 2025. https://doi.org/10.1007/s10668-025-06223-2~~

~~[31] Bacanin N, Simic V, Zivkovic M, Alrasheedi M, Petrovic A. Cloud computing load prediction by decomposition reinforced attention long short-term memory network optimized by~~

~~modified particle swarm optimization algorithm. Annals of Operations Research 2023. https://doi.org/10.1007/s10479-023-05745-0~~

27. York R, Rosa EA, Dietz T. STIRPAT, IPAT and ImPACT: analytic tools for unpacking the driving forces of environmental impacts. Ecol Econ. 2003;46(3): 351–365. doi:10.1016/S0921-8009(03)00188-5

28. Karpatne A, Atluri G, Faghmous JH, Steinbach M, Banerjee A, Ganguly A, et al. Theory-guided data science: a new paradigm for scientific discovery from data. IEEE Trans Knowl Data Eng. 2017;29(10): 2318–2331. doi:10.1109/TKDE.2017.2720168

29. Reichstein M, Camps-Valls G, Stevens B, Jung M, Denzler J, Carvalhais N, et al. Deep learning and process understanding for data-driven Earth system science. Nature. 2019;566: 195–204. doi:10.1038/s41586-019-0912-1

30. Willard J, Jia X, Xu S, Steinbach M, Kumar V. Integrating scientific knowledge with machine learning for engineering and environmental systems. ACM Comput Surv. 2022;55(4): 66. doi:10.1145/3514228

31. Ritchie H, Rosado P, Roser M. CO2 and greenhouse gas emissions. Our World in Data; 2023 [cited 2026 Jul 26]. Available from: https://ourworldindata.org/co2-and-greenhouse-gas-emissions

32. Energy Institute. Statistical Review of World Energy 2025. 74th ed. London: Energy Institute; 2025. Available from: https://www.energyinst.org/statistical-review

33. U.S. Energy Information Administration. International Energy Data. Washington (DC): EIA; 2026 [cited 2026 Jul 26]. Available from: https://www.eia.gov/international/data/world

34. Bolt J, van Zanden JL. Maddison-style estimates of the evolution of the world economy: a new 2023 update. J Econ Surv. 2025;39(2): 631–671. doi:10.1111/joes.12618

35. Hoerl AE, Kennard RW. Ridge regression: biased estimation for nonorthogonal problems. Technometrics. 1970;12(1): 55–67. doi:10.1080/00401706.1970.10488634

36. Drucker H, Burges CJC, Kaufman L, Smola A, Vapnik V. Support vector regression machines. Adv Neural Inf Process Syst. 1997;9: 155–161.

37. Breiman L. Random forests. Mach Learn. 2001;45(1): 5–32. doi:10.1023/A:1010933404324

38. Chen T, Guestrin C. XGBoost: a scalable tree boosting system. Proceedings of the 22nd ACM SIGKDD International Conference on Knowledge Discovery and Data Mining. 2016: 785–794. doi:10.1145/2939672.2939785

39. Ke G, Meng Q, Finley T, Wang T, Chen W, Ma W, et al. LightGBM: a highly efficient gradient boosting decision tree. Adv Neural Inf Process Syst. 2017;30: 3146–3154.

40. Hochreiter S, Schmidhuber J. Long short-term memory. Neural Comput. 1997;9(8): 1735–1780. doi:10.1162/neco.1997.9.8.1735

41. Cho K, van Merri ënboer B, Gulcehre C, Bahdanau D, Bougares F, Schwenk H, et al. Learning phrase representations using RNN encoder–decoder for statistical machine translation. Proceedings of the 2014 Conference on Empirical Methods in Natural Language Processing (EMNLP). 2014: 1724–1734. doi:10.3115/v1/D14-1179

42. Bahdanau D, Cho K, Bengio Y. Neural machine translation by jointly learning to align and translate. 3rd International Conference on Learning Representations (ICLR); 2015. Available from: https://arxiv.org/abs/1409.0473

43. Wilcoxon F. Individual comparisons by ranking methods. Biometrics Bull. 1945;1(6): 80–83. doi:10.2307/3001968

44. Cameron AC, Gelbach JB, Miller DL. Bootstrap-based improvements for inference with clustered errors. Rev Econ Stat. 2008;90(3): 414–427. doi:10.1162/rest.90.3.414

45. Honaker J, King G. What to do about missing values in time-series cross-section data. Am J Polit Sci. 2010;54(2): 561–581. doi:10.1111/j.1540-5907.2010.00447.x

[32]46. Gal Y, Ghahramani Z. Dropout as a Bayesian approximation: representing model uncertainty in deep learning. Proceedings of the 33rd International Conference on Machine Learning (ICML). 2016;48: 1050–1059.

47. Vovk V, Gammerman A, Shafer G. Algorithmic learning in a random world. New York: Springer; 2005.

[33]48. Fontana M, Zeni G, Vantini S. Conformal prediction: a unified review of theory and new challenges. Bernoulli. 2023;29(1): 1–23. https://doi.org/10.3150/21-BEJ1447

[34] Jensen V, Bianchi FM, Anfinsen SN. Ensemble conformalized quantile regression for probabilistic time series forecasting. IEEE Transactions on Neural Networks and Learning Systems 2024;35(7):9014-9025. https://doi.org/10.1109/TNNLS.2022.3217694

[35]49. Barber RF, Candès EJ, Ramdas A, Tibshirani RJ. Conformal prediction beyond exchangeability. Annals Ann of Statistics Stat. 2023;51(2): 816–845. https://doi.org/10.1214/23-AOS2276

[36] Lundberg SM, Lee S-I. A unified approach to interpreting model predictions. Advances in Neural Information Processing Systems 2017;30:4765-4774.

[37] Ağbulut Ü. Forecasting of transportation-related energy demand and CO2 emissions in Turkey with different machine learning algorithms. Sustainable Production and Consumption 2022;29:141-157. https://doi.org/10.1016/j.spc.2021.10.001

[38] Janhuaton T, Ratanavaraha V, Jomnonkwao S. Forecasting Thailand's transportation CO2 emissions: a comparison among artificial intelligent models. Forecasting 2024;6(2):462-484. https://doi.org/10.3390/forecast6020026

[39] Li S, Tong Z, Haroon M. Estimation of transport CO2 emissions using machine learning algorithm. Transportation Research Part D 2024;133:104276. https://doi.org/10.1016/j.trd.2024.104276

[40] Jha R, Jha R, Islam M. Forecasting US data center CO2 emissions using AI models: emissions reduction strategies and policy recommendations. Frontiers in Sustainability 2025;5:1507030. https://doi.org/10.3389/frsus.2024.1507030

[41] Tian L, Zhang Z, He Z, Yuan C, Xie Y, Zhang K, Jing R. Predicting energy-based CO2 emissions in the United States using machine learning: a path toward mitigating climate change. Sustainability 2025;17(7):2843. https://doi.org/10.3390/su17072843

[42] A framework for carbon footprint computation and forecasting for Nigeria's industrial decarbonization plan (NIDP). Frontiers in Energy Research 2026;13:1717733. https://doi.org/10.3389/fenrg.2026.1717733

[43] Kaya identity-based decomposition and machine-learning perspectives on US energy-related CO2 emissions 2024. U.S. Energy Information Administration, Washington DC, 2025.

[44] LSTM hyperparameter tuning via modified metaheuristics: an application in renewable energy forecasting. In: Lecture Notes in Networks and Systems. Springer, Cham, 2025. https://doi.org/10.1007/978-3-031-98304-7_74

[45] Vovk V, Gammerman A, Shafer G. Algorithmic Learning in a Random World. Springer, New York, 2005.

[46] Comparative analysis of conformal prediction techniques and machine learning models for very short-term solar power forecasting. Energy and AI 2025. https://doi.org/10.1016/j.egyai.2025.100513

[47] Renkema Y, Visser L, AlSkaif T. Enhancing the reliability of probabilistic PV power forecasts using conformal prediction. Solar Energy Advances 2024;4:100059. https://doi.org/10.1016/j.seja.2024.100059

[48]50. Oliveira RI, Orenstein P, Ramos T, Romano JV. Split conformal prediction and non-exchangeable data. JournalJ ofMach MachineLearn Learning ResearchRes. 2024;25(225): 1–38.

[49] An interpretable machine learning-based framework for CO2 emission prediction and optimization: a case study of a university campus. Sustainability 2025;17(23):10432. https://doi.org/10.3390/su172310432

[50] Deep learning model based prediction of vehicle CO2 emissions with explainable AI integration for sustainable environment. Scientific Reports 2025;15. https://doi.org/10.1038/s41598-025-87233-y

[51] A multi-stage feature selection and explainable machine learning framework for forecasting transportation CO2 emissions. Energies 2025;18(15):4184. https://doi.org/10.3390/en18154184

[52] Chen T, Guestrin C. XGBoost: a scalable tree boosting system. Proceedings of the 22nd ACM SIGKDD International Conference on Knowledge Discovery and Data Mining 2016:785-794.

[53] Ke G, Meng Q, Finley T, Wang T, Chen W, Ma W, Ye Q, Liu T-Y. LightGBM: a highly efficient gradient boosting decision tree. Advances in Neural Information Processing Systems 2017;30:3146-3154.

[54] Hochreiter S, Schmidhuber J. Long short-term memory. Neural Computation 1997;9(8):1735-1780.

[55] Bahdanau D, Cho K, Bengio Y. Neural machine translation by jointly learning to align and translate. International Conference on Learning Representations (ICLR) 2015.

Nomenclature

Symbol	Description	Symbol	Description
[equation removed or moved to S1 Text]	Per-capita CO_2 (t person^{-1} yr^{-1})	[equation removed or moved to S1 Text]	Attention weight at window step τ
[equation removed or moved to S1 Text]	[equation removed or moved to S1 Text] , modelling target	[equation removed or moved to S1 Text]	Attention context vector
[equation removed or moved to S1 Text]	Energy, GDP, population	[equation removed or moved to S1 Text]	Kaya-head (ridge) coefficients
[equation removed or moved to S1 Text]	Fuel shares of CO_2	[equation removed or moved to S1 Text]	Deep residual branch
[equation removed or moved to S1 Text]	Low-carbon share of primary energy	[equation removed or moved to S1 Text]	PSO inertia, cognitive, social constants

Symbol	Description	Symbol	Description
[equation removed or moved to S1 Text]	8-year covariate window (W × F)	[equation removed or moved to S1 Text]	PSO particle position, velocity
[equation removed or moved to S1 Text]	Window length (8), feature count (6/7)	[equation removed or moved to S1 Text]	MC-dropout forward passes (60)
[equation removed or moved to S1 Text]	Fold-wise standardization moments	[equation removed or moved to S1 Text]	Conformal quantile
[equation removed or moved to S1 Text]	LSTM forget/input/output gates	[equation removed or moved to S1 Text]	Normalized nonconformity score
[equation removed or moved to S1 Text]	GRU update/reset gates	[equation removed or moved to S1 Text]	SHAP value of feature j

Symbol	Description	Symbol	Description
[equation removed or moved to S1 Text]	Hidden state at step τ	$MASE_h$	Mean absolute scaled error, horizon h
[equation removed or moved to S1 Text]	Conv1D kernel, layer l	PICP/MPIW	Interval coverage / mean width

Appendix A. Algorithms

Algorithm A1: PSO hyperparameter optimization of KG-CBA.

Input: bounds [l,u] for (filters, kernel, units, dropout, log-lr, batch); swarm size J=6, iterations K=5; country-disjoint train/val split of fold 1 Initialize positions p_j ~ U(l,u), velocities v_j ~ U(−1,1)·0.15(u−l) for k = 1..K: for j = 1..J: hp <- decode(p_j) # round ints, exponentiate lr model <- build_KG_CBA(hp); warm-start Kaya head at last-step ridge solution train with early stopping (patience 6, <=60 epochs) on inner val f_j <- min validation MSE; checkpoint state to disk update personal best (p_j*, f_j*) and global best (g*, f*) v_j <- 0.7 v_j + 1.5 r1 (p_j* − p_j) + 1.5 r2 (g* − p_j); clip to 0.3(u−l) p_j <- clip(p_j + v_j, l, u) Output: g* = (37, 4, 40, 0.231, 9.96e-4, 48), f* = 0.0522

Algorithm A2: KG-CBA training with ridge warm start (per fold).

Input: fold-k training countries; inner split by country (6/7 train, 1/7 val) 1. Standardize windows with training moments (mu_tr, sigma_tr) 2. Fit ridge (lambda=1) on LAST-STEP covariates -> (beta, beta0) 3. Build network: Dense(l) Kaya head initialized at (beta, beta0); Conv1D x2 -> BiLSTM -> dropout -> additive attention -> Dense(32) -> residual head initialized ~ N(0, 1e-3); output = Kaya head + residual 4. Train jointly, Adam(lr), MSE loss, early stopping patience 9, <=90 epochs 5. Save model and scaler; predict held-out countries of fold k

Algorithm A3: Country-clustered comparison of two models.

Input: out-of-fold APE vectors a, b over N country-years; country labels g 1. p_obs <- Wilcoxon signed-rank(a, b) # reported for contrast only 2. per-country medians A_c = median(a | g=c), B_c likewise p_country <- Wilcoxon signed-rank(A, B) over the C countries 3. cluster bootstrap: for r = 1..10000 draw C country labels with replacement delta_r <- median(pooled a − b over drawn countries) CI <- 2.5th and 97.5th percentiles of delta p_boot <- 2 * min(P(delta<=0), P(delta>=0)) Output: (p_obs, p_country, delta, CI, p_boot); conclude from p_boot

Appendix B. Extended tables

Table B1. PSO search space and converged configuration.

Hyperparameter	Range	Best
Conv1D filters	[16, 64]	37
Kernel size	{2, 3, 4}	4
BiLSTM units	[16, 96]	40
Dropout	[0.05, 0.45]	0.231
Learning rate	$[2\times10^{-4}, 3\times10^{-3}]$ (log)	9.96×10^{-4}
Batch size	{48, 64, 96, 128}	48

Table B2. Per-fold MdAPE (%) of the leading models, Experiment A. Fold heterogeneity motivates the block-level uncertainty treatment.

Fold	Ridge	GRU	PSO-KG-CBA	Ensemble
1	11.6	15.7	11.7	10.1
2	23.2	29.5	24.9	25.6
3	18.1	18.4	17.7	18.3
4	15.8	14.2	17.6	13.7
5	13.8	17.3	13.0	10.7

Table B6. Covariate ablation, ridge regression under identical LOCO folds.

Covariate set	RMSE	MAE	MAPE (%)	MdAPE (%)	R^2
All six (energy, GDP, population, three fuel shares)	3.056	1.443	25.2	16.3	0.800
Energy, GDP, population only (no CO_2-derived inputs)	3.244	1.567	26.7	16.9	0.775
Log energy per capita only	3.595	1.690	28.5	17.9	0.723

Table B3. Positioning against representative prior studies of ML-based CO_2 modelling.

Study	Task	Country-disjoint CV	Naive baseline	Clustered inference	Physics guidance	UQ
Ajala et al. 2025 [11]	Daily forecasting, 4 regions	No	Partial	No	No	No
MAKE 2026 [12]	Multivariate forecasting	No	No	No	No	No
AQAH 2026 [14]	Daily national forecasting	No	No	No	No	No
CCPE 2026 [15]	Plant-level prediction	No	No	No	No	No
Applied Sciences 2025 [16]	Maritime emissions	No	No	No	No	No
Africa studies [7-10]	Sectoral prediction	No	No	No	No	No

Study	Task	Country-disjoint CV	Naive baseline	Clustered inference	Physics guidance	UQ
Sci. Data 2026 [6]	Subnational gap-filling	Partial	No	No	No	Partial
This study	**LOCO inventory-gap estimation**	**Yes (grouped 5-fold)**	**Yes (persistence, drift, MASE)**	**Yes (cluster bootstrap)**	**Yes (Kaya head)**	**Yes (conformal MC-dropout)**

Columns record whether each study's *published* protocol includes the listed safeguard, as reported in its methods section; a "No" indicates the safeguard is not described rather than that it was tested and omitted.

Table B4. The 112 economies of the main panel: AFG, ALB, ARE, ARG, ARM, AUS, AUT, AZE, BEL, BEN, BGD, BGR, BIH, BLR, BOL, BRA, BWA, CAN, CHE, CHL, CHN, CMR, COD, COL, CUB, CYP, CZE, DEU, DNK, DOM, DZA, EGY, ESP, EST, FIN, FRA, GBR, GEO, GHA, GRC, GTM, HKG, HRV, HUN, IDN, IND, IRL, IRN, IRQ, ISL, ISR, ITA, JAM, JOR, JPN, KAZ, KGZ, KOR, KWT, LBN, LBY, LTU, LUX, LVA, MAR, MDA, MEX, MKD, MLT, MMR, MOZ, MYS, NER, NGA, NLD, NOR, NZL, OMN, PAK, PAN, PER, PHL, POL, PRT, QAT, ROU, RUS, RWA, SAU, SEN, SGP, SLV, SRB, SVK, SVN, SWE, SYR, THA, TJK, TKM, TUN, TUR, TWN, TZA, UKR, URY, USA, UZB, VEN, VNM, YEM, ZAF. The 76-country enriched sub-panel comprises those with ≥90% low-carbon-share coverage. Per-country imputation status is given in table_imputation_runs.csv.

Table B5. Countries with the longest filled covariate runs (top 12 of the 25 exceeding three years).

ISO	Longest filled run (yr)	ISO	Longest filled run (yr)
SLV	31	RWA	25
NER	29	BEN	24
PAN	28	BOL	21
GHA	26	LBY	15
JAM	26	SEN	12
GTM	26	TUN	11
CMR	26	TZA	9

51. Dunn R, Wasserman L, Ramdas A. Distribution-free prediction sets for two-layer hierarchical models. J Am Stat Assoc. 2023;118(544): 2491–2502. doi:10.1080/01621459.2022.2060112

52. Lundberg SM, Lee S-I. A unified approach to interpreting model predictions. Adv Neural Inf Process Syst. 2017;30: 4765–4774.

53. Lundberg SM, Erion G, Chen H, DeGrave A, Prutkin JM, Nair B, et al. From local explanations to global understanding with explainable AI for trees. Nat Mach Intell. 2020;2: 56–67. doi:10.1038/s42256-019-0138-9

54. Molnar C, König G, Herbinger J, Freiesleben T, Dandl S, Scholbeck CA, et al. General pitfalls of model-agnostic interpretation methods for machine learning models. In: Holzinger A, Goebel R, Fong R, Moon T, Müller K-R, Samek W, editors. xxAI – Beyond Explainable AI. Cham: Springer; 2022. pp. 39–68. doi:10.1007/978-3-031-04083-2_4

55. Fisher A, Rudin C, Dominici F. All models are wrong, but many are useful: learning a variable's importance by studying an entire class of prediction models simultaneously. J Mach Learn Res. 2019;20(177): 1–81.

56. Abadi M, Barham P, Chen J, Chen Z, Davis A, Dean J, et al. TensorFlow: a system for large-scale machine learning. Proceedings of the 12th USENIX Symposium on Operating Systems Design and Implementation (OSDI). 2016: 265–283.

57. Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, et al. Scikit-learn: machine learning in Python. J Mach Learn Res. 2011;12: 2825–2830.

58. Grinsztajn L, Oyallon E, Varoquaux G. Why do tree-based models still outperform deep learning on typical tabular data? Adv Neural Inf Process Syst. 2022;35: 507–520.

59. Shwartz-Ziv R, Armon A. Tabular data: deep learning is not all you need. Inf Fusion. 2022;81: 84–90. doi:10.1016/j.inffus.2021.11.011

60. Zeng A, Chen M, Zhang L, Xu Q. Are transformers effective for time series forecasting? Proc AAAI Conf Artif Intell. 2023;37(9): 11121–11128. doi:10.1609/aaai.v37i9.26317

61. Gelman A, Hill J. Data analysis using regression and multilevel/hierarchical models. Cambridge: Cambridge University Press; 2007.

Supporting information

S1 Text. Supplementary methods. Notation; equations of the recurrent, convolutional and attention layers with all symbols defined; particle swarm update; algorithms for the particle swarm search, the training of the hybrid network and the country-clustered comparison of two models; per-fold results.

S2 Text. Experiment B: temporal forecasting against naive persistence. Protocol, results for two evaluation periods (Table A) and scaled errors by horizon (Fig A).

S1 Table. Countries and data audit. For each of the 112 countries: continent, membership of the 76-country panel, number of filled country-years and longest filled run for each variable, and per-country median absolute percentage error of ridge regression. (XLSX)

S2 Table. Sensitivity analyses and additional results. Results of the missing-data and atypical-economy analyses for ridge regression and the five classical learners; paired tests of ridge regression against the re-estimated classical learners; results by continent; covariate importance; conformal coverage for ridge regression. (XLSX)

S1 Fig. Estimated against observed per-capita CO_2 emissions on held-out countries. Ridge regression, GRU, PSO-KG-CBA and the exploratory ensemble in Experiment A; both axes are logarithmic.

S2 Fig. Correlations between the target and the covariates. Pearson correlation coefficients between the log-transformed target and the six covariates over 3,808 country-years.

S3 Fig. Squared error by country for the exploratory ensemble in Experiment A. Bars show the sum of squared errors of the fifteen countries with the largest contributions (left axis) and the line the cumulative share of the total (right axis). Dark bars mark countries that are among the six worst-fitted under each of the six models examined, and light orange bars the other countries among the six worst-fitted of this model.

S4 Fig. Convergence of the particle swarm search. Validation mean squared error of each of the 30 evaluated configurations and the best value found so far.

S5 Fig. Training and inner-validation loss of the four deep networks with fixed settings in the first fold.

S6 Fig. Mean attention weight by year of the window for the hybrid network of Experiment C. The dotted line marks uniform weights.