Debrief

Run by

Stage 13 dry run (rehearsal, chromium)

Scenario	Stab wound: tamponade, PEA and pericardiocentesis
Patient	M 30 y 80 kg
Started	10/12/2026, 8:52:55 AM
Duration	15:04 simulated · 3 min 47 s on the clock

Objectives


1. **– Not assessable** Name the tamponade within 180 s of arrival from the monitor: tachycardia, narrow pulse pressure, CVP above 15 mmHg (instructor-assessed)
   This objective has no rule: the instructor judges it.
2. **✕ Not met** Start CPR within 10 s of the pulse loss
   CPR 14 s after the pulse loss (07:48 → 08:01; within 10 s asked).
3. **✕ Not met** Pericardiocentesis within 120 s of starting CPR
   The state "CPR: pericardium drained, catheter left in" 2 min 02 s after CPR (08:01 → 10:03; within 2 min 00 s asked).
4. **✓ Met** Stop CPR (“Pulse back”) within 30 s of the pulse returning on the arterial line and the capnogram
   The state "Return of circulation: ventilated, catheter in the pericardium" 26 s after the return of circulation (11:39 → 12:05; within 30 s asked).
5. **✕ Not met** Better: pericardiocentesis before the pulse loss, while the systolic pressure is above 60 mmHg — then no PEA follows
   The pulse loss at 07:48

Measured facts


Lowest SpO_2	95 % at 00:28
Lowest ART M / NIBP M	19 mmHg at 07:58
Highest $EtCO_2$	39 mmHg at 13:05
Highest T1 / T2	36.8 °C at 00:01
Pulseless	3 min 51 s in all; first lost at 07:48
Return of circulation	3 min 51 s after the pulse loss (11:39)
Labs (the model): last (lowest–highest)	K^+ 4.3 mmol/L (4.2–4.3) · Lactate 2.3 mmol/L (1.0–2.4) · pH 7.34 (7.26–7.41) · Glucose 6.0 mmol/L (5.6–6.0) · Tcore 36.7 °C (36.7–36.8)

Trends


HR bpm




ART M / NIBP M mmHg




SpO_2 %




$EtCO_2$ mmHg



awRR / RR /min





Drugs and fluids

Time	Given
12:37	Crystalloid 500 mL over 5 min

Timeline

Time	Kind	Event
00:00	Scenario	Scenario loaded: Stab wound: tamponade, PEA and pericardiocentesis
00:00	Scenario	Scenario state: Arrival: tachycardic, narrow pulse pressure, raised CVP
00:00	Scenario	Breathing spontaneously
00:00	Scenario	Cardiac tamponade: severe
00:01	Action	Speed ×4
05:59	Alarm	SpO_2: low perfusion
06:13	Scenario	Scenario state: Deteriorating: systolic below 80
07:31	Alarm	ART: no pulsatile pressure
07:48	Patient	Pulse lost (the model)
07:49	Scenario	Scenario state: PEA: organised rhythm, no pulse; the pericardium holds about 370 mL
07:49	Scenario	Cardiac tamponade: severe
07:58	Alarm	SpO_2: no pulse detected
08:01	Scenario	Scenario state: CPR, ventilated with FiO_2 1.0
08:01	Scenario	CPR started, 110 /min
08:01	Scenario	Tracheal tube in place
08:01	Scenario	Ventilator: RR 12 /min, VT 500 mL, PEEP 5 cmH_2O, FiO_2 100 %
08:01	Patient	Ventilated breaths began
08:18	Alarm	SpO_2: low perfusion
10:03	Scenario	Scenario state: CPR: pericardium drained, catheter left in
10:03	Scenario	Cardiac tamponade: severe
11:39	Patient	Circulation returned (the model)
12:05	Scenario	Scenario state: Return of circulation: ventilated, catheter in the pericardium
12:05	Scenario	CPR stopped
12:37	Scenario	Scenario state: Post-ROSC: 500 mL crystalloid running
12:37	Scenario	Crystalloid 500 mL over 5 min

Bookmarks

No bookmarks.

Notes

Appendix: one row per minute

Time	HR bpm	ART M / NIBP M mmHg	SpO_2 %	$EtCO_2$ mmHg	awRR / RR /min	T1 / T2 °C	K^+ mmol/L	Lactate mmol/L	pH	Glucose mmol/L	Tcore °C
00:00	72	68	96	36	24	36.8	4.2	1.0	7.41	5.6	36.8
01:00	98	93	98	32	14	36.8	4.2	1.0	7.39	5.6	36.8
02:00	104	90	99	30	14	36.8	4.2	1.0	7.39	5.6	36.8
03:00	112	89	99	29	14	36.8	4.2	1.0	7.38	5.6	36.8
04:00	124	85	99	27	15	36.8	4.2	1.1	7.38	5.6	36.8
05:00	128	80	99	26	15	36.8	4.2	1.1	7.37	5.6	36.8
06:00	135	73	99	24	15	36.8	4.2	1.2	7.37	5.6	36.8
07:00	147	60	99	21	16	36.8	4.2	1.4	7.37	5.6	36.8
08:00	42	20	99	11	16	36.8	4.2	1.6	7.35	5.6	36.8
09:00	115	28	99	8	12	36.8	4.2	1.9	7.31	5.6	36.8
10:00	107	28	99	9	12	36.8	4.2	2.1	7.29	5.7	36.8
11:00	105	46	99	17	12	36.8	4.2	2.3	7.27	5.8	36.8
12:00	121	83	99	25	12	36.8	4.2	2.4	7.26	5.9	36.8
13:00	63	91	99	38	13	36.8	4.3	2.4	7.27	5.9	36.8
14:00	63	91	99	38	22	36.8	4.3	2.3	7.33	6.0	36.8
15:00	62	90	99	39	20	36.8	4.3	2.3	7.34	6.0	36.7