

Glomerular filtration rate

Renal functions include maintaining an acid–base balance; regulating fluid balance; regulating sodium, potassium, and other electrolytes; clearing toxins; absorption of glucose, amino acids, and other small molecules; regulation of blood pressure; production of various hormones, such as erythropoietin; and activation of vitamin D.

The kidney has many functions, which a well-functioning kidney realizes by filtering blood in a process known as glomerular filtration. A major measure of kidney function is the **glomerular filtration rate (GFR)**. The glomerular filtration rate is the flow rate of filtered fluid through the kidney. The **creatinine clearance rate** (C_{Cr} or $CrCl$) is the volume of blood plasma that is cleared of creatinine per unit time and is a useful measure for approximating the GFR. Creatinine clearance exceeds GFR due to creatinine secretion,^[1] which can be blocked by cimetidine. Both GFR and C_{Cr} may be accurately calculated by comparative measurements of substances in the blood and urine, or estimated by formulas using just a blood test result (**eGFR** and **e C_{Cr}**). The results of these tests are used to assess the excretory function of the kidneys. Staging of chronic kidney disease is based on categories of GFR as well as albuminuria and cause of kidney disease.^[2]

Estimated GFR (eGFR) is recommended by clinical practice guidelines and regulatory agencies for routine evaluation of GFR whereas measured GFR (mGFR) is recommended as a confirmatory test when more accurate assessment is required.^[3]

Definition

Glomerular filtration rate (GFR) is the volume of fluid filtered from the renal (kidney) glomerular capillaries into the Bowman's capsule per unit time.^[4]




Diagram showing a schematic nephron and its  blood supply. The basic physiologic mechanisms of handling fluid and electrolytes by the nephron—filtration, secretion, reabsorption, and excretion—are labelled.

GFR is equal to the renal clearance rate when any solute is freely filtered and is neither reabsorbed nor secreted by the kidneys. The rate therefore measured is the quantity of the substance in the urine that originated from a calculable volume of blood. Relating this principle to the below equation – for the substance used, the product of urine concentration and urine flow equals the mass of substance excreted during the time that urine has been collected. This mass equals the mass filtered at the glomerulus as nothing is added or removed in the nephron. Dividing this mass by the plasma concentration gives the volume of plasma which the mass must have originally come from, and thus the volume of plasma fluid that has entered Bowman's capsule within the aforementioned period of time. The GFR is typically recorded in units of *volume per time*, e.g., milliliters per minute (mL/min). Compare to filtration fraction.

$$GFR = \frac{\text{Urine concentration} \times \text{urine flow}}{\text{Plasma concentration}}$$

There are several different techniques used to calculate or estimate the glomerular filtration rate (GFR or eGFR). The above formula only applies for GFR calculation when it is equal to the clearance rate.

Regulation

Central to the physiologic maintenance of GFR is the differential vasoconstriction of the afferent (input) and efferent (output) arterioles. Vasodilation of the afferent or vasoconstriction of the efferent arteriole increases GFR. Conversely, vasoconstriction of the afferent or vasodilation of the efferent will decrease GFR.^[5]

There are multiple mechanisms that the body has to regulate the constriction or dilation of these arterioles: local feedback within the kidney, hormonal regulation, and sympathetic nervous control. These systems are closely intertwined. The kidney can maintain a relatively constant GFR even as mean arterial pressure changes.^[6]

Local feedback within the kidney (renal autoregulation) includes the myogenic response and tubuloglomerular feedback.^[6] In the myogenic response, increased stretch of the afferent arteriole due to higher blood pressure results in afferent vasoconstriction, thereby preventing an increase in GFR.^[5] With tubuloglomerular feedback, the macula densa cells at the downstream nephron tubule senses changes in ion levels. If sodium chloride sodium levels in the urinary filtrate are too high, the macula densa signals to constrict the afferent arteriole; if too low, the signal is to vasodilate the afferent arteriole and to release renin (activating the RAAS system).^[5]

The RAAS system is the major hormonal control of GFR. Low blood pressure activates the RAAS system, increasing the amount of the hormone angiotensin II. Angiotensin II binds to receptors on the efferent arteriole; the resulting efferent vasoconstriction increases GFR.^[5]

There is sympathetic innervation of both afferent and efferent arterioles at the glomerulus.^[7] Sympathetic activation, for instance during a state of stress, causes vasoconstriction of both the afferent and efferent arterioles. This decreases renal blood flow and GFR. Sympathetic activation

also causes the release of renin.^[7]

Measurement

Creatinine

In clinical practice, **creatinine clearance** or estimates of creatinine clearance based on the serum creatinine level are used to measure GFR. Creatinine is produced naturally by the body (creatinine is a breakdown product of creatine phosphate, which is found in muscle). It is freely filtered by the glomerulus, but also actively secreted by the peritubular capillaries in very small amounts such that creatinine clearance overestimates actual GFR by 10–20%. This margin of error is acceptable, considering the ease with which creatinine clearance is measured. Unlike precise GFR measurements involving constant infusions of inulin, creatinine is already at a steady-state concentration in the blood, and so measuring creatinine clearance is much less cumbersome. However, creatinine estimates of GFR have their limitations. All of the estimating equations depend on a prediction of the 24-hour creatinine excretion rate, which is a function of muscle mass which is quite variable. One of the equations, the Cockcroft and Gault equation (see below) does not correct for race. With a higher muscle mass, serum creatinine will be higher for any given rate of clearance.

Inulin

The GFR can be determined by injecting inulin or the inulin analog sinistrin into the blood stream. Since both inulin and sinistrin are neither reabsorbed nor secreted by the kidney after glomerular filtration, their rate of excretion is directly proportional to the rate of filtration of water and solutes across the glomerular filter. Incomplete urine collection is an important source of error in inulin clearance measurement.^[8] Using inulin to measure kidney function is the "gold standard" for comparison with other means of estimating glomerular filtration rate.^[9] In 2018, the French pharmacovigilance agency withdrew inulin and sinistrin-based products from the market after some patients experienced hypersensitivity reactions including a fatal outcome.^[10] Consequently, the contrast agents iohexol and iothalamate have become more popular alternatives to determine GFR and are considered to show sufficient accuracy to determine GFR.^[11]

Radioactive tracers

GFR can be accurately measured using radioactive substances, in particular chromium-51 and technetium-99m. These come close to the ideal properties of inulin (undergoing only glomerular filtration) but can be measured more practically with only a few urine or blood samples.^[12] Measurement of renal or plasma clearance of ⁵¹Cr-EDTA is widely used in Europe but not available in the United States, where ^{99m}Tc-DTPA may be used instead.^[13] Renal and plasma clearance ⁵¹Cr-EDTA has been shown to be accurate in comparison with the gold standard, inulin.^{[14][15][9]} Use of ⁵¹Cr-EDTA is considered a reference standard measure in UK guidance.^[16]

Cystatin C

Problems with creatinine (varying muscle mass, recent meat ingestion (much less dependent on the diet than urea), etc.) have led to evaluation of alternative agents for estimation of GFR. One of these is cystatin C, a ubiquitous protein secreted by most cells in the body (it is an inhibitor of cysteine protease).^[17]

Cystatin C is freely filtered at the glomerulus. After filtration, Cystatin C is reabsorbed and catabolized by the tubular epithelial cells, with only small amounts excreted in the urine. Cystatin C levels are therefore measured not in the urine, but in the bloodstream.

Equations have been developed linking estimated GFR to serum cystatin C levels.^[18] Most recently, some proposed equations have combined sex, age, adjusted cystatin C and creatinine. In 2022, the National Kidney Foundation (NKF) and American Society of Nephrology (ASN)'s Joint Task Force on Reassessing the Inclusion of Race in Diagnosing Kidney Diseases recommended national efforts to facilitate increased, routine, and timely use of cystatin C. They noted that cystatin C would be useful particularly to confirm estimated GFR in adults who are at risk for or have chronic kidney disease. They suggested that combining filtration markers (creatinine and cystatin C) is more accurate and would support better clinical decisions than either marker alone.^[19]

Calculation

More precisely, GFR is the fluid flow rate between the glomerular capillaries and the Bowman's capsule:

$$Q = \frac{dV}{dt} = K_f \times (P_G - P_B - \Pi_G + \Pi_B)^{[20][21]}$$

Where:

- Q is the GFR.
- K_f is called the *filtration constant* and is defined as the product of the hydraulic conductivity and the surface area of the glomerular capillaries.
- P_G is the hydrostatic pressure within the glomerular capillaries.
- P_B is the hydrostatic pressure within the Bowman's capsule.
- Π_G is the colloid osmotic pressure within the glomerular capillaries.
- and Π_B is the colloid osmotic pressure within the Bowman's capsule.

K_f

Because this constant is a measurement of hydraulic conductivity multiplied by the capillary surface area, it is almost impossible to measure physically. However, it can be determined experimentally. Methods of determining the GFR are listed in the above and below sections and it

is clear from our equation that K_f can be found by dividing the experimental GFR by the net filtration pressure:^[20]

$$K_f = \frac{\text{GFR}}{\text{Net Filt. Pressure}} = \frac{\text{GFR}}{(P_G - P_B - \Pi_G + \Pi_B)}$$

P_G

The hydrostatic pressure within the glomerular capillaries is determined by the pressure difference between the fluid entering immediately from the afferent arteriole and leaving through the efferent arteriole. The pressure difference is approximated by the product of the total resistance of the respective arteriole and the flux of blood through it:^[21]

$$P_a - P_G = R_a \times Q_a$$

$$P_G - P_e = R_e \times Q_e$$

Where:

- P_a is the afferent arteriole pressure.
- P_G is the hydrostatic pressure within the glomerular capillaries.
- P_e is the efferent arteriole pressure.
- R_a is the afferent arteriole resistance.
- R_e is the efferent arteriole resistance.
- Q_a is the afferent arteriole flux.
- And, Q_e is the efferent arteriole flux.

P_B

The pressure in the Bowman's capsule and proximal tubule can be determined by the difference between the pressure in the Bowman's capsule and the descending tubule:^[21]

$$P_B - P_d = R_d \times (Q_a - Q_e)$$

Where:

- P_d is the pressure in the descending tubule.
- And, R_d is the resistance of the descending tubule.

Π_G

Blood plasma has a good many proteins in it and they exert an inward directed force called the osmotic pressure on the water in hypotonic solutions across a membrane, i.e., in the Bowman's capsule. Because plasma proteins are virtually incapable of escaping the glomerular capillaries, this oncotic pressure is defined, simply, by the ideal gas law:^{[20][21]}

$$\Pi_G = RTc$$

Where:

- R is the universal gas constant
- T is the temperature.
- And, c is concentration in mol/L of plasma *proteins* (remember the solutes can freely diffuse through the glomerular capsule).

$$\Pi_B$$

This value is almost always taken to be equal to zero because in a healthy nephron, there should be no proteins in the Bowman's capsule.^[20]

Clearance and filtration fraction

Filtration fraction

The filtration fraction is the amount of plasma that is actually filtered through the kidney. This can be defined using the equation:

$$FF = \frac{GFR}{RPF}$$

- FF is the filtration fraction
- GFR is the glomerular filtration rate
- RPF is the renal plasma flow

Normal human FF is 20%.

Renal clearance

$$C_x = (U_x) \frac{V}{P_x}$$

- C_x is the clearance of X (normally in units of mL/min).
- U_x is the urine concentration of X.
- P_x is the plasma concentration of X.
- V is the urine flow rate.

Estimation

In clinical practice, however, **creatinine clearance** or estimates of creatinine clearance based on the serum creatinine level are used to measure GFR. Creatinine is produced naturally by the body (creatinine is a breakdown product of creatine phosphate, which is found in muscle). It is freely filtered by the glomerulus, but also actively secreted by the peritubular capillaries in very small amounts such that creatinine clearance overestimates actual GFR by 10–20%. This margin of error is acceptable, considering the ease with which creatinine clearance is measured. Unlike precise GFR measurements involving constant infusions of inulin, creatinine is already at a steady-state concentration in the blood, and so measuring creatinine clearance is much less cumbersome. However, creatinine estimates of GFR have their limitations. All of the estimating equations depend on a prediction of the 24-hour creatinine excretion rate, which is a function of muscle mass which is quite variable. The Cockcroft-Gault and CKD-EPI 2021 equations (see below) do not correct for race. With a higher muscle mass, serum creatinine will be higher for any given rate of clearance.^[22]

A common mistake made when just looking at serum creatinine is the failure to account for muscle mass. Hence, an older woman with a serum creatinine of 1.4 mg/dL may actually have a moderately severe chronic kidney disease, whereas a young muscular male can have a normal level of renal function at this serum creatinine level. Creatinine-based equations should be used with caution in cachectic patients and patients with cirrhosis. They often have very low muscle mass and a much lower creatinine excretion rate than predicted by the equations below, such that a cirrhotic patient with a serum creatinine of 0.9 mg/dL may have a moderately severe degree of chronic kidney disease.

Estimated GFR (eGFR) is recommended by clinical practice guidelines and regulatory agencies for routine evaluation of GFR whereas measured GFR (mGFR) is recommended as a confirmatory test when more accurate assessment is required.^[3]

Creatinine clearance C_{Cr}

One method of determining GFR from creatinine is to collect urine (usually for 24 h) to determine the amount of creatinine that was removed from the blood over a given time interval. If one removes 1440 mg in 24 h, this is equivalent to removing 1 mg/min. If the blood concentration is 0.01 mg/mL (1 mg/dL), then one can say that 100 mL/min of blood is being "cleared" of creatinine, since, to get 1 mg of creatinine, 100 mL of blood containing 0.01 mg/mL would need to have been cleared.

Creatinine clearance (C_{Cr}) is calculated from the creatinine concentration in the collected urine sample (U_{Cr}), urine flow rate (Vdt), and the plasma concentration (P_{Cr}). Since the product of urine concentration and urine flow rate yields creatinine excretion rate, which is the rate of removal from the blood, creatinine clearance is calculated as removal rate per min ($U_{Cr} \times Vdt$) divided by the plasma creatinine concentration. This is commonly represented mathematically as

$$C_{Cr} = \frac{U_{Cr} \times \dot{V}}{P_{Cr}}$$

Example: A person has a plasma creatinine concentration of 0.01 mg/mL and in 1 hour produces 60 mL of urine with a creatinine concentration of 1.25 mg/mL.

$$C_{Cr} = \frac{1.25 \text{ mg/mL} \times \frac{60 \text{ mL}}{60 \text{ min}}}{0.01 \text{ mg/mL}} = \frac{1.25 \text{ mg/mL} \times 1 \text{ mL/min}}{0.01 \text{ mg/mL}} = \frac{1.25 \text{ mg/min}}{0.01 \text{ mg/mL}} = 125 \text{ mL/min}$$

The common procedure involves undertaking a 24-hour urine collection, from empty-bladder one morning to the contents of the bladder the following morning, with a comparative blood test then taken. The urinary flow rate is still calculated per minute, hence:

$$C_{Cr} = \frac{U_{Cr} \times 24\text{-hour volume}}{P_{Cr} \times 24 \times 60 \text{ min}}$$

To allow comparison of results between people of different sizes, the C_{Cr} is often corrected for the body surface area (BSA) and expressed compared to the average sized man as mL/min/1.73 m². While most adults have a BSA that approaches 1.7 m² (1.6 m² to 1.9 m²), extremely obese or slim patients should have their C_{Cr} corrected for their *actual* BSA.

$$C_{Cr\text{-corrected}} = \frac{C_{Cr} \times BSA}{1.73}$$

BSA can be calculated on the basis of weight and height.

Twenty-four-hour urine collection to assess creatinine clearance is no longer widely performed, due to difficulty in assuring complete specimen collection. To assess the adequacy of a complete collection, one always calculates the amount of creatinine excreted over a 24-hour period. This amount varies with muscle mass and is higher in young people/old, and in men/women. An unexpectedly low or high 24-hour creatinine excretion rate voids the test. Nevertheless, in cases where estimates of creatinine clearance from serum creatinine are unreliable, creatinine clearance remains a useful test. These cases include "estimation of GFR in individuals with variation in dietary intake (vegetarian diet, creatine supplements) or muscle mass (amputation, malnutrition, muscle wasting), since these factors are not specifically taken into account in prediction equations."^[23]

A number of formulae have been devised to estimate GFR or C_{Cr} values on the basis of serum creatinine levels. Where not otherwise stated serum creatinine is assumed to be stated in mg/dL, not μmol/L—divide by 88.4 to convert from μmol/L to mg/dL.

Cockcroft–Gault formula

A commonly used surrogate marker for the estimation of creatinine clearance is the Cockcroft–Gault (CG) formula, which in turn estimates GFR in mL/min:^[24] It is named after the scientists, the asthmologist Donald William Cockcroft (b. 1946) and the nephrologist Matthew Henry Gault

(1925–2003), who first published the formula in 1976, and it employs serum creatinine measurements and a patient's weight to predict the creatinine clearance.^{[25][26]} The formula, as originally published, is:

$$eC_{Cr} = \frac{(140 - \text{Age}) \times \text{Mass (in kilograms)} \times [0.85 \text{ if Female}]}{72 \times [\text{Serum Creatinine (in mg/dL)}]}$$

This formula expects weight to be measured in kilograms and creatinine to be measured in mg/dL, as is standard in the US. The resulting value is multiplied by a constant of 0.85 if the patient is female. This formula is useful because the calculations are simple and can often be performed without the aid of a calculator.

When serum creatinine is measured in $\mu\text{mol/L}$:

$$eC_{Cr} = \frac{(140 - \text{Age}) \times \text{Mass (in kilograms)} \times \text{Constant}}{[\text{Serum Creatinine (in } \mu\text{mol/L)}]}$$

Where *Constant* is 1.23 for men and 1.04 for women.

One interesting feature of the Cockcroft and Gault equation is that it shows how dependent the estimation of CCr is based on age. The age term is $(140 - \text{age})$. This means that a 20-year-old person $(140 - 20 = 120)$ will have twice the creatinine clearance as an 80-year-old $(140 - 80 = 60)$ for the same level of serum creatinine. The C-G equation assumes that a woman will have a 15% lower creatinine clearance than a man at the same level of serum creatinine.

Modification of diet in renal disease (MDRD) formula

Another formula for calculating the GFR is the one developed by the *Modification of Diet in Renal Disease Study Group*.^[27] Most laboratories in Australia,^[28] and the United Kingdom calculate and report the estimated GFR along with creatinine measurements and this forms the basis of diagnosis of chronic kidney disease.^{[29][30]} The adoption of the automatic reporting of MDRD-eGFR has been widely criticised.^{[31][32][33]}

The most commonly used formula is the "4-variable MDRD", which estimates GFR using four variables: serum creatinine, age, ethnicity, and gender.^[34] The original MDRD used six variables with the additional variables being the blood urea nitrogen and albumin levels.^[27] The equations have been validated in patients with chronic kidney disease; however, both versions underestimate the GFR in healthy patients with GFRs over 60 mL/min.^{[35][36]} The equations have not been validated in acute renal failure.

For creatinine in $\mu\text{mol/L}$:

$$eGFR = 32788 \times [\text{Serum Creatinine}]^{-1.154} \times \text{Age}^{-0.203} \times [1.212 \text{ if Black}] \times [0.742 \text{ if Female}]$$

For creatinine in mg/dL:

$$eGFR = 186 \times [\text{Serum Creatinine}]^{-1.154} \times \text{Age}^{-0.203} \times [1.212 \text{ if Black}] \times [0.742 \text{ if Female}]$$

Creatinine levels in $\mu\text{mol/L}$ can be converted to mg/dL by dividing them by 88.4. The 32788 number above is equal to $186 \times 88.4^{1.154}$.


A more elaborate version of the MDRD equation also includes serum albumin and blood urea nitrogen (BUN) levels:

$$\text{eGFR} = 170 \times [\text{Serum Creatinine}]^{-0.999} \times \text{Age}^{-0.176} \times [0.762 \text{ if Female}] \times [1.180 \text{ if Black}] \times \text{BUN}^{-0.170} \times \text{Albumin}^{+0.318}$$

where the creatinine and blood urea nitrogen concentrations are both in mg/dL . The albumin concentration is in g/dL .

These MDRD equations are to be used only if the laboratory has NOT calibrated its serum creatinine measurements to isotope dilution mass spectrometry (IDMS). When IDMS-calibrated serum creatinine is used (which is about 6% lower), the above equations should be multiplied by 175/186 or by 0.94086.^[37]

Since these formulae do not adjust for body size, results are given in units of $\text{mL/min per } 1.73 \text{ m}^2$, 1.73 m^2 being the estimated body surface area of an adult with a mass of 63 kg and a height of 1.7 m.



CKD-EPI formula

The CKD-EPI (Chronic Kidney Disease Epidemiology Collaboration) formula was first published in May 2009. It was developed in an effort to create a formula more accurate than the MDRD formula, especially when actual GFR is greater than 60 mL/min per 1.73 m². This is the formula recommended as of 2014 by NICE in the UK.^[30]

Researchers pooled data from multiple studies to develop and validate this new equation. They used 10 studies that included 8254 participants, randomly using 2/3 of the data sets for development and the other 1/3 for internal validation. Sixteen additional studies, which included 3896 participants, were used for external validation.

The CKD-EPI equation performed better than the MDRD (Modification of Diet in Renal Disease Study) equation, especially at higher GFR, with less bias and greater accuracy. When looking at NHANES (National Health and Nutrition Examination Survey) data, the median estimated GFR was 94.5 mL/min per 1.73 m² vs. 85.0 mL/min per 1.73 m², and the prevalence of chronic kidney disease was 11.5% versus 13.1%. Despite its overall superiority to the MDRD equation, the CKD-EPI equations performed poorly in certain populations, including black women, the elderly and the obese, and was less popular among clinicians than the MDRD estimate.^[39]

The 2009 CKD-EPI equation is:

$$\text{eGFR} = 141 \times \min(\text{SCr}/k, 1)^a \times \max(\text{SCr}/k, 1)^{-1.209} \times 0.993^{\text{Age}} \times [1.018 \text{ if Female}] \times [1.159 \text{ if Black}]$$

where SCr is serum creatinine (mg/dL), k is 0.7 for females and 0.9 for males, a is -0.329 for females and -0.411 for males, min indicates the minimum of SCr/k or 1, and max indicates the maximum of SCr/k or 1.

As separate equations for different populations: For creatinine (IDMS calibrated) in mg/dL:

Male, not black

If serum creatinine (Scr) ≤ 0.9

$$\text{eGFR} = 141 \times (\text{SCr}/0.9)^{-0.411} \times 0.993^{\text{Age}}$$

If serum creatinine (Scr) > 0.9

$$\text{eGFR} = 141 \times (\text{SCr}/0.9)^{-1.209} \times 0.993^{\text{Age}}$$

Female, not black

If serum creatinine (Scr) ≤ 0.7

$$\text{eGFR} = 144 \times (\text{SCr}/0.7)^{-0.329} \times 0.993^{\text{Age}}$$

If serum creatinine (Scr) > 0.7

$$\text{eGFR} = 144 \times (\text{SCr}/0.7)^{-1.209} \times 0.993^{\text{Age}}$$

Black male

If serum creatinine (Scr) ≤ 0.9

$$\text{eGFR} = 163 \times (\text{SCr}/0.9)^{-0.411} \times 0.993^{\text{Age}}$$

If serum creatinine (Scr) > 0.9

$$\text{eGFR} = 163 \times (\text{SCr}/0.9)^{-1.209} \times 0.993^{\text{Age}}$$

Black female

If serum creatinine (Scr) ≤ 0.7

$$\text{eGFR} = 166 \times (\text{Scr}/0.7)^{-0.329} \times 0.993^{\text{Age}}$$

If serum creatinine (Scr) > 0.7

$$\text{eGFR} = 166 \times (\text{Scr}/0.7)^{-1.209} \times 0.993^{\text{Age}}$$

This formula was developed by Levey et al.^[40]

The 2009 CKD-EPI formula was suggested to improve cardiovascular risk prediction over the MDRD Study formula in a middle-age population.^[41]

The 2021 CKD-EPI formula^[42] does not include a race coefficient (see discussion below). The 2021 CKD-EPI equation is:

$$\text{eGFR} = 142 \times \min(\text{Scr}/k, 1)^a \times \max(\text{Scr}/k, 1)^{-1.209} \times 0.9938^{\text{Age}} \times [1.012 \text{ if Female}]$$

where Scr is serum creatinine (mg/dL), k is 0.7 for females and 0.9 for males, a is -0.241 for females and -0.302 for males, min indicates the minimum of Scr/k or 1, and max indicates the maximum of Scr/k or 1.

As separate equations for different populations: For creatinine (IDMS calibrated) in mg/dL:

Male

If serum creatinine (Scr) ≤ 0.9

$$\text{eGFR} = 142 \times (\text{Scr}/0.9)^{-0.302} \times 0.9938^{\text{Age}}$$

If serum creatinine (Scr) > 0.9

$$\text{eGFR} = 142 \times (\text{Scr}/0.9)^{-1.2} \times 0.9938^{\text{Age}}$$

Female

If serum creatinine (Scr) ≤ 0.7

$$\text{eGFR} = 143 \times (\text{Scr}/0.7)^{-0.241} \times 0.9938^{\text{Age}}$$

If serum creatinine (Scr) > 0.7

$$\text{eGFR} = 143 \times (\text{Scr}/0.7)^{-1.2} \times 0.9938^{\text{Age}}$$

Mayo Quadratic formula

Another estimation tool to calculate GFR is the Mayo Quadratic formula. This formula was developed by Rule et al.,^[35] in an attempt to better estimate GFR in patients with preserved kidney function. It is well recognized that the MDRD formula tends to underestimate GFR in patients with preserved kidney function. Studies in 2008 found that the Mayo Clinic Quadratic Equation compared moderately well with radionuclide GFR, but had inferior bias and accuracy than the MDRD equation in a clinical setting.^{[43][44]}

The equation is:

$$\text{eGFR} = \exp(1.911 + 5.249/[\text{Serum Creatinine}] - 2.114/[\text{Serum Creatinine}]^2 - 0.00686 \times \text{Age} - [0.205 \text{ if Female}])$$

^[44]

If Serum Creatinine < 0.8 mg/dL, use 0.8 mg/dL for Serum Creatinine.

Schwartz formula

In children, the Schwartz formula is used.^{[45][46]} This employs the serum creatinine (mg/dL), the child's height (cm) and a constant to estimate the glomerular filtration rate:

$$\text{eGFR} = \frac{k \times \text{Height}}{\text{Serum Creatinine}}$$

Where k is a constant that depends on muscle mass, which itself varies with a child's age:

In first year of life, for pre-term babies $k=0.33$ ^[47] and for full-term infants $k=0.45$ ^[46]

For infants and children of age 1 to 12 years, $k=0.55$.^[45]

The method of selection of the constant k has been questioned as being dependent upon the gold-standard of renal function used (i.e. inulin clearance, creatinine clearance, etc.) and also may be dependent upon the urinary flow rate at the time of measurement.^[48]

In 2009 the formula was updated to use standardized serum creatinine (recommend $k=0.413$) and additional formulas that allow improved precision were derived if serum *cystatin C* is measured in addition to serum creatinine.^[49]

IDMS standardization effort

One problem with any creatinine-based equation for GFR is that the methods used to assay creatinine in the blood differ widely in their susceptibility to non-specific chromogens, which cause the creatinine value to be overestimated. In particular, the MDRD equation was derived using serum creatinine measurements that had this problem. The NKDEP program in the United States has attempted to solve this problem by trying to get all laboratories to calibrate their measures of creatinine to a "gold standard", which in this case is isotope dilution mass spectrometry (IDMS). In late 2009 not all labs in the U.S. had changed over to the new system. There are two forms of the MDRD equation that are available, depending on whether or not creatinine was measured by an IDMS-calibrated assay. The CKD-EPI equation is designed to be used with IDMS-calibrated serum creatinine values only.

Normal ranges

The normal range of GFR, adjusted for body surface area, is 100–130 average 125 mL/min/1.73m² in men and 90–120 mL/min/1.73m² in women younger than the age of 40. In children, GFR measured by inulin clearance is 110 mL/min/1.73 m² until 2 years of age in both sexes, and then it progressively decreases. After age 40, GFR decreases progressively with age, by 0.4–1.2 mL/min per year.

Decreased GFR

A decreased renal function can be caused by many types of kidney disease. Upon presentation of decreased renal function, it is recommended to perform a history and physical examination, as well as performing a renal ultrasound and a urinalysis. The most relevant items in the history are medications, edema, nocturia, gross hematuria, family history of kidney disease, diabetes and polyuria. The most important items in a physical examination are signs of vasculitis, lupus erythematosus, diabetes, endocarditis and hypertension.

A urinalysis is helpful even when not showing any pathology, as this finding suggests an extrarenal etiology. Proteinuria and/or urinary sediment usually indicates the presence of glomerular disease. Hematuria may be caused by glomerular disease or by a disease along the urinary tract.

The most relevant assessments in a renal ultrasound are renal sizes, echogenicity and any signs of hydronephrosis. Renal enlargement usually indicates diabetic nephropathy, focal segmental glomerular sclerosis or myeloma. Renal atrophy suggests longstanding chronic renal disease.

Chronic kidney disease stages

Risk factors for kidney disease include diabetes, high blood pressure, family history, older age, ethnic group and smoking. For most patients, a GFR over 60 mL/min/1.73 m² is adequate. But significant decline of the GFR from a previous test result can be an early indicator of kidney disease requiring medical intervention. The sooner kidney dysfunction is diagnosed and treated the greater odds of preserving remaining nephrons, and preventing the need for dialysis.

The severity of chronic kidney disease (CKD) is described by six stages; the most severe three are defined by the MDRD-eGFR value, and first three also depend on whether there is other evidence of kidney disease (e.g., proteinuria):

CKD stage	GFR level (mL/min/1.73 m ²)
Stage 1	≥ 90
Stage 2	60–89
Stage 3	30–59
Stage 4	15–29
Stage 5	< 15

0) Normal kidney function – GFR above 90 mL/min/1.73 m² and no proteinuria

1) CKD1 – GFR above 90 mL/min/1.73 m² with evidence of kidney damage

2) CKD2 (mild) – GFR of 60 to 89 mL/min/1.73 m² with evidence of kidney damage

3) CKD3 (moderate) – GFR of 30 to 59 mL/min/1.73 m²

4) CKD4 (severe) – GFR of 15 to 29 mL/min/1.73 m²

5) CKD5 kidney failure – GFR less than 15 mL/min/1.73 m² Some people add CKD5D for those stage 5 patients requiring dialysis; many patients in CKD5 are not yet on dialysis.

Note: others add a "T" to patients who have had a transplant regardless of stage.

Not all clinicians agree with the above classification, suggesting that it may mislabel patients with mildly reduced kidney function, especially the elderly, as having a disease.^{[50][51]} A conference was held in 2009 regarding these controversies by Kidney Disease: Improving Global Outcomes

(KDIGO) on CKD: Definition, Classification and Prognosis, gathering data on CKD prognosis to refine the definition and staging of CKD.^[52]

Removal of race from eGFR calculations

In 2017, Beth Israel Deaconess Medical Center dropped the Black race coefficient in their estimated glomerular filtration rate (eGFR) report, resulting in a steady increase in the percentage of Black patients listed before starting dialysis. Hoenig et al said on the topic "Race is a social construct that cannot be measured, can be used imprecisely and may contribute to disparities in kidney transplant access for Black patients."^[53]

In 2020, Vyas et al published an article in *The New England Journal of Medicine* criticizing the use of race in eGFR calculations which resulted in higher reported eGFR values for black-identifying patients. They noted that "The algorithm developers justified these outcomes with evidence of higher average serum creatinine concentrations among black people than among white people. Explanations that have been given for this finding include the notion that black people release more creatinine into their blood at baseline, in part because they are reportedly more muscular. Analyses have cast doubt on this claim, but the "race-corrected" eGFR remains the standard." They concluded saying "As long as uncertainty persists about the cause of racial differences in serum creatinine levels, we should favor practices that may alleviate health inequities over those that may exacerbate them."^[54]

In February 2022, the National Kidney Foundation (NKF) and American Society of Nephrology (ASN)'s Joint Task Force on Reassessing the Inclusion of Race in Diagnosing Kidney Diseases recommended use of the 2021 CKD-EPI equation which discontinues using a race modifier.^[19]

In July 2022 the OPTN Board eliminated race-based calculations for transplant candidate listing across all transplant hospitals in the US.^[55]

See also

- Clearance
- Dialysis
- Filtration fraction
- Kt/V
- Pharmacokinetics
- Renal clearance ratio
- Renal failure
- Standardized Kt/V
- Tubuloglomerular feedback
- Urea reduction ratio

References

1. Ganong (2016). "Renal Function & Micturition". *Review of Medical Physiology* (25th ed.). McGraw-Hill Education. p. 677. ISBN 978-0-07-184897-8.
2. Stevens PE, Levin A (Jun 4, 2013). "Evaluation and management of chronic kidney disease: synopsis of the kidney disease: improving global outcomes 2012 clinical practice guideline". *Annals of Internal Medicine*. **158** (11): 825–830. doi:10.7326/0003-4819-158-11-201306040-00007 (<https://doi.org/10.7326/0003-4819-158-11-201306040-00007>). ISSN 1539-3704 (<https://search.worldcat.org/issn/1539-3704>). PMID 23732715 (<https://pubmed.ncbi.nlm.nih.gov/23732715>).
3. Levey AS, Coresh J, Tighiouart H, Greene T, Inker LA (2020). "Measured and estimated glomerular filtration rate: current status and future directions". *Nat Rev Nephrol*. **16** (1): 51–64. doi:10.1038/s41581-019-0191-y (<https://doi.org/10.1038/s41581-019-0191-y>). PMID 31527790 (<https://pubmed.ncbi.nlm.nih.gov/31527790>). S2CID 202573933 (<https://api.semanticscholar.org/CorpusID:202573933>).
4. Nosek TM. "Section 7/7ch04/7ch04p11" (<https://web.archive.org/web/20160324124828/http://humanphysiology.tuars.com/program/section7/7ch04/7ch04p11.htm>). *Essentials of Human Physiology*. Archived from the original (<http://humanphysiology.tuars.com/program/section7/7ch04/7ch04p11.htm>) on 2016-03-24. – "Glomerular Filtration Rate"
5. Ogobuiro I, Tuma F (2026), "Physiology, Renal" (<http://www.ncbi.nlm.nih.gov/books/NBK538339/>), *StatPearls*, Treasure Island (FL): StatPearls Publishing, PMID 30855923 (<https://pubmed.ncbi.nlm.nih.gov/30855923>), retrieved 2026-06-20
6. Carlström M, Wilcox CS, Arendshorst WJ (April 2015). "Renal autoregulation in health and disease" (<https://pmc.ncbi.nlm.nih.gov/articles/PMC4551215/>). *Physiological Reviews*. **95** (2): 405–511. doi:10.1152/physrev.00042.2012 (<https://doi.org/10.1152/physrev.00042.2012>). ISSN 1522-1210 (<https://search.worldcat.org/issn/1522-1210>). PMC 4551215 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4551215>). PMID 25834230 (<https://pubmed.ncbi.nlm.nih.gov/25834230>).
7. Kopp UC (2011), "Neural Control of Renal Hemodynamics" (<https://www.ncbi.nlm.nih.gov/books/NBK57243/>), *Neural Control of Renal Function*, Morgan & Claypool Life Sciences, retrieved 2026-06-20
8. Rose GA (1969). "Measurement of glomerular filtration rate by inulin clearance without urine collection" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1982852>). *BMJ*. **2** (5649): 91–3. doi:10.1136/bmj.2.5649.91 (<https://doi.org/10.1136/bmj.2.5649.91>). PMC 1982852 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1982852>). PMID 5775456 (<https://pubmed.ncbi.nlm.nih.gov/5775456>).
9. Hsu CY, Bansal N (August 2011). "Measured GFR as "gold standard"--all that glitters is not gold?" (<https://doi.org/10.2215/cjn.06040611>). *Clinical Journal of the American Society of Nephrology*. **6** (8): 1813–4. doi:10.2215/cjn.06040611 (<https://doi.org/10.2215/cjn.06040611>). PMID 21784836 (<https://pubmed.ncbi.nlm.nih.gov/21784836>).
10. Bui TV, Prot-Bertoye C, Ayari H, Baron S, Bertocchio JP, Bureau C, Davis P, Blanchard A, Houillier P, Prie D, Lillo-Le Louet A, Courbebaisse M (2021). "Safety of Inulin and Sinistrin: Combining Several Sources for Pharmacovigilance Purposes" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8637630>). *Front Pharmacol*. **12** 725417. doi:10.3389/fphar.2021.725417 (<https://doi.org/10.3389/fphar.2021.725417>). PMC 8637630 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8637630>). PMID 34867328 (<https://pubmed.ncbi.nlm.nih.gov/34867328>).

11. Soveri I, Berg UB, Björk J, Elinder CG, Grubb A, Mejare I, Sterner G, Bäck SE (September 2014). "Measuring GFR: a systematic review". *Am J Kidney Dis.* **64** (3): 411–24. doi:10.1053/j.ajkd.2014.04.010 (<https://doi.org/10.1053%2Fj.ajkd.2014.04.010>). PMID 24840668 (<https://pubmed.ncbi.nlm.nih.gov/24840668>).
12. Murray AW, Barnfield MC, Waller ML, Telford T, Peters AM (8 May 2013). "Assessment of Glomerular Filtration Rate Measurement with Plasma Sampling: A Technical Review" (<https://doi.org/10.2967%2Fjnmt.113.121004>). *Journal of Nuclear Medicine Technology.* **41** (2): 67–75. doi:10.2967/jnmt.113.121004 (<https://doi.org/10.2967%2Fjnmt.113.121004>). PMID 23658207 (<https://pubmed.ncbi.nlm.nih.gov/23658207>).
13. Speeckaert M, Delanghe J (2015). "Assessment of renal function". In Giuseppe D, Winearls C, Remuzzi G (eds.). *Oxford Textbook of Clinical Nephrology* (4th ed.). Oxford: Oxford University Press. p. 44. ISBN 978-0-19-959254-8.
14. Henriksen UL, Henriksen JH (January 2015). "The clearance concept with special reference to determination of glomerular filtration rate in patients with fluid retention". *Clinical Physiology and Functional Imaging.* **35** (1): 7–16. doi:10.1111/cpf.12149 (<https://doi.org/10.1111%2Fcpf.12149>). PMID 24750696 (<https://pubmed.ncbi.nlm.nih.gov/24750696>). S2CID 44756080 (<https://api.semanticscholar.org/CorpusID:44756080>).
15. Soveri I, Berg UB, Björk J, Elinder CG, Grubb A, Mejare I, Sterner G, Bäck SE (September 2014). "Measuring GFR: A Systematic Review". *American Journal of Kidney Diseases.* **64** (3): 411–424. doi:10.1053/j.ajkd.2014.04.010 (<https://doi.org/10.1053%2Fj.ajkd.2014.04.010>). PMID 24840668 (<https://pubmed.ncbi.nlm.nih.gov/24840668>).
16. "Chronic kidney disease in adults: assessment and management" (<https://www.nice.org.uk/guidance/cg182/chapter/1-recommendations#when-highly-accurate-measures-of-gfr-are-required>). NICE. 23 July 2014. Retrieved 19 August 2016.
17. Schück O, Teplan V, Jabor A, Stollova M, Skibova J (2003). "Glomerular filtration rate estimation in patients with advanced chronic renal insufficiency based on serum cystatin C levels". *Nephron Clin Pract.* **93** (4): c146–51. doi:10.1159/000070234 (<https://doi.org/10.1159%2F000070234>). PMID 12759583 (<https://pubmed.ncbi.nlm.nih.gov/12759583>).
18. Stevens LA, Coresh J, Schmid CH, Feldman HI, Froissart M, Kusek J, Rossert J, Van Lente F, Bruce RD (March 2008). "Estimating GFR using serum cystatin C alone and in combination with serum creatinine: a pooled analysis of 3,418 individuals with CKD" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2390827>). *American Journal of Kidney Diseases.* **51** (3): 395–406. doi:10.1053/j.ajkd.2007.11.018 (<https://doi.org/10.1053%2Fj.ajkd.2007.11.018>). PMC 2390827 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2390827>). PMID 18295055 (<https://pubmed.ncbi.nlm.nih.gov/18295055>).
19. Delgado C, Baweja M, Crews DC, Eneanya ND, Gadegbeku CA, Inker LA, Mendu ML, Miller WG, Moxey-Mims MM, Roberts GV, St Peter WL, Warfield C, Powe NR (February 2022). "A Unifying Approach for GFR Estimation: Recommendations of the NKF-ASN Task Force on Reassessing the Inclusion of Race in Diagnosing Kidney Disease". *Am J Kidney Dis.* **79** (2): 268–288.e1. doi:10.1053/j.ajkd.2021.08.003 (<https://doi.org/10.1053%2Fj.ajkd.2021.08.003>). PMID 34563581 (<https://pubmed.ncbi.nlm.nih.gov/34563581>).
20. Guyton A, Hall J (2006). "26: Urine Formation by the Kidneys: I. Glomerular Filtration, Renal Blood Flow, and Their Control". In Grulow R (ed.). *Textbook of Medical Physiology* (11th ed.). Elsevier. pp. 308–325. ISBN 978-0-7216-0240-0.
21. Keener J, Sneyd J (2004). "20: Renal Physiology" (https://link.springer.com/chapter/10.1007/0-387-22706-7_20). In Marsden J (ed.). *Mathematical Physiology* (Book). Interdisciplinary Mathematics. Vol. 8 (Mathematical Biology). Sirovich, Wiggins (1st ed.). Springer. pp. 612–636. doi:10.1007/0-387-22706-7_20 (https://doi.org/10.1007%2F0-387-22706-7_20). ISBN 978-0-387-98381-3.

22. Stehlé T, Delanaye P (October 2024). "Which is the best glomerular filtration marker: Creatinine, cystatin C or both?" (<https://doi.org/10.1111%2Feci.14278>). *Eur J Clin Invest.* **54** (10) e14278. doi:10.1111/eci.14278 (<https://doi.org/10.1111%2Feci.14278>). PMID 38949475 (<https://pubmed.ncbi.nlm.nih.gov/38949475>).
23. "KDOQI CKD Guidelines" (https://web.archive.org/web/20121003072347/http://www.kidney.org/g/professionals/kdoqi/guidelines_ckd/p5_lab_g4.htm). Archived from the original (http://www.kidney.org/professionals/kdoqi/guidelines_ckd/p5_lab_g4.htm) on 2012-10-03. Retrieved 2010-08-25.
24. "GFR Calculator at [cato.at](http://www.cato.at) – Cockcroft-Gault" (https://web.archive.org/web/20040905112600/http://www.cato.at/web/service/servlet/location?goto=SERVICE_CALC&lang=EN&URL=%22calc%2Fcato_GFR_CG.htm%22). Archived from the original (http://www.cato.at/web/service/servlet/location?goto=SERVICE_CALC&lang=EN&URL=%22calc/cato_GFR_CG.htm%22) on 2004-09-05. – GFR calculation (Cockcroft-Gault formula)
25. Cockcroft DW, Gault MH (1976). "Prediction of creatinine clearance from serum creatinine". *Nephron.* **16** (1): 31–41. doi:10.1159/000180580 (<https://doi.org/10.1159%2F000180580>). PMID 1244564 (<https://pubmed.ncbi.nlm.nih.gov/1244564>).
26. Gault MH, Longerich LL, Harnett JD, Wesolowski C (1992). "Predicting glomerular function from adjusted serum creatinine". *Nephron.* **62** (3): 249–56. doi:10.1159/000187054 (<https://doi.org/10.1159%2F000187054>). PMID 1436333 (<https://pubmed.ncbi.nlm.nih.gov/1436333>).
27. Levey AS, Bosch JP, Lewis JB, Greene T, Rogers N, Roth D (March 1999). "A more accurate method to estimate glomerular filtration rate from serum creatinine: a new prediction equation. Modification of Diet in Renal Disease Study Group". *Annals of Internal Medicine.* **130** (6): 461–70. doi:10.7326/0003-4819-130-6-199903160-00002 (<https://doi.org/10.7326%2F0003-4819-130-6-199903160-00002>). PMID 10075613 (<https://pubmed.ncbi.nlm.nih.gov/10075613>). S2CID 1902375 (<https://api.semanticscholar.org/CorpusID:1902375>).
28. Mathew TH, Johnson DW, Jones GR (October 2007). "Chronic kidney disease and automatic reporting of estimated glomerular filtration rate: revised recommendations" (http://www.mja.com.au/public/issues/187_08_15/10/07/mat10525_fm.html). *The Medical Journal of Australia.* **187** (8): 459–63. doi:10.5694/j.1326-5377.2007.tb01357.x (<https://doi.org/10.5694%2Fj.1326-5377.2007.tb01357.x>). PMID 17937643 (<https://pubmed.ncbi.nlm.nih.gov/17937643>). S2CID 14920030 (<https://api.semanticscholar.org/CorpusID:14920030>).
29. Joint Specialty Committee on Renal Disease (June 2005). "Chronic kidney disease in adults: UK guidelines for identification, management and referral" (<https://web.archive.org/web/20071019043929/http://www.renal.org/CKDguide/full/UKCKDfull.pdf>) (PDF). Archived from the original (<http://www.renal.org/CKDguide/full/UKCKDfull.pdf>) (PDF) on 2007-10-19.
30. www.nice.org.uk (July 2014). "Chronic kidney disease in adults: assessment and management" (<https://www.nice.org.uk/guidance/cg182/chapter/1-Recommendations#investigations-for-chronic-kidney-disease-2>).
31. Davey RX (January 2006). "Chronic kidney disease and automatic reporting of estimated glomerular filtration rate" (http://www.mja.com.au/public/issues/184_01_02/01/06/matters_arising_020106_fm-6.html). *The Medical Journal of Australia.* **184** (1): 42–3, author reply 43. doi:10.5694/j.1326-5377.2006.tb00098.x (<https://doi.org/10.5694%2Fj.1326-5377.2006.tb00098.x>). hdl:2440/34660 (<https://hdl.handle.net/2440%2F34660>). PMID 16398632 (<https://pubmed.ncbi.nlm.nih.gov/16398632>). S2CID 9648508 (<https://api.semanticscholar.org/CorpusID:9648508>).
32. Twomey PJ, Reynolds TM (November 2006). "The MDRD formula and validation" (<https://doi.org/10.1093%2Fqjmed%2Fhcl108>). *QJM.* **99** (11): 804–5. doi:10.1093/qjmed/hcl108 (<https://doi.org/10.1093%2Fqjmed%2Fhcl108>). PMID 17041249 (<https://pubmed.ncbi.nlm.nih.gov/17041249>).

33. Kallner A, Ayling PA, Khatami Z (2008). "Does eGFR improve the diagnostic capability of S-Creatinine concentration results? A retrospective population based study" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2204044>). *International Journal of Medical Sciences*. **5** (1): 9–17. doi:10.7150/ijms.5.9 (<https://doi.org/10.7150%2Fijms.5.9>). PMC 2204044 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2204044>). PMID 18219370 (<https://pubmed.ncbi.nlm.nih.gov/18219370>). S2CID 14970724 (<https://api.semanticscholar.org/CorpusID:14970724>).
34. National Kidney Foundation (February 2002). "K/DOQI clinical practice guidelines for chronic kidney disease: evaluation, classification, and stratification". *American Journal of Kidney Diseases*. **39** (2 Suppl 1): S1–266. doi:10.1016/S0272-6386(02)70081-4 (<https://doi.org/10.1016%2FS0272-6386%2802%2970081-4>). PMID 11904577 (<https://pubmed.ncbi.nlm.nih.gov/11904577>).
35. Rule AD, Larson TS, Bergstralh EJ, Slezak JM, Jacobsen SJ, Cosio FG (December 2004). "Using serum creatinine to estimate glomerular filtration rate: accuracy in good health and in chronic kidney disease". *Annals of Internal Medicine*. **141** (12): 929–37. doi:10.7326/0003-4819-141-12-200412210-00009 (<https://doi.org/10.7326%2F0003-4819-141-12-200412210-00009>). PMID 15611490 (<https://pubmed.ncbi.nlm.nih.gov/15611490>). S2CID 30342139 (<https://api.semanticscholar.org/CorpusID:30342139>).
36. Levey AS, Coresh J, Greene T, Stevens LA, Zhang YL, Hendriksen S, Kusek JW, Van Lente F, Chronic Kidney Disease Epidemiology Collaboration (August 2006). "Using standardized serum creatinine values in the modification of diet in renal disease study equation for estimating glomerular filtration rate". *Annals of Internal Medicine*. **145** (4): 247–54. doi:10.7326/0003-4819-145-4-200608150-00004 (<https://doi.org/10.7326%2F0003-4819-145-4-200608150-00004>). PMID 16908915 (<https://pubmed.ncbi.nlm.nih.gov/16908915>). S2CID 37149831 (<https://api.semanticscholar.org/CorpusID:37149831>).
37. "GFR MDRD Calculator for Adults" (<https://www.niddk.nih.gov/health-information/community-health-outreach/information-clearinghouses/nkdep?dkrd=lgdmw0002>). *National Kidney Disease Education Program*. United States: National Institutes of Health. Archived (https://web.archive.org/web/20120226204408/http://nkdep.nih.gov/professionals/gfr_calculators/idms_con.htm) from the original on 2012-02-26. Retrieved 2009-11-16.
38. Based on formulas from http://ovid.com/jnls/urol/fulltext/10.4103/ua.ua_7_24~race-modified-estimated-glomerular-filtration-rate
39. Hougardy JM, Delanaye P, Le Moine A, Nortier J (2014). "Estimation of the glomerular filtration rate in 2014 by tests and equations: strengths and weaknesses". *Rev Med Brux* (in French). **35** (4): 250–7. PMID 25675627 (<https://pubmed.ncbi.nlm.nih.gov/25675627>).
40. Levey AS, Stevens LA, Schmid CH, Zhang YL, Castro A, Feldman HI, Kusek JW, Eggers P, Van Lente F (May 2009). "A new equation to estimate glomerular filtration rate" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2763564>). *Annals of Internal Medicine*. **150** (9): 604–12. doi:10.7326/0003-4819-150-9-200905050-00006 (<https://doi.org/10.7326%2F0003-4819-150-9-200905050-00006>). PMC 2763564 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2763564>). PMID 19414839 (<https://pubmed.ncbi.nlm.nih.gov/19414839>).
41. Matsushita K, Selvin E, Bash LD, Astor BC, Coresh J (April 2010). "Risk implications of the new CKD Epidemiology Collaboration (CKD-EPI) equation compared with the MDRD Study equation for estimated GFR: the Atherosclerosis Risk in Communities (ARIC) Study" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2858455>). *American Journal of Kidney Diseases*. **55** (4): 648–59. doi:10.1053/j.ajkd.2009.12.016 (<https://doi.org/10.1053%2Fj.ajkd.2009.12.016>). PMC 2858455 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2858455>). PMID 20189275 (<https://pubmed.ncbi.nlm.nih.gov/20189275>).

42. Inker LA, Eneanya ND, Coresh J, Tighiouart H, Wang D, Sang Y, Crews DC, Doria A, Estrella MM, Froissart M, Grams ME, Greene T, Grubb A, Gudnason V, Gutiérrez OM, Kalil R, Karger AB, Mauer M, Navis G, Nelson RG, Poggio ED, Rodby R, Rossing P, Rule AD, Selvin E, Seegmiller JC, Shlipak MG, Torres VE, Yang W, Ballew SH, Couture SJ, Powe NR, Levey AS (November 2021). "New Creatinine- and Cystatin C-Based Equations to Estimate GFR without Race" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8822996>). *N Engl J Med.* **385** (19): 1737–49. doi:10.1056/NEJMoa2102953 (<https://doi.org/10.1056%2FNEJMoa2102953>). PMC 8822996 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8822996>). PMID 34554658 (<https://pubmed.ncbi.nlm.nih.gov/34554658>).
43. Saleem M, Florkowski CM, George PM (2008). "Comparison of the Mayo Clinic Quadratic Equation with the Modification of Diet in Renal Disease equation and radionuclide glomerular filtration rate in a clinical setting". *Nephrology.* **13** (8): 684–8. doi:10.1111/j.1440-1797.2008.01045.x (<https://doi.org/10.1111%2Fj.1440-1797.2008.01045.x>). ISSN 1320-5358 (<https://search.worldcat.org/issn/1320-5358>). PMID 19154321 (<https://pubmed.ncbi.nlm.nih.gov/19154321>). S2CID 45943783 (<https://api.semanticscholar.org/CorpusID:45943783>).
44. Fontserè N, Bonal J, Salinas I, de Arellano MR, Rios J, Torres F, Sanmarti A, Romero R (2008). "Is the New Mayo Clinic Quadratic Equation Useful for the Estimation of Glomerular Filtration Rate in Type 2 Diabetic Patients?" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2584175>). *Diabetes Care.* **31** (12): 2265–7. doi:10.2337/dc08-0958 (<https://doi.org/10.2337%2Fdc08-0958>). PMC 2584175 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2584175>). PMID 18835955 (<https://pubmed.ncbi.nlm.nih.gov/18835955>). S2CID 24211196 (<https://api.semanticscholar.org/CorpusID:24211196>).
45. Schwartz GJ, Haycock GB, Edelmann CM, Spitzer A (August 1976). "A simple estimate of glomerular filtration rate in children derived from body length and plasma creatinine". *Pediatrics.* **58** (2): 259–63. doi:10.1542/peds.58.2.259 (<https://doi.org/10.1542%2Fpeds.58.2.259>). PMID 951142 (<https://pubmed.ncbi.nlm.nih.gov/951142>). S2CID 38437700 (<https://api.semanticscholar.org/CorpusID:38437700>).
46. Schwartz GJ, Feld LG, Langford DJ (June 1984). "A simple estimate of glomerular filtration rate in full-term infants during the first year of life". *The Journal of Pediatrics.* **104** (6): 849–54. doi:10.1016/S0022-3476(84)80479-5 (<https://doi.org/10.1016%2FS0022-3476%2884%2980479-5>). PMID 6726515 (<https://pubmed.ncbi.nlm.nih.gov/6726515>).
47. Brion LP, Fleischman AR, McCarton C, Schwartz GJ (October 1986). "A simple estimate of glomerular filtration rate in low birth weight infants during the first year of life: noninvasive assessment of body composition and growth". *The Journal of Pediatrics.* **109** (4): 698–707. doi:10.1016/S0022-3476(86)80245-1 (<https://doi.org/10.1016%2FS0022-3476%2886%2980245-1>). PMID 3761090 (<https://pubmed.ncbi.nlm.nih.gov/3761090>).
48. Haenggi MH, Pelet J, Guignard JP (February 1999). "Estimation of glomerular filtration rate by the formula $GFR = K \times T/Pc$ ". *Archives de Pédiatrie* (in French). **6** (2): 165–72. doi:10.1016/S0929-693X(99)80204-8 (<https://doi.org/10.1016%2FS0929-693X%2899%2980204-8>). PMID 10079885 (<https://pubmed.ncbi.nlm.nih.gov/10079885>).
49. Schwartz GJ, Muñoz A, Schneider MF, Mak RH, Kaskel F, Warady BA, Furth SL (March 2009). "New equations to estimate GFR in children with CKD" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2653687>). *Journal of the American Society of Nephrology.* **20** (3): 629–37. doi:10.1681/ASN.2008030287 (<https://doi.org/10.1681%2FASN.2008030287>). PMC 2653687 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2653687>). PMID 19158356 (<https://pubmed.ncbi.nlm.nih.gov/19158356>).
50. Bauer C, Melamed ML, Hostetter TH (2008). "Staging of Chronic Kidney Disease: Time for a Course Correction" (<http://jasn.asnjournals.org/content/19/5/844.full>). *Journal of the American Society of Nephrology.* **19** (5): 844–46. doi:10.1681/ASN.2008010110 (<https://doi.org/10.1681%2FASN.2008010110>). PMID 18385419 (<https://pubmed.ncbi.nlm.nih.gov/18385419>).

51. Eckardt KU, Berns JS, Rocco MV, Kasiske BL (June 2009). "Definition and Classification of CKD: The Debate Should Be About Patient Prognosis—A Position Statement From KDOQI and KDIGO" (https://web.archive.org/web/20110725131557/http://www.kdigo.org/meetings_events/pdf/KDOQI-KDIGO_Editorial_on_CKD_Classification.pdf) (PDF). *American Journal of Kidney Diseases*. **53** (6): 915–920. doi:10.1053/j.ajkd.2009.04.001 (<https://doi.org/10.1053%2Fj.ajkd.2009.04.001>). PMID 19406541 (<https://pubmed.ncbi.nlm.nih.gov/19406541>). Archived from the original (http://www.kdigo.org/meetings_events/pdf/KDOQI-KDIGO_Editorial_on_CKD_Classification.pdf) (PDF) on 2011-07-25.
52. "KDIGO Controversies Conference: Definition, Classification and Prognosis in CKD, London, October 2009" (https://web.archive.org/web/20101124202928/http://www.kdigo.org/meetings_events/CKD_Controversies_Conference.php). Kidney Disease: Improving Global Outcomes (KDIGO). 2009. Archived from the original (http://www.kdigo.org/meetings_events/CKD_Controversies_Conference.php) on 2010-11-24.
53. Hoenig MP, Mann A, Pavlakis M (February 2022). "Removal of the Black race coefficient from the estimated glomerular filtration equation improves transplant eligibility for Black patients at a single center". *Clinical Transplantation*. **36** (2) e14467. doi:10.1111/ctr.14467 (<https://doi.org/10.1111%2Fctr.14467>). PMID 34605076 (<https://pubmed.ncbi.nlm.nih.gov/34605076>). S2CID 238256526 (<https://api.semanticscholar.org/CorpusID:238256526>).
54. Vyas DA, Eisenstein LG, Jones DS (27 August 2020). "Hidden in Plain Sight — Reconsidering the Use of Race Correction in Clinical Algorithms" (<https://doi.org/10.1056%2FNEJMms2004740>). *New England Journal of Medicine*. **383** (9): 874–882. doi:10.1056/NEJMms2004740 (<https://doi.org/10.1056%2FNEJMms2004740>). PMID 32853499 (<https://pubmed.ncbi.nlm.nih.gov/32853499>).
55. "OPTN Board approves elimination of race-based calculation for transplant candidate listing - OPTN" (<https://optn.transplant.hrsa.gov/news/optn-board-approves-elimination-of-race-based-calculation-for-transplant-candidate-listing/>). *optn.transplant.hrsa.gov*. UNOS News Bureau. Retrieved 12 December 2023.

External links

Reference links

- National Kidney Disease Education Program website. (<https://web.archive.org/web/20120224110305/http://www.nkdep.nih.gov/professionals/index.htm>) Includes professional references and GFR calculators
- eGFR at Lab Tests Online (<https://www.testing.com/tests/estimated-glomerular-filtration-rate-egfr/>)

Retrieved from "https://en.wikipedia.org/w/index.php?title=Glomerular_filtration_rate&oldid=1372412934"