

# Design and Analysis of Algorithms

---

CSE 5311

Lecture 2 Asymptotic Notation and Solving Recurrences

Song Jiang, Ph.D.

Department of Computer Science and Engineering



# Asymptotic notation

$O$ -notation (upper bounds):

We write  $f(n) = O(g(n))$  if there exist constants  $c > 0$ ,  $n_0 > 0$  such that  $0 \leq f(n) \leq cg(n)$  for all  $n \geq n_0$ .



# Asymptotic notation

$O$ -notation (upper bounds):

We write  $f(n) = O(g(n))$  if there exist constants  $c > 0$ ,  $n_0 > 0$  such that  $0 \leq f(n) \leq cg(n)$  for all  $n \geq n_0$ .

**EXAMPLE:**  $2n^2 = O(n^3)$  ( $c = 1$ ,  $n_0 = 2$ )



# Asymptotic notation

$O$ -notation (upper bounds):

We write  $f(n) = O(g(n))$  if there exist constants  $c > 0$ ,  $n_0 > 0$  such that  $0 \leq f(n) \leq cg(n)$  for all  $n \geq n_0$ .

**EXAMPLE:**  $2n^2 = O(n^3)$  ( $c = 1$ ,  $n_0 = 2$ )

*functions,  
not values*



# Asymptotic notation


$O$ -notation (upper bounds):

We write  $f(n) = O(g(n))$  if there exist constants  $c > 0$ ,  $n_0 > 0$  such that  $0 \leq f(n) \leq cg(n)$  for all  $n \geq n_0$ .

**EXAMPLE:**  $2n^2 = O(n^3)$  ( $c = 1$ ,  $n_0 = 2$ )

*functions,  
not values*

*funny, “one-way”  
equality*



# Set definition of O-notation

$O(g(n)) = \{ f(n) : \text{there exist constants } c > 0, n_0 > 0 \text{ such that } 0 \leq f(n) \leq cg(n) \text{ for all } n \geq n_0 \}$



# Set definition of O-notation

$O(g(n)) = \{ f(n) : \text{there exist constants } c > 0, n_0 > 0 \text{ such that } 0 \leq f(n) \leq cg(n) \text{ for all } n \geq n_0 \}$

**EXAMPLE:**  $2n^2 \in O(n^3)$



# Macro substitution

***Convention:*** A set in a formula represents an anonymous function in the set.





# Macro substitution

***Convention:*** A set in a formula represents an anonymous function in the set.

**EXAMPLE:**  $f(n) = n^3 + O(n^2)$

means

$$f(n) = n^3 + h(n)$$

for some  $h(n) \in O(n^2)$  .



# Macro substitution

**Convention:** A set in a formula represents an anonymous function in the set.

**EXAMPLE:**  $n^2 + O(n) = O(n^2)$

means

for any  $f(n) \in O(n)$ :


$$n^2 + f(n) = h(n)$$

for some  $h(n) \in O(n^2)$  .



# $\Omega$ -notation (lower bounds)

$O$ -notation is an *upper-bound* notation. It makes no sense to say  $f(n)$  is at least  $O(n^2)$ .



# $\Omega$ -notation (lower bounds)

$O$ -notation is an *upper-bound* notation. It makes no sense to say  $f(n)$  is at least  $O(n^2)$ .

$$\Omega(g(n)) = \{ f(n) : \text{there exist constants } c > 0, n_0 > 0 \text{ such that } 0 \leq cg(n) \leq f(n) \text{ for all } n \geq n_0 \}$$



# $\Omega$ -notation (lower bounds)

$O$ -notation is an *upper-bound* notation. It makes no sense to say  $f(n)$  is at least  $O(n^2)$ .

$$\Omega(g(n)) = \{ f(n) : \text{there exist constants } c > 0, n_0 > 0 \text{ such that } 0 \leq cg(n) \leq f(n) \text{ for all } n \geq n_0 \}$$

**EXAMPLE:**  $\sqrt{n} = \Omega(\lg n)$  ( $c = 1, n_0 = 16$ )



# $\Theta$ -notation (tight bounds)

$$\Theta(g(n)) = O(g(n)) \cap \Omega(g(n))$$



# $\Theta$ -notation (tight bounds)

$$\Theta(g(n)) = O(g(n)) \cap \Omega(g(n))$$

**EXAMPLE:**  $\frac{1}{2}n^2 - 2n = \Theta(n^2)$



# $o$ -notation and $\omega$ -notation

$O$ -notation and  $\Omega$ -notation are like  $\leq$  and  $\geq$ .  
 $o$ -notation and  $\omega$ -notation are like  $<$  and  $>$ .

$o(g(n)) = \{ f(n) : \text{for any constant } c > 0, \\ \text{there is a constant } n_0 > 0 \\ \text{such that } 0 \leq f(n) < cg(n) \\ \text{for all } n \geq n_0 \}$

**EXAMPLE:**  $2n^2 = o(n^3)$  ( $n_0 = 2/c$ )





# **$O$ -notation and $\omega$ -notation**

$O$ -notation and  $\Omega$ -notation are like  $\leq$  and  $\geq$ .  
 $o$ -notation and  $\omega$ -notation are like  $<$  and  $>$ .

$\omega(g(n)) = \{ f(n) : \text{for any constant } c > 0, \\ \text{there is a constant } n_0 > 0 \\ \text{such that } 0 \leq cg(n) < f(n) \\ \text{for all } n \geq n_0 \}$



# Solving recurrences

- The analysis of merge sort from *Lecture 1* required us to solve a recurrence.
- Recurrences are like solving integrals, differential equations, etc.
  - Learn a few tricks.
- *Lecture 3*: Applications of recurrences to divide-and-conquer algorithms.



# Substitution method

*The most general method:*

- 1. *Guess*** the form of the solution.
- 2. *Verify*** by induction.
- 3. *Solve*** for constants.



# Substitution method

*The most general method:*

- 1. *Guess*** the form of the solution.
- 2. *Verify*** by induction.
- 3. *Solve*** for constants.

**EXAMPLE:**  $T(n) = 4T(n/2) + n$


- [Assume that  $T(1) = \Theta(1)$ .]
- Guess  $O(n^3)$ . (Prove  $O$  and  $\Omega$  separately.)
- Assume that  $T(k) \leq ck^3$  for  $k < n$ .
- Prove  $T(n) \leq cn^3$  by induction.



# Example of substitution

$$\begin{aligned}T(n) &= 4T(n/2) + n \\&\leq 4c(n/2)^3 + n \\&= (c/2)n^3 + n \\&= cn^3 - ((c/2)n^3 - n) \leftarrow \textit{desired} - \textit{residual} \\&\leq cn^3 \leftarrow \textit{desired}\end{aligned}$$

whenever  $(c/2)n^3 - n \geq 0$ , for example,  
if  $c \geq 2$  and  $n \geq 1$ .  
 $\nwarrow$   
*residual*



# Example (continued)

- We must also handle the initial conditions, that is, ground the induction with base cases.
- **Base:**  $T(n) = \Theta(1)$  for all  $n < n_0$ , where  $n_0$  is a suitable constant.
- For  $1 \leq n < n_0$ , we have “ $\Theta(1)$ ”  $\leq cn^3$ , if we pick  $c$  big enough.



# Example (continued)

- We must also handle the initial conditions, that is, ground the induction with base cases.
- **Base:**  $T(n) = \Theta(1)$  for all  $n < n_0$ , where  $n_0$  is a suitable constant.
- For  $1 \leq n < n_0$ , we have “ $\Theta(1)$ ”  $\leq cn^3$ , if we pick  $c$  big enough.

---

---

***This bound is not tight!***



# A tighter upper bound?

We shall prove that  $T(n) = O(n^2)$ .





# A tighter upper bound?

We shall prove that  $T(n) = O(n^2)$ .

Assume that  $T(k) \leq ck^2$  for  $k < n$ :

$$\begin{aligned} T(n) &= 4T(n/2) + n \\ &\leq 4c(n/2)^2 + n \\ &= cn^2 + n \\ &= O(n^2) \end{aligned}$$



# A tighter upper bound?

We shall prove that  $T(n) = O(n^2)$ .

Assume that  $T(k) \leq ck^2$  for  $k < n$ :

$$\begin{aligned} T(n) &= 4T(n/2) + n \\ &\leq 4c(n/2)^2 + n \\ &= cn^2 + n \end{aligned}$$

~~$= O(n^2)$~~  **Wrong!** We must prove the I.H.



# A tighter upper bound?

We shall prove that  $T(n) = O(n^2)$ .

Assume that  $T(k) \leq ck^2$  for  $k < n$ :

$$\begin{aligned} T(n) &= 4T(n/2) + n \\ &\leq 4c(n/2)^2 + n \\ &= cn^2 + n \end{aligned}$$

~~$= O(n^2)$~~  **Wrong!** We must prove the I.H.

$$= cn^2 - (-n) \quad [\text{desired} - \text{residual}]$$

$$\leq cn^2 \quad \text{for } \textit{no} \text{ choice of } c > 0. \text{ Lose!}$$



# A tighter upper bound!

**IDEA:** Strengthen the inductive hypothesis.

- *Subtract* a low-order term.

*Inductive hypothesis:*  $T(k) \leq c_1 k^2 - c_2 k$  for  $k < n$ .



# A tighter upper bound!

**IDEA:** Strengthen the inductive hypothesis.

- *Subtract* a low-order term.

*Inductive hypothesis:*  $T(k) \leq c_1 k^2 - c_2 k$  for  $k < n$ .

$$\begin{aligned} T(n) &= 4T(n/2) + n \\ &= 4(c_1(n/2)^2 - c_2(n/2)) + n \\ &= c_1 n^2 - 2c_2 n + n \\ &= c_1 n^2 - c_2 n - (c_2 n - n) \\ &\leq c_1 n^2 - c_2 n \quad \text{if } c_2 \geq 1. \end{aligned}$$



# A tighter upper bound!

**IDEA:** Strengthen the inductive hypothesis.

- *Subtract* a low-order term.

*Inductive hypothesis:*  $T(k) \leq c_1 k^2 - c_2 k$  for  $k < n$ .


$$\begin{aligned} T(n) &= 4T(n/2) + n \\ &= 4(c_1(n/2)^2 - c_2(n/2)) + n \\ &= c_1 n^2 - 2c_2 n + n \\ &= c_1 n^2 - c_2 n - (c_2 n - n) \\ &\leq c_1 n^2 - c_2 n \quad \text{if } c_2 \geq 1. \end{aligned}$$

Pick  $c_1$  big enough to handle the initial conditions.



# Recursion-tree method


- A recursion tree models the costs (time) of a recursive execution of an algorithm.
- The recursion-tree method can be unreliable, just like any method that uses ellipses (...).
- The recursion-tree method promotes intuition, however.
- The recursion tree method is good for generating guesses for the substitution method.



# Example of recursion tree

Solve  $T(n) = T(n/4) + T(n/2) + n^2$ :






# Example of recursion tree


Solve  $T(n) = T(n/4) + T(n/2) + n^2$ :

$$T(n)$$



# Example of recursion tree


Solve  $T(n) = T(n/4) + T(n/2) + n^2$ :






# Example of recursion tree


Solve  $T(n) = T(n/4) + T(n/2) + n^2$ :






# Example of recursion tree


Solve  $T(n) = T(n/4) + T(n/2) + n^2$ :






# Example of recursion tree


Solve  $T(n) = T(n/4) + T(n/2) + n^2$ :






# Example of recursion tree


Solve  $T(n) = T(n/4) + T(n/2) + n^2$ :





# Example of recursion tree


Solve  $T(n) = T(n/4) + T(n/2) + n^2$ :





# Example of recursion tree

Solve  $T(n) = T(n/4) + T(n/2) + n^2$ :







# The master method

The master method applies to recurrences of the form

$$T(n) = a T(n/b) + f(n) ,$$

where  $a \geq 1$ ,  $b > 1$ , and  $f$  is asymptotically positive.




# Three common cases

Compare  $f(n)$  with  $n^{\log_b a}$ :

1.  $f(n) = O(n^{\log_b a - \varepsilon})$  for some constant  $\varepsilon > 0$ .

- $f(n)$  grows polynomially slower than  $n^{\log_b a}$  (by an  $n^\varepsilon$  factor).

***Solution:***  $T(n) = \Theta(n^{\log_b a})$ .



# Three common cases

Compare  $f(n)$  with  $n^{\log_b a}$ :

1.  $f(n) = O(n^{\log_b a - \varepsilon})$  for some constant  $\varepsilon > 0$ .

- $f(n)$  grows polynomially slower than  $n^{\log_b a}$  (by an  $n^\varepsilon$  factor).

**Solution:**  $T(n) = \Theta(n^{\log_b a})$ .

2.  $f(n) = \Theta(n^{\log_b a} \lg^k n)$  for some constant  $k \geq 0$ .

- $f(n)$  and  $n^{\log_b a}$  grow at similar rates.

**Solution:**  $T(n) = \Theta(n^{\log_b a} \lg^{k+1} n)$ .



# Three common cases (cont.)

Compare  $f(n)$  with  $n^{\log_b a}$ :

3.  $f(n) = \Omega(n^{\log_b a + \varepsilon})$  for some constant  $\varepsilon > 0$ .

- $f(n)$  grows polynomially faster than  $n^{\log_b a}$  (by an  $n^\varepsilon$  factor),

*and*  $f(n)$  satisfies the **regularity condition** that  $a f(n/b) \leq c f(n)$  for some constant  $c < 1$ .

**Solution:**  $T(n) = \Theta(f(n))$ .



# Examples

**Ex.**  $T(n) = 4T(n/2) + n$   
 $a = 4, b = 2 \Rightarrow n^{\log_b a} = n^2; f(n) = n.$   
**CASE 1:**  $f(n) = O(n^{2-\varepsilon})$  for  $\varepsilon = 1.$   
 $\therefore T(n) = \Theta(n^2).$



# Examples

**Ex.**  $T(n) = 4T(n/2) + n$   
 $a = 4, b = 2 \Rightarrow n^{\log_b a} = n^2; f(n) = n.$   
**CASE 1:**  $f(n) = O(n^{2-\varepsilon})$  for  $\varepsilon = 1.$   
 $\therefore T(n) = \Theta(n^2).$

**Ex.**  $T(n) = 4T(n/2) + n^2$   
 $a = 4, b = 2 \Rightarrow n^{\log_b a} = n^2; f(n) = n^2.$   
**CASE 2:**  $f(n) = \Theta(n^2 \lg^0 n)$ , that is,  $k = 0.$   
 $\therefore T(n) = \Theta(n^2 \lg n).$



# Examples

**Ex.**  $T(n) = 4T(n/2) + n^3$

$$a = 4, b = 2 \Rightarrow n^{\log_b a} = n^2; f(n) = n^3.$$

**CASE 3:**  $f(n) = \Omega(n^{2 + \epsilon})$  for  $\epsilon = 1$

*and*  $4(n/2)^3 \leq cn^3$  (reg. cond.) for  $c = 1/2$ .

$$\therefore T(n) = \Theta(n^3).$$



# Examples

**Ex.**  $T(n) = 4T(n/2) + n^3$

$$a = 4, b = 2 \Rightarrow n^{\log_b a} = n^2; f(n) = n^3.$$


**CASE 3:**  $f(n) = \Omega(n^{2+\epsilon})$  for  $\epsilon = 1$   
*and*  $4(n/2)^3 \leq cn^3$  (reg. cond.) for  $c = 1/2$ .  
 $\therefore T(n) = \Theta(n^3)$ .

**Ex.**  $T(n) = 4T(n/2) + n^2/\lg n$

$$a = 4, b = 2 \Rightarrow n^{\log_b a} = n^2; f(n) = n^2/\lg n.$$


Master method does not apply. In particular,  
for every constant  $\epsilon > 0$ , we have  $n^\epsilon = \omega(\lg n)$ .






# Idea of master theorem


*Recursion tree:*






# Idea of master theorem


*Recursion tree:*





# Idea of master theorem


*Recursion tree:*





# Idea of master theorem


*Recursion tree:*





# Idea of master theorem


*Recursion tree:*





# Idea of master theorem


*Recursion tree:*





# Idea of master theorem

*Recursion tree:*





# Appendix: geometric series

$$1 + x + x^2 + \cdots + x^n = \frac{1 - x^{n+1}}{1 - x} \quad \text{for } x \neq 1$$

$$1 + x + x^2 + \cdots = \frac{1}{1 - x} \quad \text{for } |x| < 1$$

Return to last  
slide viewed.

