

# Supplementary materials for Analysis of Cohort Studies with Multivariate, Partially Observed, Disease Classification Data

BY NILANJAN CHATTERJEE

Division of Cancer Epidemiology and Genetics,  
National Cancer Institute, NIH, DHHS. Rockville, MD 20852, USA.  
chattern@mail.nih.gov

SAMIRAN SINHA

Texas A&M University, College Station, TX 77843, USA.  
sinha@stat.tamu.edu

W. RYAN DIVER AND HEATHER SPENCER FEIGELSON

Department of Epidemiology and Surveillance Research,  
American Cancer Society, Atlanta, GA 30303, USA.

## *Derivation of derivatives of the estimating function*

Define

$$\begin{aligned}
 S_{\theta\theta}^{(2)}(V_i, y^{or}) &= \frac{1}{n} \sum_{j=1}^n I(V_j \geq V_i) \mathcal{E}_{y_i^{or}}(\mathcal{B}_Y \mathcal{B}_Y^\top | X_j) \otimes X_j X_j^\top \omega_{y_i^{or}}(X_j), \\
 S_{\xi\xi}^{(2)}(V_i, u) &= \frac{1}{n} \sum_{j=1}^n I(V_j \geq V_i) \mathcal{E}_u(\mathcal{A}_Y \mathcal{A}_Y^\top | X_j) \omega_u(X_j), \\
 S_{\theta\xi}^{(2)}(V_i, y^{or}) &= \frac{1}{n} \sum_{j=1}^n I(V_j \geq V_i) \mathcal{E}_{y_i^{or}}(\mathcal{B}_Y \mathcal{A}_Y^\top | X_j) \otimes X_j \omega_{y_i^{or}}(X_j), \\
 S_{\xi\theta}^{(2)}(V_i, u) &= \frac{1}{n} \sum_{j=1}^n I(V_j \geq V_i) \mathcal{E}_u(\mathcal{A}_Y \mathcal{B}_Y^\top | X_j) \otimes X_j \omega_u(X_j), \\
 S_{\xi}^{(1)}(V_i, y^{or}) &= \frac{1}{n} \sum_{j=1}^n I(V_j \geq V_i) \mathcal{E}_{y_i^{or}}(\mathcal{A}_Y | X_j) \omega_{y_i^{or}}(X_j), \\
 S_{\theta}^{(1)}(V_i, u) &= \frac{1}{n} \sum_{j=1}^n I(V_j \geq V_i) \mathcal{E}_u(\mathcal{B}_Y | X_j) \otimes X_j \omega_u(X_j),
 \end{aligned}$$

and let  $s_{\theta\theta}^{(2)}(V_i, y^{or})$ ,  $s_{\xi\xi}^{(2)}(V_i, u)$ ,  $s_{\theta\xi}^{(2)}(V_i, y^{or})$ ,  $s_{\xi\theta}^{(2)}(V_i, u)$ ,  $s_{\xi}^{(1)}(V_i, y^{or})$  and  $s_{\theta}^{(1)}(V_i, u)$  denote the corresponding population expectations. Now we can write

$$\frac{\partial S_{\theta}}{\partial \theta^T} = \sum_r \sum_{\Delta_i=1, R_i=r} \left[ \nu_{y_i^{or}}(\mathcal{B}_Y | X_i) \otimes X_i X_i^\top - \frac{S_{\theta\theta}^{(2)}(V_i, y^{or})}{S_{\theta\theta}^{(0)}(V_i, y^{or})} + \frac{S_{\theta}^{(1)}(V_i, y^{or})}{S_{\theta}^{(0)}(V_i, y^{or})} \left\{ \frac{S_{\theta}^{(1)}(V_i, y^{or})}{S_{\theta}^{(0)}(V_i, y^{or})} \right\}^\top \right],$$

$$\begin{aligned}
\frac{\partial S_\theta}{\partial \xi^T} &= \sum_r \sum_{\Delta_i=1, R_i=r} \left[ \mathcal{C}_{y_i^{or}}(\mathcal{B}_Y, \mathcal{A}_Y | X_i) \otimes X_i - \frac{S_{\theta\xi}^{(2)}(V_i, y^{or})}{S^{(0)}(V_i, y^{or})} + \frac{S_\theta^{(1)}(V_i, y^{or})}{S^{(0)}(V_i, y^{or})} \left\{ \frac{S_\xi^{(1)}(V_i, y^{or})}{S^{(0)}(V_i, y^{or})} \right\}^\top \right], \\
\frac{\partial S_\xi}{\partial \xi^T} &= \sum_r \sum_{\Delta_i=1, R_i=r} \left[ \mathcal{V}_{y_i^{or}}(\mathcal{A}_Y | X_i) - \frac{S_{\xi\xi}^{(2)}(V_i, u)}{S^{(0)}(V_i, u)} + \frac{S_\xi^{(1)}(V_i, u)}{S^{(0)}(V_i, u)} \left\{ \frac{S_\xi^{(1)}(V_i, u)}{S^{(0)}(V_i, u)} \right\}^\top \right], \\
\frac{\partial S_\xi}{\partial \theta^T} &= \sum_r \sum_{\Delta_i=1, R_i=r} \left[ \mathcal{C}_{y_i^{or}}(\mathcal{A}_Y, \mathcal{B}_Y | X_i) \otimes X_i - \frac{S_{\theta\xi}^{(2)}(V_i, u)}{S^{(0)}(V_i, u)} + \frac{S_\xi^{(1)}(V_i, u)}{S^{(0)}(V_i, u)} \left\{ \frac{S_\theta^{(1)}(V_i, u)}{S^{(0)}(V_i, u)} \right\}^\top \right],
\end{aligned}$$

where  $\mathcal{V}_{y_i^{or}}$  and  $\mathcal{C}_{y_i^{or}}$  denote variances and covariances with respect to the conditional distribution  $Q_{y_o^r}^{y_m^r}$ . Thus, the components of the matrix  $\mathcal{I} = \lim_{n \rightarrow \infty} (1/n) \partial T_n / \partial \eta$  can now be obtained by replacing  $S$  by  $s$  throughout the above equations and then taking the expectations of the corresponding i.i.d. sums.

#### *Regularity Conditions*

In the following, let  $|\theta|_p$  denote the sum of the absolute values of the  $\theta$ -parameters associated with the vector of covariates  $X_p$ .

- (A1)  $\pi^{(r)}(T, S) > 0$  almost surely in  $(T, S)$  for  $r = (1, 1, \dots, 1)$ .
- (A2) The elements of the second-stage design matrices  $\mathcal{B}$  and  $\mathcal{A}$  remain uniformly bounded in absolute value by constants, say  $c_B$  and  $c_A$ , respectively.
- (A3) Assume the function  $X^{\otimes 3} \exp \left( c_B \sum_{p=1}^P |\theta|_p X_p \right)$  can be bounded by an integrable function of  $X$  uniformly in a open neighborhood of  $\theta_0$ .
- (A4) The functions  $E_{V,X} I(V \geq v) \exp \left( -c_A - c_B \sum_{p=1}^P |\theta|_p X_p \right)$  are bounded away from zero uniformly in  $v$  and  $\eta = (\theta^T, \xi^T)^T$  in a open neighborhood of  $\eta_0$ .
- (A5) The matrix  $\mathcal{I}$  is positive definite.