


SECTION 1 (Maximum Marks: 12)

- This section contains **FOUR (04)** questions.
- Each question has **FOUR** options (A), (B), (C) and (D). **ONLY ONE** of these four options is the correct answer.
- For each question, choose the option corresponding to the correct answer.
- Answer to each question will be evaluated **according to the following marking scheme**:
Full Marks : +3 If **ONLY** the correct option is chosen;
Zero Marks : 0 If none of the options is chosen (i.e. the question is unanswered);
Negative Marks : -1 In all other cases.

- Q.1 An ideal gas (0.5 mol), initially at 2 bar pressure, is compressed at a constant temperature of 600 K in two steps: first, against a constant external pressure of P bar ($2 < P < 8$), and then against constant external pressure of 8 bar. At each step, the compression is stopped only when the pressure of the gas becomes equal to the external pressure. The total work done on the gas in these steps is W . Considering all possible values of P ($2 < P < 8$) and taking the gas constant as R (in $\text{J K}^{-1} \text{mol}^{-1}$), the minimum value of $|W|$ (in J) is

(A)	207R	(B)	600R	(C)	630R	(D)	900R
-----	------	-----	------	-----	------	-----	------


- Q.2 For a reversible reaction $\text{R} \rightleftharpoons \text{P}$, at constant temperature, both the forward and the backward reactions are first order elementary reactions with rate constants k_f and k_b , respectively. At time zero, the concentration of R is $[\text{R}]_0$ and the concentration of P is zero. At any given time, $[\text{R}]$ and $[\text{P}]$ are the concentrations of R and P , respectively. If $k_b = 4k_f$, the correct graphical representation of the reaction is



Q.3 The correct order of dipole moments for the given species is

- | | |
|-----|---|
| (A) | $\text{BF}_3 = \text{NH}_4^+ < \text{NF}_3 < \text{NH}_3$ |
| (B) | $\text{BF}_3 < \text{NH}_4^+ < \text{NF}_3 < \text{NH}_3$ |
| (C) | $\text{NH}_4^+ < \text{BF}_3 < \text{NH}_3 < \text{NF}_3$ |
| (D) | $\text{BF}_3 < \text{NH}_4^+ < \text{NH}_3 < \text{NF}_3$ |

Q.4 Considering LiBH_4 reduces an ester group to the corresponding alcohol and does not reduce a carboxylic acid group, the correct statement about the major products **P**, **Q**, **R** and **S** is



- | | |
|-----|---|
| (A) | P & Q are identical, and R & S are diastereomers. |
| (B) | P & Q are diastereomers, and R & S are identical. |
| (C) | P & Q are diastereomers, and R & S are diastereomers. |
| (D) | P & Q are identical, and R & S are identical. |




SECTION 2 (Maximum Marks: 16)

- This section contains **FOUR (04)** questions.
- Each question has **FOUR** options (A), (B), (C) and (D). **ONE OR MORE THAN ONE** of these four option(s) is(are) correct answer(s).
- For each question, choose the option(s) corresponding to (all) the correct answer(s).
- Answer to each question will be evaluated **according to the following marking scheme**:
Full Marks : +4 **ONLY** if (all) the correct option(s) is(are) chosen;
Partial Marks : +3 If all the four options are correct but **ONLY** three options are chosen;
Partial Marks : +2 If three or more options are correct but **ONLY** two options are chosen, both of which are correct;
Partial Marks : +1 If two or more options are correct but **ONLY** one option is chosen and it is a correct option;
Zero Marks : 0 If none of the options is chosen (i.e. the question is unanswered);
Negative Marks : -1 In all other cases.
- For example, in a question, if (A), (B) and (D) are the **ONLY** three options corresponding to correct answers, then
 choosing **ONLY** (A), (B) and (D) will get +4 marks;
 choosing **ONLY** (A) and (B) will get +2 marks;
 choosing **ONLY** (A) and (D) will get +2 marks;
 choosing **ONLY** (B) and (D) will get +2 marks;
 choosing **ONLY** (A) will get +1 mark;
 choosing **ONLY** (B) will get +1 mark;
 choosing **ONLY** (D) will get +1 mark;
 choosing no option (i.e. the question is unanswered) will get 0 marks; and
 choosing any other combination of options will get -1 marks.

- Q.5 The 2s and the 2p orbital energies of hydrogen atom are $E_{2s}(\text{H})$ and $E_{2p}(\text{H})$, respectively. The 2s and the 2p orbital energies of lithium atom are $E_{2s}(\text{Li})$ and $E_{2p}(\text{Li})$, respectively. The correct option(s) about the orbital energies is(are)

(A)	$E_{2s}(\text{Li}) < E_{2p}(\text{Li})$
(B)	$E_{2s}(\text{H}) = E_{2p}(\text{H})$
(C)	$E_{2p}(\text{H}) < E_{2s}(\text{Li})$
(D)	$E_{2s}(\text{H}) > E_{2s}(\text{Li})$

Q.6 Correct statement(s) about the compounds **X**, **Y** and **Z** is(are)




(A)	X is used for sterilizing drinking water.
(B)	Y has a planar structure.
(C)	Z is used in the enrichment of ^{235}U .
(D)	Y is a stronger Lewis base than ammonia.

Q.7 Reaction of PtF_6 with oxygen (O_2) gas results in the formation of an ionic compound, X^+Y^- .
Correct statement(s) is(are)

(A)	The bond order of X^+ is 1.5.
(B)	Valence <i>d</i> -orbitals of the metal ion in X^+Y^- has 5 electrons.
(C)	PtF_6 acts as an oxidant in this reaction.
(D)	PtF_6 acts as a fluorinating agent in this reaction.

Q.8

In the following reaction sequence, **Q**, **R**, **S** and **T** are the major products.



The correct statement(s) about **Q**, **R**, **S** and **T** is(are)

(A)	S on warming with ammoniacal AgNO_3 results in the formation of silver mirror.
(B)	Q on treatment with Cl_2 (excess)/UV gives gammaxane.
(C)	T is a heterocyclic compound.
(D)	R on acid catalyzed intramolecular cyclization followed by treatment with Zn-Hg/HCl gives 9,10-dihydroxyanthracene.

SECTION 3 (Maximum Marks: 16)



- This section contains **FOUR (04)** questions.
- The answer to each question is a **NUMERICAL VALUE**.
- For each question, enter the correct numerical value corresponding to the answer in the designated place using the mouse and the on-screen virtual numeric keypad.
- If the numerical value has more than two decimal places, **truncate/round-off** the value to **TWO** decimal places.
- Answer to each question will be evaluated **according to the following marking scheme:**
Full Marks : +4 If ONLY the correct numerical value is entered in the designated place;
Zero Marks : 0 In all other cases.

- Q.9 Two cylinders, both fitted with frictionless pistons, are filled with mixtures of He and Ar gases. In the first cylinder, the masses of He and Ar are m_1 and m_2 , respectively. In the second cylinder, the masses of He and Ar are m_2 and m_1 , respectively. The molar mass of Ar is 10 times the molar mass of He. The external pressure applied by the piston on the first cylinder needs to be 5 times that on the second cylinder so that the volume of the gas mixtures in both the cylinders are equal at the same temperature. Assuming He and Ar behave like ideal gases, the value of (m_1/m_2) is ____.

- Q.10 The total number of all possible isomers for the square planar complex with formula $K[M(NCS)(NO_2)(gly)]$ is ____.

(M = metal ion and gly = $NH_2CH_2COO^-$)

- Q.11 The sum of total number of carbonyl groups ($>C=O$) present in the major products **X** and **Y** in the following reactions is ____.



- Q.12 Treatment of buta-1,3-diyne with NaNH_2 (2 equivalents), followed by reaction with excess of *trans*- $\text{CH}_3\text{-CH=CH-CH}_2\text{-Br}$ gives **X** as the major product. The maximum number of carbon atoms that are collinear (in a straight line) in **X** is ____.

SECTION 4 (Maximum Marks: 16)

- This section contains **FOUR (04)** Matching List Sets.
- Each set has **ONE** Multiple Choice Question.
- Each set has **TWO** lists: **List-I** and **List-II**.
- **List-I** has **Four** entries (P), (Q), (R) and (S) and **List-II** has **Five** entries (1), (2), (3), (4) and (5).
- **FOUR** options are given in each Multiple Choice Question based on **List-I** and **List-II** and **ONLY ONE** of these four options satisfies the condition asked in the Multiple Choice Question.
- Answer to each question will be evaluated **according to the following marking scheme**:
Full Marks : +4 **ONLY** if the option corresponding to the correct combination is chosen;
Zero Marks : 0 If none of the options is chosen (i.e. the question is unanswered);
Negative Marks : -1 In all other cases.

- Q.13 **List-I** contains various physical/chemical processes, and **List-II** contains combinations of changes in enthalpy (ΔH) and entropy (ΔS). Match each entry in **List-I** to the appropriate entry in **List-II**, and choose the correct option.

List-I

(P) Physisorption

(Q) Diamond $\rightarrow$ Graphite


(R) Denaturation of protein

(S) Propene $\rightarrow$ Cyclopropane**List-II**(1) $\Delta H > 0$ and $\Delta S > 0$ (2) $\Delta H < 0$ and $\Delta S < 0$ (3) $\Delta H < 0$ and $\Delta S = 0$ (4) $\Delta H > 0$ and $\Delta S < 0$ (5) $\Delta H < 0$ and $\Delta S > 0$

(A)	P $\rightarrow$ 2; Q $\rightarrow$ 3; R $\rightarrow$ 5; S $\rightarrow$ 4
(B)	P $\rightarrow$ 4; Q $\rightarrow$ 3; R $\rightarrow$ 5; S $\rightarrow$ 1
(C)	P $\rightarrow$ 2; Q $\rightarrow$ 5; R $\rightarrow$ 1; S $\rightarrow$ 4
(D)	P $\rightarrow$ 2; Q $\rightarrow$ 5; R $\rightarrow$ 1; S $\rightarrow$ 3

Q.14

Consider the following species:



List-I contains different molecular shapes and **List-II** contains total number of species with the same molecular shapes from the given species. Match each entry in **List-I** with the appropriate entry in **List-II** and choose the correct option.

List-I

(P) See-saw

(Q) T-Shaped

(R) Trigonal Planar

(S) Square Pyramidal

List-II

(1) one

(2) two

(3) three

(4) four

(5) zero


(A)	$P \rightarrow 1; Q \rightarrow 2; R \rightarrow 5; S \rightarrow 3$
(B)	$P \rightarrow 5; Q \rightarrow 4; R \rightarrow 2; S \rightarrow 3$
(C)	$P \rightarrow 3; Q \rightarrow 2; R \rightarrow 1; S \rightarrow 4$
(D)	$P \rightarrow 1; Q \rightarrow 3; R \rightarrow 5; S \rightarrow 4$

Q.15


The **List-II** contains products obtained from the reaction of compounds in **List-I** with $\text{O}_3/\text{Zn-H}_2\text{O}$ followed by cyclization (via more stable enolate) in the presence of aqueous NaOH . Match each entry in **List-I** with appropriate entry in **List-II** and choose the correct option.

List-I


(P)




(Q)




(R)




(S)

**List-II**


(1)




(2)




(3)



(4)












(5)



(A)	P $\rightarrow$ 2; Q $\rightarrow$ 4; R $\rightarrow$ 1; S $\rightarrow$ 3
(B)	P $\rightarrow$ 3; Q $\rightarrow$ 4; R $\rightarrow$ 5; S $\rightarrow$ 2
(C)	P $\rightarrow$ 2; Q $\rightarrow$ 1; R $\rightarrow$ 5; S $\rightarrow$ 3
(D)	P $\rightarrow$ 3; Q $\rightarrow$ 5; R $\rightarrow$ 4; S $\rightarrow$ 2

Q.16

Match the major products obtained in the reactions given in **List-I** with the corresponding structures in **List-II** and choose the correct option.

List-I**List-II**

(A)	P → 2; Q → 1; R → 5; S → 4
(B)	P → 1; Q → 2; R → 4; S → 5
(C)	P → 1; Q → 2; R → 3; S → 4
(D)	P → 2; Q → 1; R → 3; S → 5

END OF THE QUESTION PAPER

JEE (Advanced) 2026 Chemistry Paper-1

Detailed Solutions

Q1. For isothermal irreversible compression:

$$w = -P_{\text{ext}} \Delta V$$

Step-1 work:

$$w_1 = nRT(P/2 - 1)$$

Step-2 work:

$$w_2 = nRT(8/P - 1)$$

Total work:

$$W = nRT(P/2 + 8/P - 2)$$

$$\text{Given } nRT = 0.5 \times 600R = 300R.$$

Differentiate and minimize:

$$df/dP = 1/2 - 8/P^2 = 0$$

$$P = 4 \text{ bar.}$$

Minimum value:

$$W_{\text{min}} = 300R \times 2 = 600R.$$

Answer: (B)

Q2. For reversible first-order reaction:



At equilibrium:

$$k_f[R]_{eq} = k_b[P]_{eq}$$

Given $k_b = 4k_f$,

$$[P]_{eq}/[R]_{eq} = 1/4.$$

Thus concentration of R decreases but remains greater than P at equilibrium. The graph where P rises to a smaller equilibrium value is correct.

Answer: (C)

Q3.

BF₃ is trigonal planar and symmetrical, therefore dipole moment = 0.

NH₄⁺ is tetrahedral and symmetrical, therefore dipole moment = 0.

NF₃ has small dipole moment because bond moments oppose lone pair direction.

NH₃ has larger dipole moment.

Order:

$$BF_3 = NH_4^+ < NF_3 < NH_3$$

Answer: (A)

Q4. LiBH₄ reduces ester groups but does not reduce carboxylic acid groups. Careful stereochemical analysis shows:

P and Q are identical compounds.

R and S are diastereomers.

Answer: (A)

Q5.

- (A) In lithium atom, 2s orbital is lower in energy than 2p orbital due to penetration effect.
(B) In hydrogen atom, 2s and 2p orbitals are degenerate.
(C) False.
(D) $E_{2s}(H) > E_{2s}(Li)$ because lithium 2s orbital is stabilized.

Correct options: A, B, D.

Answer: (A), (B), (D)


Q6.

X is used for sterilization of drinking water.
Z is UF_6 , used in enrichment of U-235.
Other statements are incorrect.

Answer: (A), (C)

Q7.

Reaction:



PtF_6 acts as oxidizing agent.
Oxidation state of Pt = +5.
Therefore Pt has d⁵ configuration.

Correct statements are B and C.

Answer: (B), (C)

Q8.

From reaction analysis:
S gives Tollens' test.
Q forms gammaxane upon chlorination.
T is heterocyclic.
Statement D is incorrect.

Answer: (A), (B), (C)

Q9.

Using ideal gas equation and equal volume condition:

$$n_1/n_2 = P_2/P_1 = 1/5$$


Substituting moles of He and Ar and solving gives:

$$m_1/m_2 = 2$$

Answer: 2

Q10.

Square planar complex:



Considering linkage isomerism and geometrical isomerism, total possible isomers = 6.

Answer: 6

Q11.

Counting total carbonyl groups present in both products X and Y gives:

Total carbonyl groups = 5.

Answer: 5

Q12.

The final product contains consecutive sp-hybridized carbon atoms in a straight chain.

Maximum number of collinear carbon atoms = 8.

Answer: 8

Q13.

Physisorption: $\Delta H < 0$, $\Delta S < 0$

Diamond $\rightarrow$ Graphite: $\Delta H < 0$, $\Delta S > 0$

Denaturation of protein: $\Delta H > 0$, $\Delta S > 0$

Propene $\rightarrow$ Cyclopropane: $\Delta H > 0$, $\Delta S < 0$

Correct matching corresponds to option C.

Answer: (C)

Q14.

See-saw: SOCl_2 only.

T-shaped: ClF_3 and XeF_3^+ .

Square pyramidal: ClF_5 , XeOF_4 and XeF_5^+ .

Trigonal planar: none.

Correct matching corresponds to option A.

Answer: (A)

Q15.

After ozonolysis followed by aldol cyclization through stable enolate formation, the correct matching corresponds to option B.

Answer: (B)

Q16.

Detailed reaction mechanism and product analysis give the correct matching:

$P \rightarrow 1$, $Q \rightarrow 2$, $R \rightarrow 3$, $S \rightarrow 4$

Thus correct option is C.

Answer: (C)