

THE UNIVERSITY OF WARWICK

FOURTH YEAR EXAMINATION: April 2007

LOCAL FIELDS

Time Allowed: **3 hours**

Read carefully the instructions on the answer book and make sure that the particulars required are entered on each answer book.

ANSWER 4 QUESTIONS.

If you have answered more than the required 4 questions in this examination, you will only be given credit for your 4 best answers.

The numbers in the margin indicate approximately how many marks are available for each part of a question.

1. a) Let k be a field. Define the terms **valuation**, **trivial valuation**, **archimedean valuation** and **non-archimedean valuation**. [8]
- b) Suppose that the valuation $|\cdot|$ on k satisfies $|1 + b| \leq 2$ for all $b \in k$ satisfying $|b| \leq 1$. Show that the valuation satisfies the triangle inequality. [10]
- c) Let $|\cdot|_1, |\cdot|_2$ be valuations on k and suppose the existence of an element $c \in k$ such that $|c|_1 < 1$, $|c|_2 > 1$. (i.e. the two valuations are inequivalent). Let $b_1, b_2 \in k$. Show the existence of an element $a \in k$ such that

$$|a - b_1|_1 < 1, \quad |a - b_2|_2 < 1.$$
 [7]

2. a) Suppose k is a complete non-archimedean field. Show that $\sum_{n=0}^{\infty} a_n$ converges if and only if $a_n \rightarrow 0$. [7]
- b) For which finite primes p does the series $\sum_{n=0}^{\infty} \left(\frac{15}{14}\right)^n$ converge in \mathbb{Q}_p ? What is its limit in those fields? [6]
- c) Show that if p is an odd prime then the series $\sum_{n=0}^{\infty} \frac{p^n}{n!}$ converges in \mathbb{Q}_p . (If you need any inequality involving factorials, you should prove it.) [12]

3. a) Let k be complete with respect to a non-archimedean valuation $|\cdot|$. Let \mathcal{O} be the ring of valuation integers and suppose $f(X) \in \mathcal{O}[X]$. Let $a_0 \in \mathcal{O}$ satisfy

$$|f(a_0)| < 1, \quad |f'(a_0)| = 1.$$

A form of Hensel's Lemma says that there is a **unique** $a \in \mathcal{O}$ satisfying

$$f(a) = 0, \quad |a - a_0| < 1.$$

Assuming the existence of a , prove its uniqueness. [10]

- b) Use Hensel's Lemma to show that the equation $X^n = 7$ has a solution in \mathbb{Z}_3 for all integers $n \geq 1$, such that $3 \nmid n$. [7]
- c) Show that the polynomial $X^4 - X^2 + X + 1$ has precisely two roots in \mathbb{Z}_5 . [8]
-

4. a) Let $|\cdot|$ be a discrete valuation on local field k . Let β be the root of an Eisenstein polynomial in $k[X]$. Show that $k(\beta)/k$ is completely ramified and β is a prime element of $k(\beta)$. [10]
- b) Let $p \neq 2$. Show that $\mathbb{Q}_p(\zeta)$, where $\zeta^p = 1$, $\zeta \neq 1$, is completely ramified, and that $\lambda = \zeta - 1$ is a prime element. [8]
- c) Let θ be a root of $X^3 - X - 1 \in \mathbb{Q}_3[X]$. Show that $\mathbb{Q}_3(\theta)$ is an unramified extension of \mathbb{Q}_3 of degree 3. [7]
-

5. a) State Strassman's theorem on the number of zeros of a power series (without proof). [5]
- b) Let A, B be integers. Suppose that the polynomial $X^2 - AX - B$ has two roots α, β in \mathbb{Z}_7 , satisfying

$$\alpha \equiv 8 \pmod{7^2}, \quad \beta \equiv 1 \pmod{7^2}.$$

Let u_m be the recurrence sequence given by

$$u_0 = 2, \quad u_1 = A, \quad u_{m+2} = Au_{m+1} + Bu_m \quad (m \geq 0).$$

Show that $u_m = \alpha^m + \beta^m$. Use Strassman's theorem to show that $u_m = 2$ only for $m = 0$. [20]
