

# Exam Presentation

## Life Insurance Mathematics

Sami Zweidler

ETH Zurich

20/01/2026

## 1 Markov Model

- Banach Fixed Point Theorem
- The Fixed Point Algorithm
- Fixed Point The Case Where Multiple Derivatives Are Zero at The Fixed Point

## 2 Steffensen's Acceleration Method

- Aitken's  $\Delta^2$  Method
- Steffensen's Acceleration Method

## 3 Comparison With Fixed point iteration and Steffensen's Acceleration Method

# Markov Chains

# Frame Title

# Frame Title

# Fixed Point Iteration Method

## Definition

A point  $p$  is a **fixed point** for a given function  $g$  if  $g(p) = p$ .

## Definition

A point  $p$  is a **fixed point** for a given function  $g$  if  $g(p) = p$ .

Fixed point problems and root finding problems are in fact equivalent.

- if  $p$  is a fixed point of the function  $g$ , then  $p$  is a root of the function

$$f(x) = [g(x) - x]h(x)$$

[as long as  $h(x) \in \mathbb{R}$ ]

- if  $p$  is a root of the function of  $f$ , then  $p$  is a fixed point of the function

$$g(x) = x - h(x)f(x)$$

[as long as  $h(x) \in \mathbb{R}$ ]

# Fixed Point Iteration Method

## Definition

Let  $U$  be a subset of a metric space  $X$ .

A function  $g:U \rightarrow X$  called **Lipschitz continuous** provided there exists a constant  $\lambda \geq 0$  (called Lipschitz constant)

such that  $\forall x,y \in U \ d(g(x),g(y)) \leq \lambda d(x,y)$

if  $\lambda \in [0,1]$ , then  $g$  is called **contraction** (with contraction constant  $\lambda$ ).



# Fixed Point Iteration Method

## Definition

Let  $U$  be a subset of a metric space  $X$ .

A function  $g:U \rightarrow X$  called **Lipschitz continuous** provided there exists a constant  $\lambda \geq 0$  (called Lipschitz constant)

such that  $\forall x,y \in U \ d(g(x),g(y)) \leq \lambda d(x,y)$

if  $\lambda \in [0,1]$ , then  $g$  is called **contraction** (with contraction constant  $\lambda$ ).

## Theorem (A Fixed Point Theorem)

Suppose  $g : [a, b] \rightarrow [a, b]$  is continuous. Then  $g$  has a fixed point.

# Fixed Point Iteration Method

## Lemma

A contraction has at most one fixed point.

# Fixed Point Iteration Method

## Lemma


A contraction has at most one fixed point.

## Corollary

Suppose  $g : [a, b] \rightarrow [a, b]$  is continuous and  $\lambda := \sup |g'(x)| < 1$  for  $x \in (a, b)$

Then  $g$  is a contraction with contraction constant  $\lambda$ .

# Graphical determination of the existence of a fixed point for the function $g(x) = \frac{x^2-3}{2}$



# Banach Fixed Point Theorem

## Theorem (Banach Fixed Point Theorem)

*Let  $U$  be a complete subset of a metric space  $X$ , and let  $g:U \rightarrow U$  be a contraction with contraction constant  $\lambda$ .  
Then*

# Banach Fixed Point Theorem

## Theorem (Banach Fixed Point Theorem)

*Let  $U$  be a complete subset of a metric space  $X$ , and let  $g:U \rightarrow U$  be a contraction with contraction constant  $\lambda$ .  
Then*

- $g$  has a unique fixed point, say  $p$ .*
- For any sequence  $\{x_n\}$  defined by  $x_n = g(x_{n-1})$ ,  $n=1,2,\dots$  converges to this unique fixed point  $p$ .*

# Banach Fixed Point Theorem

## Theorem (Banach Fixed Point Theorem)

*Let  $U$  be a complete subset of a metric space  $X$ , and let  $g:U \rightarrow U$  be a contraction with contraction constant  $\lambda$ . Then*

- $g$  has a unique fixed point, say  $p$ .*
- For any sequence  $\{x_n\}$  defined by  $x_n = g(x_{n-1})$ ,  $n=1,2,\dots$  converges to this unique fixed point  $p$ .*

*Moreover, we have the **a priori** error estimate*

$$|x_n - p| \leq \frac{\lambda^n}{1 - \lambda} |x_1 - x_0|$$

# Banach Fixed Point Theorem

## Theorem (Banach Fixed Point Theorem)

*Let  $U$  be a complete subset of a metric space  $X$ , and let  $g:U \rightarrow U$  be a contraction with contraction constant  $\lambda$ . Then*

- $g$  has a unique fixed point, say  $p$ .*
- For any sequence  $\{x_n\}$  defined by  $x_n = g(x_{n-1})$ ,  $n=1,2,\dots$  converges to this unique fixed point  $p$ .*

*Moreover, we have the **a priori** error estimate*

$$|x_n - p| \leq \frac{\lambda^n}{1 - \lambda} |x_1 - x_0|$$

*and the **a posteriori** error estimate*

$$|x_n - p| \leq \frac{\lambda}{1 - \lambda} |x_n - x_{n-1}|$$



# Banach Fixed Point Theorem

## Proof

For  $n > m$ , we have

$$\begin{aligned} |x_n - x_m| &= |x_n - x_{n-1} + x_{n-1} - x_{n-2} + \dots + x_{m+1} - x_m| \\ &\leq |x_n - x_{n-1}| + |x_{n-1} - x_{n-2}| + \dots + |x_{m+1} - x_m| \end{aligned}$$

by\*

$$\begin{aligned} &\leq (\lambda^{n-1} + \lambda^{n-2} + \dots + \lambda^m) |x_1 - x_0| \\ &= \lambda^m (\lambda^{n-m-1} + \lambda^{n-m-2} + \dots + 1) |x_1 - x_0| \\ &= \lambda^m \frac{1 - \lambda^{n-m}}{1 - \lambda} |x_1 - x_0| \leq \frac{\lambda^m}{1 - \lambda} |x_1 - x_0| \end{aligned}$$

so that  $x_n$  is Cauchy sequence in  $U$ .

Since  $U$  is complete,  $x_n$  converges to a point  $p \in U$

$$* |x_k - x_{k-1}| = |g(x_{k-1}) - g(x_{k-2})| \leq \lambda |x_{k-1} - x_{k-2}| \leq \dots \leq \lambda^{k-1} |x_1 - x_0|$$

**Continue.**

Now, since  $g$  being contraction is continuous, we have

$$p = \lim_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} g(x_{n-1}) = g(\lim_{n \rightarrow \infty} x_{n-1}) = g(p)$$

so that  $p$  is fixed point of  $g$ .

By the lemma  $p$  is the unique fixed point of  $g$ .

**Continue.**

Now, since  $g$  being contraction is continuous, we have

$$p = \lim_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} g(x_{n-1}) = g(\lim_{n \rightarrow \infty} x_{n-1}) = g(p)$$

so that  $p$  is fixed point of  $g$ .

By the lemma  $p$  is the unique fixed point of  $g$ .

Since

$$|x_n - x_m| \leq \frac{\lambda^m}{1 - \lambda} |x_1 - x_0|,$$

letting  $n \rightarrow \infty$

**Continue.**

Now, since  $g$  being contraction is continuous, we have

$$p = \lim_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} g(x_{n-1}) = g(\lim_{n \rightarrow \infty} x_{n-1}) = g(p)$$

so that  $p$  is fixed point of  $g$ .

By the lemma  $p$  is the unique fixed point of  $g$ .

Since

$$|x_n - x_m| \leq \frac{\lambda^m}{1 - \lambda} |x_1 - x_0|,$$

letting  $n \rightarrow \infty$

we get

$$|p - x_m| \leq \frac{\lambda^m}{1 - \lambda} |x_1 - x_0|$$

for  $y_0 = x_{n-1}$ ,  $y_1 = x_n$

$$|y_1 - p| \leq \frac{\lambda}{1 - \lambda} |y_1 - y_0|$$

# The Fixed Point Algorithm

## The Fixed Point Algorithm

If  $g$  has a fixed point  $p$ , then the fixed point algorithm generates a sequence  $\{x_n\}$  defined as

$x_0$ : arbitrary but fixed,

$x_n = g(x_{n-1})$ ,  $n=1,2,3,\dots$  to approximate  $p$ .

# Fixed Point The Case Where Multiple Derivatives Are Zero at The Fixed Point

## Theorem

Let  $g$  be a continuous function on the closed interval  $[a, b]$  with  $\alpha > 1$  continuous derivatives on the interval  $(a, b)$ . Further, Let  $p \in (a, b)$  be a fixed point of  $g$ .

# Fixed Point The Case Where Multiple Derivatives Are Zero at The Fixed Point

## Theorem

Let  $g$  be a continuous function on the closed interval  $[a, b]$  with  $\alpha > 1$  continuous derivatives on the interval  $(a, b)$ .

Further, Let  $p \in (a, b)$  be a fixed point of  $g$ .

if

$$g'(p) = g''(p) = \dots = g^{(\alpha-1)}(p) = 0$$

but  $g^{(\alpha)}(p) \neq 0$ ,

# Fixed Point The Case Where Multiple Derivatives Are Zero at The Fixed Point

## Theorem

Let  $g$  be a continuous function on the closed interval  $[a, b]$  with  $\alpha > 1$  continuous derivatives on the interval  $(a, b)$ .

Further, Let  $p \in (a, b)$  be a fixed point of  $g$ .

if

$$g'(p) = g''(p) = \dots = g^{(\alpha-1)}(p) = 0$$

but  $g^{(\alpha)}(p) \neq 0$ ,

then there exist a  $\delta > 0$  such that for any  $p_0 \in [p - \delta, p + \delta]$ , the sequence  $p_n = g(p_{n-1})$  converges to the fixed point  $p$  of order  $\alpha$  with asymptotic error constant

$$\lim_{n \rightarrow \infty} \frac{|e_{n+1}|}{|e_n|^\alpha} = \frac{|g^{(\alpha)}(p)|}{\alpha!}$$



## Proof

Let's start by establishing the existence of  $\delta > 0$  such that for any  $p_0 \in [p - \delta, p + \delta]$ , the sequence  $p_n = g(p_{n-1})$  converges to the fixed point  $p$ .

## Proof

Let's start by establishing the existence of  $\delta > 0$  such that for any  $p_0 \in [p - \delta, p + \delta]$ , the sequence  $p_n = g(p_{n-1})$  converges to the fixed point  $p$ .

Let  $\lambda < 1$ . Since  $g'(p) = 0$  and  $g'$  is continuous, it follows that there exists a  $\delta > 0$  such that  $|g'(x)| \leq \lambda < 1$  for all  $x \in I \equiv [p - \delta, p + \delta]$ . From this, it follows that  $g: I \rightarrow I$ ; for if  $x \in I$  then,

## Proof

Let's start by establishing the existence of  $\delta > 0$  such that for any  $p_0 \in [p - \delta, p + \delta]$ , the sequence  $p_n = g(p_{n-1})$  converges to the fixed point  $p$ .

Let  $\lambda < 1$ . Since  $g'(p) = 0$  and  $g'$  is continuous, it follows that there exists a  $\delta > 0$  such that  $|g'(x)| \leq \lambda < 1$  for all  $x \in I \equiv [p - \delta, p + \delta]$ . From this, it follows that  $g: I \rightarrow I$ ; for if  $x \in I$  then,

$$\begin{aligned} |g(x) - p| &= |g(x) - g(p)| \\ &= |g'(\xi)| |x - p| \\ &\leq \lambda |x - p| < |x - p| \\ &\leq \delta \end{aligned}$$

## Proof

Let's start by establishing the existence of  $\delta > 0$  such that for any  $p_0 \in [p - \delta, p + \delta]$ , the sequence  $p_n = g(p_{n-1})$  converges to the fixed point  $p$ .

Let  $\lambda < 1$ . Since  $g'(p) = 0$  and  $g'$  is continuous, it follows that there exists a  $\delta > 0$  such that  $|g'(x)| \leq \lambda < 1$  for all  $x \in I \equiv [p - \delta, p + \delta]$ . From this, it follows that  $g: I \rightarrow I$ ; for if  $x \in I$  then,

$$\begin{aligned} |g(x) - p| &= |g(x) - g(p)| \\ &= |g'(\xi)| |x - p| \\ &\leq \lambda |x - p| < |x - p| \\ &\leq \delta \end{aligned}$$

Therefore by the a fixed point theorem established earlier, the sequence  $p_n = g(p_{n-1})$  converges to the fixed point  $p$  for any

## Continue

To establish the order of convergence, let  $x \in I$  and expand the iteration function  $g$  into a Taylor series about  $x=p$ :

$$g(x) = g(p) + g'(p)(x-p) + \dots + \frac{g^{(\alpha-1)}(p)}{(\alpha-1)!}(x-p)^{\alpha-1} + \frac{g^{(\alpha)}(\xi)}{(\alpha)!}(x-p)^{\alpha}$$

where  $\xi$  is between  $x$  and  $p$ .

## Continue

To establish the order of convergence, let  $x \in I$  and expand the iteration function  $g$  into a Taylor series about  $x=p$ :

$$g(x) = g(p) + g'(p)(x-p) + \dots + \frac{g^{(\alpha-1)}(p)}{(\alpha-1)!}(x-p)^{\alpha-1} + \frac{g^{(\alpha)}(\xi)}{(\alpha)!}(x-p)^{\alpha}$$

where  $\xi$  is between  $x$  and  $p$ .

Using the hypotheses regarding the value of  $g^{(k)}(p)$  for  $1 \leq k \leq \alpha - 1$  and letting  $x = p_n$ , the Taylor series expansion simplifies to

$$p_n + 1 - p = \frac{g^{(\alpha)}(\xi)}{\alpha!}(p_n - p)^{\alpha}$$

where  $\xi$  is now between  $p_n$  and  $p$ .

## Continue.

The definitions of fixed point iteration scheme and of a fixed point have been used to replace  $g(p_n)$  with  $p_{n+1}$  and  $g(p)$  with  $p$ .



**Continue.**

The definitions of fixed point iteration scheme and of a fixed point have been used to replace  $g(p_n)$  with  $p_{n+1}$  and  $g(p)$  with  $p$ .

Finally, let  $n \rightarrow \infty$ . Then  $p_n \rightarrow p$ , forcing  $\xi \rightarrow p$  also. Hence

$$\lim_{n \rightarrow \infty} \frac{|e_{n+1}|}{|e_n|^\alpha} = \frac{|g^{(\alpha)}(p)|}{\alpha!}$$

or  $p_n \rightarrow p$  of order  $\alpha$ .





## Theorem (Aitken's $\Delta^2$ method)

*Suppose that*

- $\{x_n\}$  is a sequence with  $x_n \neq p$  for all  $n \in \mathbb{N}$
- there is a constant  $c \in \mathbb{R} \setminus \{\pm 1\}$  and a sequence  $\{\delta_n\}$  such that
  - $\lim_{n \rightarrow \infty} \delta_n = 0$
  - $x_{n+1} - p = (c + \delta_n)(x_n - p)$  for all  $n \in \mathbb{N}$

## Theorem (Aitken's $\Delta^2$ method)

Suppose that

- $\{x_n\}$  is a sequence with  $x_n \neq p$  for all  $n \in \mathbb{N}$
- there is a constant  $c \in \mathbb{R} \setminus \{\pm 1\}$  and a sequence  $\{\delta_n\}$  such that
  - $\lim_{n \rightarrow \infty} \delta_n = 0$
  - $x_{n+1} - p = (c + \delta_n)(x_n - p)$  for all  $n \in \mathbb{N}$

Then

- $\{x_n\}$  converges to  $p$  iff  $|c| < 1$
- if  $|c| < 1$ , then

$$y_n = \frac{x_{n+2}x_n - x_{n+1}^2}{x_{n+2} - 2x_{n+1} + x_n} = x_n - \frac{(x_{n+1} - x_n)^2}{x_{n+2} - 2x_{n+1} + x_n}$$

is well-defined for all sufficiently large  $n$ .

## Theorem (Aitken's $\Delta^2$ method)

Suppose that

- $\{x_n\}$  is a sequence with  $x_n \neq p$  for all  $n \in \mathbb{N}$
- there is a constant  $c \in \mathbb{R} \setminus \{\pm 1\}$  and a sequence  $\{\delta_n\}$  such that
  - $\lim_{n \rightarrow \infty} \delta_n = 0$
  - $x_{n+1} - p = (c + \delta_n)(x_n - p)$  for all  $n \in \mathbb{N}$

Then

- $\{x_n\}$  converges to  $p$  iff  $|c| < 1$
- if  $|c| < 1$ , then

$$y_n = \frac{x_{n+2}x_n - x_{n+1}^2}{x_{n+2} - 2x_{n+1} + x_n} = x_n - \frac{(x_{n+1} - x_n)^2}{x_{n+2} - 2x_{n+1} + x_n}$$

is well-defined for all sufficiently large  $n$ .

Moreover  $\{y_n\}$  converges to  $p$  faster than  $\{x_n\}$  in the sense that

$$\lim_{n \rightarrow \infty} \frac{y_n - p}{x_n - p} = 0$$

**Proof.**

Let  $e_n = x_n - p$



**Proof.**

$$\text{Let } e_n = x_n - p$$

$$y_n - p = \frac{x_{n+2}x_n - x_{n+1}^2}{x_{n+2} - 2x_{n+1} + x_n} - p$$



**Proof.**

Let  $e_n = x_n - p$

$$y_n - p = \frac{x_{n+2}x_n - x_{n+1}^2}{x_{n+2} - 2x_{n+1} + x_n} - p$$

$$y_n - p = \frac{(e_{n+2}+p)(e_n+p) - (e_{n+1}+p)^2}{(e_{n+2}+p) - 2(e_{n+1}+p) + (e_n+p)}$$



**Proof.**

$$\text{Let } e_n = x_n - p$$

$$y_n - p = \frac{x_{n+2}x_n - x_{n+1}^2}{x_{n+2} - 2x_{n+1} + x_n} - p$$

$$y_n - p = \frac{(e_{n+2}+p)(e_n+p) - (e_{n+1}+p)^2}{(e_{n+2}+p) - 2(e_{n+1}+p) + (e_n+p)}$$

$$y_n - p = \frac{e_{n+2}e_n - e_n^2}{e_{n+2} - 2e_{n+1} + e_n}$$



**Proof.**

Let  $e_n = x_n - p$

$$y_n - p = \frac{x_{n+2}x_n - x_{n+1}^2}{x_{n+2} - 2x_{n+1} + x_n} - p$$

$$y_n - p = \frac{(e_{n+2}+p)(e_n+p) - (e_{n+1}+p)^2}{(e_{n+2}+p) - 2(e_{n+1}+p) + (e_n+p)}$$

$$y_n - p = \frac{e_{n+2}e_n - e_n^2}{e_{n+2} - 2e_{n+1} + e_n}$$

since we have

$$x_{n+1} - p = (c + \delta_n)(x_n - p) \quad \text{and} \quad e_{n+1} = (c + \delta_n e_n)$$





**Proof.**

Let  $e_n = x_n - p$

$$y_n - p = \frac{x_{n+2}x_n - x_{n+1}^2}{x_{n+2} - 2x_{n+1} + x_n} - p$$

$$y_n - p = \frac{(e_{n+2}+p)(e_n+p) - (e_{n+1}+p)^2}{(e_{n+2}+p) - 2(e_{n+1}+p) + (e_n+p)}$$

$$y_n - p = \frac{e_{n+2}e_n - e_n^2}{e_{n+2} - 2e_{n+1} + e_n}$$

since we have

$$x_{n+1} - p = (c + \delta_n)(x_n - p) \quad \text{and} \quad e_{n+1} = (c + \delta_n e_n)$$

$$y_n - p = \frac{(c + \delta_{n+1})(c + \delta_n)e_n e_n - (c + \delta_n)^2 e_n^2}{(c + \delta_{n+1})(c + \delta_n)e_n - 2(c + \delta_n)e_n + e_n}$$



**Proof.**

Let  $e_n = x_n - p$

$$y_n - p = \frac{x_{n+2}x_n - x_{n+1}^2}{x_{n+2} - 2x_{n+1} + x_n} - p$$

$$y_n - p = \frac{(e_{n+2}+p)(e_n+p) - (e_{n+1}+p)^2}{(e_{n+2}+p) - 2(e_{n+1}+p) + (e_n+p)}$$

$$y_n - p = \frac{e_{n+2}e_n - e_n^2}{e_{n+2} - 2e_{n+1} + e_n}$$

since we have

$$x_{n+1} - p = (c + \delta_n)(x_n - p) \quad \text{and} \quad e_{n+1} = (c + \delta_n e_n)$$

$$y_n - p = \frac{(c + \delta_{n+1})(c + \delta_n)e_n e_n - (c + \delta_n)^2 e_n^2}{(c + \delta_{n+1})(c + \delta_n)e_n - 2(c + \delta_n)e_n + e_n}$$

$$y_n - p = \frac{(c + \delta_{n+1})(c + \delta_n) - (c + \delta_n)^2}{(c + \delta_{n+1})(c + \delta_n) - 2(c + \delta_n) + 1} (x_n - p)$$



**Proof.**

Let  $e_n = x_n - p$

$$y_n - p = \frac{x_{n+2}x_n - x_{n+1}^2}{x_{n+2} - 2x_{n+1} + x_n} - p$$

$$y_n - p = \frac{(e_{n+2}+p)(e_n+p) - (e_{n+1}+p)^2}{(e_{n+2}+p) - 2(e_{n+1}+p) + (e_n+p)}$$

$$y_n - p = \frac{e_{n+2}e_n - e_n^2}{e_{n+2} - 2e_{n+1} + e_n}$$

since we have

$$x_{n+1} - p = (c + \delta_n)(x_n - p) \quad \text{and} \quad e_{n+1} = (c + \delta_n e_n)$$

$$y_n - p = \frac{(c + \delta_{n+1})(c + \delta_n)e_n e_n - (c + \delta_n)^2 e_n^2}{(c + \delta_{n+1})(c + \delta_n)e_n - 2(c + \delta_n)e_n + e_n}$$

$$y_n - p = \frac{(c + \delta_{n+1})(c + \delta_n) - (c + \delta_n)^2}{(c + \delta_{n+1})(c + \delta_n) - 2(c + \delta_n) + 1} (x_n - p)$$

$$y_n - p = \frac{(c + \delta_n)(\delta_{n+1} - \delta_n)}{(c - 1) + c(\delta_{n+1} + \delta_n) + \delta_n(\delta_{n+1} - 2)}$$



**Proof.**

Let  $e_n = x_n - p$

$$y_n - p = \frac{x_{n+2}x_n - x_{n+1}^2}{x_{n+2} - 2x_{n+1} + x_n} - p$$

$$y_n - p = \frac{(e_{n+2}+p)(e_n+p) - (e_{n+1}+p)^2}{(e_{n+2}+p) - 2(e_{n+1}+p) + (e_n+p)}$$

$$y_n - p = \frac{e_{n+2}e_n - e_n^2}{e_{n+2} - 2e_{n+1} + e_n}$$

since we have

$$x_{n+1} - p = (c + \delta_n)(x_n - p) \quad \text{and} \quad e_{n+1} = (c + \delta_n)e_n$$

$$y_n - p = \frac{(c+\delta_{n+1})(c+\delta_n)e_n e_n - (c+\delta_n)^2 e_n^2}{(c+\delta_{n+1})(c+\delta_n)e_n - 2(c+\delta_n)e_n + e_n}$$

$$y_n - p = \frac{(c+\delta_{n+1})(c+\delta_n) - (c+\delta_n)^2}{(c+\delta_{n+1})(c+\delta_n) - 2(c+\delta_n) + 1} (x_n - p)$$

$$y_n - p = \frac{(c+\delta_n)(\delta_{n+1} - \delta_n)}{(c-1) + c(\delta_{n+1} + \delta_n) + \delta_n(\delta_{n+1} - 2)}$$

$$\text{Therefore } \lim_{n \rightarrow \infty} \frac{y_n - p}{x_n - p} = 0$$



## Steffensen's Acceleration Method

Steffensen's Method is a combination of fixed-point iteration and the Aitken's  $\Delta^2$  method:

## Steffensen's Acceleration Method

Steffensen's Method is a combination of fixed-point iteration and the Aitken's  $\Delta^2$  method:

Suppose we have a fixed point iteration:

$$x_0, x_1 = g(x_0), x_2 = g(x_1), \dots$$

## Steffensen's Acceleration Method

Steffensen's Method is a combination of fixed-point iteration and the Aitken's  $\Delta^2$  method:

Suppose we have a fixed point iteration:

$$x_0, x_1 = g(x_0), x_2 = g(x_1), \dots$$

Once we have  $x_0, x_1$ , and  $x_2$ , we can compute

$$y_0 = x_0 - \frac{(x_1 - x_0)^2}{x_2 - 2x_1 + x_0}$$

At this point we "restart" the fixed point iteration with  $x_0 = y_0$

## Steffensen's Acceleration Method

Steffensen's Method is a combination of fixed-point iteration and the Aitken's  $\Delta^2$  method:

Suppose we have a fixed point iteration:

$$x_0, x_1 = g(x_0), x_2 = g(x_1), \dots$$

Once we have  $x_0, x_1$ , and  $x_2$ , we can compute

$$y_0 = x_0 - \frac{(x_1 - x_0)^2}{x_2 - 2x_1 + x_0}$$

At this point we "restart" the fixed point iteration with

$$x_0 = y_0$$

e.g.

$$x_3 = y_0, x_4 = g(x_3), x_5 = g(x_4),$$

and compute

$$(x_4 - x_3)^2$$



# Comparison with Fixed Point Iteration and Steffensen's Acceleration Method

## EXAMPLE

Use the Fixed Point iteration method to find a solution to  $f(x) = x^2 - 2x - 3$  using  $x_0 = 0$ , tolerance  $= 10^{-1}$  and compare the approximations with those given by Steffensen's Acceleration method with  $x_0 = 0$ , tolerance  $= 10^{-2}$ .

# Comparison with Fixed Point Iteration and Steffensen's Acceleration Method

## EXAMPLE

Use the Fixed Point iteration method to find a solution to  $f(x) = x^2 - 2x - 3$  using  $x_0 = 0$ , tolerance  $= 10^{-1}$  and compare the approximations with those given by Steffensen's Acceleration method with  $x_0 = 0$ , tolerance  $= 10^{-2}$ .

- We can see that my MATLAB code while Fixed Point iteration method reaches the root by 788 iteration, Steffensen's Acceleration method reaches the root by only 3 iterations.