

COURSEWORK

IMPERIAL COLLEGE LONDON

DEPARTMENT OF COMPUTING

477 - Computational Optimisation

Author:

Jiahao Lin (CID: 00837321)

Date: November 6, 2016

1 Part 1

1.1 1

1) To prove is $\log \sum_{k=1}^{10} \exp(B_{jk})$ is convex, we define:

$$f(B_j) = \log \sum_{k=1}^{10} \exp(B_{jk}) \quad (1)$$

$$f(x) = \log \sum_{k=1}^{10} \exp(x_k) \quad (2)$$

First we need to compute the Hessian of $f(x)$:

$$\nabla f(x) = \frac{1}{1^\top Z} \times z \quad (3)$$

$$\nabla^2 f(x) = \frac{1}{1^\top Z} \times \text{diag}(Z) - \frac{Z^\top Z}{(1^\top Z)^2} \quad (4)$$

$$\text{where } Z = \sum_{k=1}^{10} \exp(x_k) \quad (5)$$

For convexity we need to prove the Hessian is positive semi-definite, we need to prove:

$$\nabla^2 f(x) \geq 0 \quad (6)$$

$$v^\top \nabla^2 f(x) v \geq 0 \quad (7)$$

$$v^\top \left(\frac{1}{1^\top Z} \times \text{diag}(Z) - \frac{Z^\top Z}{(1^\top Z)^2} \right) v \geq 0 \quad (8)$$

$$\frac{(\sum_{k=1}^{10} Z_k V_k^2) \sum_{k=1}^{10} Z_k - \sum_{k=1}^{10} (Z_k v_k)^2}{(1^\top Z)^2} \geq 0 \quad (9)$$

According to Cauchy-Schwartz inequality:

$$\sum_{k=1}^{10} (Z_k v_k)^2 \leq \sum_{k=1}^{10} Z_k \times \sum_{k=1}^{10} (Z_k V_k^2) \quad (10)$$

There for the Hessian is positive semi-definite holds, the Log-Sum-Exp function is convex.

2) Denote affine mapping on β_k as $g(\beta_k)$:

$$g(\beta_k) = x_i^\top \beta_k \quad (11)$$

And denote Log-Sum-Exp function as $f(x_k)$:

$$f(x_k) = \log \sum_{k=1}^{10} \exp(x_k) \quad (12)$$

Now we know that $f(x_k)$ is convex, and we want to prove that $f \circ g$ is convex as well, from the definition of convexity, we need to prove:

$$f \circ g(\alpha x + (1 - \alpha)y) \leq \alpha f \circ g(x) + (1 - \alpha)f \circ g(y) \quad (13)$$

$$\text{for any } x, y \text{ and } \alpha \in [0, 1] \quad (14)$$

Since we know:

$$g(\alpha x + (1 - \alpha)y) = \alpha g(x) + (1 - \alpha)g(y) \quad (15)$$

We can prove that:

$$f \circ g(\alpha x + (1 - \alpha)y) = f(g(\alpha x + (1 - \alpha)y)) \quad (16)$$

$$= f(\alpha g(x) + (1 - \alpha)g(y)) \quad (17)$$

$$\leq \alpha f \circ g(x) + (1 - \alpha)f \circ g(y) \quad (18)$$

Now we know that $\log \sum_{k=1}^{10} \exp(x_i^\top \beta_k)$ is convex as well.

3)