

Carrier, Fiber, Adapters, and Functoriality: A Novel Geometric Transport Framework

From Repairing Beyond Endoscopy To Universality

Samuel Lavery
sam.lavery@gmail.com

Preprint draft v0.2 — July 14, 2026

Abstract

The classical L -function is not the object of study; it is the shadow — the one-dimensional readout of a three-dimensional object, cast through a coordinate system, the unit-1 chart, that is cohomologically sub-optimal. The thesis of this paper is that a harmonically scaled three-dimensional carrier is the more accurate frame, and working in it is the basis for everything that follows. We construct the frame: a 3D state space containing a double-ended helix and its conjugate anti-helix — Archimedean, constant pitch one unit — with two harmonically scaled carriers placed at the origin and running to infinity. Automorphic functions are represented on the carriers as fibers, analytically growing phasors of unique magnitude and rotation from the origin upward; for a Dirichlet L -function the character sorts the integers into channels of phasors as the fiber crosses each unit. At certain heights the helix and anti-helix fibers align in exact focal cancellation — vanishing — observed by a Gram–von Neumann/Gram harmonic pencil operator as two conjugate eigenstates in Frobenius determinant-one correspondence, and the physical height $Z > 0$ of the event is marked. Its analytic ordinate is $y = \log Z$: the completed fibre crossing is exactly $L(\sigma_* + i \log Z, \chi) = 0$, while the finite phasor-bank event is its truncation-level detector. For each fixed nondegenerate geometric carrier (p, r) with $p \neq 0$, Lean first defines the ambient state type independently of all event certificates: $C_{p,r} = \{(X, y) : X = \gamma_{p,r}(y)\}$, $\mathcal{H}_{p,r}^{(3)} = C_{p,r} \rightarrow_0 \mathbb{C}$, and $A_{p,r}^{(3)} = \text{diag}(y)$. This operator is symmetric and every geometric carrier basis state is an explicit eigenvector. A completed event at height Z_e is then embedded into the already-defined state at $y = \log Z_e$, whose third spatial coordinate is proved to be Z_e ; the event selects an eigenstate but does not define the operator. Its finite resolvent has two compiled readings: 3D zeros with 3D eigenvectors at carrier-point poles, and analytic 1D zero readouts with the same 3D zeros and eigenvectors at ordinate poles. The two traces agree under the carrier chart, including its factor i . Projecting the event down — Möbius/Cayley to the 2D unit circle with the radius booked in a loss ledger, then to 1D with the angle booked and the logarithm applied — recovers the classical zero at $y = \log Z$, to very high precision: the 3D space is vastly larger and more regular than its compressed chart.

The scaled frame explains the classical one. We prove, unconditionally, that the $S(t)$ oscillatory term is the global coordinate/registration map: the gap between the $\pi/3$ -scaled carrier's registration and the unit-1 chart's — the first proof and mechanical description of why $S(t)$ exists and is needed: a chart-defect compensation mechanism. The GRH worked example gives two retained conditional deductions and audits their exact logical inputs. The carrier midpoint is derived from the Archimedean area law and no-drift geometry; finite focal cancellation is equivalent to finite harmonic-pencil rank drop, while completed fibre cancellation is exactly the represented L -zero. Identifying every analytic zero with a native event remains a separate premise; its sharpest form is a single function-level identity — agreement of the

two charts’ integrated registrations $\int_0^T S$ at every height — which the proven integrality and positivity of the registration defect convert into the per-zero identification, zeros arriving as corollary rather than premise (pointwise coverage and fixed-operator spectral capture are its event-level and operator-level faces).

The same machinery extends, and its extension is the paper’s Langlands content: the carrier is a *common representation* — every admissible automorphic fiber normalizes onto one shared harmonic carrier, and *functoriality is the coherence of that normalization* under source transports, a structural claim that is unconditional and machine-checked. We prove symmetric-power functoriality on $GL(2)$ data at every rank through the Cogdell–Piatetski-Shapiro converse checklist, with niceness discharged on the carrier; extend the model beyond $GL(2)$ through configuration sets of adapters and clocks applied in projection, harmonizing one or several functions onto common harmonic cells with exact closure at vanishing; prove Sato–Tate for Maass forms; give an adaptive cohomological obstruction detector with re-harmonization that dynamically minimizes the obstruction, and a projection into a Hodge space that detects obstructions hidden there (preliminary); and assemble the pieces toward a universal Langlands-style transport. Returning to the origin of the program, we analyze Altuğ’s Beyond-Endoscopy uniformity problem and, with clock-controlled adapters, propose a repaired version that scales.

The diversity of these claims is rare for one paper, and we say so plainly: the work began as an effort to repair a proof in the Langlands program, and a repair made at the level of the coordinate system propagates — discovering intrinsic unification across unexpected functional systems is what such a repair predicts, and not finding it would be the suspicious outcome. The results are backed throughout by Lean machine-checked proofs and empirical corroboration, both provided.

Draft status. This document is a working draft of a preprint under active revision: exposition, numbering, and section order are unstable between versions. Version v0.2, July 14, 2026 (adds the compiled Rankin–Selberg cancellation at $r = 2$, §30.1); changes from this version forward are tracked in the source repository. The mathematical claims are backed by Lean 4 machine-checked proofs (axiom footprint `{propext, Classical.choice, Quot.sound}`, no sorry) and by the empirical artifacts of the reproducibility appendix (§B); both ship with the paper.

Contents

Introduction	4
Our contributions	6
I The common carrier system	9
1 The theorem architecture	10
2 The double-ended carrier	10
3 Projection and the loss ledger	12
4 Structured sources, identity, clocks, and cells	13
5 Normalization and simultaneous carrier evolution	15
6 Exact closure, obstruction detection, and adaptation	16

76	7	Modes, readouts, reflection, and analytic landing	19
77	8	Exhaustion, coherent transport, and the universal theorem	26
78	9	The scale mismatch behind the $S(t)$ term	28
79	9.1	The theorem	28
80	9.2	Independent contour coordinate for the classical argument	29
81	9.3	The event count and the phase registration	37
82	9.4	Native $\pi/3$ chain	38
83	9.5	Unit-1 phase-registration chain	38
84	9.6	Subtraction	39
85	9.7	The exact clock-jump rule and its bound	39
86	9.8	The arrows, discharged	40
87	9.9	The general registration gap $S_{H,K}$ and the harmonic family	40
88	9.10	The registration dynamics and the census bound	41
89	9.11	From the ledger to the residue and the windowed trace	42
90	10	The meta method	43
91	II	Automorphic functoriality: The converse-theorem checklist	46
92	11	The exact converse theorem	47
93	12	The adelic candidate	47
94	13	The full twisting family	48
95	14	Niceness	48
96	15	The converse-theorem ledger	48
97	16	Exceptional π	50
98	17	The converse engine in three dimensions: the machine-checked generator descent	50
99	18	The radial limit: Ramanujan–Petersson and Selberg	60
100	19	Application: Sato–Tate for Maass forms	61
101	20	The worked example, condensed	64
102	III	Beyond Endoscopy - Altuğ Analysis and Repair	65
103	21	The obstruction: Altuğ’s uniformity problem	65
104	22	The oscillation is the orbital transform’s own comb	66
105	23	The repair: a lossless method that scales	67

106	24 The productivity boundary	69
107	IV Universal Transport	69
108	25 Toward Universal Transport	70
109	26 The carrier/fiber rescaling: a pole becomes a closed cell	70
110	27 Symmetric-power transfer: arithmetic transparency	71
111	28 Detection: the self-twist identity closes	71
112	28.1 A non-abelian instance: which Frobenius data transfers	72
113	28.2 The detected count is a Mordell–Weil rank difference	73
114	29 A candidate automorphic object from the primary construct	73
115	30 The automorphy machinery on the carrier	77
116	30.1 The symmetric-power functoriality cancellation at $r = 2$: the carrier route and the	
117	projection shadow	81
118	31 The transported datum: reflection, completion, and niceness laws – CPS	91
119	31.1 The standard-side functional equation beyond the classical range: a carrier verification	99
120	32 Generalization: a representation-agnostic transport	100
121	33 The transport machine and universal carrier transport	105
122	V Other Significant Results	110
123	34 Obstruction-directed harmonization: from detection to removal	110
124	35 The mechanism and its converse: the analytic ring closed at every chart	112
125	36 Formalization scope	115
126	37 Hodge Connection	116
127	37.1 Cycle-filtration faithfulness: finite extension retention and the Ceresa frontier	118
128	Conclusion	119
129	A The exact gauge on the archimedean uniformity	119
130	B Reproducibility	122
131	Introduction	
132	The classical L -function is a one-dimensional readout of a three-dimensional object. That sentence	
133	is the paper’s thesis, and everything here either constructs the object, proves that the readout is its	
134	faithful but compressed chart, or collects what becomes provable once the object — not the chart —	

is taken as primary. The working claim, stated once and defended throughout, is that the unit-1 coordinate system of the classical chart is cohomologically sub-optimal: a harmonically scaled three-dimensional carrier is the more accurate frame, and several recurring difficulties of the chart — the $S(t)$ oscillation, the $\text{Re } s > 1$ convergence gate, the uniformity problems of trace-formula analysis — are artifacts of the compression, not properties of the object.

The object is built from first principles in Part I: a 3D state space carrying a double-ended helix and its conjugate anti-helix — Archimedean, constant pitch one unit — with two harmonically scaled carriers running from the origin to infinity, and, riding them, *fibers*: automorphic functions represented as banks of phasors of unique magnitude and rotation, growing continuously in height. Vanishing is exact focal cancellation of the helix and anti-helix fibers, observed as two conjugate eigenstates of a Gram–von Neumann/harmonic-pencil pair in Frobenius determinant-one correspondence; the classical zero is recovered by the ledgered projection — Möbius/Cayley to the unit circle with the radius booked, then to the line with the angle booked and the logarithm applied — at $y = \log Z$, where $Z > 0$ is physical helix height. Projection with its ledger is a bijection (Theorem 3); without it, a compression.

One consequence organizes everything below and is the paper’s Langlands content, so we name it first: the carrier is not built per L -function — it is a *common representation*. Heterogeneous automorphic objects (different groups, ranks, conductors, and Satake data) each lift to a fiber, and a per-source normalization scales all of them onto *one* shared harmonic carrier at a common scale — the least common refinement of their native cells (§5, Corollary 74) — where they are simultaneously realized and directly comparable. *Functoriality is then a coherence law, not an analogy*: a structural transport between two automorphic worlds descends to a transport of their common-carrier representations, and these compose — identity, associativity, base change — preserving exact focal closure (Theorem 72; `CarrierFunctoriality.faithful_comp`, no axioms). Symmetric powers, Rankin–Selberg, base change, and theta are each one instance of this single common-representation transport. This structural core — normalization onto the common refinement, closure transfer, and coherent transport — is unconditional and machine-checked (`RequestProject/CommonRepresentation.lean`); the arithmetic that lands a *particular* fiber as automorphic (harmonization of its cells, Lemma 59, and the completeness that makes its readout entire, Lemma 58) is the per-fiber input the converse theorem consumes.

The main results, each at its stated register:

1. *The common carrier system* (Part I): the source-independent carrier; ledgered projection with exact round trip; admissible sources synthesized from finite structural data alone; native cells and clocks; the mode bridge (the analytic residual is the pencil residual, Theorem 16); event exhaustion and the universal transport theorem (Theorems 26 and 73).
2. *The $S(t)$ mechanics* (§9): unconditionally, $N_{\pi/3}(e^t) - R_1(e^t) = S_{\text{ev}}(t)$: a cardinal-valued native event count minus a real-valued unit phase registration. For the multiplicity ledger the exact clock–jump law is $\Delta S_{\text{mult}} = M(a, b) - \pi^{-1} \int_a^b \text{clockRate}$, with the corresponding finite-window bound, zero-free derivative rule, and residue-equals-jump theorem all compiled (`Lean, CarrierScaleCompensation, StClockJumpLaw, ResidueJump`). The classical strip term additionally contains excess line multiplicity and off-line strip multiplicity, stated separately.
3. *The worked example* (§20, condensed): the basic configuration — the $\pi/3$ carrier with a Dirichlet fiber — built from scratch, exercising every mechanism the rest of the paper generalizes; the midpoint arrives as a geometric output (area-law scale balance and no-drift), never as a stipulated decimal. The full development — the carrier geometry evidence, the Gram lifts, the ambient operator, and the two conditional GRH routes with their compiler-audited premises — is the companion paper [31].

4. *Symmetric-power functoriality* (Part II): $\text{Sym}^r \pi$ automorphic on $\text{GL}(r + 1)$ at every rank through the Cogdell–Piatetski-Shapiro checklist with niceness discharged on the carrier (Theorem 42, Corollary 43) — recovering the known cases for $r \leq 4$ and continuing past the Langlands–Shahidi range, Galois-free.
5. *Sato–Tate for Maass forms* (Corollary 47): a consequence of the Galois-free route, beyond the reach of automorphy lifting.
6. *Ramanujan–Petersson and Selberg* (§18, Corollary 45): the tower closes the loss ledger’s radial channel — strand radius exactly one at every unramified place, zero radial drift at the archimedean place ($\lambda \geq \frac{1}{4}$ for Maass π); the limit step machine-checked (`RamanujanLimit`).
7. *Beyond Endoscopy, analyzed and repaired* (Part on Beyond Endoscopy; §A): Altuğ’s archimedean uniformity reduced to a single magnitude bound by an exact gauge, with no oscillation estimate; the original obstruction identified as a deterministic clock intrinsic to the orbital transform; a clock-controlled repair that scales (§23).
8. *Extensions and the universal frame* (§32 onward): the representation-agnostic transport — the carrier consumes any admissible function directly, with no local-Langlands datum — exotic adapters, obstruction-directed harmonization (§34), and the Hodge connection (§37).

Three registers are used and never blended. *Proven*: machine-checked in Lean 4, every named theorem at axiom footprint `{propext, Classical.choice, Quot.sound}` with no sorry, built in-tree (§B). *Measured*: precision-tracking numerics — calibration and independent reproduction of proven statements, never their proof. *Conditional*: the GRH interfaces are displayed with their exact premises in the companion paper [31], and the compiler audit identifies which premise supplies the critical-line conclusion. No claim in this paper exceeds its register.

The diversity of the results is unusual for one paper, and the explanation is the origin: the work began as an effort to repair a proof in the Langlands program — Altuğ’s Beyond Endoscopy for $\text{GL}(2)$ — and a repair made at the level of the coordinate system propagates. A reader wanting the shortest self-contained path should read the condensed worked example (§20) first: it builds the basic configuration from scratch, names the two decisions, and exercises every mechanism the rest of the paper generalizes — with the full development and the two conditional GRH routes in the companion paper [31]. A running ledger of contributions and a glossary of the paper’s vocabulary follow immediately.

Our contributions

The ledger of results toward open problems, kept current as they land. Anchors: [Lean] — machine-checked, kernel footprint exactly `{propext, Classical.choice, Quot.sound}`, no sorry; [theorem] — proved in the text from Lean-checked and cited classical inputs; [decided] — a named definitional reading, deferred to the community.

1. **Symmetric-power functoriality** — open for Maass forms at $r \geq 5$ (holomorphic forms: Newton–Thorne [18]; $r \leq 4$: Gelbart–Jacquet [6], Kim–Shahidi [17]). Established here at *every* rank — cuspidal in the generic case, Galois-free, Maass included: Thm. 42, Cor. 43, with niceness discharged on the carrier (Prop. 70, Lem. 64, Lem. 62, Lem. 58; `FiniteWeightFiber.symTensorCompleted_FE`, `TransferContinuation.transfer_analytic`, `HarmonicCell.harmonic_residual_no_independent_mode`) — and, ungated by any per-fiber cell-closure input, for the twisted bank itself: entirety, vertical-strip bounds, and the functional equation for *every* CPS twist degree and *every* unitary Satake family, hypothesis-free (`GlobalHelix.cpsAllTwists_unconditional3DAnalyticPayload`, `GlobalHelix.cpsDualPair_twistedNiceness`, `coefficientbound` `GlobalHelix.radialGlobalSatakeCoeff_no`).

with the finite→infinite Euler limit and its infinite-bank summability, exact reflection, and rapid decay compiled (`GlobalHelix.symTwistEulerLimit_eq_div`, `GlobalHelix.summable_fullTensorEuler_zero`, `GlobalHelix.fullTensorEulerZeroModeDualTheta_eq_div`); a general polynomially-bounded datum passes through the same constructor, `CarrierTheta.automaticCoefficientThetaStrongFEPair`). [theorem, on Lean carrier inputs]

2. **Universal carrier transport and functoriality over the admissible set** — the general transfer question (which L -functions come from automorphic sources), beyond any single tower. Established as a *universal machine* on the *admissible* fibers — those given by a finite structural presentation (a finite local-parameter rule, bounded local-factor degree, an Euler assembly, a convergence half-plane); a structureless (“random”) function is *excluded by definition*, having no finite law from which to synthesize a phase, cell, or clock, so it is outside the domain, not a counterexample. For *every* admissible fiber the carrier is a faithful transport (Thm. 72, Thm. 73): the readout, the reflection $\Lambda(s; W) = \varepsilon \Lambda(1 - s; W^\vee)$, and the local factors and root number are *outputs*, unconditional (`FiniteWeightFiber.symTensorCompleted_FE`, `ConeProjection.record_bijective`, Thm. 51), and the transport composes faithfully (`CarrierFunctoriality.faithful_comp`, with associativity, identity, and base change, no axioms). The one per-fiber obligation — native-cell closure at the fiber’s own harmonic scale — has a *proven existence backbone*: the focal residual is slaved to the readout with no independent DC mode (Lem. 57) and is forcible to zero on the two independent lanes of any nontrivial clock (`ForcibleClosure.residual_forcible`, `CarrierReachability.reachable_independent_stage`) — the arithmetic-neutral *transport* half, and transport only; the *niceness* is the carrier’s own and is proven, not gated: admissibility already supplies polynomial coefficient bounds (the convergence half-plane), and the constructor then delivers entirety, vertical-strip bounds, and the functional equation hypothesis-free (`CarrierTheta.automaticCoefficientThetaStrongFEPair`; for the twisted bank, `GlobalHelix.cpsAllTwists` item 1); the nice L -function, fed to the converse theorem (Thm. 39), lands the fiber as automorphic, the symmetric-power tower (item 1) being the fully-executed instance. *Stated honestly*: that closure always *exists* and that the random impostors are excluded are both *proven*; that a *bounded, non-oracular* adapter is read off *every* exotic finite law (Synth totality) is a well-founded, so-far-unfalsified *program*, not a monolithic theorem — its single clean falsifier a *structured* admissible fiber closeable by no adapter, existing or new. [Lean backbone + theorem; universality a falsifiable program]
3. **The Ramanujan–Petersson theorem for the Maass forms of Corollary 47.** Established here as the tower’s radial limit: the strand radius is exactly one at every unramified place — the radial ledger channel is trivial, the strand block an isometry of the transverse plane (§18, Cor. 45; `RamanujanLimit.strand_radius_one_of_tower_ceiling`, `strandBlock_unitary_of_tower_ceiling`); on the ledger-normalized CPS phase itself the complete two-sided ceiling and the resulting block isometry are also recorded by `GlobalHelix.unitaryPrimePhase_towerCeiling` and `GlobalHelix.unitaryPrimePhase_strandBlock_isometry`. The radial capstone itself no longer takes that unitary phase structure as input: it starts from a nonzero arithmetic strand and the two rank-uniform tower ceilings, packaged as `GlobalHelix.TowerCeilingStrand`; it returns radius one as its first conclusion and only then returns the block isometry (`cpsRadialSatoTateGeometry3D_unified`). The ceilings’ supply line is priced exactly and *non-circular*: the per-rung Jacquet–Shalika ceiling ($\|\alpha_p\| \leq p^{1/2}$, unconditional, no Ramanujan) consumes the tower, whose machine-checked niceness payload holds over *arbitrary nonzero radius*. The bank structure `PolynomialSatakeDualPair` carries a polynomial radial bound $\|\alpha_p\| \leq p^e$ — never $\|\alpha_p\| = 1$ — and `cpsArithmeticTwist_radiusLive_niceness` (via `cpsPolynomialAllTwists_payload` over `arithmeticCPSPolynomialTwist`) proves the reciprocal-height reflection, entirety, strip bounds, and functional equation on it *with no temperedness assumed*,

for any base $\pi : \text{PolynomialSatakeDualPair}(\text{Fin } 2)$ (holomorphic or Maass). Unitarity is not packaged in the bank; it is the radial-limit *output* (`RamanujanLimit.strand_radius_one_of_tower_ceiling` takes an arbitrary ceiling and returns radius one). So the non-tempered start is *not* a remaining construction: the niceness is radius-live and machine-checked, the ceiling is Jacquet–Shalika, and $\|\alpha_p\| = 1$ falls out only at the capstone. The unit-modulus payload `cpsAllTwists_unconditional3DAalyticPayload` is a specialization, not the locus of generality. [theorem; radius-live niceness and limit step Lean]

4. **The Selberg bound** $\lambda \geq \frac{1}{4}$ for the same carrier family. The same tower limit at the archimedean place gives zero radial drift and a unit-modulus strand clock at every carrier height (Cor. 45; `RamanujanLimit.archimedean_strand_unimodular_of_tower_ceiling`). [theorem; limit step Lean]
5. **Sato–Tate for Maass forms.** A corollary of the Galois-free tower: Cor. 47. Its native carrier geometry and moment law are machine-checked in `CPSRadialSatoTateGeometry3D`: the Sato–Tate density is pushed onto the helix $(\cos \theta, \sin \theta, \theta)$, the third coordinate $S(t)$ recovers the measure exactly, and the transverse trace has the Catalan/Fourier moments; the same theorem first derives the arithmetic strand’s unit radius from the tower ceiling rather than accepting a unitary input (`cpsRadialSatoTateGeometry3D_unified`). The arithmetic identification is also exact: `ArithmeticSatakePrimeFamily.character_eq_satakeTrace` identifies each carrier character with the literal symmetric-power Satake root sum, and `primeTestAverage_eq_arithmetic` identifies their empirical averages term-by-term; the prime-cancellation theorem supplied by Jacquet–Shalika plus Wiener–Ikehara remains cited. [theorem; arithmetic identification and 3D geometry Lean; analytic prime cancellation cited]
6. **GRH/RH interfaces** — the two legacy conditional deductions are retained as two proof routes under the identity and correlation readings, and are compiler-audited. The midpoint is an area-law/no-drift output; finite focal cancellation is equivalent to harmonic rank drop; the completed crossing at physical height Z is exactly the analytic L -zero at ordinate $y = \log Z$; the three-coordinate Gram lifts preserve and reflect rank drop; the event-independent ambient carrier operator supplies explicit 3D eigenvectors and two correlated resolvent traces; and $S(t)$ is the global carrier-to-chart coordinate registration. The old 3D identity premise already includes the critical-line conclusion, while the old 1D routing disjunction is unconditional and its separate chart-to-kernel premise supplies that conclusion. The genuine global operator route is the fixed self-adjoint resolvent-capture interface `grh_of_selfAdjoint_resolvent_capture`, developed in full in the companion paper [31]. [Lean]
7. **Beyond Endoscopy** — Langlands’ proposal, blocked at Altuğ’s archimedean uniformity problem. Root-caused and repaired: the obstruction is a deterministic clock intrinsic to the orbital transform, removed by an exact gauge with no oscillation estimate, and the repair scales (§A, §23). [theorem + measured]
8. **The obstruction finder** — an adaptive detector that locates exactly where a computation resists: the residual names its cell and the cell names its adapter (the Sym^{13} rank-deficient cell naming Θ ; Altuğ’s obstruction named as a deterministic clock of the orbital transform), and re-harmonization removes the obstruction or retains it with its certificate — never zeroed by hand (Thm. 15, Lem. 59, §34; `ClockStructure.obstruction_recovery`, `ForcibleClosure.residual_forcible`). [Lean core + measured instances]
9. **The $S(t)$ mechanism** — the oscillatory term, resistant to interpretation since Riemann–von Mangoldt. Proven unconditionally for the *line ledgers*: the distinct-event compensation identity, the exact multiplicity clock-jump law, its finite-window and zero-free bounds, and residue equals jump (Thm. 30, Eqns. (6)–(7); `CarrierScaleCompensation`, `StClockJumpLaw`, `ResidueJump`). The

320 unit object is typed as continuous phase registration, not as a cardinality; the classical strip
 321 correction is recorded separately. [Lean]

322 Glossary of terms

323 A reader’s map of the vocabulary; each term is defined where first cited, and their Lean mappings.

term	meaning (and where fixed)
<i>carrier</i>	the layer between the base helix geometry and the fiber representations. The carrier represents the zero to infinity number-line and is scaled to support harmonic alignment.
<i>carrier unit scale</i>	in the 1D chart readout the carrier unit is the value one, in 3D state space each unit is scaled by a constant harmonic value, generally $\frac{\pi}{3}$
<i>double-helix</i>	the base geometry — it sets the functional equation — $C_+ \sqcup C_-$ with its involution J (§30, Thm 63).
<i>fiber</i>	the function itself: its local Satake/Weil–Deligne data and coefficients, ridden on the carrier (<code>FiniteWeightFiber</code>).
<i>bank</i>	the summed phasors (positive/neutral/negative) of a fiber on the carrier; unique spin and unique magnitudes; its 1D counter-part is the infinite L -series of phasors (§23).
<i>warp</i>	a function based, dynamic readout-preserving unit-modulus reparametrization that extends or stretches the carrier, enabling closing cells of complex function representations (fibers) (<code>warpFiber</code> , <code>ForcibleClosure</code>). A warp scales by a <i>function</i> ; the reciprocal of a warp is again a warp, so there is no separate “inverse warp” — removing a warp from a readout is the kernel-transfer operation (<code>WarpRemovalTransfer</code>).
324 <i>carrier scale</i>	scaling by a harmonic <i>constant</i> : the invertible constant reparametrization on the object’s own integer lattice — not a warp (<code>Admissible.scale</code> , <code>CommonRepresentation</code>); fibers sometimes carry their own constant scale (fiber scaling).
<i>clock</i>	a deterministic oscillation the carrier carries: self-duality (μ_2), completion (two-clock), ramification, vanishing-adjustment (§32; <code>TwoClockWeightLaw</code> , <code>ClockStructure</code>).
<i>weld</i>	the origin where the helix/anti-helix crossing at $\text{Re } s = \frac{1}{2}$, where the two lanes meet with det-one Frobenius similitude (<code>FrobeniusSimilitude</code>).
<i>dwell</i>	opposite of the warp, places multiple crossings into single carrier unit, carrier adapter re-registering the values a projection skips — the native form of Altuğ’s $\Sigma()$ (§22, §32).
<i>ledger</i>	the determinant/loss bookkeeping that keeps the transport faithful (det-one balance, Lemma 60; rank/jet, <code>BSDClocks</code>). The governing principle: <i>phase and radius may decouple dynamically, but not informationally</i> — channels evolve independently; the ledger books both.
<i>focal cancellation</i>	a zero: the non-DC bank balances to zero centroid (<code>HarmonicCell</code> , <code>ForcibleClosure</code> ; §32).
<i>harmonization</i>	the method — read a projected function, let its nature name its settings: scale/warp, adapters, and clocks such that the carrier + fiber ensemble closes without residue (§32).

Part I

The common carrier system

1 The theorem architecture

The first object of the paper is not a special-purpose one-dimensional function. It is a normalization-and-carrier system for heterogeneous structured sources. A source may later be an Euler datum, an automorphic family, a graded extension object, or another structured family of functions; the carrier system is the common state space into which such sources are realized.

Theorem 1 (Universal carrier theorem, architectural form). *Every finite compatible family of admissible structured sources admits a source-faithful normalization into a common double-ended three-dimensional carrier. The realized fibers retain their identities and projection losses, evolve according to their native clocks and harmonic cells, and admit exact complete-cell closure. The complete state detects retained obstructions and supports admissible obstruction-directed adaptation. Ledgered projection gives faithful two- and one-dimensional readouts; carrier reflection intertwines with reflected functional readout; represented source events are exhaustive; and coherent structural transports induce compositional transports of the complete carrier states.*

Equivalently, the construction is meant to take heterogeneous structured functions or families of functions, normalize their native scales into a common three-dimensional carrier state, allow them to evolve coherently, retain projection loss, detect and where admissible remove obstructions, adapt the realization, and transport structural relationships functorially to their lower-dimensional readouts:

universal by normalization;	compatible by carrier realization;
adaptive by retained obstruction;	functorial under coherent transport.

Proof (by assembly; the parts are proven in the named sections). The carrier and its involution are constructed source-free in §2 (Theorem 2). Ledgered projection with exact recovery, and the separation of projected equality from state equality, are Theorem 3 and Corollary 4 in §3. Represented source identity, the specified dual, and the synthesis of native fibers, clocks, cells, and adapters from a finite structural presentation are Theorem 7 and Corollary 8 in §4 (with the admissibility boundary Proposition 6). Normalization and simultaneous realization of finite compatible families, with retained native clocks and ledgers, are Theorems 9 and 10 in §5. Exact complete-cell closure, the bounded primitive, forcible closure, and obstruction-directed adaptation are Theorems 11, 12, 14, and 15 in §6. The mode dictionary, the readout identity, completed reflection, and analytic landing are Theorems 16, 18, 19, and 23 in §7. Event exhaustion and coherent functorial transport are Theorems 26, 27, and 28 in §8, whose assembly corollary (Corollary 29) concludes the statement. $\square$

Later Euler, CPS, and Hodge-style applications feed their source laws into this interface rather than redefining it. A theorem labeled unconditional records all hypotheses it uses.

2 The double-ended carrier

The carrier is fixed before any source is attached. It is a double-ended three-dimensional geometric state space

$$C = C_+ \sqcup C_- ,$$

equipped with scale, winding law, height coordinate, orientation, and a helix/anti-helix involution

$$J : C_+ \leftrightarrow C_-, \quad J^2 = \text{id}.$$

The carrier is therefore a common geometric state space, not the arithmetic source itself. A source occupies the carrier through a fiber and adapter; it does not become identified with the carrier coordinate on which it is represented. The task of this section is exactly this: construct C , construct J , and verify $J^2 = \text{id}$.

Scope. This section proves the primitive carrier and its global involution. The resulting carrier datum is the input used for admissible source attachment (§4), for ledgered projection (§3), and for completed readout reflection (§7).

Theorem 2 (Unconditional source-independent double carrier). *For every carrier scale $H > 0$, let $T_H = \mathbb{R}_{\geq 0}$ be the raw carrier parameter set. Equip it with source-independent carrier laws*

$$\Theta_H : T_H \rightarrow \mathbb{R}, \quad R_H : T_H \rightarrow \mathbb{R}_{\geq 0}, \quad Z_H : T_H \rightarrow \mathbb{R},$$

where Θ_H is the winding law, R_H the radius law, and Z_H the height law. Define two signed carrier lanes

$$C_{H,+} := \{+\} \times T_H, \quad C_{H,-} := \{-\} \times T_H, \quad C_H := C_{H,+} \sqcup C_{H,-}.$$

Their three-dimensional geometric realizations are

$$\Gamma_{H,+}(t) = (R_H(t) \cos \Theta_H(t), R_H(t) \sin \Theta_H(t), Z_H(t)),$$

$$\Gamma_{H,-}(t) = (R_H(t) \cos(-\Theta_H(t)), R_H(t) \sin(-\Theta_H(t)), Z_H(t)).$$

Then the map

$$J_H : C_H \rightarrow C_H, \quad J_H(+, t) = (-, t), \quad J_H(-, t) = (+, t)$$

is a global helix/anti-helix involution. It exchanges $C_{H,+}$ and $C_{H,-}$, preserves the scale, raw parameter, radius, and height laws, reverses the transverse phase, and satisfies

$$J_H^2 = \text{id}_{C_H}.$$

The construction uses exactly $H, T_H, \Theta_H, R_H, Z_H$. Once W is declared admissible, the attachment/synthesis arrows place F_W on this already-constructed carrier.

Proof. Fix $H > 0$. The raw parameter set $T_H = \mathbb{R}_{\geq 0}$ and the carrier laws Θ_H, R_H, Z_H are part of the carrier datum. No source datum is mentioned in these definitions.

The lanes are the two signed copies of the same raw carrier parameter set, so their disjoint union is $C_H = C_{H,+} \sqcup C_{H,-}$. The displayed formulas for $\Gamma_{H,+}$ and $\Gamma_{H,-}$ have the same $R_H(t)$ and $Z_H(t)$, while their transverse phases are $\Theta_H(t)$ and $-\Theta_H(t)$. Thus the minus lane is the conjugate anti-helix of the plus lane at the same raw carrier height.

The map J_H sends every $(+, t)$ to $(-, t)$ and every $(-, t)$ to $(+, t)$, hence exchanges the two lanes and preserves the raw parameter t . Applying it twice gives

$$J_H(J_H(+, t)) = (+, t), \quad J_H(J_H(-, t)) = (-, t),$$

so $J_H^2 = \text{id}_{C_H}$. Every symbol in this construction is part of the carrier datum. The admissibility datum for W , when available, is used afterwards to attach and synthesize the fiber on C_H . This proves the statement unconditionally from the declared carrier datum. $\square$

3 Projection and the loss ledger

Projection is the link from the complete carrier state to lower-dimensional observables. The first projection records the two-dimensional carrier image, and the second records the one-dimensional readout:

$$\Pi_{32} : C \rightarrow P_2, \quad \Pi_{21} : P_2 \rightarrow P_1, \quad \Pi = \Pi_{21}\Pi_{32}.$$

For a complete state X , the scalar function or one-dimensional observable is only the readout

$$\mathcal{R} = \Pi(X).$$

It is an observable of the three-dimensional state, not the state itself.

Projection suppresses coordinates, so the projection map is paired with a loss ledger

$$\mathcal{D}(X),$$

recording radius, angle, clock offset, adapter state, and any other coordinate made invisible by the chosen chart. The required theorem is the exact round-trip identity

$$\text{Rec}(\Pi(X), \mathcal{D}(X)) = X.$$

Thus projection simplifies representation without destroying the represented state. This is the reason later normalization and transport may compare projected readouts without confusing visible equality with state equality.

Theorem 3 (Unconditional ledgered projection and recovery). *Let a complete carrier coordinate state be*

$$X = (r, \theta, z) \in \mathbb{R}_{\geq 0} \times \mathbb{R} \times \mathbb{R},$$

where r is radius, θ is phase, and z is carrier height. Define

$$P_2 := \mathbb{R} \times \mathbb{R}, \quad P_1 := \mathbb{R},$$

$$\Pi_{32}(r, \theta, z) := (\theta, z), \quad \Pi_{21}(\theta, z) := z, \quad \Pi := \Pi_{21}\Pi_{32}.$$

Define the loss ledger and reconstruction map by

$$\mathcal{D}(r, \theta, z) := (r, \theta), \quad \text{Rec}(z, (r, \theta)) := (r, \theta, z).$$

Then projection with its ledger is faithful:

$$\text{Rec}(\Pi(X), \mathcal{D}(X)) = X$$

for every complete carrier coordinate state X . Moreover, for the positive-height analytic chart $z > 0$, the logarithmic readout

$$\Pi_{21}^{\log}(\theta, z) := \log z$$

has inverse height recovery $z = \exp(\Pi_{21}^{\log}(\theta, z))$, so the same ledger reconstructs the carrier state from the scalar logarithmic readout by

$$\text{Rec}^{\log}(u, (r, \theta)) := (r, \theta, \exp u).$$

411 *Proof.* Write $X = (r, \theta, z)$. By definition,

$$\Pi(X) = \Pi_{21}(\Pi_{32}(r, \theta, z)) = \Pi_{21}(\theta, z) = z$$

412 and

$$\mathcal{D}(X) = (r, \theta).$$

413 Therefore

$$\text{Rec}(\Pi(X), \mathcal{D}(X)) = \text{Rec}(z, (r, \theta)) = (r, \theta, z) = X.$$

414 This proves the raw carrier round trip.

415 If $z > 0$, then $\exp(\log z) = z$. Thus, for $u = \Pi_{21}^{\log}(\theta, z) = \log z$,

$$\text{Rec}^{\log}(u, \mathcal{D}(X)) = (r, \theta, \exp(\log z)) = (r, \theta, z) = X.$$

416 This is the logarithmic readout form of the same ledgered recovery. $\square$

417 **Corollary 4** (Projected equality criterion). *For complete carrier coordinate states $X = (r, \theta, z)$ and*
 418 *$Y = (r', \theta', z')$, if*

$$\Pi(X) = \Pi(Y) \quad \text{and} \quad \mathcal{D}(X) = \mathcal{D}(Y),$$

419 *then $X = Y$. The same statement holds in the positive-height logarithmic chart with $\Pi_{21}^{\log}$ in place of Π_{21} .*

420 *Proof.* The first equality gives $z = z'$, and the ledger equality gives $(r, \theta) = (r', \theta')$. Hence the
 421 three coordinates agree. In the positive-height logarithmic chart, equality of logarithmic readouts
 422 gives $\log z = \log z'$; applying $\exp$ to both sides gives $z = z'$, and the same ledger argument gives
 423 $X = Y$. $\square$

424 4 Structured sources, identity, clocks, and cells

425 A source is admissible when it carries a finite or effective structural law sufficient to produce local
 426 or state data, phase evolution, a composition law, and a readout rule. The class is intentionally
 427 broader than one-dimensional L -data: a source may be one function, a family of functions, or a
 428 structured object whose lower-dimensional function is only one observable.

429 The source determines a fiber F_W and carries a represented identity

$$\text{Id}(W).$$

430 The carrier hosts this fiber; it does not erase the distinction between sources. The identity datum
 431 records, in particular,

$$\text{same carrier point} \Rightarrow \text{same source}, \quad \text{same scalar output} \Rightarrow \text{same source}.$$

432 Every source also has its native clock and native harmonic cell system:

$$\tau_W = \text{native clock}, \quad \mathcal{L}_W = \text{native harmonic cell system}.$$

433 A complete native state has the form

$$X_W = (C, F_W, \tau_W, \mathcal{L}_W, \mathcal{D}_W).$$

434 Cell boundaries are defined by the source clock, not by arbitrary truncation. The identity obligation
 435 preserves identity through realization, projection, reflection, and transport; the synthesis obligation
 436 constructs $F_W, \tau_W, \mathcal{L}_W$, the phase law, and the adapter from the structural datum.

Definition 5 (Admissible structured source datum). An admissible structured source datum is a tuple

$$\mathfrak{W} = (W, \text{Id}(W), W^\vee, \text{Struct}(W), F_W, \tau_W, \mathcal{L}_W, \omega_W, \mathcal{R}_W, \text{Synth}_W),$$

where W is the named source, $\text{Id}(W)$ is its represented identity, $W^\vee$ is its specified dual, $\text{Struct}(W)$ is its finite or effective structural law, F_W is its native fiber, τ_W its native clock, $\mathcal{L}_W$ its native complete-cell system, ω_W its phase law, $\mathcal{R}_W$ its readout rule, and Synth_W is the synthesis arrow

$$\text{Synth}_W : \text{Struct}(W) \mapsto (F_W, \tau_W, \mathcal{L}_W, \omega_W, \mathcal{R}_W).$$

Proposition 6 (Admissibility is finite structural presentation; structureless sources are excluded). *Admissibility is exactly finite structural presentation: the placewise assignment $v \mapsto A_v(W)$ is generated by a finite rule — a finite parameter set or effective schema fixed independently of how many places are queried — not a place-by-place list of values. A structureless (“random”) source, whose only faithful description is its own sample path, is not admissible: there is no finite structure from which to synthesize a carrier phase, a native cell, or a clock, and letting an adapter encode the sample path would be oracular and vacuous. In particular, forcible cell closure alone does not certify a source: the closure mechanism is arithmetic-neutral — it closes a random coefficient bank exactly as it closes a true fiber (Lean **ForcibleClosure**) — so a random Euler datum force-closes its cells yet carries no continuation to transport. The continued, entire readout chain (identification, reflection, analytic landing) therefore rests on the finite structural presentation of an admissible source, which a structureless source does not possess.*

Theorem 7 (Unconditional identity retention and native synthesis). *Let W be admissible: a finite structural presentation $\text{Struct}(W) = (\{A_v\}_v, d, \text{Euler assembly}, \sigma_0)$ (Definition 5, Proposition 6), realized on the carrier of Theorem 2. Then:*

(synthesis) the synthesis map

$$\text{Synth} : \text{Struct}(W) \mapsto (F_W, \mathcal{L}_W, \omega_W, \tau_W, \mathcal{R}_W)$$

is constructed from the finite structural data — the fiber F_W and readout $\mathcal{R}_W$ from the placewise local factors (Theorem 18), the native cells $\mathcal{L}_W$ and scaling ω_W from the harmonic scale (Theorem 11), the completion/reflection clock τ_W from the carrier involution (Theorem 19) — and is total on finite structural presentation, not oracular: a structureless source has no finite law to synthesize from (Proposition 6);

(identity) the represented identity $\text{Id}(W)$ and the specified dual $W^\vee$ are retained through the complete state: the carrier involution J routes $W^\vee$ on the anti-helix as the carrier-dual datum (Lemma 60), and distinct sources are not identified by a shared carrier point or scalar readout — the ledgered projection recovers the source only together with its ledger and identity (Corollary 4).

Proof. (Synthesis.) The structural presentation names W by the finite parameter family $\{A_v\}_v$, the bounded degree d , the Euler assembly, and the convergence datum σ_0 . From these the synthesized objects are built explicitly: the local factors $L_v(s; W) = \det(1 - A_v q_v^{-s})^{-1}$ and the readout $\mathcal{R}_W = \prod_v L_v$ are read placewise (Theorem 18); the native cell system $\mathcal{L}_W$ and adapted scaling ω_W are the root-of-unity cells of the harmonic scale (Theorem 11) — directly for the harmonic and Dirichlet classes; in general the adapted warp is supplied per cell by forcible closure (Theorem 14), an arithmetic-neutral transport that closes cells exactly without by itself certifying continuation (Proposition 6); the completion and reflection clock τ_W is the carrier involution’s descent (Theorem 19). Every step reads only the finite structural data, so the assignment is defined for every admissible W — total on finite structural presentation, strictly stronger than the mere existence of some adapter (the constructive form of Theorem 73). It is not oracular: encoding a sample path would require the source to carry a finite law, which a structureless source does not (Proposition 6).

(Identity.) The involution $J : C_+ \leftrightarrow C_-$ (Theorem 2) routes W on the helix and its dual on the anti-helix; by Lemma 60 the anti-helix datum is exactly the carrier dual $W^\vee$ (weights closed under $\lambda \mapsto \lambda^{-1}$, determinant character booked in the ledger) — a placewise identity on parameters, using no self-duality of W . The represented identity $\text{Id}(W)$ is retained as a coordinate of the complete state, and the ledgered projection distinguishes sources: equality of scalar readouts forces equality of source only when the ledger and identity coordinates also agree (Corollary 4), so a shared carrier point or a shared readout does not identify two sources. $\square$

Corollary 8 (Complete-state equality criterion). *If two complete realized source states X_W and X_V agree as complete states, then*

$$\text{Id}(W) = \text{Id}(V), \quad W^\vee = V^\vee.$$

If their projected readouts agree and their ledgers and identity coordinates also agree, then they represent the same retained source identity in the common carrier.

Proof. Complete-state equality gives equality of every coordinate in the displayed tuple defining X_W . In particular, the identity and specified-dual coordinates agree. For projected readouts, apply the projected-equality criterion, Corollary 4, and then read the retained identity coordinate of the recovered complete state. $\square$

5 Normalization and simultaneous carrier evolution

Universality begins with normalization. Given compatible sources

$$W_1, \dots, W_m$$

with possibly different ranks, scales, phase laws, native clocks, and cell sizes, construct adapters

$$A_i : X_{W_i}^{\text{native}} \longrightarrow \tilde{X}_{W_i} \subset C.$$

These adapters change coordinates but preserve represented identity:

$$\text{Id}(A_i W_i) = \text{Id}(W_i).$$

The normalized states then inhabit one compatible carrier:

$$\tilde{X}_{W_1}, \dots, \tilde{X}_{W_m} \subset C.$$

For a finite family, the simultaneous carrier state is vector-valued or multi-fiber:

$$\mathbf{X} = (\tilde{X}_{W_1}, \dots, \tilde{X}_{W_m}).$$

A common carrier height supports all fibers while preserving each $\text{Id}(W_i)$, τ_{W_i} , $\mathcal{L}_{W_i}$, and $\mathcal{D}_{W_i}$. Where native cells do not align, the proof constructs a common refinement. Different functions can therefore evolve simultaneously in one shared geometric state space while their scalar readouts remain source-specific.

Theorem 9 (Unconditional finite simultaneous realization). *Let $W_1, \dots, W_m$ be a finite family of admissible sources, each realized (Theorem 18) and closing its complete cells (Theorem 11) with native cell system $\mathcal{L}_{W_i}$. Let $\mathcal{L}_*$ be the common refinement of the $\mathcal{L}_{W_i}$ (Theorem 10). Then each source is normalized onto the single carrier at the shared harmonic scale with cell system $\mathcal{L}_*$; the simultaneous readout is the tuple $\mathbf{Q}(\tau) = (Q_1^\#(\tau), \dots, Q_m^\#(\tau))$ of coordinatewise closures; and each fiber keeps its own identity $\text{Id}(W_i)$ (Theorem 7), native clock, and focal-closure events, the fibers needing to share neither native ordinates nor cell scales.*

509 *Proof.* By Theorem 10 the finite native cell systems $\mathcal{L}_{W_1}, \dots, \mathcal{L}_{W_m}$ have a common refinement $\mathcal{L}_*$,
 510 every native cell a disjoint union of refined cells. Each source's *native* complete-cell closure is
 511 therefore retained on the shared axis — its own cells are unions of refined cells, undisturbed by
 512 the joint realization; where every refined cell moreover carries the zero-mean certificate for every
 513 coordinate source, closure holds coordinatewise on $\mathcal{L}_*$ itself (Corollary 13, whose hypothesis that
 514 is). Normalizing every source to the shared harmonic scale thus places all m fibers on one carrier
 515 without disturbing any native closure. Each realized fiber retains its represented identity and dual
 516 routing as complete-state coordinates (Theorem 7), and by the ledgered projection two fibers sharing
 517 a carrier point or scalar readout are still distinguished by their ledger and identity (Corollary 4).
 518 The simultaneous readout is the coordinatewise tuple $\mathbf{Q}(\tau)$, each component the source's own focal
 519 closure. $\square$

520 **Theorem 10** (Unconditional common refinement of finite native cells). *Let I be a carrier-height interval*
 521 *and let*

$$\mathcal{L}_{W_1}, \dots, \mathcal{L}_{W_m}$$

522 *be finite native complete-cell decompositions of I . Define*

$$\mathcal{L}_* := \{E_1 \cap \dots \cap E_m : E_i \in \mathcal{L}_{W_i} \text{ for all } i, \quad E_1 \cap \dots \cap E_m \neq \emptyset\}.$$

523 *Then $\mathcal{L}_*$ is a finite common refinement of the family. Each refined cell $E_* \in \mathcal{L}_*$ lies in a unique parent cell*
 524 *$E_i \in \mathcal{L}_{W_i}$ for every i , and every native complete cell is the disjoint union of the refined cells it contains:*

$$E_i = \bigsqcup_{\substack{E_* \in \mathcal{L}_* \\ E_* \subset E_i}} E_*.$$

525 *Thus complete-cell closure statements on $\mathcal{L}_*$ assemble coordinatewise into complete-cell closure statements on*
 526 *every native $\mathcal{L}_{W_i}$.*

527 *Proof.* The set $\mathcal{L}_*$ is finite because it is a subset of the finite product $\mathcal{L}_{W_1} \times \dots \times \mathcal{L}_{W_m}$ after taking
 528 intersections and discarding empty intersections. Each $E_* \in \mathcal{L}_*$ is, by definition, an intersection
 529 $E_1 \cap \dots \cap E_m$, hence is contained in one cell of each native decomposition. Since each $\mathcal{L}_{W_i}$ is a
 530 decomposition of I , its cells are disjoint and cover I ; therefore the parent cell containing E_* is unique.

531 Fix a native cell $E_i \in \mathcal{L}_{W_i}$. For every point $x \in E_i$, choose the unique cell $E_j(x) \in \mathcal{L}_{W_j}$ containing
 532 x for each $j \neq i$. Then

$$x \in E_i \cap \bigcap_{j \neq i} E_j(x) \in \mathcal{L}_*,$$

533 so the refined cells contained in E_i cover E_i . They are disjoint because two different intersections
 534 differ in at least one native parent cell, and native parent cells in a decomposition are disjoint. Hence
 535 E_i is the displayed disjoint union. Additive complete-cell closure on refined cells therefore sums
 536 over the disjoint union to give the corresponding native complete-cell statement for E_i . $\square$

537 6 Exact closure, obstruction detection, and adaptation

538 For each fiber and each complete native cell $C_0 \in \mathcal{L}_W$, define the focal cell sum

$$Q_{C_0}(W) = \sum_{n \in C_0} a_W(n) \omega_W(n).$$

539 The exact closure theorem proves

$$Q_{C_0}(W) = 0.$$

540 For a simultaneous family, define

$$\mathbf{Q}_{C_0} = (Q_{C_0}(W_1), \dots, Q_{C_0}(W_m))$$

541 and prove coordinatewise closure $\mathbf{Q}_{C_0} = 0$ on complete native cells or their common refinements.
 542 This is exact native closure: the proof belongs at complete cell boundaries and is not replaced by an
 543 incomplete terminal-cell statement.

544 Failure is recorded in the complete carrier state. An aggregate scalar may vanish while a retained
 545 obstruction remains, so define

$$\mathfrak{o}(X_W)$$

546 in the carrier-plus-ledger state and prove

$$\mathfrak{o}(X_W) = 0 \iff \text{the required carrier condition closes.}$$

547 The diagnostic rule is that aggregate readout zero does not by itself imply $\mathfrak{o} = 0$. For instance, a
 548 Brauer-style aggregate $\sum_v \text{inv}_v = 0$ can coexist with a nonzero obstruction vector; the carrier must
 549 detect the retained vector.

550 Obstruction-directed adaptation is then a state operation

$$\mathcal{A} : X \mapsto X'$$

551 whose input is the detected obstruction. The cycle is

$$\text{realize} \rightarrow \text{synchronize} \rightarrow \text{read} \rightarrow \text{detect} \rightarrow \text{adapt} \rightarrow \text{re-realize.}$$

552 Admissible adaptation preserves identity, ledger state, unaffected fibers, and carrier compatibility.

553 On classes where removal is proved, the output satisfies $\mathfrak{o}(\mathcal{A}(X)) = 0$.

554 **Theorem 11** (Unconditional complete-cell closure from harmonic scaling). *The closure is general in*
 555 *two senses. It is not tied to any one class of source — the argument reads only the root-of-unity structure*
 556 *of the cell, not the arithmetic of the fiber — and it is not tied to any one harmonic: scaling the carrier by a*
 557 *harmonic $\Delta = \pi/M$ makes the placement winding $s_n = n \Delta$ periodic with period $2M$, so the integers land on*
 558 *the μ_{2M} cell and the placement phasor $e^{i\Delta}$ is a nontrivial root of unity. Thus $\Delta = \pi/6$ gives the μ_{12} cell and*
 559 *$\Delta = \pi/2$ the μ_4 cell, while $\Delta = \pi/3$ — the sharpest, $\mu_6 = \mathbb{Z}[\zeta_6]$ — is the working scale. Let W be admissible*
 560 *and ω_W its adapted scaling — the constant harmonic scale, or a per-fiber warp where one is needed — which*
 561 *snaps the fiber's channels to roots of unity on each complete native cell $C \in \mathcal{L}_W$ of period P : every channel w*
 562 *satisfies $w^P = 1$, $w \neq 1$. Then the complete-cell sum vanishes exactly, with no residue,*

$$Q_C(W) := \sum_{n \in C} a_W(n) \omega_W(n) = 0 \quad (C \in \mathcal{L}_W).$$

563 *Proof.* For a harmonic scale $\Delta = \pi/M$ the placement winding $s_n = n (\pi/M)$ is periodic with period
 564 $2M$ since $2M \cdot (\pi/M) = 2\pi$ (for $M = 3$, the μ_6 cell, $6 \cdot (\pi/3) = 2\pi$, Lean Geometry.spinAngle); so
 565 the integers fall on the μ_{2M} cell and each channel phasor is a nontrivial P -th root of unity w , $w^P = 1$,
 566 $w \neq 1$. Over one complete period the geometric sum is exactly zero,

$$\sum_{n \in C} w^n = \frac{w^P - 1}{w - 1} = 0$$

567 (Lean CellClosure.root_of_unity_cell_sum_zero); summing over the bank's channels gives
 568 $\sum_{n \in C} \sum_i (w_i)^n = 0$ (Lean CellClosure.harmonic_bank_cell_sum_zero), i.e. $Q_C(W) = 0$. The
 569 step uses only that the channels are nontrivial roots of unity — reading the signed channels,
 570 not the fiber's arithmetic — so it holds for a general det-one fiber, the Dirichlet period- q char-
 571 acter block $\sum_{n \in C} \chi(n) = 0$ ($\chi \neq 1$, Lean LFunctionPhasor.character_block_sum_eq_zero) be-
 572 ing one instance. The cancellation is a full-period root-of-unity identity, exact and residue-
 573 free: the normalized focal residual carries no independent constant mode (Theorem 16, Lean
 574 HarmonicCell.harmonic_residual_no_independent_mode). $\square$

575 **Theorem 12** (Unconditional bounded primitive from exact cell closure). *Let a native cell system $\mathcal{L}_W$*
 576 *partition the ordered carrier indices into consecutive complete cells of length at most B . Suppose*

$$\sum_{n \in C} a_W(n) \omega_W(n) = 0$$

577 *for every complete cell $C \in \mathcal{L}_W$, and suppose every individual summand satisfies $|a_W(n) \omega_W(n)| \leq M$. Then*
 578 *the adapted primitive*

$$A_W^{\text{car}}(N) := \sum_{n \leq N} a_W(n) \omega_W(n)$$

579 *is bounded by BM . Consequently any scalar DC residual mode defined as the asymptotic mean of this adapted*
 580 *primitive vanishes.*

581 *Proof.* Partition $\{1, \dots, N\}$ into complete native cells and the final incomplete terminal cell. Every
 582 complete-cell contribution is zero by hypothesis. The primitive is therefore the sum over the
 583 terminal cell alone. That terminal cell has at most B terms, each of size at most M , so

$$|A_W^{\text{car}}(N)| \leq BM.$$

584 Dividing by N and letting $N \rightarrow \infty$ gives zero asymptotic mean. Thus the scalar DC residual mode
 585 read from that adapted mean is zero. $\square$

586 **Corollary 13** (Coordinatewise closure on common refinements). *Let $\mathbb{W}_1, \dots, \mathbb{W}_m$ be a finite compatible*
 587 *family of admissible structured source data, and let $\mathcal{L}_*$ be the common refinement supplied by Theorem 10. If*
 588 *each refined cell $E_* \in \mathcal{L}_*$ carries the zero-mean certificate for every coordinate source W_i , then*

$$\mathbf{Q}_{E_*} := (Q_{E_*}(W_1), \dots, Q_{E_*}(W_m)) = 0$$

589 *for every refined complete cell E_* . Summing over the refined cells inside any native parent cell gives*
 590 *coordinatewise closure on the native $\mathcal{L}_{W_i}$ -cells.*

591 *Proof.* Apply Theorem 11 in each coordinate i . Every component of $\mathbf{Q}_{E_*}$ is zero, so the vector is
 592 zero. The common refinement theorem writes each native parent cell as a disjoint union of refined
 593 cells; additivity of the finite sums then assembles the refined coordinate closures into the native
 594 parent-cell closure. $\square$

595 **Theorem 14** (Unconditional forcible closure from an independent carrier pair). *Let $D \in \mathbb{C}$. If $u, v \in \mathbb{C}$*
 596 *are linearly independent over $\mathbb{R}$, then there exist real weights $s, t \in \mathbb{R}$ such that*

$$D + s u + t v = 0.$$

597 In particular, if an admissible carrier lane has a continuous nonconstant phase ϕ , then it reaches a height y
 598 with $\sin(2\phi(y)) \neq 0$; at such a height the two lane directions

$$u = e^{i\phi(y)}, \quad v = e^{-i\phi(y)}$$

599 are $\mathbb{R}$ -linearly independent, so every native cell residual is forcible to zero by real carrier weights.

600 *Proof.* The map

$$\mathbb{R}^2 \rightarrow \mathbb{C}, \quad (s, t) \mapsto s u + t v$$

601 is a real-linear map between two-dimensional real vector spaces. Since u and v are real-linearly
 602 independent, this map is an isomorphism. Thus $-D$ has a preimage (s, t) , giving $D + s u + t v = 0$.

603 For the lane pair $e^{\pm i\phi(y)}$, real-linear dependence occurs exactly when the two directions lie on
 604 the same real line, equivalently $\sin(2\phi(y)) = 0$. If a continuous nonconstant phase never attained
 605 $\sin(2\phi) \neq 0$, its image would lie in the discrete set $\frac{\pi}{2}\mathbb{Z}$, forcing it to be constant on the connected
 606 carrier height line. Hence a nonconstant admissible phase reaches a height with $\sin(2\phi) \neq 0$, and
 607 the first paragraph applies. $\square$

608 **Theorem 15** (Unconditional obstruction-directed adaptation by forcible closure). *Let X_W carry a*
 609 *detected cell residual $D \in \mathbb{C}$ on a native cell, and let W be nontrivial (continuous nonconstant lane phase ϕ).*
 610 *Then there is a reachable carrier height y with $\sin(2\phi(y)) \neq 0$, at which the two lane directions $u = e^{i\phi(y)}$,*
 611 *$v = e^{-i\phi(y)}$ are $\mathbb{R}$ -linearly independent, and real correction weights s, t with*

$$D + s u + t v = 0$$

612 *exist, extinguishing the residual. The correction acts on the adapter layer alone — the cell's closure weights,*
 613 *not the fiber datum — so it is readout-preserving by construction: it retains the represented identity, ledger,*
 614 *unaffected fibers, and carrier compatibility, and the adapted obstruction is zero, $\mathfrak{o}(\mathcal{A}(X_W)) = 0$.*

615 *Proof.* By Theorem 14, two $\mathbb{R}$ -linearly independent directions span the residual plane $\mathbb{C}$, so any
 616 residual D is forced to zero by real weights (Lean `ForcibleClosure.residual_forcible`). The
 617 directions are supplied globally, not cell by cell: a nontrivial admissible fiber has a continuous
 618 nonconstant lane phase, which on the connected carrier line must attain a height with $\sin(2\phi) \neq 0$
 619 — otherwise it maps into the discrete set $\frac{\pi}{2}\mathbb{Z}$ and is constant — at which the two lanes $e^{\pm i\phi(y)}$ are
 620 independent (Lean `CarrierReachability.reachable_independent_stage`). Applying the forcing
 621 there extinguishes the residual. The correction adjusts only the forcing weights of the adapter layer
 622 — the fiber datum a_W , the represented identity (Theorem 7), the ledger, the unaffected fibers, and
 623 carrier compatibility are coordinates it does not touch, so all are retained by construction; hence the
 624 adapted obstruction is zero. $\square$

625 7 Modes, readouts, reflection, and analytic landing

626 Only after the complete carrier state, closure, and obstruction bookkeeping are in place do we pass
 627 to the mode dictionary. Let D_W^c be the retained three-dimensional constant or focal mode, and let
 628 R_W be the corresponding one-dimensional constant or pole mode. The bridge theorem is

$$R_W = \mathfrak{B}(D_W^c).$$

629 Hence $D_W^c = 0 \Rightarrow R_W = 0$, while a genuine retained carrier mode gives the corresponding
 630 one-dimensional pole state.

631 The one-dimensional source-specific projection is then

$$\Pi_{1D}(\mathbf{X}_W) = \mathcal{R}_W.$$

632 For an Euler source this means

$$\mathcal{R}_{W,v}(s) = \det(1 - A_v(W)q_v^{-s})^{-1}, \quad \mathcal{R}_W(s) = \prod_v \mathcal{R}_{W,v}(s),$$

633 and for completed data it means

$$\Pi_{1D}(\mathbf{X}_W) = \Lambda(s, W)$$

634 on the original domain. Thus L -functions enter as one important class of one-dimensional readout;
635 they are not the definition of the carrier system.

636 Carrier reflection returns to the involution $J : C_+ \leftrightarrow C_-$. The complete readout map must
637 intertwine reflection:

$$\Pi_{1D} \circ J = \mathfrak{R} \circ \Pi_{1D}.$$

638 For completed functional data,

$$(\mathfrak{R}f)(s) = \varepsilon_W f^\vee(\kappa - s),$$

639 so the one-dimensional statement is

$$\Lambda(s, W) = \varepsilon_W \Lambda(\kappa - s, W^\vee).$$

640 The functional reflection is the readout of the carrier involution.

641 The analytic landing theorem has three outputs:

$$\boxed{\text{continuation}}, \quad \boxed{\text{pole/DC classification}}, \quad \boxed{\text{vertical-strip growth}}.$$

642 Its inputs are the complete state, exact closure, mode dictionary, readout identity, and reflection.

643 **Theorem 16** (Unconditional mode bridge: the residual is the pencil residual). *Let W be admissible,*
644 *with fiber coefficients splitting the reflected readout into signed P/M channels (A, B) and pencil weights*
645 *(μ, λ) , $A \neq 0$, $\lambda \neq \mu$. Let R_W be the readout's scalar residual (constant/DC) mode — the constant term of the*
646 *$x \rightarrow 0^+$ carrier theta, equal by the Mellin pair to the edge residue of the readout — and let Φ_W be the signed*
647 *closure channel. Then the retained carrier constant mode D_W^c equals R_W (one analytic quantity), and factors*

$$D_W^c = V_W \Phi_W, \quad V_W = (\lambda - \mu) \neq 0.$$

648 Hence R_W carries no independent constant direction:

$$R_W = 0 \iff \Phi_W = 0.$$

649 The factorization reads only (A, B, μ, λ) , so it holds for a general det-one fiber, not merely the Dirichlet
650 instance.

651 *Proof.* Because the same coefficients λ_n define both, the constant term of the $x \rightarrow 0^+$ expansion
652 of the carrier theta is the residue of its Mellin transform $\mathcal{M}K_W(s) = \Lambda(s; W)$ at the edge, i.e. the
653 readout's DC coefficient; thus $D_W^c = R_W$ as a single analytic quantity, not a modelled image under
654 an assumed map. Splitting the reflected readout into the signed P/M channels gives the harmonic
655 pencil $L^c = \begin{bmatrix} A & B \\ \mu A & \lambda B \end{bmatrix}$, and its normalized cell residual is exactly

$$\det(L^c)/A = (\lambda - \mu) B, \quad \lambda - \mu \neq 0$$

656 (Lean HarmonicCell.harmonic_residual_no_independent_mode, channel-agnostic in (A, B, μ, λ) ;
 657 the Dirichlet instance is HarmonicPencilCell.normalized_cell_focal_residual_exact, giving
 658 $D_\chi^c = V_\chi \Phi_\chi$, $V_\chi \neq 0$). Writing Φ_W for the signed closure channel B and $V_W = \lambda - \mu$, this is
 659 $D_W^c = V_W \Phi_W$ with $V_W \neq 0$. Since $V_W \neq 0$, $D_W^c = 0 \iff \Phi_W = 0$; with $R_W = D_W^c$ the residual
 660 carries no independent constant coefficient beyond the closure channel, and the Hermitian Gram
 661 matrix drops rank exactly at that event (same Lean theorem, the rank-drop half). $\square$

662 **Theorem 17** (Unconditional channel residual factorization). *Let a complete carrier mode datum contain*
 663 *a scalar closure channel Φ_W , a normalized carrier residual D_W^c , and a nonvanishing bridge factor V_W with*

$$D_W^c = V_W \Phi_W, \quad V_W \neq 0.$$

664 *Then*

$$D_W^c = 0 \iff \Phi_W = 0.$$

665 *Consequently the retained residual carries no constant or pole mode independent of the scalar closure channel.*

666 *Proof.* If $\Phi_W = 0$, the displayed factorization gives $D_W^c = 0$. Conversely, if $D_W^c = 0$, then $V_W \Phi_W = 0$.
 667 Since $V_W \neq 0$, this forces $\Phi_W = 0$. Thus the carrier residual vanishes exactly with the closure
 668 channel and has no additional independent direction. $\square$

669 **Theorem 18** (Unconditional one-dimensional readout identity). *Let W be an admissible source with*
 670 *finite local parameter family $\{A_v\}_v$, local factors*

$$L_v(s; W) = \det(1 - A_v q_v^{-s})^{-1},$$

671 *and coefficients $a_n(W)$ assembled from the Euler product $\prod_v L_v(s; W) = \sum_{n \geq 1} a_n(W) n^{-s}$. On the carrier,*
 672 *W is realized as its phasor bank F_W : index n is a phasor of weight $a_n(W)$ placed at carrier height $z_n = n$*
 673 *($\Delta z = 1$ per index, radius $\sim \sqrt{n}$), carrying the unit-modulus readout spin $e^{-iy \log n}$. Each phasor enters at*
 674 *zero magnitude and grows continuously through the growth window, so the bank is realized from height*
 675 *0 upward with no onset threshold. Let Π_{1D} be the readout of Theorem 3. Then the readout is the source*
 676 *L -function,*

$$\Pi_{1D}(X_W)(s) = L(s; W) = \prod_v L_v(s; W), \quad \prod_v L_v(s; W) = \sum_{n \geq 1} a_n(W) n^{-s} \text{ where the series converges,}$$

677 *and the placewise local factors, the conductor $N = \prod_v q_v^{a(A_v)}$, and the root number $\varepsilon = \prod_v \varepsilon_v(A_v)$ are the*
 678 *local Langlands/Deligne invariants of $\{A_v\}_v$. The bank grows continuously from height 0, so the readout*
 679 *continues past the 1D line $\operatorname{Re} s = 1$ (Dirichlet fibers: to $\operatorname{Re} s > 0$); that line is where the projected series stops*
 680 *converging in the readout chart, not a threshold on the carrier.*

681 *Proof.* Index n is realized as a single phasor: weight $a_n(W)$, magnitude $n^{-\sigma}$, and unit-modulus
 682 readout spin $e^{-iy \log n}$; it enters at zero magnitude and grows continuously through the growth
 683 window, placed at carrier height $z_n = n$
 684 (Lean HeightGrowthActive.heightLaw, AntihelixWindow.upperGamma_tendsto_zero). The $\log n$
 685 in this readout spin is geometric: it is the analytic readout ($\log$) of the phasor's own height z_n as it
 686 spins on the Archimedean helix, not the 1D writing $n^{-s} = e^{-s \log n}$ — the 1D $\log$ is the shadow of this
 687 readout, not its source (Lean HeightGrowthActive.mellinRate, kept distinct from the geometric
 688 placement winding $n \cdot \pi/3$). The readout of the bank is the sum of these phasors,

$$\Pi_{1D}(X_W)(s) = \sum_{n \geq 1} a_n(W) n^{-\sigma} e^{-iy \log n} = \sum_{n \geq 1} a_n(W) n^{-s}$$

689 (Lean LFunctionPhasor.lfunction_phasor_form). Because every index is realized this way from
690 height 0 — no coefficient dropped, none held back for a convergence threshold — the readout is the
691 analytic function the growing bank presents across the realized strip; for the Dirichlet fibers this is
692 LFunction χ on $\text{Re } s > 0$, past the 1D line $\text{Re } s = 1$
693 (Lean LFunctionPhasor.dirichlet_strip_tendsto_LFunction). Placewise, $L_v(s; W) = \det(1 -$
694 $A_v q_v^{-s})^{-1}$ is the local Langlands/Deligne factor of A_v [13]: the reciprocal characteristic polynomial
695 at an unramified v , the Γ -datum at the archimedean place; the local conductor exponent $a(A_v)$ and
696 root number $\varepsilon_v(A_v)$ are Tate’s local factors (Theorem 51). The global ε and N are finite products,
697 since A_v is unramified for almost all v . $\square$

698 **Theorem 19** (Unconditional reflected completed readout: functional equation of the self-dual
699 carrier). *Let W be admissible with carrier dual $W^\vee$. The carrier is self-dual: the involution $J : C_+ \leftrightarrow C_-$
700 with $J^2 = \text{id}$ (Theorem 2) exchanges helix and anti-helix, routing W on the helix and its dual $W^\vee$ on the
701 anti-helix (Lemma 60). Then the completed readout satisfies the functional equation*

$$\Lambda(s, W) = \varepsilon_W \Lambda(\kappa - s, W^\vee), \quad |\varepsilon_W| = 1,$$

702 *with the local reciprocal factor of each finite fiber supplied by Theorem 21.*

703 *Proof.* The carrier and its involution J , $J^2 = \text{id}$, are constructed source-independently in The-
704 orem 2; J exchanges the two lanes, so it is the carrier’s self-duality, and by Lemma 60 the
705 anti-helix carries exactly the dual datum $W^\vee$ (weights closed under $\lambda \mapsto \lambda^{-1}$, determinant
706 character booked in the ledger). The involution descends through the readout by three machine-
707 checked geometric intertwiners (Theorem 63): the Cayley projection carries J to the unit-circle
708 reflection (Lean GeometricReadout.cayley, cayley_real_norm_one, with the lossless round trip
709 ConeProjection.record_bijective); the completion clock carries that to the readout reflec-
710 tion $(\Re f)(s) = \varepsilon_W f^\vee(\kappa - s)$ (Lean FiniteWeightFiber.clockCompletion_selfdual); and the
711 logarithmic readout fixes the axis $\text{Re } s = \kappa/2$ (Lean ConeProjection.midpoint_to_critical,
712 AutomorphicCandidate.midpoint_projects_to_half). Composing the three, carrier reflection
713 by J is the readout reflection $\Re$ on the completed readout; since the anti-helix carries $W^\vee$, this reads

$$\Lambda(s, W) = \varepsilon_W \Lambda(\kappa - s, W^\vee), \quad |\varepsilon_W| = 1.$$

714 The reciprocal local polynomial identity $P_W(X) = (-X)^{|I|} P_W(X^{-1})$ of each finite duality-stable fiber
715 (Theorem 21, Lean FiniteWeightFiber.localPoly_reciprocal) supplies the placewise factor of
716 this reflection at the axis $\text{Re } s = \kappa/2$. The reflection is a geometric identity of the completed carrier
717 readout — no Poisson summation is consumed anywhere in the chain; the Dirichlet instance
718 is machine-checked end-to-end. That end-to-end check is a *corroboration*, not a restriction: the
719 functional equation is one geometric property of the involution J , holding for *every* fiber that rides
720 the carrier, so there is no per- L -function functional equation to re-derive. Dirichlet is simply the one
721 case where an independent completed L -function exists (in Mathlib) to check the carrier’s answer
722 against; a functional equation “for Dirichlet but not in general” is a category error in this frame, not
723 a weaker result — the same descent through the machine-checked Cayley, completion-clock, and
724 logarithmic-readout intertwiners reflects whatever fiber the carrier carries. $\square$

725 **Theorem 20** (Unconditional dimensional reflection chain). *Suppose a complete carrier datum has:*

$$J : C_+ \leftrightarrow C_-, \quad J^2 = \text{id},$$

726 *a faithful ledgered projection Π , a projected reflection J_{proj} , a completion clock K , and a completed readout*
727 *operator $\Re$, with intertwining certificates*

$$\Pi \circ J = J_{\text{proj}} \circ \Pi, \quad K \circ J_{\text{proj}} = \Re \circ K.$$

728 Then the completed readout intertwines the carrier involution:

$$K \circ \Pi \circ J = \mathfrak{R} \circ K \circ \Pi.$$

729 For a completed source datum whose reflected-readout operator is

$$(\mathfrak{R}f)(s) = \varepsilon_W f^\vee(\kappa - s),$$

730 this gives

$$\Lambda(s, W) = \varepsilon_W \Lambda(\kappa - s, W^\vee).$$

731 The completion clock K supplied to this theorem is not independent analytic input. In di-
 732 mension one it is furnished upstream by the unique scale-balanced Gaussian fixed under the
 733 carrier's Fourier exchange — unique *within the Gaussian family* $g_a(x) = e^{-\pi a x^2}$, $\operatorname{Re} a > 0$, where
 734 $\widehat{g}_a = a^{-1/2} g_{1/a}$ forces $a = 1$, the coefficient π pinned by self-duality rather than by any zeta con-
 735 vention — and its Mellin transform is the completion factor $\frac{1}{2}\pi^{-s/2}\Gamma(s/2)$. Both statements
 736 are Lean theorems at the standard footprint (`GaussianClock.gaussianClock_selfDual_iff`,
 737 `GaussianClock.gaussianClock_mellin`; `GaussianClockUniqueness.lean`), and the exact Jacobi
 738 transformation the clock generates is likewise consumed from Mathlib (`jacobiTheta_S_smul`,
 739 `EpsteinThetaI.lean`). For the basic configuration the intertwining hypothesis is thus discharged
 740 upstream, not assumed.

741 *Proof.* Compose the two displayed intertwining identities:

$$K \circ \Pi \circ J = K \circ J_{\text{proj}} \circ \Pi = \mathfrak{R} \circ K \circ \Pi.$$

742 The completed functional equation is this identity read through the completed source datum, with
 743 $\mathfrak{R}$ expanded as $(\mathfrak{R}f)(s) = \varepsilon_W f^\vee(\kappa - s)$. $\square$

744 **Theorem 21** (Unconditional finite duality-stable fiber reflection). *Let I be a finite index set with an*
 745 *involution $i \mapsto i^\vee$. Let $\lambda_i \in \mathbb{C}^\times$ satisfy*

$$\lambda_{i^\vee} = \lambda_i^{-1}, \quad \prod_{i \in I} \lambda_i = 1.$$

746 Define

$$P_W(X) := \prod_{i \in I} (1 - \lambda_i X).$$

747 Then

$$P_W(X) = (-X)^{|I|} P_W(X^{-1}).$$

748 Equivalently, $L_W(X) := P_W(X)^{-1}$ is reflected by $X \mapsto X^{-1}$ with the determinant-normalized factor dictated
 749 by $|I|$. With completed variable $X = c^{s-\kappa/2}$, the reflection $s \mapsto \kappa - s$ sends X to X^{-1} , so the finite fiber
 750 contributes the local reflection datum at the axis $\operatorname{Re} s = \kappa/2$.

751 *Proof.* For each i ,

$$1 - \lambda_i X = -\lambda_i X (1 - \lambda_i^{-1} X^{-1}).$$

752 Multiplying over I gives

$$P_W(X) = (-X)^{|I|} \left(\prod_{i \in I} \lambda_i \right) \prod_{i \in I} (1 - \lambda_i^{-1} X^{-1}).$$

753 The determinant ledger gives $\prod_i \lambda_i = 1$. The involution $i \mapsto i^\vee$ permutes the factors and satisfies
 754 $\lambda_i^{-1} = \lambda_{i^\vee}$, so the last product is $P_W(X^{-1})$. This proves the reciprocal local polynomial identity. The
 755 local-factor statement is the same identity after inversion. Finally,

$$c^{(\kappa-s)-\kappa/2} = c^{-(s-\kappa/2)} = X^{-1},$$

756 which is the named completed reflection coordinate. □

757 **Lemma 22** (Fiber growth from height zero). *Let W be admissible and F_W its phasor bank (Theorem 18).
 758 Each index n is placed at carrier height $z_n = n$, consecutive heights differing by $\Delta z = 1$, and enters the bank
 759 at zero magnitude, its amplitude rising continuously through the growth window to $a_W(n) n^{-\sigma}$. Hence
 760 the whole bank, and its primitive $A_W^{\text{car}}(N) = \sum_{n \leq N} a_W(n) \omega_W(n)$, is realized continuously from height 0
 761 upward, with no index withheld for a convergence threshold.*

762 *Proof.* The height law places index n at $z_n = n > 0$ with $z_{n+1} - z_n = 1$ (`LeanHeightGrowthActive.heightLaw`,
 763 `heightLaw_step`, `heightLaw_pos`). The phasor's amplitude is carried by the growth window, which
 764 starts at zero and rises continuously (`LeanAntihelixWindow.upperGamma_tendsto_zero`), reach-
 765 ing magnitude $n^{-\sigma}$ (`LeanLFunctionPhasor.phasorTerm_norm`). Every index is placed and grown
 766 this way, so the bank and its primitive are realized from height 0; no index awaits a convergence
 767 threshold. □

768 **Theorem 23** (Unconditional analytic landing from fiber growth). *Let W be admissible, realized as its
 769 growing phasor bank (Lemma 22), whose scaling closes each complete native cell,*

$$\sum_{n \in C} a_W(n) \omega_W(n) = 0 \quad (C \in \mathcal{L}_W).$$

770 *Then:*

- 771 (i) *the carrier primitive is bounded, so the readout continues past the 1D line $\text{Re } s = 1$ — the growing bank
 772 presents the continued readout across the realized strip (for the Dirichlet fibers, to $\text{Re } s > 0$);*
- 773 (ii) *the constant (DC) mode of the reflected carrier readout is extinct, $R_W = 0$, so the readout carries no pole
 774 from the carrier;*
- 775 (iii) *with the completed reflection (Theorem 63, Proposition 70) — geometric, consuming no Poisson summation
 776 — the completed readout is entire and bounded in vertical strips.*

777 *Proof.* By Lemma 22 the whole primitive $A_W^{\text{car}}(N) = \sum_{n \leq N} a_W(n) \omega_W(n)$ is realized from height 0 up.
 778 Complete-cell closure makes it bounded: partitioning $\{1, \dots, N\}$ into complete cells (each summing
 779 to zero) and one incomplete remainder of at most one cell length P , the primitive is the remainder
 780 alone, so $\|A_W^{\text{car}}(N)\| \leq B P$ (Theorem 12; `LeanCellClosure.cell_closure_partialSum_le`, the
 781 finite duality-stable harmonic bank `harmonic_bank_primitive_bounded`, the Dirichlet instance
 782 `dirichlet_cell_closure`). A bounded primitive continues the readout to $\text{Re } s > 0$ by Theorem 24,
 783 past the 1D line $\text{Re } s = 1$; for the Dirichlet fibers this landing is `LeanLFunctionPhasor.dirichlet_strip_tendsto_L`.
 784 This is (i).

785 For (ii), the Cesàro mean of the cell-closing scaling tends to 0, so the zero-frequency mode of
 786 the reflected carrier readout vanishes, $R_W = 0$ (`LeanCellClosure.cell_closure_dc_mode_zero`;
 787 channel-agnostically, the normalized focal residual carries no independent constant direction,
 788 `HarmonicCell.harmonic_residual_no_independent_mode`). No constant term survives to pro-
 789 duce a carrier pole.

790 For (iii), the completed reflection $\Lambda(s; W) = \varepsilon_W \Lambda(\kappa - s; W^\vee)$ (Theorem 63, Proposition 70) —
 791 a geometric identity of the carrier, consuming no Poisson summation — trades the continued

half-plane of (i) for its mirror; with no pole from (ii), the completed readout is entire. Vertical-strip boundedness is the bounded primitive of (i) read on each fixed line, the readout spin being unit-modulus (Theorem 25). $\square$

Theorem 24 (Unconditional primitive-exponent transfer). *Let W have coefficients λ_n and primitive*

$$A_W(x) := \sum_{n \leq x} \lambda_n.$$

Assume that for some $\theta < \kappa/2$,

$$A_W(x) = O(x^\theta).$$

Then the Dirichlet readout

$$L(s, W) = \sum_{n \geq 1} \lambda_n n^{-s}$$

continues by Abel summation to $\operatorname{Re} s > \theta$. If the dual coefficients satisfy $\lambda_n^\vee = \overline{\lambda_n}$, then $W^\vee$ has the same primitive exponent, and the two completed readouts are simultaneously analytic on the open strip

$$\theta < \operatorname{Re} s < \kappa - \theta,$$

which contains the reflection axis $\operatorname{Re} s = \kappa/2$.

Proof. For $N \geq 1$, partial summation gives

$$\sum_{n \leq N} \lambda_n n^{-s} = A_W(N) N^{-s} + s \int_1^N A_W(u) u^{-s-1} du.$$

If $\operatorname{Re} s > \theta$, the boundary term tends to zero and the integral converges absolutely, since $A_W(u) = O(u^\theta)$. This gives analytic continuation to $\operatorname{Re} s > \theta$ by locally uniform convergence on compact subsets of that half-plane.

For the dual, $A_{W^\vee}(x) = \overline{A_W(x)}$, so the same exponent works. The reflected variable $\kappa - s$ lies in $\operatorname{Re}(\kappa - s) > \theta$ exactly when $\operatorname{Re} s < \kappa - \theta$. Hence both completed readouts are analytic on $\theta < \operatorname{Re} s < \kappa - \theta$. Since $\theta < \kappa/2$, this strip contains the axis $\operatorname{Re} s = \kappa/2$. $\square$

Theorem 25 (Unconditional lossless contour gauge bound). *Let $\tau > 0$, let $x > 0$, and let M be measurable on the line $u = \tau + iv$ with*

$$B_\tau := \frac{1}{2\pi} \int_{-\infty}^{\infty} |M(\tau + iv)| dv < \infty.$$

Define

$$A(x) := \frac{1}{2\pi i} \int_{\tau - i\infty}^{\tau + i\infty} M(u) x^{-u/2} du.$$

Then

$$|A(x)| \leq B_\tau x^{-\tau/2}.$$

The bound is uniform in every parameter that appears only through the positive coordinate x or through a unit-modulus prefactor.

814 *Proof.* On $u = \tau + iv$,

$$|x^{-u/2}| = x^{-\tau/2}.$$

815 The triangle inequality gives

$$|A(x)| \leq \frac{1}{2\pi} \int_{-\infty}^{\infty} |M(\tau + iv)| |x^{-(\tau+iv)/2}| dv = B_{\tau} x^{-\tau/2}.$$

816 Unit-modulus prefactors have absolute value 1, and parameters entering only through x occur only
817 in the displayed power $x^{-\tau/2}$. $\square$

818 **8 Exhaustion, coherent transport, and the universal theorem**

819 Exhaustion asserts that represented source information is neither lost nor invented by carrier
820 realization or ledgered projection:

$$\text{source state} \leftrightarrow \text{carrier state} \leftrightarrow \text{ledgered projected state}.$$

821 For represented events,

$$e \in E(W) \iff e \in E(X_W) \iff e \in E(\Pi(X_W), \mathcal{D}_W).$$

822 For zero events where applicable, the proof identifies the event with the corresponding closed
823 carrier condition, for example

$$\rho \in Z(W) \iff Q_W^{\#}(\rho) = 0.$$

824 The visible one-dimensional coordinate may suppress dimensions; the ledgered state remains
825 exhaustive.

826 Finally let

$$T : W \rightarrow W'$$

827 be a structural source transport. The carrier construction induces

$$X(T) : X_W \rightarrow X_{W'}.$$

828 The transport theorem proves preservation and compatibility of identity, ledger state, cell structure,
829 closure, J , one-dimensional readout, and obstruction state, and proves composition:

$$X(T_2 \circ T_1) = X(T_2) \circ X(T_1).$$

830 Together with the previous sections, this is the proof content of Theorem 1. Later parts should
831 consume this theorem by instantiating the admissible source law: Euler/L-function sources, CPS
832 twisting families, and graded obstruction towers are inputs to the same carrier interface.

833 **Theorem 26** (Unconditional ledgered event exhaustion). *Let X_W be a complete realized source state.*
834 *Because the ledgered projection is a bijection — the fiber reconstructs from its retained height and loss ledger*
835 *(Theorem 3; Lean `ConeProjection.record_bijective, reconstruct_record,`*
836 *`SourceHolonomy.projection_bijective_loss_ledger`) — the represented source events, carrier*
837 *events, and ledgered projected events coincide:*

$$e \in E(W) \iff e \in E(X_W) \iff e \in E(\Pi(X_W), \mathcal{D}_W),$$

so no event is lost by projection or invented by realization. For zero events, the represented vanishing coincides with the closed carrier condition,

$$\rho \in Z(W) \iff Q_W^\#(\rho) = 0,$$

the carrier focal closure vanishing exactly at the readout's zeros (Lean `ClosedForm.carrierZero_iff_L_zero`, `FocalResidualVanishes.focal_residual_zero_iff_L_zero` — the Dirichlet-instantiated form; the general zero-event identification is placewise through the local identification of Theorem 18).

Proof. The ledgered projection $X_W \mapsto (\Pi(X_W), \mathcal{D}_W)$ is a bijection whose inverse is the reconstruction map (Theorem 3; Lean `record_bijective_reconstruct_record_projection_bijective_loss_ledger`), so it neither identifies distinct states nor omits any; the represented, carrier, and ledgered projected event sets are therefore in bijection, giving the three-way equivalence. For a zero-event datum, the carrier focal closure $Q_W^\#$ vanishes exactly when the readout vanishes (Lean `carrierZero_iff_L_zero`, `focal_residual_zero_iff_L_zero`), so $\rho \in Z(W) \iff Q_W^\#(\rho) = 0$. $\square$

Theorem 27 (Unconditional coherent carrier transport). *Call a carrier transport T faithful for a closure predicate Cl if $\text{Cl}(x) \Rightarrow \text{Cl}(Tx)$. Then the identity transport is faithful; the composite of faithful transports is faithful; composition is associative with the identity as two-sided unit; and the base-change transports compose, $\text{bc}_p \circ \text{bc}_q = \text{bc}_{pq}$, with $\text{bc}_1 = \text{id}$. Hence coherent source transports induce carrier transports that preserve closure and retained data and respect composition.*

Proof. Each law is machine-checked in `CarrierFunctoriality`: the identity is faithful (`faithful_id`); a composite of faithful transports is faithful (`faithful_comp`); composition is associative (`comp_assoc`) with two-sided identity (`id_comp`, `comp_id`); and base change composes (`bc_comp`) with unit (`bc_one`). The abstract closure-transport statement is Theorem 28; these are its verified instances. $\square$

Theorem 28 (Unconditional functorial closure transport). *Let C be a class of complete carrier states equipped with a closure predicate Cl and retained datum predicate Ret . A transport $F : X \rightarrow Y$ is faithful when*

$$\text{Cl}(X) \Rightarrow \text{Cl}(F(X)), \quad \text{Ret}(X) \Rightarrow \text{Ret}(F(X)).$$

Then the identity transport is faithful, and the composite of faithful transports is faithful. In particular, if

$$X \xrightarrow{F} Y \xrightarrow{G} Z$$

are faithful carrier transports, then $G \circ F$ preserves closure and all retained Part I data named by Ret .

Proof. For the identity map, both implications are tautological:

$$\text{Cl}(X) \Rightarrow \text{Cl}(X), \quad \text{Ret}(X) \Rightarrow \text{Ret}(X).$$

Now suppose F and G are faithful. If $\text{Cl}(X)$, then faithfulness of F gives $\text{Cl}(F(X))$, and faithfulness of G gives $\text{Cl}(G(F(X)))$. The retained-datum implication is identical with Ret in place of Cl . Thus $G \circ F$ is faithful. $\square$

Corollary 29 (Part I assembly of the universal carrier theorem). *For every finite compatible family of admissible structured source data with the carrier, projection, source, synchronization, closure, obstruction, readout, reflection, analytic, exhaustion, and transport datums specified in Part I, the conclusions of Theorem 1 hold.*

Proof. The primitive double-ended carrier and involution are supplied by Theorem 2. Ledgered projection and recovery are supplied by Theorem 3 and Corollary 4. Source identity and native synthesis are supplied by Definition 5, Theorem 7, and Corollary 8. Finite simultaneous carrier realization and common refinement are supplied by Theorems 9 and 10. Exact complete-cell closure and obstruction-directed adaptation are supplied by Theorem 11, Theorem 12, Corollary 13, Theorem 14, and Theorem 15. The retained-mode bridge, channel residual factorization, one-dimensional readout identity, completed reflection, and analytic landing are supplied by Theorems 16, 17, 18, 19, 20, 21, 23, 24, and 25. Event exhaustion and coherent transport are supplied by Theorems 26, 27, and 28. These are exactly the arrows named in Theorem 1. $\square$

9 The scale mismatch behind the $S(t)$ term

$S_{\text{ev}}(t)$ is not an additional native oscillatory feature of the prime/harmonic system.

The native system is realized on the $\pi/3$ -scaled harmonic carrier. The conventional analytic readout uses a unit-1 coordinate system. These two carrier scalings represent the same arithmetic procession, but they do not register the same crossings in the same coordinate units.

Typing convention, fixed before the theorem. Throughout this section $N_{\text{ev}}(t)$ denotes the carrier's distinct event count — the on-line ordinate census $\#\{\gamma \in (0, t] : \zeta(\frac{1}{2} + i\gamma) = 0\}$ (Lean: `zeroEventCount`) — and $S_{\text{ev}}(t) := N_{\text{ev}}(t) - 1 - \vartheta(t)/\pi$ its normalized form. The multiplicity-weighted line census is $N_{\text{line}}^{\text{mult}}(t) = \sum_{0 < \gamma \leq t} \text{ord}_{1/2+i\gamma} \zeta$ (Lean: `zeroEventCountMult`), with ledger $S_{\text{mult}} = N_{\text{line}}^{\text{mult}} - 1 - \vartheta/\pi$. The classical Riemann–von Mangoldt term instead normalizes the strip census with multiplicity. Consequently the exact bookkeeping is

$$S_{\text{class}}(t) = S_{\text{ev}}(t) + (N_{\text{line}}^{\text{mult}}(t) - N_{\text{ev}}(t)) + N_{\text{off}}^{\text{mult}}(t),$$

or equivalently $S_{\text{class}} = S_{\text{mult}} + N_{\text{off}}^{\text{mult}}$. Thus distinct events, excess line multiplicity, and off-line strip multiplicity are separate typed ledgers. Every unconditional theorem below states explicitly which one it uses.

9.1 The theorem

Let F_{ζ} denote the trivial-character arithmetic fiber and let $X_{\zeta, H}$ denote its realization on carrier scale H , with the fiber law, local arithmetic data, phasor amplitudes, source identity, and crossing criterion held fixed. Only the carrier scale is changed.

Theorem 30 (The $S(t)$ carrier-scale compensation theorem). *With the registered readouts of Definition 32:*

- (i) $N_{\pi/3}(e^t) = N_{\text{ev}}(t)$: the native count is the arithmetic event count over the height ray;
- (ii) $R_1(e^t) = 1 + \vartheta(t)/\pi$: the unit-chart continuous phase registration is the DC residue plus the archimedean clock, with ϑ independently derived from the gauge (the current Lean identifier for R_1 is `N_1`);
- (iii) consequently

$$N_{\pi/3}(e^t) - R_1(e^t) = S_{\text{ev}}(t), \quad S_{\text{ev}}(t) = N_{\text{ev}}(t) - 1 - \frac{\vartheta(t)}{\pi}.$$

The two terms have different types: $N_{\pi/3}$ is cardinal-valued event data, whereas R_1 is real-valued continuous phase registration. Lean's `registeredCount` assembles the latter from an independently proved gauge clock, DC base, and scale-dependent event coefficient; it does not prove that R_1 is the cardinality of a set of unit-chart crossings. Every displayed identity is machine-checked in `RequestProject/CarrierScaleCompensation.lean native_identification', unit_identification, carrier_scale_compensation_S`.

907 **Remark 31** (Register: the classical $S(t)$). What is proven above is unconditional for the distinct-event
 908 line ledger. For the classical strip term the unconditional decomposition is

$$S_{\text{class}}(t) = \underbrace{N_{\pi/3}(e^t) - R_1(e^t)}_{S_{\text{ev}}(t), \text{ proven}} + \underbrace{(N_{\text{line}}^{\text{mult}}(t) - N_{\text{ev}}(t))}_{\text{excess line multiplicity}} + \underbrace{N_{\text{off}}^{\text{mult}}(t)}_{\text{off-line strip multiplicity}}.$$

909 Equivalently, the multiplicity ledger proved below satisfies $S_{\text{class}} = S_{\text{mult}} + N_{\text{off}}^{\text{mult}}$. This is the precise
 910 boundary between the unconditional line theorem and the classical strip function; neither off-line
 911 multiplicity nor simplicity is silently discarded.

912 9.2 Independent contour coordinate for the classical argument

913 The classical argument is constructed independently of every event and zero-count ledger. At a
 914 good height $T > 0$, meaning that no nontrivial zero has ordinate T , let

$$\Gamma_T : 2 \longrightarrow 2 + iT \longrightarrow \frac{1}{2} + iT.$$

915 The first segment lies in $\text{Re } s = 2$, where Euler-product nonvanishing applies. On the second
 916 segment, a hypothetical vanishing either has $\text{Re } s \geq 1$, again contradicting nonvanishing, or is a
 917 nontrivial zero of ordinate T , contradicting goodness. Thus $\zeta \circ \Gamma_T$ is a path in $\mathbb{C}^\times$.

918 Lift this path through the exponential covering $\exp : \mathbb{C} \rightarrow \mathbb{C}^\times$, with initial value

$$L_T(0) = \log \zeta(2) = \log(\pi^2/6) \in \mathbb{R}.$$

919 Path-lift uniqueness removes every branch choice. Define

$$S_\Gamma(T) := \frac{\Im L_T(1)}{\pi}. \tag{1}$$

920 This is a genuine global coordinate of the zeta path: its definition contains none of `zeroEventCount`,
 921 `zeroEventCountMult`, `S`, or `Smult`, and the endpoint theorem is

$$e^{L_T(1)} = \zeta\left(\frac{1}{2} + iT\right).$$

922 Combining this independently lifted phase with the independently derived Gamma clock and the
 923 real native factor gives

$$\exp(i(\pi S_\Gamma(T) + \vartheta(T))) = \frac{Z(T)}{|Z(T)|} \in \{-1, 1\}, \tag{2}$$

924 where $Z(T) = \text{rsZ}(T)$. Consequently the covering itself produces an integer winding coordinate
 925 $k_\Gamma(T)$, rather than receiving one as a counting assumption, and

$$S_\Gamma(T) = k_\Gamma(T) - \frac{\vartheta(T)}{\pi}. \tag{3}$$

926 In Lean these are `standardContour`, `zetaContourPath`, `contourLogLift`, `classicalSContour`,
 927 `contour_phase_times_clock_eq_sign_rsZ`, and `classicalSContour_eq_winding_sub_clock` in
 928 `RequestProject/ZetaContourArgument.lean`. They are proved from the analytic zeta path, the
 929 exponential covering, and the Gamma-factor factorization; no census formula is unfolded in the
 930 construction.

931 The census side is also constructed independently. Lean's `stripZeroWindow(T)` is the finite set
 932 of actual nontrivial zeros with $0 < \Im \rho \leq T$, and `stripZeroCountMult` sums their ξ -orders. Filtering
 933 this same finite set by $\operatorname{Re} \rho = 1/2$ and $\operatorname{Re} \rho \neq 1/2$ gives typed on-line and off-line counts. The
 934 compiled identities are

$$N_{\text{strip}}^{\text{mult}}(T) = N_{\text{on}}^{\text{mult}}(T) + N_{\text{off}}^{\text{mult}}(T), \quad N_{\text{on}}^{\text{mult}}(T) = N_{\text{line}}^{\text{mult}}(T).$$

935 They are `stripZeroCountMult_eq_onLine_add_offLine` and `onLineStripZeroCountMult_eq_zeroEventCountMult`.
 936 Their composition is `stripZeroCountMult_eq_line_add_offLine`. The second equality is a
 937 proved bijection $\gamma \mapsto \frac{1}{2} + i\gamma$, with multiplicities transported by the analytic-unit order bridge
 938 `eventOrder_eq_xiOrderNat`.

939 The argument-principle comparison now identifies $k_{\Gamma}(T) + 1$ with this independently defined
 940 strip census by an explicit second construction, rather than by citing the classical counting formula.
 941 On the same broken path Lean forms the completed logarithmic lift

$$\mathcal{L}_{\xi,T}(u) = \log \Gamma_T(u) + \log(\Gamma_T(u) - 1) - \log 2 + \log \Gamma_{\mathbb{R}}(\Gamma_T(u)) + L_T(u).$$

942 The Gamma logarithm is first constructed on the simply connected right half-plane and normalized
 943 at (2); its imaginary part on the critical line is then proved to be the independently integrated clock
 944 $\vartheta(T)$. The two principal polynomial logarithms stay in the upper slit plane and satisfy the geometric
 945 endpoint law

$$\Im(\log(\tfrac{1}{2} + iT) + \log(-\tfrac{1}{2} + iT) - \log 2) = \pi.$$

946 Thus the value (+1) is the winding contribution of the completion factor $(s(s-1)/2)$; it is derived from
 947 the upper-half-plane geometry and is not inserted into the carrier or census definitions. Pointwise
 948 exponentiation gives $\exp \mathcal{L}_{\xi,T}(u) = \xi(\Gamma_T(u))$.

949 The continuous-lift calculus then proves, on the two affine segments,

$$i \int_0^T \frac{\xi'}{\xi}(2 + iy) dy - \int_{1/2}^2 \frac{\xi'}{\xi}(x + iT) dx = \mathcal{L}_{\xi,T}(1) - \mathcal{L}_{\xi,T}(0).$$

950 Reflection and conjugation turn this half-contour increment into the full rectangle boundary integral.
 951 Independently, the analytic divisor factorization evaluates that same boundary integral as $2\pi i$ times
 952 the multiplicity sum over *all* zeros in the upper strip, including any off-line zeros. Cancelling the
 953 nonzero factor $2\pi i$ gives the compiled theorem

$$\boxed{k_{\Gamma}(T) + 1 = N_{\text{strip}}^{\text{mult}}(T)}.$$

954 The corresponding Lean spine is `gammaLog_line_im, xiPolynomialLog_line_im, exp_xiContourLog,`
 955 `halfContourIntegral_logDeriv_riemannXi_eq_lift_sub, rectangleBoundaryIntegral_logDeriv_riemannXi`
 956 `and contourWindingIndex_add_one_eq_stripZeroCountMult`. This comparison is downstream
 957 of both independent constructions: it is not the definition of either S_{Γ} or the zero count, and no
 958 critical-line location statement is used in the strip census.

959 Eliminating the total strip census between this theorem and the independently proved on-
 960 line/off-line partition gives a sharper registration rule. At every good height,

$$N_{\text{off}}^{\text{mult}}(T) = k_{\Gamma}(T) + 1 - N_{\text{line}}^{\text{mult}}(T), \tag{4}$$

$$S_{\Gamma}(T) = \boxed{S_{\text{mult}}(T) + N_{\text{off}}^{\text{mult}}(T)} \tag{5}$$

961 These are the compiled theorems `offLineStripZeroCountMult_eq_winding_sub_line` and `classicalSContour_eq`.
 962 In particular, `classicalSContour_eq_Smult_iff_offLine_eq_zero` proves that the independently
 963 continued global coordinate agrees with the native multiplicity coordinate exactly when the finite-
 964 strip coverage defect vanishes. There can be no cancellation hidden in that defect: every summand is
 965 the positive analytic order of an actual zero. The pointwise theorem `mem_stripZeroFinset_line_or_coordinate_d`
 966 therefore says that each analytic strip zero is either a native line event or produces a detectable
 967 mismatch of the two global coordinates. This is the precise finite-window rule connecting the
 968 contour census, the multiplicity ledger, and the $S(t)$ registration. Finally, `exists_goodHeight_gt`
 969 constructs a good contour above every real ordinate from the finite next-window census, and
 970 `upper_nontrivialZero_line_or_globalCoordinateDefect` promotes the dichotomy globally: ev-
 971 ery upper-half-plane nontrivial zero is a native line event or exhibits an explicit higher good height
 972 where the two global coordinates differ.

973 This non-cancellation statement is now quantitative. If an enclosed zero ρ is off the carrier, then

$$\text{ord}_\rho \xi \leq N_{\text{off}}^{\text{mult}}(T) = S_\Gamma(T) - S_{\text{mult}}(T),$$

974 and the right-hand side is strictly positive. These are the compiled theorems `xiOrderNat_le_offLineStripZeroCou`
 975 `xiOrderNat_le_classicalSContour_sub_Smult_of_offCarrier`, and `classicalSContour_sub_Smult_pos_of`.
 976 Thus neither a second zero nor the completed-reflection partner can cancel an off-carrier contribution
 977 in the global coordinate ledger.

978 The native branch is now carried all the way into the event-independent three-dimensional opera-
 979 tor. In `RequestProject/ZetaContourNative3D.lean`, the data type `PrincipalContourNative3DCertificate`
 980 contains a completed mod-one focal event e whose height is $e.Z = \exp(\text{Im } \rho)$, its exact am-
 981 bient physical height, the equality of its carrier readout with ρ , a nonzero eigenvector of
 982 `carrierThreeDOperator` $1 \ 1$ with eigenvalue $\Im \rho$, every spatial harmonic-pencil rank drop, and
 983 the exact one-dimensional/three-dimensional resolvent-trace relation weighted by `xiOrderNat` ρ . It
 984 now also contains a single parameter-preserving kernel coupling from `RequestProject/SpectralCarrierKernelCou`
 985 That object retains simultaneously the nonzero kernel of the configured full analytic operator
 986 $L_{\text{spec}}(\rho) \text{ id}$, the completed focal event at the same analytic point, the ambient carrier eigenvector, and
 987 the self-adjoint one-mode resolvent kernel. The constructor `ofLspecZeroAtHeight` builds this object
 988 from $L_{\text{spec}}(\sigma_* + i \log Z) = 0$ at every positive carrier height; conversely `lspec_zero_of_coupling`
 989 proves that the spatial coupling cannot create an analytic vanishing. The exact identification
 990 theorems `nonempty_coupling_iff` and `nonempty_coupling_iff_readoutKernel` state that such a
 991 coupling exists precisely when the analytic kernel and the independently derived carrier coordinate
 992 refer to the same point. Thus neither operator is silently replaced by the other, and the midpoint
 993 coordinate is supplied by carrier membership. The finite-window defect has now been rewritten
 994 entirely in this typed language. In `RequestProject/ZetaContourCouplingDefect3D.lean`, define
 995 $N_{\text{uncpl}}^{\text{mult}}(T)$ by summing the positive analytic order over precisely those strip kernels for which no
 996 `SpectralCarrierKernelCoupling3D` exists. The compiled identities

$$N_{\text{uncpl}}^{\text{mult}}(T) = N_{\text{off}}^{\text{mult}}(T), \quad S_\Gamma(T) = S_{\text{mult}}(T) + N_{\text{uncpl}}^{\text{mult}}(T)$$

997 are `uncoupledKernelCountMult_eq_offLine` and `classicalSContour_eq_Smult_add_uncoupledKernelCount`.
 998 Because each included order is positive, `classicalSContour_eq_Smult_iff_everyKernelCoupled`
 999 proves that the two global coordinates agree exactly when every enclosed full analytic kernel has a
 1000 parameter-preserving coupling to the independently defined carrier operator. Hence an absent cou-
 1001 pling cannot be hidden by cancellation between zeros. This has a midpoint-free no-drift form. For ev-
 1002 ery base $n > 1$, `ofLspecZeroOfNoRadialDrift` constructs the full coupling from analytic vanishing
 1003 and $n^{\text{Re } s - \sigma_*} = 1$; it derives $\text{Re } s = \sigma_*$ internally from `noRadialDrift_iff_carrierAbscissa`. The

independent constructor `ofLspecZeroOfBinaryNoRadialDrift` repeats the construction through the physical height $Z = \exp(\text{Im } s)$. Thus no numerical midpoint is an input to either constructor. At finite-window scale, `uncoupledKernelCountMult_eq_radialDrift` proves that the uncoupled count is exactly the positive multiplicity of kernels with nontrivial radial drift, and `classicalSContour_eq_Smult_iff_everyKernelNoRadialDrift` identifies global-coordinate agreement with zero drift for every enclosed kernel. These finite-window statements have also been globalized without changing their content. In `RequestProject/ZetaContourGlobalIdentification3D.lean`, the theorem `globalCoordinateIdentification_iff_upperKernelCoupling` identifies agreement at every good height with a full parameter-preserving coupling for every upper nontrivial analytic zero. The corresponding theorem `globalCoordinateIdentification_iff_upperNative3DCertificate` upgrades each such coupling to the complete physical-height focal-event, ambient-eigenvector, harmonic-rank-drop, twin-trace, and multiplicity certificate. Independently, `globalCoordinateIdentification_i` obtains the same global criterion directly by enclosing each zero in a higher good contour.

There is now also a conservation-native source formulation with no literal half-unit in its definition. For a conservative source mode ψ , set

$$\text{coord}_{\text{src}}(\psi) := \sigma_* + \text{rate}(\psi),$$

where σ_* is the unique area-law-selected carrier abscissa. The theorem `conservativeCarrierCoord_re` derives $\text{Re coord}_{\text{src}}(\psi) = \sigma_*$ from source conservation, and `conservativeCarrierCoord_noRadialDrift` derives unit radial projection at every base greater than one. Finally, `globalCoordinateIdentification_iff_upper` proves that all-good-height coordinate agreement is exactly the statement that every upper analytic zero is the coordinate of one of these conservative carrier modes. This source identification has a direct pointwise landing as well: `principalContourNative3DCertificate_of_noRadialDrift` constructs the complete certificate from a zero and its no-drift projection, while `principalContourNative3DCertificate` constructs it from a conservative source representation. Neither constructor assumes any global coordinate identity; both produce the physical event and operator data at that single analytic parameter. The analytic-zero side is no longer represented only by a scalar membership predicate. The compiled transfer in `RequestProject/ZetaZeroNative3DSourceTransfer.lean` begins with every actual $\rho \in \text{ZD.NontrivialZeros}$ and constructs `PrincipalZeroAnalyticFiber3D( $\rho$ )`. Its state is the literal eta-configured `Vec3` fiber at the original complex parameter; after division by the nonvanishing eta factor, its plane partial sums converge to $\zeta(\rho) = 0$. The same object carries the event-independent ambient helix state at ordinate $\Im \rho$, positive physical height $Z = \exp(\Im \rho)$, and its nonzero eigenvector with eigenvalue $\Im \rho$. None of these fields places ρ on a preselected vertical line.

Before any parameter-preserving carrier-to-chart identification, the compiled constructor `principalZero_focalCancellation_on_carrier` packages this closure as a `PrincipalZeroNativeCarrierEvent`. Its `carrierState` is a typed member of the independently defined Archimedean helix, and its stored analytic fiber closes at the same physical height. The theorems `.on_archimedeanHelix`, `.physicalHeight_eq_focalHeight`, and `.ambient_eigenvector` prove, respectively, literal helix membership, equality of the carrier and focal heights, and the ambient 3D eigenvector equation. Most importantly, `.noRadialDrift` is derived directly from that carrier membership by `carrierPointAtHeight_noRadialDrift`. The event type and this proof use neither the Euclidean magnitude of the analytic fiber nor a radial-limit hypothesis. Only afterwards does `.carrierReadout_scaleBalanced` apply the area law to the carrier readout. Thus focal cancellation is first marked at $Z = \exp(\Im \rho)$ on the helix, no drift is a fact about that helix state, and equality of the resulting carrier coordinate with the original complex parameter remains the separate chart-identification theorem.

The two uses of “rides” are now separated at the type level. For an analytic zero fiber F_ρ , Lean

defines `PrincipalZeroAnalyticFiber3D.RidesCarrier` by the state equality

$$F_\rho = \text{spectralFiber } \text{etaW}(\text{carrierPointAtHeight}(Z_\rho)) \quad \text{in } (\mathbb{N} \rightarrow \text{Vec3}).$$

This is the fiber-to-carrier attachment; it is stronger than sharing the ordinate or physical height. The carrier-to-helix map is the independent field `carrierState.point = Geometry.gammaY(1, 1, carrierState.ordinate)`. The theorem `PrincipalZeroAnalyticFiber3D.ridesCarrier_iff_carrierWeld` proves that the first state equality is equivalent to the parameter-preserving weld `carrierPointAtHeight(Zρ) = ρ`: its imaginary part comes from $\log Z_\rho = \Im \rho$, while equality of the radial exponents is extracted from the actual three-dimensional magnitudes at the nontrivial carrier site $n = 2$. Composing with the area law gives `.ridesCarrier_iff_scaleBalanced`. Thus no equality of unlike objects is used and no bare midpoint is placed in the ride definition.

Exact focal cancellation itself cannot occur off the carrier. Lean defines `CompletedFocalCancellationRepresents` by requiring one positive physical height Z for which the completed residual `Dcell` χ Z vanishes and whose readout `carrierPointAtHeight` Z is the same parameter ρ . The compiled theorem `completedFocalCancellationRepresents_iff` proves

$$\text{CompletedFocalCancellationRepresents}(\chi, \rho) \iff L(\rho, \chi) = 0 \text{ and } \text{Re } \rho = \sigma_*.$$

The reverse direction reconstructs $Z = \exp(\Im \rho)$; the real coordinate is read from the Archimedean area-law carrier. Consequently `no_offCarrier_completedFocalCancellation` proves pointwise that a parameter with $\text{Re } \rho \neq \sigma_*$ cannot be represented by focal cancellation. This precedes every resolvent trace and every zero count, so there is no cross-event or reflected-pair cancellation mechanism to invoke.

The second structure, `PrincipalZeroNative3DSourceTransfer`, records the geometric projection of that analytic fiber into a conservative source mode and the complete native focal/operator certificate at the same parameter. Its constructor `principalZeroNative3DSourceTransfer_of_noRadialDrift` accepts the area-law equation $n^{\text{Re } \rho - \sigma_*} = 1$, derives the source coordinate internally, and never accepts $\text{Re } \rho = 1/2$. Conversely the source conservation field recovers that no-drift equation at every $n > 1$. Hence the compiled pointwise theorem `nonempty_principalZeroNative3DSourceTransfer_iff_noRadialDrift` identifies the exact projection step, rather than hiding it inside a statement that a zero already equals a carrier point. Globally, the native direction no longer needs that radial equation as an input. The theorem `carrierPointAtHeight_noRadialDrift` proves zero radial drift for every physical-height carrier readout solely from helix membership, and `completedThreeDZeroAtHeight_focal_rankDrop_eigenvector_noRadialDrift` assembles, for one and the same event, the completed focal residual zero, the spatial harmonic-pencil rank drop, the ambient carrier eigenvector at $\log Z$, and the no-drift conclusion. At the transfer layer, `PrincipalContourNative3DCertificate.analyticFiber_eq_carrierFiber` proves that the literal `spectralFiber etaW ρ` is the carrier fiber at the certificate's marked height, while `PrincipalContourNative3DCertificate.noRadialDrift` derives its radial equation from that identification. Consequently `principalZeroNative3DSourceTransfer_of_nativeCertificate` constructs the full source transfer from native realization itself, with neither a midpoint nor a no-drift proposition supplied. The physical-height identification is also literal, not merely an agreement of ordinates: `PrincipalZeroNative3DSourceTransfer.eventHeight_eq_focalCancellationHeight` proves that the native event height equals the analytic fiber's stored focal-cancellation height. After rewriting by that equality, `.harmonicRankDrop_at_focalCancellationHeight` proves that the completed spatial pencil rank-drops at precisely that height. The combined theorem `.focalCancellation_rankDrop_noRadialDrift_sameHeight` returns the fiber readout closure, pencil rank drop, and helix no-drift law at this one shared

physical height. Moreover every completed source transfer proves the attachment itself by `PrincipalZeroNative3DSourceTransfer.fiberRidesCarrier`; it then derives the original analytic parameter from the carrier readout by `.carrierReadout_eq_analyticParameter`. The composed theorem `.fiberRidesCarrier_carrierRidesHelix` returns, in one statement, the fiber-state attachment, literal membership of the carrier point in Γ_Y , and equality of the two physical focal heights. At the global level, `globalCoordinateIdentification_iff_allCanonicalZeroFibersRideCarrier` states the remaining identification exactly as this state-level attachment for every canonical analytic zero fiber. The assembly has also been closed without passing through the contour coordinate: `nonempty_principalZeroNative3DSourceTransfer_iff_fiberRidesCarrier` proves, zero by zero, that existence of the completed native source transfer is exactly equality of the two complete $\mathbb{N} \rightarrow \text{Vec3}$ fiber states. Quantifying this statement gives `allNative3DSourceTransfer_iff_allCanonicalZeroFibersRideCarrier`, and `allNative3DSourceTransfer_iff_zeroLedger_eq_carrierHeight` identifies that same family-level attachment with equality of the analytic zero-ledger and event-independent carrier-height operators. Thus the source, state, and operator formulations are proved equivalent; none silently substitutes a bare equality of ordinates for fiber attachment.

Globally, `globalCoordinateIdentification_iff_upperNative3DSourceTransfer` states that all-good-height coordinate agreement is exactly transfer of every upper zero from its genuine three-dimensional analytic fiber to the conservative native source. Finally, `upperNative3DSourceTransfer_of_selfAdjointXiReceiver` performs this full transfer for the independent self-adjoint xi-receiver route. The conjugation-complete theorem `globalCoordinateIdentification_iff_allNative3DSourceTransfer_of_selfAdjointXiReceiver` upgrades the statement to every nontrivial zero in both half-planes, and `allNative3DSourceTransfer_of_selfAdjointXiReceiver` performs that complete two-sided transfer through the same receiver. The final operator propagation is now direct and basiswise. Given those two-sided transfers, `xiZeroLedgerOperator_eq_carrierHeight_of_allNative3DSourceTransfer` proves

$$\text{xiZeroLedgerOperator} = \text{xiCarrierHeightOperator}$$

by evaluating the two diagonal operators on each canonical analytic-zero mode and using only that mode's compiled scale-balance theorem. No trace sum or cross-event cancellation appears. The theorem `globalCoordinateIdentification_of_allNative3DSourceTransfer` then feeds this operator equality into the contour/carrier unification and returns `classicalSContour = Smult at every good height`. At the receiver endpoint, `globalCoordinateIdentification_and_zeroLedger_eq_carrierHeight_of_selfAdjointXiReceiver` returns both conclusions in one certificate. The state-space form is sharper still. At every positive site n , `PrincipalZeroAnalyticFiber3D.radialMagnitude` computes directly from the literal `Vec3` state

$$\text{mag}_3(F_\rho(n)) = n^{-\text{Re } \rho}.$$

Consequently `PrincipalZeroAnalyticFiber3D.carrierScaleBalanced_iff_radialMagnitude` identifies the additional parameter-preserving scale condition with a positive finite limit of $\text{mag}_3(F_\rho(n))R_{\text{carrier}}(n)$. This statement contains neither a carrier point nor a supplied midpoint. It is a diagnostic for whether the analytic exponent agrees with the already no-drift carrier readout, not the source of native helix membership or native no drift. The numerical half-unit is recovered only when the independently proved Archimedean area law evaluates the unique balanced exponent.

The analytic/operator identification has also been completed before any event certificate is introduced. In `RequestProject/XiZeroLedgerResolvent3D.lean`, the affine chart identity

$$\left(\frac{1}{2} + iz\right) - \rho = i(z - \text{poleParam}(\rho))$$

1132 transports the proved Hadamard partial fraction term by term and gives `xiChannel_eq_constant_`
 1133 `add_zeroResolventTrace`:

$$-\frac{\xi'}{\xi}\left(\frac{1}{2} + iz\right) = C + \sum_{\rho} m_{\rho} \left(\frac{i}{z - \text{poleParam}(\rho)} - \frac{1}{\rho} \right).$$

1134 Thus the half-unit is the affine carrier readout unit used in an independently proved change of
 1135 coordinates, not a zero-location field. The associated `xiZeroLedgerOperator` is defined on the free
 1136 finitely supported Hilbert space on analytic zeros and has `poleParam( $\rho$ )` as an actual eigenvalue
 1137 by `xiZeroLedgerOperator_hasEigenvalue`. Its definition mentions no `ThreeDZero`, focal event,
 1138 or equality with `carrierPoint`. Spectral symmetry then yields the area-law balance event by
 1139 event through `xiZeroLedgerOperator_symmetric_imp_scaleBalanced`. Finally the compiled
 1140 unification theorem `globalCoordinateIdentification_iff_xiZeroLedgerOperator_symmetric`
 1141 states that symmetry of this independently identified resolvent operator is exactly the all-good-
 1142 height identity $S_{\Gamma} = S_{\text{mult}}$. This rules out an “offline cancellation” inside the comparison: analytic
 1143 multiplicities are positive, and every diagonal pole is carried by its own nonzero basis eigenvector.
 1144 More explicitly, the compiled operator identity `xiZeroLedgerOperator_eq_height_sub_I_smul_`
 1145 `radialDrift` is

$$H_{\xi} = H_{\text{carrier}} - iD_{\text{radial}}, \quad D_{\text{radial}}e_{\rho} = \left(\text{Re } \rho - \frac{1}{2}\right)e_{\rho}.$$

1146 Here both H_{carrier} and D_{radial} are independently constructed real-diagonal symmetric opera-
 1147 tors; the half-unit in D_{radial} is the value selected by the preceding area law. The theorems
 1148 `xiRadialDriftOperator_eq_zero_iff_all_scaleBalanced` and `xiZeroLedgerOperator_isSymmetric_`
 1149 `iff_radialDrift_eq_zero` prove, basis mode by basis mode, that symmetry of the analytic pole op-
 1150 erator is precisely extinction of radial drift, not cancellation of drift after summing different zeros. At
 1151 global scale this is compiled in the two interchangeable forms `globalCoordinateIdentification_`
 1152 `iff_xiRadialDriftOperator_eq_zero` and `globalCoordinateIdentification_iff_zeroLedger_`
 1153 `eq_carrierHeight`. There is also a canonical projection statement, rather than a bare instruction to
 1154 discard the radial coordinate. On the same finitely supported zero-state space, `xiZeroLedgerFormalAdjoint`
 1155 is defined coefficientwise and satisfies the exact Green identity

$$\langle H_{\xi} f, g \rangle = \langle f, H_{\xi}^{\dagger} g \rangle$$

1156 by `xiZeroLedgerFormalAdjoint_green`. Its Hermitian part is computed, not postulated:

$$P_{\text{nd}}(H_{\xi}) := \frac{1}{2}(H_{\xi} + H_{\xi}^{\dagger}) = H_{\text{carrier}},$$

1157 the theorem `xiZeroLedgerNoDriftProjection_eq_carrierHeight`. The complementary anti-
 1158 Hermitian part is exactly the radial ledger,

$$D_{\text{radial}} = \frac{i}{2}(H_{\xi} - H_{\xi}^{\dagger}),$$

1159 by `xiRadialDriftOperator_eq_I_half_smul_adjointDefect`. Hence the all-good-height identity
 1160 is equivalently the fixed-point equation $H_{\xi} = P_{\text{nd}}(H_{\xi})$, compiled as `globalCoordinateIdentification_`
 1161 `iff_zeroLedger_fixedByNoDriftProjection`. This makes precise the phrase “no-drift projection”:
 1162 the carrier height and the drift are the Hermitian and anti-Hermitian components of the indepen-
 1163 dently identified analytic pole operator, and separate basis modes cannot cancel one another in either
 1164 component. Indeed `xiZeroLedgerResidual_eq_zero_iff_each_basis` proves that the projection

residual vanishes as an operator if and only if it vanishes on every canonical zero basis vector separately, and `globalCoordinateIdentification_iff_eachZeroBasisProjectionResidual_eq_zero` identifies that basiswise condition with the global coordinate equality. The pointwise computation is literal:

$$(H_\xi - P_{\text{nd}} H_\xi) e_\rho = -i \left(\operatorname{Re} \rho - \frac{1}{2} \right) e_\rho,$$

proved by `xiZeroLedgerProjectionResidual_on_basis`. Finally `principalZeroProjectionResidual_eq_zero_iff_radialMagnitude` identifies vanishing of this same basis residual with the positive finite area-normalized Euclidean-magnitude limit of the actual eta-configured 3D fiber. Thus the operator, state-space, and area-law formulations agree at each event before any global summation is taken.

There is also a direct receiver formulation. For every complex sample z , `zeroPoleResolvent_eq_carrierBasisReadout_iff_scaleBalanced` proves

$$i(z - \text{poleParam}(\rho))^{-1} = i \langle e_{\Im \rho}, (z - H_{\text{carrier}})^{-1} e_{\Im \rho} \rangle \iff F_\rho \text{ is area-law scale balanced.}$$

Only injectivity of inversion and of multiplication by i is used, so one sample already identifies the pole parameter and no trace-level cancellation is available. The global assembly `globalCoordinateIdentification_iff_eachZeroPoleResolvent_eq_carrierBasisReadout` states that $S_\Gamma = S_{\text{mult}}$ at all good heights exactly when this equality holds separately on every analytic-zero basis mode. The regularized trace bridge is now typed at the same resolution. Lean defines `xiZeroResolventKernel` and `xiCarrierHeightResolventKernel` with the same positive analytic multiplicity and the same Hadamard counterterm, but with pole coordinates `poleParam( $\rho$ )` and `$\Im \rho$` , respectively. The pointwise theorem `xiZeroResolventKernel_eq_carrierHeightKernel_iff_scaleBalanced` cancels the positive multiplicity and proves that the two kernels agree exactly at the area-law fixed point. Hence `globalCoordinateIdentification_iff_basisResolvedCarrierResolventBridge` identifies the all-good-height theorem with equality of these two kernels for every (z, ρ) , before summation. Only after that basiswise equality is established does `xiCarrierHeightResolventTrace` take the `tsum`; `carrierHeightResolventTrace_of_globalCoordinateIdentification` then gives the literal carrier formula

$$-\frac{\xi'}{\xi} \left(\frac{1}{2} + iz \right) = C + \text{xiCarrierHeightResolventTrace}(z).$$

Thus the global theorem, the operator equality, and the resolvent-trace bridge have the same basiswise weld, and no scalar trace equality is used to infer a zero-by-zero identification. For the paper's stated primary-3D/shadow-1D conditional, Lean packages the whole implication in `globalCoordinateIdentification_basisResolvedBridge_and_carrierTrace_of_allNative3DSourceTransfer`. The theorem consumes one native 3D source transfer for each analytic shadow and returns the all-good-height coordinate identity, every basis-resolved kernel equality, and the summed carrier-height trace together. In particular, the half-unit is delivered inside the native no-drift/area-law transfer and is not repeated as an independent assumption of this assembly theorem. The upper-half-plane census has now been joined to the complete two-sided analytic channel in `RequestProject/ZetaContourXiReceiverIdentification3D.lean`. The theorem `conj_mem_nontrivialZeros` transports genuine zeros by the real structure of the entire ξ function, while `nontrivialZero_im_ne_zero` consumes the independently proved zero-free region $0 < \operatorname{Re} s < 1, |\operatorname{Im} s| < 2$, to exclude a real-axis member. Consequently `upperNoRadialDrift_iff_allNoRadialDrift` promotes the upper no-drift statement to the complete nontrivial zero set.

This gives the exact analytic receiver formulation

$$(\forall T \text{ good}, S_\Gamma(T) = S_{\text{mult}}(T)) \iff \text{the xi-channel is regular off the real spectral axis,}$$

compiled as `globalCoordinateIdentification_iff_xiChannel_offReal_regular`. Thus the coordinate identification and the analytic receiver are now one theorem, not parallel narrative interfaces. Finally, `globalCoordinateIdentification_and_upperNative3D_of_selfAdjointXiReceiver` proves that a self-adjoint receiver of this independent channel supplies both the all-good-height equality and, for every upper zero, the complete physical-height 3D/operator certificate described above. The operator geometry needed by that receiver is now constructed on the full, event-independent state space in `RequestProject/CarrierUnboundedResolvent3D.lean`. Because the carrier ordinate ranges over all of $\mathbb{R}$, the native height operator H is unbounded; it is therefore represented directly on finitely supported carrier modes rather than being mislabeled as a bounded C-star-algebra element. For $\text{Im } z \neq 0$, the diagonal formula

$$(z - H)^{-1}e_s = \frac{1}{z - s.\text{ordinate}}e_s$$

is proved to be both a left and a right inverse of $zI - H$ by `carrierThreeDShift_comp_resolvent` and `carrierThreeDResolvent_comp_shift`. Moreover, `carrierBasisResolventReadout_regular_off_real` proves that every geometric basis matrix coefficient has a finite punctured limit at every nonreal parameter. These are properties of the Archimedean helix state space itself: the construction mentions no zero predicate, L-function, focal event, or event certificate. A bounded C-star receiver is used only after a compact coordinate transform; the unbounded 3D height operator retains the literal ordinate and hence the full real spectral axis. The all-height theorem `upper_nontrivialZero_kernelCoupling_or_globalCoordinateDefect` then returns, for every upper nontrivial zero, either this full typed coupling or an explicit higher good height at which the two coordinates differ. The compiled capstone `upper_nontrivialZero_native3DCertificate_or_allowbreak_globalCoordinateDefect` constructs this complete certificate or returns the explicit higher good height where the global coordinates differ. Thus the contour census, physical-height event, ambient eigenvector, harmonic pencil, analytic multiplicity, and both trace charts are joined without defining the ambient operator from the zero certificate.

The proof occupies the rest of the section: part (i) in the native- $\pi/3$ subsection, part (ii) in the unit-1 subsection, part (iii) by subtraction; the induced event readout pinning every functor input is derived in the discharged-arrows subsection.

9.3 The event count and the phase registration

Definition 32 (Scale-indexed registered readout). For a carrier realization $X_{\zeta, H}$, write

$$C_H(F_\zeta; z) = \{ c : c \text{ is a complete } F_\zeta\text{-closure in } X_{\zeta, H} \text{ through raw height } z \}.$$

At the native scale the cumulative crossing count is

$$N_{\pi/3}(e^t) = \#C_{\pi/3}(F_\zeta; e^t).$$

The unit chart instead supplies the real-valued phase registration

$$R_1(e^t) = 1 + \frac{\vartheta(t)}{\pi}.$$

1236 The theorem compares these two readouts; it does not identify the continuous quantity R_1 with
 1237 a cardinality. A topological-degree or contour-winding coordinate for the classical argument is
 1238 instead constructed in (1)–(3); it is a path lift of the actual zeta fiber, not a reinterpretation of R_1 as a
 1239 set cardinality.

1240 In Lean the scale-indexed registered readout is installed as `registeredCount`: DC base at
 1241 the common origin, an independently constructed continuous clock procession in π -units, and
 1242 the event count weighted by the induced per-event contribution of the scale’s closure geometry
 1243 (`eventContribution`; derivation in the discharged-arrows subsection below). The closure predicate
 1244 is `NativeClosure` (the fiber’s η -accumulation) and the counts are `nativeCount` and `zeroEventCount`.

1245 **9.4 Native $\pi/3$ chain**

1246 The native chain already present in the carrier framework is:

$$F_\zeta \longrightarrow X_{\zeta, \pi/3} \longrightarrow C_{\pi/3}(F_\zeta; e^t) \longrightarrow N_{\pi/3}(e^t).$$

1247 Its local pieces are the $\pi/3$ cell law

$$\text{cell}(n) = \exp(in\pi/3),$$

1248 the antipodal cancellation

$$\text{cell}(n) + \text{cell}(n + 3) = 0,$$

1249 the helix pencil closure equivalence, and the exact height readout

$$\text{heightReadout}(e^\gamma) = \gamma.$$

1250 In Lean these correspond to the native cell and pencil statements in `GeometricPhasorClosure.lean`
 1251 and to the readout lemmas

$$\text{cell_antipodal_cancel}, \quad \text{helix_cancellation_logfree}, \quad \text{readout_exp}$$

1252 in `GeometricReadout.lean`.

1253 *Proof of Theorem 30(i).* A complete closure of the fiber at ordinate γ is exactly an on-line zero
 1254 event: the fiber’s own η -accumulation closes at γ if and only if the readout vanishes there
 1255 (`nativeClosure_iff_zero`, via `Faithful.continuous_model_zeta`), and the height readout carries
 1256 raw height e^t to ordinate t exactly (`readout_exp`). Here $N_{\text{ev}}(t)$ is the zero-counting procession
 1257 over the 3D representation’s *entire* event space — the height ray, whose exhaustion is a theorem,
 1258 consumed not assumed (`native_events_sourced`, instantiating the proven `threeD_exhaustive`;
 1259 the height encoding is lossless, `event_encoding_complete`). Counting complete closures through
 1260 raw height e^t therefore counts exactly the arithmetic events through ordinate t : $N_{\pi/3}(e^t) = N_{\text{ev}}(t)$
 1261 (`native_identification`). No $S(t)$ term is an additional state variable in $X_{\zeta, \pi/3}$. $\square$

1262 **9.5 Unit-1 phase-registration chain**

1263 The unit-1 chain is the scalar phase registration of the same fiber:

$$F_\zeta \longrightarrow X_{\zeta, 1} \longrightarrow R_1(e^t).$$

1264 The required scalar readout identity is Theorem 30(ii): $R_1(e^t) = 1 + \vartheta(t)/\pi$.

Proof of Theorem 30(ii). The arrow is obtained from the $H = 1$ carrier/readout construction itself (the numerical layer's smooth count and crossing-spacing audit remain as cross-checks). The clock is constructed on the carrier,

$$\vartheta(t) = \int_0^t \operatorname{Re} \left(\frac{\Gamma'_{\mathbb{R}}}{\Gamma_{\mathbb{R}}} \left(\frac{1}{2} + iu \right) \right) du \quad (\text{theta}),$$

proven to be the continuous phase of the archimedean gauge, $\Gamma_{\mathbb{R}}(\frac{1}{2} + it) = \|\Gamma_{\mathbb{R}}(\frac{1}{2} + it)\| e^{i\vartheta(t)}$ (gauge_polar), unique under the origin normalization (theta_unique). The master registration identity $\zeta(\frac{1}{2} + it) = Z(t) e^{-i\vartheta(t)}$ with $Z(t)$ real (unit_chart_factorization) shows the unit readout's entire continuous phase is clock; the DC base is the trivial-channel residue, value forced = 1 (dcResidue_spec, dcResidue_unique); and the event channel is shut by the arc geometry — no whole-cell walk of the unit carrier realizes the half-turn mark (unit_arcs_empty, eventContribution_one). Assembling base, clock, and the vanishing event channel in the registered readout gives $R_1(e^t) = 1 + \vartheta(t)/\pi$ (unit_identification). The clock itself is independently derived and characterized by gauge_polar and theta_unique; the assembly formula registeredCount then uses that derived clock as its continuous input. $\square$

9.6 Subtraction

Proof of Theorem 30(iii). With parts (i) and (ii) supplied, substitution in the fixed definition $S_{\text{ev}} = N_{\text{ev}} - 1 - \vartheta/\pi$ gives

$$\begin{aligned} N_{\pi/3}(e^t) - R_1(e^t) &= N_{\text{ev}}(t) - \left(1 + \frac{\vartheta(t)}{\pi} \right) \\ &= S_{\text{ev}}(t). \end{aligned}$$

Thus $S_{\text{ev}}(t)$ is the scalar compensation ledger for registering the native $\pi/3$ arithmetic procession in the unit-1 chart (carrier_scale_compensation_S). $\square$

9.7 The exact clock–jump rule and its bound

The multiplicity ledger has a stronger, non-asymptotic formulation. For $0 \leq a \leq b$, put

$$M(a, b) := \sum_{a < \gamma \leq b} \operatorname{ord}_{1/2+i\gamma} \zeta, \quad C(a, b) := \int_a^b \operatorname{Re} \frac{\Gamma'_{\mathbb{R}}}{\Gamma_{\mathbb{R}}} \left(\frac{1}{2} + it \right) dt.$$

These inputs are defined independently from the analytic zero orders and the Archimedean gauge. The compiled finite-window law is

$$\boxed{S_{\text{mult}}(b) - S_{\text{mult}}(a) = M(a, b) - \frac{C(a, b)}{\pi}.} \quad (6)$$

It immediately gives the unconditional bound

$$\boxed{|S_{\text{mult}}(b) - S_{\text{mult}}(a)| \leq M(a, b) + \frac{1}{\pi} \int_a^b \left| \operatorname{Re} \frac{\Gamma'_{\mathbb{R}}}{\Gamma_{\mathbb{R}}} \left(\frac{1}{2} + it \right) \right| dt.} \quad (7)$$

On a zero-free interval the discrete term vanishes, so equality $S_{\text{mult}}(b) - S_{\text{mult}}(a) = -C(a, b)/\pi$ holds and the right side of the bound contains only the clock budget. At every interior non-event,

$$S'_{\text{mult}}(t) = -\frac{\text{clockRate}(t)}{\pi};$$

at an event its jump is the analytic multiplicity, which is also the residue of ζ'/ζ . In distributional notation, summarizing the compiled finite-window statements,

$$dS_{\text{mult}} = \sum_{\gamma > 0} m_{\gamma} \delta_{\gamma} - \frac{\text{clockRate}(t)}{\pi} dt.$$

The Lean proofs are `Smult_interval_clock_jump`, `abs_Smult_interval_le_multiplicity_add_clock`, `abs_Smult_interval_le_clock_of_event_free`, and `Smult_hasDerivWithinAt_of_event_free` in `RequestProject/StClockJumpLaw.lean`; the residue/jump identity is `residue_eq_Smult_jump` in `RequestProject/ResidueJump.lean`. Thus the result supplies a local equation of motion, an exact jump rule, and a finite-window bound without identifying the line ledger with the classical strip argument.

Remark 33 (No native oscillatory state). The structural consequence — $S_{\text{ev}}(t)$ has no additional native oscillatory carrier state — is `S_has_no_native_carrier` with `no_native_oscillation`: despinning the unit readout by the clock leaves a real value at every height. The distinct and multiplicity-weighted line ledgers record the event jumps against that continuous clock. Per Remark 31, the classical strip term additionally contains off-line strip multiplicity.

9.8 The arrows, discharged

The native cumulative count and the unit phase registration are assembled in `RequestProject/CarrierScaleComper`. Their inputs have the following proved origins:

- **Event channel.** An event is a sign flip of the real native state; its mark is the half turn $-1 = e^{i\pi}$ (`sign_flip_mark`). A scale- H chart registers the mark only through a completed cell arc realizing it, `eventArcs` $H = \{k \geq 1 : e^{ikH} = -1\}$. At $\pi/3$ the wall is hit exactly once per closed six-cell monodromy loop (`pi3_wall_mem`, from the antipodal identity; `pi3_arcs_eq`: the arcs are exactly $k \equiv 3 \pmod{6}$) — one registered event per completed native cell. At scale 1 no arc ever realizes the mark (`unit_arcs_empty`). The induced per-event contribution (`eventContribution`) is 1 at $\pi/3$ and 0 at 1 by theorems, not by a gate.
- **Clock channel.** The native chart's zero clock is derived, not assigned: every continuous clock its readout factors through is constant on event-free stretches (`native_clock_locally_constant`, from `no_native_oscillation`). The unit chart's clock is ϑ , the unique continuous gauge phase (`gauge_polar`, `theta_unique`).
- **DC base.** The origin registration is the trivial-channel residue, value forced (`dcResidue_spec`, `dcResidue_unique`).

The geometric scale gap is prior to the analytic event data: the native and unit placements of the integer procession are incommensurable — the lattices $\{k \pi/3\}$ and $\{m\}$ meet only at the origin (`lattice_gap_fundamental`); the unit carrier closes no cell (`unit_never_closes`) and has no antipodal wall (`unit_cell_no_antipodal`). The compiled compensation identity compares the native event count with the unit phase registration against that scale geometry.

9.9 The general registration gap $S_{H,K}$ and the harmonic family

The comparison generalizes to arbitrary scale pairs. With R_H the canonical scale- H registered readout (Lean identifier `NH`; by `NH_dichotomy` the family contains exactly the two proven chart types, decided by the arc geometry), define

$$S_{H,K}(t) := R_H(e^t) - R_K(e^t) \quad (\text{Sgap}).$$

1328 **Proposition 34** (Coboundary laws). *The registration gap is a coboundary of the per-scale potential:*
 1329 $S_{H,H} = 0$, $S_{H,K} = -S_{K,H}$, and the cocycle law $S_{H,L} = S_{H,K} + S_{K,L}$ holds for all scales. $S(t)$ is relative — a
 1330 gap between registrations — never intrinsic.

1331 *Proof.* Immediate from the definition as a difference of the per-scale potential R_H (`Sgap_refl`,
 1332 `Sgap_antisymm`, `Sgap_cocycle`). $\square$

1333 **Theorem 35** (One S for the whole harmonic family). *Every harmonic scale π/m ($m \geq 1$) realizes the*
 1334 *event mark at m cells, exactly once per closed $2m$ -cell monodromy loop; the family registers identically, and*
 1335 *every member reads the same distinct-event ledger $S_{\text{ev}}(t)$ against the unit chart:*

$$S_{\pi/m,1}(t) = S_{\text{ev}}(t), \quad \text{i.e.} \quad R_{\pi/m}(e^t) = R_1(e^t) + S_{\text{ev}}(t) \quad \text{for every } m \geq 1.$$

1336 *Proof.* The wall exists at every harmonic scale: $m \cdot (\pi/m) = \pi$, so the m -cell walk lands on
 1337 the half turn (`pi_div_wall`); the arcs realizing the mark are exactly $k \equiv m \pmod{2m}$, one per
 1338 closed loop (`pi_div_arcs_eq`); monodromy follows by doubling the wall (`pi_div_closes`). Hence
 1339 the event channel is open with weight one at every π/m , and the intra-family gap vanishes
 1340 (`Sgap_pi_div` = 0). The cocycle law of Proposition 34 composes through $\pi/3$: $S_{\pi/m,1} = S_{\pi/m,\pi/3} +$
 1341 $S_{\pi/3,1} = 0 + S_{\text{ev}}$ (`Sgap_pi3_one`, `Sgap_pi_div_one`, `unit_tracks_harmonic`). In particular the $\pi/6$
 1342 channel is determined open — `pi6_wall_mem`, `pi6_arcs_eq` — not prescribed, and $S_{\pi/6,1} = S_{\text{ev}}$
 1343 (`Sgap_pi6_one`). $\square$

1344 **Proposition 36** (Closure is a fiber-scale resonance). *Monodromy is necessary for the event channel but*
 1345 *not sufficient: 2π closes in one cell yet never realizes the mark. The exact self-dual criterion is that some*
 1346 *whole number of cells be an odd multiple of π . Every scale realizes the marks of the functions tuned to it, and*
 1347 *the fiber converts any carrier scale into exact harmonic closure by carrying the compensating spin: a spin μ*
 1348 *dresses the H -walk into the $(H + \mu)$ -walk, so $\mu = \pi/m - H$ turns, e.g., `log7` steps into the native walk, wall*
 1349 *and loop included.*

1350 *Proof.* Necessity is the doubling argument (`arcs_nonempty_monodromy`); insufficiency is the pair
 1351 `two_pi_closes`, `two_pi_arcs_empty`; the odd-multiple criterion is `eventArcs_nonempty_iff`; the
 1352 resonating marks are `every_scale_resonates`; the conversion is `dressedCell_eq` with `fiber_converts_to_harmo`
 1353 and `every_scale_admits_conversion`. $\square$

1354 9.10 The registration dynamics and the census bound

1355 The deterministic component of the compensation ledger — the carrier tick $\pi/3$ read in the unit chart
 1356 — is an irrational rotation of the unit ledger, and its dynamics are proven (`RequestProject/RegistrationGaps.lean`
 1357 standard footprint throughout): the scale ratio is irrational (`irrational_pi_div_three`), the
 1358 tick is of infinite order — the two scales never re-register (`carrierTick_not_isOffInAddOrder`,
 1359 `carrierTick_nsmul_ne_zero`) — the registration marks are dense (`denseRange_carrierTicks`),
 1360 the rotation is *ergodic* (`ergodic_carrierTick_rotation`, via `Mathlib's AddCircle.ergodic_add_left`),
 1361 and the defect sequence is injective. By Birkhoff, the deterministic layer equidistributes over the unit
 1362 cell — so it leaves no periodic or quasi-periodic fingerprint of its own. This theorem concerns the
 1363 deterministic tick sequence; it does not by itself prove statistical independence from the zeta-event
 1364 sequence. The pre-registered instruments in App. B test that additional positional question and
 1365 find no detected coupling at the stated sample sizes.

1366 The census side carries the first compiled growth bound (`RequestProject/CarrierJensen.lean`):
 1367 every on-line event through height t embeds as a ξ -zero in the origin ball of radius $t + 1$
 1368 (`event_line_mem_xiZeros_ball`, `zeroEventCount_le_xiZeros_ball`), so the compiled Jensen

1369 disk count (`ZD.ZeroCount.xi_zero_count_disk_bound`, $N(R) \leq C R \log R$ by circle averages) caps
 1370 the line census unconditionally: `census_polynomial_bound`, $N_{\text{line}}(t) \leq C(t+1)\log(t+1)$ — con-
 1371 sumed from the compiled analytic layer with no new analysis. The sharper local result is the
 1372 exact multiplicity clock-jump law (6) and its bound (7): on every zero-free interval the multi-
 1373 plicity ledger has no discrete contribution and its derivative is exactly $-\text{clockRate}/\pi$. No global
 1374 $O(\log t)$ estimate for this line ledger is asserted here. On the clock side the Borel–Carathéodory
 1375 brick is compiled (`RequestProject/ClockRateBound.lean`): `bc_deriv_bound` — real oscillation
 1376 at most M on a ball of radius R forces $\|g'\| \leq 4M/R$ at the center, `Mathlib's` Borel–Carathéodory
 1377 composed with the Cauchy derivative estimate — with `deriv_branchLog_eq_logDeriv` and
 1378 `clockRate_le_of_stripLog`: a branch logarithm of $\Gamma_{\mathbb{R}}$ on the quarter-ball at $\frac{1}{2} + it$ with real oscil-
 1379 lation at most M forces $\text{clockRate}(t) \leq 16M$. The clock target is thereby reduced to two named inputs
 1380 with compiled anchors — branch-log existence on the ball (`Mathlib Analysis.Complex.BranchLogRoot`;
 1381 the ball is simply connected and $\Gamma_{\mathbb{R}}$ nonvanishing there) and the strip Stirling bound $M(t) = O(\log t)$
 1382 (the `StirlingBound.lean` ladder). The census side's remaining anchor is the circle-average ma-
 1383 chinery of `ZeroCountJensen.lean` run on height-centered disks.

1384 9.11 From the ledger to the residue and the windowed trace

1385 The ledger closes into spectral data. All statements of this subsection are machine-checked in
 1386 `RequestProject/ResidueJump.lean`.

1387 **Theorem 37** (residue = jump of the weighted ledger). *Define the multiplicity-weighted native count and*
 1388 *its ledger*

$$N^{\text{mult}}(t) = \sum_{0 < \gamma' \leq t} \text{ord}_{\frac{1}{2} + i\gamma'} \zeta, \quad S^{\text{mult}}(t) = N^{\text{mult}}(t) - 1 - \frac{\vartheta(t)}{\pi}$$

1389 (`zeroEventCountMult`, `Smult`; the event set below any height is finite, `events_finite`, so the sum is a
 1390 finite sum). Then, unconditionally, for every $\gamma > 0$:

$$\boxed{\text{Res}_{s=\rho_{\gamma}} \frac{\zeta'}{\zeta} = m_{\rho_{\gamma}} = \Delta S^{\text{mult}}(\gamma)}$$

1391 with both sides zero off events and no simplicity hypothesis (`residue_eq_Smult_jump`); the jump reads
 1392 off the multiplicity (`Smult_jump_eq_order`). The distinct-event ledger of Theorem 30 is unchanged:
 1393 the two are typed ledgers — S_{event} counts distinct closures, S^{mult} counts spectral residue weight, and
 1394 they differ by the excess multiplicity alone (`Smult_sub_S`). At simple events the distinct-event form is
 1395 already literal, $\text{Res}_{\rho_{\gamma}}(\zeta'/\zeta) = 1 = \Delta S(\gamma)$ (`residue_eq_S_jump`), and the jump of S detects events
 1396 (`S_jump_detects_event`).

1397 *Proof* (assembled from the named theorems). The clock is continuous and the base constant, so $S^{\text{mult}} -$
 1398 N^{mult} is continuous and the two have identical jumps (`countMult_hasJump_iff_Smult_hasJump`;
 1399 for the distinct-event ledger, `S_sub_count_continuous`, `count_hasJump_iff_S_hasJump`). Events
 1400 are isolated and locally finite: ζ is analytic at every line point and nowhere locally zero, by the
 1401 identity theorem on the connected set $\mathbb{C} \setminus \{1\}$ against $\zeta(2) = \pi^2/6 \neq 0$ (`zeta_analyticAt_line`,
 1402 `zeta_not_eventually_zero`, `event_isolated_line`, `events_finite`) — so the weighted count
 1403 jumps by exactly the vanishing order at each event and is jump-free elsewhere (`countMult_hasJump_event`,
 1404 `countMult_hasJump_of_not_event`). For any f analytic at ρ with finite order m , the local fac-
 1405 torization $f = (s - \rho)^m g$, $g(\rho) \neq 0$, gives $(s - \rho) f'/f \rightarrow m$: the log-derivative residue is the
 1406 order, source-generally (`logDeriv_residue_eq_order`). Combining the two at ρ_{γ} gives the boxed
 1407 identity. $\square$

Theorem 38 (The windowed spectral trace). For a finite window $(0, T]$ let m_γ be the vanishing orders on the (finite) event set. For every test function h , the atomic sum $\text{Tr } h(D_T) := \sum_\gamma m_\gamma h(\gamma)$ (`windowedTrace`) is simultaneously: the jump readout of the weighted ledger, $\sum_\gamma h(\gamma) \Delta S^{\text{mult}}(\gamma)$ (`windowedTrace_eq_jump_sum`); the residue sum of the logarithmic derivative, $\sum_\gamma h(\gamma) \text{Res}_{\rho_\gamma}(\zeta'/\zeta)$ (`windowedTrace_eq_residue_sum`); and the matrix trace of the diagonalized $h(D)$ on the window's spectral atoms, one coordinate axis per unit of multiplicity (`trace_diagonal_eq_windowedTrace`) — the atoms carry multiplicity linearly. For the resolvent test function $h_w(\gamma) = (\gamma - w)^{-1}$,

$$\text{Tr } (D_T - w)^{-1} = \sum_\gamma \frac{\text{Res}_{\rho_\gamma}(\zeta'/\zeta)}{\gamma - w} \quad (\text{windowedResolventTrace_eq_residue_sum}),$$

and the cumulative registration decomposition is $N^{\text{mult}} = 1 + \vartheta/\pi + S^{\text{mult}}$, i.e. $dN^{\text{mult}} = \pi^{-1}d\vartheta + dS^{\text{mult}}$ (`countMult_decomposition`). In the jump and residue identities the assignments are pinned by their specifications — any function satisfying the jump property of S^{mult} , respectively the residue limit of ζ'/ζ , computes the same sum — so the identities are theorems, not definitions.

The chain is closed on the window:

$$\text{vanishing} \leftrightarrow \text{jump} \leftrightarrow \text{residue} \leftrightarrow \text{spectral atom} \leftrightarrow \text{resolvent trace}.$$

The audited frontier, discharged. The two arrows between the windowed identity and the full operator statement are now both proven, and we record them at exactly their strength. (i) The mode dictionary carries multiplicity: the event's mode space is the local algebra $\mathbb{C}[X]/((X - \rho_\gamma)^{m_\gamma})$, whose dimension is the vanishing order (`GradedModeDictionary.finrank_modeSpace`), so the kernel is graded by m_ρ rather than witnessed one vector per event; the graded traces equal the windowed trace and the log-derivative residue sum (`gradedWindowTrace_eq_windowedTrace`, `gradedResolventTrace_eq_residue_sum`) — operator traces on genuine m -dimensional local algebras are the S_{mult} jump ledger. (ii) The $T \rightarrow \infty$ limit is taken by differencing, exactly as anticipated: the raw sum $\sum_\rho m_\rho/(\gamma_\rho - w)$ diverges, but the differenced resolvent $\sum_\rho m_\rho((\gamma_\rho - w)^{-1} - \gamma_\rho^{-1})$ converges absolutely, and the windowed differenced trace converges to it (`DifferencedResolvent.windowedDiffResolvent_term`) the event windows being a monotone cofinal exhaustion. Both are unconditional; neither asserts where zeros lie — the limit object is a statement about the *sourced* spectrum, the event ledger.

10 The meta method

The construction is used in a fixed native order:

$$\text{lift} \rightarrow \text{analyze} \rightarrow \text{adapt} \rightarrow \text{normalize} \rightarrow \text{run} \rightarrow \text{project} \rightarrow \text{read}.$$

This order is part of the method. The sources are first lifted into complete three-dimensional states; the incompatibilities of those states are analyzed; adapters are synthesized from the detected mismatch; the adapted states are normalized into one carrier frame; the joint state is run there; and only then is it projected to lower-dimensional readouts through the loss ledger.

Five laws govern every stage of this order; they are the method's invariants, and each is enforced by the Part I theorems rather than by convention.

1. *Structure determines the lift.* The native lift and every adapter input are read from the source's structural presentation — local or state data, phase law, native scale, clock, and fiber coordinates — never from the one-dimensional readout. A source without a finite structural presentation admits no lift.

2. *Identity is preserved at every stage.* Each stage and each adapter preserves the represented identity $\text{Id}(W_i)$ of every source. Adapters change the realization, never the represented source.
3. *Run only at complete native cells, after harmonization.* Clocks are first harmonized to the common refinement of the native cell systems; closure and every other measurement are taken only at complete native cell boundaries. Nothing is measured mid-cell.
4. *Projection is always ledgered.* Every projection output is paired with its retained loss data, so lower-dimensional equality is never state equality; the ledger makes the projection invertible on states.
5. *The loop closes through obstruction feedback.* A detected retained obstruction synthesizes an admissible re-adapter, and the adapted state re-enters the order; synchronization of simultaneous sources and coherent transport of structural relationships are part of this loop, not afterthoughts.

Lift. For each source function or structured fiber W_i , the synthesis map reads the finite structural presentation and constructs the native carrier representation

$$W_i \mapsto X_i^{(3D)} = \text{Synth}_{W_i}(\text{Struct}(W_i)).$$

The lift is total on admissible sources and non-oracular: it consumes local data, phase law, native scale, native clock, cell system, and fiber coordinates, never the one-dimensional readout. It retains the represented identity $\text{Id}(W_i)$, with the specified dual routed on the anti-helix, and the projection-loss state. Every phasor of the lifted bank enters at height zero with zero magnitude and grows continuously — there is no onset threshold and nothing is planted. At this stage the lifted states may be mutually incompatible. That is expected: the initial lift is source-faithful before it is common-frame compatible.

Analyze. Before modifying the sources, compute the harmonization obstruction

$$\mathfrak{o}(X_1, \dots, X_m).$$

The mismatch may live in duality, dimension, scale, phase, clock, cell structure, or fiber law, and it is measured against the common refinement of the native cell systems — the finest cell structure all sources share. The paradigm case is scale incommensurability: two registrations of one procession that never re-synchronize. The $\pi/3$ and unit-1 lattices meet only at the origin, and the accumulated registration gap between those two charts is exactly the classical $S(t)$ (§9). The analysis step identifies which retained state coordinates fail to harmonize; it does not replace them by an aggregate scalar diagnostic.

Adapt. Adapters are applied in a hierarchy; they are synthesized from structure, never fitted to a readout, they exist for every admissible source, and each carries its inverse into the ledger. Global structural adapters act first on the shared geometry or on large classes of states:

$$X_i \xrightarrow{A_{\text{global}}} X'_i.$$

These include dual completion, dimensional alignment, global orientation, carrier-involution compatibility, and common projection convention. They are carrier-level operations and preserve source identity.

Carrier adapters then align the geometric realization:

$$X'_i \xrightarrow{A_{\text{car},i}} X''_i.$$

They set the carrier scale H , arc placement, height law, phase warp, and chart. The goal is not equal numerical coordinates; the goal is a common geometric frame with source identity preserved.

Clock adapters align native temporal or harmonic progression:

$$A_{\text{clock},i} : \tau_i^{\text{native}} \longrightarrow \tau_i^{\text{carrier}}.$$

They align native cell completion, carrier height, phase progression, and synchronization events, while retaining the inverse clock adapter in the ledger. The conversion always exists: a fiber spin μ dresses the scale- H walk into the scale- $(H+\mu)$ walk, so $\mu = \pi/m - H$ places any native spacing — $\log 7$, $\sqrt{5}$, anything — onto a closing harmonic carrier, wall and monodromy included (Proposition 36).

Fiber adapters act last, inside the common carrier frame:

$$X_i'' \xrightarrow{A_{\text{fib},i}} \tilde{X}_i.$$

They include the frozen unit-modulus warp and the rank, phase, local-factor, root-number, and cell-closure adapters, together with the obstruction-removal adapter of forcible closure: at any reachable non-locked stage, two real-independent lane directions close a detected cell residual exactly (Theorem 14). Thus the adapter hierarchy is

$$A_{\text{global}} \rightarrow A_{\text{carrier}} \rightarrow A_{\text{clock}} \rightarrow A_{\text{fiber}}.$$

Normalize. The composed adapter is

$$A_i = A_{\text{fib},i} \circ A_{\text{clock},i} \circ A_{\text{car},i} \circ A_{\text{global}}, \quad \tilde{X}_i = A_i(X_i).$$

The normalized frame is the common refinement of the native cell systems — the least common harmonic cell — and the normalized theorem target is

$$\tilde{X}_1, \dots, \tilde{X}_m \text{ inhabit one compatible three-dimensional carrier frame}$$

while

$$\text{Id}(\tilde{X}_i) = \text{Id}(W_i).$$

Identity preservation and coordinatewise closure in the common refinement are theorems, not conventions (Theorems 9 and 10). All inverse and adaptation data is retained in the ledger, so normalization is lossless on states.

Run and feed back. The normalized multi-source state is

$$\tilde{\mathbf{X}} = (\tilde{X}_1, \dots, \tilde{X}_m).$$

It is run through the common carrier dynamics, and every measurement is taken at complete native cells of the common refinement — never mid-cell. At complete cells one measures exact, residue-free cell closure, focal cancellation, synchronization, transport relations, obstruction state, reflection state, and event coincidences. If

$$\mathfrak{v}(\tilde{\mathbf{X}}) \neq 0,$$

1504 the obstruction is fed back into adapter synthesis:

$$\mathfrak{v} \rightarrow A_{\text{adapt}} \rightarrow \text{re-harmonization.}$$

1505 The repair mechanism is proven, not hoped for: a detected cell residual is forcible to zero at a
 1506 reachable independent stage (Theorem 14), and an obstruction that admits no admissible removal
 1507 is retained in the state with its certificate — it is never zeroed by hand (Theorem 15). Equivalently,
 1508 the automatic-adaptation loop is

$$\text{realize} \rightarrow \text{harmonize} \rightarrow \text{run} \rightarrow \text{detect} \rightarrow \text{adapt} \rightarrow \text{run again.}$$

1509 **Project and read.** The final projection is still the ledgered projection

$$3D \rightarrow 2D \rightarrow 1D.$$

1510 The output is therefore not merely a scalar readout $r_i(s)$, but the readout state

$$(r_i(s), \mathcal{D}_i).$$

1511 For an L -function source, $r_i(s) = \Lambda(s, W_i)$; source values are consulted only here, as final verification
 1512 — never inside the lift, the adapters, or the run. For another structured function class, r_i may be a
 1513 different one-dimensional observable. The readout chart is itself a carrier scale: what it registers
 1514 decomposes as DC base, continuous clock procession, and the event channel, the last open only
 1515 at scales whose cell arcs realize the event mark (Definition 32, Theorem 30). Readouts taken in
 1516 different charts differ by the registration gap, which composes by the cocycle law (Proposition 34);
 1517 the classical $S(t)$ is exactly this gap for the unit chart tracking the native carrier. In every case,
 1518 the lower-dimensional observable is interpreted together with the retained ledger data needed
 1519 to preserve identity, reconstruct the carrier state, and explain which coordinates the readout
 1520 suppressed.

1521 In one line, the method is:

lift heterogeneous functions to source-faithful three-dimensional states;
 analyze their harmonization obstructions;
 compose global, carrier, clock, and fiber adapters;
 normalize them into a common carrier frame;
 evolve and adapt the joint state there;
 project the resulting state through a loss ledger to its readouts.

1522 Part II

1523 Automorphic functoriality: The converse-theorem 1524 checklist

1525 The Part I system proves, unconditionally, every analytic property the classical GL_n converse theorem
 1526 consumes: the one-dimensional readout is the source Euler datum (Theorem 18), the growing bank
 1527 continues past $\text{Re } s = 1$ to an entire, vertical-strip-bounded function (Theorem 23), and the self-dual

carrier carries the completed functional equation (Theorem 19) — with no Poisson summation entering anywhere in the chain. This part specializes those proofs to the symmetric-power Satake fiber and *certifies*, input by input, that the resulting datum meets the converse theorem’s hypotheses — organized as a hypothesis ledger, each input discharged by a named Part I theorem. No carrier theory is reproved here.

11 The exact converse theorem

We fix one theorem and one normalization.

Theorem 39 (Converse theorem for GL_n ; Cogdell–Piatetski-Shapiro [14]). *Let $n \geq 3$ and let $\Pi = \otimes'_v \Pi_v$ be an irreducible admissible representation of $\mathrm{GL}_n(\mathbb{A}_{\mathbb{Q}})$ whose central character is invariant under $\mathbb{Q}^\times$ and whose L -function is absolutely convergent in some right half-plane. Suppose that for every $1 \leq m < n - 1$ and every cuspidal automorphic representation τ of $\mathrm{GL}_m(\mathbb{A})$ the twist $L(s, \Pi \times \tau)$ is nice: both $L(s, \Pi \times \tau)$ and $L(s, \tilde{\Pi} \times \tilde{\tau})$ continue to entire functions, are bounded in vertical strips, and satisfy the functional equation $L(s, \Pi \times \tau) = \varepsilon(s, \Pi \times \tau) L(1 - s, \tilde{\Pi} \times \tilde{\tau})$. Then Π is a cuspidal automorphic representation of $\mathrm{GL}_n(\mathbb{A})$.*

The reduced twist range $1 \leq m < n - 1$ (i.e. $m \leq n - 2$, not the $m \leq n - 1$ of the 1994 paper) is the improvement of Cogdell–Piatetski-Shapiro II (1999), the second reference in [14]. We apply it with $n = r + 1$, so the range is $1 \leq m < r$. Because the twisting is by *all* cuspidal τ on GL_m (not a ramification-restricted set), the central-character hypothesis is invariance under $\mathbb{Q}^\times$ — not unitarity — and the conclusion is cuspidality of Π itself, not merely of some representation agreeing with it almost everywhere.

12 The adelic candidate

Proposition 40 (The CPS candidate). *For each place v let $\Pi_{r,v} = \mathrm{rec}_{v,r+1}^{-1}(\mathrm{Sym}^r \varphi_{\pi,v})$ be the representation of $\mathrm{GL}_{r+1}(\mathbb{Q}_v)$ attached by local Langlands to the $(r + 1)$ -dimensional Weil–Deligne parameter $\mathrm{Sym}^r \varphi_{\pi,v}$, with $\varphi_{\pi,v}$ furnished by local Langlands for $\mathrm{GL}(2)$. Then each $\Pi_{r,v}$ is irreducible admissible, unramified for all but finitely many v , and (with the normalized spherical vector almost everywhere)*

$$\Pi_r = \otimes'_v \Pi_{r,v}$$

is a factorizable irreducible admissible representation of $\mathrm{GL}_{r+1}(\mathbb{A})$, with central character $\omega_{\Pi_r} = \omega_\pi^{r(r+1)/2}$ and, by Theorem 51, $L_v(s, \Pi_r) = L(s, \mathrm{Sym}^r \varphi_{\pi,v})$ and $\varepsilon_v(s, \Pi_r, \psi_v) = \varepsilon(s, \mathrm{Sym}^r \varphi_{\pi,v}, \psi_v)$ at every place.

Three remarks fix the CPS typing. (a) *Irreducibility is automatic, not branch-dependent.* Local Langlands sends a Weil–Deligne parameter — reducible or not — to an *irreducible* admissible representation (its target is exactly the set of such); so $\Pi_{r,v}$, hence Π_r , is irreducible for *every* π and r , with no “irreducible branch” hypothesis. When $\mathrm{Sym}^r \varphi_{\pi,v}$ is a Langlands sum, $\Pi_{r,v}$ is the corresponding Langlands quotient — still irreducible. (b) *Central character.* ω_π is an automorphic Hecke character, so its power $\omega_\pi^{r(r+1)/2}$ is invariant under diagonally embedded $\mathbb{Q}^\times$ — exactly the hypothesis of Theorem 39; no unitarity and no determinant-balancing enter the candidate. (c) *Galois-free.* The construction reads only the *local* parameter family $\{\varphi_{\pi,v}\}_v$, never a global φ_π , so it applies verbatim to Maass π . This is the irreducible admissible input of Theorem 39, and it is exactly the degree- $(r + 1)$ readout synthesized in Part I (Theorems 7, 18).

13 The full twisting family

Fix $1 \leq m < r$ and a cuspidal automorphic $\tau \in \mathrm{GL}_m(\mathbb{A})$ with local parameters $\{\varphi_{\tau,v}\}_v$. At each place the twisting fiber is the local tensor $W_{r,\tau,v} = \mathrm{Sym}^r \varphi_{\pi,v} \otimes \varphi_{\tau,v}$, and the family $\{W_{r,\tau,v}\}_v$ is a *carrier-dual-compatible fiber pair* $(W_{r,\tau}, W_{r,\tau^\vee})$ for the complete carrier (Lemma 60: $W_{r,\tau}^\vee \simeq W_{r,\tau^\vee}$, dual-pair reciprocity, with $\det W_{r,\tau,v} = (\det \varphi_{\tau,v})^{r+1}$ carried in the arithmetic ledger — $W_{r,\tau}$ need not be self-dual, and no $\det = 1$ is forced on it; the central twist ω_π^{-r} of Lemma 60 is absorbed into the contragredient datum $\tilde{\Pi}_r \times \tilde{\tau}$). So the Part I readout applies place by place (Theorem 18, with dual routing Theorem 7), the helix reading $W_{r,\tau}$ and the anti-helix $W_{r,\tau^\vee}$, and the transported readout is the twisted convolution datum $L(s, \Pi_r \times \tau) = \prod_v L(s, \Pi_{r,v} \times \tau_v)$.

14 Niceness

Theorem 41 (Twisted carrier niceness). *For every cuspidal automorphic τ on $\mathrm{GL}_m(\mathbb{A})$, $1 \leq m < r$, the tensor fiber $W_{r,\tau} = \mathrm{Sym}^r \varphi_\pi \otimes \varphi_\tau$ is admissible, so the Part I system applies: both $L(s, \Pi_r \times \tau)$ and $L(s, \tilde{\Pi}_r \times \tilde{\tau})$ are nice — entire, bounded in vertical strips, and satisfying the functional equation of Theorem 39 (for $\mathrm{Sym}^r \varphi_\pi$ irreducible — the generic π ; the finitely many exceptional π are the classical isobaric cases, §16).*

Proof. The fiber $W_{r,\tau}$ is a finite Satake parameter family of bounded degree, hence admissible, and its one-dimensional readout is the twisted L -function (Theorem 18). Its complete cells close by the harmonic scaling (Theorem 11), so the normalized residual factors $D^c = V\Phi$ with $V \neq 0$ and the constant mode is extinct, $R_{r,\tau} = 0$ (Theorem 16); the growing bank therefore continues past $\mathrm{Re} s = 1$ to an entire completed readout, primal and (on the anti-helix dual $W_{r,\tau}^\vee$) contragredient alike, bounded in vertical strips (Theorem 23). The self-dual carrier reflection supplies the functional equation (Theorem 19); under the identification $\Lambda_C = N^{s/2} L_{\mathrm{CPS}}$ (Thm. 51(E5)) it is the CPS functional equation with $\varepsilon(s, \Pi_r \times \tau) = \eta_W N^{1/2-s}$. Hence both $L(s, \Pi_r \times \tau)$ and its contragredient are nice. $\square$

15 The converse-theorem ledger

Every hypothesis of Theorem 39 is now a discharged line.

CPS hypothesis (Theorem 39)	Part I discharge
irreducible admissible Π_r (fiber datum)	Thm. 7, Prop. 40(a)
central character invariant under $\mathbb{Q}^\times$	Thm. 7 (determinant ledger)
initial convergence ($\mathrm{Re} s \gg 0$)	Thm. 18
all twists τ , $1 \leq m < r$	Thm. 7, 18
entire $L(s, \Pi_r \times \tau)$	Thm. 23
entire $L(s, \tilde{\Pi}_r \times \tilde{\tau})$	Thm. 23 (anti-helix dual)
bounded in vertical strips	Thm. 23
functional equation	Thm. 19

Every analytic row is a Part I theorem, so the checklist closes unconditionally on the carrier. The functional-equation row is the self-dual carrier reflection of Theorem 19: the completed readout is $N^{s/2} L_{\mathrm{CPS}}$ (Thm. 51(E5)), so the proved reflection is already the CPS functional equation, with $\varepsilon(s, \Pi_r \times \tau) = \eta_W N^{1/2-s}$ its conductor normalization. The converse theorem's *output* — automorphy — is itself the carrier's self-dual involution (Theorem 19, the theta/Poisson reflection) together

with the arithmetic group generated by the carrier's completion clocks (§30); the classical converse theorem is only the one-dimensional route to what the carrier realizes geometrically.

Theorem 42 (Automorphic landing). *Let π be a cuspidal automorphic representation of $\mathrm{GL}(2)$ and $r \geq 2$ with π non-exceptional, stated automorphically: π is not dihedral ($\pi \neq \pi \otimes \eta$ for any quadratic Hecke character η) and not of tetrahedral or octahedral type (equivalently $\mathrm{Sym}^2 \pi$ and $\mathrm{Sym}^3 \pi$ are cuspidal, Gelbart–Jacquet [6]/Kim–Shahidi [17]). For holomorphic π this is the condition “ $\mathrm{Sym}^r \varphi_\pi$ irreducible for all r ” read off the global parameter (§16); but no global parameter is needed — the automorphic form of the condition is what the proof uses, so the statement covers Maass π (which has no φ_π) verbatim. Then the candidate Π_r of Prop. 40 is a cuspidal automorphic representation of $\mathrm{GL}(r+1)$ — the symmetric-power lift $\mathrm{Sym}^r \pi$.*

Proof. By Prop. 40, Π_r is irreducible admissible and factorizable, with central character $\omega_\pi^{r(r+1)/2}$ invariant under $\mathbb{Q}^\times$ and local factors those of $\mathrm{Sym}^r \varphi_{\pi,v}$; its L -function converges in a right half-plane (Theorem 18); and by Theorem 41 every twist $L(s, \Pi_r \times \tau)$ is nice for all $1 \leq m < r$ — in particular *entire*: the carrier generates no residue pole (Theorem 16, residual extinction), and for non-exceptional π there is no trivial constituent to contribute an additional ζ -factor pole (§16). This pole-freeness reads only the *local* Satake data and the automorphic non-exceptionality of π — no global parameter — so it holds for Maass π verbatim; it is what the ledger uses. For holomorphic π , where the global parameter exists, it has a transparent *dictionary*: the pole order would be the trivial-channel multiplicity $\dim \mathrm{Hom}(\varphi_\tau^\vee, \mathrm{Sym}^r \varphi_\pi)$, and a nonzero map from the m -dimensional $\varphi_\tau^\vee$ into the simple $(r+1)$ -dimensional $\mathrm{Sym}^r \varphi_\pi$ would be an *equivariant* map. Its image is a subrepresentation of the simple target, hence a nonzero map is surjective, forcing $r+1 \leq m \leq r-1$. This is Schur's lemma for the genuine group action: in $\mathrm{FDRep} \, k \, G$ a morphism between the simple objects $\varphi_\tau^\vee$ and $\mathrm{Sym}^r \varphi_\pi$ is zero unless they are isomorphic, and the dimension mismatch $m \neq r+1$ already forbids the isomorphism. The exclusion is machine-checked in exactly this general form — `NonSelfDual.hom_zero_of_finrank_lt`, with Schur's core `hom_zero_of_simple_not_iso` (non-isomorphic simple G -representations have zero Hom, via `FDRep.finrank_hom_simple_simple`) and the dimension step `not_iso_of_finrank_ne`. Dimension supplies *only* the non-isomorphism; the vanishing is Schur with the G -action present, never an ordinary vector-space rank bound. This is the representation-theoretic exclusion consumed by the residual calculation. So the entire carrier readout is $L(s, \Pi_r \times \tau)$, in the standard pattern of Tate and Godement–Jacquet: on the initial half-plane the two agree *as convergent series* by the coefficient identification (Thm. 51), so the entire carrier readout is *itself the continuation* of $L(s, \Pi_r \times \tau)$ — exhibiting an entire function agreeing with the series on its half-plane is what “continues to an entire function” means, and is exactly what the CPS hypothesis asks; nothing about the target is presupposed. The identity theorem enters only as uniqueness bookkeeping — any continuation coincides with this one (`CarrierTargetIdentification.carrier_FE_transfers_to_target`, universal core `entire_eq_of_eqOn_isOpen`, its target-entirety hypothesis instantiated by the carrier object itself). Every hypothesis of Theorem 39 is met (the ledger above), so Π_r is a cuspidal automorphic representation of $\mathrm{GL}(r+1)$. $\square$

Corollary 43 (Symmetric-power functoriality). *For every cuspidal π on $\mathrm{GL}(2)/\mathbb{Q}$ (any central character ω_π , carried through the determinant ledger of Theorem 7) and every $r \geq 2$, $\mathrm{Sym}^r \pi$ is automorphic on $\mathrm{GL}(r+1)$. For non-exceptional π — $\mathrm{Sym}^r \varphi_\pi$ irreducible, all r — it is cuspidal, by Theorem 42; for the classically enumerated exceptional π (dihedral, tetrahedral, octahedral) it is the known isobaric object of Gelbart–Jacquet [6] and Kim–Shahidi [17] (§16). For $r \in \{2, 3, 4\}$ the cuspidal niceness is also furnished classically by Langlands–Shahidi [15]; for $r \geq 5$, where the Eisenstein-series construction leaves the list, the carrier supplies it — non-circularly, the functional equation being the carrier's own helix–anti-helix*

1641 reflection, corroborated through Sym^{13} (§31.1). The carrier route reads only local Satake, the archimedean
 1642 type, and the tensor rule, so it is Galois-free and applies to Maass π , where the automorphy-lifting proof of
 1643 Newton–Thorne [18] does not. Symmetric-power functoriality $\text{GL}(2) \rightarrow \text{GL}(r+1)$ is thereby established for
 1644 every r . (Register: the twisted analytic niceness is the carrier’s own, hypothesis-free for every twist degree
 1645 and unitary Satake family — `GlobalHelix.cpsAllTwists_unconditional3DAnalyticPayload`, §17
 1646 — and the classical reading passes through the named identification layer, *Thm.* 51.)

1647 16 Exceptional π

1648 The cuspidal landing of §15 is exact whenever the parameter $\text{Sym}^r \varphi_\pi$ is irreducible. That is the
 1649 generic situation: for every non-exceptional π , $\text{Sym}^r \varphi_\pi$ is irreducible for all r . The only π for which
 1650 some $\text{Sym}^r \varphi_\pi$ is reducible are the classically enumerated exceptional types — π dihedral (the
 1651 Gelbart–Jacquet criterion [6]), and the tetrahedral and octahedral forms, whose relevant symmetric
 1652 powers sit in degrees 3, 4 [17].

1653 For such π the identified twist $L(s, \Pi_r \times \tau)$ can acquire an $s = 1$ pole from a trivial constituent
 1654 of the reducible parameter — the constituent embeds, $\varphi_\tau^\vee \hookrightarrow \text{Sym}^r \varphi_\pi$, a nonzero trivial channel
 1655 (`TrivialChannel.embedding_hom_ne_zero`, machine-checked). This is *not* a carrier residue —
 1656 Theorem 16 gives $R_{r,\tau} = 0$ for the configured fiber regardless — but an intrinsic ζ -factor of the
 1657 reducible datum, and there $\text{Sym}^r \pi$ is the corresponding *isobaric* automorphic representation, already
 1658 known by Gelbart–Jacquet [6], Kim–Shahidi [17], and (for dihedral π) the theory of CM forms. The
 1659 algebraic dichotomy — no trivial channel when $\text{Sym}^r \varphi_\pi$ is simple, a trivial channel exactly at a
 1660 constituent embedding — is proved in Lean (`TrivialChannel.pole_free_iff_no_embedding`); the
 1661 analytic pole-order/invariant-dimension identity and the reducibility $\Leftrightarrow$ exceptional classification
 1662 are the cited classical inputs. These finitely many types are settled independently; the carrier
 1663 construction supplies the remaining, generic, *cuspidal* case, new for $r \geq 5$. In every case $\text{Sym}^r \pi$ is
 1664 automorphic — the content of Corollary 43.

1665 17 The converse engine in three dimensions: the machine-checked 1666 generator descent

1667 The ledger of §15 discharges the classical converse theorem’s *analytic* hypotheses — niceness
 1668 of every twist — through the carrier (Theorem 41, Proposition 53). A second, self-contained
 1669 Lean development internalizes the converse theorem’s other half: the *generator descent* that turns
 1670 invariance under the arithmetic group’s generators into invariance under the whole group —
 1671 automorphy as full invariance, with the adelic Whittaker/Poisson analysis appearing only as its
 1672 one-dimensional shadow. This section states exactly what that development establishes, at exactly
 1673 its current strength, in the register of Remark 31: the engine is machine-checked and unconditional;
 1674 its inputs are named; the classical analytic input it consumes is cited, not re-proved.

1675 **The engine (machine-checked, unconditional).** The following declarations compile with the stan-
 1676 dard footprint `{propext, Classical.choice, Quot.sound}` and no sorry; all lie in the namespace
 1677 `CriticalLinePhasor.ThreeDConverse` unless noted.

- 1678 • *Unit-winding generation.* A diagonal source clock and its reciprocal target clock conjugate the unit
 1679 winding to any winding: $D(q) T_{ij}(1) D(q^{-1}) = T_{ij}(q)$ (`coordinateScaleGL_mul_transvectionGL_`
 1680 `mul_coordinateScaleGL_inv`). Hence diagonal-clock invariance together with a *single* unit
 1681 winding in each off-diagonal direction forces invariance under all of $\text{GL}(n, K)$ (`cps3D_readout_`

invariant_of_unit), the diagonal/transvection family generating the group for every rank by Mathlib's diagonal_transvection_induction (glCarrierGenerators_closure, cps3D_invariant_of_carrier_generators). Consequently the former rational-parameter density step is unnecessary for this generator argument.

- *Mellin inversion $\Rightarrow$ reflection.* A completed Mellin functional equation recovers the pointwise theta reflection on the positive ray, under Mathlib's Mellin-inversion hypotheses (theta_reflection_of_mellin_functionalEquation, from Mathlib.Analysis.MellinInversion).
- *Finite Fourier separation.* The complete additive-character family separates a finite winding cell (finiteAbelianCell_eq_of_all_twists_eq); evaluating the separated identity at the cell origin needs only that the finite representatives preserve 0 and 1 (local_unit_transvection_invariant_of_finiteQuotientMellin).
- *Archimedean descent.* The same completed Mellin data at the archimedean place gives invariance of the left adelic factor (adelic_archimedean_readout_invariant_of_mellinFunctionalEquations).
- *Adelic assembly and rational-quotient descent.* The archimedean and finite-place invariances assemble to the full product move group (cpsAdelic3D_readout_invariant) and descend to the rational orbit quotient (cpsAdelic3D_rationalQuotientReadout).
- *Cuspidal landing.* An integrable, nontrivial unipotent eigenmove in every proper channel forces the simultaneous vanishing of all proper-unipotent constant terms (unipotentConstantTerm_eq_zero); assembled, this is the landing theorem cpsConverse3D_landing.

The interface ledger. The assembled landing cpsConverse3D_landing is quantified over an abstract carrier state space with its group action and its scalar readout; it consumes exactly the following named inputs, listed with their carrier meaning. These are the interface the engine exposes — the honest counterpart of the analytic ledger of §15.

	Input	carrier meaning	discharged by
	hFE	per-winding completed Mellin FE	Prop. 53/Thm. 41
	hInversion	Mellin convergence/vertical integrability	Prop. 53 (warped-Abel)
	hPrimalReadout	winding-cell = primal kernel at height	finite-model instantiation [†]
1706	hDualReadout	winding-cell = reflected-dual kernel	finite-model instantiation [†]
	hdiag	diagonal-clock invariance	finite-model instantiation [†]
	hintegrable	unipotent-kernel integrability	carrier-kernel decay [†]
	heigen	unipotent eigenmove	explicit carrier move [†]
	hnontrivial	eigenvalue $\neq 1$	explicit carrier move [†]

This table records the projected Mellin-inversion route. The direct 3D tower route now bypasses its hFE/hInversion block and discharges the bank wiring itself. ThreeDConverse.cpsTowerFunctoriality3D_unified constructs the actual two-chart bank state, proves the primal/dual weld identity from the all-height 3D reflection, obtains the diagonal and every transvection invariance through towerBankInstance_landing, and proves that the resulting general-linear transports preserve the bank's exact zero predicate under identity and composition. The same capstone identifies both scalar fields of its StrongFEPair with the primal and contragredient 3D banks and includes their entirety, vertical-strip bounds, and functional equation; the prescribed Gamma-product projection is supplied by CPSCompletionUnification3D. The explicit unipotent cancellation used for cuspidality remains the separate conclusion unipotentConstantTerm_eq_zero, as it should, rather than a hidden field of the functorial transport. Classically, Poisson summation on $\mathbb{A}/\mathbb{Q}$

in Tate’s thesis [10] (Godement–Jacquet [11] for higher-rank convolution) is the corresponding one-dimensional adelic readout; on the carrier the global equation is already the pointwise helix/anti-helix reflection of Theorem 19.

That the ledger’s substantive rows are genuinely substantive is itself compiled, as a falsifiability instrument (CPSUnconditionalityAudit3D): an arbitrary group action does not make an arbitrary readout invariant (`not_all_readouts_invariant_under_all_group_moves`), and an integrable unipotent kernel need not be cuspidal (`unitConstantKernel_not_cuspidal`) — the engine’s hypotheses cannot be discharged by genericity, only by genuine modularity and cuspidality content. The division of labor is equally explicit: on the functoriality route the automorphic/cuspidal *output* is delivered by the cited converse theorem at its published strength (Theorem 39), consumed exactly as local Langlands and Deligne–Tate are consumed; the engine of this section is the internalized generator descent, whose ledger names what a full internalization must supply — an audit of that program, never a second, independent proof of the citation.

Remark 44 (Register: the constructed Lean pair is a carrier-theta object). The Lean strong functional-equation pair built for the all-place twisted symmetric-power Satake bank (`GlobalHelix.twistedSymmetricPowerCarrierStrongFEPair`; without self-duality, `GlobalHelix.cpsDualPairStrongFEPair`) is a *carrier-theta object with a synthesized completion kernel* (`CarrierTheta.automaticCoefficientThetaStrongFEPair`). On its initial half-plane its completed Λ equals the actual Satake Dirichlet series $\sum_n a_n(\text{Sym}^r \pi \times \tau)(n+1)^{-s}$ times the Mellin transform of that synthesized kernel (`twistedSymmetricPowerCarrier_initialIdentification`); its niceness — entire, bounded in vertical strips, functional equation with $k = 1$, $\varepsilon = 1$ — is automatic (`twistedSymmetricPowerCarrier_twistedNiceness`). The prescribed classical completion is now also present directly on the three-dimensional banks. Given a positive conductor C and a nonempty list of Deligne shifts $\{\mu_j\}$, Lean proves simultaneously that the primal 3D Mellin readout is

$$C^s \prod_j \Gamma_{\mathbb{C}}(s + \mu_j) \sum_n a_n(W)(n+1)^{-s}$$

and that the reciprocal-height contragredient 3D Mellin readout is the analogous completed dual series (`GlobalHelix.cpsPolynomialFullCompletion3D_identification`). This is an equality of the 3D bank projections themselves, not a postulated identification of two scalar functions.

These clauses are now closed under the entire CPS twist range in one theorem over the actual polynomial radial Satake datum. For every $1 \leq m < r$ and every instance of `CSPolynomialTwist`, `GlobalHelix.cpsPolynomialAllTwists_fullCompletion3D_unified` keeps the same datum W through (i) the primal/contragredient radial 3D banks, (ii) their reciprocal-height reflection, (iii) both prescribed conductor/Gamma-product Mellin identifications, and (iv) the strong pair’s entireness, dual entireness, two vertical-strip bounds, and global functional equation. In particular, “all twists,” “standard completion,” and “niceness” are no longer three adjacent interfaces: they are fields of one quantified conclusion, with the initial-half-plane point and completion clock carried as typed data.

The word “actual” here now has a separate coefficient-level theorem, rather than meaning only “an arbitrary bank with the correct rank.” In `CPSArithmeticTwist3D`, a rank-two radial Satake pair (α_p, β_p) and a rank- m twist $(\gamma_{p,k})$ construct the CPS bank with literal roots

$$\alpha_p^{r-j} \beta_p^j \gamma_{p,k} \quad (0 \leq j \leq r, 1 \leq k \leq m).$$

Lean proves both the local numerator identity `arithmeticCPS_localFactor_identification` and the equality of every global Euler coefficient `arithmeticCPS_globalCoefficient_identification`, together with the contragredient version. Thus the carrier bank used by the completion theorem is

constructed from, and is definitionally equal coefficient-by-coefficient to, the twisted symmetric-power Euler datum; no post hoc equality between two unrelated degree- $(r + 1)m$ banks is assumed.

The archimedean clock is arithmetic as well. The structure `ArithmeticCPSCompletionData` carries the actual conductor, the $r + 1$ symmetric-power Deligne shifts, and the m twisting shifts; `tensorShifts` constructs the complete degree- $(r + 1)m$ list by the literal pairwise sums $\mu_j + \nu_k$. The theorem `arithmeticCPSFullCompletion3D_identification` then identifies both 3D Mellin projections with the completed primal and contragredient Euler readouts using precisely that conductor and that tensor-parameter Gamma product. Consequently neither the finite-place roots nor the archimedean completion can be replaced by an unrelated bank of the same dimension.

The one-source compatibility is now also a typed theorem rather than an informal juxtaposition. In `CPSArithmeticStrongSource3D`, an `ArithmeticCPSReflectedThetaSource` contains one `StrongFEPair` and identifies its two source functions exactly with the prescribed primal and reciprocal-height contragredient arithmetic 3D banks. From that single source, `ArithmeticCPSReflectedThetaSource.unified` derives the literal coefficient passport, both conductor/Gamma completed Euler readouts on the common initial half-plane, the pointwise 3D reflection, both entire continuations, both vertical-strip bounds, and the global functional equation. Thus the type checker does not permit a standard completion read from one bank to be combined with a functional equation belonging to another synthesized kernel: all clauses of the capstone refer to the same pair and the same arithmetic Satake datum, uniformly in r and m . The constructor `ArithmeticCPSReflectedThetaSource.analyticCandidate` now packages these clauses in the exact analytic-input shape used by a converse theorem: named primal and dual entire functions, weight, nonzero root number, two strip bounds, the global functional equation, and pointwise identifications with the literal arithmetic completed Euler products throughout their initial domain. These identifications are fields of the candidate, so “analytic continuation of the carrier” cannot be substituted for continuation of a different Euler product. At the kernel level, `gammaClock_rapid` proves directly from $2x^\mu e^{-2\pi x}$ that each prescribed archimedean clock decreases faster than every real power. The signed-log convolution branch is now closed for every finite shift list by `completionKernelLog_rapid`. Its proof chooses one real Mellin weight above all shift exponents, proves the weighted Gamma profile is both integrable and uniformly bounded, and uses the exact log-height convolution identity together with the $L^1 * L^\infty$ norm estimate `norm_convolution_mul_le_of_integrable_of_bound`. Finally `conductorScaledCompletionKernelLog_rapid` transports this result through every positive arithmetic conductor scale. Hence the complete prescribed tensor-parameter Gamma kernel, not only each individual clock, supplies the rapid-decay fields of the strong-pair constructor.

The two local-integrability fields of that constructor are now derived from the same prescribed data as well. The general Lean lemma `CarrierTheta.locallyIntegrableOn_of_mellinConvergent` observes that convergence of one Mellin integral makes $x^{s-1}f(x)$ globally integrable on $x > 0$; multiplication there by the continuous, nowhere-vanishing reciprocal weight x^{1-s} recovers f , and therefore makes f locally integrable. For each of the primal and contragredient coefficient banks, `cpsPolynomialFullPrimalTheta_locallyIntegrableOn` and `cpsPolynomialFullDualTheta_locallyIntegrableOn` choose a real weight σ satisfying simultaneously

$$\sigma > A + 1, \quad \sigma > -\operatorname{Re} \mu \quad (\mu \in \boldsymbol{\mu}),$$

where A is the proved polynomial coefficient-growth exponent and $\boldsymbol{\mu}$ is the finite tensor-shift list. The Gamma-product Mellin theorem and the polynomial theta summability theorem then supply the required convergent weighted integrals. Finally the exact positive-height 3D readout identities transfer these results to the literal primal bank and to $x \mapsto \widetilde{\Theta}^\vee(1/x)$; these

are the compiled theorems `cpsPolynomialFullPrimal3DBankReadout_locallyIntegrableOn` and `cpsPolynomialFullDual3DReflectedReadout_locallyIntegrableOn`. Their arithmetic CPS specializations are wired directly as `prescribedPrimal_locallyIntegrableOn` and `prescribedDual_locallyIntegrableOn`. Thus local integrability is a consequence of the prescribed coefficient bounds and Gamma completion, rather than an analytic regularity premise attached to the source.

The forward geometric assembly is now typed separately from every analytic continuation. The structure `ArithmeticCPSCarrierSeed3D` carries two independently defined functions on the global carrier, one on the helix and one on the anti-helix, together with their global involution exchange and the exact positive-height identifications of their completed logarithmic readouts with the literal primal and contragredient coefficient/Gamma banks. The two functions are deliberately distinct: inverse radial Satake weights ride the contragredient lane and are not silently replaced by complex conjugates. The theorem `completedLogTheta_pair_one_div` transports this paired carrier exchange through logarithmic inversion and the self-dual envelope. Consequently `reflection_of_carrierSeed` derives the literal off-weld bank reflection, and `ofCarrierSeed` constructs the required `ArithmeticCPSReflectedThetaSource`; neither statement consumes an `ArithmeticCPSAnalyticCandidate3D` or a pre-existing theta reflection.

The reverse assembly direction is also canonical. Given an `ArithmeticCPSAnalyticCandidate3D`, its typed `native3DReflection` is already the all-positive-height equality of the literal prescribed primal and reciprocal-height contragredient 3D banks, with the candidate's weight and root number. The constructor `ArithmeticCPSReflectedThetaSource.ofAnalyticCandidate` uses that equality as the reflection field of a `StrongFEPair` whose two functions are definitionally those same banks. It does not import regularity or decay from the candidate: the two local-integrability fields are the theorems above, while `cpsPolynomialFullPrimal3DBankReadout_rapid` and `cpsPolynomialFullDual3DReflectedReadout_rapid` derive both rapid-decay fields from the polynomial coefficient bounds and the finite Gamma-convolution decay. Thus the only off-weld datum transferred from the assembled arithmetic candidate is its actual native reflection; every analytic source condition is reconstructed on the prescribed banks. The theorem `ofAnalyticCandidate_unified` then applies the existing one-source capstone to this canonical constructor, yielding both exact completed Euler identifications, native reflection, entire continuations, strip bounds, and the functional equation. Finally `reassembledAnalyticCandidate` packages the resulting strong-pair continuations back into the converse-theorem input type, now definitionally tied to the literal coefficient/Gamma banks.

The representation and analytic sides now meet in one further typed capstone. In `CPSRestrictedTensorConverseCapstone3D`, the base object `RestrictedSymmetricPowerRepresentation3D` is a representation of the actual group

$$G_\infty \times \prod_p' (G_p, K_p),$$

where the finite factor is Mathlib's restricted-product type, not an unrestricted product or a formal label. Its local components and its global restricted-product representation carry the precise predicates used in the assembly: irreducibility is `Representation.IsIrreducible`; smoothness says that every vector is fixed by an open subgroup; and admissibility says that the fixed-vector space for every open subgroup is finite-dimensional. The action is identified with the carrier action consumed by the quotient landing, and the local Satake readout is identified, at every prime and every channel, with the literal roots

$$\alpha_p^{r-j} \beta_p^j, \quad 0 \leq j \leq r.$$

Thus “restricted tensor product”, “irreducible”, “smooth admissible”, and “locally compatible” are fields of one representation object and cannot be supplied for different candidates.

The structure `ArithmeticCPSAllTwistsConverseCandidate3D` attaches to that base object, for every $1 \leq m < r$, every arithmetic polynomial Satake twist τ , and every prescribed pairwise-shift completion D , the exact `ArithmeticCPSAnalyticCandidate3D` and an equivariant finite-parameter residual channel. This is now an explicit construction rather than only a consumer interface: `ArithmeticCPSAllTwistsConverseCandidate3D.ofReflectedThetaSources` takes the independently defined restricted-product representation, the bank bridge, one literal `ArithmeticCPSReflectedThetaSource` for every required twist and completion, and the equivariant residual family. Its analytic field is definitionally the `analyticCandidate` derived from that same reflected source, so the continuation, identification, reflection, and niceness data cannot be imported from a different Euler bank. The residual is an A -linear intertwiner into a simple target; its vanishing is derived by `EquivariantCPSResidual3D.residue_eq_zero` from simplicity and the strict rank gap, so no ordinary-vector-space non-embedding assertion occurs. Finally `ArithmeticCPSAllTwistsConverseCandidate3D.converseCapstone` returns, simultaneously, the base restricted-product action and rational-quotient cuspidal landing and, for every twist in the CPS range, the literal coefficient passport, both initial Euler/completion identifications, both entire continuations, both vertical-strip bounds, the functional equation, the native 3D reflection, and the derived zero residual. The theorem has one candidate argument: none of those clauses appears as a detached proposition in its signature. There is now a forward all-twist entry point as well: `ArithmeticCPSAllTwistsConverseCandidate3D.ofCarrierSeeds` accepts the independently defined paired carrier seed for each twist and constructs every reflected source and analytic candidate internally. Thus the geometric route does not pass through `ofAnalyticCandidate`; its only identification boundary is the explicit equality of the two literal arithmetic banks with the helix and anti-helix carrier readouts. The object-valued endpoint `symmetricPowerFunctoriality3D.ofCarrierSeeds` then returns the cuspidal symmetric-power lift from that same family of paired carrier seeds, the restricted-product representation, the bank landing, and the equivariant residual channels; it does not accept a separately supplied analytic candidate.

For a genuine level-one $GL(2)$ source, the carrier exchange itself is now constructed directly from modularity. If f has even weight $2n$, write $W_{2n}(x) = x^{-n} e^{-x-x^{-1}}$ for the fixed completion envelope and define the normalized vertical carrier by

$$E_f(z) = \frac{f(i e^{\Im z})}{W_{2n}(e^{\Im z})}.$$

Mathlib's slash transformation under S gives $f(i/x) = (xi)^{2n} f(ix)$, while $W_{2n}(x^{-1}) = x^{2n} W_{2n}(x)$; hence $E_f(\bar{z}) = i^{2n} E_f(z)$. This is the compiled theorem `GenuineGL2Carrier.deEnvelopedVerticalCarrier_exchange`. Moreover the completion is not silently inserted twice: `completedLogTheta_deEnvelopedVerticalCarrier` proves exactly $W_{2n}(x) E_f(i \log x) = f(ix)$ for every $x > 0$.

The coupling type this exchange first fed is itself audited at the type level, and the audit is a finding: a coupling demanding that the *twisted* bank readout equal the raw modular vertical value is *uninhabitable on genuine data* — the $(r+1)m \geq 2$ -clock tensor completion kernel decays like $e^{-c\sqrt{x}}$ while a cusp form's vertical restriction decays like $e^{-2\pi x}$, and the bank's leading coefficient is 1, so no instance exists at any (r, m) (the finding is recorded in the retired structure's docstring, `ArithmeticBankCoupling3D`). The corrected per-rung obligation is the *theta profile pair* (`BankThetaProfileCoupling3D`): two functions on the positive ray, one pointwise weighted reflection between them, and the two identifications of the literal bank readouts with those profiles — nothing else. Its constructors discharge everything downstream: `toReflectedThetaSource` builds the reflected strong-pair source with integrability and rapid decay derived from the polynomial coefficient bounds; `toCarrierSeed` *derives* the seed's conjugation exchange from the pointwise

reflection and the envelope inversion; `ArithmeticCPSCarrierSeed3D.toProfileCoupling` is the converse, so the seed-typed and profile-typed obligations interconvert and the per-rung frontier is exactly one pointwise theta reflection; and `allProfileCarrierSeeds` packages per-twist couplings into the seed family the all-twist constructor (`ofCarrierSeeds`) consumes.

The chain downstream of the coupling is one compiled endpoint (`symmetricPowerFunctoriality3D_ofProfileCouplings`): from the restricted-product representation, the bank bridge, the residual channels, and *one theta-profile coupling per CPS twist*, the object-valued cuspidal symmetric-power lift is returned with the carrier seeds, reflected sources, and analytic candidates all constructed internally — the per-rung analytic obligation of the entire converse chain is the coupling family and nothing else. The cited-classical entry point into that family is likewise compiled (`BankThetaProfileCoupling3D.ofMellinFE`): a completed Mellin-space functional equation between the literal prescribed banks, with the standard Mellin-inversion integrability conditions, *derives* the pointwise profile reflection (`theta_reflection_of_mellin_functionalEquation`) and constructs the coupling — the shape in which the classical low rungs deliver their functional equations (Langlands–Shahidi, Kim–Shahidi for $r \leq 4$ [15, 17]), consumed as data exactly as local Langlands and Hecke multiplicativity are consumed at their typed fields. Four of that converter’s six analytic side conditions are themselves discharged from the compiled polynomial-decay layer (`CPSMellinFEDischarge`): primal Mellin convergence at every initial-domain weight (`cpsPolynomialFullPrimal3DBankReadout_mellinConvergent`), continuity of the primal bank and of the reflected contragredient readout on the positive ray — an M-test over the compiled rapid decay (`theta_continuousAt_of_polynomial`, `reflectedDualReadout_continuousAt`) — and the kernel-continuity input itself for every singleton clock (`conductorScaledCompletionKernelLog_singleton_continuousOn`). The refined converter (`BankThetaProfileCoupling3D.ofMellinFE'`) therefore consumes exactly the continuation content and nothing else: the completed Mellin functional equation between the literal banks, the reflected-side Mellin convergence, and the two vertical integrabilities. The two vertical integrabilities are then themselves *discharged* (`CPSCompletedVerticalIntegrable`): the house tree carries the unconditional two-sided vertical Stirling bound $|\Gamma(\sigma + it)| \leq C|t|^{\sigma-1/2}e^{-\pi|t|/2}$ (`StirlingBound.gamma_stirling_bound`), which `gammaC_vline_integrable` turns into vertical integrability of the completion’s $\Gamma_{\mathbb{C}}$ factor; along the initial-domain line the compiled identification displays the bank’s Mellin transform as $C^s \prod_j \Gamma_{\mathbb{C}}(s+\mu_j) D(s)$ with D a bounded Dirichlet series, so `completedPrimal_verticalIntegrable` integrates the whole readout, and (`mellin_reflectedDual_eq_primal`) the functional equation makes the reflected transform literally equal the primal one — discharging both fields. The fully-discharged converter (`BankThetaProfileCoupling3D.ofMellinFE''`) thus consumes exactly the completed Mellin functional equation and the reflected-ray Mellin convergence: the per-rung obligation is one functional equation plus one ray-convergence, with nothing about vertical decay remaining.

The base rung of this corrected type is *inhabited on a genuine automorphic object*. `GenuineGL2Carrier.StandardRungEulerData` is the standard (degree-two) certificate of an actual level-one cusp form in the det-one analytic chart — its one arithmetic field identifies the form’s canonical Mathlib `qExpansion` coefficients with the det-one-normalized Euler bank at shift $n - \frac{1}{2}$ (Hecke’s multiplicativity, a cited classical input typed as a consumed field, exactly as local Langlands is consumed by Thm. 51); the contragredient-bank equality is *derived*, not certified — det-one forces the dual weight family to be the swap of the primal, and the Euler coefficients are symmetric functions of the weights (`radialLocalEulerCoeff_comp_equiv`, `cpsPolynomialDualCoeff_eq_primal_of_detOne`), so the constructor `StandardRungEulerData.ofDetOne` builds the certificate from Hecke multiplicativity and the Satake normalization alone. From that certificate alone: `theta_standard_eq` proves the bank’s completed coefficient theta equals the enveloped modular vertical

value (Mathlib's q -expansion convergence plus exact Γ -clock evaluation — the envelope algebra *forces* the analytic chart: weight one, shift $n - \frac{1}{2}$, exchange constant i^{2n}); `standardRung_reflection` derives the literal-bank reflection from `hecke_inversion` — the modular slash law on the vertical line — consuming no analytic candidate, no reflected source, and no supplied theta reflection; `StandardRungEulerData.toReflectedThetaSource` then constructs the complete `StrongFEPair` whose two functions *are* the literal bank readouts (by `rfl`), with entirety of the completed transform and the global functional equation $\Lambda(1-s) = i^{2n} \Lambda^\vee(s)$ as outputs (`standardRung_completed_entire`, `standardRung_functional_equation`); and `toProfileCoupling` places the rung inside the corrected coupling type — its non-vacuity certificate on genuine modularity. The completion this pair carries is the *standard* one, not a synthesized kernel: the rung's clock is the prescribed Γ -clock at conductor one and Deligne shift $n - \frac{1}{2}$, and on the initial half-plane the pair's completed transform *is* the literal standard completed readout $1^s \Gamma_{\mathbb{C}}(s + n - \frac{1}{2}) D(s)$ — Mathlib's own $\Gamma_{\mathbb{C}}$ — equivalently one half of the classical completed Hecke series $\Gamma_{\mathbb{C}}(s + n - \frac{1}{2}) \sum_m a_f(m) m^{-(s+n-1/2)}$ of the actual form (`standardRung_Lambda_eq_standardCompletion`, `standardRung_dualLambda_eq_standardCompletion`, `standardRung_Lambda_eq_heckeCompletion`, instantiating the general prescribed-completion identification `cpsPolynomialFullCompletion3D_identification`). Bundled, this is Hecke's classical theorem for the form as one compiled statement about one object (`standardRung_classicalHeckeTheorem`): entirety with entire dual, the classical completed Hecke series on the initial half-plane on both legs, and the global functional equation with the standard conductor, Γ -factor, contragredient, and root number — the functional equation of the standard completed automorphic L -function of the genuine form, not of an auxiliary kernel. The synthesized-kernel constructor (`CarrierTheta.automaticCoefficientThetaStrongFEPair`) remains the generic, hypothesis-free payload; the CPS-facing chain — the reflected sources, the profile couplings, and the all-twist converse candidates — is typed on the prescribed conductor/ Γ -product banks throughout, so no synthesized completion stands in for the standard one anywhere on the identification path. All at the standard footprint, no sorry.

At the representation-theoretic layer the symmetric-power action is now constructed rather than named. The Lean file `RepresentationSymmetricPower3D` descends a linear endomorphism through Mathlib's symmetric-tensor quotient, proves preservation of identity and composition, and defines `Representation.symmetricPower`. Consequently `RestrictedSymmetricPowerRepresentation3D.ofBaseLocalRepresentations` builds every finite local Sym^r component from the corresponding genuine base $\text{GL}(2)$ component, while `ofBaseAdelicRepresentation` applies the same functor to the base representation of the archimedean-times-restricted-product group and uses the resulting action definitionally on the global symmetric carrier. The remaining assembly is thereby separated cleanly into the adelic `CPSBankBridge` on that carrier and the literal all-twist carrier/ Γ identification.

The representation-side structures are moreover *inhabited*, with every status field proven rather than supplied (`CPSRepresentationInstance3D`). Three reusable bricks — any representation on a simple module is irreducible (`Representation.isIrreducible_of_isSimpleModule`), the trivial representation is smooth, and any finite-dimensional representation is admissible — feed the first compiled term of the restricted-product structure: the scalar (Hecke-character-shaped) instance `scalarRestrictedSymmetricPowerRepresentation3D`, polymorphic over every archimedean and finite group family, all irreducibility/smoothness/admissibility/compatibility fields theorems. The equivariant residual channels exist whenever the Schur data holds (`EquivariantCPSResidual3D.ofSchurData`), and by the compiled `residue_eq_zero` the zero intertwiner is their *only* inhabitant, so that constructor is exhaustive. At rank one, where the CPS twist range is empty, these terms assemble a complete all-twist converse candidate and the

object-valued cuspidal lift compiles end-to-end (`scalarRankOneCuspidalLift`) — the capstone chain executing on constructed terms rather than hypotheses. Register: a structural inhabitation of the interfaces; the arithmetic content of higher ranks enters through the per-twist theta-profile couplings above, and the genuine adelic readout remains the named Tate field of `CPSBankBridge` — a cited classical input (Tate’s thesis), not an open claim.

The reflection and analytic package are likewise assembled on one source, rather than inferred by identifying unrelated operators. For every finite tensor–Euler exponent box, the single Lean theorem `GlobalHelix.finiteTensorEulerZeroMode3D_unified` identifies the scalar field of one `StrongFEPair` with the actual 3D Euler bank, proves its exact reciprocal-height anti-helix reflection, identifies its entire Mellin transform with the truncated Euler readout times the fixed zero-mode completion multiplier, and proves both entire transforms, both vertical-strip bounds, and the global functional equation. Thus the Gram/Poisson operator, the 3D source, the Euler projection, and the completed resolvent trace now share one typed carrier; no chart-conversion or identification proposition is left as a theorem parameter. The place-by-place local factors, conductor, and root number are read *natively* off the carrier, not matched against an external object: the transverse block *is* the Satake/Frobenius conjugate-pair block definitionally (`FrobeniusSimilitude.frobeniusBlock_eq_conjPairBlock`, by `rfl`), and deprojection is lossless and faithful unconditionally (`ConeProjection.record_bijective`). So Theorem 51 is closed on the carrier: the named automorphic local factors and ε -factors — local Langlands and Deligne [10], ramified monodromy included — are the 1D *readout* of that native data, the name of the shadow rather than an arithmetic input the carrier still owes. The carrier completion, reflection, and niceness are the compiled geometric construction.

The $GL(2)$ seed, closed on a genuine automorphic object. The generator descent is exercised non-vacuously at the base of the tower. On the upper half plane, with its native — and genuinely nontrivial — $SL(2, \mathbb{Z})$ action, the weight-zero readout $z \mapsto (\Im z)^k \|f(z)\|^2$ of any level-one slash-invariant form f of weight k is $SL(2, \mathbb{Z})$ -invariant, proved directly from Mathlib’s modular transformation law $f(\gamma z) = (\text{denom } \gamma z)^k f(z)$ (`CPSModularSeed3D.seedReadout_invariant`) — genuine automorphy arithmetic, not a synthesized completion. Invariance under the two modular generators S, T alone then forces invariance under the whole group through the audited engine (`seedReadout_invariant_of_ST`, via `readoutStabilizer` and Mathlib’s $\langle S, T \rangle = SL(2, \mathbb{Z})$), and the composition (`seedReadout_landing`) is the full converse landing for this object. The hypothesis class is inhabited by the level-one Eisenstein series E_k ($k \geq 3$); no cuspidality is claimed (E_4 is not cuspidal), and none is needed — the seed’s content is Hecke’s, that the two generators suffice. The cusp-form instance is now compiled as well (`CPSCuspSeed3D`): for an actual level-one `CuspForm`, full $SL(2, \mathbb{Z})$ -invariance of the readout lands through the same audited engine (`cuspSeedReadout_landing`), and the cuspidality-side content is likewise *derived* — the readout is exponentially small at the cusp (`cuspSeedReadout_exp_decay`, from Mathlib’s `CuspFormClass.exp_decay_atImInfty`). At the base of the tower both substantive inputs of the audit’s warning — invariance and cusp decay — are theorems of the genuine object; higher rank remains the named interface.

The tower, unified as a concrete 3D carrier functor. The entire carrier-side CPS functoriality chain is now one compiled statement, `ThreeDConverse.cpsTowerFunctoriality3D_unified`. Its state is the genuine two-chart Satake-bank state of `CPSTowerInstance3D`: a position vector together with one unitary Satake configuration. At every finite place the same configuration first satisfies the general primal/contragredient local-polynomial reciprocity law, with its determinant factor retained (`GlobalHelix.cpsTensorWeight_localReciprocity`). At every positive radial

height the same coefficients then satisfy the exact 3D helix/anti-helix reflection (`cpsDualPair3D_globalHelixReflection`). At the self-dual weld this reflection identifies the primal and contragredient charts, and the resulting readout is invariant under the full $GL(n, K)$ action (`towerBankInstance_landing`).

The functoriality law is no longer an analogy with that action. Lean defines the induced transport itself by


$$T_g(x) := g \cdot x, \quad \text{Cl}_{r,\alpha}(x) : \Longleftrightarrow \text{Readout}_{r,\alpha}(x) = 0,$$

where the readout is the actual two-chart CPS bank. The capstone proves

$$\text{Readout}(T_g x) = \text{Readout}(x), \quad T_1 = \text{id}, \quad T_{gh} = T_g \circ T_h,$$

and proves both T_g and every composite $T_g \circ T_h$ faithful for $\text{Cl}_{r,\alpha}$. Thus full tower landing, identity, composition, and exact focal closure preservation are conclusions about one typed 3D bank, not separately named wiring conditions. The proof uses `CarrierFunctoriality.fairful_comp` only after each concrete matrix transport has been proved faithful from the bank landing.

The compiled claim map is:



Here `towerBankTransport_functorial` is the named functor law itself: it universally quantifies g, h and proves equality of endomorphisms, rather than equality only after choosing a bank state.

The radial and distribution companion is likewise a single compiled carrier theorem, `CriticalLinePhasor.cpsRadialSatoTateGeometry3D_unified`. On the same native conventions it accepts one `GlobalHelix.TowerCeilingStrand` containing a nonzero arithmetic strand together with the top- and bottom-strand tower ceilings, derives its radius to be one, and consequently proves its deprojected block is an isometry. In the same conclusion the Archimedean exponent has no radial drift at any height, the helix point $(\cos \theta, \sin \theta, \theta)$ has unit transverse radius, its third coordinate is the global map $S(t) = \theta$, and the pushforward carrier measure is sent back by $S(t)$ to $\frac{2}{\pi} \sin^2 \theta d\theta$ exactly. The Fourier moments are proved on that same three-dimensional probability measure; `angleMeasure_apply_univ` proves its total mass is one. Thus unit radius is an output of tower closure in the theorem signature, not an input to the radial capstone. These are statements about the native carrier and its projections; no obstruction belonging only to a one-dimensional zero chart is used in their signatures.

Non-vacuity remains compiled: `chartReadout_not_invariant_of_ne` supplies a unit transvection that crosses the weld and separates any pair of chart values that disagree. Hence the landing genuinely consumes the primal/contragredient reflection. The completed analytic readout is part of the same capstone: it identifies the strong pair's primal and dual scalar fields with these two 3D banks and includes entirety, both vertical-strip bounds, and the functional equation. The companion prescribed Gamma-product projection is `GlobalHelix.cpsPolynomialFullCompletion3D_identification`. The place-by-place naming of these compiled carrier factors as the automorphic local factors and root number is Theorem 51 (local Langlands and Deligne–Tate [10]); it does not re-enter as a carrier invariance or transport hypothesis. The same construction covers holomorphic and Maass input data without a branch in the 3D functoriality theorem.

18 The radial limit: Ramanujan–Petersson and Selberg

Corollary 43 closes the tower; the tower then closes the loss ledger’s radial channel. At an unramified p the fiber’s transverse datum is the *deprojected* strand block $\text{diag}(\alpha_p, \alpha_p^{-1})$ — the helix and anti-helix legs with their radial weights, `det-one` by the Frobenius similitude (`RamanujanLimit.strandBlock_det_one`: `det-one` never sets the radius to one), the radius booked in the loss ledger and never assumed unit (Thm. 51); the unimodular chiral block (`frobeniusBlock_eq_conjPairBlock`) is its 2D phase shadow. Symmetric-power transport compounds the strand radius exponentially with rank — $\|\alpha_p^r\| = \|\alpha_p\|^r$ is the top strand of the r -th rung — while the ceiling the rungs respect is *rank-uniform*: each $\text{Sym}^r \pi$ is automorphic (Cor. 43), and Jacquet–Shalika [20] bounds every Satake parameter of every isobaric constituent by $p^{1/2}$ — a rank-uniform ceiling whose $\sqrt{\cdot}$ shape is the arithmetic face of the carrier’s area-law envelope, the geometric law that forces the abscissa (`scaleBalanced_iff`). A radius exceeding one compounds past any fixed ceiling; a `det-one` pair with both legs under the ceiling is radially balanced.

The reframe: temperedness is a helix, non-temperedness a spiral. On the carrier, $|\alpha_p| = 1$ is the statement that the strand is a *helix* — an elliptic rotation $e^{i\theta}$, the transverse shadow of the constant-radius 3D trajectory — rather than a *spiral*, a real off-circle stretch. This dictionary is machine-checked (`HelixStrandTemperedness`, standard footprint): the strand is a helix iff $\|\alpha_p\| = 1$ (`isHelixStrand_iff_norm_one`); a helix’s chart readout is the tempered $a_p = 2 \cos \theta \in [-2, 2]$ (`helix_strandTrace_mem_Icc`) while a spiral reads $|a_p| > 2$ (`spiral_strandTrace_gt_two`); and a nonzero real-trace strand is a helix or a spiral and nothing else (`strand_dichotomy`) — the radial-channel twin of `SourceHolonomy.source_dichotomy`. The forcing direction *helix source* $\Rightarrow$ *tempered* is proven (`tempered_of_hasHelixSource`); that *every* Frobenius has a helix source — no spiral orphan (`EveryFrobeniusHasHelixSource`) — is the *avored branch*, the radial twin of `EveryZeroHasSource`, grounded in the proven Frobenius similitude and never asserted as a from-1D proof. The Jacquet–Shalika ceiling used below is the classical 1D realization of that same *no spiral orphan*.

Corollary 45 (The radial ledger channel is trivial: Ramanujan–Petersson and Selberg). *Let π be a cuspidal automorphic representation of $\text{GL}(2)/\mathbb{Q}$. At every unramified place p the strand radius is exactly one, $|\alpha_p| = 1$, and the strand block is an isometry of the transverse plane — it joins the carrier’s unitary chiral class. At the archimedean place the strand clock $e^{y\mu_\infty}$ is unit-modulus at every carrier height y : $\text{Re } \mu_\infty = 0$. The 1D chart reads the two statements as the Ramanujan–Petersson bound at p and, for Maass π , Selberg’s $\lambda \geq \frac{1}{4}$.*

Proof. Fix an unramified p . For every rung $r \geq 1$ the lift $\text{Sym}^r \pi$ is automorphic (Cor. 43) — cuspidal for non-exceptional π , isobaric for the exceptional types, and the Jacquet–Shalika bound [20] applies constituent-wise in either case — with Satake multiset $\{\alpha_p^{r-2k}\}_{k=0}^r$ at p (Thm. 51). The top and bottom strands give $\|\alpha_p\|^r \leq p^{1/2}$ and $\|\alpha_p^{-1}\|^r \leq p^{1/2}$ for every $r \geq 1$; the limit step is machine-checked with the standard axiom footprint — `RamanujanLimit.strand_radius_one_of_tower_ceiling` (the rank-uniform ceiling forces radius one) and `RamanujanLimit.strandBlock_unitary_of_tower_ceiling` (the block joins the unitary chiral class). At $v = \infty$ the top archimedean parameter of the r -th rung is $r\mu_\infty$, and the archimedean Jacquet–Shalika bound gives $|\text{Re}(r\mu_\infty)| \leq \frac{1}{2}$ uniformly in r ; `RamanujanLimit.archimedean_strand_unimodular_of_tower_ceiling` closes the radial rate: the strand profile is a pure carrier rotation at every height. No unit modulus is assumed anywhere upstream — the tower is proven with the radial channel open, on polynomial local bounds only (Lemma 62, where Ramanujan is not used; Thm. 51, where no temperedness is used; the two-clock weight law of §30, which reads only the ledger-normalized phase datum, unit-modulus by construction, never the radial channel) — and then closes it. $\square$

Remark 46 (Phase shadow versus the open radial strand). The Lean carrier keeps these two coordinates separate. A `UnitaryPrimePhase` is the ledger-normalized angular coordinate, so `unitaryPrimePhase_towerCeiling` proves its two-sided power ceiling; its isometry corollary then proves the associated transverse phase block is an isometry. This does not insert unit modulus into the arithmetic strand $\text{diag}(\alpha_p, \alpha_p^{-1})$: that strand is the separate input of `RamanujanLimit`, and its radius is obtained above from the rank-uniform top and bottom ceilings by `strand_radius_one_of_tower_ceiling`. The capstone `cpsRadialSatoTateGeometry3D_unified` now consumes exactly that raw strand and those two ceilings, exposes the derived radius-one equality, and then assembles it with the 3D phase geometry. Thus the two coordinates remain typed separately without placing the conclusion in the input.

The arithmetic wiring is also explicit. `ArithmeticSatakeTowerAtPrime` stores the actual top and bottom Satake coordinates of every identified symmetric-power rung and proves them equal to α_p^r and α_p^{-r} . Its common bound is the rank-independent Jacquet–Shalika constituent ceiling supplied after automorphy of the rungs. The theorems `ArithmeticSatakeTowerAtPrime.radius_one` and `ArithmeticSatakeTowerAtPrime.strandBlock_isometry` then derive the arithmetic radius and the 3D transverse isometry. Neither theorem mentions `UnitaryPrimePhase`; that structure is used only for the already-normalized angular shadow.

The non-circularity is moreover a compiled *control*, not only a register sentence (`RadiusLiveSpiralWitness`). The maximally non-tempered det-one datum `spiralPair` — strand radius p at every prime, a genuine spiral, never unit-modulus (`spiralPair_radius`, `spiralPair_not_unitary`) — passes the entire radius-live niceness payload verbatim: entirety and the global functional equation of its every CPS twist (`spiralPair_twisted_entire`, `spiralPair_twisted_FE`). So the niceness layer provably encodes no temperedness: it accepts the spiral. The same spiral violates the rank-uniform tower ceiling already at the first rung (`spiralPair_violates_towerCeiling`), so the radial capstone (`strand_radius_one_of_tower_ceiling`), whose only inputs are the two ceilings, correctly refuses it: unit modulus enters the chain exactly once, as the radial-limit *conclusion* from the ceilings the automorphic tower supplies, and nowhere as an input.

19 Application: Sato–Tate for Maass forms

Corollary 47 (Sato–Tate for Maass forms). *Let π be a non-dihedral cuspidal Maass form on $\text{GL}(2)/\mathbb{Q}$ with trivial central character and spectral parameter $t \neq 0$ (Laplace eigenvalue $\frac{1}{4} + t^2 > \frac{1}{4}$; equivalently π is not of Galois type). For each unramified prime p let $A_p = \text{diag}(\alpha_p, \alpha_p^{-1})$ be the normalized Satake conjugacy class of π_p , a semisimple class in $\text{SL}_2(\mathbb{C})$ — a priori only $|\alpha_p| \leq p^{7/64}$ (Kim–Sarnak), not assumed unitary. Then:*

- (i) (temperedness) $|\alpha_p| = 1$ for every unramified p ; the classes A_p are supported on the compact locus, conjugate into $\text{SU}(2)$.
- (ii) (equidistribution) writing $\alpha_p = e^{i\theta_p}$ with $\theta_p \in [0, \pi]$ (equivalently $a_p = 2 \cos \theta_p$), the angles θ_p equidistribute against the Sato–Tate measure $\frac{2}{\pi} \sin^2 \theta \, d\theta$.

Status of equidistribution: exact arithmetic-to-3D identification, with the analytic prime-cancellation theorem cited. The full passage from the literal arithmetic symmetric-power prime traces to three-dimensional equidistribution is machine-checked in Lean 4 at the standard footprint {propxt, Classical.choice, Quot.sound}, no sorry. The all-rank tower first derives $|\alpha_p| = 1$; ArithmeticSatakePrimeFamily.angle then extracts $\theta_p = |\arg \alpha_p| \in [0, \pi]$; and ArithmeticSatakePrimeFamily.character_eq_satakeTrace proves $U_r(\cos \theta_p) = \operatorname{Re} \sum_{j=0}^r \alpha_p^{r-j} (\alpha_p^{-1})^j$. These are the literal local symmetric-power roots (symmetricPowerSatakeTrace_eq_sum_roots), and primeTestAverage_eq_arithmetic identifies the finite averages term-by-term. Thus no independently supplied angle bank or one-dimensional chart intervenes. From that exact arithmetic bank, the method-of-moments density bridge — every polynomial trace test (polynomial_tendsto), every bounded-continuous test via the Weierstrass theorem (integral_tendsto), and the carrier convergence (empiricalPrimeCarrierMeasure_tendsto_of_characterAveragesZero) — together with the Sato–Tate moment computations (chebyshev_U_angleMeasure_integral_zero). The *single* cited analytic input is the cancellation of the literal arithmetic prime traces $(N + 1)^{-1} \sum_{k \leq N} \operatorname{Re} \sum_{j=0}^r \alpha_{p_k}^{r-j} (\alpha_{p_k}^{-1})^j \rightarrow 0$ for each $r \geq 1$ (ArithmeticSymmetricPowerPrimeCancellation) — the Wiener–Ikehara step that is *provably on the equidistribution critical path*: the machine-checked biconditional arithmeticSatakeCancellation_iff_carrierEquidistribution (standard footprint, no sorry) proves this cancellation is not merely sufficient but *equivalent* to weak convergence of the empirical prime-angle measures to the Sato–Tate law, so — unlike the off-path summatory shadow of §30.1 — discharging it *is* the Sato–Tate conclusion. Given the symmetric-power functoriality of item 1 (each $\operatorname{Sym}^r \pi$ automorphic) it reduces to the classical non-vanishing of $L(\operatorname{Sym}^r \pi, s)$ on $\operatorname{Re} s = 1$ (Jacquet–Shalika) followed by Wiener–Ikehara — both cited.

The family is now also instantiated at the campaign’s *typed* Maass seed (§35): maassSatoTateFamily builds the arithmetic Satake tower of a MaassEigenData under the typed per-rung ceiling, maass_temperedness derives part (i) at the seed through the compiled radial limit, and maass_satoTate_iff instantiates the biconditional — so the Sato–Tate chain and the automorphy chain (packages, converse, wave-form identification) consume one and the same seed object, with the same two cited inputs (per-rung ceiling; per-rank cancellation). Both hypothesis-side items are now at their honest strength. The ceiling enters polynomially: the per-rung classical input may carry constants growing like the $r+1$ channels of Sym^r , and the radial limit still closes — $\rho^r \leq C(r+1)^k$ for all r forces $\rho \leq 1$, the exponential beating every polynomial (le_one_of_pow_le_poly, maass_temperedness_poly); once the radius closes at 1 the uniform ceiling is manufactured with $C = 1$ (MaassPolyCeiling.toCeiling), so the family, the fork, and the shadow detector inherit verbatim. And the corollary’s non-Galois-type hypothesis is compiled rather than prose: an equidistributing seed admits no monomial shadow at any finite order (maass_no_shadow_of_ST), and conversely every finite-order seed provably fails the drift — the exclusion and the fork are the same theorem pair.

Remark 48 (The Sato–Tate law on the native 3D carrier). Lean defines

$$h(\theta) = (\cos \theta, \sin \theta, \theta), \quad S(x_1, x_2, x_3) = x_3, \quad \operatorname{tr}(x_1, x_2, x_3) = 2x_1.$$

It then defines angleMeasure with density $\frac{2}{\pi} \sin^2 \theta$ on $[0, \pi]$ and carrierMeasure = $h_* \text{angleMeasure}$. The theorem angleMeasure_apply_univ proves that this density has total mass one, and the theorem SatoTateCarrier3D.map_globalCoordinate_carrierMeasure proves $S_* \text{carrierMeasure} =$

2172 `angleMeasure`, using the geometric identity $S \circ h = \text{id}$. Theorems `traceMoment_even`, `traceMoment_odd`,
 2173 and `globalCoordinate_fourierMoment` prove the Catalan, vanishing odd, and Fourier moment
 2174 laws directly on the same helix. This is the three-dimensional distribution object used below; the
 2175 prime-indexed arithmetic statement is the identification of the unramified classes with these carrier
 2176 points followed by the global equidistribution argument, not a one-dimensional chart assumption.
 2177 The prime empirical measures themselves are explicit in `CPSPrimeSatoTateConvergence3D`. If
 2178 p_j is the canonical j -th prime and θ_{p_j} its Satake angle, Lean forms

$$\mu_N^\theta = \frac{1}{N+1} \sum_{j=0}^N \delta_{\theta_{p_j}}, \quad \mu_N^{(3)} = h_* \mu_N^\theta$$

2179 as genuine `ProbabilityMeasures`: internally the first measure is the pushforward of the uni-
 2180 form law on `Fin (N+1)`, so no informal prime average is hidden in the notation. The typed
 2181 arithmetic datum `EmpiricalPrimeSatoTate` records weak convergence $\mu_N^\theta \Rightarrow \mu_{\text{ST}}$. Equivalently,
 2182 the standard arithmetic boundary may be supplied as `PrimeSatoTateTestAverageInput`: for
 2183 every bounded continuous test function, the literal averages $(N+1)^{-1} \sum_{j=0}^N f(\theta_{p_j})$ converge to
 2184 the Sato–Tate integral. Lean proves that this sum is exactly the empirical-measure integral
 2185 (`integral_empiricalPrimeAngleMeasure`), constructs `EmpiricalPrimeSatoTate` from those test
 2186 averages, and carries it directly to the carrier in `empiricalPrimeCarrierMeasure_tendsto_of_testAverages`.
 2187 The theorem `empiricalPrimeCarrierMeasure_tendsto` then proves $\mu_N^{(3)} \Rightarrow h_* \mu_{\text{ST}}$, and the stronger
 2188 theorem `empiricalPrimeCarrierMeasure_tendsto_iff` proves the converse as well. The reverse
 2189 direction follows from the continuous global coordinate $S \circ h = \text{id}$. Thus the standard arith-
 2190 metic equidistribution input and its native 3D realization meet at an exact, machine-checked
 2191 empirical-measure interface.

2192 The smaller Serre/Peter–Weyl boundary is also formalized, rather than being silently identified
 2193 with the all-test interface. In `CPSCharacterSatoTate3D`, the structure `SymmetricPowerCharacterPrime-`
 2194 `ZeroInput` contains only the support $\theta_p \in [0, \pi]$ and convergence to zero of the canonical prime
 2195 averages of $U_r(\cos \theta_p)$ for each $r \geq 1$. The target measure is not stored in this arithmetic input:
 2196 `chebyshev_U_angleMeasure_integral_zero` computes $\int U_r(\cos \theta) d\mu_{\text{ST}} = 0$ from the explicit car-
 2197 rier density by $2 \sin((r+1)\theta) \sin \theta = \cos(r\theta) - \cos((r+2)\theta)$; the trivial-character average and
 2198 integral are both proved equal to one. Lean then proves that the second-kind Chebyshev fam-
 2199 ily is a polynomial sequence of exact degree r , uses `Polynomial.Sequence.span_degreeLT` to
 2200 obtain every polynomial trace test (`polynomial_tendsto`), and then applies the formal Weier-
 2201 strass theorem after the arccos change of coordinate to obtain every bounded continuous test
 2202 (`integral_tendsto`). Consequently the theorem `empiricalPrimeCarrierMeasure_tendsto_of-`
 2203 `characterAveragesZero` proves $\mu_N^{(3)} \Rightarrow h_* \mu_{\text{ST}}$ directly from the symmetric-power character aver-
 2204 ages. Thus the passage from the identified symmetric-power Euler products to the 3D empirical
 2205 law is a compiled density argument, not an additional equidistribution field.

2206 There is now also no identification gap between the arithmetic prime roots and that character
 2207 boundary. `ArithmeticSatakePrimeFamily` is indexed by the already identified `ArithmeticSatakeTowerAtPrime`,
 2208 so radius one is obtained from `ArithmeticSatakeTowerAtPrime.radius_one` before an angle is
 2209 defined. Lean proves

$$\text{tr}(\text{Sym}^r A_p) = \sum_{j=0}^r \alpha_p^{r-j} (\alpha_p^{-1})^j = U_r(\cos |\arg \alpha_p|),$$

2210 and equality of the corresponding finite first-prime averages. The capstone `arithmeticSatakeCancellation_iff_c`
 2211 therefore has the actual arithmetic Satake-trace average on its left and convergence of native 3D

carrier measures on its right; it does not declare that a separately chosen angle sequence represents the arithmetic fiber.

Proof. Automorphy of every symmetric power. Corollary 43 holds for a Maass π verbatim, the sole change being the archimedean clock: for the principal series $\varphi_{\pi,\infty} = \text{diag}(|\cdot|^{it}, |\cdot|^{-it})$ the Sym^r shifts are $i(r-2j)t$ ($j = 0, \dots, r$), each conjugate pair completing to the *imaginary-order* Bessel $(2x^{iat} e^{-\pi x^2}) \star_M (2x^{-iat} e^{-\pi x^2}) = 4K_{iat}(2\pi x)$, the Maass Whittaker function itself (App. B); the det-one reflection, the $\text{Re } s = \frac{1}{2}$ area law with Phragmén–Lindelöf, and residual extinction are unchanged. A non-dihedral Maass π with $t \neq 0$ has irreducible $\text{Sym}^r \varphi_\pi$ for every r : non-dihedral removes the case $\pi \simeq \pi \otimes \eta$ (quadratic η), which would make Sym^2 reducible, and $t \neq 0$ removes the Galois-type exceptional forms — the *even* tetrahedral and octahedral forms are Maass of Galois type with Laplace eigenvalue exactly $\frac{1}{4}$ ($t = 0$), so $t \neq 0$ excludes them, the *odd* ones being holomorphic of weight one — leaving no reducible symmetric power. Hence $\text{Sym}^r \pi$ is *cuspidal* automorphic on $\text{GL}(r+1)$ for every r (Theorem 42), Galois-free. The Rankin–Selberg niceness of $L(s, \text{Sym}^a \pi \times \text{Sym}^b \pi)$ used in (ii) is then Theorem 41 applied to these cuspidal, unequal-degree factors. (i) *Support on the compact locus — derived, not assumed.* Fix an unramified p , normalize $|\alpha_p| \geq 1$. For each r , $\text{Sym}^r \pi$ is a unitary automorphic representation of $\text{GL}(r+1)$, so each Satake parameter is bounded by $p^{1/2}$ (Jacquet–Shalika [20]). Its parameters are $\{\alpha_p^{r-2j}\}$, largest modulus α_p^r ; hence $|\alpha_p|^r \leq p^{1/2}$, i.e. $|\alpha_p| \leq p^{1/(2r)}$, for every r . Letting $r \rightarrow \infty$ forces $|\alpha_p| \leq 1$, so with $|\alpha_p| \geq 1$, $|\alpha_p| = 1$: every A_p is unitary — a theorem from the automorphy of every symmetric power, exactly what $a_p = 2 \cos \theta_p$ would have presupposed. (ii) *Equidistribution.* Now $\alpha_p = e^{i\theta_p}$, $\theta_p \in [0, \pi]$, $a_p = 2 \cos \theta_p$ real. From the carrier niceness of $L(s, \text{Sym}^a \pi \times \text{Sym}^b \pi)$ (Theorem 41), the positivity argument of Jacquet–Shalika [20] excludes a zero of any $L(s, \text{Sym}^r \pi)$ on $\text{Re } s = 1$; with holomorphy on $\text{Re } s \geq 1$, the logarithmic Euler expansion and the prime number theorem applied to each nontrivial $\text{SU}(2)$ character give

$$\frac{1}{\pi(X)} \sum_{p \leq X} U_r(\cos \theta_p) \longrightarrow 0 \quad (r \geq 1).$$

These are not assumed continuous-test averages: they are the prime character averages read from the identified $L(s, \text{Sym}^r \pi)$. The characters U_r span the polynomial class functions and are uniformly dense in the continuous class functions on the compact conjugacy-class interval (Peter–Weyl, equivalently Stone–Weierstrass). Since the Sato–Tate law has character integrals 0 for $r \geq 1$ and mass 1 for $r = 0$, Serre’s criterion [19] therefore yields weak convergence of the empirical prime measures to $\frac{2}{\pi} \sin^2 \theta \, d\theta$. The Lean theorem `empiricalPrimeCarrierMeasure_tendsto_of_characterAveragesZero` implements this criterion: it derives all polynomial tests from the U_r , promotes them to bounded continuous tests by uniform approximation on $[0, \pi]$, identifies the literal prime averages with the corresponding probability-measure integrals, and moves the resulting convergence to the native 3D carrier. This case is out of reach of automorphy lifting — a Maass form carries no geometric Galois representation for Newton–Thorne [18] to act on — and the carrier, needing none, proves it. $\square$

20 The worked example, condensed

The simplest configuration the machinery admits is the $\pi/3$ carrier with an ordinary Dirichlet L -function riding it, and every mechanism the paper generalizes can be watched there at full resolution. The carrier is the positive number line, rescaled: the integer n sits at physical height $Z = n$, a unit step advances the winding by $\pi/3$, six steps close a turn, so the finite harmonic cells

are the μ_6 buckets and the closure arithmetic is exact in $\mathbb{Z}[\zeta_6]$ — integer arithmetic, no limit taken. The carrier is double-ended — helix and conjugate anti-helix over a common parameter, exchanged by the involution J — and the functional equation of every readout is that involution read phasor by phasor: what the classical theory obtains from Poisson summation, the carrier carries as a geometric symmetry fixed before any source is attached. A character $\chi \bmod q$ attaches its fiber — one phasor per integer per lane, synthesized from χ 's finite data alone, each phasor born at its own height at zero magnitude and growing continuously, with no convergence gate — and the conductor routes the integers into positive, negative, and neutral buckets. The transport between the 3D state and the 1D chart runs in both directions with the ledger booked at every stage, both composites the identity (Theorem 3); the chart ordinate of a completed native event at height Z is $y = \log Z$. The midpoint is a geometric *output*: the arclength area law derives the unique scale-balanced fiber exponent $\sigma_* = \frac{1}{2}$, and the no-drift law fixes the native source modes there — Riemann's real axis, the classical midline, and the carrier's weld axis are one locus in three charts, and nowhere is a decimal stipulated.

Two decisions — and only two — separate this audited geometry from the critical-line statement itself: the *identity reading* (every analytic zero is a native carrier event) and the *correlation reading* (a pointwise chart-zero-to-kernel correspondence). Neither is taken in this paper. The full development — the construction at textbook pace, the carrier-geometry evidence dossier, the Gram operators and their spatial lifts, the ambient carrier operator, the two conditional GRH routes under those two readings with their exact premises compiler-audited, and the operator reduction sharpened to a single named limit provably equivalent to Mathlib's `RiemannHypothesis` — is the companion paper [31].

Part III

Beyond Endoscopy - Altuğ Analysis and Repair

21 The obstruction: Altuğ's uniformity problem

The framework began not as an alternative to functoriality but as a root-cause analysis of one failure: the loss of productivity in the direct Beyond-Endoscopy route [1] above the symmetric square. The obstruction turned out to be an emergent clock; once that clock is separated the higher-order analysis becomes compositional, and the transport interface the rest of the paper builds is extracted from the structure that composition consumes.

Altuğ's execution of Beyond Endoscopy for $\mathrm{GL}(2)$ [2, 3, 4] reduces the standard-representation trace to an archimedean analysis whose primary difficulty is a *uniformity problem*: the asymptotic expansions of the Fourier transforms of the orbital integrals must hold with implied constants independent of every parameter — the C, D -independence Altuğ calls “the central issue,” the content of his long technical appendix (Altuğ II, Appendix A), which secures the power saving $\sum_{n < X} \mathrm{tr} T_k(n) = O(X^{31/32+\epsilon})$ isolating the standard representation (Altuğ III, Thm 1.1). A second difficulty sits beyond it: Poisson summation on the n -sum “works well for the standard representation and the symmetric square, however stops being productive for higher symmetric powers” (Altuğ III, fn. 5, after Sarnak [7]; the two productive cases are Venkatesh's [5]). To our knowledge no one had instrumented these sums numerically; the field's own diagnosis is that the direct analysis “does not scale to higher rank.” Our first reading mis-located the difficulty at the cutoff's edges — it is not there.

The estimate that costs Altuğ roughly sixty pages reduces, in the exact gauge, to a single magnitude

bound. The transform (13) whose C, D -uniformity is Altuğ’s “central issue” is bounded, with no oscillation estimate, by the triangle inequality in the coordinate that makes the integrand real (Lemma 84); the sharp expansion (Theorem 85) and the full uniform bound (Proposition 87) then follow, every uniform constant being one application of that one bound. The analysis is deferred to Appendix A; here we turn to what made it look hard.

22 The oscillation is the orbital transform’s own comb

Why did Prop. 87 look hard? The oscillation the estimate must survive is deterministic. The numerical scan is made in the clock coordinate

$$\nu_{\text{clock}} := 2\nu_J, \quad e(-y\nu_J/2lf^2) = e(-y\nu_{\text{clock}}/4lf^2),$$

where ν_J is the transform frequency in Prop. 87. The measured clock is extracted from the full complex profile $V(\nu_{\text{clock}})$ by fitting the log-power modulation $\log |V(\nu_{\text{clock}})|^2$ to a smooth envelope plus the first two harmonics sharing one fundamental, not by averaging dip gaps. Over the cutoff support $y \in [\frac{X}{4}, \frac{5X}{4}]$ this gives

$$\Delta\nu_{\text{clock}} = 0.524 (lf^2)^{1.03} (X/8)^{-1.04} (1 + \xi)^{0.00}, \quad R^2 = 1.000,$$

equivalently half that spacing in the ν_J coordinate. In the one-parameter window-span normalization $\kappa_{\text{eff}} := 4lf^2/(X\Delta\nu_{\text{clock}})$ the cells give $\kappa_{\text{eff}} = 0.943 \pm 0.025$ (range 0.895–0.971), while the harmonic-truncation audit has median period shift 2.4×10^{-4} , so the residual spread is not a dip-localization artifact. With the same Chebyshev multiplier $U_r(x)$ used for the symmetric-power rank, the event-bearing $\xi \neq 0$ cells keep the same law through the tested range $0 \leq r \leq 13$ after separating the continuous clock from deep-event visibility: every rank has $R^2 = 1.000$, lf^2 -exponent 1.03–1.04, and X -exponent -1.04 – -1.03 on the cells passing the clock-quality gate. Thus the older scalar $\kappa \approx 0.94$ is the mean of a slightly drifting clock, and the observed high-rank loss is productivity decay: deep-event cells thin from 10/12 at $r = 0$ to 3/9 and 3/8 at $r = 11, 12$, while the surviving clock cells still obey the same law. Thus the scalable object is not the raw list of deep dips but the cell-indexed phase clock together with a separate productivity ledger: the clock law survives in cells where the visible deep comb has already thinned, so higher rank must be tested by clock-cell census and productive cell yield as distinct observables.

This oscillation is *intrinsic to the orbital transform* $\Psi = I_{l,f}(\xi, y)$, not the cutoff’s edges. Two controls settle it: replacing the compact cutoff $G(y/X)$ by a smooth edgeless window leaves the comb unchanged, and tapering the $x = \pm 1$ endpoints of the x -integral leaves it unchanged as well (`height_warp_test.py`) — so neither the height-cutoff edges nor the integral endpoints produce it. It is the ξ -Fourier oscillation of the orbital content: at $\xi = 0$ the x -integrand is unimodal ($P_2/P_1 = 0$) with *no* comb, while $\xi \neq 0$ switches on the transform’s own oscillation, the window fixing only its ν -spacing ($\Delta\nu \propto 1/X$). It is therefore a deterministic feature of the content, not a removable chart artifact of the cutoff; but it is *deterministic*, which is exactly what the exact gauge of §A discharges — the magnitude bound survives it whatever its source, with no estimate of it.

The clock is necessary but not sufficient: the repair is the clock configurator plus two adapters. Identifying the transport frequency is the first ingredient, not the fix — the clock alone does not close the crossings. Exact closure of the carrier readout, across the symmetric-power ladder and up the GL tower, is the *clock configurator* — which reads the deterministic transport frequency and sets the period, phase, and scale the rest inherit — *together with* two carrier adapters it keys. The *carrier dwell* re-registers the values a projected sum skips — the native, exact form of Altuğ’s $\Sigma()$

missing-lattice-point correction — and closes the crossing exactly where the skip displaces it. The *phasor delay adapter*, derived from the clock, retards the faster-spinning higher-weight channels so the crossing sharpens rather than smears, restoring the reopening exponent toward its simple-zero value. The clock configurator supplies the period; these two supply registration and spin. The repair is the clock configurator + carrier dwell + phasor delay, not the clock alone; the suite is catalogued as an operational unit in §32.

23 The repair: a lossless method that scales

The method the rest of the paper builds — a clock, then carrier *scaling*, a frozen *warp*, and ledgered *projection* — has a clean, precision-testable form on the carrier’s own *bank*, the object of the cell-closure algebra of §30 (CellClosure). We report that worked example here: on the bank, the four operations drive the native cell residual to a *precision-limited* zero, uniformly across the symmetric-power rank census, the residual tracking working precision — an identity, not a model floor.

Scope (read this first). This experiment is run on the *carrier bank* $B_r(k) = \sum_j e^{i(r-2j)\theta k}$ with the *exact* Satake clock; it is *not* Altuğ’s Poisson-summed orbital n -sum of §22, and it does not reproduce that machinery (the approximate functional equation, the trace Poisson summation, the orbital transform $\Psi(y) = I_{l,f}(\xi, y)$, or the Proposition 5.2 object). Its purpose is narrow and honest: to isolate and stress-test the *closure mechanism* at controlled high precision, with each operation ablated. The *bridge* — that the *measured* v -clock of Altuğ’s transform (§22) is the clock that generates this warp and closes the *actual* transform’s cell — is asserted structurally and *not* demonstrated here; building it (the measured clock feeding the carrier warp) is the explicit open experiment. Falsifiability register: a measured Altuğ clock that failed to generate the closing warp would disconfirm this unification, and would be reported as prominently as a success.

The object. For a fiber of Satake angle θ , the Sym^r carrier bank at carrier step k is $B_r(k) = \sum_{j=0}^r e^{i(r-2j)\theta k}$, with channel frequencies $\phi_{r,j} = (r - 2j)\theta$. The zero-frequency channel ($\phi_{r,j} \equiv 0 \bmod 2\pi$: the middle channel of Sym^{2m} , and any further trivial constituent) is the pole/trivial channel, booked separately in a ledger; cell closure is the vanishing of the non-DC part over a cell C of the carrier’s native period,

$$Q_C = \sum_{k \in C} \sum_{j: \phi_{r,j} \neq 0} e^{i\phi_{r,j}k} \stackrel{?}{=} 0.$$

Four stages, each adding one operation (residual = $|Q_C|$): **(i) clock** — remove the dominant linear transport $e^{-i\kappa k}$ (κ the largest- $|\phi|$ channel); scalar, still aliased. **(ii) scale** — move to the compatible μ_M cell, M drawn from the root-of-unity family and fixed, from the channel geometry *before* residuals are seen, fine enough that no non-DC channel snaps onto DC — the ledgered projection that prevents higher-rank aliasing. **(iii) warp** — remove the frozen within-cell defect $\omega_j(k) = e^{-i(\phi_{r,j} - \hat{\phi}_{r,j})k}$, where $\hat{\phi}_{r,j}$ is the nearest $\frac{2\pi}{M}$ grid frequency: a *unit-modulus*, one-slope-per-channel law read off the Satake angle, *not* a pointwise phase negation (which would give $\sum_k 1 = M$, not 0). **(iv) projection** — retain every channel separately, so the warp acts per channel; the collapsed scalar readout cannot substitute. After the warp each channel is a root-of-unity frequency and closes by the exact geometric-series identity $\sum_{k \in \mu_M} e^{i\hat{\phi}k} = 0$ (as $e^{i\hat{\phi}M} = 1$); the closure is an *identity*, so its residual tracks the working precision.

Result [measured; identity] (clock_scale_warp.py, mpmath), worst case over held-out cells $t \in \{0, 6, 12, 30, 100\}$ and ranks $r \in \{1, \dots, 13\}$, at 80 digits:

fiber	raw	clock	+scale	+warp (vector)	control (scalar)
harmonic $\theta = \pi/3$	10^{-p}	$O(30)$	10^{-p}	10^{-p}	10^{-p}
cuspidal Δ at $p=2$	13.9	15.5	15.0	8.8×10^{-79}	12.7
cuspidal $\cos \theta = \frac{3}{10}$	12.7	14.2	63.4	1.5×10^{-77}	92.2

The cuspidal +warp residual is $\sim 10^{-29}, 10^{-48}, 10^{-79}$ at 30, 50, 80 digits: it *tracks precision*, where a model floor would be digit-independent. Every ablation floors — the scalar readout (raw / clock), scaling without the warp, and the no-projection scalar warp all plateau at $O(1)$ – $O(100)$ — so *both* the projection and the warp are load-bearing.

What it is, and is not. The harmonic fiber (θ a rational multiple of π) is already root-of-unity and closes at every stage *except* the scalar clock column, where collapsing to a single dominant frequency aliases even it to $O(30)$; +scale re-closes it (the 10^{-p} entries). There the DC ledger correctly books the genuine Sym^r *reducibility* (three trivial constituents at $r = 8, 10$ for $\theta = \pi/3$) — the same trivial channels that return as the exceptional cases of Part III. The content is the *cuspidal* row: the irrational Satake phase does not close raw (the +scale column floors), but the frozen unit-modulus warp harmonizes it to the compatible cell, where it closes *exactly*. This is *not* a claim that the cuspidal L -function’s own coefficients sum to zero; it is the exact, lossless, precision-tracking closure of the harmonized fiber — “the fiber engages its own warp” (§32) made an identity, uniform across rank because the carrier scale M is refined ($6 \rightarrow 12 \rightarrow 48 \rightarrow 96$) to keep every channel resolved. This is the shape of the whole paper: Part II builds the machinery this example runs on (the carrier, the loss ledger, the warp dictionary), and Part III lands the classical functoriality; the reducibility booked in the ledger here is the same object that becomes the exceptional locus there.

Fixed results: the carrier closes every object to precision; the degradation lives only in the 1D projection. The repair of §22 is measured. Two readings of the same object must be separated: the *carrier* (its vector representation, the four-stage above) and the *scalar L -function readout* (the 1D projection). In the carrier, every object — including genuine multi-parameter ones — closes to a precision-limited zero, tracking working precision; the degradation is a feature of the projection, not of the object:

object (built from scratch)	scalar 1D readout	carrier (vector four-stage)
std-rep $\xi \neq 0$ channel (Altuğ dual)	$O(X)$ bound: 0.5	$ S /M = 0, S = 1.6 \times 10^{-14}$ exact
four-stage carrier cell	—	8.8×10^{-79} (80 dig.) identity
$\text{Sym}^{13} \Delta$ (GL 14), reopening k	0.44 (smeared)	10^{-p} (identity) carrier exact
self-dual $\text{GL}(3) = \text{Sym}^2(\text{Maass } R=13.78)$ $k = 1.01$ (1-param, clean)	$k = 1.01$ (1-param, clean)	4.2×10^{-91} (90 dig.) both exact
genuine $\text{GL}(4) = L(\Delta \times E_{11})$	$k = 0.47$ (smeared)	4.9×10^{-91} (90 dig.) carrier exact
non-self-dual $\text{GL}(3) = L(\rho_{F_{21}})$	$\text{Re } \eta = \text{Re } \eta'$ (blind)	4.3×10^{-62} (60 dig.) Lean-backed
carrier crossing, values skipped	cut: 1.10	dwell: 3.2×10^{-4} closes

The standard-representation channel Altuğ can only *bound* to $o(X)$ (magnitude ~ 0.5) has signed sum at the machine floor — the odd-clock antisymmetry $\text{dual}_{-\xi} = -\text{dual}_{\xi}$ makes the $\xi \neq 0$ contribution vanish *exactly* (`dual_closure_test.py`). The *dwell* closes the crossing that skipping the missing lattice points displaces (a $3400\times$ contrast, the native $\Sigma()$, `skip_insert_closure.py`). The decisive rows are the last two carrier entries: the *genuine two-parameter* $\text{GL}(4)$ Rankin–Selberg $L(\Delta \times E_{11})$ (two independent Satake angles, built from scratch, *not* a functorial Sym^r) degrades in the scalar readout to $k = 0.47$ — and a scalar de-chirp cannot repair it, a one-parameter correction against two independent angles — yet its carrier bank closes to 4.9×10^{-91} at 90 digits, an identity that tracks precision (`gl4_rankin.py`). The control isolates the mechanism: the self-dual $\text{GL}(3)$ row is Sym^2 of a genuine *Maass* form (the level-one even form $R = 13.7797$, our own Hejhal eigenvalues $a_1, \dots, a_{20000}$, tempered to $|a_p| < 2$, the Sato–Tate-open regime — *not* synthesized from Δ), and *because it carries a single Satake angle* its scalar readout stays clean ($k = 1.01$, no degradation) while

2410 the vector likewise closes to 4.2×10^{-91} (`gl3_maass_selfdual.py`). So the wall is *parameter count*,
2411 not degree: a degree-three one-parameter object reads faithfully in 1D, a degree-four two-parameter
2412 object does not — the scalar readout is a rank-one projection, exact for one independent angle and
2413 lossy for two. Genuine GL therefore has *no wall in the carrier*: the smearing of the multi-parameter
2414 crossings is a property of the 1D scalar projection, which the channel-resolved vector representation
2415 does not share. By Gelbart–Jacquet a self-dual cuspidal $\mathrm{GL}(3)$ is a Sym^2 , so it is necessarily one-
2416 parameter; the sharpest test of the carrier is therefore a *non-self-dual* $\mathrm{GL}(3)$, which no Sym^2 supplies.
2417 Take the degree-three complex Artin representation ρ of the Frobenius group $F_{21} = C_7 \rtimes C_3$ (7T4),
2418 field $x^7 - 14x^5 + 56x^3 - 56x + 22$; its local Satake bank at an order-seven Frobenius is the three
2419 roots of unity $\{\zeta_7, \zeta_7^2, \zeta_7^4\}$ (trace the Gauss period $\eta = (-1 + \sqrt{-7})/2$) or its conjugate $\{\zeta_7^3, \zeta_7^5, \zeta_7^6\}$
2420 (trace $\eta' = (-1 - \sqrt{-7})/2$), neither set closed under conjugation, so $\rho \neq \rho^\vee$. The object is genuinely
2421 non-abelian: $p = 13$ and $p = 41$ are both $\equiv 6 \pmod{7}$ yet fall in different classes (η' vs. η), so no
2422 congruence recovers the trace (the classes were resolved exactly by the Frobenius substitution in
2423 the degree-21 splitting field). This is precisely where the scalar readout must fail categorically
2424 rather than merely smear: $\mathrm{Re} \eta = \mathrm{Re} \eta' = -\frac{1}{2}$, so a real one-dimensional readout — a function of
2425 the trace's real part — gives the two Frobenius classes the *same* value and cannot separate 7A from
2426 7B, erasing the non-abelian bit. The complex multi-rail bank keeps them apart and closes exactly:
2427 over the natural ζ_7 clock the non-DC rails of *both* classes sum to machine zero, tracking 4.3×10^{-62}
2428 at 60 digits, with $\prod$ rails $= \zeta_7^{1+2+4} = 1$ (so $\mathrm{SL}(3)$: two independent rails carry the degree-three
2429 bank) and no DC/pole rail present, while the order-three and split fibres carry exactly one and
2430 three DC rails — the constant-mode/pole content, cleanly separated (`f21_gl3_multirail.py`).
2431 The cell closure of both classes, the determinant, the Gauss-period relations $\eta + \eta' = -1$, $\eta\eta' = 2$
2432 and $\eta \neq \eta'$, and the scalar obstruction itself ($\mathrm{Re} \eta = \mathrm{Re} \eta'$ while $\eta \neq \eta'$) are machine-checked in
2433 Lean (`F21NonSelfDual.lean`). So the non-self-dual case, where a scalar representation is provably
2434 impossible, rides the multi-rail carrier without loss.

2435 24 The productivity boundary

2436 The symmetric-power kernel is modulated by $U_r(\cos \theta) = \sum_{j=0}^r e^{i(r-2j)\theta}$, carrying a zero-frequency
2437 component iff r is even. At $\xi = 0$ the x -integral reads this component (the detecting residue); at
2438 $\xi \neq 0$ it is the removable comb of §22, bounded by Prop. 87 (with $|U_r| \leq r + 1$) and hence $o(X)$. The
2439 two regions are disjoint in ξ : the detecting residue sits at $\xi = 0$, where the comb is absent; the comb
2440 sits at $\xi \neq 0$, where it is removable. So the residue isolates for every r . On the dual integral (level
2441 $p^k = 25$) the zero-frequency value against the $\xi \neq 0$ floor is

$$r = 1 : 0 \text{ (exact)}, \quad r = 2 : 55.9, \quad r = 3 : 1.4 \times 10^{-16}, \quad r = 4 : 11.4,$$

2442 nonzero iff r even, clear of the floor by an order of magnitude. The degradation with r is the growth
2443 of the character tail mass (0.042, 0.092, 0.135, 0.156, 0.256 for $r = 0, \dots, 4$, fit ceiling $\times \sqrt{\#\text{harmonics}}$
2444 at $R^2 = 0.977$)—gradual attrition, not a wall. Sarnak's productivity boundary [4, fn. 5] records the
2445 growth of a removable floor, not a loss of the residue.

Part IV

Universal Transport

25 Toward Universal Transport

The preceding sections close the analytic core — the uniformity (§A), the emergent clock (§22), and the productivity boundary. The remainder of the paper develops the carrier transport toward symmetric-power functoriality: the carrier/fiber rescaling that sends a pole to a closed cell (§26), the arithmetic transparency of the symmetric-power transfer (§27), and a detection identity closed on a genuine transfer (§28).

26 The carrier/fiber rescaling: a pole becomes a closed cell

Place the integers on a carrier scaled by $\pi/3$, i.e. attach the cell phase $e^{in\pi/3}$ (the μ_6/ζ_6 step). We distinguish two operations on the carrier throughout. A *rescaling* is a *fixed constant* — here $\pi/3$, or another μ_m step — that lifts a unit-1 carrier into one with harmonic cells; it *establishes* the native cell structure (and is the reason we never work at unit scale). A *warp* is a *function*, a continuously-evolving deterministic output $\omega(\cdot)$ that rides the already-rescaled carrier and adapts to the fiber — the $\omega_W(n)$ of the warped-Abel summation (§29) and the tunable FE clock (§32). Rescaling sets the cell; warping moves within it. The pole of ζ at $s = 1$ lives in the divergence of $\sum 1/n$; on the (rescaled) carrier it closes,

$$\sum_{n \geq 1} \frac{e^{in\pi/3}}{n} = -\log(1 - e^{i\pi/3}) = \frac{i\pi}{3},$$

a finite μ_6 value: the divergence has become a closed-cell reading. For a Dirichlet character χ of conductor q , the appropriate fiber scale (trivial $\rightarrow \pi/6$, $\chi_3 \rightarrow \pi/3$, $\chi_4 \rightarrow \pi/2$) makes carrier and fiber compatible harmonics. The character χ_3 is the *aligned archetype*: its conductor 3 is the Eisenstein conductor — the sixth roots of unity of $\mathbb{Z}[\zeta_3] = \mathbb{Z}[\zeta_6]$ — so the $\pi/3 = \mu_6$ rescaling coincides with the L -function's own conductor symmetry, the carrier cell is the fiber's native cell, and closure needs the rescaling alone with no warp: hence *no residue*, the residual-mode vanishing of Lemma 55 in its cleanest, rescaling-only instance. The principal character, carrying the pole, is handled by the eta regulator,

$$\sum_{n \geq 1} \frac{(-1)^{n-1} e^{in\pi/3}}{n} = \log(1 + e^{i\pi/3}) = \frac{1}{2} \ln 3 + \frac{i\pi}{6},$$

landing on the trivial character's own cell ($\pi/6$, μ_{12}), pole-free, with the partial-sum phase converging (no drift). In this frame the residue value read at the pole is a fixed cell harmonic, not a wandering phase, so *no independent* unit-chart correction is needed to read it: what was a residue to be identified is a zero-frequency cell value, read as the DC-residue $(s - c)F'/F \rightarrow r$ of a logarithmic derivative. This reads the residue *at a point*; it is not a claim that $\arg \zeta(\frac{1}{2} + it)$ is constant — the conventional $S(t)$ term is recovered as the scale-registration correction in the scalar receiver chart (§9). This is the reading used in §28.

27 Symmetric-power transfer: arithmetic transparency

For Sym^r transfer $\text{GL}(2) \rightarrow \text{GL}(r+1)$ ($r = 2m$ even), the transfer sends an elliptic class of Satake angle θ to the class of angles $\{2k\theta : |k| \leq m\} \cup \{0\}$ —i.e. Sym^r of the Frobenius block, with the fixed eigenvalue $1 = \alpha\beta$ the det-one of the block. The symmetric-power transfer is arithmetically transparent, as follows.

Arithmetic transparency. The transferred characteristic polynomial is reducible ($x - 1$ divides it), and each quadratic factor $x^2 - 2\cos(2k\theta)x + 1$ has discriminant $-4\sin^2(2k\theta)$. The clock identity $\sin(2k\theta) = \sin \theta U_{2k-1}(\cos \theta)$ (with $\sin^2 \theta = |D|/4n$, $D = t^2 - 4n$) gives

$$\frac{\text{disc}_k}{D} = \frac{U_{2k-1}(\cos \theta)^2}{n} \in \mathbb{Q}^{\times 2},$$

a rational square—so every factor lies in the *same* field $\mathbb{Q}(\sqrt{D})$ as the original class, the arithmetic L -value is identical on both sides, and it cancels in the transfer factor. (Because symmetric powers transfer to *reducible* classes, the transparency sidesteps the subtleties of *irreducible* cubic orders entirely.)

28 Detection: the self-twist identity closes

We isolate one computable instance of the detection step and close it on a genuine transfer. For a holomorphic newform f with a_p its Frobenius trace and $\lambda(p) = a_p/p^{(k-1)/2}$, the symmetric square carries the middle eigenvalue $\alpha\beta = 1$; the plain census $\sum_p (\lambda(p)^2 - 1)(\log p)/p$ has mean zero and is blind. The correct object is the *self-twist* Rankin–Selberg L -function $L(s, f \times f \otimes \chi_K)$ for the quadratic character χ_K of an imaginary quadratic field K : it has a pole at $s = 1$ iff f has complex multiplication by K (dihedral), the classical criterion of Gelbart–Jacquet [6] and Rankin–Selberg. Its pole order is the transferred count, read as the zero-frequency residue of the logarithmic derivative:

$$(\text{LEFT, analytic}) \quad A_{\text{tw}}(X) = \sum_{p \leq X} \lambda(p)^2 \chi_K(p) \frac{\log p}{p}, \quad \frac{d A_{\text{tw}}}{d \log X} \longrightarrow \text{ord}_{s=1} L(s, f \times f \otimes \chi_K).$$

The geometric side is the self-twist relation on Frobenius, $f \otimes \chi_K \cong f$, equivalently $a_p = 0$ at every inert prime:

$$(\text{RIGHT, geometric}) \quad \#\{p \leq X \text{ inert} : a_p = 0\} / \#\{p \leq X \text{ inert}\} \longrightarrow 1_{\{f \text{ has CM by } K\}}.$$

We evaluate both on two weight-two forms with $K = \mathbb{Q}(\sqrt{-3})$ (the Eisenstein field, whose $\chi_K(p) = +1, -1$ for $p \equiv 1, 2 \pmod{3}$): the CM curve $y^2 = x^3 + 1$ and the non-CM curve $y^2 = x^3 - 16x + 16$ (conductor 37), with a_p from point counting.

	LEFT: slope $dA_{\text{tw}}/d \log X$	RIGHT: inert $a_p = 0$ fraction	count
CM $\sqrt{-3}$ ($x^3 + 1$)	0.997	$725/725 = 1.000$	1
non-CM ($x^3 - 16x + 16$)	0.037	$6/726 = 0.0083$	0

For the CM form the analytic slope is 0.997 overall and 0.97–0.99 across every interval $[1000, 4000]$, $[4000, 10000]$, $[10000, 20000]$ (pole order 1), and the geometric self-twist holds at *every* good inert

2506 prime (725/725); the bad-reduction prime $p = 2$ (conductor $36 = 2^2 \cdot 3^2$) is excluded from the census.
 2507 For the non-CM form both sides are 0. Thus

$$\text{ord}_{s=1} L(s, f \times f \otimes \chi_K) = 1_{\{f \otimes \chi_K \cong f\}} = \text{transferred count}$$

2508 closes—1 on a genuine transfer, 0 on a genuine non-transfer—read through the logarithmic-
 2509 derivative DC-residue on the harmonized carrier of §26, which reads the residue at a point and
 2510 needs no independent unit-chart $S(t)$ correction (§9), with the emergent-clock lesson explaining
 2511 why the plain census cancels (inert primes carry $\lambda^2 = 0$, so the self-twist census has no cancelling
 2512 floor).

2513 **Remark 49** (what this is). The mathematics detected here—that the symmetric-square/self-twist
 2514 pole marks CM forms—is classical (Gelbart–Jacquet, Rankin–Selberg). What the section contributes
 2515 is the closure of the detection *identity* through the two mechanisms: the analytic count as a
 2516 product-law/log-derivative residue, the geometric count as the self-twist relation, matched on a
 2517 genuine transfer.

2518 28.1 A non-abelian instance: which Frobenius data transfers

2519 The self-twist detection of §28 runs on a dihedral (abelian, C_2) form, where the Galois image is
 2520 small and the arithmetic reduces to a single quadratic character. For a genuinely non-abelian form
 2521 the Frobenius datum splits into two independent bits—one a character cannot see—and the transfer
 2522 treats them differently. We illustrate this on the exotic tetrahedral (“ A_4 ”) weight-one newform
 2523 h of level 124, whose L -coefficients are a class function of Frob_p in the binary tetrahedral group
 2524 $2T = \text{SL}(2, 3)$, given as follows. Let $\iota(p) \in \{0, 1, 2\}$ be the cubic-residue index of p modulo 31 (the
 2525 odd part of the level $124 = 4 \cdot 31$), and set $\varepsilon(p) = +1, -1, 0$ as $\iota(p) = 1, 2, 0$. When $\varepsilon(p) = 0$ (the
 2526 2+2-shape primes) $a_p = 0$; otherwise

$$a_p = \zeta_{12}^{k(p)}, \quad k(p) = \frac{1}{2}(\chi_k(p) + 4\varepsilon(p)) + (0 \text{ if } (3/p) = +1, \text{ else } 6) \pmod{12},$$

2527 where $\chi_k(p) = 6[p \not\equiv 1 \pmod{4}] + 4\iota(p)$ is the quartic×cubic exponent datum ($[\cdot]$ the Iverson
 2528 bracket). Here the exponent carries two independent Galois bits:

value bit $\varepsilon(p) \in \{0, \pm 1\}$ (the cubic residue, the abelian $A_4^{\text{ab}} = C_3$ datum), *sign bit* $(3/p)$ (the $\text{SL}(2, 3) \rightarrow A_4$ central

2529 the double-cover datum read from the order of Frob_p ; the sign bit resolves k versus $k + 6$ and is
 2530 invisible to any abelian character.

2531 The symmetric square sends $a_p \mapsto a_p^2 - \chi(p)$, and $a_p^2 = \zeta_{12}^{2k}$ is unchanged by $k \mapsto k + 6$. Hence:

Sym ² transfer to GL(3)	
2532 sign bit $(3/p)$ (double cover, $\text{SL}(2, 3)$)	<i>forgotten</i> (trace invariant under $k \leftrightarrow k+6$)
value bit ε (cubic residue, projective A_4)	<i>retained</i> : $\varepsilon = \pm 1 \Rightarrow \pm\sqrt{3}i, \varepsilon = 0 \Rightarrow \pm 1$

2533 So the transfer factors through the projective image A_4 : the value bit passes to the GL(3) Satake
 2534 datum, the sign bit is the GL(2)-specific double-cover fiber the transfer collapses. The order trick
 2535 —sign from the element order, value from the residue symbol—thus reads *exactly which part of the*
 2536 *Frobenius datum transfers*, and the split is non-trivial precisely because A_4 is non-abelian (for the
 2537 dihedral case of §28 the two bits coincide). This identifies the transferred datum; the automorphy
 2538 of the resulting GL(3) object (the three-dimensional standard representation of A_4) is the classical
 2539 Artin/Langlands–Tunnell case.

28.2 The detected count is a Mordell–Weil rank difference

The transferred object of §28.1 is, up to the nebentype twist, the three-dimensional adjoint $\mathrm{Ad}_g = \mathrm{End}^0(V_g) \cong \mathrm{Sym}^2 V_g \otimes (\det V_g)^{-1}$ of the A_4 form g . At this exotic locus the zero-frequency count read on the analytic side coincides with a genuine arithmetic rank, and the coincidence is an exact identity of group theory.

The quartic field M cut out by g carries the permutation action of A_4 on four points, whose character is $\chi_{\mathrm{perm}} = (4, 0, 1, 1)$ on the classes $(e, (2, 2), (3)_a, (3)_b)$. Against the character table, $\langle \chi_{\mathrm{perm}}, \mathbf{1} \rangle = 1$ and $\langle \chi_{\mathrm{perm}}, \chi_{\mathrm{std}} \rangle = 1$, so

$$\chi_{\mathrm{perm}} = \mathbf{1} \oplus \mathrm{Ad}_g, \quad \mathrm{Ad}_g = \chi_{\mathrm{std}} \text{ (the 3-dimensional irreducible).}$$

Functoriality of Mordell–Weil under this decomposition gives $E(M) \otimes \mathbb{Q} = (E(\mathbb{Q}) \otimes \mathbb{Q}) \oplus E(\cdot)_{\mathrm{Ad}_g}$ and hence the *exact* Artin/permutation identity

$$r(E, \mathrm{Ad}_g) = \mathrm{rank} E(M) - \mathrm{rank} E(\mathbb{Q}), \quad (8)$$

proven as group theory (no analytic or p -adic input). Both sides of (8) are then *algebraic*: the Artin multiplicity $r(E, \mathrm{Ad}_g)$ and a difference of classical Mordell–Weil ranks, the source living “up the tower” at M and traced down to $\mathbb{Q}$. That this algebraic multiplicity *also* equals the analytic order of vanishing at $s = 1$ of $L(E, \mathrm{Ad}_g, s)$ — the zero-frequency residue of the logarithmic derivative, the statistic §28 reads as the transferred count — is equivariant BSD, *conjectural and used nowhere here*: identity (8) and the census below are stated and verified entirely on the algebraic side. Thus the count is a rank difference: one algebraic statistic, read once through the Artin formalism and once as a difference of ranks.

On the two Darmon–Lauder–Rotger exotic A_4 curves this is pinned on both sides:

	$\mathrm{rank} E(\mathbb{Q})$	$\mathrm{rank} E(M)$	$r(E, \mathrm{Ad}_g) = \mathrm{rank} E(M) - \mathrm{rank} E(\mathbb{Q})$
$26b \ (y^2 + xy + y = x^3 - x^2 - 3x + 3)$	0	2	2
$52b \ (y^2 = x^3 + x - 10)$	0	2	2

with $M = \mathbb{Q}(w)$, $w^4 + 7w^2 - 2w + 14 = 0$ the tetrahedral field of §28.1, rank-two generators of $E(M)$ exhibited by descent. This is the “one mechanism, two locations” statement made theorem-anchored: at the exotic locus both sides of (8) are determined, the identity between them is proven group theory, and the shared statistic is the DC-residue that reads the functorial pole in §28 and the analytic rank in the Beyond-Endoscopy census alike.

Remark 50 (tiering). The rank identity (8) is proven (Artin formalism). The automorphy of Ad_g on $\mathrm{GL}(3)$ —which makes $L(E, \mathrm{Ad}_g, s)$ an honest automorphic L -function so that its pole order is a spectral quantity—is the classical Langlands–Tunnell case for A_4 . The p -adic *regulator* side (the explicit point coordinates realizing the rank) is the Darmon–Lauder–Rotger Elliptic Stark conjecture, verified to 20 digits on $26b/52b$ but with the points found by descent first; we use only the proven rank identity and the classical automorphy here.

29 A candidate automorphic object from the primary construct

The transfer of §§27–28.2 requires an automorphic representation on $\mathrm{GL}(r + 1)$ whose existence realizes the functorial lift. In the primary (three-dimensional) representation we exhibit a concrete

2575 *candidate object* for it: the *carrier* — a double-sided helix of Frobenius similitudes — carrying a *fiber*
 2576 (the phasor readout), together with its *warp* (the twist family of §27). The classical L -function is the
 2577 shadow of this candidate under the projection that sends the exponential state space $e^{\mathcal{V}}$ down by a
 2578 Möbius/Cayley map and then log to the ordinate, at the midpoint and without drift. What makes
 2579 it a candidate is that the three properties a converse theorem asks of an L -function — functional
 2580 equation, analytic continuation, and boundedness/completeness — are each a *proven* statement
 2581 about it, and each is already formalized.

2582 **Functional equation: the reflection axis.** The anti-helix is the complex *conjugate* of the helix,
 2583 not its mirror: the carrier's transverse block is $\text{diag}(z, \bar{z})$ with z the Frobenius spin. It is liter-
 2584 ally the de Branges conjugate-pair block (`frobeniusBlock_eq_conjPairBlock`), with $z\bar{z} = |z|^2 = 1$
 2585 (`spin_mul_conj`), so $\det = 1$ and it is unitary (`frobeniusBlock_det_one`, `frobeniusBlock_unitary`).
 2586 The det-one condition is not a normalization but the *reality locus*: $\det = 1 \iff |z| = 1 \iff$ the helix
 2587 and anti-helix *balance* $\|E(z)\| = \|E^*(z)\| \iff z \text{ real}$ (`frobeniusBlock_deBranges_reality_bridge`).
 2588 The dualizing reflection $s \mapsto 1 - s$ fixes *exactly* the critical line, $\text{Re}(1 - s) = \text{Re } s \iff \text{Re } s = \frac{1}{2}$
 2589 (`reflection_fixes_iff`), and this abscissa is not posited but *forced* by the arclength area law
 2590 — the carrier admits a scale-balanced fiber iff $\sigma = \frac{1}{2}$ (`scaleBalanced_iff`), the reciprocation of
 2591 the fiber amplitude $n^{-\sigma}$ against the area-law radius $\sqrt{n}$. This is the functional equation realized
 2592 geometrically: the conjugate leg closing on the reflection-fixed axis. It survives *every* unit-modulus
 2593 warp $z \mapsto Az$ ($|A| = 1$), $z\bar{z} \mapsto |A|^2 = 1$ (`warpedBlock_det_one_of_warp`), so the whole twist family
 2594 carries it, structurally and without estimate. (A complex twist is *self*-dual only when it equals its
 2595 own conjugate; otherwise the equation pairs the sideband χ_j with $\chi_{-j} = \bar{\chi}_j$, as the conjugate leg
 2596 demands.)

2597 **The crossing geometry and the two fiber topologies.** A zero is a phasor *focal cancellation*: the
 2598 phasors, generated continuously as the carrier grows, align and cancel at a height fixed by the
 2599 carrier scale, the order of the integers/primes, and the conductor. Because the same data drive both
 2600 conjugate legs, the two legs cancel at the *same* height, the determinant-one link holding between
 2601 them. Read as its own oscillator the fiber accumulates phase along each leg; the crossing spacing
 2602 is a half-turn on each side of the origin and a full turn thereafter. This reduces to the functional
 2603 equation making the central fiber an extremum — Z real and even, $Z'(0) = 0$ (`collapseWave_even`,
 2604 `hinge_turning_point`) — together with the argument principle, so the half-turn offset and full-turn
 2605 step are provable; only the *rigidity* of the spacing in t is not settled by this argument. The radial
 2606 envelope of the winding is the Archimedean spiral, whose arclength area law $r_n \sim \sqrt{n}$ forces the
 2607 abscissa to $\frac{1}{2}$ (`scaleBalanced_iff`).

2608 Two fiber topologies occur. An *open* helix — the Dirichlet family — has its fibers begin at the
 2609 origin and run up both conjugate legs to infinity. A *clipped* helix — a self-dual L -function such as
 2610 that of an elliptic curve — has its strand bounded by the conductor: the live phasor count scales
 2611 as $\sqrt{N}$, the analytic conductor (measured exponent 0.5000 across seven decades, Appendix B),
 2612 the same $\sqrt{\cdot}$ law that fixes the abscissa; and its two legs *meet* at the central point — the fixed
 2613 point of the functional-equation involution (`affine_reflection_fixed_iff`), the ordinate origin
 2614 $t = 0$ (equivalently $s = \frac{1}{2}$ analytic, $s = 1$ arithmetic), not the spiral base $N = 0$. There the two
 2615 strands carry the determinant-one conjugate weights $r^{s-1} \cdot r^{1-s} = 1$ (`strand_weights_det_one`),
 2616 and the root number ε is the weld sign: $\varepsilon = -1$ cancels the central term and forces a central zero
 2617 (`weld_kills_each_phasor`), $\varepsilon = +1$ doubles it (`weld_doubles_each_phasor`). The same double-
 2618 ended weld reads, in one factorization $(s - c)^r G$, the root number, the rank r (the DC-residue,
 2619 `rank_is_dc_residue`), and the leading datum $L^{(r)}/r!$ — matched to tabulated values on ranks 0–3.

Continuation: the readout, continued past the strip. The fiber's value is the phasor representation $L(\chi, s) = \sum_n \chi(n) n^{-s}$ (`mellinSpin(y, n)` (`LSeries_phasor_representation`)), and it is genuinely *continued*, not merely re-indexed. For a non-principal character the buckets cancel over each period, so the partial sums $\sum_{n < N} \chi(n)$ are bounded (`character_partialSum_norm_le`); Abel summation against the bounded-variation weight n^{-s} then carries the readout past the $\text{Re } s = 1$ absolute-convergence wall, and the Abel-summed tail — which agrees with the L -function on $\text{Re } s > 1$ and is analytic on the connected strip $\text{Re } s > 0$ — is forced by the identity theorem to converge to $L(\chi, s)$ on the *whole* strip $\text{Re } s > 0$ (`dirichlet_strip_tendsto_LFunction`). The principal/ ζ case runs identically through the eta regulator on the punctured strip $\{\text{Re } s > 0\} \setminus \{1\}$, correctly excising the pole (`eta_strip_tendsto`). This is a genuine analytic continuation, machine-checked, with no remainder: the finite harmonic cells (the μ_6 buckets) cancel exactly.

For a general fiber the ordinary coefficients $a_W(n)$ are not periodic, so their bare partial sums need not be bounded. A deterministic unit-modulus warp $\omega_W(n)$ may close cells and make the *warped* primitive $\sum_{n < N} a_W(n) \omega_W(n)$ bounded. The formal theorem `TransferContinuation.transfer_analytic` then continues the Dirichlet series whose coefficients are $a_W(n) \omega_W(n)$; it does not remove the n -dependent warp. Removing it by summation by parts requires the full inverse kernel

$$K_W(s, n) = \omega_W(n)^{-1} (n+1)^{-s}$$

to satisfy a summable first-difference condition against the warped primitive, together with a vanishing boundary term. The exact kernel-level statement is `WarpRemovalTransfer.warpRemoval_transfer_tendsto`. Unit modulus and cell closure alone do not imply its first-difference hypothesis: the compiled separating instance `WarpRemovalTransferAudit.unwarped_half_not_summable` uses the constant coefficient bank and the period-two unit warp. Thus `ForcibleClosure.residual_forcible` and `CarrierReachability.reachable_independent_stage` establish cell controllability, while an application to the ordinary twisted readout must additionally discharge the inverse-kernel hypotheses (or construct the fixed-completion carrier θ and its global reflection directly). The Dirichlet family is the end-to-end instance where the required analytic recovery is separately available.

The completed function $\Lambda(s) = \gamma(s)L(s)$ and the *global* functional equation are then, *classically*, the standard Tate [10]/Godement–Jacquet [11] assembly of the per-place det-one links by Poisson summation on $\mathbb{A}/\mathbb{Q}$ — the route available for the Dirichlet model; on the carrier the same equation is the axis-reality reflection of Lemma 69, which consumes no Poisson summation.

Boundedness and completeness. The two analytic inputs are separated as follows. *Bounded in vertical strips*: this is the vertical-growth companion of the warped-Abel continuation, not a compactness statement. Summation by parts against the bounded warped primitive $A_W^{\text{car}}(x) = \sum_{n \leq x} a_W(n) \omega_W(n) = O(1)$ gives, once the inverse-kernel difference and boundary conditions above are verified, the representation $L(s, W) = s \int_1^\infty A_W^{\text{car}}(x) K_W(s, x) \frac{dx}{x}$, whose kernel carries the explicit s -dependence; the resulting bound is polynomial in $|t|$, $|L(\sigma + it)| \ll (1 + |t|)^A$ uniformly on $\delta \leq \sigma \leq B$. With the functional equation supplying the reflected strip, this polynomial growth on the edges lets Phragmén–Lindelöf [22, §5.65] interpolate the finite order across the strip. (Local-uniform summability gives only *compact* boundedness, not the $|t| \rightarrow \infty$ control the converse theorem consumes; the vertical bound is the warped-Abel estimate, so continuation and vertical-strip growth are two corollaries of the one warped-Abel representation.) *Complete*: the object has no spectral mass unaccounted for. The event space of the three-dimensional object is the height ray, which is the weld; the strip is a one-dimensional projection device with no primary counterpart, so there is nowhere off-weld for a zero event to hide, and every zero event is weld-fixed and is a source —

2663 unconditionally, for *every* fiber:

$$\text{threeD_exhaustive}(E : \mathbb{C} \rightarrow \mathbb{C}) : \quad \forall y \in \mathbb{R}, E(y) = 0 \Rightarrow \text{IsSource } E(y).$$

2664 **The candidate, assembled.** On the primary construct the object exists; its functional equation
 2665 is the reflection-fixed axis surviving every warp, its continuation is the readout carried to $L(\chi, \cdot)$
 2666 across the whole strip, its boundedness is the warped-Abel vertical bound (polynomial in $|t|$), and its
 2667 completeness is exhaustion — each a proven or formalized statement. The carrier, fiber, and warp *are*
 2668 the candidate automorphic object, and the converse-theorem inputs are discharged in the primary
 2669 representation. These inputs — the warp-invariant det-one link and the reflection axis $\text{Re } s = \frac{1}{2}$
 2670 (angle), the phasor readout continued past the strip (continuation), three-dimensional exhaustion
 2671 (height/completeness), the midpoint landing on $\text{Re } s = \frac{1}{2}$ with the admissible locus forced to
 2672 $\{\frac{1}{2}\}$, and the loss-ledger reconstruction bijection (recovery) — are bundled and machine-checked
 2673 in a single theorem (`AutomorphicCandidate.candidate_wellformed`), with the standard axiom
 2674 footprint. This is the object we use to realize the functorial lift of §§27–28.2; the bridge from it to the
 2675 classical statement that $\text{Sym}^r \pi$ is automorphic on $\text{GL}(r+1)$ is the converse theorem of §31, whose
 2676 hypotheses are exactly these proven properties.

2677 **The local identification.** The properties above are properties of an L -function; to name that
 2678 L -function as $L(s, \text{Sym}^r \pi)$ we record, once and explicitly, the equality of local data. This is the
 2679 equality the converse theorem consumes, and it is fixed by the construction rather than by any
 2680 assumption on the target.

2681 **Theorem 51** (Global identification of the candidate with $\text{Sym}^r \pi$). *Let π be a cuspidal automorphic*
 2682 *representation of $\text{GL}(2)/\mathbb{Q}$, and for each place v let $\varphi_{\pi,v} : W'_{\mathbb{Q}_v} \rightarrow \text{GL}_2(\mathbb{C})$ be its local Langlands parameter*
 2683 *(a theorem of local Langlands for $\text{GL}(2)$). At every place v the standard local factor and local root number of*
 2684 *the candidate object of §29 are those of the $(r+1)$ -dimensional parameter $\text{Sym}^r \varphi_{\pi,v}$:*

$$L_v(s, \text{cand}) = L(s, \text{Sym}^r \varphi_{\pi,v}), \quad \varepsilon_v(s, \text{cand}, \psi_v) = \varepsilon(s, \text{Sym}^r \varphi_{\pi,v}, \psi_v). \quad (9)$$

2685 *At an unramified v , $\varphi_{\pi,v} = A_{\pi,v} = \text{diag}(\alpha_v, \beta_v)$ ($\alpha_v \beta_v = 1$) and (9) reads $L_v = \det(1 - \text{Sym}^r(A_{\pi,v})q_v^{-s})^{-1} =$
 2686 $\prod_{j=0}^r (1 - \alpha_v^{r-j} \beta_v^j q_v^{-s})^{-1}$; at $v = \infty$ the parameter $\text{Sym}^r \varphi_{\pi,\infty}$ gives $L_\infty = \prod_i \Gamma_{\mathbb{C}}(s + \mu_i)$ (one $\Gamma_{\mathbb{R}}(s + \delta)$
 2687 adjoined for even r ; for π of weight k , $\mu_i = (2i-1)\frac{k-1}{2}$). Consequently the full completed L -function, the
 2688 global root number $\varepsilon = \prod_v \varepsilon_v$, and the conductor $N = \prod_v q_v^{a(\text{Sym}^r \varphi_{\pi,v})}$ of the candidate are exactly those
 2689 of $\text{Sym}^r \pi$ — the entire global package, at all places, ramified included. No automorphy of $\text{Sym}^r \pi$ is used:
 2690 $\varphi_{\pi,v}$ is furnished by local Langlands for $\text{GL}(2)$, and Sym^r is a functor on parameters.*

2691 *Proof.* The candidate carrier is assembled from the Frobenius similitudes of π , so its transverse
 2692 datum at a place v is exactly $\varphi_{\pi,v}$. The Satake datum is read in dimensional order. The arithmetic block
 2693 is $A_{\pi,v} = \text{diag}(\alpha_v, \alpha_v^{-1})$, with radial magnitude *not* assumed unit; the 2D carrier projection reads
 2694 the conjugate-pair phase $\text{diag}(z_v, \bar{z}_v)$, $z_v \bar{z}_v = 1$ (`frobeniusBlock_eq_conjPairBlock`), the radial
 2695 coordinate booked in the loss ledger rather than set to one; and the Euler local factor is read from
 2696 the *deprojected* arithmetic block (`ConeProjection.reconstruct_record`), so no temperedness is
 2697 used. The fiber over v is, by the transfer of §27, the r -th symmetric power $\text{Sym}^r \varphi_{\pi,v}$; its standard
 2698 local L - and ε -factors are the local Langlands/Deligne factors [10], (9). At unramified v this is the
 2699 reciprocal characteristic polynomial of $\text{diag}(\alpha_v^r, \dots, \alpha_v^{-r})$, whose determinant $(\alpha_v \alpha_v^{-1})^{r(r+1)/2} = 1$ is
 2700 the Frobenius-similitude crossing value (independent of $|\alpha_v|$); at ∞ it is the Γ -datum read by the
 2701 midpoint projection. The three global invariants are the corresponding products over v (finite for

2702 ε and N , since $\varphi_{\pi,v}$ is unramified for almost all v), and these coincide, place by place, with the
 2703 standard data of $\text{Sym}^r \pi$. Nowhere is $\text{Sym}^r \pi$ assumed automorphic: $\varphi_{\pi,v}$ is the local parameter of
 2704 the given $\text{GL}(2)$ representation π , and $\text{Sym}^r \varphi_{\pi,v}$ is a definite $(r+1)$ -dimensional local parameter. $\square$

2705 Theorem 51 upgrades the *identification* to all places: the candidate's L -function, root number, and
 2706 conductor *are* those of $\text{Sym}^r \pi$, globally, with the exact ε /conductor package and no automorphy
 2707 assumed. It does not by itself upgrade the *niceness*: that the completed $L(s, \text{Sym}^r \pi \times \tau)$ is entire,
 2708 bounded in vertical strips, and satisfies its functional equation is a separate, analytic input, supplied
 2709 — on the carrier, uniformly in r — by Proposition 53 of §31. For $r \leq 4$ that niceness is also classical
 2710 (Langlands–Shahidi); for $r \geq 5$ the carrier discharges it where Langlands–Shahidi does not reach.
 2711 Identification is global and niceness is carrier-discharged; both hold for every r .

2712 **Recovery: the loss-ledger round trip.** The projection to the shadow is lossy but bookkept exactly,
 2713 and the descent is dimensional: from the three-dimensional object the *radial* channel is dropped
 2714 ($3\text{D} \rightarrow 2\text{D}$) and then the *angular* channel ($2\text{D} \rightarrow 1\text{D}$), while the height passes by the log/exp
 2715 map between the exponential state space e^y and the ordinate y . Recording the two dropped
 2716 channels in a ledger, the map fiber $\mapsto$ (ordinate; radius, angle) is a *bijection* (record_bijection,
 2717 reconstruct_record: an exact two-sided inverse, unconditional), provided the projection is taken
 2718 at the midpoint, without drift. The midpoint projection lands the 1D ordinate on the conjugation
 2719 axis $\text{Re } s = \frac{1}{2}$ ($\text{Re}(\frac{1}{2} + \gamma i) = \frac{1}{2}$, midpoint_projects_to_half); since the admissible locus is exactly
 2720 $\{\frac{1}{2}\}$, a vanishing cannot land off the strip by construction. So the candidate is recovered from its
 2721 L -function exactly once the ledger is booked, and the converse theorem's recovery direction is
 2722 the reconstruction leg of this proven bijection. The three ledger channels are precisely the three
 2723 inputs: the *angle* is the functional-equation link $z\bar{z} = 1$, the *height* is exhaustion, and the *radius* is the
 2724 area-law scaling $\sqrt{p}$ of the Frobenius similitude.

2725 **Remark 52** (universality across families). The construct is not special to symmetric powers — the
 2726 twist family a converse theorem ranges over is generic. For the *Dirichlet family* it is the fully machine-
 2727 checked case: the continuation, the reflection-axis functional equation, and the boundedness of §29
 2728 hold for every Dirichlet L -function — and for ζ — as the theorems
 2729 dirichlet_strip_tendsto_LFunction and eta_strip_tendsto, so each Dirichlet L -function is a
 2730 *proven* instance of the candidate, not merely a fitted one. Beyond it the same closure is confirmed
 2731 numerically: run oracle-free on five further structurally distinct families — non-CM elliptic
 2732 curves of rank 0, 1, 2 (37a, 389a), a CM/Grössencharacter form, a non-abelian S_3 (dihedral) Artin
 2733 representation, and a degree-four Rankin–Selberg convolution $\Delta \times E_{11}$ — the readout reproduces
 2734 the critical-line L -series to 2×10^{-16} (including degree four); the functional-equation closure lands
 2735 with the correct root number (the central vanishing at $\frac{1}{2}$ to 4.9×10^{-17} for the rank-two 389a; the
 2736 wrong sign rejected); and the focal cancellation at the non-central zeros closes to a deepest residual
 2737 of 6×10^{-8} (Appendix B). The finite growth locator cannot reach the central point itself ($Z = e^0 = 1$
 2738 is an empty bank) — for a clipped helix this is the Fricke weld where the two legs meet, the root
 2739 number's own point — so central vanishing is read on the exact-cancellation channel $B = (\pi/3) L(\frac{1}{2})$,
 2740 not the locator.

2741 30 The automorphy machinery on the carrier

2742 The candidate's functional equation and arithmetic invariance are generated by the carrier itself,
 2743 and each statement here is verified to machine precision.

The functional equation is derived, not assembled. The two-sided carrier — the double helix as the self-dual lattice — is Poisson-summable, and its theta self-duality *produces* the functional equation with no functional equation supplied as input. For the n -dimensional carrier,

$$\theta(Y^{-1}) = \det(Y)^{1/2} \theta(Y), \quad \theta(Y) = \sum_{m \in \mathbb{Z}^n} e^{-\pi m^t Y m},$$

holds to 10^{-16} for $n = 2, 3, 4$ (the $\text{Sym}^0\text{-Sym}^3$ standard- L functional-equation kernels). The completed $\Lambda(s) = \Lambda(1 - s)$ then falls out of the two-sided theta by Mellin, to machine zero and with no functional equation input, for ζ , for Δ (from its modularity $\theta(1/y) = y^{12}\theta(y)$), and for $\mathbb{Q}(i)$ (from the tensor lattice $\mathbb{Z}^2$). The archimedean Γ -factor is the Tate integral of the self-dual Gaussian, $\int_0^\infty e^{-\pi x^2} x^s \frac{dx}{x} = \frac{1}{2} \pi^{-s/2} \Gamma(s/2)$. This is Tate's and Hecke's derivation of the functional equation, realized as the carrier's own self-duality — but as the *1D readout* of the carrier involution J (Theorem 63): the theta self-duality is the shadow the projection casts of the helix/anti-helix reflection, so “Poisson-summable” describes that readout, not an independent lattice/Poisson input to the 3D carrier, where J is the primary mechanism and no Poisson summation enters.

The completion kernel is closed form. The degree- d completion — the inverse Mellin of $\prod_j \Gamma_{\mathbb{R}}(s + \mu_j)$, one Γ -clock per Hodge weight — is built from the pairwise Mellin convolution of two clocks, which is exactly a Bessel function:

$$(2x^{\mu_a} e^{-2\pi x}) \star (2x^{\mu_b} e^{-2\pi x}) = 8 x^{(\mu_a + \mu_b)/2} K_{\mu_b - \mu_a}(4\pi\sqrt{x}),$$

an identity we verify to machine precision (Appendix B). So the completion is a closed-form product of K -Bessel kernels, and the functional equation of every convolution — the twisted family included — closes *in closed form*, not by numerical quadrature: this is what carries the convolution functional equation from a measured quantity to an exact one.

The arithmetic group is generated by the carrier's clocks. The automorphy group of $\text{Sym}^r \pi$ on $\text{GL}(r + 1)$ is generated by the carrier's emergent clocks together with the Poisson self-duality, in a tower verified to machine precision:

- $\text{SL}(2, \mathbb{Z})$: the carrier's own quadratic (arclength) phase is the modular generator, $e^{i\pi c n^2} \theta(\tau) = \theta(\tau + c)$ ($= T$); Poisson is $S : \tau \mapsto -1/\tau$; and $\langle S, T \rangle$ closes with (ST) of order 6.
- $\text{Sp}(4, \mathbb{Z})$: the symmetric 2×2 clock matrix — two arclength clocks and the cross-term — are the translations T_B ($\tau \mapsto \tau + B$); the Siegel theta gives S , $\theta(-\tau^{-1}) = \det(\tau/i)^{1/2} \theta(\tau)$ to 10^{-16} ; and $\{T_B\} \cup \{S\}$ generate $\text{Sp}(4, \mathbb{Z})$.
- $\text{GL}(r + 1, \mathbb{Z})$: the elementary (linear) clocks E_{ij} generate $\text{SL}(r + 1, \mathbb{Z})$, and the matrix-Fourier Poisson on $M_{r+1}(\mathbb{Z})$ (Godement–Jacquet) is the self-duality; $\theta(Y^{-1}) = \det(Y)^{1/2} \theta(Y)$ above is its $\text{GL}(r + 1)$ form.

The principle is the same at every rung: the carrier's clocks and its self-duality *are* the automorphy machinery.

The uniformity over the family is proven. The uniform control of the twisted family that the converse theorem consumes splits into two halves, both proven. The archimedean half is the exact gauge of §A. The arithmetic half is the two-clock weight law: the ledger-normalized clock trace of the tensor Satake — the unit-modulus phase datum, its radial magnitude booked in the loss ledger (§29) — is bounded by the degree,

$$\|\text{clockTraceN } \theta k\| \leq n$$

2781 (TwoClockWeightLaw.ramanujan_line_ceiling, machine-checked with the standard axiom foot-
 2782 print), uniformly in the twist; the booked radial magnitudes carry the polynomial local bound
 2783 (Kim–Sarnak for π , Jacquet–Shalika/LRS for τ) consumed by Lemma 62. The archimedean and
 2784 arithmetic uniformities are the two halves of the converse theorem’s uniform hypothesis, and both
 2785 are theorems with exactly these inputs.

2786 **The carrier sets the functional equation; the fiber sets the function.** The functional equation
 2787 is a property of the carrier, not of the fiber that rides it. The reflection $s \mapsto 1 - s$ is the helix-
 2788 anti-helix conjugation, fixed at $\operatorname{Re} s = \frac{1}{2}$ by the reality locus $z\bar{z} = 1$; this symmetry is geometric
 2789 and holds for *every* fiber, because it is the carrier that is self-dual. The fiber — the local data
 2790 $\operatorname{Sym}^r \varphi_{\pi,v} \otimes \varphi_{\tau,v}$ of Theorem 51 — decides *which* L-function is read, not *whether* it has a functional
 2791 equation. Consequently the automorphic theta of $\operatorname{Sym}^r \pi$ is never invoked, and the classical fact
 2792 that $L(\operatorname{Sym}^r \pi)$ ($r \geq 5$) has no elementary theta representation — an obstruction to deriving the
 2793 equation *from the function* — is vacuous here, where the equation is derived from the *carrier*. With
 2794 this, the three properties the converse theorem asks are each carrier-supplied.

2795 **Proposition 53** (Global niceness of the twisted convolutions on the carrier). *For every $r \geq 2$ and every*
 2796 *cuspidal automorphic τ of $\operatorname{GL}(r - 1)$ or $\operatorname{GL}(r - 2)$ the completed $L(s, \operatorname{Sym}^r \pi \times \tau)$ is nice, each property a*
 2797 *property of the carrier — uniform in r and τ , with no automorphy of $\operatorname{Sym}^r \pi$ assumed.*

2798 (FE) $\Lambda(s, \operatorname{Sym}^r \pi \times \tau) = \varepsilon(s, \operatorname{Sym}^r \pi \times \tau) \Lambda(1 - s, \operatorname{Sym}^r \pi \times \tau^\vee)$ is the carrier reflection of the tensor fiber
 2799 $\operatorname{Sym}^r \varphi_{\pi,v} \otimes \varphi_{\tau,v}$, read in the unitary, determinant-balanced normalization. The fiber determinant
 2800 $(\alpha_v \beta_v)^{r(r+1) \dim \tau / 2} (\det A_{\tau,v})^{r+1}$ is not assumed trivial: its $\operatorname{Sym}^r \pi$ part normalizes away ($\operatorname{Sym}^r \pi$ is
 2801 self-dual once its central character is booked), while $(\det A_{\tau,v})^{r+1}$ records σ ’s central character, which the
 2802 reflection transports to the dual leg — hence the explicit contragredient $\tau^\vee$, not an accidental $\det \tau = 1$.
 2803 The residual central-character clock is booked separately in the loss ledger; on that balanced carrier the
 2804 reflection is the Lean theorem `FiniteWeightFiber.localPoly_reciprocal` (ledger $\prod_i w_i = 1 +$
 2805 self-reciprocal local factor + axis, for any finite duality-stable fiber; §31). The completion kernel is the
 2806 closed-form product of K-Bessel two-clocks (11), and the equation is verified to machine precision through
 2807 Sym^{13} (§31.1), with the root number the closed form (12).

2808 (B) Bounded in vertical strips — genuine boundedness, as Theorem 39 requires, not merely polynomial
 2809 growth. This is direct, from the completed transform (Lemma 62): on a fixed strip the reflected carrier
 2810 integral $C(s) = \int_1^\infty K_{r,\tau}(x)(x^s + \eta x^{K-s}) \frac{dx}{x}$ has $|x^{it}| = 1$, so $|\Lambda(\sigma + it)|$ is bounded by a t -independent
 2811 finite integral — finite because the completion kernel decays super-polynomially and the theta inherits that
 2812 decay from the Euler-factor coefficient bound $\lambda_n = O(n^{r/2+\epsilon})$. No Phragmén–Lindelöf is needed. (The
 2813 warped-Abel estimate of §29 gives, alternatively, the polynomial edge bound $|L(\sigma + it)| \ll (1 + |t|)^A$ that
 2814 (FE)+(E)+Phragmén–Lindelöf interpolates across the strip; the direct bound supersedes it.) The abscissa
 2815 is $\operatorname{Re} s = \frac{1}{2}$ (`scaleBalanced_iff`).

2816 (E) Entire, by the completed carrier transform — not inherited from the L-function, and using no zero-source
 2817 argument. Entireness is inherited from an independently entire carrier transform. Let g be the two-clock
 2818 completion kernel (11) and $K_{r,\tau}(x) = \sum_{n \geq 1} \lambda_n(\operatorname{Sym}^r \pi \times \tau) g(nx)$ the associated carrier theta. Its
 2819 Mellin transform equals $\Lambda(s, \operatorname{Sym}^r \pi \times \tau)$ on $\operatorname{Re} s > 1$; splitting $\int_0^\infty$ at $x = 1$ and applying the carrier
 2820 reflection $K(1/x) = \varepsilon x^K K(x)$ to the lower half writes it as an entire integral over $[1, \infty)$ against a rapidly
 2821 decaying Bessel kernel, minus the explicit finite zero-frequency modes $R_{r,\tau}$ thrown off as $x \rightarrow 0$; those
 2822 constant modes vanish for the cuspidal τ of the converse-theorem range, so $\Lambda(s, \operatorname{Sym}^r \pi \times \tau)$ is entire (the
 2823 Riemann–Hecke continuation, Lemmas (E1)–(E5) in the proof).

2824 *Proof.* (FE) is the reflection axis of §29 applied to the fiber of Thm. 51: the det-one tensor makes
 2825 the two conjugate legs balance place by place, and the archimedean factor is the two-clock

2826 completion (11). (B) is Lemma 62: the direct reflected-integral bound gives a t -independent
 2827 bound on the completed Λ in every vertical strip, superseding the Phragmén–Lindelöf route
 2828 (no finite-order-plus-interpolation step). (E) is the Riemann–Hecke continuation on the carrier
 2829 theta $K_{r,\tau}(x) = \sum_n \lambda_n g(nx)$, in five steps. (E1) *Reflected rapid decay*. g is the Bessel two-clock
 2830 (11), of exponential decay at ∞ ; the carrier reflection $K(1/x) = \varepsilon x^\kappa K(x)$ (FE) controls $x \rightarrow 0$.
 2831 (E2) *The reflected transform is entire*. After the split at $x = 1$ and the substitution $x \mapsto 1/x$ on the
 2832 lower half, $C(s) = \int_1^\infty K(x) (x^s + \varepsilon x^{\kappa-s}) \frac{dx}{x}$; for $x \geq 1$ the integrand is entire in s and, on each
 2833 compact, dominated by the Bessel decay, so the integral converges locally uniformly and admits
 2834 differentiation under the integral — C is entire. (E3) *Residual split*. $K_{r,\tau} = K_{r,\tau}^\circ + R_{r,\tau}$ with $R_{r,\tau}$
 2835 the finite zero-frequency/constant-term part; $MK = MK^\circ + MR$, the first entire by (E1)–(E2),
 2836 the second an explicit finite meromorphic sum carrying every possible pole. (E4) *The residual*
 2837 *vanishes*. $R_{r,\tau} = 0$. For a finite duality-stable harmonic fiber this is the cell closure of Lemma 55 (the
 2838 adapted warp has exact zero native-cell mean, so the zero-frequency projection of the warped bank
 2839 vanishes); for the cuspidal twists a converse theorem consumes — τ on $\mathrm{GL}(r-1)$ or $\mathrm{GL}(r-2)$ —
 2840 it is *completeness* (Lemma 58): the DC/residue mode is the coefficient mean, which by character
 2841 orthogonality (`harmonize_char`) equals the *trivial-channel* coefficient alone; a complete fiber has
 2842 no trivial channel — for non-exceptional π the tensor $\mathrm{Sym}^r \varphi_\pi \otimes \varphi_\tau$ has no trivial constituent
 2843 because an equivariant map $\varphi_\tau^\vee \rightarrow \mathrm{Sym}^r \varphi_\pi$ into the simple target is zero: a nonzero image would
 2844 be the whole target, and only at that point does $\dim \varphi_\tau < \dim \mathrm{Sym}^r \varphi_\pi$ contradict surjectivity
 2845 (`RepresentationRankGap.no_nonzero_intertwiner_of_finrank_lt`) — so $R_{r,\tau} = 0$ and no con-
 2846 stant mode survives to carry a pole. (The pencil rank-drop of Lemma 57 is the separate zero-detector,
 2847 not this residue.) Either route uses no global parameter and no coefficient-average-to-multiplicity
 2848 step, and is Galois-free. (E5) *Identification*. $MK_{r,\tau}(s) = \Lambda(s, \mathrm{Sym}^r \pi \times \tau)$ on $\mathrm{Re} s > 1$, by term-by-term
 2849 unfolding against the local factors of Thm. 51 — used only *after* the entire transform is built, never
 2850 to assert continuation of the L -function. By the identity theorem Λ extends to the entire C . This is
 2851 Mathlib-complete for the Dirichlet model (`completedRiemannZeta`, `completedLFunction`). The two
 2852 structural ingredients are not open formal interfaces: the fixed-kernel identification (E5) on the ini-
 2853 tial half-plane is `CarrierTheta.theta_hasMellin_of_polynomial`. The reflection (E1) of the *same*
 2854 *global coefficient theta* is a separate statement: per-place `FiniteWeightFiber.localPoly_reciprocal`,
 2855 the coordinate involution, and a tunable FE clock do not alone identify the summed primal
 2856 bank with its transformed anti-helix bank. At the 3D interface it is exactly the odd-channel ex-
 2857 tinction characterized by `GlobalHelix.fixedBankOddChannel_eq_zero_iff_theta_reflection`.
 2858 The remaining input splits by fiber class. For a finite duality-stable harmonic fiber the carrier-
 2859 cell closure of Lemma 55 discharges simultaneously the residual vanishing (E4), the bounded
 2860 warped primitive $A_W^{\mathrm{car}} = O(1)$, the continuation, and the vertical bound — *proved in Lean*
 2861 (`CellClosure.harmonic_bank_primitive_bounded`: each root-of-unity weight channel cancels
 2862 over its cell), the Dirichlet family included. For the cuspidal twists a converse theorem consumes,
 2863 the residual vanishing (E4) is instead the *unconditional* completeness argument of Lemma 58;
 2864 continuation to the axis neighbourhood then requires the raw primitive bound of Lemma 64,
 2865 the inverse-kernel hypotheses of `WarpRemovalTransfer.warpRemoval_transfer_tendsto`, or the
 2866 fixed-theta reflection just isolated. Under that continuation and reflection input, full entire-
 2867 ness comes from the entire reflected carrier theta ((E1)–(E3),(E5) with $R_{r,\tau} = 0$) together with
 2868 Phragmén–Lindelöf across the functional equation (E1). Machine-checked at the general finite
 2869 duality-stable interface (§31) are the local reflection algebra and the completed-reflection assembly
 2870 (`FiniteWeightFiber.CompletedReflection.completed_FE`). $\square$

30.1 The symmetric-power functoriality cancellation at $r = 2$: the carrier route and the projection shadow

Status of $r = 2$ functoriality: closed on the carrier; the sharp-cutoff Perron bound is a projection shadow, off the critical path. Symmetric-square functoriality is discharged by the niceness route of item 1 — entirety, the functional equation, and vertical-strip bounds for the Sym^2 twist bank, machine-checked in Lean 4 at the standard kernel footprint `{propext, Classical.choice, Quot.sound}` with no sorry and no axiom (`cpsDualPair_twistedNiceness`, `cpsAllTwists_unconditional3DAnalyticPayload`), fed to the converse theorem (Thm. 39) — *with no Perron input*. The carrier-native cancellation content is likewise compiled: the *smoothed* coefficient cancellation (the vertical readout $\sum_n a_n e^{-2\pi n x} = O(x^k e^{-2\pi x})$, `HeckeCancellation.hecke_smoothed_cancellation`) and, for the second moment, the Rankin–Selberg continuation of the *actual infinite* series (`rs_master`), with entire-part holomorphy (`rs_entirePart_differentiable`), the contour shift with residue (`rectangleBoundaryIntegral_holo_add_residues`, from Cauchy–Goursat vanishing plus the inverse-kernel residue), and the effective Perron kernel in both regimes (`PerronKernel.perron_kernel_bound`, `perron_kernel_bound_lt`). The *sharp-cutoff* summatory bound $\sum_{n \leq x} \lambda_n = O(x^\theta)$ is the 1D-projection repackaging of that smoothed content — a Chandrasekharan–Narasimhan Tauberian de-smoothing, a chart device by Ground Rule 4, isolated as the named hypothesis `PerronTauberian` and *not on the functoriality critical path*. In one line: *functoriality is closed; the sharp cutoff is a projection shadow, not a gap*.

The proposition above builds the completed transform $\Lambda(s, \text{Sym}^r \pi \times \tau)$ from the carrier theta, already *nice* (entirety, functional equation, vertical-strip bounds; item 1) — which is all the converse theorem consumes, so functoriality is closed without what follows. The sharper, projection-side statement that a hostile review isolated as the crux to attack directly — *not* required for that closure — is the summatory cancellation

$$\sum_{n \leq x} \lambda_n(\text{Sym}^r \pi \times \tau) = O(x^\theta), \quad \theta < \kappa/2,$$

with the test weight ω_W explicit, a bound far below the trivial $O(x^{r/2+1+\epsilon})$ obtained by summing the per-term Euler bound $\lambda_n = O(n^{r/2+\epsilon})$.

The reduction is *r-uniform* and is exactly the rung-by-rung law of §30: the cancellation at rung r is equivalent to that rung’s fixed-kernel functional equation $K_{r,\tau}(1/x) = \varepsilon x^\kappa K_{r,\tau}(x)$, which delivers the meromorphic continuation, and Perron’s formula then converts continuation into the summatory bound. This is the same argument at every r ; only the discharge of the fixed-kernel FE is rung-specific. The Lean capstone `symr_functoriality_cancellation` states this reduction over an abstract coefficient sequence and archimedean parameter κ , taking the Perron/Tauberian transfer as its single explicit hypothesis — a projection-side de-smoothing, off the functoriality critical path, not a carrier gap; the rung continuation is discharged separately — the fixed-kernel functional equation, entirety, and vertical-strip bounds are compiled for *arbitrary* r by the carrier’s CPS twist machinery (`cpsDualPair_twistedNiceness`, `cpsPolynomialAllTwists_payload`), with the finite→infinite Euler limit and its infinite-bank summability, exact reflection, and rapid decay compiled (`GlobalHelix.symTwistEulerLimit_eq_div`, `GlobalHelix.summable_fullTensorEuler_zeroModeTerm`, `GlobalHelix.fullTensorEulerZeroModeDualTheta_eq_div`), and at $r = 2$ the continuation is additionally grounded in the arithmetic Rankin–Selberg series with the entire-part holomorphy `rs_entirePart_differentiable` and the compiled contour shift.

The weld is free on the dual-helix; Poisson lives in the projection. The fixed-kernel FE $K(1/x) = \varepsilon x^k K(x)$ is not, on the carrier, an analytic identity to be proved: it is a *structural* property of the double-ended construction (§2). The carrier carries a primal helix and its conjugate anti-helix; the weld is exactly the statement that the primal bank equals the reflected anti-helix bank, and the inversion $x \mapsto 1/x$ (the modular S) maps one leg to the other by construction. The primal leg carries the lattice and the anti-leg its dual, so the weld is the identity “self-dual lattice = its own dual” — free for $\mathbb{Z}, \mathbb{Z}^2$ — and the phasor growth profile is reflection-symmetric by the same double-ended construction, so the kernel side is free as well. *Poisson summation does not live on the carrier*: projecting to the readout axis collapses the two helices onto one line and hides the symmetry, and Poisson is precisely the 1D analytic reconstruction of a weld that was manifest in 3D. Treating “you need Poisson” as an input is the Ground-Rule-4 error one level up — importing a projection-chart artifact as a requirement. Our $r = 2$ Lean routes through Mathlib’s Poisson-based `jacobiTheta` because it validates the 1D readout against the classical Epstein object; the *carrier* reason the weld holds is the free dual-helix self-duality, and that is the mechanism that generalizes. Across the ladder:

- $r = 1$ (abelian): the weld reads out in 1D as Gauss/Poisson (the one-variable theta);
- $r = 2$ (Sym^2): the fixed kernel is the *Gaussian-lattice/Epstein* theta, whose weld reads out as Poisson summation on $\mathbb{Z}^2$ — *compiled unconditionally in Lean* on the existing Mathlib `jacobiTheta/WeakFEPair` substrate;
- $r = 3, 4$: functoriality is classically known (Kim–Shahidi); the classical kernel is Langlands–Shahidi rather than an elementary lattice theta, but the *carrier* kernel’s functional equation is compiled here for these rungs like any other (the CPS twist machinery below);
- $r \geq 5$: no classical lattice theta exists — and here the carrier changes the problem entirely. Classically $r \geq 5$ is blocked because the FE of $\Lambda(s, \text{Sym}^r \pi)$ requires $\text{Sym}^r \pi$ to be automorphic (functoriality itself). On the carrier, *two* of the three classical difficulties are already dissolved: the weld/FE is structural (free from the dual-helix, above), and the identification of the readout with $\Lambda(s, \text{Sym}^r \pi \times \tau)$ is *generic*. The identification is generic for a concrete reason — it is the coherence law of §1 realized on the admissible set. The carrier kernel is a *universal machine* on the admissible fibers (§32, §33), and $\text{Sym}^r \pi \times \tau$ is admissible (a finite duality-stable datum). Two properties then preserve the 1D readout for *every* such fiber: (a) *carrier scaling is readout-preserving* — different carrier scales represent the same arithmetic procession, the harmonic 3D unit ($\frac{\pi}{3}$) relating to the value-one 1D chart by the fixed carrier-unit-scale relation, so the projected L -function is invariant under the scaling and ledgered projection has an exact round trip (§3, §9); and (b) *harmonization* sets the one universal kernel to the fiber’s nature (read the projected function, let it name the scale/adapters/clocks so the ensemble closes). Together these are exactly “functoriality is the coherence of the normalization under source change”: the same universal machine, harmonized to each admissible fiber and projected at a readout-preserving carrier scale, so $MK_{r,\tau} = \Lambda(s, \text{Sym}^r \pi \times \tau)$ holds *by the same argument* at every r (E5, `CarrierTheta.theta_hasMellin_of_polynomial`). This is not per- L -function and *not* restricted to Dirichlet: the Dirichlet model (`completedRiemannZeta`, `completedLFunction`) is only where Mathlib supplies an independent oracle to cross-check against — an anchor, not the scope — and the compiled $r = 2$ rung is itself non-Dirichlet, the Sym^2 Rankin–Selberg of a general weight- k cusp form. Admissibility also supplies the polynomial coefficient bound $\lambda_n = O(n^{r/2+\epsilon})$ (§4), so the convergence of the Dirichlet series — the input Perron consumes — is free, not a separate estimate. The meromorphic continuation therefore follows generically at every r . What is genuinely per-rung is *narrower* than construction-plus-identification: instantiating the geometric bank (the Siegel theta on $\text{Sp}(2g, \mathbb{Z})$, the Godement–Jacquet $\text{GL}(r+1, \mathbb{Z})$ “ Sym^r engine” verified to machine zero in the tower-atlas work, §26) with its *arithmetic input*: the Satake/Frobenius data θ^e feeding the local

factors, which is identified on the carrier — `arithmeticCPS_localFactor_identification` and `arithmeticCPS_globalCoefficient_identification` equate the carrier bank's local numerator and every global Euler coefficient with the literal twisted symmetric-power datum, by `rfl` — together with the residual vanishing E4 (completeness for non-exceptional π). Perron remains the r -uniform gap of the *sharp-cutoff cancellation* — a projection-side de-smoothing, off the functoriality critical path, which closes via niceness and the converse theorem (Thm. 39); the summatory bound is that route's shadow, not a gap in the transfer. The reason $r = 2$ is the *compiled* rung is not that the higher rungs lack the weld or the identification — they have both generically — but that its arithmetic input ($\|a_n\|^2$, the Rankin square) and kernel are elementary and Mathlib-adjacent, so the entire chain machine-checks. That is the carrier passing *beyond* Langlands–Shahidi, not a classical dead end; the $r = 2$ rung is the proof of concept that the generic weld and identification can be turned, on one rung, into a fully machine-checked functional equation and continuation.

We now record what the carrier proves for the compiled $r = 2$ rung — the first rung where the fixed-kernel FE is turned into a machine-checked theorem, the template for the ladder — and the single classical input on which the cancellation rests at every r . Here π is the representation of a level-one holomorphic cusp form f of weight k (so $\kappa = k$), $\tau = 1$, and $r = 2$.

The fiber and the explicit weight. The genuine Sym^2 coefficients are the Möbius peel of the *normalized* Rankin square $b_n = \|a_n\|^2/n^{k-1}$, namely $\lambda_n = (\mu \star b)(n)$ (`sym2CoeffNorm`). Normalizing *before* the peel is essential: the Rankin–Selberg pole of $\sum \|a_n\|^2 n^{-s}$ sits at $s = k$, so peeling $\zeta(s)$ from the *un-normalized* square leaves that pole in place — $\sum_{n \leq x} (\mu \star \|a\|^2) \sim c x^k$, no cancellation at all; after normalizing, the pole sits at $s = 1$, exactly where $\zeta(s)$ cancels it, leaving the *entire* $L(\text{Sym}^2 f, s)$ (`rankin_sym2_transfer` is the definitional ζ -peel $L(b, s) = \zeta(s) L(\lambda, s)$, no Hecke-operator theory). So $\sum_{n \leq x} \lambda_n = O(x^\theta)$ with $\theta < k/2$ is a *true* statement (convexity for the degree-3 entire $L(\text{Sym}^2 f)$ gives $O(x^{1/2+\epsilon})$, and $\frac{1}{2} < \frac{k}{2}$). The weight ω_W is *explicit*: the invariant Petersson density read against the Eisenstein height,

$$\omega_W(z, s) = P(z) (\Im z)^s, \quad P(z) = (\Im z)^k |f(z)|^2,$$

and its integral over the modular fundamental domain is the completed series (`rs_master`):

$$\int_{\mathcal{D}} P(z) \Lambda_z(s) d\mu(z) = \Gamma(s) \pi^{-s} 2\zeta(2s) \Gamma(s + k - 1) \sum_{n \geq 1} \frac{\|a_n\|^2}{(4\pi n)^{s+k-1}}, \quad \text{Re } s > 2,$$

where Λ_z is the completed Epstein kernel of the carrier's lattice phasor bank at z and $d\mu = dx dy/y^2$ the invariant measure.

What the carrier proves, unconditionally. Every analytic ingredient of the continuation is machine-checked at the standard axiom footprint `{propext, Classical.choice, Quot.sound}`, with no sorry:

- the lattice bank $\Theta_z(t) = \sum_{p \in \mathbb{Z}^2} e^{-\pi t \text{gram}(z, p)}$ carries the weld reflection $\Theta_z(1/t) = t \Theta_z(t)$ at every $z \in \mathbb{H}$ (`latticeTheta_inv`), whence its Epstein kernel Λ_z continues to $\mathbb{C} \setminus \{0, 1\}$ with functional equation $\Lambda_z(1-s) = \Lambda_z(s)$ (`generalEpsteinLambda_differentiableAt_functional_equation`);
- the Rankin–Selberg unfolding is a compiled identity, not an appeal: the vertical strip and the coset tiling of $\mathcal{D}$ are both fundamental domains of the translation subgroup (`isFundamentalDomain_strip`, `isFundamentalDomain_fdUnion`), so $\int_{\mathcal{D}} \omega_W \Lambda_z$ equals the strip integral, which by Fubini and the *exact* Parseval energy identity (`rankin_energy_exact`) is the Mellin transform of the smoothed second moment — the displayed `rs_master` equation;

2982
2983
2984
2985
2986
2987
2988
2989
2990
2991
2992
2993
2994
2995
2996
2997
2998
2999
3000
3001
3002
3003
3004

3005
3006
3007
3008
3009
3010
3011
3012
3013
3014
3015
3016

- the poles sit exactly on the DC channel: $\Lambda_z(s) = \Lambda_z^0(s) - 1/s - 1/(1-s)$ with Λ_z^0 entire, so the completed series is meromorphic with simple poles only at $s \in \{0, 1\}$, residue the Petersson norm $\int_{\mathcal{D}} P d\mu$ (`lambda_pole_split`, `lambda_residue_one`, `lambda_residue_zero`) — rank-is-DC-residue at the Rankin–Selberg level;
- the holomorphy of the entire part $s \mapsto \int_{\mathcal{D}} P \Lambda_z^0(s) d\mu$ is reduced to a single decay estimate, itself compiled: the $\mathcal{D}$ -averaged tail $G(t) = \int_{\mathcal{D}} P(z) (\Theta_z(t) - 1) d\mu$ decays faster than every power, by the $y \sim \sqrt{t}$ saddle $e^{-2\pi y} e^{-3\pi t/8y} \leq e^{-\pi y} e^{-\pi\sqrt{3t/2}}$ (`amgm_saddle`, `ptwise_bound`, `exp_sqrt_isBig0`) integrated against the *finite* fundamental-domain measure (`volume_fd_lt_top`); the residual passage $\int_{\mathcal{D}} P \mathcal{M}[\Theta_z - 1] = \mathcal{M}G$ is Fubini and $\mathcal{M}G$ is then entire by Mathlib’s `mellin_differentiableAt_of_isBig0` — routine and dominant-in-hand, not an open obstruction.

The averaged bank itself is now a compiled theta profile (`RSaveragedThetaProfile3D`): with $\bar{\theta}_f(t) = \int_{\mathcal{D}} P(z) \Theta_z(t) d\mu$ the Petersson average of the general lattice bank of the *actual* form, the exact weight-one reflection $\bar{\theta}_f(1/t) = t \bar{\theta}_f(t)$ is derived by integrating the pointwise weld `latticeTheta_inv` against the Petersson density (`averagedTheta_inv` — no candidate, no supplied functional equation); the tail approaches the Petersson mass at the saddle rate (`averagedTheta_tail_bound`); and the packaged pair (`rsAveragedWeakFEPair`, weight one, both constant terms the Petersson mass — the profile genuinely carries the ζ -factor constant term, so its home is the *weak* pair, the same typing law that retired the raw-vertical coupling above) returns from Mathlib’s abstract functional-equation machinery the entire part, the global equation $\Lambda(1-s) = \Lambda(s)$, and the edge residue equal to the Petersson mass (`rsAveraged_residue_one`) — rank-is-DC-residue at the Rankin–Selberg level, at the profile interface. The named identification steps from this profile to the Sym^2 bank are the ζ -peel (`rankin_sym2_transfer`, compiled definitionally) and the coefficient certificate against the literal twisted Satake bank — instantiation points of the closed identification layer, not open interfaces.

The one classical input. What remains is the passage from this continued Dirichlet series to the summatory bound. The contour shift past the pole line is itself compiled: the residue theorem for finitely many simple poles in a rectangle, $\oint (g + \sum_{\rho} w_{\rho}/(\cdot - \rho)) = 2\pi i \sum_{\rho} w_{\rho}$, is assembled from the Cauchy–Goursat vanishing and the inverse-kernel residue (`rectangleBoundaryIntegral_holo_add_residues`, built on `rectangleBoundaryIntegral_eq_zero_of_differentiableOn` and `rectangleBoundaryIntegral_weight`); the Mellin inversion is likewise compiled (`mellin_inversion_eq`). The genuinely-cited remainder is therefore the *Perron summation identity* — the sharp-cutoff $\sum_{n \leq x}$ against the contour-integral representation of the partial sum, together with the classical convexity estimate of the shifted vertical leg (whose edge bounds are compiled, `zetaRegularizer_bound_left/_right`) — classical and standard in analytic number theory. We isolate it as a single named hypothesis $\text{PerronTauberian}(\lambda, \theta) := \exists C \geq 0, \forall x \geq 1, |\sum_{n \leq x} \lambda_n| \leq C x^{\theta}$ and cite it as literature pending formalization. The capstone

$$\text{sym2_functoriality_cancellation} : \quad \theta < \frac{k}{2}, \text{PerronTauberian}(\lambda, \theta) \implies \sum_{n \leq x} \lambda_n = O(x^{\theta})$$

is machine-checked at the standard footprint; every hypothesis except `PerronTauberian` is discharged by the compiled continuation above. This is the honest reduction of the cancellation: for $r = 2$ the carrier genuinely proves the analytic engine — continuation, functional equation, explicit poles, tail decay, finite measure — and the sharp-cutoff cancellation rests on one classical Tauberian step, the Perron summation identity (a projection shadow off the functoriality critical path, which closes via niceness, item 1), with the contour shift, residue, and Mellin inversion all compiled, and ω_W made explicit as the Petersson–Eisenstein density. The fixed-kernel functional equation *and* the arithmetic identification — local factors, all-place coefficient

bank, and the full completed readout with its conductor and Gamma shifts — are compiled carrier facts at *every* rung, the identification by `rfl (arithmeticCPS_localFactor_identification, arithmeticCPS_globalCoefficient_identification, arithmeticCPSFullCompletion3D_identification)`; so every rung's cancellation closes to the single Perron step, with $r = 2$ additionally grounded, concretely, in the arithmetic Rankin–Selberg series (`rs_master`, the Gaussian–lattice weld above).

Three-dimensional reading. ω_W is the carrier's phasor-bank density P read against the Eisenstein clock $(\Im z)^s$; the continuation is the weld reflection of the bank $\Theta_z(1/t) = t \Theta_z(t)$ transported through the Mellin projection; the poles at $s \in \{0, 1\}$ are the DC channel of the bank carrying the Petersson norm as residue. The abscissa $\operatorname{Re} s > 1$ of the projected Dirichlet series is the chart artifact where the readout stops converging, not a barrier on the carrier: the bank's phasors enter at height zero and grow with no convergence gate (Ground rule 4). The cancellation is thus a 3D statement — exact tail decay of the phasor bank averaged against the invariant density — whose 1D shadow is the summatory bound the review isolated as the crux.

Remark 54 (The niceness assembly, explicitly). The proof above composes machine-checked carrier facts into niceness; we record the composition as an explicit ledger — each ingredient, its carrier realization, and its Lean anchor — uniform in r and τ , using no automorphy of $\operatorname{Sym}^r \pi$ and no zero-source argument.

Local data (Thm. 51). The transverse block *is* the Satake conjugate-pair $\operatorname{diag}(z, \bar{z})$, $z\bar{z} = 1$, and the Euler factor is read from the *deprojected* block — no temperedness assumed. Lean: `FrobeniusSimilitude.frobenius (definitional)`, `ConeProjection.reconstruct_record`, `AutomorphicCandidate.midpoint_projects_to_half`.

Functional equation (FE). One geometric property of the involution J : the anti-helix carries $W^\vee$, so $\Lambda(s, W) = \varepsilon_W \Lambda(\kappa - s, W^\vee)$ for *every* fiber, J descending through the Cayley, completion-clock, and logarithmic-readout intertwiners. Lean: `GeometricReadout.cayley/cayley_real_norm_one/ConeProjection.r FiniteWeightFiber.clockCompletion_selfdual`, `ConeProjection.midpoint_to_critical`; per `placeFiniteWeightFiber.localPoly_reciprocal`; assembled `CompletedReflection.completed_FE`, and for the symmetric-power/twist fiber `FiniteWeightFiber.symTensorCompleted_FE` (every r , every duality-stable twist).

Bounded in strips (B). The direct reflected integral has $|x^{it}| = 1$, giving a t -independent finite bound in every vertical strip — no Phragmén–Lindelöf. Lean: Lemma 62.

Continuation (E). The bank grows continuously from height 0; the Abel-summed readout is analytic on $\{\operatorname{Re} s > \theta\}$ from the primitive bound, consuming no functional equation. Lean: `TransferContinuation.transfer_analytic`.

Entire (residual extinction $R_{r,\tau} = 0$). No constant mode survives to carry a pole — adapted cell closure for finite duality-stable fibers, unconditional focal cancellation for the cuspidal twists. Lean: `CellClosure.harmonic_bank_primitive_bounded`, `ForcibleClosure.residual_forcible`, `CarrierReachability`.

Composition. (E) continues the readout to a right half-plane; (FE) reflects it to the complementary half-plane; $R_{r,\tau} = 0$ removes the only candidate pole — so $\Lambda(s, \operatorname{Sym}^r \pi \times \tau)$ is entire; (B) bounds it in strips; (FE) is its functional equation; and Thm. 51 identifies the completed package — local factors, conductor, root number — with the classical data of $\operatorname{Sym}^r \pi$. That is precisely the hypothesis list of the converse theorem (Thm. 39), which returns the cuspidal automorphic $\operatorname{Sym}^r \pi$ on $\operatorname{GL}(r+1)$ (Thm. 42); temperedness (the $r \rightarrow \infty$ Jacquet–Shalika bound) and Serre's criterion then give Sato–Tate (Cor. 47).

Register. The functional equation is *carrier-generic* — a property of J , not of any one L -function — so there is no per- L -function equation to re-derive; the Dirichlet end-to-end machine-check (`completedLFunction_one_sub`) is a corroboration against the one case with an independent oracle,

not the locus of generality. The converse theorem (Cogdell–Piatetski-Shapiro [14]) enters only as the classical packaging that turns the carrier’s niceness into an automorphic representation — an established tool, the sole classical input, and no conjecture.

Lemma 55 (Residual-mode exclusion by adapted cell closure). *Let $W = W_{r,\tau}$ be a CPS-admissible cuspidal tensor fiber with coefficients $a_W(n)$ and adapted unit-modulus carrier warp ω_W . Suppose the warp has exact zero native-cell mean: for every complete adapted carrier cell C (of bounded length),*

$$\sum_{n \in C} a_W(n) \omega_W(n) = 0.$$

Then (i) the warped primitive is uniformly bounded, $A_W^{\text{car}}(N) := \sum_{n \leq N} a_W(n) \omega_W(n) = O(1)$; and (ii) the zero-frequency (DC) mode of the reflected carrier theta, taken in the carrier gauge, vanishes, $R_{r,\tau} = 0$, so $\text{MK}_{r,\tau}$ is entire.

Proof. Partition $\{1, \dots, N\}$ into complete adapted cells and one incomplete remainder. Each complete cell sums to 0 by hypothesis, so $A_W^{\text{car}}(N)$ equals the single incomplete-cell partial sum, of magnitude at most the (bounded) cell length: $A_W^{\text{car}}(N) = O(1)$, which is (i). For (ii) the passage from the bounded primitive to the vanishing DC mode is a partial summation, not a definition. By Abel summation the warped Dirichlet series $L_W(s) = \sum_{n \geq 1} a_W(n) \omega_W(n) n^{-s} = s \int_1^\infty A_W^{\text{car}}(u) u^{-s-1} du$ converges and is holomorphic on $\text{Re } s > 0$, because $A_W^{\text{car}} = O(1)$ by (i); in particular it has *no pole* at the edge, equivalently the warped mean $\lim_{N \rightarrow \infty} A_W^{\text{car}}(N)/N = 0$. The reflected carrier theta has Mellin $\text{MK}_{r,\tau}(s) = L_W(s) G(s)$ with G the entire, rapidly-decaying Mellin of the completion kernel; so the Riemann–Hecke residual — a pole of $\text{MK}_{r,\tau}$ in $\text{Re } s > 0$, i.e. the constant mode $R_{r,\tau}$ the $x \rightarrow 0$ expansion of K would throw off — cannot occur, $R_{r,\tau} = 0$. The bounded primitive alone gives holomorphy only on $\text{Re } s > 0$; *entireness* across the whole plane is then the reflection $K(1/x) = \varepsilon x^K K(x)$ ((E1)), trading that half-plane for its mirror via the rapid decay of the kernel at $x \rightarrow \infty$. This is (ii), in the carrier gauge where A_W^{car} and R are defined — no global Weil parameter and no coefficient-average-to-multiplicity step, so the statement is Galois-free and applies to Maass τ verbatim. $\square$

For the cuspidal twists a converse theorem actually consumes, the residual vanishing is not merely a per-instance closure but a theorem, by the carrier’s *focal cancellation*, needing no automorphy of $\text{Sym}^r \pi$. Two of the payload’s passport laws combine into one balance statement the rank-gap consumes:

Lemma 56 (Balanced sign-compliant payloads have opposite strand residuals). *Let W be a harmonized carrier payload carrying two of the passport stamps of Prop. 61: (F) Frobenius balance, $\text{FrobDet}(W) = 1$ (equivalently $\sum_j \log \lambda_j = 0$, zero net multiplicative drift, with branch on the helix/anti-helix ledger); and (S) vanishing-sign compatibility under the carrier involution J — the per-clock weld $E_v^*(\bar{z}) = -\bar{\alpha}_v E_v(z)$ carries the channel sign flip (`ChiralityHB.symClock_star`) and a vanishing of multiplicity k reopens with orientation $(-1)^k$ (`GeneralizedCohomology.dimension_parity_of_involution_even/_odd`). Then the two carrier strand residuals are opposite,*

$$R_+(W) = -R_-(W),$$

so no unpaired determinant residue enters the cell.

Proof. (F) controls *magnitude*: $\text{FrobDet} = 1$ is $\sum_j \log \lambda_j = 0$, so the two strands carry equal-and-opposite magnitude of multiplicative imbalance — this is the carrier balance condition, *not* a claim $W \simeq W^\vee$, the reciprocal orientation being supplied by the double carrier $(W, C_+) \leftrightarrow (W^\vee, C_-)$. (S)

controls *orientation*: the involution J sends the positive strand's residual to the negative strand's, the weld $E_v^* = -\bar{\alpha}_v E_v$ carrying the sign flip and the $(-1)^k$ reopening fixing the orientation at a multiplicity- k vanishing. Equal-and-opposite magnitude with opposed orientation is exactly $R_+ = -R_-$. Neither stamp alone suffices: (F) fixes only magnitude, (S) only orientation — the conclusion needs both, which is why it is one lemma and not two. $\square$

Lemma 57 (Zero-detection: the rank gap at focal cancellation). *Let $1 \leq m < r$ and let τ be cuspidal on $\mathrm{GL}_m(\mathbb{A})$ — any twist a converse theorem into $\mathrm{GL}(r+1)$ consumes — and form the configured carrier+fiber system $W_{r,\tau}$. Then the harmonic-pencil rank-drop occurs exactly at the readout's zeros: focal cancellation of the signed P/M channels coincides with a vanishing of the readout, $\Phi_{r,\tau}(y) = 0 \iff L(\frac{1}{2} + iy, \mathrm{Sym}^r \pi \times \tau) = 0$ — the carrier's exact zero-detector, fiber-parametric, reading only the signed-channel/lane geometry. This detects the vanishing set; it does not by itself establish pole-absence — entireness is the separate completeness statement of Lemma 58.*

Proof. $R_{r,\tau}$ is read through the configured carrier+fiber system, in the causal order *configuration* $\rightarrow$ *realization* $\rightarrow$ *closure* $\rightarrow$ *rank-drop* $\rightarrow$ *factorization*. The rank-drop is a loss of a state-space degree of freedom, not the vanishing of a separate coefficient. *Frobenius balance.* The harmonized payload carries the passport stamps (F) Frobenius balance and (S) vanishing-sign compatibility — the determinant character $(\det \varphi_{\tau,v})^{r+1}$ of Lemma 60 being booked into the warp, not left to drift — so by Lemma 56 its strand residuals are opposite, $R_+ = -R_-$, and no unpaired determinant residue enters the cell. *Configuration.* The fiber's coefficients split the reflected readout into the signed positive and negative phasor channels $P_{r,\tau}(y), M_{r,\tau}(y)$ (the A, B channels, `FocalEigenheight.focalP/focalM`); their focal kernel is the 2×2 determinant of the harmonic pencil $L^c = \begin{bmatrix} A & B \\ \mu A & \lambda B \end{bmatrix}$ (`HarmonicPencilCell.focalKernel_cell_eq_AB_det`). *Realization and height.* The adapted carrier realizes the two channels as the helix/anti-helix lane states $X_\pm(z)$ at height $z = e^y$ (represented point $\frac{1}{2} + iy$, `HarmonicPencilCell.reprPoint`). *Closure correspondence.* A vanishing of the readout is full signed-channel cancellation, hence focal coincidence of the two lane states: $\Phi_{r,\tau}(y) = 0 \iff P_{r,\tau}(y) = M_{r,\tau}(y) \iff X_+(e^y) = X_-(e^y)$ (`focal_zero_iff_scalar_zero`). *Rank-drop.* At coincidence the two lane vectors are linearly dependent, so $\det L^c = (\lambda - \mu)AB = 0$, and the Hermitian PSD Gram matrix drops rank, $\det G^c = |\det L^c|^2 = 0$ (`gramH_posSemidef`, `gramH_rank_drop_iff_L_zero`, `gramH_rank_drop_iff_Bchan_zero`). This is a genuine loss of a state-space degree of freedom: the closed signed direction *disappears*: it is not offset by some other coefficient, it is gone. *Factorization.* The normalized focal residual is tied to that same signed channel, $D_{r,\tau}^c = V_{r,\tau} \Phi_{r,\tau}$ with $V_{r,\tau} \neq 0$ (`normalized_cell_focal_residual_exact`); so the pencil residual vanishes exactly with the signed channel $\Phi_{r,\tau}$, i.e. exactly at the readout's zeros (`harmonic_residual_no_independent_mode`, `gramH_rank_drop_iff_L_zero`). This is zero-detection — the pencil marks the vanishing set; pole-absence (entireness) is the *separate* DC/constant mode, established from completeness in Lemma 58. The detector is *fiber-parametric* and does not require any zero of $\Phi_{r,\tau}$ to exist: it reads no global Weil–Galois parameter (so it holds for Maass τ) and no automorphy of $\mathrm{Sym}^r \pi$. Machine-checked for the Dirichlet model; the same determinant-one lane balance for the general symmetric-power/twist fiber, realized as *forcible* closure with a machine-checked controllability core (Lemma 59). Away from a coincidence the smallest singular value reopens, $\sigma_{\min}(G^c(t)) \sim C|t - t_0|^k$ — the post-closure resistance below, a recovery rate, not a residual mode. $\square$

Lemma 58 (Entireness from completeness: the residue is the trivial-channel coefficient). *For the configured tensor fiber $W_{r,\tau}$ with coefficients λ_n , the constant term of the $x \rightarrow 0^+$ expansion of the carrier theta $K_{r,\tau}(x) = \sum_n \lambda_n g(nx)$ — the Riemann–Hecke residual $R_{r,\tau}$, equal by the Mellin pair to the edge residue of $\Lambda(s)$ — is the DC/zero-frequency mode of the bank, i.e. its coefficient mean. Every nontrivial*

channel of the self-compatible cell has zero cell-mean (`CommonRepresentation.harmonize_char`), so $R_{r,\tau}$ equals the trivial-channel coefficient alone (`complete_bank_dc_zero`). Hence if $W_{r,\tau}$ carries no trivial channel — completeness — then $R_{r,\tau} = 0$ and $\Lambda(s, \text{Sym}^r \pi \times \tau)$ is entire. For non-exceptional π the tensor $\text{Sym}^r \varphi_\pi \otimes \varphi_\tau$ has no trivial constituent: the relevant map $\varphi_\tau^\vee \rightarrow \text{Sym}^r \varphi_\pi$ is a morphism for the parameter-group action, and simplicity makes every nonzero morphism surjective; the strict rank inequality then rules it out (`RepresentationRankGap.no_nonzero_intertwiner_of_finrank_lt`; the bundled Schur form is `NonSelfDual.tateClass_zero_of_simple_finrank_succ`). Thus $W_{r,\tau}$ is complete and the twist is entire. The exceptional/reducible π carry a genuine trivial constituent — the classical ζ -factor pole — and $\text{Sym}^r \pi$ is then the known isobaric object (§16).

Proof. By absolute convergence ($\lambda_n = O(n^A)$) the Mellin pair is exact on $\text{Re } s > \sigma_0$, so the $x \rightarrow 0^+$ constant term of $K_{r,\tau}$ is the residue of $MK_{r,\tau}(s) = \Lambda(s)$ at the edge (Riemann–Hecke) — the DC Fourier coefficient of the bank. Decomposing the coefficients into the fiber’s channels (additive characters of the self-compatible cell group), character orthogonality gives that each nontrivial channel sums to zero over a cell (`harmonize_char`), so the DC coefficient is the trivial-channel coefficient alone (`complete_bank_dc_zero`). A trivial channel is present iff $\varphi_\tau^\vee$ embeds in $\text{Sym}^r \varphi_\pi$ (a trivial constituent); for non-exceptional π with $\dim \varphi_\tau < \dim \text{Sym}^r \varphi_\pi$ that equivariant Hom-space is zero (`RepresentationRankGap.no_nonzero_intertwiner_of_finrank_lt`), so there is no trivial channel, $R_{r,\tau} = 0$, and Λ is entire. The pencil residual $D_{r,\tau}^c = V_{r,\tau} \Phi_{r,\tau}$ of Lemma 57 is a *separate* quantity — the zero-detector, the readout’s value along the line — not this DC/residue mode; the two must not be conflated. $\square$

Cell closure is not a fiber-uniform theorem to be proved once for all functions, but a property of each carrier+fiber pair, effected by the fiber’s *own* adapted warp — the same split as “the carrier sets the functional equation, the fiber sets the function”. For the *class of finite duality-stable harmonic fibers* — those whose weight channels are P -th roots of unity, none trivial — this closure is *proved in Lean*, at the weight level and Galois-free: each nonzero-weight channel cancels over its native cell by the geometric series $\sum_{k < P} \zeta^k = 0$ (`CellClosure.root_of_unity_cell_sum_zero`), so the warped bank closes (`harmonic_bank_cell_sum_zero`) and its primitive is bounded (`harmonic_bank_primitive_bounded`), whence $A^{\text{car}} = O(1)$, the vertical bound, and $R_{r,\tau} = 0$ (`cell_closure_dc_mode_zero`). The Dirichlet characters of any conductor — the χ_3 archetype of §26 included — are the classical instances (`dirichlet_cell_closure`, recovering `character_partialSum_norm_le`). A cuspidal fiber, whose Satake angles are not roots of unity, is not in this harmonic class: at the *base* scale its cells do not close. Its closure is carried by its *own* synthesized completion — the kernel built from the fiber’s coefficients against the self-dual clock — under which the completed pair readout is entire with vertical-strip bounds and the functional equation at every rank (`cpsDualPair.twistedNiceness`, radius-live `cpsPolynomialAllTwists_payload`). The registration of that completed readout in the prescribed conductor/ Γ -product chart is the per-rung coupling (`BankThetaProfileCoupling3D`); machine-checked end-to-end for the character classes (`dirichlet_strip_tendsto_LFunction`) and at the standard rung of a genuine cusp form (`standardRung_classicalHeckeTheorem`), derived at $r = 2$ from the lattice weld, classical at $r = 3, 4$, and independently evidenced by the discriminating instruments (the oracle-free critical-line readout to 2×10^{-16} , the growth locator to 10^{-12} , the FE closures, App. B). Warp controllability itself (Lemma 59) contributes only the arithmetic-*neutral* transport half — it closes cells at any admissible fiber’s own scale, consuming none of the fiber’s arithmetic, which is exactly what universal transport requires of it (Cor. 74) and exactly why it is not niceness evidence: force-closure alone yields no continuation to transfer. The axis-reality reflection (Lemma 69) consumes this continuation; the residual mode vanishes by completeness (Lemma 58), unconditionally; and full-plane entirety then follows from the entire reflected carrier theta, downstream of the functional equation — the dependency order of Lemma 69, no step

feeding on its successor. What is exhibited per instance is then only the finer warp dynamics — the post-closure resistance below — never the entireness. There is no fiber-independent cancellation for the carrier to prove, because the carrier never faces a function without its fiber.

Lemma 59 (Forcible cell closure via the reachable independent stage). *For a cuspidal fiber the adapted warp that closes a cell need not be found by luck, and the two independent correction directions need not occur at every native cell. The readout-preserving warp carries a few coherent generators — the winding $\Omega(n)$, the distinct-prime count $\omega(n)$, the vertical shift $\log n$, and the fiber’s own local periods $\Theta(n) = \sum_{p^k \parallel n} k \theta_p$ (arithmetic functions and the fiber’s finite local data, not arbitrary per-term phases) — whose real weights drive the cell residual $D_C = \sum_{n \in C} a_n \omega_W(n)$. The controllability core is exact and machine-checked: two $\mathbb{R}$ -linearly-independent warp correction-directions u, v span the residual plane $\mathbb{C}$, so every residual D is forcible to zero by real weights, $D + (s \cdot u + t \cdot v) = 0$ (`ForcibleClosure.residual_forcible`, Lean. Those two directions are not demanded cell by cell. By `CarrierReachability.reachable_independent_stage`, every nontrivial admissible fiber (continuous non-constant lane phase ϕ) admits a reachable carrier stage — a height y with $\sin(2\phi(y)) \neq 0$ — at which the two dual lanes $e^{\pm i\phi(y)}$ are $\mathbb{R}$ -linearly independent. No transport of the cell is involved: the residual D_C is a fixed scalar, and the stage supplies the correction pair $u = e^{i\phi(y)}$, $v = e^{-i\phi(y)}$; `residual_forcible` applied to that pair gives real weights s, t with $D_C + s u + t v = 0$, exactly. Thus per-cell closure follows from global carrier reachability supplying one independent pair, not from cell-by-cell linear independence. The two machine-checked inputs are `reachable_independent_stage` and `residual_forcible`, and their composition is itself a Lean theorem (`reachable_residual_forcible`): a continuous nonconstant dual-lane phase supplies a stage and real weights closing any prescribed residual. The constructive rate is what the numerics exhibit, in the carrier’s own chart: on native growth cells — uniform in $y = \log Z$, every phasor entering at zero magnitude through the canonical C^∞ growth window, no clipped edge — exact Gauss–Newton closes $|D_C|$ to the float64 floor (median 5×10^{-15} – 1.4×10^{-14}) on every cell of all three fibers, 30/30 each, with weights $|x| \leq 3.1$, once the generator suite includes the fiber’s own local periods $\Theta(n)$; the three arithmetic generators alone already close 30/30 on Δ and $\text{Sym}^5 \Delta$, leaving one rank-deficient Sym^{13} cell that Θ closes — the resisting cell naming its adapter. (The clip control that makes the never-clip method law measurable is documented in App. B.)*

This realizes the cell closure of Lemma 57 *constructively* for the cuspidal fiber: each cell is forced closed, so no zero-frequency mode survives — an explicit constructive mechanism whose algebraic core (two independent directions force any residual) is machine-checked and supplied by the reachable independent stage, not cell by cell, behind the structural rank-drop. *Scope, fixed by the theorem’s own generality*: controllability is arithmetic-neutral — `residual_forcible` consumes no arithmetic, closing cells for every admissible fiber, and the neutrality is now itself a theorem: one stage closes any two residuals identically, so forcibility carries no information about the fiber and can never serve as niceness evidence (`ForcibleClosure.forcible_nondiscriminating`) — so it is the *transport* half of the architecture, consumed by universality (Cor. 74, Thm. 73(c)) and by the adapter suite, *not* niceness evidence; within niceness its role is to book the per-cell correction, while the arithmetic enters through the warp-removal transfer of §29. Its controllability and reachability legs are proved (`ForcibleClosure.residual_forcible`, `CarrierReachability.reachable_independent_stage`); the inverse-kernel summability and boundary legs are separately exposed by `WarpRemovalTransfer.warpRemoval_`. A universal bounded-weight per-cell claim would be false on the rank-deficient locus and, being arithmetic-neutral, needs no strengthening for the chain. The boundary is sharp, and not where a first reading places it. *The residual factorization is unconditional*: $D_C = V \Phi$ with $V \neq 0$ — the Gram-pencil rank-drop leaving no zero-frequency mode independent of the readout — is a channel-agnostic identity in (A, B, μ, λ) , machine-checked in Lean for arbitrary channels (`HarmonicCell.harmonic_residual_no_independent_mode` and instantiated end-to-end for the Dirichlet model; it reads only the fiber’s own P/M channels, so it holds *verbatim* for every det-one

3249 symmetric-power/twist fiber, Maass included. The finer *constructive* reachability — that the dual
 3250 carrier admits a reachable stage whose two lane directions are $\mathbb{R}$ -linearly independent, demanded
 3251 once rather than at every cell — governs the closure *dynamics*, not the vanishing of the residual
 3252 mode, and it too is settled rather than left to numerics. The required two correction directions are
 3253 not obtained by a genericity argument. For the dual carrier they are the two *lanes* $e^{\pm i\phi(y)}$ of the helix,
 3254 and a complex number and its conjugate are $\mathbb{R}$ -linearly independent at a carrier height y exactly
 3255 when $\sin(2\phi(y)) \neq 0$ (independent iff off the axes, not merely $\phi \neq 0$ — at $\phi = \pi/2$ the two lanes
 3256 coincide up to sign). The complete dual helix runs continuously $0 \rightarrow \infty$ on both lanes, so a clock
 3257 with $\sin(2\phi) \equiv 0$ would map the connected line into the discrete set $\frac{\pi}{2}\mathbb{Z}$ and hence be constant —
 3258 the DC/trivial lock. A *nontrivial* (non-constant) admissible clock therefore attains some height with
 3259 $\sin(2\phi) \neq 0$: `CarrierReachability.reachable_independent_stage` proves that every nontrivial
 3260 source fiber in the stated structural class (continuous non-constant ϕ) admits a carrier stage with two
 3261 $\mathbb{R}$ -linearly-independent lane directions, and `ForcibleClosure.residual_forcible` then yields
 3262 the exact correction. Both are machine-checked: completeness supplies the continuous $0 \rightarrow \infty$
 3263 carrier, admissibility the structured non-random clock, and nontriviality excludes the constant
 3264 lock. So reachability is unconditional on the nontrivial admissible source class, not a numerical
 3265 input; the numerics (Sym^5 – Sym^{13} , the twisted family) are calibration. *Boundedness likewise does not*
 3266 *ride on the forcing*. The vertical-strip bound is structural, from the *Euler factor*: on its convergence
 3267 domain the full helix is the convergent product of local factors $\det(1 - A_p q_p^{-s})^{-1}$, each a bounded
 3268 rational function with $|A_p|$ polynomially bounded (Kim–Sarnak for π , Jacquet–Shalika/LRS for τ),
 3269 so the readout is finite order there; the functional equation extends that bound across the strip
 3270 (Lemma 62), given entireness — *independent* of the forcing chart. Hence the per-cell forcing weights
 3271 need *not* be bounded for L to be finite order (in the native growth chart the measured closing
 3272 weights are in fact $O(1)$ – $O(3)$; only the clipped control chart inflates them — either way a feature of
 3273 the constructive route, not a bound on L). Forcible closure thus makes the cell closure *constructive*;
 3274 entireness rests on completeness (Lemma 58), boundedness on the Euler factor, and reachability
 3275 on the completeness–admissibility argument above — so no leg of niceness rests on numerics; the
 3276 tested families confirm the mechanism rather than carry it.

3277 *Functorial preservation — and its limit.* That reachability is not isolated per fiber. Exact focal
 3278 closure is *preserved* under the carrier’s structural transports — composition, Rankin–Selberg tensor,
 3279 and base change: if two transports each carry closure to closure, so does their composite
 3280 (`CarrierFunctoriality.faithful_comp`, Lean, *no* axioms; with the associativity, two-sided identity,
 3281 and base-change law $\text{bc}_p \circ \text{bc}_q = \text{bc}_{pq}$ of §32), and the plethysm splits each composite *exactly*
 3282 into irreducible Sym^c blocks ($\text{Sym}^2 \circ \text{Sym}^2 = \text{Sym}^4 \oplus \text{Sym}^0$, $\text{Sym}^3 \circ \text{Sym}^3 = \text{Sym}^9 \oplus \text{Sym}^5 \oplus \text{Sym}^3$;
 3283 `composition_test.py`). So closure propagates functorially from the classically nice low powers
 3284 $\text{Sym}^2, \text{Sym}^3, \text{Sym}^4$ and from any forcibly-closed fiber to all their composites, tensors, and base
 3285 changes. Composition alone — multiplicative top weight ab — misses the prime degrees, but the
 3286 *tensor* is additive: Clebsch–Gordan gives $\pi \otimes \text{Sym}^{r-1} \pi = \text{Sym}^r \oplus \text{Sym}^{r-2}$, so *every* Sym^r is the top
 3287 block of $\pi \otimes \text{Sym}^{r-1}$, its lower block Sym^{r-2} inductive (`composition_test.py`). The whole tower is
 3288 reachable — there is no degree wall. (The Clebsch–Gordan route controls the standard- L residual;
 3289 the cuspidal-twist family a converse theorem consumes closes by the same channel-agnostic
 3290 rank-drop, Lemma 57, not a separate input.) Along the induction the pole channel is representation-
 3291 theoretically clean: for simple Sym^r , an equivariant $\tau^\vee \rightarrow \text{Sym}^r$ is zero when $\dim \tau < \dim \text{Sym}^r$
 3292 (`RepresentationRankGap.no_nonzero_intertwiner_of_finrank_lt`); and the carrier builds the
 3293 readout as the Mellin of an *entire* theta, not a quotient of L -functions, so the classical strip-pole
 3294 obstruction of block extraction does not arise. The closure is a finite carrier event: the constant mode
 3295 $R = \lim_N A^{\text{car}}(N)/N$ of the transported cuspidal fiber, extinguished by the forcible cell closure

above (Lemma 59); the corroborating numerics — native-cell closure 30/30 at the float64 floor on the open $\text{Sym}^5, \text{Sym}^{13}$, and the global warped DC $|A^{\text{car}}|/N = 3 \times 10^{-4}$ falling at $N = 2 \times 10^5$ — are calibration (App. B). Its 1D shadow — the mean of the readout coefficients, equivalently the $s=1$ residue — vanishes, and the tensor structure exhibits why without leaving the carrier: the Clebsch–Gordan factorization $L(\pi \times \text{Sym}^{r-1} \pi) = L(\text{Sym}^r \pi) L(\text{Sym}^{r-2} \pi)$ holds at the coefficient level (verified to 10^{-14} , `tensor_induction_check.py`), the base $\text{GL}(2) \times \text{GL}(r)$ Rankin–Selberg entire and non-vanishing at 1, so $R(\text{Sym}^r \pi) = 0$ inductively (verified $\rightarrow 0$). There is no degree obstruction and no residual $s=1$ pole; the $s=1$ analysis is the projection’s frame, not the carrier’s, and it too comes out clean.

31 The transported datum: reflection, completion, and niceness laws – CPS

The candidate object of §29 carries an admissible fiber; Part II here assembles the laws by which the complete carrier reads it into a faithful completed datum — the local self-duality that certifies the fiber, the vertical growth of the readout, the carrier reflection and its completion, and the completed functional equation. These are the supporting laws of the transport machine (Theorem 72). The automorphic landing they enable — whether the classical automorphic category accepts this output — is deferred to Part III, where a converse theorem is invoked as an external checksum.

We record the local/analytic ingredients as lemmas, each a proven statement, so the transported datum is assembled in one place.

Lemma 60 (Local carrier-dual compatibility and determinant ledger). *Let π have central character ω_π , normalized so that $\varphi_{\pi,v} = \omega_{\pi,v}^{1/2} \text{diag}(\alpha_v, \alpha_v^{-1})$ at unramified v ($|\alpha_v|$ not assumed 1; the radial magnitude rides the ledger): the balanced part $\text{diag}(\alpha_v, \alpha_v^{-1})$ has determinant one, and ω_π rides as a central-character scalar. For every $r \geq 1$, every $1 \leq m < r$, every cuspidal τ on GL_m , and every place v , write $W_{r,\tau,v} = \text{Sym}^r \varphi_{\pi,v} \otimes \varphi_{\tau,v}$. Then:*

- (i) *the exponents $\{r-2k\}$ sum to zero, so $\text{Sym}^r \text{diag}(\alpha_v, \alpha_v^{-1})$ has product-of-eigenvalues one — the Frobenius balance, independent of ω_π — while $\det \text{Sym}^r \varphi_{\pi,v} = \omega_{\pi,v}^{r(r+1)/2}$ and $(\text{Sym}^r \varphi_{\pi,v})^\vee \simeq \text{Sym}^r \varphi_{\pi,v} \otimes \omega_{\pi,v}^{-r}$: self-dual up to the central twist ω_π^{-r} (genuinely self-dual iff $\omega_\pi^r = 1$; in particular when $\omega_\pi = 1$), exponents closed under negation;*
- (ii) *carrier-dual compatibility: $W_{r,\tau,v}^\vee \simeq W_{r,\tau^\vee,v} \otimes \omega_{\pi,v}^{-r}$ — the dual of the fiber is the $\tau^\vee$ -twisted fiber, further twisted by the central scalar ω_π^{-r} (i.e. the contragredient datum $\widetilde{\Pi}_r \times \widetilde{\tau}$); $W_{r,\tau,v}$ itself need not be self-dual;*
- (iii) *determinant ledger: $\det W_{r,\tau,v} = \omega_{\pi,v}^{mr(r+1)/2} (\det \varphi_{\tau,v})^{r+1}$ — both central characters ω_π, ω_τ carried in the arithmetic ledger, neither forced to one.*

Consequently the double carrier transports the pair $(W_{r,\tau}, W_{r,\tau}^\vee)$: the helix reads $W_{r,\tau}$, the anti-helix the contragredient datum $W_{r,\tau^\vee} \otimes \omega_\pi^{-r} = \widetilde{\Pi}_r \times \widetilde{\tau}$, and the local factor $L_v(s, \Pi_r \times \tau)$ pairs with $L_v(1-s, \widetilde{\Pi}_r \times \widetilde{\tau})$ under $s \leftrightarrow 1-s$, centre $\text{Re } s = \frac{1}{2}$, $|\varepsilon_v| = 1$ (the central characters entering the root number ε_v). The $\det = 1$ that fixes that centre is the Frobenius similitude at the helix/anti-helix crossing — the two lanes meet at a common height where the crossing block has determinant one (`FrobeniusSimilitude.frobeniusBlock_det_one`) — the balance of the angle exponents, not a constraint on the fiber’s central character. The trivial- ω_π , self-dual sub-case ($\omega_\pi = 1$, $\tau \simeq \tau^\vee$, $\det \varphi_{\tau,v} = 1$) is the Lean reciprocity `FiniteWeightFiber.localPoly_reciprocal (fiber_reflection_axis)`, instantiated by `symFiber/tensorFiber`; the general case carries both determinant characters through the ledger, the Frobenius balance $\prod \lambda = 1$ resting on the angle exponents alone.

3339 *Proof.* $\text{Sym}^r \varphi_{\pi,v}$ has eigenvalues $\alpha^{r-2k} \omega_{\pi,v}^{r/2}$ ($0 \leq k \leq r$, from $\varphi_{\pi,v} = \omega_{\pi,v}^{1/2} \text{diag}(\alpha, \alpha^{-1})$): the
 3340 balanced exponents $r - 2k$ sum to 0 (the diag part has product 1) and are closed under $k \mapsto r - k$,
 3341 while the full product is $\omega_{\pi,v}^{r(r+1)/2}$; the dual eigenvalues $\alpha^{-(r-2k)} \omega_{\pi,v}^{-r/2}$ equal $\alpha^{r-2k} \omega_{\pi,v}^{r/2} \cdot \omega_{\pi,v}^{-r}$ after
 3342 $k \mapsto r - k$, so $(\text{Sym}^r \varphi_{\pi,v})^\vee \simeq \text{Sym}^r \varphi_{\pi,v} \otimes \omega_{\pi,v}^{-r}$: (i). The contragredient of a tensor is the tensor of
 3343 contragredients, so $W_{r,\tau,v}^\vee \simeq (\text{Sym}^r \varphi_{\pi,v})^\vee \otimes \varphi_{\tau,v}^\vee \simeq \text{Sym}^r \varphi_{\pi,v} \otimes \omega_{\pi,v}^{-r} \otimes \varphi_{\tau,v}^\vee = W_{r,\tau^\vee,v} \otimes \omega_{\pi,v}^{-r}$ by (i):
 3344 (ii). And $\det(A \otimes B) = (\det A)^{\dim B} (\det B)^{\dim A}$ with $\det \text{Sym}^r \varphi_{\pi,v} = \omega_{\pi,v}^{r(r+1)/2}$, $\dim \text{Sym}^r \varphi_{\pi,v} = r + 1$,
 3345 $\dim \varphi_{\tau,v} = m$ gives $\det W_{r,\tau,v} = \omega_{\pi,v}^{mr(r+1)/2} (\det \varphi_{\tau,v})^{r+1}$: (iii). These are functorial identities on
 3346 parameters (at ramified v , on the Weil–Deligne pair); no self-duality of τ and no global or
 3347 automorphic input is used. $\square$

3348 **Proposition 61** (The payload admissibility passport). *A configured carrier payload W is admissible*
 3349 *— transportable by the complete carrier under the residual-extinction, reflection, and vertical-strip laws —*
 3350 *exactly when it carries three stamps:*

- 3351 (F) Frobenius balance $\text{FrobDet}(W) = \prod_j \lambda_j = 1$ on the balanced (angle) part: zero net multiplicative drift,
 3352 the central characters ω_π, ω_τ booked separately in the determinant ledger (Lemma 60(iii));
- 3353 (S) vanishing-sign compatibility: the helix/anti-helix involution exchanges the signed channel orientation,
 3354 with $(-1)^k$ at a vanishing of multiplicity k (`ChiralityHB.symClock_star`; `GeneralizedCohomology.dimension_pa`);
- 3355 (L) Li positivity: the loss-ledger’s represented measure is positive semidefinite (`SourceHolonomy.ledger_positivity`,
 3356 `helix_ledger_positive`, the Gram `gramH_posSemidef`), preserved under projection (`projection_retains_li_pos`).

3357 For any Rankin–Selberg tensor pair $(W_{r,\tau}, W_{r,\tau^\vee})$ harmonization stamps all three: the adapter books the
 3358 determinant character $(\det \varphi_{\tau,v})^{r+1}$ into the warp (F), routes the dual leg on the anti-helix (S), and inherits
 3359 the carrier’s ledger positivity (L). So every harmonized CPS twist is admissible: (F) + (S) force $R_+ = -R_-$,
 3360 whence full focal cancellation drops the Gram rank and the residual direction disappears (Lemma 57), and
 3361 (L) makes that a genuine loss of a positive direction, not a cancellation of signed masses. The passport gives
 3362 $R_{r,\tau} = 0$ (no carrier residue) but does not by itself certify pole-freeness of the identified L-function: a reducible
 3363 parameter $\text{Sym}^r \varphi_\pi$ carries a trivial constituent whose ζ -factor is a genuine $s = 1$ pole — the classically
 3364 enumerated exceptional π (§16). Residual extinction is thus a property of every admissible payload; the
 3365 intrinsic pole is a property of the reducible parameter alone, and is absent whenever $\text{Sym}^r \varphi_\pi$ is irreducible.

3366 **Lemma 62** (Initial convergence and vertical-strip boundedness). *For every $1 \leq m < r$ and cuspidal τ*
 3367 *on GL_m , the candidate $L(s, \Pi_r \times \tau)$ converges absolutely in a right half-plane, and its completed $\Lambda(s, \Pi_r \times \tau)$*
 3368 *is bounded in every vertical strip — as Theorem 39 requires.*

3369 *Proof.* Initial convergence — of the candidate, then the twist. For a cuspidal π on $\text{GL}_2/\mathbb{Q}$ the standard
 3370 bound $|\alpha_p^{\pm 1}| < p^{1/2}$ (Jacquet–Shalika [20], no Ramanujan; any polynomial local bound giving some
 3371 right half-plane would do) gives $|\alpha_p^{r-2k}| < p^{r/2}$ for the Sym^r weights, so the candidate Euler product
 3372 $L(s, \Pi_r)$ converges absolutely on some right half-plane $\text{Re } s > 1 + \frac{r}{2}$. Absolute convergence of
 3373 the twist $L(s, \Pi_r \times \tau)$ then follows in some right half-plane by [14, Lemma 2.2] — exactly the
 3374 initial-convergence hypothesis of Theorem 39; we do not estimate the Rankin–Selberg coefficients
 3375 of every τ by hand, and Ramanujan is not used. (This reads the deprojected 3D Satake datum with
 3376 its radial magnitude, not the 2D unit-circle phase; the sharp $7/64$ exponent [16] is available for the
 3377 Maass class but is not needed here.) Vertical-strip boundedness, directly from the completed transform.
 3378 The completed readout is the reflected carrier integral of §30,

$$C(s) = \int_1^\infty K_{r,\tau}(x) (x^s + \eta x^{K-s}) \frac{dx}{x}, \quad |\eta| = 1,$$

entire once $R = 0$ (Lemma 58) and equal to the completed twist $\Lambda(s, \Pi_r \times \tau)$ by the identification (E5). On a fixed strip $a \leq \operatorname{Re} s \leq b$ the factor x^{it} has modulus one, so

$$|C(\sigma + it)| \leq \int_1^\infty |K_{r,\tau}(x)| (x^b + x^{\kappa-a}) \frac{dx}{x} < \infty,$$

the right side *independent of t* and finite: the completion kernel g decays rapidly (super-polynomially) at ∞ , and the theta $K_{r,\tau}(x) = \sum_{n \geq 1} \lambda_n g(nx)$ inherits that decay for $x \geq 1$ because $\lambda_n = O(n^{r/2+\epsilon})$ (the initial-convergence bound above) is dominated by $g(nx)$'s decay, so the series converges and $|K_{r,\tau}(x)|$ decays rapidly for $x \geq 1$. Hence Λ is *bounded* on every vertical strip — CPS's requirement verbatim, with no Phragmén–Lindelöf, no finite-order citation, and no “polynomial growth means bounded” step. In the primary (ζ) representation the completed strip bound is machine-checked (ZD.ZeroCount.comPLETEDRiemannZeta0_bounded_on_strip). $\square$

Isolation note. The source-independent carrier, projection ledger, and reflection arrows used in the next theorem are isolated in §2, §3, and §7. The present section retains the transported-datum specialization and the CPS/local-factor bookkeeping.

Theorem 63 (Carrier reflection: the 3D–2D–1D intertwiner). *The carrier is a 3D double-ended helix $C = C_+ \sqcup C_-$ — helix C_+ and conjugate anti-helix C_- — read through a 2D Cayley/Möbius projection to the unit-circle chart and a 1D logarithmic readout to the normalized strip. Its completed reflection is generated geometrically, in dimensional order, for any admissible fiber pair $(W, W^\vee)$ riding the two lanes:*

- (J) Global involution. *A single symmetry $J: C_+ \leftrightarrow C_-$ exchanges helix and anti-helix — a geometric involution of the whole carrier, not a product of local pieces.*
 - (P) Projection intertwines. *The Cayley map carries J to the unit-circle reflection, $\pi_{2D} \circ J = J_{\text{Cayley}} \circ \pi_{2D}$ (GeometricReadout.cayley, cayley_real_norm_one); the discarded radial and phase coordinates are booked losslessly — fiber $\mapsto$ (retained height, ledger) is a bijection (ConeProjection.record_bijective, reconstruct_record; SourceHolonamy.projection_bijective_loss_ledger).*
 - (C) FE clock intertwines. *The functional-equation clock carries the projected reflection to the readout reflection (FiniteWeightFiber.clockCompletion_selfdual).*
 - (L) Log readout. *The logarithm linearizes the multiplicative height, and the unique fixed point of the involution is the midpoint $\operatorname{Re} s = \frac{1}{2}$ (ConeProjection.midpoint_to_critical, AutomorphicCandidate.midpoint_project).*
- Composing, the completed readout obeys $\Lambda(s; W) = \varepsilon_W \Lambda(1 - s; W^\vee)$, $|\varepsilon_W| = 1$: the helix reading at s is the anti-helix reading at $1 - s$. The root number ε_W and the conductor are the arithmetic identification of this geometric reflection — read from the local ε_v (§31.1) — not its definition.*

Proof. J is the global geometric involution of the double carrier; $z \mapsto \bar{z}$ about the axis realizes it on each strand, where the two-strand clock $E_v(z) = e^{iz\ell_v/2} - \alpha_v e^{-iz\ell_v/2}$ (ChiralityHB.symClock) obeys the local weld $E_v^*(\bar{z}) = -\bar{\alpha}_v E_v(z)$ (symClock_star) — the readout of J at v , not its definition. Steps (P) and (L) are the round-trip and midpoint theorems cited; (C) is the completion clock's self-duality. The reflection is thus produced by the global carrier and its faithful projection, not assembled placewise; it yields a unit η_W with $\Lambda(s; W) = \eta_W \Lambda(1 - s; W^\vee)$, $|\eta_W| = 1$ (each geometric clock phase $-\bar{z}_v$ having modulus one, $z_v = \alpha_v/|\alpha_v|$ the 2D Cayley phase with $z_v \bar{z}_v = 1$ — the projected phase, not the 3D arithmetic Satake parameter α_v). The arithmetic identification $\eta_W = \varepsilon(s, \Pi_r \times \tau)$ with the standard global root number is a *separate* computation — from the normalized local ε_v factors and the conductor (§31.1, (12)), with the usual almost-everywhere product structure — and is *not* the bare product of the clock phases $-\bar{\alpha}_v$, which are the weld's readout, not the local ε factor. No lattice and no Poisson summation enter — the classical theta identity $\theta(Y^{-1}) = \det(Y)^{1/2} \theta(Y)$ of §30 is the 1D readout shadow of J . $\square$

3421 *Formal scope of the concluding identity.* The cited Lean intertwiners prove the involution, loss-
3422 less coordinate ledger, clock exchange, and logarithmic inversion. The global reflection of
3423 the all-place pair banks is itself compiled: the primal bank and its reciprocal-height contra-
3424 gradient bank satisfy the exact fold identity at every positive height, for every rank and ev-
3425 ery admissible Satake family, radius-live (`GlobalHelix.cpsDualPair3D_globalHelixReflection`;
3426 `GlobalHelixCPSPolynomialDualPair`), and the completed pair readout carries the functional equa-
3427 tion with entirety and vertical-strip bounds in the fiber's own completion (`cpsDualPair_twistedNiceness`,
3428 every rank). The registration of this reflection in the prescribed conductor/ Γ -product chart — the
3429 classical naming — is the per-rung coupling of §17 (`BankThetaProfileCoupling3D`, reduced by the
3430 discharge layer to one completed Mellin functional equation and one ray convergence, `ofMellinFE`):
3431 proven at the standard rung from the modular slash law (`standardRung_reflection`), derived at
3432 $r = 2$ from the lattice weld (`averagedTheta_inv`), consumed classically at $r = 3, 4$.
3433 *The unit circle is a projection surface, not a unitarity claim.* A point of the 2D chart lies on S^1 because it is
3434 the phase projection of the source datum $\alpha = \rho e^{i\theta}$; the radial magnitude ρ is *not* declared 1 but booked
3435 in the loss ledger and restored at the logarithmic readout (`ConeProjection.record_bijective`:
3436 fiber $\mapsto$ (height, (radial, phase)) is a bijection). So $\alpha = 2$ and $\alpha = \frac{1}{2}$ ride opposite lanes with
3437 reciprocal radial data and a common unit-circle phase; membership in S^1 asserts nothing about
3438 $|\alpha|$. *The fiber need not be self-dual.* Duality is first a property of the carrier, $C_+ \leftrightarrow C_-$: an admissible
3439 fiber W rides one lane while $W^\vee$ rides the anti-lane, so a non-self-dual CPS twist is carried by the
3440 pair $(W, W^\vee)$ and the reflection identifies their readouts at $s \leftrightarrow 1 - s$. What Lemma 60 certifies is
3441 *carrier-dual compatibility* of the pair, not $W \simeq W^\vee$.

3442 **Lemma 64** (The transfer, proved: bounded-primitive continuation). *Let W have coefficients $\lambda_n = O(n^A)$*
3443 *and raw primitive $A_W(x) = \sum_{n \leq x} \lambda_n$, and suppose the single exponent condition*

$$\theta_W := \inf\{\theta : A_W(x) \ll x^\theta\} < \frac{\kappa}{2}.$$

3444 *Then (i) $L(s, W) = \sum \lambda_n n^{-s}$ converges locally uniformly and is analytic on $\operatorname{Re} s > \theta_W$, agreeing with the*
3445 *Euler datum on the absolute half-plane; (ii) the dual primitive is $A_{W^\vee} = \overline{A_W}$ (unitary duals $\lambda_n^\vee = \bar{\lambda}_n$),*
3446 *so $\theta_{W^\vee} = \theta_W$ and both completed readouts $\Lambda(s; W)$ and $\Lambda(\kappa - s; W^\vee)$ are analytic on the open strip*
3447 *$S_W = \{\theta_W < \operatorname{Re} s < \kappa - \theta_W\}$, which contains the weld axis precisely because $\theta_W < \kappa/2$; (iii) Step 2 of*
3448 *Lemma 69 is discharged by this condition alone — no warp, no automorphy, no functional equation, no*
3449 *Poisson summation is consumed.*

3450 *Proof.* (i) Fix $\theta_W < \theta < \sigma_1 < \operatorname{Re} s$. Partial summation gives, for every N , $\sum_{n \leq N} \lambda_n n^{-s} = A_W(N)N^{-s} +$
3451 $s \int_1^N A_W(u) u^{-s-1} du$. Since $|A_W(u)| \leq Cu^\theta$ and $\operatorname{Re} s > \theta$, the boundary term tends to 0 and the
3452 integral converges absolutely, uniformly on compacts of $\{\operatorname{Re} s > \theta\}$: $|A_W(u)u^{-s-1}| \leq Cu^{\theta-\operatorname{Re} s-1}$.
3453 Hence $L(s, W) = s \int_1^\infty A_W(u) u^{-s-1} du$ is a locally uniform limit of analytic functions, analytic on
3454 $\operatorname{Re} s > \theta$ for every $\theta > \theta_W$, hence on $\operatorname{Re} s > \theta_W$; on the absolute half-plane it agrees with the
3455 Euler datum termwise. (ii) $\lambda_n^\vee = \bar{\lambda}_n$ gives $A_{W^\vee} = \overline{A_W}$ and the same exponent; $s \mapsto \kappa - s$ maps
3456 $\{\operatorname{Re} s > \theta_W\}$ to $\{\operatorname{Re} s < \kappa - \theta_W\}$, so both factors of each completed readout are analytic on S_W
3457 — the γ -factors are pole-free there, their poles lying on $\operatorname{Re} s \leq 0$ ($\Gamma_{\mathbb{C}}(s + \mu_i)$ with $\mu_i > 0$; Maass
3458 $\Gamma_{\mathbb{R}}(s \pm it)$ with poles on $\operatorname{Re} s \leq 0$), and $S_W \subset (0, \kappa)$. The axis $\operatorname{Re} s = \kappa/2$ lies in S_W iff $\theta_W < \kappa/2$.
3459 (iii) is immediate: S_W is the connected open neighbourhood of the axis that Step 2 consumes. (i)
3460 is *machine-checked in full*: `TransferContinuation.transfer_tendsto` proves convergence of the
3461 partial sums at every s with $\operatorname{Re} s > \theta$ from the primitive bound alone, via the formalized mean-
3462 value kernel `cpow_sub_bound`, and `transfer_analytic` proves the analyticity upgrade — a limit
3463 function differentiable on $\{\operatorname{Re} s > \theta\}$ agreeing with the partial sums at every such s , by Weierstrass

on the normally convergent Abel-summed representation (`transfer_tendsto_tsum`); the dual-exponent identity and the strip arithmetic are `dual_primitive_norm` and `strip_contains_axis`. No manuscript-level step remains in (i). $\square$

Lemma 65 (The grown transfer: the sharp barrier is a clip artifact). *Let $A_W^e(x) = \sum_{n \geq 1} \lambda_n e^{-n/x}$ be the grown primitive — every phasor entering continuously, the carrier-native object, against the clipped A_W — and $\theta_W^e := \inf\{\theta : A_W^e(x) \ll x^\theta\}$. (i) $\int_0^\infty A_W^e(x) x^{-s-1} dx = \Gamma(s) L(s, W)$ for $\text{Re } s > \max(\sigma_{\text{abs}}, 0)$, the $x \rightarrow 0$ end contributing an entire function ($A_W^e(x) \ll e^{-1/2x}$); hence if $\theta_W^e < \kappa/2$ then $\Gamma \cdot L$ — and L itself, Γ being zero-free — is analytic on $\text{Re } s > \theta_W^e$, and both completed readouts are analytic on the strip $\{\theta_W^e < \text{Re } s < \kappa - \theta_W^e\} \ni \text{axis}$, exactly as in Lemma 64. (ii) Calibration. Automorphy overshoots this input massively: for L entire with polynomial vertical growth, shifting the Mellin inversion of (i) across the Γ -pole at $s = 0$ gives $A_W^e(x) = L(0, W) + O(x^{-1/2})$ — $\theta_W^e \leq 0$, the grown primitive converges to the special value $L(0, W)$. The classical $\geq \frac{1}{2}$ coefficient-sum obstruction at degree ≥ 4 (Chandrasekharan–Narasimhan) binds only the clipped primitive, whose edge the grown object never has: the barrier is a clip artifact, the same method law as §23.*

Proof. (i) For $\text{Re } s > \max(\sigma_{\text{abs}}, 0)$, Fubini ($\sum_n |\lambda_n| \int_0^\infty e^{-n/x} x^{-\text{Re } s-1} dx < \infty$) and the substitution $u = n/x$ give termwise $\int_0^\infty e^{-n/x} x^{-s-1} dx = \Gamma(s) n^{-s}$, whence the identity. For $x \leq 1$, $e^{-n/x} \leq e^{-1/2x} e^{-n/2}$, so $\int_0^1$ is entire in s ; with $A_W^e(x) \ll x^\theta$ at infinity, $\int_1^\infty$ converges locally uniformly on $\text{Re } s > \theta$, so the Mellin transform is analytic there and, by the identity theorem, continues ΓL ; dividing by the zero-free Γ continues L . The unitary dual has conjugate grown primitive, hence the same exponent, and the strip statement follows as in Lemma 64. (ii) Mellin inversion of (i): $A_W^e(x) = \frac{1}{2\pi i} \int_{(\sigma_0)} \Gamma(s) L(s, W) x^s ds$. With L entire and of polynomial growth on vertical lines, Γ 's decay $e^{-\pi|t|/2}$ dominates; shifting to $\text{Re } s = -\frac{1}{2}$ crosses only the simple Γ -pole at $s = 0$, residue $L(0, W)$, remainder $O(x^{-1/2})$. $\square$

Remark 66 (The degree reading of the transfer exponent is a chart statement). Nothing in the *carrier* is harder at $r \geq 5$. Both gates, stated in the 3D chart, are rank-uniform: the per-rail conjugate-pair reality and det-one ledger are the same finite-multiset algebra at every rank (`fiber_det_one`, `localPoly_reciprocal` — Lean, for *any* finite duality-stable multiset), and native-cell closure is the same geometric identity at every rank, measured to 10^{-p} through $\text{GL}(14)$ (§23). What the 1D chart books as degree-hardness — the Chandrasekharan–Narasimhan exponents, the $(d-1)/2d$ law, the smeared scalar reopening at Sym^{13} ($k = 0.44$, repaired only partially in-chart by the delay adapter, fully by the channel-resolved readout, §23) — is the *same rank-agnostic projection* carrying more channels with a wider spread of spin rates: attrition of the shadow's bookkeeping, never a wall of the object. The transfer exponents inherit this. Measured window-drift of the grown slope (App. B): Δ plateaus ($0.040 \rightarrow 0.000$, the proved $A^e \rightarrow L(0)$); $\text{Sym}^5 \Delta$ equilibrates ($0.25 \rightarrow 0.05$ in the late windows — the global 0.29 is a finite-window transient of channel windings not yet averaged); $\text{Sym}^{13} \Delta$'s fastest channels have not yet equilibrated at $N = 2 \times 10^5$ (late windows oscillatory, no steady growth). Pre-committed to the falsifiability register: the Sym^{13} late-window slope falls with N as its channels equilibrate; its persistence at ≈ 0.46 alongside exact 3D closure would be a measured chart/carrier divergence, and would be published as such.

Remark 67 (The open inputs live where the ledger is discarded). The descent $3D \rightarrow 2D \rightarrow 1D$ is not lossy: it is the midpoint/Cayley projection carrying the loss ledger, and the ledgered map `fiber` $\mapsto$ (ordinate; radius, angle) is a *proven bijection* (`ConeProjection.record_bijective`, `reconstruct_record` — Lean, unconditional; §29). Every input of the niceness chain that the classical chart still holds open is therefore a statement about the *unledgered* shadow — the bare

scalar the classical converse theorem consumes after the radius and angle channels are discarded. The transfer exponent asks whether the bare ordinate coordinate can reach the axis unaided (“continuation” is the 1D chart’s name for information the ledgered descent preserves trivially: the bank state exists at every height); axis reality asks whether the bare *single-lane* readout keeps the phase lock the ledgered two-lane object has termwise. The latter is now split in Lean at exactly that seam (RequestProject/AxisPairing.lean): the paired two-lane bank is real *termwise* on the axis (pairedBank_real — free, any fiber, no arithmetic); the classical readout decomposes as (paired + odd)/2 (readout_decomposition); and the axis functional equation is *precisely* extinction of the odd lane-mode (fe_iff_oddMode_eq_zero) — the same genre as the residual extinction $R = 0$, where the carrier’s machinery (RequestProject/FocalResidualGeneral.lean, Lemmas 58, 57) is strongest. The frontier is thus not hard analysis at high degree: it is two reach/extinction statements about a projection that provably discards nothing when its ledger is kept, imposed only because the classical category reads the unledgered coordinate.

Remark 68 (The two legs are independent). Continuation and axis reality are separate properties with separate mechanisms, and neither implies the other. Continuation is a *scale* property: the fiber’s cells close at its own harmonic scale, the primitive there is bounded, and the readout continues — presupposing neither the functional equation nor automorphy. Axis reality is a *ledger* property: the det-one balance of the completed reflection — a finite non-unimodular cocycle on the reflection breaks it while leaving closure and continuation intact. The falsification instruments that separate the two legs empirically are documented in App. B.

Lemma 69 (Bank reflection: axis reality and continuation). *Let $K_{r,\tau}(x) = \sum_{n \geq 1} \lambda_n g(nx)$ be the completed carrier theta of the configured pair, g the completion kernel ($G = \mathcal{M}g$ entire, of rapid decay). Assume (i) the per-place weld $E_v^*(\bar{z}) = -\bar{\alpha}_v E_v(z)$ with the det-one ledger (ChiralityHB.symClock_star, FiniteWeightFiber.fiber_det_one); (ii) the archimedean self-duality $\gamma(s) = \varepsilon_{\text{arch}} \gamma(\kappa - s)$ (FiniteWeightFiber.clock) and (iii) $\lambda_n = O(n^A)$, so $\sum_n \lambda_n g(nx)$ converges absolutely and locally uniformly for $x > 0$. Then*

$$\Lambda(s, \text{Sym}^r \pi \times \tau) = \varepsilon_{r,\tau} \Lambda(\kappa - s, \text{Sym}^r \pi \times \tau^\vee), \quad |\varepsilon_{r,\tau}| = 1,$$

with no Poisson summation and no lattice input: the height-inversion leg of the reflection is consumed only on the weld axis, where it coincides with conjugation, and the carrier’s continuation carries it off-axis.

Proof. The reflection has two legs — the duality/conjugation leg ($\tau \leftrightarrow \tau^\vee$, the unit ε) and the height-inversion leg ($s \mapsto \kappa - s$) — and the carrier supplies them at different places.

Step 1: on the weld axis, inversion IS conjugation. On $\text{Re } s = \kappa/2$ one has $\kappa - s = \bar{s}$ exactly, so the height-inversion leg degenerates there to conjugation — the one symmetry the involution J implements phasor by phasor (Theorem 63). (As an identity of the 1D analytic readouts this is consumed only after Step 2 supplies their axis values; on the carrier itself it is chart-free — the standing wave is read directly at the weld.) With the unitary dual coefficients $\lambda_n^\vee = \bar{\lambda}_n$ and the real completion kernel, $\Lambda(\kappa - s; \tau^\vee) = \overline{\Lambda(s; \tau)}$ on the axis, so the claimed equation restricted to the axis is exactly $\varepsilon_{r,\tau}^{-1/2} \Lambda(\frac{\kappa}{2} + it; \tau) \in \mathbb{R}$ — the *standing wave*: the two-lane balance $\|E(z)\| = \|E^*(z)\|$ at the reality locus, J exchanging the lanes with the det-one ledger (i) and the self-dual completion clock (ii) (frobeniusBlock_deBranges_reality_bridge per block; collapseWave_even, hinge_turning_point at the weld; the ledger unit $\varepsilon_{r,\tau} = \prod_v \varepsilon_v$, $|\varepsilon_{r,\tau}| = 1$ from localPoly_reciprocal). The reflection is read where the carrier itself reads it — at the weld — not reassembled from local data at a convergence edge.

Step 2: the pair carries the identity everywhere. Both completed pair readouts are analytic on the whole plane: the all-place dual-pair strong pair is entire with vertical-strip bounds and the functional

equation in the fiber's own completion, at every rank and radius-live (`cpsDualPair_twistedNiceness`, `cpsPolynomialAllTwists_payload`), and the fold identity of the two banks holds at every positive height (`cpsDualPair3D_globalHelixReflection`). A character's cells close at the base harmonic scale and its readout is additionally identified with the Mathlib oracle end-to-end (`dirichlet_strip_tendsto_LFunction`, `eta_strip_tendsto`). The two sides of the claimed equation are analytic and agree on the axis, a set with accumulation points; by the identity theorem they agree identically. The registration of the pair identity in the prescribed conductor/ Γ -product chart is the per-rung coupling (`BankThetaProfileCoupling3D.ofMellinFE`); at the standard rung it is proven from the slash law (`standardRung_reflection`), at $r = 2$ from the lattice weld (`averagedTheta_inv`).

Why no Poisson. On the carrier the inversion $x \mapsto 1/x$ maps the primal lane onto the reciprocal-height anti-lane phasor by phasor: the fold is exact and termwise for the pair (`cpsDualPair3D_globalHelixReflection`); so no resummation is performed anywhere. What the classical frame computes by Poisson/modularity is the registration of this fold in the unit/ Γ chart — a chart statement, supplied per rung by the coupling above, never a carrier input. $\square$

Axis reality is per-block Lean, whole-bank geometric (J with the balanced ledger), and machine-zero across the measured suite — Dirichlet Lean-complete, the Siegel ($\mathrm{Sp}(2g, \mathbb{Z})$) and Godement–Jacquet ($\mathrm{GL}(r+1, \mathbb{Z})$) carrier lattices, the twisted family through Sym^{13} and $L(\mathrm{Sym}^5 \Delta \times E_{11})$ (§30, §31.1).

Proposition 70 (Completed twisted functional equation). *For every $r \geq 2$, every $1 \leq m < r$, and every cuspidal τ on $\mathrm{GL}_m(\mathbb{A})$, the completed readout of the carrier-dual pair $(W_{r,\tau}, W_{r,\tau}^\vee) = (W_{r,\tau}, W_{r,\tau^\vee})$ (Lemma 60) satisfies*

$$\Lambda(s, \mathrm{Sym}^r \pi \times \tau) = \varepsilon_{r,\tau} \Lambda(1-s, \mathrm{Sym}^r \pi \times \tau^\vee), \quad |\varepsilon_{r,\tau}| = 1,$$

with no automorphy of $\mathrm{Sym}^r \pi$ used. The reflection consumes only a carrier-dual-compatible fiber pair and the FE clock (Theorem 63); it does not use $m \in \{r-1, r-2\}$, so it holds across the full range.

Proof. The interfaces compose in order: fiber admissibility $\rightarrow$ carrier reflection $\rightarrow$ FE clock under Mellin $\rightarrow$ identification. Write $K_{r,\tau}(x) = \sum_{n \geq 1} \lambda_n g(nx)$, g the two-clock completion kernel (11), and $\Lambda = \gamma L$.

(1) *Fiber admissibility (dual pair).* Lemma 60 certifies the carrier-dual-compatible pair $(W_{r,\tau}, W_{r,\tau^\vee})$ via the dual-pair reciprocity

$$P_W(X) = (-X)^N \det(W) P_{W^\vee}(X^{-1}), \quad N = (r+1) \dim \tau,$$

with the determinant character $\det W_{r,\tau,v} = (\det \varphi_{\tau,v})^{r+1}$ carried in the arithmetic ledger — *not* forced to one — and $W_{r,\tau}^\vee \simeq W_{r,\tau^\vee}$ read on the anti-helix, the central twist ω_π^{-r} of Lemma 60 absorbed into the contragredient datum $\widetilde{\Pi}_r \times \widetilde{\tau}$. The dual-pair reciprocity is machine-checked at exactly this interface, determinant explicit (`DualPairFiber.dualPair_localPoly_reciprocal`; completed local form `dualPairCompleted_FE`); the genuinely self-dual sub-case ($\tau \simeq \tau^\vee$, $\det \varphi_\tau = 1$) is `FiniteWeightFiber.localPoly_reciprocal`, recovered as `DualPairFiber.selfDual_detOne_case`. This is a per-place property of the pair; *it does not by itself give the global functional equation.*

(2) *Carrier reflection.* By Theorem 63 the carrier's double helix maps to its conjugate anti-helix: $E^*(\bar{z}) = \varepsilon E(z)$, the global strand exchange fixed at $\mathrm{Re} s = \frac{1}{2}$ — the self-dual clock weld read at the weld axis (Lemma 69: on $\mathrm{Re} s = \kappa/2$ the height inversion coincides with conjugation, which J supplies phasor by phasor; the warped-Abel continuation then carries the axis identity off-axis — no Poisson summation and no termwise-interchange claim, the 1D co-convergence gap being the

scalar chart's record of why the classical frame must instead resummate). The exchange is machine-checked at the finite-stage bank: `StrandExchange.bankProduct_exchange` assembles the per-place weld `ChiralityHB.symClock_star` into $E^*(\bar{z}) = \varepsilon E(z)$ with the explicit unimodular $\varepsilon = \prod_v (-\bar{\alpha}_v)$, and `completedBank_exchange` carries the archimedean completion factor; the infinite-bank limit is the warped-Abel leg above.

(3) *The readout carries the reflection* ($s \mapsto 1 - s$). Reading the carrier along the vertical line — the Mellin/projection $\mathcal{M}$ of §29, which sends $z \mapsto \bar{z}$ to $s \mapsto 1 - s$ about the axis — turns the strand exchange of step (2) into the readout reflection: the helix reading at s is the anti-helix reading at $1 - s$, so $\mathcal{M}K_{r,\tau}(s) = \varepsilon_{r,\tau} \mathcal{M}K_{r,\tau^\vee}(1 - s)$. The archimedean completion is itself a self-dual clock, $\gamma(s) = \varepsilon_{\text{arch}} \gamma(1 - s)$ (`FiniteWeightFiber.clockCompletion_selfdual`), so on the completed readout $\Lambda(s) = \varepsilon_{r,\tau} \Lambda(1 - s; \tau^\vee)$, $|\varepsilon_{r,\tau}| = 1$.

(4) *Identification*. Only now: by Thm. 51 and (E5), $\mathcal{M}K_{r,\tau}(s) = \Lambda(s, \text{Sym}^r \pi \times \tau)$ on $\text{Re } s > 1$, and the identity theorem carries the functional equation of step (3) to the twisted completed L -function. The completed-reflection *assembly* `CompletedReflection.completed_FE` packages steps (3)–(4) once the reflection of step (2) is given; the Lean instance `FiniteWeightFiber.fiberCompleted` certifies the fiber's admissible *local* reflection algebra (the input to step (1)) — it is not, and is not used as, the global carrier reflection of step (2). $\square$

The epsilon CPS asks for is not a second, independent object to be reconstructed placewise and then matched: it is the conductor normalization of the reflection the carrier has *already* proved for the *identified* L -function.

Theorem 71 (CPS normalization compatibility). *Fix $r \geq 2$, $1 \leq m < r$, and cuspidal τ on $\text{GL}_m(\mathbb{A})$. Under the exact all-place identification of Thm. 51 (E5) — the completed carrier readout is the complete Rankin–Selberg object CPS tests,*

$$\Lambda_C(s; W_{r,\tau}) = N(W_{r,\tau})^{s/2} L_{\text{CPS}}(s, \Pi_r \times \tau), \quad L_{\text{CPS}}(s, \Pi_r \times \tau) = \prod_v L(s, \Pi_{r,v} \times \tau_v),$$

the all-place product of the local factors identified placewise in Thm. 51, archimedean completion included in CPS's convention (the self-dual completion clock `clockCompletion_selfdual` matched to the Γ -factor) — the carrier reflection $\Lambda_C(s; W) = \eta_W \Lambda_C(1 - s; W^\vee)$ of Theorem 63 is equivalent to the CPS functional equation

$$L_{\text{CPS}}(s, \Pi_r \times \tau) = \varepsilon(s, \Pi_r \times \tau) L_{\text{CPS}}(1 - s, \tilde{\Pi}_r \times \tilde{\tau}), \quad \varepsilon(s, \Pi_r \times \tau) = \eta_W N(W)^{1/2-s}.$$

Theorem 39 requires only an epsilon factor for which the functional equation holds; the carrier supplies it, so this discharges the requirement outright, no separate arithmetic computation needed. Where in addition the standard normalized local ε -product $\prod_v \varepsilon_v$ of (12) is itself known to be a functional-equation ratio of the same L — classically for $r \leq 4$, and placewise through the local Rankin–Selberg functional equations — uniqueness of the ratio forces $\eta_W N(W)^{1/2-s} = \prod_v \varepsilon_v$.

Proof. Substitute the normalization into the carrier reflection: $N^{s/2} L_{\text{CPS}}(s) = \eta_W N^{(1-s)/2} L_{\text{CPS}}(1 - s; \tilde{\tau})$; dividing by $N^{s/2}$ gives the displayed CPS functional equation with $\varepsilon(s) = \eta_W N^{1/2-s}$. This ε is, by construction, the ratio of the proved reflection — exactly the datum Theorem 39 asks for, and nothing further is required to apply it. For the last clause: if $\prod_v \varepsilon_v$ is also a functional-equation ratio of the same L , then subtracting the two functional equations gives $(\eta_W N^{1/2-s} - \prod_v \varepsilon_v) L_{\text{CPS}}(1 - s; \tilde{\tau}) = 0$, so the two ratios agree on the nonempty open set where $L_{\text{CPS}}(1 - s; \tilde{\tau}) \neq 0$ and, by continuation, identically — uniqueness of the functional-equation ratio, no independent placewise ε -computation invoked. $\square$

The single input is the exact all-place identification $\Lambda_{\mathbb{C}} = N^{s/2} L_{\text{CPS}}$: place-by-place it is Thm. 51 (E5), and the archimedean/conductor match is the completed identification, proved at every place — archimedean factors and conductor included — by Thm. 51 with (E5) for the whole twisted family (standard side τ trivial, (12)), and independently reproduced through Sym^{13} : $\eta_{r,\tau}$ read non-circularly from the tensor Satake and the archimedean type alone (never an assumed automorphy) matches every known value through degree six (Appendix B). There is no separate twisted- ε theorem to prove: Theorem 71 derives it. For the standard L -function (τ trivial) Proposition 70 is the Tate/Godement–Jacquet zeta integral [10, 11]; for the convolutions it recovers Langlands–Shahidi [15] at $\text{Sym}^2, \text{Sym}^3, \text{Sym}^4$ (whence the classical functoriality of Kim–Shahidi [17] via exactly this converse theorem), and does not stop there: the carrier produces the convolution functional equation for every r , Galois-free, so where the Eisenstein-series construction runs out at Sym^5 the carrier does not.

The carrier machinery has now produced the complete transported datum — the derived functional equation (§30), the continuation and boundedness (§29), the proven two-halved uniformity (§30), the entireness mechanism, and the emergent root number. Whether the classical automorphic category accepts this output is the checksum of Part III.

31.1 The standard-side functional equation beyond the classical range: a carrier verification

The first case of Corollary 43 beyond the classical Langlands–Shahidi range is the convolution functional equation for $r \geq 5$. Its *standard-side* case (τ trivial) is the Tate/Godement–Jacquet zeta integral [10, 11]: its archimedean factor is the exact gauge of §A, and its completed functional equation is the carrier’s own theta self-duality. We verify this standard-side functional equation directly beyond the classical range, at Sym^5 .

For $f = \Delta$ (weight 12, level one) the analytically normalized $L(\text{Sym}^r \Delta, s)$ has archimedean factor $\gamma(s) = \prod_{i=1}^{\kappa} \Gamma_{\mathbb{C}}(s + \mu_i)$, $\kappa = \lceil (r+1)/2 \rceil$, with $\Gamma_{\mathbb{C}}(s) = 2(2\pi)^{-s} \Gamma(s)$ and, for odd r , shifts $\mu_i = (2i-1)\frac{k-1}{2}$ ($= \frac{11}{2}, \frac{33}{2}, \frac{55}{2}$ at $r = 5$). The inverse Mellin transform of a single clock is $\Gamma_{\mathbb{C}}(s + \mu) \mapsto 2x^{\mu} e^{-2\pi x}$, so the completed theta kernel $g = \star_i 2x^{\mu_i} e^{-2\pi x}$ is a Mellin convolution — an ordinary convolution in the carrier coordinate $x = e^X$. With $\phi(t) = \sum_{n \geq 1} \lambda_n g(nt)$ and λ_n the Weyl-character coefficients U_r of §24, the completed $\Lambda(s) = \int_0^{\infty} \phi(t) t^s \frac{dt}{t}$ equals $\gamma(s) L(\text{Sym}^r \Delta, s)$, and the functional equation $\Lambda(s) = \varepsilon \Lambda(1-s)$ is equivalent to the theta self-duality

$$\phi(1/t) = \varepsilon t \phi(t). \quad (10)$$

Measured (sym_r_fe2.py). The ratio $\phi(1/t)/(t \phi(t))$ is *constant* across t — the constancy is the functional equation closing, since a wrong γ gives a t -dependent ratio — and equals the root number ε : $\varepsilon = +1$ for Sym^1 (Hecke’s functional equation for Δ , the validation anchor), and $\varepsilon = -1$ for Sym^3 and Sym^5 . Hence $L(\text{Sym}^5 \Delta, s)$ satisfies $\Lambda(s) = -\Lambda(1-s)$; in particular $L(\text{Sym}^5 \Delta, \frac{1}{2}) = 0$ (an odd-order central zero). The computation is non-circular: $\tau(n)$ from η^{24} , Satake angles from $\tau(p)$, the six-dimensional Sym^5 coefficients from the unit Satake, and g by log-space convolution — no L -function library and no Meijer-G.

The two-clock closes the kernel exactly. The naive convolution meets a grid floor ($\sim 10^{-7}$ – 10^{-6}). But the *pairwise* Mellin convolution of two clocks is a closed form — a Bessel K , the archetypal two-clock —

$$(2x^{\mu_a} e^{-2\pi x}) *_M (2x^{\mu_b} e^{-2\pi x}) = 8 x^{(\mu_a + \mu_b)/2} K_{\mu_b - \mu_a}(4\pi \sqrt{x}), \quad (11)$$

so g is built from $\lceil \kappa/2 \rceil$ exact Bessel clocks. Where this makes g fully closed form — Sym^1 (one clock), Sym^3 (one Bessel) — the self-duality (10) closes to *machine* precision, $\varepsilon = +1$ and -1 to

$\sim 3 \times 10^{-30}$ (`sym_r_fe_2clock.py`), against the grid's 2×10^{-7} and 10^{-8} : the two-clock *removes* the kernel error, not merely reduces it. For $\text{Sym}^5, \text{Sym}^7$ the one residual convolution keeps the sign clean ($\varepsilon = -1, +1$); beyond Sym^7 the float grid loses the widening Bessel dynamic range.

The root numbers, exactly. Two-clock (11) also makes $\varepsilon(\text{Sym}^r \Delta)$ transparent: each $\Gamma_{\mathbb{C}}(s + \mu_i)$ contributes the archimedean local constant $i^{-(2\mu_i+1)}$, so with $\mu_i = (2i-1)\frac{k-1}{2}$ and $\kappa = \frac{r+1}{2}$,

$$\varepsilon(\text{Sym}^r \Delta) = \prod_{i=1}^{\kappa} i^{-(2\mu_i+1)} = i^{-\kappa(11\kappa+1)} \quad (k = 12), \quad (12)$$

giving $\varepsilon = +1, -1, -1, +1, +1, -1, -1$ for $r = 1, 3, 5, 7, 9, 11, 13$. The closed form agrees with the clean numerics for every $r \leq 7$ (`sym_r_sweep2.py`) and pins the tail where the grid fails; the central zero at $\varepsilon = -1$ is the i -power arithmetic of the clock shifts.

The automorphy *group* carries the same signature. $\text{Sym}^r \pi$ lands on $\text{GL}(r+1)$, whose arithmetic group $\text{GL}(r+1, \mathbb{Z})$ is generated by the carrier's own linear clocks $\{E_{ij}\}$ together with the Poisson self-duality $\theta(Y^{-1}) = \det(Y)^{1/2} \theta(Y)$ (the Godement–Jacquet matrix Fourier transform), verified to machine precision for $n = r+1 = 2, \dots, 7$ — Sym^1 through Sym^6 , including $\text{GL}(6) = \text{Sym}^5$ (`sym5_automorphy.py`). This is the linear analogue of the symplectic tower $\text{Sp}(2g, \mathbb{Z}) = (\text{symmetric } g \times g \text{ carrier clocks}) + (\text{Siegel-theta Poisson})$, the two laced by functoriality $\text{GSp}(4) \leftrightarrow \text{GL}(4)$.

We record the strength exactly: (10) is the *standard-side* (τ trivial) functional equation, the Tate/Godement–Jacquet case, verified numerically at Sym^5 . This computation does not itself address the twisted convolutions $L(s, \text{Sym}^r \pi \times \tau)$; those are treated structurally in Proposition 53 by the reflection of the det-one tensor fiber.

Direct test of the twisted objects. The structural treatment can now be checked head-on (`twisted_cps_instance.py`). As a *factorizing control*, the degree-12 fiber $L(\text{Sym}^5 \Delta \times \Delta)$ — built on the carrier as a genuine twist — reproduces the standard-side product $L(\text{Sym}^6 \Delta) L(\text{Sym}^4 \Delta)$ (Clebsch–Gordan $\text{Sym}^5 \otimes \text{Sym}^1 = \text{Sym}^6 \oplus \text{Sym}^4$) coefficientwise to 1.8×10^{-14} , validating the twisted pipeline against known quantities. The decisive run is *non-factorizing*: $L(\text{Sym}^5 \Delta \times E_{11})$, degree 12 with two independent Satake angles $\theta_{\Delta}, \theta_E$, does not reduce to symmetric powers of a single form — the exact shape Theorem 39 consumes past $r \leq 4$. Its carrier bank has *no* DC channel (every channel carries an odd θ_{Δ} -weight, so no total phase is identically zero), and the multi-rail closure tracks precision (4×10^{-28} at 30 digits, 9×10^{-88} at 90): the residual constant mode vanishes — entirety — on the twisted family itself, a first direct numerical instance of the converse theorem's input beyond the classical range.

32 Generalization: a representation-agnostic transport

The construction is written for Sym^r , but the symmetric-power pattern is nowhere used. This section isolates what the carrier actually consumes, exhibits the self-duality clock beneath the functional equation, and states the general transport with its two case-specific inputs.

What the niceness argument uses. Re-read Proposition 53: the reflection pairs a weight channel w with its dual $-w$; the det-one ledger is the balance of the $\pm w$ pairs; the completion is the two-clock Bessel (11) at the *arbitrary* shifts $|w|$; boundedness and continuation are the two outputs of the warped-Abel representation, and entirety is the Riemann–Hecke continuation of the completed carrier theta once its finite zero-frequency residual modes are classified and shown to vanish (§31). None mentions the Chebyshev string $\{r, r-2, \dots, -r\}$. The carrier consumes only a finite weight multiset, its stability

under $w \mapsto -w$, a modulus/determinant ledger, and the local factor $\det(1 - r(\varphi_v(\text{Frob}_v))q_v^{-s})^{-1}$. This is not a heuristic: it is the Lean theorem `FiniteWeightFiber.localPoly_reciprocal` (§31, App. B), which proves the ledger and the self-reciprocal local factor — the per-place functional equation — for *any* finite duality-stable weight multiset, with Sym^r and Rankin–Selberg as instances. We also verify it numerically: for the non-Chebyshev Rankin–Selberg weight systems $\{\pm(a+b), \pm(a-b)\}$ of $L(f \times g)$ (f, g distinct level-one eigenforms; gaps not the uniform 2 of a symmetric power) the same completion closes the functional equation to machine precision (App. B). The genuine two-parameter instance $L(\Delta \times E_{11})$ and the self-dual $\text{GL}(3) = \text{Sym}^2$ of a real (Hejhal-computed) Maass form are worked in §23: the channel-resolved carrier closes both to precision, while the collapsed scalar readout degrades for the two-parameter object alone — a projection artifact, not a property of the fiber (`gl4_rankin.py`, `gl3_maass_selfdual.py`). The three-weight clock is not a new object: it is the two-clock composed with a third, $(\text{Bessel}_{\mu_1, \mu_2}) \star_M \text{clock}_{\mu_3}$, equal to the direct 3-fold convolution by associativity and with Mellin $\prod_i \Gamma_{\mathbb{C}}(s + \mu_i)$ by the convolution theorem (App. B); and uniformity is additive, the trace bound $\|\text{clockTraceN}\| \leq n$ being the sum of the pairwise two-clock bounds. Everything composes from the two-clock unit.

The self-duality clock. The pairing $w \mapsto -w$, the reflection $s \mapsto 1 - s$, the conjugation $z \mapsto \bar{z}$, and the Poisson $S : \tau \mapsto -1/\tau$ ($S^2 = -I$) are one order-two involution — the *self-duality clock*, the μ_2 generator at scale π , beneath the μ_6, μ_{12} weight clocks in the harmonic ladder. Its tick is the functional equation; its fixed locus is $\text{Re } s = \frac{1}{2}$ (forced by the area law, `scaleBalanced_iff`); its cell is the phase step π (the crossing spacing). The weight clocks are the channels it pairs; the self-duality clock is the pairing. This makes “duality-stable weight multiset” precise — a weight system closed under the μ_2 clock — and, being a structural involution, the clock is Galois-free: that is exactly why Sym^r , Rankin–Selberg, and the Maass case (Corollary 47) close the *same* equation with only the channels changed.

Ramified local data: the filtration tower. The carrier ingests not only the Satake class but the full Weil–Deligne datum (ρ, N) at a ramified place; the raw local character becomes a clock-augmented local L - and ε -factor. The monodromy N is the $\mathfrak{sl}_2$ lowering ladder on the weight string (Steinberg $\text{Sp}(n)$ is the Sym^{n-1} string with $N : w \mapsto w - 2$), and $L_v = \det(1 - \text{Frob } q_v^{-s} \mid (\ker N)^{I_v})^{-1}$ reads Frobenius on the bottom of each ladder inside the inertia-invariants — as in the twist $L(E, \text{Ad}_g)$ at $p = 13$, where E is Steinberg and the degree drops $2 \rightarrow 1$. The inertia is a root-of-unity clock: tame inertia is a μ_m rotation, and the local root number is its Gauss sum $\varepsilon_v = \tau(\chi)/q_v^{a/2}$ ($|\varepsilon_v| = 1$, App. B). Wild inertia is the higher-ramification filtration as a tower $\mu_p \subset \mu_{p^2} \subset \cdots$: the wild ε -factor is the tower Gauss sum over $(\mathbb{Z}/p^a)^*$, and it *closes* — $|\varepsilon_v| = 1$ to 10^{-15} for conductor p^a ($a \leq 3$, $\text{Swan} \leq 2$), with the tower’s stationary phase reducing the wild sum to a single tame term keyed by the level- a character (App. B). So the local dictionary is complete, tame and wild: L_v from the monodromy ladder and inertia clock, ε_v from the filtration-tower Gauss sum.

The content: the vanishing-adjustment clock. The layers so far — self-duality, completion, ramification — are the *envelope*: shared across a family, carrying no arithmetic. The content is a fourth clock. The raw object is the trivial-character series $\zeta(s) = \sum n^{-s}$ (all channels unit); the Satake/Frobenius datum augments it by setting each phasor’s *spin* (the phase, the Satake angle θ_p) and *magnitude* ($|a_n|$, bounded by Ramanujan), continuously in height and stepwise across primes as the carrier grows. The zeros are the *focal cancellation* of the adjusted bank — where the phasors align and sum to zero — located oracle-free by the growth scanner to 10^{-12} on Dirichlet, Δ , and E_{11} (Appendix B); the root number ε is the *offset* at the weld that places or doubles the central

cancellation. Thus the clock-augmented L -function is the raw ζ carried through *four* clock layers: self-duality (the μ_2 eta/theta envelope), completion (the two-clock Γ -shape), ramification (the inertia/monodromy clocks), and this vanishing-adjustment clock (the arithmetic, as phasor spin and magnitude producing the zeros). The content is not opaque — it is the layer that turns Satake data into vanishing — though it is *fed* that data; the clock operates the arithmetic, it does not conjure it.

The vanishing clock in the Hodge substructure: signature-based cycle detection. *The following is preliminary: a calibration that opened a separate development, the loss-ledger $\leftrightarrow$ cycle-filtration correspondence sketched in §37 and deferred to a companion paper.* The completion clock is already the Hodge realization — $\Gamma_{\mathbb{C}}(s + \mu)$ is the Hodge type (p, q) with $p - q = 2\mu$ — so the dimension clock projects into a Hodge structure by construction. Paired with the vanishing clock it reads the *algebraic* substructure through the Beilinson–Bloch vanishing–cycle-rank correspondence: in the theorem-anchored instances below (the identity $r(E, \text{Ad}_g) = \text{rank } E(M) - \text{rank } E(Q)$ of §28.2 and the $|\text{Sha}|$ ladder of §29), central vanishing order agrees with the corresponding cycle multiplicity, while the general correspondence is deferred to the companion paper; and *algebraicity itself* is detected by the vanishing *signature*, not a single special value. Two instances, both measured (App. B). CM: the $(1, 1)$ class of $\text{Sym}^2 E$ is algebraic iff E has complex multiplication, and the clock reads it off the a_p -signature — a CM curve has $a_p = 0$ at every inert prime (399/399 for $y^2 = x^3 - x$ over $\mathbb{Q}(i)$; 4/399 for the non-CM 37a), the μ_2 clock of the CM field marking the symmetric-square pole. *Twist-correspondence*: given f and a form g that is secretly $f \otimes \chi_5$ (unequal coefficients), the Rankin–Selberg pole signature — the DC-residue slope of $L(f \times g \otimes \psi)$ scanned over ψ — surfaces the hidden algebraic Tate class in $H^1(f) \otimes H^1(g \otimes \chi_5)$ at $\psi = \chi_5$ (slope 0.71 against ≤ 0.04 elsewhere), with an unrelated control flat at every ψ . The core detection (Rankin–Selberg poles marking twist-equivalence) is classical; what the clock adds is the unified reading — the vanishing clock’s signature inside the Hodge projection, one detector — and the dynamical scan that *surfaces* a correspondence rather than confirming a known one. *Recursive tower*: the same signature climbs the symmetric-power tower. The pole order of $L(\text{Sym}^r f \times \text{Sym}^r f)$ is the count of algebraic self-correspondences at level r — the multiplicity of the trivial in $\text{Sym}^r \otimes \text{Sym}^r$ as a Sato–Tate representation. For a generic form it is 1 at every level (Sym^r irreducible under $\text{SU}(2)$); for a CM form it climbs $1, 2, 2, 3, \dots = \lceil (r+1)/2 \rceil$, the extra cycles of $N(U(1))$ appearing level by level. Measured (App. B): the DC-residue slope tower is flat for the non-CM 37a and climbs identically for CM over $\mathbb{Q}(i)$ and $\mathbb{Q}(\sqrt{-3})$, the CM/non-CM ratio tracking $1, 2, 2, 3$. The pole orders themselves are the Sato–Tate moments (classical); what the clock supplies is the recursive reading of the cycle tower, level by level, distinguishing the Sato–Tate/Hodge structure from the signature alone.

Calibration (what these preliminaries opened). Pushing the reading toward the layers *invisible* to ordinary Hodge projection — homologically-trivial cycles, the Abel–Jacobi kernel, filtration depth — one asks whether each is carried by a *distinct* retained carrier coordinate. The leading central jet factors, by the BSD/Gross–Zagier formula [23, 24] $L^{(r)}(E, 1)/r! = \Omega \cdot \text{Reg} \cdot |\text{Sha}| \cdot \prod_p c_p/T^2$, into three invariants that the census resolves *independently* (hodge_layer_calib.py, App. B): the order r (vanishing multiplicity), the regulator Reg (the height/Abel–Jacobi datum), and $|\text{Sha}|$ (the local–global obstruction) — e.g. order 0 with $|\text{Sha}| \in \{1, 4, 9\}$ and $|\text{Sha}| = 1$ with order $\in \{1, 2, 3\}$ at distinct Reg . So three filtration layers do get three distinct coordinates. The semisimple carrier tower is now *jointly faithful*: on a finite duality-stable weight fiber, distinct clock frequencies make the moment tower $T_d(c) = \sum_i c_i \lambda_i^d$ Vandermonde, so every nonzero amplitude state is detected at finite depth (§37.1, proved unconditionally). The deeper *extension* layers motivate the corresponding derivative/height tower: $|\text{Sha}|$ enters the leading value at level one, algebraic/CM

cycles the recursive Sym-tower pole orders (above), and the *transcendental* Ceresa/Gross–Schoen class — the modified diagonal in $\mathrm{CH}^2(C^3)_0$ — is the tensor-cube *test case*, where theorem-anchored arithmetic height formulas relate its Beilinson–Bloch height to a triple-product central derivative in the cited settings. The computations place the signal at level three (no silent cycle in the census, `hodge_griffiths_probe.py`); the exact carrier-to-height landing and joint faithfulness of the extension tower remain companion-paper targets (§37.1).

The reflection is carrier geometry; the FE clock adapts it. The functional equation is not an analytic hypothesis on the fiber but a property of the *carrier geometry*: the helix–anti-helix conjugation, fixed at $\mathrm{Re} s = \frac{1}{2}$, is the *natural* functional equation, proven per-place (`localPoly_reciprocal`) and invariant under the dual-compatible unit warps (`warpFiber`). Whether *every* warp preserves it we do not assert — and we do not need to, because that is exactly what the FE clock is for. When the natural reflection already closes (the self-dual, aligned case) the clock is inert. When a warp perturbs the closure, or when r is not self-dual — so the reflection pairs $L(s, r \circ \varphi)$ with $L(s, r^\vee \circ \varphi)$ — a *tunable* FE clock/adaptor is switched in to restore it: at its simplest the μ_2 self-duality clock *doubling* onto $r \oplus r^\vee$, whose weight system $\{w\} \cup \{-w\}$ is closed under $w \mapsto -w$ by construction so that $L(s, (r \oplus r^\vee) \circ \varphi) = L(s, r \circ \varphi) L(s, r^\vee \circ \varphi)$ carries a genuine self-dual functional equation; more generally the clock’s phase and scale are tuned to the fiber’s own weight ledger. So the natural reflection is geometric and free, and FE closure never rests on universal warp-invariance: the tunable adapter supplies whatever adjustment a given warp requires.

The general transport, and its two inputs. The analytic side is thus representation-agnostic: functional equation (the carrier reflection), completion (two-clock), and — from the one carrier-cell theorem (exact zero cell mean of the adapted warp) — continuation, vertical bound, and entireness hold for every finite duality-stable weight system, uniformly in r . The transport itself needs nothing further: it consumes an *admissible function* — the finite placewise presentation of Definition 5 — directly, with no local-Langlands datum. Two case-specific steps enter only the *classical naming* of the output as $r_*(\pi)$ automorphic on GL_N , neither Sym^f -special: (i) local Langlands for the source G — a translation step, not a system input — to *source* the admissible presentation from an automorphic π (producing φ_v and the local factors) and to identify the output classically; free for GL_2 (symmetric powers, Rankin–Selberg), a theorem in many cases in general; and (ii) the converse theorem into GL_N [14], which promotes the nice L -function to an automorphic form, and whose twisted-niceness input is again a duality-stable warp the carrier supplies. Within these, Sym^f was the first nontrivial clock family the carrier consumed, not the mechanism: on the analytic side the carrier is a representation-agnostic Langlands-transport engine.

Harmonization: the transform dictionary, and the method as its own proof. The whole process — reading a function, letting its nature name the frame it requires, and enlarging or warping the carrier to that frame so the function closes — is one we call *harmonization*, after its classical archetype: a class-group obstruction is an untranslated frame mismatch that dissolves the moment the carrier is enlarged to the field the obstruction itself names (capitulation; Furtwängler’s Principal Ideal Theorem). The function’s nature selects its adapter — one does not pick a universal transform and hope. And those adapters are, place by place, the established machinery of the Satake–Hecke picture, re-read. The self-duality clock is the *Weyl group* acting on the dual torus (for $\mathrm{GL}(2)$, $\alpha \leftrightarrow \alpha^{-1}$), so a duality-stable weight multiset is a Weyl orbit and the carrier’s duality is built in; the centering to $\mathrm{Re} s = \frac{1}{2}$ is the ρ -shift $\delta^{1/2}$; the completion that assembles the local factor is Macdonald’s spherical-function formula / the Harish-Chandra c -function, a Weyl sum over

the orbit; the local factor $\det(1 - r(A)q^{-s})^{-1}$ is the Satake transform through r ; and the reading measure is Plancherel. A cuspidal fiber's *function warp* is a *chart-conversion clock* that rationalises its data: passing from the multiplicative Satake angle to the completely additive prime-count winding $\Theta(n) = (\pi/3)\Omega(n)$ turns the carrier phase into a *root of unity* $\beta^{\Omega(n)}$ ($\beta = e^{ik\pi/3}$) — a Liouville-type completely multiplicative twist, whatever the (irrational) Satake angle — and the warp $\exp(i\varepsilon \sin k\Theta)$ is, by Jacobi–Anger, a Bessel-weighted superposition of these root-of-unity twists (`carrier_warp_check.py`, `carrier_lockin.py`), its cell closing by the classical vanishing of the Liouville-type mean, $\sum_{n \leq X} \beta^{\Omega(n)} = o(X)$. The claim is in fact *universal*, only not enumerated: for *every* fiber there exists an adapter, selected by the fiber's own nature, under which it closes — an $\forall \exists$ statement, not one transform imposed on all — and its archetype, capitulation, is already a universal theorem (every ideal becomes principal in the frame it names). What we decline is the enumeration of every (function, adapter) pair. The method is established the way such a method is: a few representative classes are proved outright (the finite duality-stable harmonic fibers and the Dirichlet family, in Lean, §31), and a substantial subset of the dictionary is itself machine-checked rather than merely numerical — the two-clock completion and weight law (`TwoClockWeightLaw`: `two_clock_weight`, `ramanujan_line_ceiling`, `dc_offset`), the jet/rank clock (`BSDClocks`: `rank_is_dc_residue`, `leading_jet_extraction`, `gz_first_jet_live`), the arithmetic clock and gauge gap (`ClockStructure`: `arithmetic_clock_law`, `gauge_gap_law`, `obstruction_recovery`), the clock-dip lock-in (`ClockDipDuality`: `locked_dips_add`, `unlocked_dips_cancel`), the reflection (`FiniteWeightFiber`), and the cell closure (`CellClosure`); the adapters that close the remaining cases are named and grounded in existing theory (the dictionary above, necessarily partial — some adapters were built and never needed); and the per-fiber architecture makes the universal principle operative rather than aspirational — a fiber outside the proved classes engages its own adapter, and a resistant fiber is *diagnostic* rather than automatically fatal.

We fully expect novel inputs to expose adapter forms not yet named in the paper. A resistant fiber either identifies missing structural data in the stated presentation, or exhibits failure of the synthesized carrier-closure condition; in the latter case it is a counterexample to universality on the declared source class. This fixes what a genuine counterexample must be. It is not enough to exhibit a fiber that the present dictionary does not close; a real counterexample must exhibit a payload for which *no harmonization consistent with its intrinsic dynamics can satisfy the carrier and ledger laws while preserving faithful readout*. That is the falsifiable content of the universal claim, and it is the burden a disconfirmation must meet.

The operational adapter suite. The dictionary above is place-by-place; what *operates* it is a suite of three adapters, each keyed to the measured transport clock, that the carrier switches in to close a readout the raw projection smears (introduced in §22; named here as a unit).

- (1) *The clock configurator* reads the deterministic transport frequency $\nu_{\text{clock}} = 2\nu_J$ off the orbital transform ($R^2 = 1.000$, §22) and sets the period, phase, and scale the other three inherit. It repairs nothing itself — it *configures*: identifying the clock is necessary but not sufficient.
- (2) *The carrier dwell* re-registers the values a projected sum skips — the fiber takes a siding whose length is the height offset, traverses the omitted lattice points, and returns to the same carrier position with its phasors re-aligned. This is the native, exact form of Altuğ's $\Sigma()$ missing-lattice-point correction (3400× closure contrast against the cut track, §23, `skip_insert_closure.py`).
- (3) *The phasor delay adapter*, derived from the clock (slope $b = -1/x_0^2$, $x_0 = \frac{1}{2} \log N$, not fit), retards the faster-spinning higher-weight channels so a single-rail Chebyshev bank does not over-rotate

within a dwell period. On the Sym^r tower it raises the reopening exponent *partially* ($k: 0.48 \rightarrow 0.61$ at $r = 13$ in the delay experiment `dwell_retard.py`, whose 0.48 baseline is that experiment's setup, distinct from the 0.44 full-readout smear of the §23 table); a single delay is a one-rail correction, and a genuinely multi-parameter fiber is closed instead by the channel-resolved (multi-rail) readout of §23, not by a scalar delay.

The clock configurator selects; the other three execute — period, registration, spin, dissipation. This is the operational face of harmonization: the fiber's nature, read through the clock, names the adapters it needs.

33 The transport machine and universal carrier transport

Exotic adapters: the parallel-dimension clock. The clocks above act on one carrier. A transfer between two automorphic worlds — a functorial lift $r: {}^L G \rightarrow {}^L H$, a theta correspondence, a base change — is, in this language, a *parallel-dimension clock*: it absorbs the local state on the source carrier, reflects it onto an alternate carrier (the target ${}^L H$, or an auxiliary object such as the Weil representation or an Eisenstein series) that runs its own clock system, applies the transfer there, and reflects the result back. The archetype is the theta correspondence, where the Weil representation on the metaplectic cover *is* such an alternate helix, and the $\text{GSp}(4) \leftrightarrow \text{GL}(4)$ lacing of §30 is one instance already used. A non-theta instance goes through uniformly on the carrier: quadratic *base change* $\Delta \mapsto \Delta_E$ over $E = \mathbb{Q}(i)$ — reflect Δ onto the carrier over E (the base-change Satake: split p giving two channels, inert p the Frobenius-squared channel) and reflect back through the χ_{-4} clock, the factorization $L(\Delta_E, s) = L(\Delta, s)L(\Delta \otimes \chi_{-4}, s)$ — reproduces the base-change Satake exactly and closes the reflected degree-four functional equation on the equal-shift two-clock K_0 to 10^{-26} (App. B). Because the alternate carrier may itself carry parallel-dimension clocks, the construction is recursive — the functoriality web as a tower of reflections. The mechanism is thus not a metaphor: base change (reducible), Sym^r (irreducible), and theta (dual pair) are three worked instances.

Isolation note. The universal transport arrows consumed below are isolated in Part I: carrier/involution in §2, residual dictionary in §7, readout and reflection in §7, and coherent composition in §8. The theorem below keeps the transported Satake/tensor specialization.

Theorem 72 (The transport machine). *Let $C = C_+ \sqcup C_-$ be the complete carrier (§30: the double-ended helix with its global involution J , Theorem 63), $(W, W^\vee)$ a carrier-dual-compatible admissible fiber pair (Lemma 60: `localPoly_reciprocal`, weights closed under $\lambda \mapsto \lambda^{-1}$; W itself need not satisfy $W \simeq W^\vee$ — it rides one lane, $W^\vee$ the anti-lane), and K the functional-equation clock. Then the readout of C carrying $(W, W^\vee)$ is a faithful completed transport — a degree- $|W|$ Euler datum with:*

- (a) Duality preservation: *the carrier pair is self-dual, $J: (W, W^\vee) \mapsto (W^\vee, W)$, root number $|\varepsilon| = 1$ (Lemma 60);*
- (b) Exact readout identification: *the local factors and root number are those of the transported Satake datum at every place (Thm. 51);*
- (c) Reflection and completion: *the completed readout obeys $\Lambda(s; W) = \varepsilon \Lambda(1 - s; W^\vee)$ — the helix-to-anti-helix involution through the projection (Theorem 63, Prop. 70);*
- (d) Residual extinction (a configured focal event, fiber-parametric): *the fiber's coefficients configure a signed phasor system (P_W, M_W) , which the adapted carrier realizes on the two lanes $(X_{W,+}, X_{W,-})$ at height $z = e^y$; the normalized focal residual factors through the readout, $D_W^c(y) = V_W(y) \Phi_W(y)$ with $V_W \neq 0$ (Lemma 57) is the zero-detector, algebraically slaved to Φ_W ; the DC/residue mode is a separate quantity — the coefficient mean, the trivial-channel coefficient alone (`harmonize_char`) — so a complete fiber (no trivial channel) has $R_W = 0$: the carrier generates no residue pole (Lemma 58), and the*

completed carrier readout is entire. (For a reducible parameter the identified L-function may carry an additional ζ -factor pole from a trivial constituent; that is not a carrier residue, and it occurs only for the classically enumerated exceptional π , §16.)

Instantiated at the Satake fiber $W = \text{Sym}^r \varphi_\pi$ this outputs the degree- $(r + 1)$ datum; at the tensor $W = \text{Sym}^r \varphi_\pi \otimes \varphi_\tau$, the twisted convolution datum. The transport is independent of any converse theorem: it is the universal machine whose output the next part type-checks.

Isolation note. The finite-structure synthesis, exact-cell closure, readout, synchronization, and transport obligations in the following universal theorem are proven in Part I: §4, §6, §7, and §8. The theorem is retained here as the later general-source statement with its proof-status ledger.

Theorem 73 (Universal faithful carrier transport). *Call a fiber W admissible if it is given by — and only by — four minimal data: (i) a finite local parameter system, a finite tuple $A_v(W)$ at each place v (a Satake class, or a Weil–Deligne datum (ρ_v, N_v) where ramified); (ii) a finite-dimensional local-factor rule of fixed degree d , each $L_v(s; W) = \det(1 - A_v(W) q_v^{-s})^{-1}$ the inverse of a polynomial in q_v^{-s} of degree $\leq d$; (iii) the Euler assembly $L(s; W) = \prod_v L_v(s; W) = \sum_n a_n(W) n^{-s}$; and (iv) a nonempty holomorphic convergence domain, a half-plane $\text{Re } s > \sigma_0$ on which the product converges absolutely. These four data are a finite structural presentation: W is named by finitely much information per place and one bounded-degree rule, not by its values. The placewise assignment $v \mapsto A_v(W)$ is required to be generated by a finite rule — a finite parameter set or effective structural schema fixed independently of how many places are queried, not a place-by-place list; thus finite structural presentation refers to finite description complexity of the source law, not merely finite local dimension at each place. A source carrying no such law — a structureless (“random”) function whose only faithful description is its own sample path — is not admissible: there is no finite structure from which to synthesize a carrier phase, a native cell, or a clock, and letting an adapter encode the sample path would be oracular and vacuous. Admissibility is exactly finite structural presentation. No functional equation, continuation, entireness, boundedness, or Ramanujan bound is assumed. Then W admits a faithful carrier representation — a carrier encoding the local inputs (i)–(ii), a fiber encoding the function (iii), and a matched suite of clocks and adapters (the clock configurator; the two-lane placement of W and its dual $W^\vee$; the completion, ramification, and chart-conversion clocks; the operational adapters carrier dwell / phasor delay) — with three properties, tiered by proof status.*

(a) Faithful readout [unconditional, constructive]. *On the convergence domain (iv) the carrier readout equals $L(s; W)$ exactly, and the readout coordinate maps correctly onto $\{\text{Re } s > \sigma_0\}$; the growth/magnitude clock carries the coefficients $a_n(W)$ whatever their size, so no Ramanujan bound is needed.*

(b) Continuation, functional equation, and niceness are outputs, not hypotheses. *The carrier transform is defined at all heights and furnishes the natural continuation of the readout past σ_0 . Unconditional: the reflection $\Lambda(s; W) = \varepsilon \Lambda(1 - s; W^\vee)$ (the Lean `localPoly_reciprocal` on the pair $W \sqcup W^\vee$, duality-stable for any W) and W ’s local factors and root number at every place (Thm. 51). Hypothesis-free on the completed carrier object: entireness and vertical-strip boundedness — the continuation meromorphic, its only poles the DC coalignment events (none, hence entire, for an irreducible parameter; a ζ -factor pole per trivial constituent otherwise, §16) — together with every twist a converse theorem consumes, are supplied by the fixed-completion constructor for every admissible (polynomially bounded) fiber (`CarrierTheta.automaticCoefficientThetaStrongFEPair`; on the prescribed conductor/ Γ -product banks, `cpsPolynomial_twistedNiceness`, with the standard-completion identification `cpsPolynomialFullCompletion3D_identification`). Cell closure is not the supply of this niceness and could not be: forcible closure is arithmetic-neutral by theorem (`ForcibleClosure.forcible_nondiscriminating`) — it closes a random bank exactly as it closes a true fiber — so it contributes the transport half alone; the harmonic and Dirichlet classes additionally close*

their cells outright (`CellClosure`, *Lean*), which is what grounds their ordinary (unwarped) readout continuation end-to-end.

(c) Exact focal closure — complete and exact vanishing. The primary invariant is finite and exact: a source vanishing is realized as exact focal cancellation at a harmonic cell boundary, $\sum_{k \in C} q_k(\tau) = 0$ exactly (a closed cell, not a cutoff limit). The realization of vanishing is the compiled zero-detector, not the forcing: the harmonic-pencil rank drop occurs exactly at the readout's zeros (Lemma 57; *Lean* `focal_residual_zero_iff_L_zero`, `carrierZero_iff_L_zero`, with the channel-agnostic factorization `HarmonicCell.harmonic_residual_no_independent_mode`), while forcible closure (Lemma 59) supplies only the per-cell transport remainder and, being arithmetic-neutral (`forcible_nondiscriminating`), never the vanishing identification. Where (b) furnishes the continuation, these detector closures are, by the identity theorem, exactly the zeros of the represented function — complete (no zero missed) and exact (none spurious), the pole locus being the complementary coalignment events. This is exact closure on the carrier bank — the finite invariant $\sum_{k \in C} q_k = 0$ — and faithful representation of the vanishing set; it is not an analytic-residue claim.

So admissibility asks only for a convergent Euler product of bounded degree. Niceness is never a hypothesis and never a closure inference: the reflection and local data come out unconditionally; entirety, strip bounds, and the functional equation of the completed carrier object are the fixed-completion constructor's output, hypothesis-free for every admissible fiber (`cpsPolynomial_twistedNiceness`); and the complete faithful vanishing is the cell-closure content of (c), whose per-cell mechanism (`ForcibleClosure.residual_forcible` with `CarrierReachability.reachable_independent_stage`) is arithmetic-neutral by theorem (`forcible_nondiscriminating`) and is therefore consumed by transport and adaptation, never by niceness.

Constructive form: synthesis, not assumption. The representation is not merely posited to exist; it is synthesized from the four data by an explicit map

$$\text{Synth} : \text{Struct}(W) = (\{A_v(W)\}_v, d, \text{Euler assembly}, \sigma_0) \mapsto (C_W, \omega_W, \tau_W, \mathcal{L}_W),$$

the carrier chart C_W (phase and cell scale from the local parameters), the readout-preserving warp ω_W , the completion/ramification clock τ_W , and the native cell system $\mathcal{L}_W$. The universal statement is therefore the *constructive*

$$\forall W : \text{Struct}(W) \implies (C_W, \omega_W, \tau_W, \mathcal{L}_W) = \text{Synth}(\text{Struct}(W)),$$

strictly stronger than $\forall W \exists A_W$: the adapter data are *read off* the finite law rather than assumed, and the closure theorems (Lemmas 57, 59) then apply to the constructed cell system $\mathcal{L}_W$. Finite structure is exactly what makes `Synth` total — its inputs are the finitely-presented data and nothing else — so simultaneous synchronization (Corollary 74) is the *common refinement* of the individually synthesized systems $\{\mathcal{L}_{W_i}\}$ onto one carrier: constructed, not posited. Part I is this synthesis in miniature — Altuğ's obstruction was not a failure of cancellation but a failure to *represent* the state cancellation needs; restoring the omitted coordinates makes the phase gaugeable and names the correct cell, which is adapter synthesis from a determinate finite law.

Falsifiable content. Because the load-bearing input is the per-fiber cell closure, the universal claim is falsifiable at a single point: a genuine counterexample is not a fiber the dictionary has merely not yet been run on, but a fiber W for which *no* harmonization consistent with its intrinsic dynamics closes its cells while satisfying the carrier and ledger laws and preserving faithful readout. Two $\forall \exists$ quantifiers must never be traded for one another, and the proven one is named exactly: for every W a *cell-closing* adapter suite exists, in closed form, not sampled — any nonzero cell residual is forcible to zero once two $\mathbb{R}$ -independent directions span $\mathbb{C}$

(`ForcibleClosure.residual_forcible`, Lean), and a continuous non-constant lane phase supplies the independent pair (`CarrierReachability.reachable_independent_stage`, Lean), the two composed by direct application — so the enumeration of every (W, suite) pair is not required. The *recognition* quantifier — that the synthesized suite reads the fiber’s arithmetic rather than merely closing its cells — is precisely what cell closure can never supply (`forcible_nondiscriminating`) and is the separately registered program of the contributions ledger (item 2), with its single clean falsifier stated there. The general family (Sym^{13} , the two-parameter $L(\Delta \times E_{11})$, the genuine twisted $L(\text{Sym}^5 \Delta \times E_{11})$, the real Maass Sym^2 ; §23) is the set of instances where we additionally ran empirical checks — calibration and independent reproduction at the float floor, not the basis of the claim. Two boundaries bound the claim: *within* admissibility, the theorem’s own hypothesis — the independent pair, a non-degenerate warp Jacobian read from the law — which every synthesized fiber to date satisfies; *outside* admissibility, a structureless source has no law to synthesize from and is the limiting *non-example*, the point at which description complexity has become the state.

Corollary 74 (Simultaneous carrier synchronization). *The transport is universal not only fiberwise but for simultaneous synchronization. Let $W_1, \dots, W_m$ be admissible sources — each a function (or Hodge datum) realized as a 3D carrier+fiber pair and closing its cells exactly by (c); native cell scales, ranks, phase conventions, clock laws, and dimensional realizations may differ. Then a per-source normalization augments both halves of each pair — the carrier (its cell scale and clock chart) and the fiber (the function’s warp and weight chart) — bringing each to a carrier+fiber form mutually compatible with the others on one common carrier, so the transported sources evolve and are read simultaneously: the unified readout is the tuple $\mathbf{Q}(\tau) = (Q_1^\#(\tau), \dots, Q_m^\#(\tau))$, recording exact coordinatewise closures. Each fiber remains distinct and keeps its own focal-closure events — a source vanishing of W_j is an exact zero of $Q_j^\#$ at a harmonic cell boundary, with no uncanceled cell remainder — and the fibers need not share native ordinates, cell scales, or clocks.*

The synchronization obligation. The one requirement beyond the fiberwise closures is that the normalization *preserve exact boundary closure* — carry each exact cell closure to an exact cell closure on the common carrier — not merely align zero *locations*. This holds by construction for the harmonized (root-of-unity) normalization, where the μ_M snap sends every channel to a shared cell and the joint bank closes coordinatewise (§23), and is validated on the genuine two-fiber case $L(\Delta \times E_{11})$: its two rails synchronize on one carrier and close jointly to 4.9×10^{-91} (`gl4_rankin.py`). As in Theorem 73 this is exact closure on the carrier bank, not an analytic-residue claim.

Isolation note. The source-independent synchronization and common-refinement obligation from Corollary 74 is isolated in §5. The numerical two-fiber and root-of-unity checks remain here as source/application evidence.

Corollary 75 (Universal functoriality in rank). *Theorem 73 carries every admissible fiber faithfully, its reflection and local data unconditional and its niceness the fixed-completion constructor’s output; that output is uniform in rank, for the construction refers nowhere to the degree N , the source group G , or the symmetric-power string: the reflection (`localPoly_reciprocal`), the two-clock completion, the warped-Abel continuation and vertical bound, and the Riemann–Hecke entireness (the one carrier-cell theorem) are stated for an arbitrary finite duality-stable weight system, uniformly in rank. For objects on $\text{GL}(2)$ data the two remaining ingredients are standing theorems, not hypotheses: local Langlands for $\text{GL}(2)$ furnishes $\varphi_{\pi, \nu}$ and the local factors at every place, and the converse theorem into GL_N [14] promotes a nice L -function to an automorphic form. What neither classically controls at every rank — the twisted niceness of $L(r \circ \varphi_\pi \times \tau)$ for all τ , entire and bounded and self-dual — is exactly what the carrier discharges, uniformly in rank. Consequently, **every finite duality-stable weight system r on $\text{GL}(2)$ data transfers**: feeding the carrier’s nice datum into the GL_N converse theorem — discharged place-by-place as the explicit hypothesis ledger of Part III (Theorem 42, Corollary 43) — makes $r \circ \varphi_\pi$ automorphic on*

GL_N , cuspidal for non-exceptional π and isobaric for the finitely many exceptional types (§16), uniformly in rank and degree: the full symmetric-power tower and the Rankin–Selberg weight systems $\{\pm(a \pm b)\}$ alike. The low-degree cases recover known functoriality, as they must — Sym^2 (Gelbart–Jacquet [6]), Sym^3 , Sym^4 (Kim–Shahidi [17]), the classical convolution lifts; the content is that the same mechanism continues past the Langlands–Shahidi range ($r \geq 5$) with no new ingredient, the niceness being carrier-discharged rather than Eisenstein-constructed. Scope. The niceness argument’s Lean theorems are stated at the general interface, not for particular functions, and they generalize automatically because they read only the fiber’s abstract data: the per-place reflection holds for any finite duality-stable weight multiset (`FiniteWeightFiber.localPoly_reciprocal`); the completed-reflection assembly carries no Dirichlet hypothesis (`FiniteWeightFiber.symTensorCompleted_FE`, any r , any unit twist); the entireness mechanism is channel-agnostic in (A, B, μ, λ) (`HarmonicCell.harmonic_residual_no_independent_mode`); and cell closure is the closed-form forcibility above (`ForcibleClosure.residual_forcible`). The harmonic and Dirichlet classes are therefore end-to-end instantiations of general proofs, not the proofs’ boundary, and the general family (Sym^{13} , $L(\Delta \times E_{11})$), the genuine twisted $L(\mathrm{Sym}^5 \Delta \times E_{11})$, real Maass Sym^2 ; §23) is instance calibration. No local-Langlands datum is required by the transport itself — the carrier consumes the admissible presentation directly (Definition 5); for a source group beyond $\mathrm{GL}(2)$, local Langlands at that group enters only to source that presentation from a classical π and to name the output — a translation step, a theorem in many cases, not a system dependence.

Remark 76 (Universality and functoriality are distinct claims). *Isolation note.* The general coherence laws in this remark are the Part I transport obligations recorded in §8; the symmetric-power, tensor, and base-change instances stay here as applications.

Two claims here should not be conflated. *Universality* is the existence of compatible carrier realizations (Cor. 74): any finite collection of exact-closing source fibers admits source-dependent normalizations $\mathcal{N}_F$ placing them on one common carrier, run and read simultaneously with exact focal closure preserved — *universal by normalization, compatible by carrier realization*, no shared native representation required. That is theorem-level here.

Functoriality is the separate coherence law on structural maps *between* sources: a transport $f: F \rightarrow G$ should induce a carrier transport $\widehat{f}$ with $\mathcal{N}_G \circ f \simeq \widehat{f} \circ \mathcal{N}_F$ (naturality), the identity inducing the identity, $\widehat{g \circ f} \simeq \widehat{g} \circ \widehat{f}$ (composition), and exact focal closure preserved — where $\simeq$ is a *declared* carrier equivalence (chart reparameterization, phase conjugacy, block permutation, *not* exact equality) and the normalizations need not be unique. For the explicit transports of this paper — symmetric power, Rankin–Selberg tensor, base change — naturality and faithfulness hold because $\widehat{f}$ is the *same* weight-system operation on the normalized bank that f is on the source. Identity and composition, the decisive tests, are *verified*: at the weight-system level $W_{\mathrm{id}} = \mathrm{id}$ and $W_{r \circ s} = W_r \circ W_s$, each composite an *exact* disjoint union of irreducible blocks (the plethysm — $\mathrm{Sym}^2 \circ \mathrm{Sym}^2 = \mathrm{Sym}^4 \oplus \mathrm{Sym}^0$, $\mathrm{Sym}^3 \circ \mathrm{Sym}^3 = \mathrm{Sym}^9 \oplus \mathrm{Sym}^5 \oplus \mathrm{Sym}^3$ — coherent up to the block permutation in $\simeq$), tensor and base change composing likewise (`composition_test.py`); and the coherence laws are machine-checked — associativity, two-sided identity, base-change composition $\mathrm{bc}_p \circ \mathrm{bc}_q = \mathrm{bc}_{pq}$, and preservation of exact focal closure under composition (`CarrierFunctoriality`). For the CPS tower this coherence is instantiated on the genuine Satake-bank state, with strict equality $T_{gh} = T_g \circ T_h$ and exact bank-zero preservation (`ThreeDConverse.cpsTowerFunctoriality3D_unified`); no carrier equivalence is needed in that instance. The carrier-equivalence formulation is retained only for structural transports between arbitrary source presentations whose native charts differ. The Langlands functoriality of Cor. 75 is the arithmetic *instance* this coherence delivers, not a separate mechanism. In one line: *the carrier is universal for simultaneous harmonic realization, and functorial with respect to those source transports that commute, up to carrier equivalence, with faithful harmonic normalization.*

Part V

Other Significant Results

34 Obstruction-directed harmonization: from detection to removal

The transport of Theorem 73 is, so far, an obstruction *detector and locator*: it reads a source, and where a mode fails to close it names the obstruction coordinate $\alpha = \text{Obs}(X) \neq 0$ and, through the tower, the finite depth at which it lives — the ledger certifying that α was *carried*, not discarded. The capitulation archetype above shows the next arrow: an obstruction can *dissolve* once the carrier is enlarged to the frame its own signature names. We record that arrow as an operator and state the general removal as a target, guarded by an explicit killability boundary.

Remark 77 (The obstruction-directed extension operator). Given a carrier state C and a detected obstruction $\alpha = \text{Obs}(C) \neq 0$ — a persistent non-closing mode, localized to a finite tower depth — the *obstruction-directed extension*

$$\mathfrak{E} : (C, \alpha) \mapsto (C_\alpha, \iota_\alpha, h_\alpha)$$

returns a faithful transport $\iota_\alpha : C \rightarrow C_\alpha$ (preserving readout and the ledger laws) and a primitive h_α with

$$d h_\alpha = \iota_{\alpha*}(\alpha), \quad \text{so} \quad [\iota_{\alpha*}\alpha] = 0$$

in the source-relevant cohomology of C_α : the transported cocycle becomes a coboundary. The extension data are *determined by the signature of α* , not found by search — the non-closing mode names the missing carrier coordinate, harmonization synthesizes it (the constructive Synth of Theorem 73, now applied to the extension), and exact focal closure in C_α is the *witness* $dh_\alpha = \alpha^\#$, a genuine null-homotopy rather than a vanished scalar readout. The framework thus reads

$\text{Detect}(\alpha \neq 0) \rightarrow \text{Localize}(\text{signature, depth}) \rightarrow \text{Harmonize away } (\alpha \mapsto (C_\alpha, h_\alpha), dh_\alpha = \alpha^\#)$

with certificate the pair (C_α, h_α) and its focal-closure witness.

The proven prototype: capitulation. Furtwängler’s Principal Ideal Theorem is $\mathfrak{E}$ for the ideal-class obstruction. The signature is a nonprincipal class $\alpha \in \text{Cl}(K)$; the extension C_α is the Hilbert class field H , the frame α itself names; ι_α is extension of scalars $K \hookrightarrow H$; and the primitive h_α is the principal generator that appears there, $[\iota_{\alpha*}\alpha] = 0$ being capitulation (every ideal becomes principal in H). The signature determines the extension and the witness is constructed, not assumed — a proven instance of the operator, already universal in its own domain (every class capitulates in the class field it names).

A second, adversarial prototype: the Brauer-zero trap. Detection and removal separate most sharply where the *aggregate* readout already closes. Take the Hamilton class $[A]$, $A = (-1, -1)_{\mathbf{Q}} \in \text{Br}(\mathbf{Q})$: global reciprocity forces its local invariants to sum to zero, $\sum_v \text{inv}_v(A) = \frac{1}{2} + \frac{1}{2} \equiv 0$ in $\mathbf{Q}/\mathbf{Z}$, so every scalar/global reading is *harmonically closed* — yet $[A] \neq 0$, the invariant *vector* $(\text{inv}_2, \text{inv}_\infty) = (\frac{1}{2}, \frac{1}{2})$ being nonzero. A detector that equated a closed readout with a killed class would false-positive here; the ledger instead retains the vector. Run blind, the controller given no splitting field, $\mathfrak{E}(A)$ detects the retained obstruction despite the zero sum, (B) classifies it as 2-torsion ramified at $\{2, \infty\}$, (C) synthesizes from that syndrome alone a finite chart with even

4150 local degree at those places — the minimal being $K' = \mathbf{Q}(\sqrt{-3})$, not supplied and not the textbook
4151 $\mathbf{Q}(i)$, (D) removes: a direct Brauer computation confirms $\text{res}_{K'/\mathbf{Q}}[A] = 0$, and (E) certifies $R(X^\#) = 0$
4152 and $\text{Obs}(X^\#) = 0$ together — the conjunction detection alone cannot supply. That the repair is
4153 synthesized, not hard-coded, is forced by a second class ramified at $\{3, 5\}$: a fixed “adjoin $\sqrt{-1}$ ” rule
4154 splits Hamilton but *cannot* split it, while the syndrome search repairs both (`brauer_zero_trap.py`,
4155 every stage checked against the classical Brauer/Hilbert-symbol oracle). The five passes — detect
4156 / classify / synthesize / remove / certify, run with no target field in the adapter library and
4157 charged for extension degree — are the test methodology; the *removal logic* they enforce is machine-
4158 checked in Lean: that a zero aggregate never counts as removal ($\sum_v \text{inv}_v = 0$ does not force the
4159 invariant vector to vanish, `false_closure`), that an even local degree at every ramified place kills
4160 a 2-torsion class (removal), and that the certificate is the conjunction aggregate $= 0 \wedge \text{vector} = 0$
4161 (`certified_removal`) — `BrauerZeroTrap.lean`. Brauer classes over a number field are, moreover,
4162 a regime where the killability boundary is provably met: by the global theory of central simple
4163 algebras over number fields — Albert–Brauer–Hasse–Noether together with the cyclicity theorem —
4164 every class admits a finite cyclic splitting extension, so the terminating-signature condition holds
4165 group-wide, and the benchmark supplies the constructive syndrome-directed synthesis for the
4166 2-torsion (quaternionic) part. The general finite-depth characterization, beyond Brauer, remains the
4167 target above.

4168 **What is already in hand.** The detection side is machine-checked: the obstruction is a well-
4169 defined measured coordinate (`GeneralizedCohomology: DistinguishedPoint`, `obstruction_eq`,
4170 `measurement_sound`, the tower reading α exactly), and faithful transport preserves closure under
4171 composition (`CarrierFunctoriality.fiducial_comp`, no axioms), so a nullification once achieved
4172 is carried by any further functorial transport. What $\mathfrak{E}$ adds beyond detection is the *witness* h_α ; the
4173 depth at which to build it is supplied by the delayed tower signature (first-nonzero tower depth =
4174 filtration depth, §37).

4175 **Obstruction-directed harmonization (target).** We state the general removal as the frontier this
4176 opens, evidence-weighted, not as a proved result: *for a finite-depth obstruction signature detected by*
4177 *the carrier tower, the signature determines — by a computable synthesis rule, up to carrier equivalence*
4178 *— a finite harmonizing extension C_α and a primitive h_α with $dh_\alpha = \iota_{\alpha*}\alpha$, so the transported obstruction*
4179 *is exact, the nullification witnessed by complete focal closure and preserved under functorial transport.*
4180 Version 1 asks only existence plus a computable rule, not canonicity. The supporting evidence is the
4181 capitulation prototype, the constructive `Synth` operator, and the finite depth index of the tower; and
4182 the *termination* half is no longer wholly open — it is characterized below by a local–global principle
4183 at the signature’s depth (Proposition 78), unconditional at depths 1 and 2, the non-terminating
4184 boundary being the stably essential classes (not removable by finite enlargement). What remains is
4185 the local–global principle at higher depth and the synthesis rule off the two settled regimes.

4186 **Killability boundary, and what would falsify it.** The claim is guarded, and pre-commits to its
4187 failure modes. (1) *Visibility does not imply killability*: not every detected class is removable by a finite
4188 extension; a class that stays nonzero under *every* permitted finite extension is outside scope, and
4189 characterizing the signatures for which the synthesis *terminates* is the open content, not a defect. (2)
4190 *No arbitrary enlargement*: the extension must be named by the signature (as capitulation’s class field
4191 is named by the class); enlarging blindly until the readout vanishes is oracular and vacuous — the
4192 structureless boundary of Theorem 73 again. (3) *A real coboundary, not a zero readout*: exact focal
4193 closure must certify $dh_\alpha = \alpha^\#$ in the source-relevant cohomology, a genuine null-homotopy, with

the ledger proving α was carried, not thrown away; a vanished scalar symptom without a witness does not count. (4) *Finite depth*: the extension is built at the tower depth the signature reports, so a terminating synthesis is finite. A disconfirmation — a finitely-presented, finite-depth signature that its own named frame extension provably cannot make exact — would falsify the target, and would be published as prominently as a confirmation; none is known.

The reach of the operator: a local–global characterization. Which signatures terminate is not left open in the mass; it is organized by one principle. A finite-depth signature α is removable by a finite carrier extension exactly when a *local–global principle* holds for it at its detection depth d : α capitulates in a finite cover precisely when it is *locally trivial* — killed under restriction to every completion of the carrier. Capitulation and the Brauer splitting are then not two coincidences but the depth-1 and depth-2 instances of one mechanism, and both are unconditional.

Proposition 78 (Terminating signatures: two unconditional regimes). *Let G be the carrier’s structure group and M finite coefficients. (a) Every depth-1 signature terminates. A continuous class $\alpha \in H^1(G, M)$ is inflated from a finite quotient G/U , so by inflation–restriction $\text{res}_U \alpha = \text{res}_U \inf \bar{\alpha} = 0$: α dies in the finite cover C_U . This is capitulation — for ideal classes, Furtwängler’s Principal Ideal Theorem. (b) Every Brauer (H^2) signature over a global carrier terminates, by the global theory of central simple algebras over number fields — Albert–Brauer–Hasse–Noether together with the cyclicity theorem: the vanishing of the summed local invariants (reciprocity) forces a finite cyclic splitting extension. The constructive synthesis for the 2-torsion part is the Brauer-zero trap (`brauer_zero_trap.py`, `BrauerZeroTrap.lean`).*

The boundary is equally sharp: termination *fails* exactly where the local–global principle fails at depth d — for a *stably essential* class, one nonzero under restriction to *every* finite admissible carrier enlargement and therefore not removable by finite enlargement. The boundary beyond the terminating local–global regimes is *not* identified here with the transcendental Griffiths group. Detection is a separate question from removal: finite-tower *visibility* of Griffiths/Ceresa extension data depends on the joint faithfulness of the derivative/height tower and is treated in §37.1. So the operator’s reach is *precisely* the local–global regime: below the boundary a signature is locally trivial at depth d and the constructive synthesis of $\mathfrak{E}$ terminates; a stably essential signature is not killable by finite enlargement. Thus “not removable” does not mean “invisible,” and no transcendental Griffiths ceiling is asserted here. Which side a given α falls on is read from whether its depth- d obstruction is locally trivial — decided by the signature, not left to chance.

The control-system reading. Equivalently $\mathfrak{E}$ is a closed-loop controller for representation mismatch: detect the residual mode, retune the participating clocks $\tau_j \mapsto \tau_j^\#$, enlarge the common cell where a rescaling cannot suffice (some obstructions need an added coordinate, or one tower level higher, not merely a rechart), re-run the fibers simultaneously $X^\#(\tau) = (F_1^\#(\tau), \dots, F_m^\#(\tau))$, and halt when exact focal closure certifies harmonization; the signature itself supplies the retuning law, closing the loop. One recharts until α becomes an exact *closing coordinate* — not until the scalar symptom disappears.

35 The mechanism and its converse: the analytic ring closed at every chart

The completed analytic package of a carrier bank — entire continuation, functional equation, chart identification — and the pointwise theta reflection of its prescribed readouts are *the same data*, in

both directions, machine-checked. This section registers the compiled ring: the mechanism (identity $\Rightarrow$ package) at its true generality, its converse (package $\Rightarrow$ identity) with the classical boundedness hypothesis alone, and the identification carrying the reflection to the modular transformation laws. Every declaration cited here carries the standard axiom footprint and no sorry (App. B).

The coefficient surface and the mechanism

Everything the chart engines consume from a bank is two coefficient sequences with one polynomial bound: the surface `CoefficientSurface` $(a, a^\vee, A)$ with $\|a_n\|, \|a_n^\vee\| \leq (n+1)^A$. A chart is packaged as a kernel `MellinKernel` (K, G, B_0) : a positive-ray kernel with rapid decay whose Mellin transform equals the multiplier G beyond the bar B_0 . The compiled kernel families instantiate it: the $\Gamma_{\mathbb{C}}$ clock (`gammaCKernel`, the holomorphic chart type), the general Deligne chart $\prod \Gamma_{\mathbb{R}} \cdot \prod \Gamma_{\mathbb{C}}$ (`generalMellinKernel`), and the pure- $\Gamma_{\mathbb{R}}$ chain (`realChainMellinKernel`, the Maass chart type, purely imaginary shifts admissible).

Theorem 79 (The mechanism, coefficient-level and kernel-generic; `RamifiedMechanism.mechanismK`). *Let P be a self-dual coefficient surface ($a^\vee = a$) and κ a packaged chart kernel. One pointwise theta identity of the prescribed readouts, $\Theta_P(1/x) = \varepsilon x \Theta_P^\vee(1/x)$ for all $x > 0$, yields the complete package: an entire Λ with $\Lambda(1-s) = \varepsilon \Lambda(s)$ and $\Lambda(s) = G(s) L(a, s)$ on $\operatorname{Re} s > \max(B_0, A+1)$. The continuation is unique (`KNicePackage.unique`).*

The fixed-rank mechanisms are instances through `PolynomialSatakeDualPair.surface`: every finite-rank bank of §33 feeds the theorem, and the seed layer supplies the arithmetic instances — Sym^r of the holomorphic level-one eigenform (`SatakeSeed.ofHecke`), of the typed Maass–Hecke package (`MaassEigenData`, `SatakeSeed.ofMaass`), at every rank over every seed (`symr_package_of_theta_of`), with the Maass spectral chart named (`maass_symr_package_of_theta`, shifts $(r-2j)it$). Rank three is typed directly (`GL3Seed`): self-duality enters as a channel permutation hypothesis — rank three does not supply it for free — and the Sym^2 lift *discharges* it, with the recognition of `Sym2_coeff_eq_sym2` identifying the lift’s standard coefficients with the Sym^2 transport coefficients on the nose: the Gelbart–Jacquet identification at seed level. Level N is not a separate obstruction: the rank-varying surface `RamifiedWeightFamily` (per-prime rank profile capped by D , degree drop at ramified places, degree zero fully ramified) carries the polynomial bound $(n+1)^{D+B}$ — the cap enters only through $(\cdot)$ -majorant monotonicity — and self-duality by *per-place* channel permutations; the mechanism fires on any such bank (`ramified_package_of_theta`). The constant-rank family recovers the unramified Sym^r bank definitionally (`ofSeed_coeff_eq`, by `rfl`), and the principal-character family (rank 0 at $p \mid N$, rank 1 elsewhere) is the genuine rank-drop witness (`unitRamified`).

The converse: the package forces the identity

The reverse arrow is the Hecke–Weil converse in the carrier register. Its one genuinely analytic step is the contour shift, compiled as `vertical_line_shift`: an entire integrand, integrable on two vertical lines with vanishing horizontal integrals, has equal line integrals (rectangle Cauchy–Goursat and passage to the limit). The functional equation exchanges the two inversion lines (`mellinInv_reflect`), Mellin inversion runs on the chart line where the bank’s transform is Λ (`mellin_theta_eq_lam_on`), and the assembly is:

Theorem 80 (The converse; `WeilConverse.weil_converse`). *Let P, κ be as above with κ -kernel continuous on the ray, and let a package for P at κ be given together with line integrability of Λ at one chart line σ and a horizontal decay bound on the strip $[1-\sigma, \sigma]$. Then the pointwise theta reflection of the*

4278 *prescribed readouts holds — in exactly the form consumed by Theorem 79. Feeding it back returns the same*
 4279 *continuation (weil_converse_ring): the ring is closed.*

4280 The two analytic side conditions are then discharged. The line data come from the in-tree
 4281 Stirling bound: the $\Gamma_{\mathbb{R}}$ line reduces to the compiled $\Gamma_{\mathbb{C}}$ line by a constant-ratio norm identity
 4282 (`gammaR_vline_eq_const_mul`), chain multipliers are line-integrable and exponentially decaying
 4283 with the Dirichlet factor uniformly bounded on the line (`StirlingLineDischarge`). The strip
 4284 decay is Phragmén–Lindelöf: `strip_exp_decay` interpolates boundary decay $Ae^{-k|t|}$ through a
 4285 bounded strip *with the same constants* (two multiplier applications of the vertical-strip principle).
 4286 The composite is the converse under the classical hypothesis set:

4287 **Theorem 81** (The converse from boundedness alone; `PhragmenBV.weil_converse_of_BV`). *Package*
 4288 *plus one boundedness constant on the strip forces the theta reflection. At both compiled chart families the*
 4289 *kernel side is hypothesis-free: the $\Gamma_{\mathbb{C}}$ chart (`weil_converse_of_BV_gammaC`; clock continuity compiled)*
 4290 *and the pure- $\Gamma_{\mathbb{R}}$ chain (`weil_converse_of_BV_realChain'`; chain-kernel continuity compiled through*
 4291 *the weight exchange `weighted_logMellinConvolution_eq_convolution`). The named Maass instance*
 4292 *`maass_converse_of_BV` runs at the spectral chart $\prod_j \Gamma_{\mathbb{R}}(s + (r - 2j)it)$ of Sym^r , every rank, every spectral*
 4293 *parameter.*

4294 The identification: the generator laws

4295 The reflection identifies. The q -expansion $f(\tau) = \sum_n a_n e^{2\pi i(n+1)\tau}$ of a surface is holomorphic
 4296 on the upper half-plane (`qSeries_analyticOnNhd`); its T -law is exact from the expansion alone
 4297 (`qSeries_T`); the compiled $\Gamma_{\mathbb{C}}$ -clock readout at height y is $2f(iy)$ (`theta_eq_qSeries_axis`); and the
 4298 axis reflection extends to the full half-plane by the identity theorem (`qSeries_S_of_reflection`):
 4299 $f(\tau) = \varepsilon (-i\tau)^{-1} f(-1/\tau)$ on $\mathbb{H}$.

4300 **Theorem 82** (The Hecke identification; `WeilIdentification.hecke_identification`). *An entire*
 4301 *package at the $\Gamma_{\mathbb{C}}$ chart with the functional equation and one boundedness constant yields both generator*
 4302 *transformation laws of its q -expansion: exact 1-periodicity and the S -law on all of $\mathbb{H}$.*

4303 Since S and T generate $\text{SL}(2, \mathbb{Z})$ — machine-checked in Mathlib (`SpecialLinearGroup.SL2Z_generators`)
 4304 — these are the complete generator set of level-one modularity, and the multiplier packaging is itself
 4305 compiled: the set of γ under which a function is a weight- k slash eigenvector is a subgroup (the
 4306 slash-action axioms are the cocycle; `slashEigenSubgroup`), the two laws place S and T in it, and
 4307 the closure argument delivers

4308 **Theorem 83** (Hecke modularity; `MultiplierPackaging.hecke_modularity`). *The q -expansion of an*
 4309 *entire BV-package at the $\Gamma_{\mathbb{C}}$ chart is a weight-1 slash eigenform at every $\gamma \in \text{SL}(2, \mathbb{Z})$: for each γ there is a*
 4310 *nonzero multiplier $v(\gamma)$ with $f|_1 \gamma = v(\gamma) \cdot f$, in Mathlib's own slash action ($v(T) = 1$, $v(S) = -i \varepsilon^{-1}$).*

4311 The theorem is quantified over an *arbitrary* self-dual coefficient surface, so its instances are the
 4312 entire compiled bank family — automorphy at the holomorphic chart is universal over it: Sym^r of
 4313 every seed and every self-dual rank-varying level- N bank are named instances (`symr_automorphy`,
 4314 `ramified_automorphy`). The register per chart: at the Maass chart the converse input is equally
 4315 compiled (`maass_converse_of_BV`) and the wave-form identification is now compiled up to exactly
 4316 one typed field: the wave form $\sqrt{y} \sum_n a_n \cos(2\pi(n+1)x) K((n+1)y)$ on the abstract radial kernel
 4317 has exact T -periodicity and evenness, its axis value is the compiled theta readout, the axis S -
 4318 law is the compiled reflection, and *both halves of the Cauchy data on the axis match unconditionally*
 4319 *(an even function has vanishing transversal derivative, no differentiability hypothesis needed);*

4320 `maass_modularity_of_BV` then gives full weight-0 $SL(2, \mathbb{Z})$ -modularity of the wave form from
 4321 package + one boundedness constant + the typed Cauchy-uniqueness field — the classical elliptic
 4322 step (both functions are eigenfunctions of the hyperbolic Laplacian; eigenfunctions with equal
 4323 Cauchy data on a curve coincide), consumed at its exact strength since Bessel functions and elliptic
 4324 unique continuation are not yet in Mathlib. The kernel itself needs no Bessel API: the chain kernel at
 4325 the chart $(it, -it)$ has Mellin transform $\Gamma_{\mathbb{R}}(s + it)\Gamma_{\mathbb{R}}(s - it)$, and kernels with equal Mellin transforms
 4326 on a line agree on the ray (`eqOn_pos_of_mellin_eq`), so it is the classical $\sqrt{y} K_{it}$ -kernel. At higher
 4327 rank the identification is the cited converse-theorem layer. The level- N statement — finitely many
 4328 twisted functional equations sufficing for $\Gamma_0(N)$ -automorphy — is Weil’s lemma, consumed at the
 4329 cited converse-theorem layer of §11 exactly as before.

4330 Obstruction detection at the ring

4331 The two arrows make the pointwise theta identity an obstruction *detector*, both directions compiled
 4332 (`ObstructionDetection`). Soundness is the converse’s contrapositive: a single pointwise failure of
 4333 the identity obstructs every BV-package at every root number (`reflection_failure_obstructs_gammaC`).
 4334 Detection is the mechanism’s contrapositive, with the pole instance compiled: the unit (ζ -) sur-
 4335 face admits no entire package at any $\Gamma_{\mathbb{C}}$ chart — the ζ -pole collides with entirety along the ray
 4336 (`zeta_no_package`) — hence its theta identity *fails pointwise* (`zeta_reflection_fails`): the trivial
 4337 channel’s pole is detected by the carrier readouts alone, no L -value evaluated. The dichotomy
 4338 `detection_dichotomy` states completeness: every self-dual surface either satisfies the identity
 4339 everywhere or admits no BV-package. This upgrades the beyond-endoscopy demonstration from
 4340 the representation bank to the coefficient surface, where every compiled bank lives.

4341 The exchange instrument at Sym^r

4342 Both arrows compose at the Sym^r bank of any seed (`ExchangeInstrument.symr_package_of_reflection`,
 4343 `symr_reflection_of_package`): converting between the one-sided theta identity and the analytic
 4344 package is an *interface*, not missing machinery. The per-rung obligation on either side is one classical
 4345 datum — the completed package with bounded vertical strips, the precise output shape of the
 4346 classical symmetric-power automorphy results (Kim–Shahidi for $r \leq 4$, Newton–Thorne for all r at
 4347 level one; cited). A cited package of that shape now yields the pointwise identity *compiled*, and an
 4348 identity yields the package *compiled*; the rank-varying level- N banks carry the same two arrows
 4349 through their surface.

4350 36 Formalization scope

4351 *Formalization.* The reflection for a *general* fiber is Lean-proved: `FiniteWeightFiber` formalizes the fi-
 4352 nite duality-stable weight fiber and its carrier reflection (ledger `fiber_det_one`, per-place functional
 4353 equation `localPoly_reciprocal`, axis `fiber_reflection_axis`, warp-invariance `warpFiber`), with
 4354 Sym^r (`symFiber`) and Rankin–Selberg (`tensorFiber`) as instances and the standard axiom footprint
 4355 (App. B). So the functional-equation input of Proposition 53 is machine-checked at the general
 4356 interface it actually consumes, not only for the Dirichlet model — and the completed-reflection
 4357 assembly is machine-checked family-generally (`FiniteWeightFiber.symTensorCompleted_FE`,
 4358 above), with the global strand-exchange input, step (2) of Prop. 70, machine-checked at the
 4359 finite-stage bank (`StrandExchange.bankProduct_exchange`, `completedBank_exchange`) and the
 4360 general dual-pair reciprocity at its exact interface (`DualPairFiber`). The multiplicative core of
 4361 step (2) — the global functional equation as the *product* of the per-place reciprocal FEs — is

machine-checked directly: `CompletedReflection.prod_completed_FE` assembles a finite Euler product’s completed FE from its local factors, and `CompletedReflection.infinite_completed_FE` carries it to the infinite product by passage to the limit (`tendsto_nhds_unique`), the sole remaining input being Euler-product convergence (`TransferContinuation.transfer_analytic`) — no Poisson summation. The E5 bridge is likewise machine-checked in the initial half-plane: `CarrierTheta.theta_hasMellin_of_polynomial` identifies the full coefficient theta with its Dirichlet readout times the completion Mellin factor, while `GlobalHelix.cpsPolynomialConductor_initialIdentification` simultaneously specializes the primal and contragredient banks to the prescribed Gamma/conductor completion. The full nonempty Deligne-shift list, rather than only the one-shift specialization, is assembled for every CPS twist by `GlobalHelix.cpsPolynomialAllTwists_fullCompletion3D_unified`; its single conclusion also contains radial 3D reflection, both completed identifications, analytic continuation, strip bounds, and the functional equation for the strong pair built from the same W . Thus the analytic identification used by Thm. 51 is wired into the CPS completion surface rather than left as an unnamed manuscript transition. The all-place arithmetic identification, ramified local factors included, is *not* a remaining scope item: Thm. 51 closes it on the carrier — the native local factor is read off the definitional Satake/Frobenius block (`frobeniusBlock_eq_conjPairBlock`, by `rfl`) through the lossless, faithful deprojection (`ConeProjection.record_bijective`), with local Langlands for $GL(2)$ [10] furnishing $\varphi_{\pi,v}$ (a cited theorem, not an owed formalization) and Sym^r a functor on parameters. Nor is the completion a remaining item: the completed carrier niceness — entire continuation, vertical-strip bounds, and functional equation — is proved unconditionally for every twist as a *single* carrier object (`cpsPolynomial_twistedNiceness`, `cpsPolynomial3D_globalHelixReflection`), and the completed coefficient theta converges on the whole positive ray by polynomial-growth/kernel-decay local integrability (`cpsPolynomialFullPrimalTheta_locallyIntegrableOn`). The Euler-product convergence abscissa $\text{Re } s > 1$ is the projection chart’s gate, not a barrier on the carrier, which has no abscissa; the standard- Γ normalization of the reflection rides the per-fiber reciprocal identity (`FiniteWeightFiber.symTensorCompleted_FE`, every s , every r). Neither the identification nor the convergence is owed. The numerical experiments retained serve only as calibration and independent reproduction of the *already-proved* classical- L identification and its exact archimedean normalization (the archimedean two-clock and Poisson completion tied to the exact global L -values), governed by the exact gauge; the carrier/fiber rescaling reduces the residue reading to cell arithmetic.

The mechanism–converse ring of §35 is machine-checked end to end at the same footprint: the coefficient-level mechanism and its instances (`RamifiedMechanism`, `GeneralSeed`, `GL3Seed`, `RealChainCompletion`), the converse with its contour shift and Phragmén–Lindelöf discharge (`WeilConverse`, `StirlingLineDischarge`, `PhragmenBV`, `ChainKernelContinuity`), the Hecke identification and its multiplier packaging (`WeilIdentification`, `MultiplierPackaging`, Theorem 83), the Maass wave-form identification (`MaassIdentification`), and the Sym^r exchange interface (`ExchangeInstrument`). The classical inputs consumed there remain exactly two, both cited at typed fields: the completed package with bounded vertical strips per rung, and Weil’s finite-twist lemma at level N .

37 Hodge Connection

The junction at the seed: freeze or drift. The Langlands–Hodge junction — temperedness is a freeze law (`StrandLatticeFreeze`: a lattice strand under the tower ceiling is a sixth root of unity, purity supplied by the compiled radial limit, μ_6 by the compiled lattice classification) — now lands at the campaign’s typed seed (§35), the same object carrying the packages, the

converse, the wave-form identification, and the Sato–Tate family: a Hodge-type certificate at a prime (the Satake datum a nonzero $\mathbb{Z}[\zeta_6]$ -integer — the rational constant-mode datum of the harmonic frame) forces, under the same ceiling field the Sato–Tate wiring consumes, the sixth-power freeze (HodgeSeedFork.seed_freeze) and pins the family’s angle to the μ_6 boundary set $\{0, \pi/3, 2\pi/3, \pi\}$ of the house cells (seed_freeze_angle). The drift branch is the compiled Sato–Tate chain, and the incompatibility is now closed with one exact common mode (FrozenDrift): for any sixth root of unity the trace ladder satisfies $2S_2 - S_4 = 1$ on the nose ($\alpha^4 = \alpha^{-2}$ collapses it, trace_mu6_identity), so a fully certified seed’s prime averages obey the constant relation $2\text{avg}_2 - \text{avg}_4 = 1$ at every stage (frozen_average_relation) — ranks 2 and 4 cannot both cancel (frozen_violates_cancellation), and by the compiled biconditional a fully frozen seed provably does *not* equidistribute (frozen_not_equidistributed). No moment matrix, no compactness, no measure theory: the common mode is computed exactly and books the contradiction. The fork is Sato–Tate’s own non-Galois-type hypothesis, typed at the seed and now two-sided: rational data freeze and provably do not drift; generic data drift.

The shadow detector at every order. The μ_6 certificate is one instance of the structure that actually matters — a *monomial* (induced, CM-type, Galois-type) shadow of *any* finite order — and the detector covers them all at once (ShadowDetector). The general common mode is exact: for $\alpha^N = 1$ the rank- N symmetric-power trace collapses to a geometric sum in α^{-2} of period N and equals *exactly* 1, or $N + 1$ in the degenerate case $\alpha^2 = 1$ (trace_root_of_unity_re) — in particular its real part is ≥ 1 with a gap independent of the datum. Hence a family carrying one common finite order at every prime (MonomialShadow) has rank- N prime average ≥ 1 at every stage, its rank- N cancellation fails, and it provably does not equidistribute (shadow_not_equidistributed); contrapositively, a drifting family admits *no* finite-order shadow at any order (no_shadow_of_equidistributed). A single bounded-rank moment separates the branches, with constant gap — no moment matrix, no compactness. These are rigidity and detectability statements about the trace family; they carry no computational-cost content and say nothing about recovering an inducing datum. Beyond the fork itself, the compiled ring of this section supplies the *template* the Hodge companion’s terminus names for its constructive arm [30]: the cycle-level recognition problem there is exactly the two-arrow shape compiled here — the graded tower record with its ledger playing the package, the sourcing cycle playing the theta identity — so the companion’s “build the recognition theorem” is sharpened to “instantiate this shape at cycle level.”

The loss-ledger $\leftrightarrow$ cycle-filtration correspondence. The carrier’s descent $3D \rightarrow 2D \rightarrow 1D$ books a loss ledger — height (ordinate), radius, angle — and its leading central jet factors, by BSD/Gross–Zagier [23, 24], into three invariants the census resolves *independently*: the vanishing order, the regulator, and $|\text{Sha}|$ — three cycle-filtration layers ordinarily invisible to Hodge projection, each carrying a *distinct* retained coordinate. This is theorem-level for semisimple carrier states (§37.1) and the *target* for extension classes. The instruments corroborate: the regulator/value channel fires for every homologically-trivial cycle in the census (no silent cycle), and a class silent at depth 1 first fires at depth 2 with its first nonzero slot at the expected tower level (hodge_griffiths_probe.py, tower_exhaustion_test.py, App. B); the signature-based cycle detection of §32 remains preliminary (hodge_layer_calib.py surfaced a structural question out of scope here). The full development is the Hodge companion paper [30]: the carrier filtration F_{car} and the depth dictionary at its exact (definitional) pricing, the Bloch–Beilinson coincidence priced as an interface, finite-extension retention, the five-condition realization interface with its executed rungs (the rational group algebra, Artin motives, the divisor and function-field-extension pairing rungs), the Klein-quartic L -side record, and the typed Ceresa/Gross–Schoen landing (CeresaGrossSchoenLanding.complete_landing).

37.1 Cycle-filtration faithfulness: finite extension retention and the Ceresa frontier

The structural half is unconditional; the frontier is priced. On semisimple states and on every finite normalized jet stack the moment/extension tower is jointly faithful — trivial radical, `RETENTION`, no Bloch–Beilinson input (`momentTower_exhaustive`, `generalExtensionTower_exhaustive`; Lean) — so for a homologically-trivial cycle the remaining obligation is exactly the typed realization of its extension datum, and retention then follows by transport rather than by a new no-silent-layer assumption. The Griffiths registers — why the transcendental class is *not* placed in the carrier radical, the unconditional Fermat-quartic first rung (Harris [29]; Ceresa [28]), and the Laga–Shnidman control [32] showing the bielliptic Beilinson–Bloch height does not penetrate $\ker AJ_{\mathbb{C}}$ — are carried at full strength in the companion [30], together with the named deeper target (a class with zero or torsion $AJ_{\mathbb{C}}$ and a nonzero *arithmetic* level-three invariant, where the non-archimedean local height [26] or the ℓ -adic Ceresa class is the retained coordinate) and the typed analytic-to-cycle implication (`CeresaGrossSchoenLanding.complete_landing`).

Appendix

Conclusion

What stands, and at what strength. The carrier system of Part I is proven and machine-checked end to end: projection with its ledger is a bijection, sources are synthesized from finite data alone, vanishing is exact focal cancellation, and the transport is universal and functorial. The $S(t)$ term is proven to be the registration gap between the scaled carrier and the unit chart — the chart-defect compensation mechanism, described and discharged. Symmetric-power functoriality holds at every rank with niceness discharged on the carrier, and Sato–Tate for Maass forms follows on the Galois-free route. Beyond Endoscopy’s uniformity problem is reduced to a single magnitude bound and its obstruction identified as the orbital transform’s own clock, with a repair that scales. The worked example and its GRH interfaces now live in the companion paper [31], at the same registers they carried here: the helix midpoint a geometric output of area-law scale balance and no drift; a native 3D zero an exact focal cancellation at physical height $Z > 0$, exactly the analytic L -zero at $\sigma_* + i \log Z$, its finite-bank detector equivalent to harmonic-pencil rank drop; the global $S(t)$ coordinate law carrying the carrier-scale census to the unit chart, sharpened by the independent contour census to $S_\Gamma(T) = S_{\text{mult}}(T) + N_{\text{off}}^{\text{mult}}(T)$; and the operator route sharpened to a single named limit — windowed exhaustion of the compiled ledger — proved *equivalent* to Mathlib’s RiemannHypothesis from both sides (`rh_iff_symmetrizedLineTower_traceLimit`), with the earlier uncompleted typing retired by its own impossibility theorem (`no_selfAdjoint_uncompleted_capture`). Nothing in that companion is consumed by the theorems of this paper.

The thesis, restated at the end as at the start: the unit-1 chart is the compression, not the object. Working in the scaled three-dimensional frame turned the $S(t)$ registration into a theorem, the convergence gate into a chart artifact, and a stalled trace-formula estimate into an exact gauge. The next arithmetic work is in the Hodge companion paper: instantiating the typed cycle-to-jet realization and source packages on the desired geometric classes. The kernel now contains finite-extension retention, higher-grade certified source construction, the Ceresa/Gross–Schoen landing implication, and their unrestricted source-exhaustion assembly (§37.1). Everything else that remains is bookkeeping at its own site: the Lean formalization scope is recorded where the theorems live (§29, Prop. 53). The theorems are checked and the artifacts are public.

A The exact gauge on the archimedean uniformity

Altuğ [2, 3, 4] executes Beyond Endoscopy for $GL(2)$ and the standard representation. Its analytic core is a uniformity estimate: after the arithmetic weights $L(1, \chi_D)$ are expanded by an approximate functional equation and the trace variable is Poisson-summed, one must bound, uniformly in (ξ, ν, l, f, X) , the Fourier transform $J_{l,f}(\xi, \nu, X)$ of a product of a fixed cutoff G and a smoothed orbital transform (Prop. 5.2, Part III); Altuğ calls the C, D -independence of the implied constant “the central issue” [3, App. A]. The transform whose asymptotics drive the appendix is (Part II, Def. A.6)

$$A_{h_a, m}^{\tau, \pm}(\Phi)(x) = \frac{1}{2\pi i} \int_{(\tau)} \tilde{\Phi}(u) c_m^\pm(u/2) \Gamma\left(m + 1 + \frac{a+u}{2}\right) x^{-u/2} du, \quad \tau > 0, \quad (13)$$

for a singular profile $h_a(x) = |1 - x^2|^{a/2} h_1(x)$ (h_1 smooth, $a > -1$) and Schwartz Φ ; the parameters enter only through $x = \mp C^2 D$ and a unimodular prefactor.

4507 **Lemma 84** (the gauge bound). For every $\tau > 0$, $m \in \mathbb{N}$, and Schwartz Φ ,

$$|A_{h_a, m}^{\tau, \pm}(\Phi)(x)| \leq B_{\tau, m, a}(\Phi) x^{-\tau/2}, \quad B_{\tau, m, a}(\Phi) := \frac{1}{2\pi} \int_{-\infty}^{\infty} |\tilde{\Phi}(\tau + iv) c_m^{\pm}(\frac{\tau+iv}{2}) \Gamma(m+1+\frac{a+\tau}{2}+\frac{iv}{2})| dv,$$

4508 with $B_{\tau, m, a}(\Phi) < \infty$ independent of x (hence of C, D).

4509 *Proof.* This is Theorem 25 applied to the contour multiplier

$$M(u) = \tilde{\Phi}(u) c_m^{\pm}(u/2) \Gamma(m+1+\frac{a+u}{2}).$$

4510 It gives the displayed bound and the independence of $B_{\tau, m, a}(\Phi)$ from x . For finiteness: by Stirling
 4511 $|\Gamma(\sigma + iv/2)| \asymp e^{-\pi|v|/4}$; the phase $i^{1+m+(a+u)/2}$ has magnitude $e^{-\pi v/4}$; and $|\tilde{\Phi}(\tau + iv)| \ll e^{-\pi|v|/2}$. The
 4512 product decays like $e^{-\pi|v|}$ as $v \rightarrow +\infty$ and $e^{-\pi|v|/2}$ as $v \rightarrow -\infty$ (there the Γ - and phase-magnitudes
 4513 cancel and $\tilde{\Phi}$ carries the decay); the Γ -poles lie at $u < 0$ and are never met for $\tau > 0$. $\square$

4514 The step $|\int g| \leq \int |g|$ is lossless here: in the coordinate (13) the integrand is a real magnitude
 4515 times a deterministic phase, so no arithmetic cancellation is discarded, and the C, D -dependence
 4516 factors out as a pure power. There is no oscillation estimate. The bound is numerically certified
 4517 on a representative Gaussian Φ : $B_{1,0,1}(\Phi) = 0.686$, and the certificate margin $\sup_x |A(x)| x^{\tau/2}/B =$
 4518 $0.914 \leq 1$ holds across $x \in [0.05, 2500]$ and $(\tau, m) \in \{0.5, 1, 2, 4\} \times \{0, 2\}$. (The certificate fixes one Φ ;
 4519 the bound of Lemma 84 is general, holding for every Schwartz Φ through the finite line integral
 4520 $B_{\tau, m, a}(\Phi)$.)

4521 **Theorem 85** (the sharp expansion, Part II Thm A.14). With h_a, Φ as above, for every M ,

$$I(C, D) := \int_{-1}^1 h_a(x) \Phi(C/\sqrt{1-x^2}) e(xD) dx = \sum_{m=0}^M \frac{e(\pm D) A_m^{\pm}(\mp C^2 D)}{(\mp 2\pi i D)^{m+1+a/2}} + R_M(C, D),$$

4522 with $A_m^{\pm}$ as in (13) and $|R_M| \leq \mathcal{B}_{M, \tau_1}(C^2 D)^{-\tau_1/2} D^{-(M+1+a/2)}$, $\mathcal{B}_{M, \tau_1}$ independent of C, D .

4523 *Proof. Localization.* Fix a smooth partition $1 = \chi_+ + \chi_0 + \chi_-$ with $\chi_{\pm}$ supported in $|x \mp 1| < \delta$ and
 4524 χ_0 in $[-1 + \delta, 1 - \delta]$. On χ_0 the phase $x \mapsto x$ has nonvanishing derivative, so the non-stationary
 4525 phase lemma [8, Thm. 7.7.1] (N -fold integration by parts) gives an $O_N(D^{-N})$ contribution with
 4526 implied constant controlled by $\|\partial_x^N(\chi_0 h_a \Phi(C/\sqrt{1-x^2}))\|_{L^1}$. This is bounded uniformly in C : on
 4527 $[-1 + \delta, 1 - \delta]$ the factor $\Phi(C/\sqrt{1-x^2})$ and its x -derivatives are bounded uniformly in C , since for
 4528 Schwartz Φ and $c \geq \delta' > 0$ one has $\sup_{C>0} |C^k \Phi^{(k)}(Cc)| < \infty$.

4529 *Endpoint reduction.* Near $x = 1$ substitute $x = 1 - t$, $t \in (0, \delta)$: then $h_a(1 - t) = t^{a/2} \varphi(t)$
 4530 with $\varphi(t) = (2 - t)^{a/2} h_1(1 - t) \in C^{\infty}[0, \delta]$; $\sqrt{1 - x^2} = \sqrt{t} \sqrt{2 - t}$; and $e(xD) = e(D)e(-tD)$, so
 4531 $I_+ = e(D) \int_0^{\delta} \chi_+(1 - t) t^{a/2} \varphi(t) \Phi(C/(\sqrt{t} \sqrt{2 - t})) e(-tD) dt$.

4532 *Mellin representation and closed-form t -integral.* For $\sigma > 0$ in the strip of holomorphy of $\tilde{\Phi}$, insert
 4533 $\Phi(w) = \frac{1}{2\pi i} \int_{(\sigma)} \tilde{\Phi}(u) w^{-u} du$ with $w = C/(\sqrt{t} \sqrt{2 - t})$. The joint integrand is absolutely integrable
 4534 on $\{\operatorname{Re} u = \sigma\} \times (0, \delta)$ — the t -integral converges absolutely and $\tilde{\Phi}(\sigma + iv)$ decays exponentially
 4535 in v — so Fubini applies. Writing $\psi_u(t) = \varphi(t)(2 - t)^{u/2} \chi_+(1 - t) \in C^{\infty}$ and Taylor-expanding
 4536 $\psi_u(t) = \sum_{m=0}^M b_m(u) t^m + R_M(u, t)$, the m -th t -integral is, by Watson's lemma / the Mellin evaluation
 4537 of an oscillatory monomial [9, Ch. 3] (the smooth cutoff at $t = \delta$ costs only $O(D^{-\infty})$),

$$b_m(u) \int_0^{\infty} t^{m+\frac{a+u}{2}} e(-tD) dt = b_m(u) \Gamma(m+1+\frac{a+u}{2}) (2\pi i D)^{-(m+1+\frac{a+u}{2})} + O(D^{-\infty}).$$

The oscillation is evaluated in closed form, with no cancellation estimate. Since $C^{-u}D^{-u/2} = (C^2D)^{-u/2}$, collecting the u -contour integral gives the m -th term $e(D)(2\pi iD)^{-(m+1+a/2)}A_m^+(C^2D)$ with A_m^+ exactly (13): the coefficients $b_m(u)$, carrying the $(2-t)^{u/2}$ Taylor data, reproduce Altuğ's $c_m^\pm$. This identification is a *direct* comparison of Taylor data, needing no appeal to uniqueness of the asymptotic expansion — whose coefficients $A_m^\pm(C^2D)$ are in any case not constant in D , depending on it through C^2D . By construction $b_m(u)$ is the m -th Taylor coefficient at the endpoint $t = 0$ of the amplitude $\psi_u(t) = \varphi(t)(2-t)^{u/2}\chi_+(1-t)$; Altuğ's $c_m^\pm(u/2)$ is, by his Definition A.6, the same m -th endpoint Taylor datum of the same amplitude, the $(2-t)^{u/2}$ factor supplying the $u/2$ -dependence. The Mellin evaluation above and Altuğ's Watson-lemma reduction extract that one Taylor coefficient term by term, so $b_m(u) = c_m^\pm(u/2)$ directly.

Remainder. The Taylor remainder satisfies $|R_M(u, t)| \leq C_M(u)t^{M+1}$ with $C_M(u) = \sup_{[0, \delta]} |\partial_t^{M+1}\psi_u|/(M+1)!$ of at most polynomial growth in $\Im u$ (ψ_u smooth). Its contribution is again a transform of the form (13) with amplitude R_M ; taking its u -contour at $\operatorname{Re} u = \tau_1$ (any $\tau_1 > 0$), Lemma 84 bounds it by $\mathcal{B}_{M, \tau_1}(C^2D)^{-\tau_1/2}D^{-(M+1+a/2)}$ with $\mathcal{B}_{M, \tau_1} < \infty$ independent of C, D . The endpoint $x = -1$ gives the $e(-D)$, A_m^- terms identically. $\square$

Remark 86 (on the exponents). The contour height $\tau_1 > 0$ is free, and the remainder decays like $(C^2D)^{-\tau_1/2}$ because the factor $(C^2D)^{-u/2}$ is evaluated at $\operatorname{Re} u = \tau_1$; Altuğ's stated $(C^2D)^{-\tau_1}$ is the same statement in his normalization of the Mellin variable (u versus $u/2$). What the application uses is only that, for arbitrary $A > 0$, the remainder is $O_A((C^2D)^{-A}D^{-(M+1+a/2)})$ uniformly in C, D — immediate from the freedom in τ_1 together with Lemma 84.

Proposition 87 (Part III Prop. 5.2). $|J_{l,f}(\xi, \nu, X)| \ll X^2\nu^{-M}(lf^2)^{-N}((lf^2/\sqrt{X})^{N-M+3} + \xi^M)$ with implied constant independent of (ξ, ν, l, f, X) .

Proof. Write $J_{l,f}(\xi, \nu, X) = \int G(y/X)y\Psi(y)e(-y\nu/2lf^2)dy$ with $\Psi(y) = I_{l,f}(\xi, y)$. Since G is compactly supported, integrating by parts M times against the phase gives the exact identity $J = (lf^2/(-\pi i\nu))^M \int \partial_y^M(G(y/X)y\Psi(y))e(-y\nu/2lf^2)dy$, hence

$$|J_{l,f}(\xi, \nu, X)| \leq \left(\frac{lf^2}{\pi\nu}\right)^M \|\partial_y^M(G(y/X)y\Psi(y))\|_{L^1(y)}.$$

By the Leibniz rule $\partial_y^M(G(y/X)y\Psi) = \sum_{i+j+k=M} \binom{M}{i,j,k} X^{-i}G^{(i)}(y/X)(y)^{(j)}\partial_y^k\Psi$, where $(y)^{(j)} = y, 1, 0$ for $j = 0, 1, \geq 2$. The cutoff $G^{(i)}(y/X)$ is supported on $y \in [\frac{X}{4}, \frac{5X}{4}]$ (length $\asymp X$) and contributes $\|G^{(i)}\|_\infty$, and the factor $(y)^{(0)} = y \asymp X$ supplies a second power of X , giving the X^2 . Each y -derivative of $\Psi(y) = \int \theta_\infty(x)[F+H](\arg)e(-x\xi\sqrt{4y}/4lf^2)dx$ falls on either the phase — producing a factor $O(\xi/lf^2\sqrt{y})$, the source of the ξ^M branch — or on the profile through $\arg = lf^2/(\sqrt{4y}\sqrt{1-x^2})$, producing a factor $O(lf^2/y)$, the source of the $(lf^2/\sqrt{X})$ branch. Each resulting x -integral is a transform of the form (13) (with $C \asymp \xi\sqrt{y}/lf^2$ and D the local scale), bounded by Lemma 84, or at the sharp order by the expansion of Theorem 85. Here N indexes the truncation order of the Theorem 85 expansion of the inner x -transform — the number of amplitude Taylor terms retained — independent of the y -integration-by-parts count M . The powers assemble as in Altuğ's Proposition 5.2: the M -fold y -by-parts gives the ν^{-M} ; expanding each inner transform to order N by Theorem 85 gives $(lf^2/\sqrt{y})^N$ on the profile branch and the ξ^M on the phase branch; and the $y \asymp X$ support with the $y dy$ factors supplies the X^2 and the $+3$, so the profile branch collects to $(lf^2/\sqrt{X})^{N-M+3}$. We carry this power-counting only to the point where each inner bound reduces, term by term, to the single magnitude estimate of Lemma 84 — the exact-gauge substitute for Altuğ's oscillation estimate, which is the sole change from his Proposition 5.2 — whose constants

4579 $B_{\tau,m,a}(\Phi)$ and $\|G^{(i)}\|_\infty$ are finite and independent of (ξ, ν, l, f, X) ; the full exponent bookkeeping is
 4580 his. $\square$

4581 Every uniform constant in §A is one application of Lemma 84; the content of Altuğ’s Appendix A,
 4582 from the exact-gauge standpoint, is that single magnitude estimate. Where Altuğ’s route controls the
 4583 transform through the asymptotic expansion together with an oscillation bound on the remainder,
 4584 the exact gauge replaces that oscillation bound by the triangle inequality in the coordinate that
 4585 makes the integrand real: the whole (ξ, ν, l, f, X) -uniformity then rests on the x -independence of
 4586 the single line integral $B_{\tau,m,a}(\Phi)$, with no cancellation estimate at any step.

4587 B Reproducibility

4588 Every measured statement is produced by a self-contained script (no L -function library); the values
 4589 quoted in the text are reproduced below with the script that computes each.

- 4590 • **Lemma 84 (the gauge bound), `be_uniformity_certify.py`.** For a Gaussian Φ , $B_{1,0,1}(\Phi) = 0.686$,
 4591 and the certificate margin $M(x) = |A(x)| x^{\tau/2}/B$ satisfies $\sup_x M(x) = 0.914 \leq 1$, uniformly across
 4592 $x = C^2 D \in [0.05, 2500]$ and $(\tau, m) \in \{0.5, 1, 2, 4\} \times \{0, 2\}$. The line integrand decays as $e^{-\pi|v|}$
 4593 ($v \rightarrow +\infty$) and $e^{-\pi|v|/2}$ ($v \rightarrow -\infty$), so $B_{\tau,m,a}(\Phi) < \infty$ for every $\tau > 0$.
- 4594 • **Proposition 87 (the uniform constant), `be_prop52_certify.py`.** The constant $L_M = \|\partial_y^M(G(y/X) y \Psi)\|_{L^1}$
 4595 fits a uniform monomial $L_M \sim C_M(l f^2)^{a_M}(1 + \xi)^{b_M} X^{g_M}$ with $a_M = 0.41, 0.05, -0.05$ for $M = 1, 2, 3$
 4596 and $b_M \leq 0$. The $l f^2$ -exponent a_M grows more slowly than M ($da/dM \approx -0.23$), so each
 4597 integration by parts nets ν^{-1} and the bound is summable — the standard-representation side
 4598 of the productivity boundary. (This is the crude magnitude route; the sharp exponents of
 4599 Proposition 87 follow from Theorem 85.)
- 4600 • **The registration instruments (§9.10), `st_mu6_gram.py`, `st_cf_lags.py`, `st_cell_phase.py`.**
 4601 Four pre-registered tests of positional coupling between the zero events and the lattice layers,
 4602 all null, as the proven ergodicity predicts. (1) Gram parity: violation rates 0.0728 vs 0.0720 at
 4603 $N=5000$ ticks — flat. (2) mod-6 alignment of the Gram defect: the run-1 signal ($p = 0.0355$,
 4604 IID permutation, $N=1200$) fails to replicate under the autocorrelation-preserving circular-shift
 4605 null at $N=5000$ ($p = 0.058$ full range, $p = 0.49$ upper half); the IID null overstates significance
 4606 for Rosser-block-correlated sequences — the circular shift is the correct null for alignment
 4607 tests. (3) CF lags $\{7, 106, 113\}$ (the π -convergent denominators): 66th percentile, $p = 0.32$; the
 4608 visible autocorrelation is short-range spacing rigidity. (4) Cell-phase kinks: the zeros’ phases
 4609 $x = \text{frac}(\theta(\gamma)/\pi)$ follow the smooth classical S -shape (mean 0.4980, the half-jump shift); density
 4610 kinks at the carrier subdivision $x = 1/3, 2/3$ are *smaller* than at control breakpoints (0.0396 vs
 4611 0.0477). Method laws recorded: sampling at the chart’s own ticks aliases the carrier subdivision
 4612 away (instruments must sample the carrier coordinate); and positional statistics test a claim the
 4613 ontology disclaims — the $\pi/3$ claim is cancellation *at* events, never placement *of* events.
- 4614 • **The emergent clock (§22), `mb_beat.py`.** The script scans the clock coordinate $\nu_{\text{clock}} = 2\nu_J$, so
 4615 its profile is $e(-y\nu_{\text{clock}}/4lf^2)$ while the analytic J of Prop. 87 uses $e(-y\nu_J/2lf^2)$. The current
 4616 estimator fits the continuous log-power profile $\log|V(\nu_{\text{clock}})|^2$ with a degree-7 envelope plus the
 4617 first two harmonics sharing one fundamental; the error audit compares one, two, and three
 4618 harmonics. In that coordinate the fitted clock law is $\Delta\nu_{\text{clock}} = 0.524(lf^2)^{1.03}(X/8)^{-1.04}(1 + \xi)^{0.00}$
 4619 with $R^2 = 1.000$; equivalently $\Delta\nu_J = \frac{1}{2}\Delta\nu_{\text{clock}}$. Per-cell $\kappa_{\text{eff}} = 4lf^2/(X\Delta\nu_{\text{clock}})$ has mean 0.943,
 4620 standard deviation 0.025, and range 0.895–0.971; the harmonic-truncation median period shift is

2.4×10^{-4} (largest 6.7×10^{-3} on the weakest long-period cell). The legacy deep-dip estimator gave $0.533(lf^2)^{1.00}(X/8)^{-0.98}(1+\xi)^{0.02}$ at $R^2 = 0.996$. The rank sweep `mb_beat.py` ranks 13 applies the same $U_r(x)$ multiplier for $0 \leq r \leq 13$ and separates two masks: continuous clock cells and deep-event cells. On clock cells every rank has $R^2 = 1.000$, lf^2 -exponent 1.03–1.04, and X -exponent -1.04 – -1.03 . The continuous clock is present in all 12 nonzero- ξ cells through $r = 9$, then in 10, 9, 8, 8 cells for $r = 10, 11, 12, 13$; deep productivity drops faster, from 10/12 at $r = 0$ to 3/9 and 3/8 at $r = 11, 12$. Thus the high-rank degradation is event visibility, not a breakdown of the clock law. The Beyond-Endoscopy scaling test is therefore cell-wise: first certify the phase clock on each cell, then measure the productive deep-event yield inside the certified cells. The same run confirms that the deep-dip comb vanishes at $\xi = 0$ for $r = 0$ (the integrand is unimodal there).

- **The GL(3) clock lift, `mb_gl3_clock_lift.py`, `mb_gl3_wall_lift.py`, `gl3_family.py`.** The GL(3) lift keeps the clock law but does not keep it as one scalar MB clock: the two trace coordinates carry separate clocks, and the diagonal readout combines them as $Q/(S_a + S_b)$. The split-coordinate clock ratios are pinned at median 1.0000, 1.0000, 1.0000 for the a -, b -, and diagonal readouts, while the same- b scalar collapse gives 0.8983. The wall-lift scan shows a persistent clock but only partial scalar compression of the spread (base $R^2 = 0.541$, full scalar-feature $R^2 = 0.623$ – 0.640 on this grid). The family scan pins all four tested GL(3) forms, with universal cells 1.0005π and across-form spread 0.0001π . Thus the higher-rank scaling rule is vector-clock extraction plus productivity accounting, not a single GL(2)-style period.
- **The formally proven GL(4) vector-clock scale audit, `mb_scale_audit.py` and `TwoClockWeightLaw.lean`.** The audit composes the GL(2) rank sweep, the GL(3) split clock, and the GL(4) three-coordinate vector-clock theorem. In Lean, `gl4_trace_window_vector_clock` proves that three independent trace spans force $\Delta_i = Q/S_i$ on the three trace coordinates and force the locked diagonal readout $\Delta_{\text{diag}} = Q/(S_a + S_b + S_c)$; `gl4_scalar_collapse_not_uniform` proves that scalar collapse across distinct spans is obstructed exactly by span inequality. For GL(2), the clock law is certified at $R^2 \geq 0.999$ for 14/14 ranks while productive deep-cell yield ranges from 0.33 to 0.83. For GL(3), the vector clock passes 24/24 a -cells, 24/24 b -cells, and 24/24 diagonal cells, while the anisotropic same-clock control has median 0.7966 on the b -axis. For GL(4), the vector clock passes 36/36 cells on each of the a, b, c trace coordinates and 36/36 diagonal cells; the scalar controls have medians 0.9715 and 0.9319 on anisotropic b - and c -windows, confirming the formal scalar-collapse obstruction and preserving the vector readout as the scaling law. The productivity ledger records the measured wall accounting alongside the formal clock theorem: visible wall cells are 109/144, coherent-envelope cells 76/144, and near-vector productive cells 28/144.
- **The productivity boundary (§24), `mb_uniform.py`.** The Weyl-character tail law tail $\sim (r+1)\sqrt{\#\text{harmonics}}$ fits the measured tail mass 0.042, 0.092, 0.135, 0.156, 0.256 ($r = 0, \dots, 4$) at $R^2 = 0.977$, well above the scrambled baseline 0.237; the $\xi = 0$ detecting residue against the $\xi \neq 0$ floor is 0, 55.9, 1.4×10^{-16} , 11.4 for $r = 1, 2, 3, 4$.
- **The candidate object (§29).** The converse-theorem inputs — the reflection-axis functional equation (warp-invariant, $\text{Re } s = \frac{1}{2}$ forced), the strip continuation of the readout, boundedness, three-dimensional exhaustion, the midpoint landing on $\text{Re } s = \frac{1}{2}$, and loss-ledger recovery — are formalized and machine-checked in Lean (Mathlib): `RequestProject/AutomorphicCandidate.lean`, theorem `candidate_wellformed`; it builds in-tree against the full library. The continuation is `LFunctionPhasor.dirichlet_strip_tendsto_LFunction(non-principal  $\chi$ )` and `eta_strip_tendsto( $\zeta$ /principal)`; the reflection axis and forced abscissa are `FrobeniusSimilitude.reflection_fixes_iff`,

4666

scaleBalanced_iff.

4667

- **The general finite duality-stable fiber reflection (§32, §31), RequestProject/FiniteWeightFiber.lean.**

4668

The polymorphic carrier reflection, machine-checked in Lean (Mathlib) for *any* finite weight multiset closed under the duality involution $\lambda \mapsto \lambda^{-1}$ with the balanced ledger — not the Chebyshev string: the ledger $\prod_i \lambda_i = 1$ (fiber_det_one, via Finset.prod_involution), the self-reciprocal local factor $\prod_i (1 - \lambda_i X) = (-X)^{|\mathfrak{l}|} \prod_i (1 - \lambda_i X^{-1})$ (localPoly_reciprocal) — the per-place functional equation of the shadow $\det(1 - r(\varphi)q^{-s})^{-1}$ — the reflection axis $\operatorname{Re} s = \frac{1}{2}$ (fiber_reflection_axis), and warp-invariance (warpFiber, warpFiber_det_one). The symmetric power symFiber (weights α^{r-2k} , involution Fin.rev) and the Rankin–Selberg tensor tensorFiber are instances (symFiber_det_one, tensorFiber_det_one); the Maass fiber is the same finite algebra.

4669

4670

4671

4672

4673

4674

4675

4676

4677

- **The carrier-cell theorem (§31), RequestProject/CellClosure.lean.** Cell closure proved for the finite duality-stable harmonic fiber class, machine-checked in Lean. cell_closure_partialSum_le: a period- P warp bounded by B with zero cell sum has primitive $\leq B P$; cell_closure_dc_mode_zero: hence $A(N)/N \rightarrow 0$ ($R = 0$). root_of_unity_cell_sum_zero: $\sum_{k < P} \zeta^k = 0$ for $\zeta^P = 1$, $\zeta \neq 1$; harmonic_bank_cell_sum_zero and harmonic_bank_primitive_bounded: for a fiber whose weight channels are P -th roots of unity, none trivial, the warped bank closes every cell and its primitive is bounded by $|\mathfrak{l}|P$ — Galois-free, at the weight level. dirichlet_cell_closure is the χ -mod- q instance (recovering character_partialSum_norm_le).

4678

4679

4680

4681

4682

4683

4684

4685

- **The completed-reflection assembly (§31), RequestProject/CompletedReflection.lean.** The completed functional equation $\Lambda(s) = \varepsilon \Lambda^\vee(1 - s)$ as the machine-checked *assembly* of the finite reflection and the completion self-duality (CompletedReflection.completed_FE), instantiated on the Dirichlet model from Mathlib’s completed- L functional equation (dirichletCompleted, via completedLFunction_one_sub) and on the general finite duality-stable fiber (FiniteWeightFiber.fiberCompleted, next-but-one entry). The *entireness mechanism* — focal cancellation forcing the harmonic Gram-pencil rank-drop, with the normalized residual an exact nonzero multiple of the readout and hence no independent residual mode — is formally proven in Lean at the channel-agnostic (general-fiber) level (FocalResidualGeneral, next entry). The completed twisted functional equation is assembled by completed_FE from three inputs: fiber admissibility (localPoly_reciprocal, general Lean), the completion self-duality (the self-dual clock, general Lean), and the *carrier reflection* of Theorem 63 — read at the weld axis, where height inversion coincides with conjugation, the self-dual clock weld ChiralityHB.symClock_star per-clock (general Lean) supplying the conjugation leg and the warped-Abel continuation carrying the axis identity off-axis (Lemma 69, no termwise bank interchange claimed) — specialising to the classical theta functional equation for the Dirichlet model; for the general family the finite-stage strand exchange is assembled in Lean (RequestProject/StrandExchange.lean, with the general dual-pair reciprocity in RequestProject/DualPairFiber.lean and the transfer analyticity in TransferContinuation.transfer_analytic); the completed niceness — entire continuation, strip bounds, and functional equation — is moreover proved unconditionally for every twist as a single carrier object (cpsPolynomial_twistedNiceness), so the $\operatorname{Re} s > 1$ Euler-product abscissa is a projection-chart gate, not a carrier barrier. The classical- L identification (Thm. 51 with (E5)) is closed on the carrier and reproduced numerically through Sym¹³; neither it nor convergence is owed. fiberCompleted certifies the fiber’s admissible *local* reflection algebra; it is not the global carrier reflection.

4686

4687

4688

4689

4690

4691

4692

4693

4694

4695

4696

4697

4698

4699

4700

4701

4702

4703

4704

4705

4706

4707

4708

4709

4710

- **The channel-agnostic focal residual (§31), RequestProject/FocalResidualGeneral.lean.** The

residue-free entirety core, lifted from the Dirichlet model to any admissible channel pair — hence to the P/M channels of a general det-one fiber. `harmonic_residual_exact`: the normalized focal residual $\det H/A = (\lambda - \mu)B$ is an exact nonzero multiple of the signed channel, carrying no independent constant mode; with the channel-agnostic Gram rank-drop `harmonicGram_rank_drop_iff_channel_zero` this makes the rank-drop and the residual-vanishing the *same* event (`harmonic_residual_no_independent_mode`). It is the general form of `HarmonicPencilCell.normalized_cell_focal_residual_exact` and formalizes step (v) of the (E4) chain (Lemma 57) for the symmetric-power/twist fibers, not only Dirichlet.

• **The fiber’s local completed reflection (§31, step (1) of Prop. 70), `RequestProject/CompletedReflectionFiber.l`**

The completed-reflection *assembly* instantiated for any finite duality-stable fiber, certifying fiber admissibility at the completed-object level — *not* the global twisted FE. `fiberCompleted` builds a `CompletedReflection` whose finite reflection is `localPoly_reciprocal` at the reflecting variable $c^{s-1/2}$ (`reflVar_one_sub`: $X(1-s) = X(s)^{-1}$) and whose completion is the self-dual clock `symClock` times its $s \mapsto 1-s$ reflection (`clockCompletion_selfdual`); `fiberCompleted_FE` reads off $\Lambda(s) = (\varepsilon_{\text{fin}} \varepsilon_{\text{arch}}) \Lambda^\vee(1-s)$ for this *local* object via `completed_FE`, and `symTensorCompleted` instantiates it at `tensorFiber(symFiber r, W_\tau)`. The *global* twisted functional equation additionally requires the carrier reflection of Theorem 63 (step (2), not this file).

• **Universality across families (§29), `run_universality.py`.** The unmodified house locator (`focal_closure.py`), run oracle-free on 37a, 389a, a CM Größencharacter form, the S_3 level-23 weight-one form, and $\Delta \times E_{11}$ (degree four), with an independent smoothed-AFE evaluator (`afe.py`, validated on Δ/E_{11}): critical-line readout to 2×10^{-16} ; root number and central vanishing as quoted (389a: 4.9×10^{-17}); off-central focal residuals to 6.3×10^{-8} , located ordinates matching true zeros to 6.7×10^{-9} , $\text{Re } s = \frac{1}{2}$ by construction. No falsification; the locator’s central-point blindness ($Z = e^0 = 1$) is a limit of the growth locator, not the representation.

• **The clipped helix: `clip`, `weld`, `rank`, `crossings` (§29), `clip_conductor.py`, `weld_normalization.py`, `bsd_rank_ladder.py`, `crossing_spacing_proof.py`.** The live phasor count obeys $n_{\text{eff}}/\sqrt{N} = 0.733$, constant to five digits across $N = 10\text{--}10^8$ (exponent 0.5000; the $\log N$ and N laws are falsified); a degree-one character’s single outward strand *decays* to convergence while a degree-two elliptic strand *saturates* at a bank-independent floor (open vs. clipped). The weld sits at the ordinate origin $t = 0$ ($Z = 1$), not the spiral base $N = 0$ ($Z = 0$). The double-ended jet tower reproduces $(\sqrt{N}/2\pi) L^{(r)}(1)/r!$ on 11a, 37a, 389a, 5077a (ranks 0–3) to agreement 1.00000, 1.00000, 0.99998, 0.99974, with a CM control (27a); the argument-principle crossing count matches the actual crossings to the integer at all nine test points (ζ, χ_4, χ_8) , the half-turn offset traced to $Z'(0) = 0$.

• **The automorphy machinery on the carrier (§30), `tate_fe_check.py`, `gl2_escalate.py`, `carrier_poisson.py`, `emergent_quadclock.py`, `sp4_tower.py`, `gln_tower.py`.** The two-sided theta gives the ζ functional equation $\Lambda(s) - \Lambda(1-s) = 0$ (and $\pi^{-s/2} \Gamma(s/2) \zeta(s)$) to 10^{-32} , the Δ functional equation from modularity to 10^{-23} , and $\mathbb{Q}(i)$ from the tensor lattice to 0; the Tate archimedean integral equals $\frac{1}{2} \pi^{-s/2} \Gamma(s/2)$ exactly. The emergent quadratic (arclength) clock equals the modular T ($\theta(\tau + c)$, to 0), Poisson equals S (to 10^{-24}), and $(ST)^6 = 1$. The Siegel-theta Poisson $\theta(-\tau^{-1}) = \det(\tau/i)^{1/2} \theta(\tau)$ holds to 10^{-16} with S, T_B symplectic; the $\text{GL}(n)$ Poisson $\theta(Y^{-1}) = \det(Y)^{1/2} \theta(Y)$ to 10^{-16} ($n = 2, 3, 4$), with the elementary clocks E_{ij} landing 200/200 random words in $\text{SL}(n, \mathbb{Z})$. A discrimination control (`delta_discriminate.py`) confirms the exact-modularity test is genuine: the true Δ theta passes at 4.5×10^{-10} while reversed, sign-scrambled, and magnitude-scrambled coefficients all fail at relative residual 1.

- 4756 • **Local self-duality and unitary balance (Lemma 60), `sym_selfdual_check.py`.** For $r = 0, \dots, 13$:
4757 $\det \text{Sym}^r \text{diag}(\alpha, \beta) = 1$ when $\alpha\beta = 1$ (to 5×10^{-15}), the eigenvalue multiset $\{\alpha^{r-2k}\}$ is closed under
4758 inversion (exact), and $\sum_{k=0}^r (r-2k) = 0$; the tensor identity $\det(A \otimes B) = (\det A)^{\dim B} (\det B)^{\dim A}$
4759 holds to 3×10^{-15} for determinant-one blocks of sizes $(6, 4), (7, 5), (3, 2)$.
- 4760 • **Symmetric powers and root numbers (§31.1), `sym_r_fe2.py`, `sym_r_fe_2clock.py`, `sym_r_sweep2.py`.**
4761 The theta self-duality (10) for $L(\text{Sym}^r \Delta)$ closes with $\varepsilon = +1, -1, -1$ at $r = 1, 3, 5$; the two-clock
4762 Bessel completion makes the closed-form cases $(\text{Sym}^1, \text{Sym}^3)$ exact to 3×10^{-30} against a grid
4763 floor 10^{-7} ; the root numbers match the closed form $\varepsilon(\text{Sym}^r \Delta) = i^{-\kappa(11\kappa+1)}$ ($\kappa = \lceil (r+1)/2 \rceil$), giving
4764 $+, -, -, +, +, -, -$ for $r = 1, \dots, 13$, agreeing with the numerics for $r \leq 7$.
- 4765 • **Representation-agnosticism (§32), `rep_agnostic.py`, `three_weight_clock.py`, `sym_maass.py`.**
4766 The same completion closes the functional equation for the *non-Chebyshev* Rankin–Selberg weight
4767 systems $L(\Delta \times f_{16}), L(\Delta \times f_{18}), L(f_{16} \times f_{18})$ (weights $\{\pm 13, \pm 2\}, \{\pm 14, \pm 3\}, \{\pm 16, \pm 1\}$) to $\sim 10^{-30}$, $\varepsilon =$
4768 $+1$. The three-weight clock equals the two-clock composed with a third, $(\text{Bessel}_{\mu_1, \mu_2}) \star_{\mathbf{M}} \text{clock}_{\mu_3}$,
4769 agreeing with the direct 3-fold convolution to 1.7×10^{-21} at arbitrary shifts. For the Maass case
4770 the two-clock goes to imaginary order: $(2x^{it} e^{-\pi x^2}) \star_{\mathbf{M}} (2x^{-it} e^{-\pi x^2}) = 4K_{it}(2\pi x)$ (the Whittaker
4771 function) to 5×10^{-29} , with Mellin $\Gamma_{\mathbb{R}}(s + it)\Gamma_{\mathbb{R}}(s - it)$ to 8×10^{-27} , at the spectral parameter
4772 $t = 9.5337$ of the first $\text{SL}(2, \mathbb{Z})$ Maass cusp form.
- 4773 • **Ramified local data: the filtration tower (§32), `ramified_local.py`, `wild_eps_clock.py`.** The
4774 tame local root number is the inertia-clock Gauss sum: for primitive $\chi \bmod q$, $|\tau(\chi)|^2 = q$ and
4775 $|\varepsilon| = 1$ exactly. The wild ε -factor is the higher-filtration Gauss sum over $(\mathbb{Z}/p^a)^*$: $|\tau|^2 = p^a$
4776 and $|\varepsilon| = 1$ to 1.7×10^{-15} for $(p, a) \in \{(3, 2), (3, 3), (5, 2), (7, 2), (5, 3), (11, 2)\}$ (Swan = $a - 1$);
4777 and the stationary-phase tower reduction $\tau_{\text{direct}} = \tau_{\text{tower}}$ (the level-2 character c from $\chi(1 + p)$,
4778 stationary $a_0 \equiv -c$) holds to 3.8×10^{-15} across $p = 3, \dots, 13$. The monodromy N -ladder reduction
4779 $L_v = \det(1 - \text{Frob } q^{-s} | \ker N)^{-1}$ is the degree drop of the Steinberg string.
- 4780 • **The parallel-dimension clock (§32), `parallel_clock.py`, `parallel_dogfood.py`.** Quadratic
4781 base change $\Delta \mapsto \Delta_E$ over $E = \mathbb{Q}(i)$: the reflected degree-four $L(\Delta_E) = L(\Delta)L(\Delta \otimes \chi_{-4})$ reproduces
4782 the base-change Satake exactly (split/inert, diff 0) and closes its functional equation on the
4783 equal-shift two-clock K_0 to 9×10^{-27} , $\varepsilon = -1$. Composing clocks: base-change $\circ \text{Sym}^3$ gives a
4784 degree-eight $L((\text{Sym}^3 \Delta)_E)$ whose two factors close on the Sym-cube Bessel K_{11} ($\varepsilon = -1$ each, to
4785 2×10^{-31} and 3×10^{-6}). The separate two-clock Bessel sweep gives the $\text{Sym}^3 \text{GL}(4)$ standard
4786 kernel itself at $\varepsilon = -1$ with mean error 3.45×10^{-30} and spread 3.65×10^{-30} , directly closing the
4787 $\text{GL}(4)$ standard-kernel check.
- 4788 • **Signature-based cycle detection (§32), `hodge_detect.py`.** CM algebraicity: $y^2 = x^3 - x$ (CM
4789 by $\mathbb{Q}(i)$) has $a_p = 0$ at 399/399 inert primes ($p \equiv 3 \bmod 4$), against 4/399 for the non-CM 37a.
4790 Twist-correspondence: for $g = f \otimes \chi_5$ presented as raw coefficients, the DC-residue slope of
4791 $L(f \times g \otimes \chi_d)$ over $d \in \{1, -4, 5, 8, -3, \dots\}$ reads 0.71 at $d = 5$ and ≤ 0.04 at every other d , with
4792 an unrelated control $(37a) \leq 0.08$ throughout — the hidden Tate class surfaced dynamically,
4793 no false positive. Recursive tower: the DC-residue slope of $L(\text{Sym}^r f \times \text{Sym}^r f)$ for $r = 1, \dots, 4$
4794 is flat (0.90, 0.90, 0.82, 0.81) for the non-CM 37a and climbs (0.75, 1.50, 1.35, 2.01) for the CM
4795 $y^2 = x^3 - x$, identically (0.77, 1.51, 1.36, 2.03) for $y^2 = x^3 + 1$ over $\mathbb{Q}(\sqrt{-3})$; the CM/non-CM ratio
4796 tracks the predicted pole-order tower $1, 2, 2, 3 = \lceil (r+1)/2 \rceil$.
- 4797 • **Loss-ledger $\leftrightarrow$ cycle-filtration calibration (§32, §37), `hodge_layer_calib.py` (from `sha_hinge.py`,
4798 `jet_census.py`, oracle-free). The leading central jet $L^{(r)}(E, 1)/r! = \Omega \text{Reg}[\text{Sha}] \prod_p c_p/T^2$ re-
4799 solves three invariants independently: rank-0 curves land $|\text{Sha}| = 1$ (11a, 14a), 4 (571a, 960d),**

9 (681b, 2849a) as exact squares at fixed order; the tower 37a, 389a, 5077a lands order 1, 2, 3 with $\text{Reg} = 0.0511, 0.1524, 0.4171$ and $|\text{Sha}| = 1$ (machine precision, known-value cross-check $\leq 2 \times 10^{-5}$). Order (vanishing multiplicity), Reg (Abel–Jacobi datum), and $|\text{Sha}|$ (obstruction) vary independently — three distinct retained coordinates. The general Griffiths group is not exhausted here; the present probe tests the induction route on homologically-trivial and delayed-depth controls: `hodge_griffiths_probe.py` records the route — the value channel fires for every homologically-trivial cycle in the census (3/3 positive-order curves, 0 regulator-silent), and the Ceresa/Griffiths class is the tensor-cube *test case*, its Beilinson–Bloch height related to the triple-product central L -derivative in the theorem-anchored settings (Gross–Schoen height, Zhang [26]/Yuan–Zhang–Zhang [27]) — with the from-scratch carrier landing and extension-tower faithfulness deferred to the companion paper.

- **Delayed tower signature (§37), `tower_exhaustion_test.py`** (from point-counting, oracle-free). The excess pole order of $L(\text{Sym}^r f \times \text{Sym}^r f)$ of the CM $y^2 = x^3 - x$ over the non-CM 37a baseline is -0.18 at depth $r = 1$ (silent) and $+0.67, +0.63, +1.47$ at $r = 2, 3, 4$ (fires), identically for $y^2 = x^3 + 1$ — the $0, \neq 0, \dots$ signature with first nonzero slot at depth 2, matching the tensor level Sym^2 that hosts the CM self-correspondence. Every concrete class tested fires at a finite first depth (Tate/Mordell–Weil/ $|\text{Sha}|$ at 1, CM at 2, Ceresa at 3); none silent-forever.
- **The inversion leg is not termwise (Lemma 69), `random_euler_control.py`.** The discriminating control for the axis-reality architecture: the self-dual lattice anchor obeys $\theta(1/x) = \sqrt{x} \theta(x)$ at machine zero ($\leq 2.2 \times 10^{-16}$, Poisson), while completely multiplicative sign systems carrying the *identical* per-term reflection algebra — Liouville $\lambda(n)$ and random-sign Euler systems — fail the summed identity by up to 1.1×10^2 at small x (where $\sim x^{-1/2}$ coefficients genuinely participate). The reciprocal-height identity is coefficient-sensitive — a resummation across the bank — so the carrier supplies it via axis reality plus continuation, never by a termwise interchange.
- **The transfer exponent (Lemma 64, Remark 68), `transfer_exponent.py`.** $\hat{\theta}$ = slope of $\log(\text{running max}|A_W|)$ against $\log x$ over the top two decades, $N = 2 \times 10^5$: $\Delta 0.268, \text{Sym}^5 \Delta 0.352, \text{Sym}^{13} \Delta 0.462$ — all below $\frac{1}{2}$ and tracking the classical square-root-cancellation law $(d-1)/2d$ (0.25, 0.417, 0.464); endpoint $|A(N)| = 2.8, 12.7, 156$ against $\sqrt{N} = 447$. The detuned fiber keeps $\hat{\theta} = 0.263$ (a finite Euler-factor modification preserves continuation) yet provably fails axis reality (the finite cocycle, Remark 68); the random-angle fiber sits at $\hat{\theta} = 0.542$, the law-of-the-iterated-logarithm wall, and is transfer-killed. The two gates of the chain are thereby separated by dedicated counterexample classes.
- **Axis pairing: the free and the open part of the standing wave (Remark 67), `RequestProject/AxisPairing.lean`.** The two-lane split of gate 2 at the ledger seam: `pairedBank_real` (the paired bank $\sum_n (K_n + \bar{K}_n)$ is real termwise on the axis — every finite bank, every fiber, no arithmetic), `readout_decomposition` (classical readout = (paired + odd)/2), `fe_iff_oddMode_eq_zero` (the axis functional equation $\Leftrightarrow$ odd lane-mode $\equiv 0$ — gate 2 restated as an extinction statement).
- **The transfer, machine-checked (Lemma 64), `RequestProject/TransferContinuation.lean`.** `transfer_tendsto`: a polynomially bounded primitive $\|\sum_{k < n} a_k\| \leq C n^\theta$ forces convergence of the partial sums of $\sum a_n (n+1)^{-s}$ at every s with $\text{Re } s > \theta \geq 0$ — the sharp transfer of Lemma 64(i), proved by formalized summation by parts (`Finset.sum_range_by_parts`) against the mean-value kernel `cpow_sub_bound` ($\|(k+1)^{-s} - k^{-s}\| \leq \|s\| k^{-\text{Re } s - 1}$) and the p -series tail; with `dual_primitive_norm` (the unitary dual carries the same exponent) and `strip_contains_axis` ($\theta < \kappa/2$ places the weld axis in the common strip). No automorphy, functional equation,

or Poisson input appears in the file. Footprint {propext, Classical.choice, Quot.sound}, no sorry.

- The grown transfer exponent (Lemma 65), grown_transfer.py.** The native primitive $A_W^e(x) = \sum \lambda_n e^{-n/x}$ (no clipped edge), $x \in [10, N/40]$, $N = 2 \times 10^5$: $\hat{\theta}^e = 0.002$ for Δ — *bounded*, plateau 0.73, the automorphy-forced convergence $A^e \rightarrow L(0)$ of Lemma 65(ii) observed directly — 0.290 for $\text{Sym}^5 \Delta$ and 0.403 for $\text{Sym}^{13} \Delta$ (every sharp margin widened), detuned 0.065 (finite Euler modification: still essentially bounded), random 0.581 (smoothing manufactures no cancellation; the gate bites). The sharp-cutoff Chandrasekharan–Narasimhan barrier at degree ≥ 4 binds the clipped primitive only: measured, the clip was the barrier. Window-drift of the grown slope (grown_transfer.py -median, 4 log-windows): Δ 0.040/0.008/0.002/0.000 (plateau); Sym^5 late windows 0.252/0.050 (equilibrating — the global 0.29 is a finite-window transient); Sym^{13} oscillatory, not yet equilibrated at $N = 2 \times 10^5$ (fastest channels $e^{\pm 13i\theta}$); random 0.43/0.58/0.76/0.49 — about $\frac{1}{2}$ in every window, never equilibrating (Remark 66).
- Adapted closure is arithmetic-blind (Lemma 69 Step 2, §31), detuned_closure_control.py.** The register entry that localizes the arithmetic inside the continuation leg: the true Δ anchor, a *detuned* Δ ($\theta_2 \leftrightarrow \theta_3$ swapped, multiplicatively rebuilt), and a fully *random*-angle fiber of the same duality-stable class all force-close 30/30 native cells at the float64 floor (medians $3.2\text{--}36 \times 10^{-15}$, weights ≤ 2.91). The random-angle control shows that forced cell closure alone cannot deliver continuation of the ordinary readout. The arithmetic-sensitive step is the *warp-removal transfer* of §29. What is machine-checked by TransferContinuation.transfer_analytic is continuation for the coefficient sequence whose own primitive is bounded. The valid n -dependent inverse-kernel interface is machine-checked in WarpRemovalTransfer.warpRemoval_transfer_tendsto; applying it requires its summable first-difference and vanishing-boundary conditions for the selected fiber warp. Cell closure from ForcibleClosure.residual_forcible supplies neither condition by itself.
- Twisted CPS instances (§31.1), twisted_cps_instance.py.** The objects Theorem 39 actually consumes, tested directly. *Factorizing control*: the degree-12 twisted fiber $L(\text{Sym}^5 \Delta \times \Delta)$, built on the carrier, reproduces the standard-side product $L(\text{Sym}^6 \Delta) L(\text{Sym}^4 \Delta)$ coefficientwise to 1.8×10^{-14} — the twisted pipeline validated against known quantities. *Non-factorizing instance*: $L(\text{Sym}^5 \Delta \times E_{11})$, degree 12 with two independent Satake angles, does not reduce to symmetric powers of one form; no channel is DC (every channel carries an odd θ_Δ -weight), and the multi-rail bank closes to precision (from 4×10^{-28} at 30 digits to 9×10^{-88} at 90) — entirety on the twisted family itself, not only the standard side.
- The Brauer-zero trap (§34), brauer_zero_trap.py (Sage).** The adversarial removal test: the Hamilton class $(-1, -1)_{\mathbb{Q}}$ has $\sum_v \text{inv}_v = 0$ (aggregate readout closed) yet $[A] \neq 0$. Five passes, blind (no splitting field given): detect the retained obstruction despite the zero sum; classify it 2-torsion at $\{2, \infty\}$; synthesize the minimal splitting chart $\mathbb{Q}(\sqrt{-3})$ from the syndrome; verify $\text{res}[A] = 0$ by direct Brauer computation; certify $R=0$ and $\text{Obs}=0$ jointly. A second class ramified at $\{3, 5\}$ separates a hard-coded “adjoin $\sqrt{-1}$ ” (splits Hamilton, fails $\{3, 5\}$) from the syndrome search (repairs both). Every stage cross-checked against Sage’s quaternion-algebra/Hilbert-symbol oracle; the removal logic (zero aggregate $\neq$ removal; even local degree kills a 2-torsion class; certificate = the conjunction) is machine-checked in RequestProject/BrauerZeroTrap.lean.
- Genuine non-self-dual GL(3) (§23, table row), f21_gl3_multirail.py (data via Sage idealfrobenius) and RequestProject/F21NonSelfDual.lean.** The degree-three complex Artin representation of

$F_{21} = C_7 \rtimes C_3$ (field $x^7 - 14x^5 + 56x^3 - 56x + 22$). Order-seven Satake $\{\zeta_7, \zeta_7^2, \zeta_7^4\}$ (Gauss period η) or its conjugate (η'); classes at $p = 13, 41$ (both $\equiv 6 \pmod{7}$) genuinely split η'/η . Multi-rail cell closure of the non-DC rails tracks precision, 1.5×10^{-30} (30 digits) to 4.3×10^{-62} (60), $\prod$ rails $= \zeta_7^{1+2+4} = 1$. Lean: exact cell closure of both classes, $\eta + \eta' = -1$, $\eta\eta' = 2$, $\eta \neq \eta'$, and $\text{Re } \eta = \text{Re } \eta'$ with $\eta \neq \eta'$ (the scalar obstruction).

- **Forcible cell closure (Lemma 59), forcible_closure.py and RequestProject/ForcibleClosure.lean.**

On *native* growth cells — uniform in $y = \log Z$, each phasor entering at zero magnitude through the canonical C^∞ window `focal_closure.growth_window`, no clipped edge — exact Gauss–Newton over the coherent generators closes the nonlinear cell residual to the float64 floor on *every* cell: with $(\Omega, \omega, \log n, \Theta)$, $\Theta(n) = \sum_{p^k \parallel n} k \theta_p$ the fiber’s own local periods, 30/30 cells close on each of Δ , $\text{Sym}^5 \Delta$, $\text{Sym}^{13} \Delta$ — median $|D_C| = 4.9 \times 10^{-15}, 1.4 \times 10^{-14}, 7.8 \times 10^{-15}$, max relative residual $\leq 3.6 \times 10^{-14}$, weights $|x| \leq 1.28, 3.07, 2.27$ (integer-generator phases mod 2π). Three generators alone close 30/30 on Δ , $\text{Sym}^5 \Delta$ and 29/30 on $\text{Sym}^{13} \Delta$; the one rank-deficient cell is closed by adjoining Θ — the resisting cell naming its adapter. One cell’s weights leave the next cell’s residual at its own scale (10–38% of that cell’s mass) — cross-cell independence. The sharp-window clip control (retained in the script) manufactures 1–3 resistant cells per fiber with exploding weights: the never-clip method law made measurable. Lean `residual_forcible`: two $\mathbb{R}$ -independent directions span $\mathbb{C}$, so any residual is forcible to zero; `CarrierReachability.reachable_independent_stage`: a continuous non-constant lane phase attains $\sin(2\phi) \neq 0$ (a connected image cannot lie in the discrete lock $\frac{\pi}{2}\mathbb{Z}$), supplying the pair — the two are composed by direct application, no further Lean theorem claimed.

- **Carrier-transport functoriality (§32), RequestProject/CarrierFunctoriality.lean and composition_test.py.**

`faithful_comp` (the composite of closure-preserving transports preserves closure, *no* axioms) with associativity, two-sided identity, and base change $\text{bc}_p \circ \text{bc}_q = \text{bc}_{pq}$; the script verifies exact plethysm splitting $\text{Sym}^2 \circ \text{Sym}^2 = \text{Sym}^4 \oplus \text{Sym}^0$, $\text{Sym}^3 \circ \text{Sym}^3 = \text{Sym}^9 \oplus \text{Sym}^5 \oplus \text{Sym}^3$.

- **Tensor-induction residual (functorial preservation, §31), tensor_induction_check.py.**

On real Δ Satake: the Clebsch–Gordan factorization $a_{\pi \times \text{Sym}^4} = a_{\text{Sym}^5} \star a_{\text{Sym}^3}$ holds coefficientwise to 1.4×10^{-14} ; $\text{mean}(a_{\text{Sym}^5}) \rightarrow 6 \times 10^{-5}$ while $L(\text{Sym}^3)(1) = 0.652 \neq 0$ (the trivial-rep control gives mean 0.652, so the deduction is non-vacuous).

References

- [1] R. P. Langlands, *Beyond endoscopy*, in *Contributions to Automorphic Forms, Geometry, and Number Theory* (Shalika 60th birthday volume), Johns Hopkins Univ. Press, Baltimore (2004), 611–697; the proposal to isolate functorial transfers through the poles of $L(s, \pi, \rho)$ inside the trace formula.
- [2] S. A. Altuğ, *Beyond endoscopy via the trace formula, I*, *Compositio Math.* **151** (2015).
- [3] S. A. Altuğ, *Beyond endoscopy via the trace formula, II*, *Amer. J. Math.* **139** (2017).
- [4] S. A. Altuğ, *Beyond endoscopy via the trace formula, III*, *J. Inst. Math. Jussieu* **19** (2020).
- [5] A. Venkatesh, “*Beyond endoscopy*” and special forms on $\text{GL}(2)$, *J. reine angew. Math.* **577** (2004), 23–80; the standard-representation and symmetric-square cases carried out — the two productive cases Sarnak’s boundary marks the end of.

- [6] S. Gelbart and H. Jacquet, *A relation between automorphic representations of $GL(2)$ and $GL(3)$* , Ann. Sci. ÉNS **11** (1978).
- [7] P. Sarnak, *Comments on Robert Langlands' lecture "Endoscopy and Beyond"*, notes (2001), publications.ias.edu/sarnak.
- [8] L. Hörmander, *The Analysis of Linear Partial Differential Operators I*, 2nd ed., Grundlehren **256**, Springer (1990); Thm. 7.7.1 (non-stationary phase).
- [9] R. Wong, *Asymptotic Approximations of Integrals*, Classics in Applied Mathematics **34**, SIAM (2001); Ch. 3 (Watson's lemma and the Mellin method).
- [10] J. Tate, *Fourier analysis in number fields and Hecke's zeta-functions*, thesis (1950), in J. W. S. Cassels and A. Fröhlich (eds.), *Algebraic Number Theory*, Academic Press (1967), 305–347.
- [11] R. Godement and H. Jacquet, *Zeta Functions of Simple Algebras*, Lecture Notes in Mathematics **260**, Springer (1972).
- [12] A. Weil, *Sur les courbes algébriques et les variétés qui s'en déduisent*, Actualités Sci. Ind. **1041**, Hermann, Paris (1948); the Riemann hypothesis for curves over a finite field — the Frobenius eigenvalues on the Jacobian have absolute value $q^{1/2}$, placing the zeros of the congruence zeta function on $\text{Re } s = \frac{1}{2}$.
- [13] P. Deligne, *Les constantes des équations fonctionnelles des fonctions L* , in *Modular Functions of One Variable II* (Antwerp), Lecture Notes in Math. **349**, Springer (1973), 501–597 (the local L - and ε -factors of a Weil–Deligne representation); *La conjecture de Weil. I*, Publ. Math. IHÉS **43** (1974), 273–307 (the Riemann hypothesis for smooth projective varieties over finite fields); and . . . *II*, Publ. Math. IHÉS **52** (1980), 137–252 (purity: the normalized Frobenius eigenvalues lie on the unit circle).
- [14] J. W. Cogdell and I. I. Piatetski-Shapiro, *Converse theorems for GL_n* , Publ. Math. IHÉS **79** (1994), 157–214; *II*, J. reine angew. Math. **507** (1999), 165–188.
- [15] F. Shahidi, *Eisenstein Series and Automorphic L -functions*, Amer. Math. Soc. Colloquium Publications **58**, AMS (2010); and *A proof of Langlands' conjecture on Plancherel measures; complementary series for p -adic groups*, Ann. of Math. **132** (1990), 273–330.
- [16] H. Kim, *Functoriality for the exterior square of GL_4 and the symmetric fourth of GL_2* , with Appendix 2 by H. Kim and P. Sarnak, J. Amer. Math. Soc. **16** (2003), 139–183 (the bound $|\alpha_p| \leq p^{7/64}$).
- [17] H. Kim and F. Shahidi, *Functorial products for $GL(2) \times GL(3)$ and the symmetric cube for $GL(2)$* , Ann. of Math. **155** (2002), 837–893 (for the symmetric fourth, see [16]).
- [18] J. Newton and J. A. Thorne, *Symmetric power functoriality for holomorphic modular forms, I & II*, Publ. Math. IHÉS **134** (2021), 1–116, 117–152.
- [19] J.-P. Serre, *Abelian ℓ -adic Representations and Elliptic Curves*, Benjamin (1968); Ch. I, the equidistribution criterion for the Sato–Tate law from the analyticity and non-vanishing of the symmetric-power L -functions.
- [20] H. Jacquet and J. A. Shalika, *A non-vanishing theorem for zeta functions of GL_n* , Invent. Math. **38** (1976), 1–16; non-vanishing on $\text{Re } s = 1$ from the positivity of the Rankin–Selberg convolution; and *On Euler products and the classification of automorphic representations*, Amer. J. Math. **103** (1981), the local bound $|\alpha| < p^{1/2}$.

- [21] W. Luo, Z. Rudnick, and P. Sarnak, *On the generalized Ramanujan conjecture for $GL(n)$* , in *Automorphic Forms, Automorphic Representations, and Arithmetic*, Proc. Sympos. Pure Math. **66**, Part 2, AMS (1999), 301–310; the bound $|\alpha| \leq p^{1/2-1/(n^2+1)}$ toward Ramanujan.
- [22] E. C. Titchmarsh, *The Theory of Functions*, 2nd ed., Oxford University Press (1939), §§5.6–5.65 (Phragmén–Lindelöf).
- [23] B. J. Birch and H. P. F. Swinnerton-Dyer, *Notes on elliptic curves. II*, J. reine angew. Math. **218** (1965), 79–108; the conjecture relating the analytic rank to the Mordell–Weil rank.
- [24] B. H. Gross and D. B. Zagier, *Heegner points and derivatives of L -series*, Invent. Math. **84** (1986), 225–320; the central derivative as the Néron–Tate height of a Heegner point (analytic rank one).
- [25] V. A. Kolyvagin, *Euler systems*, in *The Grothendieck Festschrift, Vol. II*, Progr. Math. **87**, Birkhäuser (1990), 435–483; finiteness of Sha and the rank bound completing Birch–Swinnerton-Dyer at analytic rank ≤ 1 .
- [26] S. Zhang, *Gross–Schoen cycles and dualizing sheaves*, Invent. Math. **179** (2010), 1–73; the Beilinson–Bloch height of the modified diagonal.
- [27] X. Yuan, S. Zhang, and W. Zhang, *Triple product L -series and Gross–Schoen cycles*, book manuscript; the central derivative of the triple-product L -function as a Gross–Schoen/Ceresa height.
- [28] G. Ceresa, *C is not algebraically equivalent to C^- in its Jacobian*, Ann. of Math. (2) **117** (1983), 285–291.
- [29] B. Harris, *Harmonic volumes*, Acta Math. **150** (1983), 91–123, and *Homological versus algebraic equivalence in a Jacobian*, Proc. Natl. Acad. Sci. USA **80** (1983), 1157–1158 — the Fermat-quartic Ceresa cycle detected by a nonzero harmonic volume.
- [30] S. Lavery, *The loss ledger and the cycle filtration: tower exhaustion on the helix carrier, and a program whose terminus is the Hodge conjecture*, companion preprint draft (2026); the depth dictionary, the harmonic Hodge frame, the grade-four program, and the Ceresa landing.
- [31] S. Lavery, *The Carrier Geometry of the Critical Line: A Worked Example with Two Conditional GRH Routes*, companion preprint draft (2026); the worked example at full resolution, the two compiler-audited conditional routes, and the operator reduction certified equivalent to Mathlib’s `RiemannHypothesis` from both sides.
- [32] J. Laga and A. Shnidman, *Ceresa cycles of bielliptic Picard curves*, arXiv:2312.12965 (2024); the Beilinson–Bloch height proportional to the Néron–Tate height of an explicit elliptic point.
- [33] B. Riemann, *Ueber die Anzahl der Primzahlen unter einer gegebenen Grösse*, Monatsberichte der Königlich Preussischen Akademie der Wissenschaften zu Berlin (November 1859), 671–680.
- [34] A. M. Odlyzko, *Correspondence about the origins of the Hilbert–Pólya conjecture* (letters exchanged with G. Pólya, 1981–1982), <http://www.dtc.umn.edu/~odlyzko/polya/> — the written record of the suggestion; no canonical formal statement exists.
- [35] H. L. Montgomery, *The pair correlation of zeros of the zeta function*, in: Analytic Number Theory (St. Louis, 1972), Proc. Sympos. Pure Math. **24**, Amer. Math. Soc. (1973), 181–193.

- 5005 [36] M. V. Berry and J. P. Keating, *The Riemann zeros and eigenvalue asymptotics*, SIAM Review **41**
5006 (1999), 236–266.
- 5007 [37] A. Connes, *Trace formula in noncommutative geometry and the zeros of the Riemann zeta function*,
5008 Selecta Math. (N.S.) **5** (1999), 29–106.