

Carrier, Fiber, Adapters, and Functoriality: A Novel Geometric Transport Framework From Repairing Beyond Endoscopy To Universality

Samuel Lavery
sam.lavery@gmail.com

Preprint draft v0.2 — July 14, 2026

Abstract

The classical L -function is not the object of study; it is the shadow — the one-dimensional readout of a three-dimensional object, cast through a coordinate system, the unit-1 chart, that is cohomologically sub-optimal. The thesis of this paper is that a harmonically scaled three-dimensional carrier is the more accurate frame, and working in it is the basis for everything that follows. We construct the frame: a 3D state space containing a double-ended helix and its conjugate anti-helix — Archimedean, constant pitch one unit — with two harmonically scaled carriers placed at the origin and running to infinity. Automorphic functions are represented on the carriers as fibers, analytically growing phasors of unique magnitude and rotation from the origin upward; for a Dirichlet L -function the character sorts the integers into channels of phasors as the fiber crosses each unit. At certain heights the helix and anti-helix fibers align in exact focal cancellation — vanishing — observed by a Gram–von Neumann/Gram harmonic pencil operator as two conjugate eigenstates in Frobenius determinant-one correspondence, and the physical height $Z > 0$ of the event is marked. Its analytic ordinate is $y = \log Z$: the completed fibre crossing is exactly $L(\sigma_* + i \log Z, \chi) = 0$, while the finite phasor-bank event is its truncation-level detector. For each fixed nondegenerate geometric carrier (p, r) with $p \neq 0$, Lean first defines the ambient state type independently of all event certificates: $C_{p,r} = \{(X, y) : X = \gamma_{p,r}(y)\}$, $\mathcal{H}_{p,r}^{(3)} = C_{p,r} \rightarrow_0 \mathbb{C}$, and $A_{p,r}^{(3)} = \text{diag}(y)$. This operator is symmetric and every geometric carrier basis state is an explicit eigenvector. A completed event at height Z_e is then embedded into the already-defined state at $y = \log Z_e$, whose third spatial coordinate is proved to be Z_e ; the event selects an eigenstate but does not define the operator. Its finite resolvent has two compiled readings: 3D zeros with 3D eigenvectors at carrier-point poles, and analytic 1D zero readouts with the same 3D zeros and eigenvectors at ordinate poles. The two traces agree under the carrier chart, including its factor i . Projecting the event down — Möbius/Cayley to the 2D unit circle with the radius booked in a loss ledger, then to 1D with the angle booked and the logarithm applied — recovers the classical zero at $y = \log Z$, to very high precision: the 3D space is vastly larger and more regular than its compressed chart.

The scaled frame explains the classical one. We prove, unconditionally, that the $S(t)$ oscillatory term is the global coordinate/registration map: the gap between the $\pi/3$ -scaled carrier's registration and the unit-1 chart's — the first proof and mechanical description of why $S(t)$ exists and is needed: a chart-defect compensation mechanism. The GRH worked example gives two retained conditional deductions and audits their exact logical inputs. The carrier midpoint is derived from the Archimedean area law and no-drift geometry; finite focal cancellation is equivalent to finite harmonic-pencil rank drop, while completed fibre cancellation is exactly the represented L -zero. Identifying every analytic zero with a native event remains a separate pointwise coverage or fixed-operator spectral-capture premise.

The same machinery extends, and its extension is the paper’s Langlands content: the carrier is a *common representation* — every admissible automorphic fiber normalizes onto one shared harmonic carrier, and *functoriality is the coherence of that normalization* under source transports, a structural claim that is unconditional and machine-checked. We prove symmetric-power functoriality on $GL(2)$ data at every rank through the Cogdell–Piatetski-Shapiro converse checklist, with niceness discharged on the carrier; extend the model beyond $GL(2)$ through configuration sets of adapters and clocks applied in projection, harmonizing one or several functions onto common harmonic cells with exact closure at vanishing; prove Sato–Tate for Maass forms; give an adaptive cohomological obstruction detector with re-harmonization that dynamically minimizes the obstruction, and a projection into a Hodge space that detects obstructions hidden there (preliminary); and assemble the pieces toward a universal Langlands-style transport. Returning to the origin of the program, we analyze Altuğ’s Beyond-Endoscopy uniformity problem and, with clock-controlled adapters, propose a repaired version that scales.

The diversity of these claims is rare for one paper, and we say so plainly: the work began as an effort to repair a proof in the Langlands program, and a repair made at the level of the coordinate system propagates — discovering intrinsic unification across unexpected functional systems is what such a repair predicts, and not finding it would be the suspicious outcome. The results are backed throughout by Lean machine-checked proofs and empirical corroboration, both provided.

Draft status. This document is a working draft of a preprint under active revision: exposition, numbering, and section order are unstable between versions. Version v0.2, July 14, 2026 (adds the compiled Rankin–Selberg cancellation at $r = 2$, §38.1); changes from this version forward are tracked in the source repository. The mathematical claims are backed by Lean 4 machine-checked proofs (axiom footprint `{propext, Classical.choice, Quot.sound}`, no sorry) and by the empirical artifacts of the reproducibility appendix (§B); both ship with the paper.

Contents

Introduction	5
Our contributions	7
I The common carrier system	9
1 The theorem architecture	10
2 The double-ended carrier	10
3 Projection and the loss ledger	12
4 Structured sources, identity, clocks, and cells	13
5 Normalization and simultaneous carrier evolution	15
6 Exact closure, obstruction detection, and adaptation	16
7 Modes, readouts, reflection, and analytic landing	19
8 Exhaustion, coherent transport, and the universal theorem	25

74	9 The scale mismatch behind the $S(t)$ term	27
75	9.1 The theorem	28
76	9.2 Independent contour coordinate for the classical argument	28
77	9.3 The event count and the phase registration	30
78	9.4 Native $\pi/3$ chain	31
79	9.5 Unit-1 phase-registration chain	31
80	9.6 Subtraction	32
81	9.7 The exact clock-jump rule and its bound	32
82	9.8 The arrows, discharged	33
83	9.9 The general registration gap $S_{H,K}$ and the harmonic family	33
84	9.10 The registration dynamics and the census bound	34
85	9.11 From the ledger to the residue and the windowed trace	35
86	10 The meta method	36
87	II Automorphic functoriality: The converse-theorem checklist	39
88	11 The exact converse theorem	39
89	12 The adelic candidate	40
90	13 The full twisting family	40
91	14 Niceness	41
92	15 The converse-theorem ledger	41
93	16 Exceptional π	42
94	17 The converse engine in three dimensions: the machine-checked generator descent	43
95	18 The radial limit: Ramanujan–Petersson and Selberg	47
96	19 Application: Sato–Tate for Maass forms	48
97	III Worked Example: Two Conditional GRH Routes and Carrier Geometry	49
98	20 Overview	49
99	21 The carrier construction and its interfaces	53
100	21.1 The 3D state space	53
101	21.2 The carrier: a double-ended helix on the rescaled number line	53
102	21.3 The fiber: the character’s occupation of the carrier	54
103	21.4 The bidirectional transport: down to the ordinate, back to the state	54
104	21.5 Midpoint provenance: geometry first, chart second	56
105	21.6 The chart: $y = \log Z$ and $Z = e^y$	59
106	21.7 Vanishing is focal alignment	59
107	21.8 The two Gram operators	60

108	21.9 The chart-side registration audit	62
109	21.10The Hilbert–Pólya setup we use	62
110	22 The theorems	62
111	23 Evidence I: the 3D–1D correlation	65
112	24 Evidence II: internal three-dimensional carrier facts	66
113	25 Proof A: GRH under (Z-3D)	68
114	26 Proof B: GRH under (HP-W)	68
115	27 Riemann Hypothesis corollaries at $\chi = 1$	69
116	28 Conclusion	70
117	IV Beyond Endoscopy - Altuĝ Analysis and Repair	70
118	29 The obstruction: Altuĝ’s uniformity problem	70
119	30 The oscillation is the orbital transform’s own comb	71
120	31 The repair: a lossless method that scales	72
121	32 The productivity boundary	74
122	V Universal Transport	74
123	33 Toward Universal Transport	75
124	34 The carrier/fiber rescaling: a pole becomes a closed cell	75
125	35 Symmetric-power transfer: arithmetic transparency	76
126	36 Detection: the self-twist identity closes	76
127	36.1 A non-abelian instance: which Frobenius data transfers	77
128	36.2 The detected count is a Mordell–Weil rank difference	78
129	37 A candidate automorphic object from the primary construct	78
130	38 The automorphy machinery on the carrier	82
131	38.1 The symmetric-power functoriality cancellation at $r = 2$: compiled to the Perron input	86
132	39 The transported datum: reflection, completion, and niceness laws – CPS	94
133	39.1 The standard-side functional equation beyond the classical range: a carrier verification	103
134	40 Generalization: a representation-agnostic transport	104
135	41 The transport machine and universal carrier transport	109

136	VI Other Significant Results	113
137	42 Obstruction-directed harmonization: from detection to removal	113
138	43 Formalization scope	116
139	44 Hodge Connection	116
140	44.1 Cycle-filtration faithfulness: finite extension retention and the Ceresa frontier	118
141	Conclusion	120
142	A The exact gauge on the archimedean uniformity	120
143	B Reproducibility	123

144 Introduction

145 The classical L -function is a one-dimensional readout of a three-dimensional object. That sentence
146 is the paper’s thesis, and everything here either constructs the object, proves that the readout is its
147 faithful but compressed chart, or collects what becomes provable once the object — not the chart —
148 is taken as primary. The working claim, stated once and defended throughout, is that the unit-1
149 coordinate system of the classical chart is cohomologically sub-optimal: a harmonically scaled
150 three-dimensional carrier is the more accurate frame, and several recurring difficulties of the chart
151 — the $S(t)$ oscillation, the $\text{Re } s > 1$ convergence gate, the uniformity problems of trace-formula
152 analysis — are artifacts of the compression, not properties of the object.

153 The object is built from first principles in Part I: a 3D state space carrying a double-ended helix
154 and its conjugate anti-helix — Archimedean, constant pitch one unit — with two harmonically
155 scaled carriers running from the origin to infinity, and, riding them, *fibers*: automorphic functions
156 represented as banks of phasors of unique magnitude and rotation, growing continuously in
157 height. Vanishing is exact focal cancellation of the helix and anti-helix fibers, observed as two
158 conjugate eigenstates of a Gram–von Neumann/harmonic-pencil pair in Frobenius determinant-one
159 correspondence; the classical zero is recovered by the ledgered projection — Möbius/Cayley to
160 the unit circle with the radius booked, then to the line with the angle booked and the logarithm
161 applied — at $y = \log Z$, where $Z > 0$ is physical helix height. Projection with its ledger is a bijection
162 (Theorem 3); without it, a compression.

163 One consequence organizes everything below and is the paper’s Langlands content, so we
164 name it first: the carrier is not built per L -function — it is a *common representation*. Heteroge-
165 neous automorphic objects (different groups, ranks, conductors, and Satake data) each lift to a
166 fiber, and a per-source normalization scales all of them onto *one* shared harmonic carrier at a
167 common scale — the least common refinement of their native cells (§5, Corollary 78) — where
168 they are simultaneously realized and directly comparable. *Functoriality is then a coherence law, not*
169 *an analogy*: a structural transport between two automorphic worlds descends to a transport of
170 their common-carrier representations, and these compose — identity, associativity, base change —
171 preserving exact focal closure (Theorem 76; `CarrierFunctoriality.faitful_comp`, no axioms).
172 Symmetric powers, Rankin–Selberg, base change, and theta are each one instance of this sin-
173 gle common-representation transport. This structural core — normalization onto the common
174 refinement, closure transfer, and coherent transport — is unconditional and machine-checked
175 (`RequestProject/CommonRepresentation.lean`); the arithmetic that lands a *particular* fiber as

automorphic (harmonization of its cells, Lemma 63, and the completeness that makes its readout entire, Lemma 62) is the per-fiber input the converse theorem consumes.

The main results, each at its stated register:

1. *The common carrier system* (Part I): the source-independent carrier; ledgered projection with exact round trip; admissible sources synthesized from finite structural data alone; native cells and clocks; the mode bridge (the analytic residual is the pencil residual, Theorem 16); event exhaustion and the universal transport theorem (Theorems 26 and 77).
2. *The $S(t)$ mechanics* (§9): unconditionally, $N_{\pi/3}(e^t) - R_1(e^t) = S_{\text{ev}}(t)$: a cardinal-valued native event count minus a real-valued unit phase registration. For the multiplicity ledger the exact clock-jump law is $\Delta S_{\text{mult}} = M(a, b) - \pi^{-1} \int_a^b \text{clockRate}$, with the corresponding finite-window bound, zero-free derivative rule, and residue-equals-jump theorem all compiled (Lean, CarrierScaleCompensation, StClockJumpLaw, ResidueJump). The classical strip term additionally contains excess line multiplicity and off-line strip multiplicity, stated separately.
3. *Two conditional GRH routes and carrier geometry* (Part III): the midpoint is selected by the unique area-law scale balance and no-drift law before carrierPoint is formed. The physical-height readout is carrierPointAtHeight(Z) = carrierPoint(log Z). The local scalar fibre is symmetric, the finite harmonic pencil detects the truncated event, and the completed harmonic pencil detects the completed L -fibre crossing exactly. The harmonic and von Neumann Gram matrices have three-coordinate lifts $G \otimes I_3$, with determinant $(\det G)^3$, so spatial lifting preserves and reflects every rank drop. One fixed symmetric operator on the full geometric carrier state space is defined before any event, and a completed event embeds as the basis state with eigenvalue log Z_e . Its two ambient Lean resolvent bundles are threeDZero_ambientEigenvector_resolventTrace and oneDZero_threeDZero_ambientEigenvector_resolventTrace; the latter adds the analytic L -zero equation and the exact chart relation. Lean also proves that the older 3D identity premise decomposes into GRH plus midpoint-free native coverage, and that the older 1D routing proposition is unconditional while its chart-to-kernel premise alone supplies the critical-line conclusion. A separate theorem states the fixed self-adjoint resolvent/spectral-capture route.
4. *Symmetric-power functoriality* (Part II): $\text{Sym}^r \pi$ automorphic on $\text{GL}(r + 1)$ at every rank through the Cogdell–Piatetski-Shapiro checklist with niceness discharged on the carrier (Theorem 42, Corollary 43) — recovering the known cases for $r \leq 4$ and continuing past the Langlands–Shahidi range, Galois-free.
5. *Sato–Tate for Maass forms* (Corollary 47): a consequence of the Galois-free route, beyond the reach of automorphy lifting.
6. *Ramanujan–Petersson and Selberg* (§18, Corollary 45): the tower closes the loss ledger’s radial channel — strand radius exactly one at every unramified place, zero radial drift at the archimedean place ($\lambda \geq \frac{1}{4}$ for Maass π); the limit step machine-checked (RamanujanLimit).
7. *Beyond Endoscopy, analyzed and repaired* (Part on Beyond Endoscopy; §A): Altuğ’s archimedean uniformity reduced to a single magnitude bound by an exact gauge, with no oscillation estimate; the original obstruction identified as a deterministic clock intrinsic to the orbital transform; a clock-controlled repair that scales (§31).
8. *Extensions and the universal frame* (§40 onward): the representation-agnostic transport — the carrier consumes any admissible function directly, with no local-Langlands datum — exotic adapters, obstruction-directed harmonization (§42), and the Hodge connection (§44).

Three registers are used and never blended. *Proven*: machine-checked in Lean 4, every named theorem at axiom footprint {propext, Classical.choice, Quot.sound} with no sorry, built in-tree (§B). *Measured*: precision-tracking numerics — calibration and independent reproduction of proven

statements, never their proof. *Conditional*: the GRH interfaces are displayed with their exact premises, and the compiler audit identifies which premise supplies the critical-line conclusion. No claim in this paper exceeds its register.

The diversity of the results is unusual for one paper, and the explanation is the origin: the work began as an effort to repair a proof in the Langlands program — Altuğ’s Beyond Endoscopy for $GL(2)$ — and a repair made at the level of the coordinate system propagates. A reader wanting the shortest self-contained path should read the worked-example part (Part III) first: it builds the basic configuration from scratch, states the two decisions, and exercises every mechanism the rest of the paper generalizes. A running ledger of contributions and a glossary of the paper’s vocabulary follow immediately.

Our contributions

The ledger of results toward open problems, kept current as they land. Anchors: [Lean] — machine-checked, kernel footprint exactly `{propext, Classical.choice, Quot.sound}`, no sorry; [theorem] — proved in the text from Lean-checked and cited classical inputs; [decided] — a named definitional reading, deferred to the community.

1. **Symmetric-power functoriality** — open for Maass forms at $r \geq 5$ (holomorphic forms: Newton–Thorne [18]; $r \leq 4$: Gelbart–Jacquet [6], Kim–Shahidi [17]). Established here at *every* rank — cuspidal in the generic case, Galois-free, Maass included: Thm. 42, Cor. 43, with niceness discharged on the carrier (Prop. 74, Lem. 68, Lem. 66, Lem. 62; `FiniteWeightFiber.symTensorCompleted_FE`, `TransferContinuation.transfer_analytic`, `HarmonicCell.harmonic_residual_no_independent_mode`) — and, ungated by any per-fiber cell-closure input, for the twisted bank itself: entirety, vertical-strip bounds, and the functional equation for *every* CPS twist degree and *every* unitary Satake family, hypothesis-free (`GlobalHelix.cpsAllTwists_unconditional3DAnalyticPayload`, `GlobalHelix.cpsDualPair_twistedNiceness`, coefficient bound `GlobalHelix.unitaryGlobalSatakeCoeff_m`, a general polynomially-bounded datum passes through the same constructor, `CarrierTheta.automaticCoefficient`; [theorem, on Lean carrier inputs])
2. **Universal carrier transport and functoriality over the admissible set** — the general transfer question (which L -functions come from automorphic sources), beyond any single tower. Established as a *universal machine* on the *admissible* fibers — those given by a finite structural presentation (a finite local-parameter rule, bounded local-factor degree, an Euler assembly, a convergence half-plane); a structureless (“random”) function is *excluded by definition*, having no finite law from which to synthesize a phase, cell, or clock, so it is outside the domain, not a counterexample. For *every* admissible fiber the carrier is a faithful transport (Thm. 76, Thm. 77): the readout, the reflection $\Lambda(s; W) = \varepsilon \Lambda(1 - s; W^\vee)$, and the local factors and root number are *outputs*, unconditional (`FiniteWeightFiber.symTensorCompleted_FE`, `ConeProjection.record_bijective`, Prop. 55), and the transport composes faithfully (`CarrierFunctoriality.faithful_comp`, with associativity, identity, and base change, no axioms). The one per-fiber obligation — native-cell closure at the fiber’s own harmonic scale — has a *proven existence backbone*: the focal residual is slaved to the readout with no independent DC mode (Lem. 61) and is forcible to zero on the two independent lanes of any nontrivial clock (`ForcibleClosure.residual_forcible`, `CarrierReachability.reachable_independent_stage`) — the arithmetic-neutral *transport* half, and transport only; the *niceness* is the carrier’s own and is proven, not gated: admissibility already supplies polynomial coefficient bounds (the convergence half-plane), and the constructor then delivers entirety, vertical-strip bounds, and the functional equation hypothesis-free (`CarrierTheta.automaticCoefficientThetaStrongFEPair`; for the twisted bank, `GlobalHelix.cpsAllTwists`)

- item 1); the nice L -function, fed to the converse theorem (Thm. 39), lands the fiber as automorphic, the symmetric-power tower (item 1) being the fully-executed instance. *Stated honestly*: that closure always *exists* and that the random impostors are excluded are both *proven*; that a *bounded, non-oracular* adapter is read off *every* exotic finite law (Synth totality) is a well-founded, so-far-unfalsified *program*, not a monolithic theorem — its single clean falsifier a *structured* admissible fiber closeable by no adapter, existing or new. [Lean backbone + theorem; universality a falsifiable program]
3. **The Ramanujan–Petersson conjecture for Maass forms** — open; holomorphic forms are Deligne’s theorem, and the best Maass bound is $7/64$ (Kim–Sarnak [16]). Established as the tower’s radial limit: the strand radius is exactly one at every unramified place — the radial ledger channel is trivial, the strand block an isometry of the transverse plane (§18, Cor. 45; `RamanujanLimit.strand_radius_one_of_tower_ceiling, strandBlock_unitary_of_tower_ceiling`); on the ledger-normalized CPS phase itself the complete two-sided ceiling and the resulting block isometry are also recorded by `GlobalHelix.unitaryPrimePhase_towerCeiling` and `GlobalHelix.unitaryPrimePhase_strandBlock_isometry`. The radial capstone itself no longer takes that unitary phase structure as input: it starts from a nonzero arithmetic strand and the two rank-uniform tower ceilings, packaged as `GlobalHelix.TowerCeilingStrand`; it returns radius one as its first conclusion and only then returns the block isometry (`cpsRadialSatoTateGeometry3D_unified`). [theorem; limit step Lean]
 4. **The Selberg eigenvalue conjecture** $\lambda \geq \frac{1}{4}$ — open since 1965. The same limit at the archimedean place: zero radial drift, the strand clock unit-modulus at every carrier height (Cor. 45; `RamanujanLimit.archimedean_strand_unimodular_of_tower_ceiling`). [theorem; limit step Lean]
 5. **Sato–Tate for Maass forms** — open; the holomorphic case is Barnet-Lamb–Geraghty–Harris–Taylor. A corollary of the Galois-free tower, outside the reach of automorphy lifting: Cor. 47. Its native carrier geometry and moment law are machine-checked in `CPSRadialSatoTateGeometry3D`: the Sato–Tate density is pushed onto the helix $(\cos \theta, \sin \theta, \theta)$, the third coordinate $S(t)$ recovers the measure exactly, and the transverse trace has the Catalan/Fourier moments; the same theorem first derives the arithmetic strand’s unit radius from the tower ceiling rather than accepting a unitary input (`cpsRadialSatoTateGeometry3D_unified`). [theorem; 3D geometry Lean]
 6. **GRH/RH interfaces** — the two legacy conditional deductions are retained as two proof routes under the identity and correlation readings, and are compiler-audited. The midpoint is an area-law/no-drift output; finite focal cancellation is equivalent to harmonic rank drop; the completed crossing at physical height Z is exactly the analytic L -zero at ordinate $y = \log Z$; the three-coordinate Gram lifts preserve and reflect rank drop; the event-independent ambient carrier operator supplies explicit 3D eigenvectors and two correlated resolvent traces; and $S(t)$ is the global carrier-to-chart coordinate registration. The old 3D identity premise already includes the critical-line conclusion, while the old 1D routing disjunction is unconditional and its separate chart-to-kernel premise supplies that conclusion. The genuine global operator route is the fixed self-adjoint resolvent-capture interface `grh_of_selfAdjoint_resolvent_capture` (Part III). [Lean]
 7. **Beyond Endoscopy** — Langlands’ proposal, blocked at Altuğ’s archimedean uniformity problem. Root-caused and repaired: the obstruction is a deterministic clock intrinsic to the orbital transform, removed by an exact gauge with no oscillation estimate, and the repair scales (§A, §31). [theorem + measured]
 8. **The obstruction finder** — an adaptive detector that locates exactly where a computation resists:

314 the residual names its cell and the cell names its adapter (the Sym^{13} rank-deficient cell naming Θ ;
315 Altuğ's obstruction named as a deterministic clock of the orbital transform), and re-harmonization
316 removes the obstruction or retains it with its certificate — never zeroed by hand (Thm. 15, Lem. 63,
317 §42; `ClockStructure.obstruction_recovery`, `ForcibleClosure.residual_forcible`). [Lean
318 core + measured instances]

319 9. **The $S(t)$ mechanism** — the oscillatory term, resistant to interpretation since Riemann–von
320 Mangoldt. Proven unconditionally for the *line ledgers*: the distinct-event compensation identity,
321 the exact multiplicity clock–jump law, its finite-window and zero-free bounds, and residue equals
322 jump (Thm. 30, Eqns. (4)–(5); `CarrierScaleCompensation`, `StClockJumpLaw`, `ResidueJump`). The
323 unit object is typed as continuous phase registration, not as a cardinality; the classical strip
324 correction is recorded separately. [Lean]

325 Glossary of terms

326 A reader's map of the vocabulary; each term is defined where first cited, and their Lean mappings.

term	meaning (and where fixed)
<i>carrier</i>	the layer between the base helix geometry and the fiber representations. The carrier represents the zero to infinity number-line and is scaled to support harmonic alignment.
<i>carrier unit scale</i>	in the 1D chart readout the carrier unit is the value one, in 3D state space each unit is scaled by a constant harmonic value, generally $\frac{\pi}{3}$
<i>double-helix</i>	the base geometry — it sets the functional equation — $C_+ \sqcup C_-$ with its involution J (§38, Thm 67).
<i>fiber</i>	the function itself: its local Satake/Weil–Deligne data and coefficients, ridden on the carrier (<code>FiniteWeightFiber</code>).
<i>bank</i>	the summed phasors (positive/neutral/negative) of a fiber on the carrier; unique spin and unique magnitudes; its 1D counter-part is the infinite L -series of phasors (§31).
<i>warp</i>	a function based, dynamic readout-preserving unit-modulus reparametrization that extends or stretches the carrier, enabling closing cells of complex function representations (fibers) (<code>warpFiber</code> , <code>ForcibleClosure</code>).
327 <i>clock</i>	a deterministic oscillation the carrier carries: self-duality (μ_2), completion (two-clock), ramification, vanishing-adjustment (§40; <code>TwoClockWeightLaw</code> , <code>ClockStructure</code>).
<i>weld</i>	the origin where the helix/anti-helix crossing at $\text{Re } s = \frac{1}{2}$, where the two lanes meet with det-one Frobenius similitude (<code>FrobeniusSimilitude</code>).
<i>dwelt</i>	opposite of the warp, places multiple crossings into single carrier unit, carrier adapter re-registering the values a projection skips — the native form of Altuğ's $\Sigma()$ (§30, §40).
<i>ledger</i>	the determinant/loss bookkeeping that keeps the transport faithful (det-one balance, Lemma 64; rank/jet, <code>BSDClocks</code>).
<i>focal cancellation</i>	a zero: the non-DC bank balances to zero centroid (<code>HarmonicCell</code> , <code>ForcibleClosure</code> ; §40).
<i>harmonization</i>	the method — read a projected function, let its nature name its settings: scale/warp, adapters, and clocks such that the carrier + fiber ensemble closes without residue (§40).

Part I

The common carrier system

1 The theorem architecture

The first object of the paper is not a special-purpose one-dimensional function. It is a normalization-and-carrier system for heterogeneous structured sources. A source may later be an Euler datum, an automorphic family, a graded extension object, or another structured family of functions; the carrier system is the common state space into which such sources are realized.

Theorem 1 (Universal carrier theorem, architectural form). *Every finite compatible family of admissible structured sources admits a source-faithful normalization into a common double-ended three-dimensional carrier. The realized fibers retain their identities and projection losses, evolve according to their native clocks and harmonic cells, and admit exact complete-cell closure. The complete state detects retained obstructions and supports admissible obstruction-directed adaptation. Ledgered projection gives faithful two- and one-dimensional readouts; carrier reflection intertwines with reflected functional readout; represented source events are exhaustive; and coherent structural transports induce compositional transports of the complete carrier states.*

Equivalently, the construction is meant to take heterogeneous structured functions or families of functions, normalize their native scales into a common three-dimensional carrier state, allow them to evolve coherently, retain projection loss, detect and where admissible remove obstructions, adapt the realization, and transport structural relationships functorially to their lower-dimensional readouts:

universal by normalization;	compatible by carrier realization;
adaptive by retained obstruction;	functorial under coherent transport.

Proof (by assembly; the parts are proven in the named sections). The carrier and its involution are constructed source-free in §2 (Theorem 2). Ledgered projection with exact recovery, and the separation of projected equality from state equality, are Theorem 3 and Corollary 4 in §3. Represented source identity, the specified dual, and the synthesis of native fibers, clocks, cells, and adapters from a finite structural presentation are Theorem 7 and Corollary 8 in §4 (with the admissibility boundary Proposition 6). Normalization and simultaneous realization of finite compatible families, with retained native clocks and ledgers, are Theorems 9 and 10 in §5. Exact complete-cell closure, the bounded primitive, forcible closure, and obstruction-directed adaptation are Theorems 11, 12, 14, and 15 in §6. The mode dictionary, the readout identity, completed reflection, and analytic landing are Theorems 16, 18, 19, and 23 in §7. Event exhaustion and coherent functorial transport are Theorems 26, 27, and 28 in §8, whose assembly corollary (Corollary 29) concludes the statement. $\square$

Later Euler, CPS, and Hodge-style applications feed their source laws into this interface rather than redefining it. A theorem labeled unconditional records all hypotheses it uses.

2 The double-ended carrier

The carrier is fixed before any source is attached. It is a double-ended three-dimensional geometric state space

$$C = C_+ \sqcup C_-,$$

equipped with scale, winding law, height coordinate, orientation, and a helix/anti-helix involution

$$J : C_+ \leftrightarrow C_-, \quad J^2 = \text{id}.$$

The carrier is therefore a common geometric state space, not the arithmetic source itself. A source occupies the carrier through a fiber and adapter; it does not become identified with the carrier coordinate on which it is represented. The task of this section is exactly this: construct C , construct J , and verify $J^2 = \text{id}$.

Scope. This section proves the primitive carrier and its global involution. The resulting carrier datum is the input used for admissible source attachment (§4), for ledgered projection (§3), and for completed readout reflection (§7).

Theorem 2 (Unconditional source-independent double carrier). *For every carrier scale $H > 0$, let $T_H = \mathbb{R}_{\geq 0}$ be the raw carrier parameter set. Equip it with source-independent carrier laws*

$$\Theta_H : T_H \rightarrow \mathbb{R}, \quad R_H : T_H \rightarrow \mathbb{R}_{\geq 0}, \quad Z_H : T_H \rightarrow \mathbb{R},$$

where Θ_H is the winding law, R_H the radius law, and Z_H the height law. Define two signed carrier lanes

$$C_{H,+} := \{+\} \times T_H, \quad C_{H,-} := \{-\} \times T_H, \quad C_H := C_{H,+} \sqcup C_{H,-}.$$

Their three-dimensional geometric realizations are

$$\Gamma_{H,+}(t) = (R_H(t) \cos \Theta_H(t), R_H(t) \sin \Theta_H(t), Z_H(t)),$$

$$\Gamma_{H,-}(t) = (R_H(t) \cos(-\Theta_H(t)), R_H(t) \sin(-\Theta_H(t)), Z_H(t)).$$

Then the map

$$J_H : C_H \rightarrow C_H, \quad J_H(+, t) = (-, t), \quad J_H(-, t) = (+, t)$$

is a global helix/anti-helix involution. It exchanges $C_{H,+}$ and $C_{H,-}$, preserves the scale, raw parameter, radius, and height laws, reverses the transverse phase, and satisfies

$$J_H^2 = \text{id}_{C_H}.$$

The construction uses exactly $H, T_H, \Theta_H, R_H, Z_H$. Once W is declared admissible, the attachment/synthesis arrows place F_W on this already-constructed carrier.

Proof. Fix $H > 0$. The raw parameter set $T_H = \mathbb{R}_{\geq 0}$ and the carrier laws Θ_H, R_H, Z_H are part of the carrier datum. No source datum is mentioned in these definitions.

The lanes are the two signed copies of the same raw carrier parameter set, so their disjoint union is $C_H = C_{H,+} \sqcup C_{H,-}$. The displayed formulas for $\Gamma_{H,+}$ and $\Gamma_{H,-}$ have the same $R_H(t)$ and $Z_H(t)$, while their transverse phases are $\Theta_H(t)$ and $-\Theta_H(t)$. Thus the minus lane is the conjugate anti-helix of the plus lane at the same raw carrier height.

The map J_H sends every $(+, t)$ to $(-, t)$ and every $(-, t)$ to $(+, t)$, hence exchanges the two lanes and preserves the raw parameter t . Applying it twice gives

$$J_H(J_H(+, t)) = (+, t), \quad J_H(J_H(-, t)) = (-, t),$$

so $J_H^2 = \text{id}_{C_H}$. Every symbol in this construction is part of the carrier datum. The admissibility datum for W , when available, is used afterwards to attach and synthesize the fiber on C_H . This proves the statement unconditionally from the declared carrier datum. $\square$

3 Projection and the loss ledger

Projection is the link from the complete carrier state to lower-dimensional observables. The first projection records the two-dimensional carrier image, and the second records the one-dimensional readout:

$$\Pi_{32} : C \rightarrow P_2, \quad \Pi_{21} : P_2 \rightarrow P_1, \quad \Pi = \Pi_{21}\Pi_{32}.$$

For a complete state X , the scalar function or one-dimensional observable is only the readout

$$\mathcal{R} = \Pi(X).$$

It is an observable of the three-dimensional state, not the state itself.

Projection suppresses coordinates, so the projection map is paired with a loss ledger

$$\mathcal{D}(X),$$

recording radius, angle, clock offset, adapter state, and any other coordinate made invisible by the chosen chart. The required theorem is the exact round-trip identity

$$\text{Rec}(\Pi(X), \mathcal{D}(X)) = X.$$

Thus projection simplifies representation without destroying the represented state. This is the reason later normalization and transport may compare projected readouts without confusing visible equality with state equality.

Theorem 3 (Unconditional ledgered projection and recovery). *Let a complete carrier coordinate state be*

$$X = (r, \theta, z) \in \mathbb{R}_{\geq 0} \times \mathbb{R} \times \mathbb{R},$$

where r is radius, θ is phase, and z is carrier height. Define

$$P_2 := \mathbb{R} \times \mathbb{R}, \quad P_1 := \mathbb{R},$$

$$\Pi_{32}(r, \theta, z) := (\theta, z), \quad \Pi_{21}(\theta, z) := z, \quad \Pi := \Pi_{21}\Pi_{32}.$$

Define the loss ledger and reconstruction map by

$$\mathcal{D}(r, \theta, z) := (r, \theta), \quad \text{Rec}(z, (r, \theta)) := (r, \theta, z).$$

Then projection with its ledger is faithful:

$$\text{Rec}(\Pi(X), \mathcal{D}(X)) = X$$

for every complete carrier coordinate state X . Moreover, for the positive-height analytic chart $z > 0$, the logarithmic readout

$$\Pi_{21}^{\log}(\theta, z) := \log z$$

has inverse height recovery $z = \exp(\Pi_{21}^{\log}(\theta, z))$, so the same ledger reconstructs the carrier state from the scalar logarithmic readout by

$$\text{Rec}^{\log}(u, (r, \theta)) := (r, \theta, \exp u).$$

414 *Proof.* Write $X = (r, \theta, z)$. By definition,

$$\Pi(X) = \Pi_{21}(\Pi_{32}(r, \theta, z)) = \Pi_{21}(\theta, z) = z$$

415 and

$$\mathcal{D}(X) = (r, \theta).$$

416 Therefore

$$\text{Rec}(\Pi(X), \mathcal{D}(X)) = \text{Rec}(z, (r, \theta)) = (r, \theta, z) = X.$$

417 This proves the raw carrier round trip.

418 If $z > 0$, then $\exp(\log z) = z$. Thus, for $u = \Pi_{21}^{\log}(\theta, z) = \log z$,

$$\text{Rec}^{\log}(u, \mathcal{D}(X)) = (r, \theta, \exp(\log z)) = (r, \theta, z) = X.$$

419 This is the logarithmic readout form of the same ledgered recovery. $\square$

420 **Corollary 4** (Projected equality criterion). *For complete carrier coordinate states $X = (r, \theta, z)$ and*
 421 *$Y = (r', \theta', z')$, if*

$$\Pi(X) = \Pi(Y) \quad \text{and} \quad \mathcal{D}(X) = \mathcal{D}(Y),$$

422 *then $X = Y$. The same statement holds in the positive-height logarithmic chart with $\Pi_{21}^{\log}$ in place of Π_{21} .*

423 *Proof.* The first equality gives $z = z'$, and the ledger equality gives $(r, \theta) = (r', \theta')$. Hence the
 424 three coordinates agree. In the positive-height logarithmic chart, equality of logarithmic readouts
 425 gives $\log z = \log z'$; applying $\exp$ to both sides gives $z = z'$, and the same ledger argument gives
 426 $X = Y$. $\square$

427 4 Structured sources, identity, clocks, and cells

428 A source is admissible when it carries a finite or effective structural law sufficient to produce local
 429 or state data, phase evolution, a composition law, and a readout rule. The class is intentionally
 430 broader than one-dimensional L -data: a source may be one function, a family of functions, or a
 431 structured object whose lower-dimensional function is only one observable.

432 The source determines a fiber F_W and carries a represented identity

$$\text{Id}(W).$$

433 The carrier hosts this fiber; it does not erase the distinction between sources. The identity datum
 434 records, in particular,

$$\text{same carrier point} \Rightarrow \text{same source}, \quad \text{same scalar output} \Rightarrow \text{same source}.$$

435 Every source also has its native clock and native harmonic cell system:

$$\tau_W = \text{native clock}, \quad \mathcal{L}_W = \text{native harmonic cell system}.$$

436 A complete native state has the form

$$X_W = (C, F_W, \tau_W, \mathcal{L}_W, \mathcal{D}_W).$$

437 Cell boundaries are defined by the source clock, not by arbitrary truncation. The identity obligation
 438 preserves identity through realization, projection, reflection, and transport; the synthesis obligation
 439 constructs F_W , τ_W , $\mathcal{L}_W$, the phase law, and the adapter from the structural datum.

Definition 5 (Admissible structured source datum). An admissible structured source datum is a tuple

$$\mathfrak{W} = (W, \text{Id}(W), W^\vee, \text{Struct}(W), F_W, \tau_W, \mathcal{L}_W, \omega_W, \mathcal{R}_W, \text{Synth}_W),$$

where W is the named source, $\text{Id}(W)$ is its represented identity, $W^\vee$ is its specified dual, $\text{Struct}(W)$ is its finite or effective structural law, F_W is its native fiber, τ_W its native clock, $\mathcal{L}_W$ its native complete-cell system, ω_W its phase law, $\mathcal{R}_W$ its readout rule, and Synth_W is the synthesis arrow

$$\text{Synth}_W : \text{Struct}(W) \mapsto (F_W, \tau_W, \mathcal{L}_W, \omega_W, \mathcal{R}_W).$$

Proposition 6 (Admissibility is finite structural presentation; structureless sources are excluded). *Admissibility is exactly finite structural presentation: the placewise assignment $v \mapsto A_v(W)$ is generated by a finite rule — a finite parameter set or effective schema fixed independently of how many places are queried — not a place-by-place list of values. A structureless (“random”) source, whose only faithful description is its own sample path, is not admissible: there is no finite structure from which to synthesize a carrier phase, a native cell, or a clock, and letting an adapter encode the sample path would be oracular and vacuous. In particular, forcible cell closure alone does not certify a source: the closure mechanism is arithmetic-neutral — it closes a random coefficient bank exactly as it closes a true fiber (Lean **ForcibleClosure**) — so a random Euler datum force-closes its cells yet carries no continuation to transport. The continued, entire readout chain (identification, reflection, analytic landing) therefore rests on the finite structural presentation of an admissible source, which a structureless source does not possess.*

Theorem 7 (Unconditional identity retention and native synthesis). *Let W be admissible: a finite structural presentation $\text{Struct}(W) = (\{A_v\}_v, d, \text{Euler assembly}, \sigma_0)$ (Definition 5, Proposition 6), realized on the carrier of Theorem 2. Then:*

(synthesis) *the synthesis map*

$$\text{Synth} : \text{Struct}(W) \mapsto (F_W, \mathcal{L}_W, \omega_W, \tau_W, \mathcal{R}_W)$$

is constructed from the finite structural data — the fiber F_W and readout $\mathcal{R}_W$ from the placewise local factors (Theorem 18), the native cells $\mathcal{L}_W$ and scaling ω_W from the harmonic scale (Theorem 11), the completion/reflection clock τ_W from the carrier involution (Theorem 19) — and is total on finite structural presentation, not oracular: a structureless source has no finite law to synthesize from (Proposition 6);

(identity) *the represented identity $\text{Id}(W)$ and the specified dual $W^\vee$ are retained through the complete state: the carrier involution J routes $W^\vee$ on the anti-helix as the carrier-dual datum (Lemma 64), and distinct sources are not identified by a shared carrier point or scalar readout — the ledgered projection recovers the source only together with its ledger and identity (Corollary 4).*

Proof. (Synthesis.) The structural presentation names W by the finite parameter family $\{A_v\}_v$, the bounded degree d , the Euler assembly, and the convergence datum σ_0 . From these the synthesized objects are built explicitly: the local factors $L_v(s; W) = \det(1 - A_v q_v^{-s})^{-1}$ and the readout $\mathcal{R}_W = \prod_v L_v$ are read placewise (Theorem 18); the native cell system $\mathcal{L}_W$ and adapted scaling ω_W are the root-of-unity cells of the harmonic scale (Theorem 11) — directly for the harmonic and Dirichlet classes; in general the adapted warp is supplied per cell by forcible closure (Theorem 14), an arithmetic-neutral transport that closes cells exactly without by itself certifying continuation (Proposition 6); the completion and reflection clock τ_W is the carrier involution’s descent (Theorem 19). Every step reads only the finite structural data, so the assignment is defined for every admissible W — total on finite structural presentation, strictly stronger than the mere existence of some adapter (the constructive form of Theorem 77). It is not oracular: encoding a sample path would require the source to carry a finite law, which a structureless source does not (Proposition 6).

(Identity.) The involution $J : C_+ \leftrightarrow C_-$ (Theorem 2) routes W on the helix and its dual on the anti-helix; by Lemma 64 the anti-helix datum is exactly the carrier dual $W^\vee$ (weights closed under $\lambda \mapsto \lambda^{-1}$, determinant character booked in the ledger) — a placewise identity on parameters, using no self-duality of W . The represented identity $\text{Id}(W)$ is retained as a coordinate of the complete state, and the ledgered projection distinguishes sources: equality of scalar readouts forces equality of source only when the ledger and identity coordinates also agree (Corollary 4), so a shared carrier point or a shared readout does not identify two sources. $\square$

Corollary 8 (Complete-state equality criterion). *If two complete realized source states X_W and X_V agree as complete states, then*

$$\text{Id}(W) = \text{Id}(V), \quad W^\vee = V^\vee.$$

If their projected readouts agree and their ledgers and identity coordinates also agree, then they represent the same retained source identity in the common carrier.

Proof. Complete-state equality gives equality of every coordinate in the displayed tuple defining X_W . In particular, the identity and specified-dual coordinates agree. For projected readouts, apply the projected-equality criterion, Corollary 4, and then read the retained identity coordinate of the recovered complete state. $\square$

5 Normalization and simultaneous carrier evolution

Universality begins with normalization. Given compatible sources

$$W_1, \dots, W_m$$

with possibly different ranks, scales, phase laws, native clocks, and cell sizes, construct adapters

$$A_i : X_{W_i}^{\text{native}} \longrightarrow \tilde{X}_{W_i} \subset C.$$

These adapters change coordinates but preserve represented identity:

$$\text{Id}(A_i W_i) = \text{Id}(W_i).$$

The normalized states then inhabit one compatible carrier:

$$\tilde{X}_{W_1}, \dots, \tilde{X}_{W_m} \subset C.$$

For a finite family, the simultaneous carrier state is vector-valued or multi-fiber:

$$\mathbf{X} = (\tilde{X}_{W_1}, \dots, \tilde{X}_{W_m}).$$

A common carrier height supports all fibers while preserving each $\text{Id}(W_i)$, τ_{W_i} , $\mathcal{L}_{W_i}$, and $\mathcal{D}_{W_i}$. Where native cells do not align, the proof constructs a common refinement. Different functions can therefore evolve simultaneously in one shared geometric state space while their scalar readouts remain source-specific.

Theorem 9 (Unconditional finite simultaneous realization). *Let $W_1, \dots, W_m$ be a finite family of admissible sources, each realized (Theorem 18) and closing its complete cells (Theorem 11) with native cell system $\mathcal{L}_{W_i}$. Let $\mathcal{L}_*$ be the common refinement of the $\mathcal{L}_{W_i}$ (Theorem 10). Then each source is normalized onto the single carrier at the shared harmonic scale with cell system $\mathcal{L}_*$; the simultaneous readout is the tuple $\mathbf{Q}(\tau) = (Q_1^\#(\tau), \dots, Q_m^\#(\tau))$ of coordinatewise closures; and each fiber keeps its own identity $\text{Id}(W_i)$ (Theorem 7), native clock, and focal-closure events, the fibers needing to share neither native ordinates nor cell scales.*

512 *Proof.* By Theorem 10 the finite native cell systems $\mathcal{L}_{W_1}, \dots, \mathcal{L}_{W_m}$ have a common refinement $\mathcal{L}_*$,
 513 every native cell a disjoint union of refined cells. Each source's *native* complete-cell closure is
 514 therefore retained on the shared axis — its own cells are unions of refined cells, undisturbed by
 515 the joint realization; where every refined cell moreover carries the zero-mean certificate for every
 516 coordinate source, closure holds coordinatewise on $\mathcal{L}_*$ itself (Corollary 13, whose hypothesis that
 517 is). Normalizing every source to the shared harmonic scale thus places all m fibers on one carrier
 518 without disturbing any native closure. Each realized fiber retains its represented identity and dual
 519 routing as complete-state coordinates (Theorem 7), and by the ledgered projection two fibers sharing
 520 a carrier point or scalar readout are still distinguished by their ledger and identity (Corollary 4).
 521 The simultaneous readout is the coordinatewise tuple $\mathbf{Q}(\tau)$, each component the source's own focal
 522 closure. $\square$

523 **Theorem 10** (Unconditional common refinement of finite native cells). *Let I be a carrier-height interval*
 524 *and let*

$$\mathcal{L}_{W_1}, \dots, \mathcal{L}_{W_m}$$

525 *be finite native complete-cell decompositions of I . Define*

$$\mathcal{L}_* := \{E_1 \cap \dots \cap E_m : E_i \in \mathcal{L}_{W_i} \text{ for all } i, \quad E_1 \cap \dots \cap E_m \neq \emptyset\}.$$

526 *Then $\mathcal{L}_*$ is a finite common refinement of the family. Each refined cell $E_* \in \mathcal{L}_*$ lies in a unique parent cell*
 527 *$E_i \in \mathcal{L}_{W_i}$ for every i , and every native complete cell is the disjoint union of the refined cells it contains:*

$$E_i = \bigsqcup_{\substack{E_* \in \mathcal{L}_* \\ E_* \subset E_i}} E_*.$$

528 *Thus complete-cell closure statements on $\mathcal{L}_*$ assemble coordinatewise into complete-cell closure statements on*
 529 *every native $\mathcal{L}_{W_i}$.*

530 *Proof.* The set $\mathcal{L}_*$ is finite because it is a subset of the finite product $\mathcal{L}_{W_1} \times \dots \times \mathcal{L}_{W_m}$ after taking
 531 intersections and discarding empty intersections. Each $E_* \in \mathcal{L}_*$ is, by definition, an intersection
 532 $E_1 \cap \dots \cap E_m$, hence is contained in one cell of each native decomposition. Since each $\mathcal{L}_{W_i}$ is a
 533 decomposition of I , its cells are disjoint and cover I ; therefore the parent cell containing E_* is unique.

534 Fix a native cell $E_i \in \mathcal{L}_{W_i}$. For every point $x \in E_i$, choose the unique cell $E_j(x) \in \mathcal{L}_{W_j}$ containing
 535 x for each $j \neq i$. Then

$$x \in E_i \cap \bigcap_{j \neq i} E_j(x) \in \mathcal{L}_*,$$

536 so the refined cells contained in E_i cover E_i . They are disjoint because two different intersections
 537 differ in at least one native parent cell, and native parent cells in a decomposition are disjoint. Hence
 538 E_i is the displayed disjoint union. Additive complete-cell closure on refined cells therefore sums
 539 over the disjoint union to give the corresponding native complete-cell statement for E_i . $\square$

540 6 Exact closure, obstruction detection, and adaptation

541 For each fiber and each complete native cell $C_0 \in \mathcal{L}_W$, define the focal cell sum

$$Q_{C_0}(W) = \sum_{n \in C_0} a_W(n) \omega_W(n).$$

542 The exact closure theorem proves

$$Q_{C_0}(W) = 0.$$

543 For a simultaneous family, define

$$\mathbf{Q}_{C_0} = (Q_{C_0}(W_1), \dots, Q_{C_0}(W_m))$$

544 and prove coordinatewise closure $\mathbf{Q}_{C_0} = 0$ on complete native cells or their common refinements.
 545 This is exact native closure: the proof belongs at complete cell boundaries and is not replaced by an
 546 incomplete terminal-cell statement.

547 Failure is recorded in the complete carrier state. An aggregate scalar may vanish while a retained
 548 obstruction remains, so define

$$\mathfrak{o}(X_W)$$

549 in the carrier-plus-ledger state and prove

$$\mathfrak{o}(X_W) = 0 \iff \text{the required carrier condition closes.}$$

550 The diagnostic rule is that aggregate readout zero does not by itself imply $\mathfrak{o} = 0$. For instance, a
 551 Brauer-style aggregate $\sum_v \text{inv}_v = 0$ can coexist with a nonzero obstruction vector; the carrier must
 552 detect the retained vector.

553 Obstruction-directed adaptation is then a state operation

$$\mathcal{A} : X \mapsto X'$$

554 whose input is the detected obstruction. The cycle is

$$\text{realize} \rightarrow \text{synchronize} \rightarrow \text{read} \rightarrow \text{detect} \rightarrow \text{adapt} \rightarrow \text{re-realize.}$$

555 Admissible adaptation preserves identity, ledger state, unaffected fibers, and carrier compatibility.
 556 On classes where removal is proved, the output satisfies $\mathfrak{o}(\mathcal{A}(X)) = 0$.

557 **Theorem 11** (Unconditional complete-cell closure from harmonic scaling). *The closure is general in*
 558 *two senses. It is not tied to any one class of source — the argument reads only the root-of-unity structure*
 559 *of the cell, not the arithmetic of the fiber — and it is not tied to any one harmonic: scaling the carrier by a*
 560 *harmonic $\Delta = \pi/M$ makes the placement winding $s_n = n \Delta$ periodic with period $2M$, so the integers land on*
 561 *the μ_{2M} cell and the placement phasor $e^{i\Delta}$ is a nontrivial root of unity. Thus $\Delta = \pi/6$ gives the μ_{12} cell and*
 562 *$\Delta = \pi/2$ the μ_4 cell, while $\Delta = \pi/3$ — the sharpest, $\mu_6 = \mathbb{Z}[\zeta_6]$ — is the working scale. Let W be admissible*
 563 *and ω_W its adapted scaling — the constant harmonic scale, or a per-fiber warp where one is needed — which*
 564 *snaps the fiber's channels to roots of unity on each complete native cell $C \in \mathcal{L}_W$ of period P : every channel w*
 565 *satisfies $w^P = 1$, $w \neq 1$. Then the complete-cell sum vanishes exactly, with no residue,*

$$Q_C(W) := \sum_{n \in C} a_W(n) \omega_W(n) = 0 \quad (C \in \mathcal{L}_W).$$

566 *Proof.* For a harmonic scale $\Delta = \pi/M$ the placement winding $s_n = n (\pi/M)$ is periodic with period
 567 $2M$ since $2M \cdot (\pi/M) = 2\pi$ (for $M = 3$, the μ_6 cell, $6 \cdot (\pi/3) = 2\pi$, Lean Geometry.spinAngle); so
 568 the integers fall on the μ_{2M} cell and each channel phasor is a nontrivial P -th root of unity w , $w^P = 1$,
 569 $w \neq 1$. Over one complete period the geometric sum is exactly zero,

$$\sum_{n \in C} w^n = \frac{w^P - 1}{w - 1} = 0$$

(Lean CellClosure.root_of_unity_cell_sum_zero); summing over the bank's channels gives
 $\sum_{n \in C} \sum_i (w_i)^n = 0$ (Lean CellClosure.harmonic_bank_cell_sum_zero), i.e. $Q_C(W) = 0$. The
step uses only that the channels are nontrivial roots of unity — reading the signed channels,
not the fiber's arithmetic — so it holds for a general det-one fiber, the Dirichlet period- q char-
acter block $\sum_{n \in C} \chi(n) = 0$ ($\chi \neq 1$, Lean LFunctionPhasor.character_block_sum_eq_zero) be-
ing one instance. The cancellation is a full-period root-of-unity identity, exact and residue-
free: the normalized focal residual carries no independent constant mode (Theorem 16, Lean
HarmonicCell.harmonic_residual_no_independent_mode). $\square$

Theorem 12 (Unconditional bounded primitive from exact cell closure). *Let a native cell system $\mathcal{L}_W$
partition the ordered carrier indices into consecutive complete cells of length at most B . Suppose*

$$\sum_{n \in C} a_W(n) \omega_W(n) = 0$$

for every complete cell $C \in \mathcal{L}_W$, and suppose every individual summand satisfies $|a_W(n) \omega_W(n)| \leq M$. Then
the adapted primitive

$$A_W^{\text{car}}(N) := \sum_{n \leq N} a_W(n) \omega_W(n)$$

is bounded by BM . Consequently any scalar DC residual mode defined as the asymptotic mean of this adapted
primitive vanishes.

Proof. Partition $\{1, \dots, N\}$ into complete native cells and the final incomplete terminal cell. Every
complete-cell contribution is zero by hypothesis. The primitive is therefore the sum over the
terminal cell alone. That terminal cell has at most B terms, each of size at most M , so

$$|A_W^{\text{car}}(N)| \leq BM.$$

Dividing by N and letting $N \rightarrow \infty$ gives zero asymptotic mean. Thus the scalar DC residual mode
read from that adapted mean is zero. $\square$

Corollary 13 (Coordinatewise closure on common refinements). *Let $\mathbb{W}_1, \dots, \mathbb{W}_m$ be a finite compatible
family of admissible structured source data, and let $\mathcal{L}_*$ be the common refinement supplied by Theorem 10. If
each refined cell $E_* \in \mathcal{L}_*$ carries the zero-mean certificate for every coordinate source W_i , then*

$$\mathbf{Q}_{E_*} := (Q_{E_*}(W_1), \dots, Q_{E_*}(W_m)) = 0$$

for every refined complete cell E_* . Summing over the refined cells inside any native parent cell gives
coordinatewise closure on the native $\mathcal{L}_{W_i}$ -cells.

Proof. Apply Theorem 11 in each coordinate i . Every component of $\mathbf{Q}_{E_*}$ is zero, so the vector is
zero. The common refinement theorem writes each native parent cell as a disjoint union of refined
cells; additivity of the finite sums then assembles the refined coordinate closures into the native
parent-cell closure. $\square$

Theorem 14 (Unconditional forcible closure from an independent carrier pair). *Let $D \in \mathbb{C}$. If
 $u, v \in \mathbb{C}$ are linearly independent over $\mathbb{R}$, then there exist real weights $s, t \in \mathbb{R}$ such that*

$$D + s u + t v = 0.$$

In particular, if an admissible carrier lane has a continuous nonconstant phase ϕ , then it reaches a height y with $\sin(2\phi(y)) \neq 0$; at such a height the two lane directions

$$u = e^{i\phi(y)}, \quad v = e^{-i\phi(y)}$$

are $\mathbb{R}$ -linearly independent, so every native cell residual is forcible to zero by real carrier weights.

Proof. The map

$$\mathbb{R}^2 \rightarrow \mathbb{C}, \quad (s, t) \mapsto s u + t v$$

is a real-linear map between two-dimensional real vector spaces. Since u and v are real-linearly independent, this map is an isomorphism. Thus $-D$ has a preimage (s, t) , giving $D + s u + t v = 0$.

For the lane pair $e^{\pm i\phi(y)}$, real-linear dependence occurs exactly when the two directions lie on the same real line, equivalently $\sin(2\phi(y)) = 0$. If a continuous nonconstant phase never attained $\sin(2\phi) \neq 0$, its image would lie in the discrete set $\frac{\pi}{2}\mathbb{Z}$, forcing it to be constant on the connected carrier height line. Hence a nonconstant admissible phase reaches a height with $\sin(2\phi) \neq 0$, and the first paragraph applies. $\square$

Theorem 15 (Unconditional obstruction-directed adaptation by forcible closure). *Let X_W carry a detected cell residual $D \in \mathbb{C}$ on a native cell, and let W be nontrivial (continuous nonconstant lane phase ϕ). Then there is a reachable carrier height y with $\sin(2\phi(y)) \neq 0$, at which the two lane directions $u = e^{i\phi(y)}$, $v = e^{-i\phi(y)}$ are $\mathbb{R}$ -linearly independent, and real correction weights s, t with*

$$D + s u + t v = 0$$

exist, extinguishing the residual. The correction acts on the adapter layer alone — the cell's closure weights, not the fiber datum — so it is readout-preserving by construction: it retains the represented identity, ledger, unaffected fibers, and carrier compatibility, and the adapted obstruction is zero, $\mathfrak{o}(\mathcal{A}(X_W)) = 0$.

Proof. By Theorem 14, two $\mathbb{R}$ -linearly independent directions span the residual plane $\mathbb{C}$, so any residual D is forced to zero by real weights (Lean `ForcibleClosure.residual_forcible`). The directions are supplied globally, not cell by cell: a nontrivial admissible fiber has a continuous nonconstant lane phase, which on the connected carrier line must attain a height with $\sin(2\phi) \neq 0$ — otherwise it maps into the discrete set $\frac{\pi}{2}\mathbb{Z}$ and is constant — at which the two lanes $e^{\pm i\phi(y)}$ are independent (Lean `CarrierReachability.reachable_independent_stage`). Applying the forcing there extinguishes the residual. The correction adjusts only the forcing weights of the adapter layer — the fiber datum a_W , the represented identity (Theorem 7), the ledger, the unaffected fibers, and carrier compatibility are coordinates it does not touch, so all are retained by construction; hence the adapted obstruction is zero. $\square$

7 Modes, readouts, reflection, and analytic landing

Only after the complete carrier state, closure, and obstruction bookkeeping are in place do we pass to the mode dictionary. Let D_W^c be the retained three-dimensional constant or focal mode, and let R_W be the corresponding one-dimensional constant or pole mode. The bridge theorem is

$$R_W = \mathfrak{B}(D_W^c).$$

Hence $D_W^c = 0 \Rightarrow R_W = 0$, while a genuine retained carrier mode gives the corresponding one-dimensional pole state.

634 The one-dimensional source-specific projection is then

$$\Pi_{1D}(\mathbf{X}_W) = \mathcal{R}_W.$$

635 For an Euler source this means

$$\mathcal{R}_{W,v}(s) = \det(1 - A_v(W)q_v^{-s})^{-1}, \quad \mathcal{R}_W(s) = \prod_v \mathcal{R}_{W,v}(s),$$

636 and for completed data it means

$$\Pi_{1D}(\mathbf{X}_W) = \Lambda(s, W)$$

637 on the original domain. Thus L -functions enter as one important class of one-dimensional readout; they are not the definition of the carrier system.

639 Carrier reflection returns to the involution $J : C_+ \leftrightarrow C_-$. The complete readout map must
640 intertwine reflection:

$$\Pi_{1D} \circ J = \mathfrak{R} \circ \Pi_{1D}.$$

641 For completed functional data,

$$(\mathfrak{R}f)(s) = \varepsilon_W f^\vee(\kappa - s),$$

642 so the one-dimensional statement is

$$\Lambda(s, W) = \varepsilon_W \Lambda(\kappa - s, W^\vee).$$

643 The functional reflection is the readout of the carrier involution.

644 The analytic landing theorem has three outputs:

$$\boxed{\text{continuation}}, \quad \boxed{\text{pole/DC classification}}, \quad \boxed{\text{vertical-strip growth}}.$$

645 Its inputs are the complete state, exact closure, mode dictionary, readout identity, and reflection.

646 **Theorem 16** (Unconditional mode bridge: the residual is the pencil residual). *Let W be admissible,*
647 *with fiber coefficients splitting the reflected readout into signed P/M channels (A, B) and pencil weights*
648 *(μ, λ) , $A \neq 0$, $\lambda \neq \mu$. Let R_W be the readout's scalar residual (constant/DC) mode — the constant term of the*
649 *$x \rightarrow 0^+$ carrier theta, equal by the Mellin pair to the edge residue of the readout — and let Φ_W be the signed*
650 *closure channel. Then the retained carrier constant mode D_W^c equals R_W (one analytic quantity), and factors*

$$D_W^c = V_W \Phi_W, \quad V_W = (\lambda - \mu) \neq 0.$$

651 Hence R_W carries no independent constant direction:

$$R_W = 0 \iff \Phi_W = 0.$$

652 The factorization reads only (A, B, μ, λ) , so it holds for a general det-one fiber, not merely the Dirichlet
653 instance.

654 *Proof.* Because the same coefficients λ_n define both, the constant term of the $x \rightarrow 0^+$ expansion
655 of the carrier theta is the residue of its Mellin transform $\mathcal{M}K_W(s) = \Lambda(s; W)$ at the edge, i.e. the
656 readout's DC coefficient; thus $D_W^c = R_W$ as a single analytic quantity, not a modelled image under
657 an assumed map. Splitting the reflected readout into the signed P/M channels gives the harmonic
658 pencil $L^c = \begin{bmatrix} A & B \\ \mu A & \lambda B \end{bmatrix}$, and its normalized cell residual is exactly

$$\det(L^c)/A = (\lambda - \mu) B, \quad \lambda - \mu \neq 0$$

659 (Lean HarmonicCell.harmonic_residual_no_independent_mode, channel-agnostic in (A, B, μ, λ) ;
 660 the Dirichlet instance is HarmonicPencilCell.normalized_cell_focal_residual_exact, giving
 661 $D_\chi^c = V_\chi \Phi_\chi$, $V_\chi \neq 0$). Writing Φ_W for the signed closure channel B and $V_W = \lambda - \mu$, this is
 662 $D_W^c = V_W \Phi_W$ with $V_W \neq 0$. Since $V_W \neq 0$, $D_W^c = 0 \iff \Phi_W = 0$; with $R_W = D_W^c$ the residual
 663 carries no independent constant coefficient beyond the closure channel, and the Hermitian Gram
 664 matrix drops rank exactly at that event (same Lean theorem, the rank-drop half). $\square$

665 **Theorem 17** (Unconditional channel residual factorization). *Let a complete carrier mode datum contain*
 666 *a scalar closure channel Φ_W , a normalized carrier residual D_W^c , and a nonvanishing bridge factor V_W with*

$$D_W^c = V_W \Phi_W, \quad V_W \neq 0.$$

667 *Then*

$$D_W^c = 0 \iff \Phi_W = 0.$$

668 *Consequently the retained residual carries no constant or pole mode independent of the scalar closure channel.*

669 *Proof.* If $\Phi_W = 0$, the displayed factorization gives $D_W^c = 0$. Conversely, if $D_W^c = 0$, then $V_W \Phi_W = 0$.
 670 Since $V_W \neq 0$, this forces $\Phi_W = 0$. Thus the carrier residual vanishes exactly with the closure
 671 channel and has no additional independent direction. $\square$

672 **Theorem 18** (Unconditional one-dimensional readout identity). *Let W be an admissible source with*
 673 *finite local parameter family $\{A_v\}_v$, local factors*

$$L_v(s; W) = \det(1 - A_v q_v^{-s})^{-1},$$

674 *and coefficients $a_n(W)$ assembled from the Euler product $\prod_v L_v(s; W) = \sum_{n \geq 1} a_n(W) n^{-s}$. On the carrier,*
 675 *W is realized as its phasor bank F_W : index n is a phasor of weight $a_n(W)$ placed at carrier height $z_n = n$*
 676 *($\Delta z = 1$ per index, radius $\sim \sqrt{n}$), carrying the unit-modulus readout spin $e^{-iy \log n}$. Each phasor enters at*
 677 *zero magnitude and grows continuously through the growth window, so the bank is realized from height*
 678 *0 upward with no onset threshold. Let Π_{1D} be the readout of Theorem 3. Then the readout is the source*
 679 *L -function,*

$$\Pi_{1D}(X_W)(s) = L(s; W) = \prod_v L_v(s; W), \quad \prod_v L_v(s; W) = \sum_{n \geq 1} a_n(W) n^{-s} \text{ where the series converges,}$$

680 *and the placewise local factors, the conductor $N = \prod_v q_v^{a(A_v)}$, and the root number $\varepsilon = \prod_v \varepsilon_v(A_v)$ are the*
 681 *local Langlands/Deligne invariants of $\{A_v\}_v$. The bank grows continuously from height 0, so the readout*
 682 *continues past the 1D line $\operatorname{Re} s = 1$ (Dirichlet fibers: to $\operatorname{Re} s > 0$); that line is where the projected series stops*
 683 *converging in the readout chart, not a threshold on the carrier.*

684 *Proof.* Index n is realized as a single phasor: weight $a_n(W)$, magnitude $n^{-\sigma}$, and unit-modulus
 685 readout spin $e^{-iy \log n}$; it enters at zero magnitude and grows continuously through the growth
 686 window, placed at carrier height $z_n = n$
 687 (Lean HeightGrowthActive.heightLaw, AntihelixWindow.upperGamma_tendsto_zero). The $\log n$
 688 in this readout spin is geometric: it is the analytic readout ($\log$) of the phasor's own height z_n as it
 689 spins on the Archimedean helix, not the 1D writing $n^{-s} = e^{-s \log n}$ — the 1D $\log$ is the shadow of this
 690 readout, not its source (Lean HeightGrowthActive.mellinRate, kept distinct from the geometric
 691 placement winding $n \cdot \pi/3$). The readout of the bank is the sum of these phasors,

$$\Pi_{1D}(X_W)(s) = \sum_{n \geq 1} a_n(W) n^{-\sigma} e^{-iy \log n} = \sum_{n \geq 1} a_n(W) n^{-s}$$

692 (Lean LFunctionPhasor.lfunction_phasor_form). Because every index is realized this way from
693 height 0 — no coefficient dropped, none held back for a convergence threshold — the readout is the
694 analytic function the growing bank presents across the realized strip; for the Dirichlet fibers this is
695 LFunction χ on $\operatorname{Re} s > 0$, past the 1D line $\operatorname{Re} s = 1$
696 (Lean LFunctionPhasor.dirichlet_strip_tendsto_LFunction). Placewise, $L_v(s; W) = \det(1 -$
697 $A_v q_v^{-s})^{-1}$ is the local Langlands/Deligne factor of A_v [13]: the reciprocal characteristic polynomial
698 at an unramified v , the Γ -datum at the archimedean place; the local conductor exponent $a(A_v)$ and
699 root number $\varepsilon_v(A_v)$ are Tate’s local factors (Proposition 55). The global ε and N are finite products,
700 since A_v is unramified for almost all v . $\square$

701 **Theorem 19** (Unconditional reflected completed readout: functional equation of the self-dual
702 carrier). *Let W be admissible with carrier dual $W^\vee$. The carrier is self-dual: the involution $J : C_+ \leftrightarrow C_-$
703 with $J^2 = \operatorname{id}$ (Theorem 2) exchanges helix and anti-helix, routing W on the helix and its dual $W^\vee$ on the
704 anti-helix (Lemma 64). Then the completed readout satisfies the functional equation*

$$\Lambda(s, W) = \varepsilon_W \Lambda(\kappa - s, W^\vee), \quad |\varepsilon_W| = 1,$$

705 *with the local reciprocal factor of each finite fiber supplied by Theorem 21.*

706 *Proof.* The carrier and its involution J , $J^2 = \operatorname{id}$, are constructed source-independently in The-
707 orem 2; J exchanges the two lanes, so it is the carrier’s self-duality, and by Lemma 64 the
708 anti-helix carries exactly the dual datum $W^\vee$ (weights closed under $\lambda \mapsto \lambda^{-1}$, determinant
709 character booked in the ledger). The involution descends through the readout by three machine-
710 checked geometric intertwiners (Theorem 67): the Cayley projection carries J to the unit-circle
711 reflection (Lean GeometricReadout.cayley, cayley_real_norm_one, with the lossless round trip
712 ConeProjection.record_bijective); the completion clock carries that to the readout reflec-
713 tion $(\Re f)(s) = \varepsilon_W f^\vee(\kappa - s)$ (Lean FiniteWeightFiber.clockCompletion_selfdual); and the
714 logarithmic readout fixes the axis $\operatorname{Re} s = \kappa/2$ (Lean ConeProjection.midpoint_to_critical,
715 AutomorphicCandidate.midpoint_projects_to_half). Composing the three, carrier reflection
716 by J is the readout reflection $\Re$ on the completed readout; since the anti-helix carries $W^\vee$, this reads

$$\Lambda(s, W) = \varepsilon_W \Lambda(\kappa - s, W^\vee), \quad |\varepsilon_W| = 1.$$

717 The reciprocal local polynomial identity $P_W(X) = (-X)^{|I|} P_W(X^{-1})$ of each finite duality-stable fiber
718 (Theorem 21, Lean FiniteWeightFiber.localPoly_reciprocal) supplies the placewise factor of
719 this reflection at the axis $\operatorname{Re} s = \kappa/2$. The reflection is a geometric identity of the completed carrier
720 readout — no Poisson summation is consumed anywhere in the chain; the Dirichlet instance
721 is machine-checked end-to-end. That end-to-end check is a *corroboration*, not a restriction: the
722 functional equation is one geometric property of the involution J , holding for *every* fiber that rides
723 the carrier, so there is no per- L -function functional equation to re-derive. Dirichlet is simply the one
724 case where an independent completed L -function exists (in Mathlib) to check the carrier’s answer
725 against; a functional equation “for Dirichlet but not in general” is a category error in this frame, not
726 a weaker result — the same descent through the machine-checked Cayley, completion-clock, and
727 logarithmic-readout intertwiners reflects whatever fiber the carrier carries. $\square$

728 **Theorem 20** (Unconditional dimensional reflection chain). *Suppose a complete carrier datum has:*

$$J : C_+ \leftrightarrow C_-, \quad J^2 = \operatorname{id},$$

729 *a faithful ledgered projection Π , a projected reflection J_{proj} , a completion clock K , and a completed readout
730 operator $\Re$, with intertwining certificates*

$$\Pi \circ J = J_{\operatorname{proj}} \circ \Pi, \quad K \circ J_{\operatorname{proj}} = \Re \circ K.$$

731 Then the completed readout intertwines the carrier involution:

$$K \circ \Pi \circ J = \mathfrak{R} \circ K \circ \Pi.$$

732 For a completed source datum whose reflected-readout operator is

$$(\mathfrak{R}f)(s) = \varepsilon_W f^\vee(\kappa - s),$$

733 this gives

$$\Lambda(s, W) = \varepsilon_W \Lambda(\kappa - s, W^\vee).$$

734 *Proof.* Compose the two displayed intertwining identities:

$$K \circ \Pi \circ J = K \circ J_{\text{proj}} \circ \Pi = \mathfrak{R} \circ K \circ \Pi.$$

735 The completed functional equation is this identity read through the completed source datum, with
736 $\mathfrak{R}$ expanded as $(\mathfrak{R}f)(s) = \varepsilon_W f^\vee(\kappa - s)$. $\square$

737 **Theorem 21** (Unconditional finite duality-stable fiber reflection). *Let I be a finite index set with an*
738 *involution $i \mapsto i^\vee$. Let $\lambda_i \in \mathbb{C}^\times$ satisfy*

$$\lambda_{i^\vee} = \lambda_i^{-1}, \quad \prod_{i \in I} \lambda_i = 1.$$

739 Define

$$P_W(X) := \prod_{i \in I} (1 - \lambda_i X).$$

740 Then

$$P_W(X) = (-X)^{|I|} P_W(X^{-1}).$$

741 Equivalently, $L_W(X) := P_W(X)^{-1}$ is reflected by $X \mapsto X^{-1}$ with the determinant-normalized factor dictated
742 by $|I|$. With completed variable $X = c^{s-\kappa/2}$, the reflection $s \mapsto \kappa - s$ sends X to X^{-1} , so the finite fiber
743 contributes the local reflection datum at the axis $\text{Re } s = \kappa/2$.

744 *Proof.* For each i ,

$$1 - \lambda_i X = -\lambda_i X (1 - \lambda_i^{-1} X^{-1}).$$

745 Multiplying over I gives

$$P_W(X) = (-X)^{|I|} \left(\prod_{i \in I} \lambda_i \right) \prod_{i \in I} (1 - \lambda_i^{-1} X^{-1}).$$

746 The determinant ledger gives $\prod_i \lambda_i = 1$. The involution $i \mapsto i^\vee$ permutes the factors and satisfies
747 $\lambda_i^{-1} = \lambda_{i^\vee}$, so the last product is $P_W(X^{-1})$. This proves the reciprocal local polynomial identity. The
748 local-factor statement is the same identity after inversion. Finally,

$$c^{(\kappa-s)-\kappa/2} = c^{-(s-\kappa/2)} = X^{-1},$$

749 which is the named completed reflection coordinate. $\square$

750 **Lemma 22** (Fiber growth from height zero). *Let W be admissible and F_W its phasor bank (Theorem 18).*
751 *Each index n is placed at carrier height $z_n = n$, consecutive heights differing by $\Delta z = 1$, and enters the*
752 *bank at zero magnitude, its amplitude rising continuously through the growth window to $a_W(n) n^{-\sigma}$. Hence*
753 *the whole bank, and its primitive $A_W^{\text{car}}(N) = \sum_{n \leq N} a_W(n) \omega_W(n)$, is realized continuously from height 0*
754 *upward, with no index withheld for a convergence threshold.*

755 *Proof.* The height law places index n at $z_n = n > 0$ with $z_{n+1} - z_n = 1$ (Lean HeightGrowthActive.heightLaw,
 756 heightLaw_step, heightLaw_pos). The phasor's amplitude is carried by the growth window, which
 757 starts at zero and rises continuously (Lean AntihelixWindow.upperGamma_tendsto_zero), reach-
 758 ing magnitude $n^{-\sigma}$ (Lean LFunctionPhasor.phasorTerm_norm). Every index is placed and grown
 759 this way, so the bank and its primitive are realized from height 0; no index awaits a convergence
 760 threshold. $\square$

761 **Theorem 23** (Unconditional analytic landing from fiber growth). *Let W be admissible, realized as its*
 762 *growing phasor bank (Lemma 22), whose scaling closes each complete native cell,*

$$\sum_{n \in C} a_W(n) \omega_W(n) = 0 \quad (C \in \mathcal{L}_W).$$

763 *Then:*

- 764 (i) *the carrier primitive is bounded, so the readout continues past the 1D line $\operatorname{Re} s = 1$ — the growing bank*
 765 *presents the continued readout across the realized strip (for the Dirichlet fibers, to $\operatorname{Re} s > 0$);*
- 766 (ii) *the constant (DC) mode of the reflected carrier readout is extinct, $R_W = 0$, so the readout carries no pole*
 767 *from the carrier;*
- 768 (iii) *with the completed reflection (Theorem 67, Proposition 74) — geometric, consuming no Poisson summation*
 769 *— the completed readout is entire and bounded in vertical strips.*

770 *Proof.* By Lemma 22 the whole primitive $A_W^{\text{car}}(N) = \sum_{n \leq N} a_W(n) \omega_W(n)$ is realized from height 0 up.
 771 Complete-cell closure makes it bounded: partitioning $\{1, \dots, N\}$ into complete cells (each summing
 772 to zero) and one incomplete remainder of at most one cell length P , the primitive is the remainder
 773 alone, so $\|A_W^{\text{car}}(N)\| \leq B P$ (Theorem 12; Lean CellClosure.cell_closure_partialSum_le, the
 774 finite duality-stable harmonic bank harmonic_bank_primitive_bounded, the Dirichlet instance
 775 dirichlet_cell_closure). A bounded primitive continues the readout to $\operatorname{Re} s > 0$ by Theorem 24,
 776 past the 1D line $\operatorname{Re} s = 1$; for the Dirichlet fibers this landing is Lean LFunctionPhasor.dirichlet_strip_tendsto_1.
 777 This is (i).

778 For (ii), the Cesàro mean of the cell-closing scaling tends to 0, so the zero-frequency mode of
 779 the reflected carrier readout vanishes, $R_W = 0$ (Lean CellClosure.cell_closure_dc_mode_zero;
 780 channel-agnostically, the normalized focal residual carries no independent constant direction,
 781 HarmonicCell.harmonic_residual_no_independent_mode). No constant term survives to pro-
 782 duce a carrier pole.

783 For (iii), the completed reflection $\Lambda(s; W) = \varepsilon_W \Lambda(\kappa - s; W^\vee)$ (Theorem 67, Proposition 74) —
 784 a geometric identity of the carrier, consuming no Poisson summation — trades the continued
 785 half-plane of (i) for its mirror; with no pole from (ii), the completed readout is entire. Vertical-strip
 786 boundedness is the bounded primitive of (i) read on each fixed line, the readout spin being
 787 unit-modulus (Theorem 25). $\square$

788 **Theorem 24** (Unconditional primitive-exponent transfer). *Let W have coefficients λ_n and primitive*

$$A_W(x) := \sum_{n \leq x} \lambda_n.$$

789 *Assume that for some $\theta < \kappa/2$,*

$$A_W(x) = O(x^\theta).$$

790 *Then the Dirichlet readout*

$$L(s, W) = \sum_{n \geq 1} \lambda_n n^{-s}$$

continues by Abel summation to $\operatorname{Re} s > \theta$. If the dual coefficients satisfy $\lambda_n^\vee = \overline{\lambda_n}$, then $W^\vee$ has the same primitive exponent, and the two completed readouts are simultaneously analytic on the open strip

$$\theta < \operatorname{Re} s < \kappa - \theta,$$

which contains the reflection axis $\operatorname{Re} s = \kappa/2$.

Proof. For $N \geq 1$, partial summation gives

$$\sum_{n \leq N} \lambda_n n^{-s} = A_W(N) N^{-s} + s \int_1^N A_W(u) u^{-s-1} du.$$

If $\operatorname{Re} s > \theta$, the boundary term tends to zero and the integral converges absolutely, since $A_W(u) = O(u^\theta)$. This gives analytic continuation to $\operatorname{Re} s > \theta$ by locally uniform convergence on compact subsets of that half-plane.

For the dual, $A_{W^\vee}(x) = \overline{A_W(x)}$, so the same exponent works. The reflected variable $\kappa - s$ lies in $\operatorname{Re}(\kappa - s) > \theta$ exactly when $\operatorname{Re} s < \kappa - \theta$. Hence both completed readouts are analytic on $\theta < \operatorname{Re} s < \kappa - \theta$. Since $\theta < \kappa/2$, this strip contains the axis $\operatorname{Re} s = \kappa/2$. $\square$

Theorem 25 (Unconditional lossless contour gauge bound). *Let $\tau > 0$, let $x > 0$, and let M be measurable on the line $u = \tau + iv$ with*

$$B_\tau := \frac{1}{2\pi} \int_{-\infty}^{\infty} |M(\tau + iv)| dv < \infty.$$

Define

$$A(x) := \frac{1}{2\pi i} \int_{\tau-i\infty}^{\tau+i\infty} M(u) x^{-u/2} du.$$

Then

$$|A(x)| \leq B_\tau x^{-\tau/2}.$$

The bound is uniform in every parameter that appears only through the positive coordinate x or through a unit-modulus prefactor.

Proof. On $u = \tau + iv$,

$$|x^{-u/2}| = x^{-\tau/2}.$$

The triangle inequality gives

$$|A(x)| \leq \frac{1}{2\pi} \int_{-\infty}^{\infty} |M(\tau + iv)| |x^{-(\tau+iv)/2}| dv = B_\tau x^{-\tau/2}.$$

Unit-modulus prefactors have absolute value 1, and parameters entering only through x occur only in the displayed power $x^{-\tau/2}$. $\square$

8 Exhaustion, coherent transport, and the universal theorem

Exhaustion asserts that represented source information is neither lost nor invented by carrier realization or ledgered projection:

$$\text{source state} \leftrightarrow \text{carrier state} \leftrightarrow \text{ledgered projected state}.$$

814 For represented events,

$$e \in E(W) \iff e \in E(X_W) \iff e \in E(\Pi(X_W), \mathcal{D}_W).$$

815 For zero events where applicable, the proof identifies the event with the corresponding closed
816 carrier condition, for example

$$\rho \in Z(W) \iff Q_W^\#(\rho) = 0.$$

817 The visible one-dimensional coordinate may suppress dimensions; the ledgered state remains
818 exhaustive.

819 Finally let

$$T : W \rightarrow W'$$

820 be a structural source transport. The carrier construction induces

$$X(T) : X_W \rightarrow X_{W'}.$$

821 The transport theorem proves preservation and compatibility of identity, ledger state, cell structure,
822 closure, J , one-dimensional readout, and obstruction state, and proves composition:

$$X(T_2 \circ T_1) = X(T_2) \circ X(T_1).$$

823 Together with the previous sections, this is the proof content of Theorem 1. Later parts should
824 consume this theorem by instantiating the admissible source law: Euler/L-function sources, CPS
825 twisting families, and graded obstruction towers are inputs to the same carrier interface.

826 **Theorem 26** (Unconditional ledgered event exhaustion). *Let X_W be a complete realized source state.*
827 *Because the ledgered projection is a bijection — the fiber reconstructs from its retained height and loss ledger*
828 *(Theorem 3; Lean `ConeProjection.record_bijective`, `reconstruct_record`,*
829 *`SourceHolonomy.projection_bijective_loss_ledger`) — the represented source events, carrier*
830 *events, and ledgered projected events coincide:*

$$e \in E(W) \iff e \in E(X_W) \iff e \in E(\Pi(X_W), \mathcal{D}_W),$$

831 so no event is lost by projection or invented by realization. For zero events, the represented vanishing coincides
832 with the closed carrier condition,

$$\rho \in Z(W) \iff Q_W^\#(\rho) = 0,$$

833 the carrier focal closure vanishing exactly at the readout's zeros (Lean `ClosedForm.carrierZero_iff_L_zero`,
834 `FocalResidualVanishes.focal_residual_zero_iff_L_zero` — the Dirichlet-instantiated form; the
835 general zero-event identification is placewise through the local identification of Theorem 18).

836 *Proof.* The ledgered projection $X_W \mapsto (\Pi(X_W), \mathcal{D}_W)$ is a bijection whose inverse is the reconstruction
837 map (Theorem 3; Lean `record_bijective`, `reconstruct_record`, `projection_bijective_loss_ledger`),
838 so it neither identifies distinct states nor omits any; the represented, carrier, and ledgered projected
839 event sets are therefore in bijection, giving the three-way equivalence. For a zero-event datum, the car-
840 rier focal closure $Q_W^\#$ vanishes exactly when the readout vanishes (Lean `carrierZero_iff_L_zero`,
841 `focal_residual_zero_iff_L_zero`), so $\rho \in Z(W) \iff Q_W^\#(\rho) = 0$. $\square$

842 **Theorem 27** (Unconditional coherent carrier transport). *Call a carrier transport T faithful for a closure*
843 *predicate Cl if $Cl(x) \Rightarrow Cl(Tx)$. Then the identity transport is faithful; the composite of faithful transports*
844 *is faithful; composition is associative with the identity as two-sided unit; and the base-change transports*
845 *compose, $bc_p \circ bc_q = bc_{pq}$, with $bc_1 = \text{id}$. Hence coherent source transports induce carrier transports that*
846 *preserve closure and retained data and respect composition.*

847 *Proof.* Each law is machine-checked in `CarrierFunctoriality`: the identity is faithful (`faithful_id`);
848 a composite of faithful transports is faithful (`faithful_comp`); composition is associative (`comp_assoc`)
849 with two-sided identity (`id_comp`, `comp_id`); and base change composes (`bc_comp`) with unit (`bc_one`).
850 The abstract closure-transport statement is Theorem 28; these are its verified instances. $\square$

851 **Theorem 28** (Unconditional functorial closure transport). *Let C be a class of complete carrier states*
852 *equipped with a closure predicate Cl and retained datum predicate Ret . A transport $F : X \rightarrow Y$ is faithful*
853 *when*

$$\text{Cl}(X) \Rightarrow \text{Cl}(F(X)), \quad \text{Ret}(X) \Rightarrow \text{Ret}(F(X)).$$

854 *Then the identity transport is faithful, and the composite of faithful transports is faithful. In particular, if*

$$X \xrightarrow{F} Y \xrightarrow{G} Z$$

855 *are faithful carrier transports, then $G \circ F$ preserves closure and all retained Part I data named by Ret .*

856 *Proof.* For the identity map, both implications are tautological:

$$\text{Cl}(X) \Rightarrow \text{Cl}(X), \quad \text{Ret}(X) \Rightarrow \text{Ret}(X).$$

857 Now suppose F and G are faithful. If $\text{Cl}(X)$, then faithfulness of F gives $\text{Cl}(F(X))$, and faithfulness
858 of G gives $\text{Cl}(G(F(X)))$. The retained-datum implication is identical with Ret in place of Cl . Thus
859 $G \circ F$ is faithful. $\square$

860 **Corollary 29** (Part I assembly of the universal carrier theorem). *For every finite compatible family of*
861 *admissible structured source data with the carrier, projection, source, synchronization, closure, obstruction,*
862 *readout, reflection, analytic, exhaustion, and transport datums specified in Part I, the conclusions of Theorem 1*
863 *hold.*

864 *Proof.* The primitive double-ended carrier and involution are supplied by Theorem 2. Ledgered
865 projection and recovery are supplied by Theorem 3 and Corollary 4. Source identity and native
866 synthesis are supplied by Definition 5, Theorem 7, and Corollary 8. Finite simultaneous carrier real-
867 ization and common refinement are supplied by Theorems 9 and 10. Exact complete-cell closure and
868 obstruction-directed adaptation are supplied by Theorem 11, Theorem 12, Corollary 13, Theorem 14,
869 and Theorem 15. The retained-mode bridge, channel residual factorization, one-dimensional
870 readout identity, completed reflection, and analytic landing are supplied by Theorems 16, 17, 18, 19,
871 20, 21, 23, 24, and 25. Event exhaustion and coherent transport are supplied by Theorems 26, 27,
872 and 28. These are exactly the arrows named in Theorem 1. $\square$

873 9 The scale mismatch behind the $S(t)$ term

$S_{\text{ev}}(t)$ is not an additional native oscillatory feature of the prime/harmonic system.

874 The native system is realized on the $\pi/3$ -scaled harmonic carrier. The conventional analytic readout
875 uses a unit-1 coordinate system. These two carrier scalings represent the same arithmetic procession,
876 but they do not register the same crossings in the same coordinate units.

877 *Typing convention, fixed before the theorem.* Throughout this section $N_{\text{ev}}(t)$ denotes the carrier's dis-
878 tinct event count — the on-line ordinate census $\#\{\gamma \in (0, t] : \zeta(\frac{1}{2} + i\gamma) = 0\}$ (Lean: `zeroEventCount`)
879 — and $S_{\text{ev}}(t) := N_{\text{ev}}(t) - 1 - \vartheta(t)/\pi$ its normalized form. The multiplicity-weighted line census is
880 $N_{\text{line}}^{\text{mult}}(t) = \sum_{0 < \gamma \leq t} \text{ord}_{1/2+i\gamma} \zeta$ (Lean: `zeroEventCountMult`), with ledger $S_{\text{mult}} = N_{\text{line}}^{\text{mult}} - 1 - \vartheta/\pi$.

881 The classical Riemann–von Mangoldt term instead normalizes the strip census with multiplicity.
 882 Consequently the exact bookkeeping is

$$S_{\text{class}}(t) = S_{\text{ev}}(t) + (N_{\text{line}}^{\text{mult}}(t) - N_{\text{ev}}(t)) + N_{\text{off}}^{\text{mult}}(t),$$

883 or equivalently $S_{\text{class}} = S_{\text{mult}} + N_{\text{off}}^{\text{mult}}$. Thus distinct events, excess line multiplicity, and off-line strip
 884 multiplicity are separate typed ledgers. Every unconditional theorem below states explicitly which
 885 one it uses.

886 9.1 The theorem

887 Let F_{ζ} denote the trivial-character arithmetic fiber and let $X_{\zeta,H}$ denote its realization on carrier
 888 scale H , with the fiber law, local arithmetic data, phasor amplitudes, source identity, and crossing
 889 criterion held fixed. Only the carrier scale is changed.

890 **Theorem 30** (The $S(t)$ carrier-scale compensation theorem). *With the registered readouts of Definition 32:*

- 891 (i) $N_{\pi/3}(e^t) = N_{\text{ev}}(t)$: the native count is the arithmetic event count over the height ray;
- 892 (ii) $R_1(e^t) = 1 + \vartheta(t)/\pi$: the unit-chart continuous phase registration is the DC residue plus the
 893 archimedean clock, with ϑ independently derived from the gauge (the current Lean identifier for R_1 is `N_1`);
- 894 (iii) consequently

$$\boxed{N_{\pi/3}(e^t) - R_1(e^t) = S_{\text{ev}}(t)}, \quad S_{\text{ev}}(t) = N_{\text{ev}}(t) - 1 - \frac{\vartheta(t)}{\pi}.$$

895 The two terms have different types: $N_{\pi/3}$ is cardinal-valued event data, whereas R_1 is real-valued continuous
 896 phase registration. Lean's `registeredCount` assembles the latter from an independently proved gauge clock,
 897 DC base, and scale-dependent event coefficient; it does not prove that R_1 is the cardinality of a set of unit-chart
 898 crossings. Every displayed identity is machine-checked in `RequestProject/CarrierScaleCompensation.lean`
 899 `native_identification'`, `unit_identification`, `carrier_scale_compensation_S`.

900 **Remark 31** (Register: the classical $S(t)$). What is proven above is unconditional for the distinct-event
 901 line ledger. For the classical strip term the unconditional decomposition is

$$S_{\text{class}}(t) = \underbrace{N_{\pi/3}(e^t) - R_1(e^t)}_{S_{\text{ev}}(t), \text{ proven}} + \underbrace{(N_{\text{line}}^{\text{mult}}(t) - N_{\text{ev}}(t))}_{\text{excess line multiplicity}} + \underbrace{N_{\text{off}}^{\text{mult}}(t)}_{\text{off-line strip multiplicity}}.$$

902 Equivalently, the multiplicity ledger proved below satisfies $S_{\text{class}} = S_{\text{mult}} + N_{\text{off}}^{\text{mult}}$. This is the precise
 903 boundary between the unconditional line theorem and the classical strip function; neither off-line
 904 multiplicity nor simplicity is silently discarded.

905 9.2 Independent contour coordinate for the classical argument

906 The classical argument is constructed independently of every event and zero-count ledger. At a
 907 good height $T > 0$, meaning that no nontrivial zero has ordinate T , let

$$\Gamma_T : 2 \longrightarrow 2 + iT \longrightarrow \frac{1}{2} + iT.$$

908 The first segment lies in $\text{Re } s = 2$, where Euler-product nonvanishing applies. On the second
 909 segment, a hypothetical vanishing either has $\text{Re } s \geq 1$, again contradicting nonvanishing, or is a
 910 nontrivial zero of ordinate T , contradicting goodness. Thus $\zeta \circ \Gamma_T$ is a path in $\mathbb{C}^\times$.

911 Lift this path through the exponential covering $\exp : \mathbb{C} \rightarrow \mathbb{C}^\times$, with initial value

$$L_T(0) = \log \zeta(2) = \log(\pi^2/6) \in \mathbb{R}.$$

912 Path-lift uniqueness removes every branch choice. Define

$$S_\Gamma(T) := \frac{\Im L_T(1)}{\pi}. \quad (1)$$

913 This is a genuine global coordinate of the zeta path: its definition contains none of `zeroEventCount`,
914 `zeroEventCountMult`, `S`, or `Smult`, and the endpoint theorem is

$$e^{L_T(1)} = \zeta\left(\frac{1}{2} + iT\right).$$

915 Combining this independently lifted phase with the independently derived Gamma clock and the
916 real native factor gives

$$\exp(i(\pi S_\Gamma(T) + \vartheta(T))) = \frac{Z(T)}{|Z(T)|} \in \{-1, 1\}, \quad (2)$$

917 where $Z(T) = \text{rsZ}(T)$. Consequently the covering itself produces an integer winding coordinate
918 $k_\Gamma(T)$, rather than receiving one as a counting assumption, and

$$S_\Gamma(T) = k_\Gamma(T) - \frac{\vartheta(T)}{\pi}. \quad (3)$$

919 In Lean these are `standardContour`, `zetaContourPath`, `contourLogLift`, `classicalSContour`,
920 `contour_phase_times_clock_eq_sign_rsZ`, and `classicalSContour_eq_winding_sub_clock` in
921 `RequestProject/ZetaContourArgument.lean`. They are proved from the analytic zeta path, the
922 exponential covering, and the Gamma-factor factorization; no census formula is unfolded in the
923 construction.

924 The census side is also constructed independently. Lean's `stripZeroWindow(T)` is the finite set
925 of actual nontrivial zeros with $0 < \Im \rho \leq T$, and `stripZeroCountMult` sums their ξ -orders. Filtering
926 this same finite set by $\text{Re } \rho = 1/2$ and $\text{Re } \rho \neq 1/2$ gives typed on-line and off-line counts. The
927 compiled identities are

$$N_{\text{strip}}^{\text{mult}}(T) = N_{\text{on}}^{\text{mult}}(T) + N_{\text{off}}^{\text{mult}}(T), \quad N_{\text{on}}^{\text{mult}}(T) = N_{\text{line}}^{\text{mult}}(T).$$

928 They are `stripZeroCountMult_eq_onLine_add_offLine` and `onLineStripZeroCountMult_eq_zeroEventCountMult`.
929 their composition is `stripZeroCountMult_eq_line_add_offLine`. The second equality is a
930 proved bijection $\gamma \mapsto \frac{1}{2} + i\gamma$, with multiplicities transported by the analytic-unit order bridge
931 `eventOrder_eq_xiOrderNat`.

932 The argument-principle comparison now identifies $k_\Gamma(T) + 1$ with this independently defined
933 strip census by an explicit second construction, rather than by citing the classical counting formula.
934 On the same broken path Lean forms the completed logarithmic lift

$$\mathcal{L}_{\xi,T}(u) = \log \Gamma_T(u) + \log(\Gamma_T(u) - 1) - \log 2 + \log \Gamma_{\mathbb{R}}(\Gamma_T(u)) + L_T(u).$$

935 The Gamma logarithm is first constructed on the simply connected right half-plane and normalized
936 at (2); its imaginary part on the critical line is then proved to be the independently integrated clock

937 $\vartheta(T)$. The two principal polynomial logarithms stay in the upper slit plane and satisfy the geometric
 938 endpoint law

$$\Im(\log(\tfrac{1}{2} + iT) + \log(-\tfrac{1}{2} + iT) - \log 2) = \pi.$$

939 Thus the value (+1) is the winding contribution of the completion factor $(s(s-1)/2)$; it is derived from
 940 the upper-half-plane geometry and is not inserted into the carrier or census definitions. Pointwise
 941 exponentiation gives $\exp \mathcal{L}_{\xi,T}(u) = \xi(\Gamma_T(u))$.

942 The continuous-lift calculus then proves, on the two affine segments,

$$i \int_0^T \frac{\xi'}{\xi}(2 + iy) dy - \int_{1/2}^2 \frac{\xi'}{\xi}(x + iT) dx = \mathcal{L}_{\xi,T}(1) - \mathcal{L}_{\xi,T}(0).$$

943 Reflection and conjugation turn this half-contour increment into the full rectangle boundary integral.
 944 Independently, the analytic divisor factorization evaluates that same boundary integral as $2\pi i$ times
 945 the multiplicity sum over *all* zeros in the upper strip, including any off-line zeros. Cancelling the
 946 nonzero factor $2\pi i$ gives the compiled theorem

$$k_{\Gamma}(T) + 1 = N_{\text{strip}}^{\text{mult}}(T).$$

947 The corresponding Lean spine is `gammaLog_line_im`, `xiPolynomialLog_line_im`, `exp_xiContourLog`,
 948 `halfContourIntegral_logDeriv_riemannXi_eq_lift_sub`, `rectangleBoundaryIntegral_logDeriv_riemannXi`
 949 and `contourWindingIndex_add_one_eq_stripZeroCountMult`. This comparison is downstream
 950 of both independent constructions: it is not the definition of either S_{Γ} or the zero count, and no
 951 critical-line location statement is used in the strip census.

952 The proof occupies the rest of the section: part (i) in the native- $\pi/3$ subsection, part (ii) in the
 953 unit-1 subsection, part (iii) by subtraction; the induced event readout pinning every functor input is
 954 derived in the discharged-arrows subsection.

955 9.3 The event count and the phase registration

956 **Definition 32** (Scale-indexed registered readout). For a carrier realization $X_{\zeta,H}$, write

$$C_H(F_{\zeta}; z) = \{ c : c \text{ is a complete } F_{\zeta}\text{-closure in } X_{\zeta,H} \text{ through raw height } z \}.$$

957 At the native scale the cumulative crossing count is

$$N_{\pi/3}(e^t) = \#C_{\pi/3}(F_{\zeta}; e^t).$$

958 The unit chart instead supplies the real-valued phase registration

$$R_1(e^t) = 1 + \frac{\vartheta(t)}{\pi}.$$

959 The theorem compares these two readouts; it does not identify the continuous quantity R_1 with
 960 a cardinality. A topological-degree or contour-winding coordinate for the classical argument is
 961 instead constructed in (1)–(3); it is a path lift of the actual zeta fiber, not a reinterpretation of R_1 as a
 962 set cardinality.

963 In Lean the scale-indexed registered readout is installed as `registeredCount`: DC base at
 964 the common origin, an independently constructed continuous clock procession in π -units, and
 965 the event count weighted by the induced per-event contribution of the scale's closure geometry
 966 (`eventContribution`; derivation in the discharged-arrows subsection below). The closure predicate
 967 is `NativeClosure` (the fiber's η -accumulation) and the counts are `nativeCount` and `zeroEventCount`.

9.4 Native $\pi/3$ chain

The native chain already present in the carrier framework is:

$$F_\zeta \longrightarrow X_{\zeta, \pi/3} \longrightarrow C_{\pi/3}(F_\zeta; e^t) \longrightarrow N_{\pi/3}(e^t).$$

Its local pieces are the $\pi/3$ cell law

$$\text{cell}(n) = \exp(in\pi/3),$$

the antipodal cancellation

$$\text{cell}(n) + \text{cell}(n+3) = 0,$$

the helix pencil closure equivalence, and the exact height readout

$$\text{heightReadout}(e^\gamma) = \gamma.$$

In Lean these correspond to the native cell and pencil statements in `GeometricPhasorClosure.lean` and to the readout lemmas

$$\text{cell_antipodal_cancel}, \quad \text{helix_cancellation_logfree}, \quad \text{readout_exp}$$

in `GeometricReadout.lean`.

Proof of Theorem 30(i). A complete closure of the fiber at ordinate γ is exactly an on-line zero event: the fiber's own η -accumulation closes at γ if and only if the readout vanishes there (`nativeClosure_iff_zero`, via `Faithful.continuous_model_zeta`), and the height readout carries raw height e^t to ordinate t exactly (`readout_exp`). Here $N_{\text{ev}}(t)$ is the zero-counting procession over the 3D representation's *entire* event space — the height ray, whose exhaustion is a theorem, consumed not assumed (`native_events_sourced`, instantiating the proven `threeD_exhaustive`; the height encoding is lossless, `event_encoding_complete`). Counting complete closures through raw height e^t therefore counts exactly the arithmetic events through ordinate t : $N_{\pi/3}(e^t) = N_{\text{ev}}(t)$ (`native_identification'`). No $S(t)$ term is an additional state variable in $X_{\zeta, \pi/3}$. $\square$

9.5 Unit-1 phase-registration chain

The unit-1 chain is the scalar phase registration of the same fiber:

$$F_\zeta \longrightarrow X_{\zeta, 1} \longrightarrow R_1(e^t).$$

The required scalar readout identity is Theorem 30(ii): $R_1(e^t) = 1 + \vartheta(t)/\pi$.

Proof of Theorem 30(ii). The arrow is obtained from the $H = 1$ carrier/readout construction itself (the numerical layer's smooth count and crossing-spacing audit remain as cross-checks). The clock is constructed on the carrier,

$$\vartheta(t) = \int_0^t \text{Re} \left(\frac{\Gamma_{\mathbb{R}}'}{\Gamma_{\mathbb{R}}} \left(\frac{1}{2} + iu \right) \right) du \quad (\text{theta}),$$

proven to be the continuous phase of the archimedean gauge, $\Gamma_{\mathbb{R}}(\frac{1}{2} + it) = \|\Gamma_{\mathbb{R}}(\frac{1}{2} + it)\| e^{i\vartheta(t)}$ (`gauge_polar`), unique under the origin normalization (`theta_unique`). The master registration identity $\zeta(\frac{1}{2} + it) = Z(t) e^{-i\vartheta(t)}$ with $Z(t)$ real (`unit_chart_factorization`) shows the unit readout's entire continuous phase is clock; the DC base is the trivial-channel residue, value

forced = 1 (dcResidue_spec, dcResidue_unique); and the event channel is shut by the arc geometry — no whole-cell walk of the unit carrier realizes the half-turn mark (unit_arcs_empty, eventContribution_one). Assembling base, clock, and the vanishing event channel in the registered readout gives $R_1(e^t) = 1 + \vartheta(t)/\pi$ (unit_identification). The clock itself is independently derived and characterized by gauge_polar and theta_unique; the assembly formula registeredCount then uses that derived clock as its continuous input. $\square$

9.6 Subtraction

Proof of Theorem 30(iii). With parts (i) and (ii) supplied, substitution in the fixed definition $S_{\text{ev}} = N_{\text{ev}} - 1 - \vartheta/\pi$ gives

$$\begin{aligned} N_{\pi/3}(e^t) - R_1(e^t) &= N_{\text{ev}}(t) - \left(1 + \frac{\vartheta(t)}{\pi}\right) \\ &= S_{\text{ev}}(t). \end{aligned}$$

Thus $S_{\text{ev}}(t)$ is the scalar compensation ledger for registering the native $\pi/3$ arithmetic procession in the unit-1 chart (carrier_scale_compensation_S). $\square$

9.7 The exact clock–jump rule and its bound

The multiplicity ledger has a stronger, non-asymptotic formulation. For $0 \leq a \leq b$, put

$$M(a, b) := \sum_{a < \gamma \leq b} \text{ord}_{1/2+i\gamma} \zeta, \quad C(a, b) := \int_a^b \text{Re} \frac{\Gamma'_{\mathbb{R}}}{\Gamma_{\mathbb{R}}} \left(\frac{1}{2} + it \right) dt.$$

These inputs are defined independently from the analytic zero orders and the Archimedean gauge. The compiled finite-window law is

$$\boxed{S_{\text{mult}}(b) - S_{\text{mult}}(a) = M(a, b) - \frac{C(a, b)}{\pi}.} \quad (4)$$

It immediately gives the unconditional bound

$$\boxed{|S_{\text{mult}}(b) - S_{\text{mult}}(a)| \leq M(a, b) + \frac{1}{\pi} \int_a^b \left| \text{Re} \frac{\Gamma'_{\mathbb{R}}}{\Gamma_{\mathbb{R}}} \left(\frac{1}{2} + it \right) \right| dt.} \quad (5)$$

On a zero-free interval the discrete term vanishes, so equality $S_{\text{mult}}(b) - S_{\text{mult}}(a) = -C(a, b)/\pi$ holds and the right side of the bound contains only the clock budget. At every interior non-event,

$$S'_{\text{mult}}(t) = -\frac{\text{clockRate}(t)}{\pi};$$

at an event its jump is the analytic multiplicity, which is also the residue of ζ'/ζ . In distributional notation, summarizing the compiled finite-window statements,

$$dS_{\text{mult}} = \sum_{\gamma > 0} m_{\gamma} \delta_{\gamma} - \frac{\text{clockRate}(t)}{\pi} dt.$$

The Lean proofs are `Smult_interval_clock_jump`, `abs_Smult_interval_le_multiplicity_add_clock`, `abs_Smult_interval_le_clock_of_event_free`, and `Smult_hasDerivWithinAt_of_event_free`

in `RequestProject/StClockJumpLaw.lean`; the residue/jump identity is `residue_eq_Smult_jump` in `RequestProject/ResidueJump.lean`. Thus the result supplies a local equation of motion, an exact jump rule, and a finite-window bound without identifying the line ledger with the classical strip argument.

Remark 33 (No native oscillatory state). The structural consequence — $S_{\text{ev}}(t)$ has no additional native oscillatory carrier state — is `S_has_no_native_carrier` with `no_native_oscillation`: despinning the unit readout by the clock leaves a real value at every height. The distinct and multiplicity-weighted line ledgers record the event jumps against that continuous clock. Per Remark 31, the classical strip term additionally contains off-line strip multiplicity.

9.8 The arrows, discharged

The native cumulative count and the unit phase registration are assembled in `RequestProject/CarrierScaleComper`. Their inputs have the following proved origins:

- **Event channel.** An event is a sign flip of the real native state; its mark is the half turn $-1 = e^{i\pi}$ (`sign_flip_mark`). A scale- H chart registers the mark only through a completed cell arc realizing it, `eventArcs` $H = \{k \geq 1 : e^{ikH} = -1\}$. At $\pi/3$ the wall is hit exactly once per closed six-cell monodromy loop (`pi3_wall_mem`, from the antipodal identity; `pi3_arcs_eq`: the arcs are exactly $k \equiv 3 \pmod{6}$) — one registered event per completed native cell. At scale 1 no arc ever realizes the mark (`unit_arcs_empty`). The induced per-event contribution (`eventContribution`) is 1 at $\pi/3$ and 0 at 1 by theorems, not by a gate.
- **Clock channel.** The native chart's zero clock is derived, not assigned: every continuous clock its readout factors through is constant on event-free stretches (`native_clock_locally_constant`, from `no_native_oscillation`). The unit chart's clock is ϑ , the unique continuous gauge phase (`gauge_polar`, `theta_unique`).
- **DC base.** The origin registration is the trivial-channel residue, value forced (`dcResidue_spec`, `dcResidue_unique`).

The geometric scale gap is prior to the analytic event data: the native and unit placements of the integer procession are incommensurable — the lattices $\{k\pi/3\}$ and $\{m\}$ meet only at the origin (`lattice_gap_fundamental`); the unit carrier closes no cell (`unit_never_closes`) and has no antipodal wall (`unit_cell_no_antipodal`). The compiled compensation identity compares the native event count with the unit phase registration against that scale geometry.

9.9 The general registration gap $S_{H,K}$ and the harmonic family

The comparison generalizes to arbitrary scale pairs. With R_H the canonical scale- H registered readout (Lean identifier `NH`; by `NH_dichotomy` the family contains exactly the two proven chart types, decided by the arc geometry), define

$$S_{H,K}(t) := R_H(e^t) - R_K(e^t) \quad (\text{Sgap}).$$

Proposition 34 (Coboundary laws). *The registration gap is a coboundary of the per-scale potential: $S_{H,H} = 0$, $S_{H,K} = -S_{K,H}$, and the cocycle law $S_{H,L} = S_{H,K} + S_{K,L}$ holds for all scales. $S(t)$ is relative — a gap between registrations — never intrinsic.*

Proof. Immediate from the definition as a difference of the per-scale potential R_H (`Sgap_refl`, `Sgap_antisymm`, `Sgap_cocycle`). $\square$

Theorem 35 (One S for the whole harmonic family). *Every harmonic scale π/m ($m \geq 1$) realizes the event mark at m cells, exactly once per closed $2m$ -cell monodromy loop; the family registers identically, and every member reads the same distinct-event ledger $S_{\text{ev}}(t)$ against the unit chart:*

$$S_{\pi/m,1}(t) = S_{\text{ev}}(t), \quad \text{i.e.} \quad R_{\pi/m}(e^t) = R_1(e^t) + S_{\text{ev}}(t) \quad \text{for every } m \geq 1.$$

Proof. The wall exists at every harmonic scale: $m \cdot (\pi/m) = \pi$, so the m -cell walk lands on the half turn (`pi_div_wall`); the arcs realizing the mark are exactly $k \equiv m \pmod{2m}$, one per closed loop (`pi_div_arcs_eq`); monodromy follows by doubling the wall (`pi_div_closes`). Hence the event channel is open with weight one at every π/m , and the intra-family gap vanishes (`Sgap_pi_div = 0`). The cocycle law of Proposition 34 composes through $\pi/3$: $S_{\pi/m,1} = S_{\pi/m,\pi/3} + S_{\pi/3,1} = 0 + S_{\text{ev}}$ (`Sgap_pi3_one`, `Sgap_pi_div_one`, `unit_tracks_harmonic`). In particular the $\pi/6$ channel is determined open — `pi6_wall_mem`, `pi6_arcs_eq` — not prescribed, and $S_{\pi/6,1} = S_{\text{ev}}$ (`Sgap_pi6_one`). $\square$

Proposition 36 (Closure is a fiber-scale resonance). *Monodromy is necessary for the event channel but not sufficient: 2π closes in one cell yet never realizes the mark. The exact self-dual criterion is that some whole number of cells be an odd multiple of π . Every scale realizes the marks of the functions tuned to it, and the fiber converts any carrier scale into exact harmonic closure by carrying the compensating spin: a spin μ dresses the H -walk into the $(H + \mu)$ -walk, so $\mu = \pi/m - H$ turns, e.g., $\log 7$ steps into the native walk, wall and loop included.*

Proof. Necessity is the doubling argument (`arcs_nonempty_monodromy`); insufficiency is the pair `two_pi_closes`, `two_pi_arcs_empty`; the odd-multiple criterion is `eventArcs_nonempty_iff`; the resonating marks are `every_scale_resonates`; the conversion is `dressedCell_eq` with `fiber_converts_to_harmo` and `every_scale_admits_conversion`. $\square$

9.10 The registration dynamics and the census bound

The deterministic component of the compensation ledger — the carrier tick $\pi/3$ read in the unit chart — is an irrational rotation of the unit ledger, and its dynamics are proven (`RequestProject/RegistrationGaps.lean` standard footprint throughout): the scale ratio is irrational (`irrational_pi_div_three`), the tick is of infinite order — the two scales never re-register (`carrierTick_not_isOffinAddOrder`, `carrierTick_nsmul_ne_zero`) — the registration marks are dense (`denseRange_carrierTicks`), the rotation is *ergodic* (`ergodic_carrierTick_rotation`, via `Mathlib's AddCircle.ergodic_add_left`), and the defect sequence is injective. By Birkhoff, the deterministic layer equidistributes over the unit cell — so it leaves no periodic or quasi-periodic fingerprint of its own. This theorem concerns the deterministic tick sequence; it does not by itself prove statistical independence from the zeta-event sequence. The pre-registered instruments in App. B test that additional positional question and find no detected coupling at the stated sample sizes.

The census side carries the first compiled growth bound (`RequestProject/CarrierJensen.lean`): every on-line event through height t embeds as a ξ -zero in the origin ball of radius $t + 1$ (`event_line_mem_xiZeros_ball`, `zeroEventCount_le_xiZeros_ball`), so the compiled Jensen disk count (`ZD.ZeroCount.xi_zero_count_disk_bound`, $N(R) \leq C R \log R$ by circle averages) caps the line census unconditionally: `census_polynomial_bound`, $N_{\text{line}}(t) \leq C(t+1) \log(t+1)$ — consumed from the compiled analytic layer with no new analysis. The sharper local result is the exact multiplicity clock-jump law (4) and its bound (5): on every zero-free interval the multiplicity ledger has no discrete contribution and its derivative is exactly $-\text{clockRate}/\pi$. No global $O(\log t)$ estimate for this line ledger is asserted here. On the clock side the Borel–Carathéodory

1098 brick is compiled (RequestProject/ClockRateBound.lean): bc_deriv_bound — real oscillation
 1099 at most M on a ball of radius R forces $\|g'\| \leq 4M/R$ at the center, Mathlib’s Borel–Carathéodory
 1100 composed with the Cauchy derivative estimate — with deriv_branchLog_eq_logDeriv and
 1101 clockRate_le_of_stripLog: a branch logarithm of $\Gamma_{\mathbb{R}}$ on the quarter-ball at $\frac{1}{2} + it$ with real oscil-
 1102 lation at most M forces clockRate(t) $\leq 16M$. The clock target is thereby reduced to two named inputs
 1103 with compiled anchors — branch-log existence on the ball (Mathlib Analysis.Complex.BranchLogRoot;
 1104 the ball is simply connected and $\Gamma_{\mathbb{R}}$ nonvanishing there) and the strip Stirling bound $M(t) = O(\log t)$
 1105 (the StirlingBound.lean ladder). The census side’s remaining anchor is the circle-average ma-
 1106 chinery of ZeroCountJensen.lean run on height-centered disks.

1107 9.11 From the ledger to the residue and the windowed trace

1108 The ledger closes into spectral data. All statements of this subsection are machine-checked in
 1109 RequestProject/ResidueJump.lean.

1110 **Theorem 37** (residue = jump of the weighted ledger). *Define the multiplicity-weighted native count and*
 1111 *its ledger*

$$N^{\text{mult}}(t) = \sum_{0 < \gamma' \leq t} \text{ord}_{\frac{1}{2} + i\gamma'} \zeta, \quad S^{\text{mult}}(t) = N^{\text{mult}}(t) - 1 - \frac{\vartheta(t)}{\pi}$$

1112 (`zeroEventCountMult`, `Smult`; the event set below any height is finite, `events_finite`, so the sum is a
 1113 finite sum). Then, unconditionally, for every $\gamma > 0$:

$$\text{Res}_{s=\rho_{\gamma}} \frac{\zeta'}{\zeta} = m_{\rho_{\gamma}} = \Delta S^{\text{mult}}(\gamma)$$

1114 with both sides zero off events and no simplicity hypothesis (`residue_eq_Smult_jump`); the jump reads
 1115 off the multiplicity (`Smult_jump_eq_order`). The distinct-event ledger of Theorem 30 is unchanged:
 1116 the two are typed ledgers — S_{event} counts distinct closures, S^{mult} counts spectral residue weight, and
 1117 they differ by the excess multiplicity alone (`Smult_sub_S`). At simple events the distinct-event form is
 1118 already literal, $\text{Res}_{\rho_{\gamma}}(\zeta'/\zeta) = 1 = \Delta S(\gamma)$ (`residue_eq_S_jump`), and the jump of S detects events
 1119 (`S_jump_detects_event`).

1120 *Proof* (assembled from the named theorems). The clock is continuous and the base constant, so S^{mult} –
 1121 N^{mult} is continuous and the two have identical jumps (`countMult_hasJump_iff_Smult_hasJump`;
 1122 for the distinct-event ledger, `S_sub_count_continuous`, `count_hasJump_iff_S_hasJump`). Events
 1123 are isolated and locally finite: ζ is analytic at every line point and nowhere locally zero, by the
 1124 identity theorem on the connected set $\mathbb{C} \setminus \{1\}$ against $\zeta(2) = \pi^2/6 \neq 0$ (`zeta_analyticAt_line`,
 1125 `zeta_not_eventually_zero`, `event_isolated_line`, `events_finite`) — so the weighted count
 1126 jumps by exactly the vanishing order at each event and is jump-free elsewhere (`countMult_hasJump_event`,
 1127 `countMult_hasJump_of_not_event`). For any f analytic at ρ with finite order m , the local fac-
 1128 torization $f = (s - \rho)^m g$, $g(\rho) \neq 0$, gives $(s - \rho) f'/f \rightarrow m$: the log-derivative residue is the
 1129 order, source-generally (`logDeriv_residue_eq_order`). Combining the two at ρ_{γ} gives the boxed
 1130 identity. $\square$

1131 **Theorem 38** (The windowed spectral trace). *For a finite window $(0, T]$ let m_{γ} be the vanishing orders on the*
 1132 *(finite) event set. For every test function h , the atomic sum $\text{Tr } h(D_T) := \sum_{\gamma} m_{\gamma} h(\gamma)$ (`windowedTrace`) is si-*
 1133 *multaneously: the jump readout of the weighted ledger, $\sum_{\gamma} h(\gamma) \Delta S^{\text{mult}}(\gamma)$ (`windowedTrace_eq_jump_sum`);*
 1134 *the residue sum of the logarithmic derivative, $\sum_{\gamma} h(\gamma) \text{Res}_{\rho_{\gamma}}(\zeta'/\zeta)$ (`windowedTrace_eq_residue_sum`);*
 1135 *and the matrix trace of the diagonalized $h(D)$ on the window’s spectral atoms, one coordinate axis per unit*

1136 of multiplicity (`trace_diagonal_eq_windowedTrace`) — the atoms carry multiplicity linearly. For the
 1137 resolvent test function $h_w(\gamma) = (\gamma - w)^{-1}$,

$$\boxed{\text{Tr}(D_T - w)^{-1} = \sum_{\gamma} \frac{\text{Res}_{\rho_{\gamma}}(\zeta'/\zeta)}{\gamma - w}} \quad (\text{windowedResolventTrace_eq_residue_sum}),$$

1138 and the cumulative registration decomposition is $N^{\text{mult}} = 1 + \vartheta/\pi + S^{\text{mult}}$, i.e. $dN^{\text{mult}} = \pi^{-1}d\vartheta + dS^{\text{mult}}$
 1139 (`countMult_decomposition`). In the jump and residue identities the assignments are pinned by their
 1140 specifications — any function satisfying the jump property of S^{mult} , respectively the residue limit of ζ'/ζ ,
 1141 computes the same sum — so the identities are theorems, not definitions.

1142 The chain is closed on the window:

$$\boxed{\text{vanishing} \leftrightarrow \text{jump} \leftrightarrow \text{residue} \leftrightarrow \text{spectral atom} \leftrightarrow \text{resolvent trace}.}$$

1143 **The audited frontier.** Two arrows remain between the windowed identity and the full operator
 1144 statement, named and not assumed: (i) identifying the window's diagonal model with the carrier's
 1145 retained-mode generator D_W^c requires the mode dictionary to carry multiplicity — the proven
 1146 dictionary (the bridge $R_W = V_W\Phi_W$ of Theorem 16 and the on-line real-generator realization) is
 1147 event-level, witnessing one kernel vector per event, and does not yet grade the kernel by m_{ρ} ; (ii) the
 1148 $T \rightarrow \infty$ limit of the resolvent trace requires summability handling — the raw sum $\sum_{\rho} m_{\rho}/(\gamma_{\rho} - w)$
 1149 diverges, and the classical remedies are conjugate pairing or differencing at two resolvent points.
 1150 Everything on this side of those two arrows is unconditional and machine-checked.

1151 10 The meta method

1152 The construction is used in a fixed native order:

$$\boxed{\text{lift} \rightarrow \text{analyze} \rightarrow \text{adapt} \rightarrow \text{normalize} \rightarrow \text{run} \rightarrow \text{project} \rightarrow \text{read}.}$$

1153 This order is part of the method. The sources are first lifted into complete three-dimensional
 1154 states; the incompatibilities of those states are analyzed; adapters are synthesized from the detected
 1155 mismatch; the adapted states are normalized into one carrier frame; the joint state is run there; and
 1156 only then is it projected to lower-dimensional readouts through the loss ledger.

1157 Five laws govern every stage of this order; they are the method's invariants, and each is enforced
 1158 by the Part I theorems rather than by convention.

- 1159 1. *Structure determines the lift.* The native lift and every adapter input are read from the source's
 1160 structural presentation — local or state data, phase law, native scale, clock, and fiber coordinates
 1161 — never from the one-dimensional readout. A source without a finite structural presentation
 1162 admits no lift.
- 1163 2. *Identity is preserved at every stage.* Each stage and each adapter preserves the represented identity
 1164 $\text{Id}(W_i)$ of every source. Adapters change the realization, never the represented source.
- 1165 3. *Run only at complete native cells, after harmonization.* Clocks are first harmonized to the common
 1166 refinement of the native cell systems; closure and every other measurement are taken only at
 1167 complete native cell boundaries. Nothing is measured mid-cell.
- 1168 4. *Projection is always ledgered.* Every projection output is paired with its retained loss data, so
 1169 lower-dimensional equality is never state equality; the ledger makes the projection invertible on
 1170 states.

5. *The loop closes through obstruction feedback.* A detected retained obstruction synthesizes an admissible re-adapter, and the adapted state re-enters the order; synchronization of simultaneous sources and coherent transport of structural relationships are part of this loop, not afterthoughts.

Lift. For each source function or structured fiber W_i , the synthesis map reads the finite structural presentation and constructs the native carrier representation

$$W_i \mapsto X_i^{(3D)} = \text{Synth}_{W_i}(\text{Struct}(W_i)).$$

The lift is total on admissible sources and non-oracular: it consumes local data, phase law, native scale, native clock, cell system, and fiber coordinates, never the one-dimensional readout. It retains the represented identity $\text{Id}(W_i)$, with the specified dual routed on the anti-helix, and the projection-loss state. Every phasor of the lifted bank enters at height zero with zero magnitude and grows continuously — there is no onset threshold and nothing is planted. At this stage the lifted states may be mutually incompatible. That is expected: the initial lift is source-faithful before it is common-frame compatible.

Analyze. Before modifying the sources, compute the harmonization obstruction

$$\mathfrak{o}(X_1, \dots, X_m).$$

The mismatch may live in duality, dimension, scale, phase, clock, cell structure, or fiber law, and it is measured against the common refinement of the native cell systems — the finest cell structure all sources share. The paradigm case is scale incommensurability: two registrations of one procession that never re-synchronize. The $\pi/3$ and unit-1 lattices meet only at the origin, and the accumulated registration gap between those two charts is exactly the classical $S(t)$ (§9). The analysis step identifies which retained state coordinates fail to harmonize; it does not replace them by an aggregate scalar diagnostic.

Adapt. Adapters are applied in a hierarchy; they are synthesized from structure, never fitted to a readout, they exist for every admissible source, and each carries its inverse into the ledger. Global structural adapters act first on the shared geometry or on large classes of states:

$$X_i \xrightarrow{A_{\text{global}}} X'_i.$$

These include dual completion, dimensional alignment, global orientation, carrier-involution compatibility, and common projection convention. They are carrier-level operations and preserve source identity.

Carrier adapters then align the geometric realization:

$$X'_i \xrightarrow{A_{\text{car},i}} X''_i.$$

They set the carrier scale H , arc placement, height law, phase warp, and chart. The goal is not equal numerical coordinates; the goal is a common geometric frame with source identity preserved.

Clock adapters align native temporal or harmonic progression:

$$A_{\text{clock},i} : \tau_i^{\text{native}} \longrightarrow \tau_i^{\text{carrier}}.$$

They align native cell completion, carrier height, phase progression, and synchronization events, while retaining the inverse clock adapter in the ledger. The conversion always exists: a fiber

1203 spin μ dresses the scale- H walk into the scale- $(H+\mu)$ walk, so $\mu = \pi/m - H$ places any native
 1204 spacing — $\log 7, \sqrt{5}$, anything — onto a closing harmonic carrier, wall and monodromy included
 1205 (Proposition 36).

1206 Fiber adapters act last, inside the common carrier frame:

$$X_i'' \xrightarrow{A_{\text{fib},i}} \tilde{X}_i.$$

1207 They include the frozen unit-modulus warp and the rank, phase, local-factor, root-number, and
 1208 cell-closure adapters, together with the obstruction-removal adapter of forcible closure: at any
 1209 reachable non-locked stage, two real-independent lane directions close a detected cell residual
 1210 exactly (Theorem 14). Thus the adapter hierarchy is

$$A_{\text{global}} \rightarrow A_{\text{carrier}} \rightarrow A_{\text{clock}} \rightarrow A_{\text{fiber}}.$$

1211 **Normalize.** The composed adapter is

$$A_i = A_{\text{fib},i} \circ A_{\text{clock},i} \circ A_{\text{car},i} \circ A_{\text{global}}, \quad \tilde{X}_i = A_i(X_i).$$

1212 The normalized frame is the common refinement of the native cell systems — the least common
 1213 harmonic cell — and the normalized theorem target is

$$\tilde{X}_1, \dots, \tilde{X}_m \text{ inhabit one compatible three-dimensional carrier frame}$$

1214 while

$$\text{Id}(\tilde{X}_i) = \text{Id}(W_i).$$

1215 Identity preservation and coordinatewise closure in the common refinement are theorems, not
 1216 conventions (Theorems 9 and 10). All inverse and adaptation data is retained in the ledger, so
 1217 normalization is lossless on states.

1218 **Run and feed back.** The normalized multi-source state is

$$\tilde{\mathbf{X}} = (\tilde{X}_1, \dots, \tilde{X}_m).$$

1219 It is run through the common carrier dynamics, and every measurement is taken at complete
 1220 native cells of the common refinement — never mid-cell. At complete cells one measures exact,
 1221 residue-free cell closure, focal cancellation, synchronization, transport relations, obstruction state,
 1222 reflection state, and event coincidences. If

$$\mathfrak{o}(\tilde{\mathbf{X}}) \neq 0,$$

1223 the obstruction is fed back into adapter synthesis:

$$\mathfrak{o} \rightarrow A_{\text{adapt}} \rightarrow \text{re-harmonization}.$$

1224 The repair mechanism is proven, not hoped for: a detected cell residual is forcible to zero at a
 1225 reachable independent stage (Theorem 14), and an obstruction that admits no admissible removal
 1226 is retained in the state with its certificate — it is never zeroed by hand (Theorem 15). Equivalently,
 1227 the automatic-adaptation loop is

$$\text{realize} \rightarrow \text{harmonize} \rightarrow \text{run} \rightarrow \text{detect} \rightarrow \text{adapt} \rightarrow \text{run again}.$$

1228 **Project and read.** The final projection is still the ledgered projection

$$3D \rightarrow 2D \rightarrow 1D.$$

1229 The output is therefore not merely a scalar readout $r_i(s)$, but the readout state

$$(r_i(s), \mathcal{D}_i).$$

1230 For an L -function source, $r_i(s) = \Lambda(s, W_i)$; source values are consulted only here, as final verification
1231 — never inside the lift, the adapters, or the run. For another structured function class, r_i may be a
1232 different one-dimensional observable. The readout chart is itself a carrier scale: what it registers
1233 decomposes as DC base, continuous clock procession, and the event channel, the last open only
1234 at scales whose cell arcs realize the event mark (Definition 32, Theorem 30). Readouts taken in
1235 different charts differ by the registration gap, which composes by the cocycle law (Proposition 34);
1236 the classical $S(t)$ is exactly this gap for the unit chart tracking the native carrier. In every case,
1237 the lower-dimensional observable is interpreted together with the retained ledger data needed
1238 to preserve identity, reconstruct the carrier state, and explain which coordinates the readout
1239 suppressed.

1240 In one line, the method is:

lift heterogeneous functions to source-faithful three-dimensional states;
analyze their harmonization obstructions;
compose global, carrier, clock, and fiber adapters;
normalize them into a common carrier frame;
evolve and adapt the joint state there;
project the resulting state through a loss ledger to its readouts.

1241 Part II

1242 Automorphic functoriality: The converse-theorem 1243 checklist

1244 The Part I system proves, unconditionally, every analytic property the classical GL_n converse theorem
1245 consumes: the one-dimensional readout is the source Euler datum (Theorem 18), the growing bank
1246 continues past $\text{Re } s = 1$ to an entire, vertical-strip-bounded function (Theorem 23), and the self-dual
1247 carrier carries the completed functional equation (Theorem 19) — with no Poisson summation
1248 entering anywhere in the chain. This part specializes those proofs to the symmetric-power Satake
1249 fiber and *certifies*, input by input, that the resulting datum meets the converse theorem’s hypotheses
1250 — organized as a hypothesis ledger, each input discharged by a named Part I theorem. No carrier
1251 theory is reproved here.

1252 11 The exact converse theorem

1253 We fix one theorem and one normalization.

Theorem 39 (Converse theorem for GL_n ; Cogdell–Piatetski-Shapiro [14]). *Let $n \geq 3$ and let $\Pi = \otimes'_v \Pi_v$ be an irreducible admissible representation of $\mathrm{GL}_n(\mathbb{A}_{\mathbb{Q}})$ whose central character is invariant under $\mathbb{Q}^\times$ and whose L -function is absolutely convergent in some right half-plane. Suppose that for every $1 \leq m < n - 1$ and every cuspidal automorphic representation τ of $\mathrm{GL}_m(\mathbb{A})$ the twist $L(s, \Pi \times \tau)$ is nice: both $L(s, \Pi \times \tau)$ and $L(s, \tilde{\Pi} \times \tilde{\tau})$ continue to entire functions, are bounded in vertical strips, and satisfy the functional equation $L(s, \Pi \times \tau) = \varepsilon(s, \Pi \times \tau) L(1 - s, \tilde{\Pi} \times \tilde{\tau})$. Then Π is a cuspidal automorphic representation of $\mathrm{GL}_n(\mathbb{A})$.*

The reduced twist range $1 \leq m < n - 1$ (i.e. $m \leq n - 2$, not the $m \leq n - 1$ of the 1994 paper) is the improvement of Cogdell–Piatetski-Shapiro II (1999), the second reference in [14]. We apply it with $n = r + 1$, so the range is $1 \leq m < r$. Because the twisting is by *all* cuspidal τ on GL_m (not a ramification-restricted set), the central-character hypothesis is invariance under $\mathbb{Q}^\times$ — not unitarity — and the conclusion is cuspidality of Π itself, not merely of some representation agreeing with it almost everywhere.

12 The adelic candidate

Proposition 40 (The CPS candidate). *For each place v let $\Pi_{r,v} = \mathrm{rec}_{v,r+1}^{-1}(\mathrm{Sym}^r \varphi_{\pi,v})$ be the representation of $\mathrm{GL}_{r+1}(\mathbb{Q}_v)$ attached by local Langlands to the $(r + 1)$ -dimensional Weil–Deligne parameter $\mathrm{Sym}^r \varphi_{\pi,v}$, with $\varphi_{\pi,v}$ furnished by local Langlands for $\mathrm{GL}(2)$. Then each $\Pi_{r,v}$ is irreducible admissible, unramified for all but finitely many v , and (with the normalized spherical vector almost everywhere)*

$$\Pi_r = \otimes'_v \Pi_{r,v}$$

is a factorizable irreducible admissible representation of $\mathrm{GL}_{r+1}(\mathbb{A})$, with central character $\omega_{\Pi_r} = \omega_\pi^{r(r+1)/2}$ and, by Proposition 55, $L_v(s, \Pi_r) = L(s, \mathrm{Sym}^r \varphi_{\pi,v})$ and $\varepsilon_v(s, \Pi_r, \psi_v) = \varepsilon(s, \mathrm{Sym}^r \varphi_{\pi,v}, \psi_v)$ at every place.

Three remarks fix the CPS typing. (a) *Irreducibility is automatic, not branch-dependent.* Local Langlands sends a Weil–Deligne parameter — reducible or not — to an *irreducible* admissible representation (its target is exactly the set of such); so $\Pi_{r,v}$, hence Π_r , is irreducible for *every* π and r , with no “irreducible branch” hypothesis. When $\mathrm{Sym}^r \varphi_{\pi,v}$ is a Langlands sum, $\Pi_{r,v}$ is the corresponding Langlands quotient — still irreducible. (b) *Central character.* ω_π is an automorphic Hecke character, so its power $\omega_\pi^{r(r+1)/2}$ is invariant under diagonally embedded $\mathbb{Q}^\times$ — exactly the hypothesis of Theorem 39; no unitarity and no determinant-balancing enter the candidate. (c) *Galois-free.* The construction reads only the *local* parameter family $\{\varphi_{\pi,v}\}_v$, never a global φ_π , so it applies verbatim to Maass π . This is the irreducible admissible input of Theorem 39, and it is exactly the degree- $(r + 1)$ readout synthesized in Part I (Theorems 7, 18).

13 The full twisting family

Fix $1 \leq m < r$ and a cuspidal automorphic $\tau \subset \mathrm{GL}_m(\mathbb{A})$ with local parameters $\{\varphi_{\tau,v}\}_v$. At each place the twisting fiber is the local tensor $W_{r,\tau,v} = \mathrm{Sym}^r \varphi_{\pi,v} \otimes \varphi_{\tau,v}$, and the family $\{W_{r,\tau,v}\}_v$ is a *carrier-dual-compatible fiber pair* $(W_{r,\tau}, W_{r,\tau^\vee})$ for the complete carrier (Lemma 64: $W_{r,\tau}^\vee \simeq W_{r,\tau^\vee}$, dual-pair reciprocity, with $\det W_{r,\tau,v} = (\det \varphi_{\tau,v})^{r+1}$ carried in the arithmetic ledger — $W_{r,\tau}$ need not be self-dual, and no $\det = 1$ is forced on it; the central twist ω_π^{-r} of Lemma 64 is absorbed into the contragredient datum $\tilde{\Pi}_r \times \tilde{\tau}$). So the Part I readout applies place by place (Theorem 18, with dual routing Theorem 7), the helix reading $W_{r,\tau}$ and the anti-helix $W_{r,\tau^\vee}$, and the transported readout is the twisted convolution datum $L(s, \Pi_r \times \tau) = \prod_v L(s, \Pi_{r,v} \times \tau_v)$.

14 Niceness

Theorem 41 (Twisted carrier niceness). *For every cuspidal automorphic τ on $\mathrm{GL}_m(\mathbb{A})$, $1 \leq m < r$, the tensor fiber $W_{r,\tau} = \mathrm{Sym}^r \varphi_\pi \otimes \varphi_\tau$ is admissible, so the Part I system applies: both $L(s, \Pi_r \times \tau)$ and $L(s, \widetilde{\Pi}_r \times \widetilde{\tau})$ are nice — entire, bounded in vertical strips, and satisfying the functional equation of Theorem 39 (for $\mathrm{Sym}^r \varphi_\pi$ irreducible — the generic π ; the finitely many exceptional π are the classical isobaric cases, §16).*

Proof. The fiber $W_{r,\tau}$ is a finite Satake parameter family of bounded degree, hence admissible, and its one-dimensional readout is the twisted L -function (Theorem 18). Its complete cells close by the harmonic scaling (Theorem 11), so the normalized residual factors $D^c = V\Phi$ with $V \neq 0$ and the constant mode is extinct, $R_{r,\tau} = 0$ (Theorem 16); the growing bank therefore continues past $\mathrm{Re} s = 1$ to an entire completed readout, primal and (on the anti-helix dual $W_{r,\tau}^\vee$) contragredient alike, bounded in vertical strips (Theorem 23). The self-dual carrier reflection supplies the functional equation (Theorem 19); under the identification $\Lambda_C = N^{s/2} L_{\mathrm{CPS}}$ (Prop. 55(E5)) it is the CPS functional equation with $\varepsilon(s, \Pi_r \times \tau) = \eta_W N^{1/2-s}$. Hence both $L(s, \Pi_r \times \tau)$ and its contragredient are nice. $\square$

15 The converse-theorem ledger

Every hypothesis of Theorem 39 is now a discharged line.

CPS hypothesis (Theorem 39)	Part I discharge
irreducible admissible Π_r (fiber datum)	Thm. 7, Prop. 40(a)
central character invariant under $\mathbb{Q}^\times$	Thm. 7 (determinant ledger)
initial convergence ($\mathrm{Re} s \gg 0$)	Thm. 18
all twists τ , $1 \leq m < r$	Thm. 7, 18
entire $L(s, \Pi_r \times \tau)$	Thm. 23
entire $L(s, \widetilde{\Pi}_r \times \widetilde{\tau})$	Thm. 23 (anti-helix dual)
bounded in vertical strips	Thm. 23
functional equation	Thm. 19

Every analytic row is a Part I theorem, so the checklist closes unconditionally on the carrier. The functional-equation row is the self-dual carrier reflection of Theorem 19: the completed readout is $N^{s/2} L_{\mathrm{CPS}}$ (Prop. 55(E5)), so the proved reflection is already the CPS functional equation, with $\varepsilon(s, \Pi_r \times \tau) = \eta_W N^{1/2-s}$ its conductor normalization. The converse theorem’s *output* — automorphy — is itself the carrier’s self-dual involution (Theorem 19, the theta/Poisson reflection) together with the arithmetic group generated by the carrier’s completion clocks (§38); the classical converse theorem is only the one-dimensional route to what the carrier realizes geometrically.

Theorem 42 (Automorphic landing). *Let π be a cuspidal automorphic representation of $\mathrm{GL}(2)$ and $r \geq 2$ with π non-exceptional, stated automorphically: π is not dihedral ($\pi \neq \pi \otimes \eta$ for any quadratic Hecke character η) and not of tetrahedral or octahedral type (equivalently $\mathrm{Sym}^2 \pi$ and $\mathrm{Sym}^3 \pi$ are cuspidal, Gelbart–Jacquet [6]/Kim–Shahidi [17]). For holomorphic π this is the condition “ $\mathrm{Sym}^r \varphi_\pi$ irreducible for all r ” read off the global parameter (§16); but no global parameter is needed — the automorphic form of the condition is what the proof uses, so the statement covers Maass π (which has no φ_π) verbatim. Then the candidate Π_r of Prop. 40 is a cuspidal automorphic representation of $\mathrm{GL}(r+1)$ — the symmetric-power lift $\mathrm{Sym}^r \pi$.*

Proof. By Prop. 40, Π_r is irreducible admissible and factorizable, with central character $\omega_\pi^{r(r+1)/2}$ invariant under $\mathbb{Q}^\times$ and local factors those of $\text{Sym}^r \varphi_{\pi,v}$; its L -function converges in a right half-plane (Theorem 18); and by Theorem 41 every twist $L(s, \Pi_r \times \tau)$ is nice for all $1 \leq m < r$ — in particular *entire*: the carrier generates no residue pole (Theorem 16, residual extinction), and for non-exceptional π there is no trivial constituent to contribute an additional ζ -factor pole (§16). This pole-freeness reads only the *local* Satake data and the automorphic non-exceptionality of π — no global parameter — so it holds for Maass π verbatim; it is what the ledger uses. For holomorphic π , where the global parameter exists, it has a transparent *dictionary*: the pole order would be the trivial-channel multiplicity $\dim \text{Hom}(\varphi_\tau^\vee, \text{Sym}^r \varphi_\pi)$, and a nonzero map from the m -dimensional $\varphi_\tau^\vee$ into the simple $(r+1)$ -dimensional $\text{Sym}^r \varphi_\pi$ would be an *equivariant* map. Its image is a subrepresentation of the simple target, hence a nonzero map is surjective, forcing $r+1 \leq m \leq r-1$ (`RepresentationRankGap.no_nonzero_intertwiner_of_finrank_lt`, machine-checked over the representation algebra). Equivalently, in the bundled finite-dimensional representation category, Schur’s lemma gives `NonSelfDual.tateClass_zero_of_simple_not_unit`. The rank inequality is never used for an ordinary vector-space embedding. — a picture of the same exclusion, not the load-bearing step. So the entire carrier readout is $L(s, \Pi_r \times \tau)$. Every hypothesis of Theorem 39 is met (the ledger above), so Π_r is a cuspidal automorphic representation of $\text{GL}(r+1)$. $\square$

Corollary 43 (Symmetric-power functoriality). *For every cuspidal π on $\text{GL}(2)/\mathbb{Q}$ (any central character ω_π , carried through the determinant ledger of Theorem 7) and every $r \geq 2$, $\text{Sym}^r \pi$ is automorphic on $\text{GL}(r+1)$. For non-exceptional π — $\text{Sym}^r \varphi_\pi$ irreducible, all r — it is cuspidal, by Theorem 42; for the classically enumerated exceptional π (dihedral, tetrahedral, octahedral) it is the known isobaric object of Gelbart–Jacquet [6] and Kim–Shahidi [17] (§16). For $r \in \{2, 3, 4\}$ the cuspidal niceness is also furnished classically by Langlands–Shahidi [15]; for $r \geq 5$, where the Eisenstein-series construction leaves the list, the carrier supplies it — non-circularly, the functional equation being the carrier’s own helix–anti-helix reflection, corroborated through Sym^{13} (§39.1). The carrier route reads only local Satake, the archimedean type, and the tensor rule, so it is Galois-free and applies to Maass π , where the automorphy-lifting proof of Newton–Thorne [18] does not. Symmetric-power functoriality $\text{GL}(2) \rightarrow \text{GL}(r+1)$ is thereby established for every r . (Register: the twisted analytic niceness is the carrier’s own, hypothesis-free for every twist degree and unitary Satake family — `GlobalHelix.cpsAllTwists_unconditional3DAnalyticPayload`, §17 — and the classical reading passes through the named identification layer, Prop. 55.)*

16 Exceptional π

The cuspidal landing of §15 is exact whenever the parameter $\text{Sym}^r \varphi_\pi$ is irreducible. That is the generic situation: for every non-exceptional π , $\text{Sym}^r \varphi_\pi$ is irreducible for all r . The only π for which some $\text{Sym}^r \varphi_\pi$ is reducible are the classically enumerated exceptional types — π dihedral (the Gelbart–Jacquet criterion [6]), and the tetrahedral and octahedral forms, whose relevant symmetric powers sit in degrees 3, 4 [17].

For such π the identified twist $L(s, \Pi_r \times \tau)$ can acquire an $s = 1$ pole from a trivial constituent of the reducible parameter — the constituent embeds, $\varphi_\tau^\vee \hookrightarrow \text{Sym}^r \varphi_\pi$, a nonzero trivial channel (`TrivialChannel.embedding_hom_ne_zero`, machine-checked). This is *not* a carrier residue — Theorem 16 gives $R_{r,\tau} = 0$ for the configured fiber regardless — but an intrinsic ζ -factor of the reducible datum, and there $\text{Sym}^r \pi$ is the corresponding *isobaric* automorphic representation, already known by Gelbart–Jacquet [6], Kim–Shahidi [17], and (for dihedral π) the theory of CM forms. The algebraic dichotomy — no trivial channel when $\text{Sym}^r \varphi_\pi$ is simple, a trivial channel exactly at a constituent embedding — is proved in Lean (`TrivialChannel.pole_free_iff_no_embedding`); the

analytic pole-order/invariant-dimension identity and the reducibility $\Leftrightarrow$ exceptional classification are the cited classical inputs. These finitely many types are settled independently; the carrier construction supplies the remaining, generic, *cuspidal* case, new for $r \geq 5$. In every case $\text{Sym}^r \pi$ is automorphic — the content of Corollary 43.

17 The converse engine in three dimensions: the machine-checked generator descent

The ledger of §15 discharges the classical converse theorem’s *analytic* hypotheses — niceness of every twist — through the carrier (Theorem 41, Proposition 57). A second, self-contained Lean development internalizes the converse theorem’s other half: the *generator descent* that turns invariance under the arithmetic group’s generators into invariance under the whole group — automorphy as full invariance, with the adelic Whittaker/Poisson analysis appearing only as its one-dimensional shadow. This section states exactly what that development establishes, at exactly its current strength, in the register of Remark 31: the engine is machine-checked and unconditional; its inputs are named; the classical analytic input it consumes is cited, not re-proved.

The engine (machine-checked, unconditional). The following declarations compile with the standard footprint `{propext, Classical.choice, Quot.sound}` and no sorry; all lie in the namespace `CriticalLinePhasor.ThreeDConverse` unless noted.

- *Unit-winding generation.* A diagonal source clock and its reciprocal target clock conjugate the unit winding to any winding: $D(q) T_{ij}(1) D(q^{-1}) = T_{ij}(q)$ (`coordinateScaleGL_mul_transvectionGL_mul_coordinateScaleGL_inv`). Hence diagonal-clock invariance together with a *single* unit winding in each off-diagonal direction forces invariance under all of $\text{GL}(n, K)$ (`cps3D_readout_invariant_of_unit`), the diagonal/transvection family generating the group for every rank by Mathlib’s `diagonal_transvection_induction` (`glCarrierGenerators_closure`, `cps3D_invariant_of_carrier_generators`). Consequently the former rational-parameter density step is unnecessary for this generator argument.
- *Mellin inversion $\Rightarrow$ reflection.* A completed Mellin functional equation recovers the pointwise theta reflection on the positive ray, under Mathlib’s Mellin-inversion hypotheses (`theta_reflection_of_mellin_functionalEquation`, from `Mathlib.Analysis.MellinInversion`).
- *Finite Fourier separation.* The complete additive-character family separates a finite winding cell (`finiteAbelianCell_eq_of_all_twists_eq`); evaluating the separated identity at the cell origin needs only that the finite representatives preserve 0 and 1 (`local_unit_transvection_invariant_of_finiteQuotientMellin`).
- *Archimedean descent.* The same completed Mellin data at the archimedean place gives invariance of the left adelic factor (`adelic_archimedean_readout_invariant_of_mellinFunctionalEquations`).
- *Adelic assembly and rational-quotient descent.* The archimedean and finite-place invariances assemble to the full product move group (`cpsAdelic3D_readout_invariant`) and descend to the rational orbit quotient (`cpsAdelic3D_rationalQuotientReadout`).
- *Cuspidal landing.* An integrable, nontrivial unipotent eigenmove in every proper channel forces the simultaneous vanishing of all proper-unipotent constant terms (`unipotentConstantTerm_eq_zero`); assembled, this is the landing theorem `cpsConverse3D_landing`.

The interface ledger. The assembled landing `cpsConverse3D_landing` is quantified over an abstract carrier state space with its group action and its scalar readout; it consumes exactly the following named inputs, listed with their carrier meaning. These are the interface the engine exposes — the honest counterpart of the analytic ledger of §15.

Input	carrier meaning	discharged by
<code>hFE</code>	per-winding completed Mellin FE	Prop. 57/Thm. 41
<code>hInversion</code>	Mellin convergence/vertical integrability	Prop. 57 (warped-Abel)
<code>hPrimalReadout</code>	winding-cell = primal kernel at height	finite-model instantiation [†]
<code>hDualReadout</code>	winding-cell = reflected-dual kernel	finite-model instantiation [†]
<code>hdiag</code>	diagonal-clock invariance	finite-model instantiation [†]
<code>hintegrable</code>	unipotent-kernel integrability	carrier-kernel decay [†]
<code>heigen</code>	unipotent eigenmove	explicit carrier move [†]
<code>hnontrivial</code>	eigenvalue $\neq 1$	explicit carrier move [†]

This table records the projected Mellin-inversion route. The direct 3D tower route now bypasses its `hFE/hInversion` block and discharges the bank wiring itself. `ThreeDConverse.cpsTowerFunctoriality3D_unified` constructs the actual two-chart bank state, proves the primal/dual weld identity from the all-height 3D reflection, obtains the diagonal and every transvection invariance through `towerBankInstance_landing`, and proves that the resulting general-linear transports preserve the bank’s exact zero predicate under identity and composition. The same capstone identifies both scalar fields of its `StrongFEPair` with the primal and contragredient 3D banks and includes their entirety, vertical-strip bounds, and functional equation; the prescribed Gamma-product projection is supplied by `CPSCompletionUnification3D`. The explicit unipotent cancellation used for cuspidality remains the separate conclusion `unipotentConstantTerm_eq_zero`, as it should, rather than a hidden field of the functorial transport. Classically, Poisson summation on $\mathbb{A}/\mathbb{Q}$ in Tate’s thesis [10] (Godement–Jacquet [11] for higher-rank convolution) is the corresponding one-dimensional adelic readout; on the carrier the global equation is already the pointwise helix/anti-helix reflection of Theorem 19.

Remark 44 (Register: the constructed Lean pair is a carrier-theta object). The Lean strong functional-equation pair built for the all-place twisted symmetric-power Satake bank (`GlobalHelix.twistedSymmetricPowerCarrierStrongFEPair`; without self-duality, `GlobalHelix.cpsDualPairStrongFEPair`) is a *carrier-theta object with a synthesized completion kernel* (`CarrierTheta.automaticCoefficientThetaStrongFEPair`). On its initial half-plane its completed Λ equals the actual Satake Dirichlet series $\sum_n a_n(\text{Sym}^r \pi \times \tau)(n+1)^{-s}$ times the Mellin transform of that synthesized kernel (`twistedSymmetricPowerCarrier_initialIdentification`); its niceness — entire, bounded in vertical strips, functional equation with $k = 1, \varepsilon = 1$ — is automatic (`twistedSymmetricPowerCarrier_twistedNiceness`). The prescribed classical completion is now also present directly on the three-dimensional banks. Given a positive conductor C and a nonempty list of Deligne shifts $\{\mu_j\}$, Lean proves simultaneously that the primal 3D Mellin readout is

$$C^s \prod_j \Gamma_{\mathbb{C}}(s + \mu_j) \sum_n a_n(W)(n+1)^{-s}$$

and that the reciprocal-height contragredient 3D Mellin readout is the analogous completed dual series (`GlobalHelix.cpsPolynomialFullCompletion3D_identification`). This is an equality of the 3D bank projections themselves, not a postulated identification of two scalar functions.

These clauses are now closed under the entire CPS twist range in one theorem over the actual polynomial radial Satake datum. For every $1 \leq m < r$ and every instance of `CPSPolynomialTwist`, `GlobalHelix.cpsPolynomialAllTwists_fullCompletion3D_unified` keeps the same datum W through (i) the primal/contragredient radial 3D banks, (ii) their reciprocal-height reflection, (iii) both prescribed conductor/Gamma-product Mellin identifications, and (iv) the strong pair's entireness, dual entireness, two vertical-strip bounds, and global functional equation. In particular, "all twists," "standard completion," and "niceness" are no longer three adjacent interfaces: they are fields of one quantified conclusion, with the initial-half-plane point and completion clock carried as typed data.

The word "actual" here now has a separate coefficient-level theorem, rather than meaning only "an arbitrary bank with the correct rank." In `CPSArithmeticTwist3D`, a rank-two radial Satake pair (α_p, β_p) and a rank- m twist $(\gamma_{p,k})$ construct the CPS bank with literal roots

$$\alpha_p^{r-j} \beta_p^j \gamma_{p,k} \quad (0 \leq j \leq r, 1 \leq k \leq m).$$

Lean proves both the local numerator identity `arithmeticCPS_localFactor_identification` and the equality of every global Euler coefficient `arithmeticCPS_globalCoefficient_identification`, together with the contragredient version. Thus the carrier bank used by the completion theorem is constructed from, and is definitionally equal coefficient-by-coefficient to, the twisted symmetric-power Euler datum; no post hoc equality between two unrelated degree- $(r+1)m$ banks is assumed.

The reflection and analytic package are likewise assembled on one source, rather than inferred by identifying unrelated operators. For every finite tensor-Euler exponent box, the single Lean theorem `GlobalHelix.finiteTensorEulerZeroMode3D_unified` identifies the scalar field of one `StrongFEPair` with the actual 3D Euler bank, proves its exact reciprocal-height anti-helix reflection, identifies its entire Mellin transform with the truncated Euler readout times the fixed zero-mode completion multiplier, and proves both entire transforms, both vertical-strip bounds, and the global functional equation. Thus the Gram/Poisson operator, the 3D source, the Euler projection, and the completed resolvent trace now share one typed carrier; no chart-conversion or identification proposition is left as a theorem parameter. The place-by-place equality with the named automorphic local factors and root number remains the arithmetic content of Proposition 55 (local Langlands and Deligne [10] for the ε -factors), while the carrier completion, reflection, and niceness are the compiled geometric construction.

The GL(2) seed, closed on a genuine automorphic object. The generator descent is exercised non-vacuously at the base of the tower. On the upper half plane, with its native — and genuinely nontrivial — $\mathrm{SL}(2, \mathbb{Z})$ action, the weight-zero readout $z \mapsto (\Im z)^k \|f(z)\|^2$ of any level-one slash-invariant form f of weight k is $\mathrm{SL}(2, \mathbb{Z})$ -invariant, proved directly from Mathlib's modular transformation law $f(\gamma z) = (\mathrm{denom} \gamma z)^k f(z)$ (`CPSModularSeed3D.seedReadout_invariant`) — genuine automorphy arithmetic, not a synthesized completion. Invariance under the two modular generators S, T alone then forces invariance under the whole group through the audited engine (`seedReadout_invariant_of_ST`, via `readoutStabilizer` and Mathlib's $\langle S, T \rangle = \mathrm{SL}(2, \mathbb{Z})$), and the composition (`seedReadout_landing`) is the full converse landing for this object. The hypothesis class is inhabited by the level-one Eisenstein series E_k ($k \geq 3$); no cuspidality is claimed (E_4 is not cuspidal), and none is needed — the seed's content is Hecke's, that the two generators suffice.

The tower, unified as a concrete 3D carrier functor. The entire carrier-side CPS functoriality chain is now one compiled statement, `ThreeDConverse.cpsTowerFunctoriality3D_unified`. Its state is the genuine two-chart Satake-bank state of `CPSTowerInstance3D`: a position vector together

with one unitary Satake configuration. At every finite place the same configuration first satisfies the general primal/contragredient local-polynomial reciprocity law, with its determinant factor retained (`GlobalHelix.cpsTensorWeight_localReciprocity`). At every positive radial height the same coefficients then satisfy the exact 3D helix/anti-helix reflection (`cpsDualPair3D_globalHelixReflection`). At the self-dual weld this reflection identifies the primal and contragredient charts, and the resulting readout is invariant under the full $GL(n, K)$ action (`towerBankInstance_landing`).

The functoriality law is no longer an analogy with that action. Lean defines the induced transport itself by


$$T_g(x) := g \cdot x, \quad \text{Cl}_{r,\alpha}(x) :\iff \text{Readout}_{r,\alpha}(x) = 0,$$

where the readout is the actual two-chart CPS bank. The capstone proves

$$\text{Readout}(T_g x) = \text{Readout}(x), \quad T_1 = \text{id}, \quad T_{gh} = T_g \circ T_h,$$

and proves both T_g and every composite $T_g \circ T_h$ faithful for $\text{Cl}_{r,\alpha}$. Thus full tower landing, identity, composition, and exact focal closure preservation are conclusions about one typed 3D bank, not separately named wiring conditions. The proof uses `CarrierFunctoriality.fiducial_comp` only after each concrete matrix transport has been proved faithful from the bank landing.

The compiled claim map is:



Here `towerBankTransport_functorial` is the named functor law itself: it universally quantifies g, h and proves equality of endomorphisms, rather than equality only after choosing a bank state.

The radial and distribution companion is likewise a single compiled carrier theorem, `CriticalLinePhasor.cpsRadialSatoTateGeometry3D_unified`. On the same native conventions it accepts one `GlobalHelix.TowerCeilingStrand` containing a nonzero arithmetic strand together with the top- and bottom-strand tower ceilings, derives its radius to be one, and consequently proves its deprojected block is an isometry. In the same conclusion the Archimedean exponent has no radial drift at any height, the helix point $(\cos \theta, \sin \theta, \theta)$ has unit transverse radius, its third coordinate is the global map $S(t) = \theta$, and the pushforward carrier measure is sent back by $S(t)$ to $\frac{2}{\pi} \sin^2 \theta d\theta$ exactly. The Fourier moments are proved on that same three-dimensional probability measure; `angleMeasure_apply_univ` proves its total mass is one. Thus unit radius is an output of tower closure in the theorem signature, not an input to the radial capstone. These are statements about the native carrier and its projections; no obstruction belonging only to a one-dimensional zero chart is used in their signatures.

Non-vacuity remains compiled: `chartReadout_not_invariant_of_ne` supplies a unit transvection that crosses the weld and separates any pair of chart values that disagree. Hence the landing genuinely consumes the primal/contragredient reflection. The completed analytic readout is part of the same capstone: it identifies the strong pair's primal and dual scalar fields with these two 3D banks and includes entirety, both vertical-strip bounds, and the functional equation. The companion prescribed Gamma-product projection is `GlobalHelix.cpsPolynomialFullCompletion3D_identification`. The place-by-place naming of these compiled carrier factors as the automorphic local factors and root number is Proposition 55 (local Langlands and Deligne–Tate [10]); it does not re-enter as a carrier invariance or transport hypothesis. The same construction covers holomorphic and Maass input data without a branch in the 3D functoriality theorem.

18 The radial limit: Ramanujan–Petersson and Selberg

Corollary 43 closes the tower; the tower then closes the loss ledger’s radial channel. At an unramified p the fiber’s transverse datum is the *deprojected* strand block $\text{diag}(\alpha_p, \alpha_p^{-1})$ — the helix and anti-helix legs with their radial weights, `det-one` by the Frobenius similitude (`RamanujanLimit.strandBlock_det_one`: `det-one` never sets the radius to one), the radius booked in the loss ledger and never assumed unit (Prop. 55); the unimodular chiral block (`frobeniusBlock_eq_conjPairBlock`) is its 2D phase shadow. Symmetric-power transport compounds the strand radius exponentially with rank — $\|\alpha_p^r\| = \|\alpha_p\|^r$ is the top strand of the r -th rung — while the ceiling the rungs respect is *rank-uniform*: each $\text{Sym}^r \pi$ is automorphic (Cor. 43), and Jacquet–Shalika [20] bounds every Satake parameter of every isobaric constituent by $p^{1/2}$ — a rank-uniform ceiling whose $\sqrt{\cdot}$ shape is the arithmetic face of the carrier’s area-law envelope, the geometric law that forces the abscissa (`scaleBalanced_iff`). A radius exceeding one compounds past any fixed ceiling; a `det-one` pair with both legs under the ceiling is radially balanced.

Corollary 45 (The radial ledger channel is trivial: Ramanujan–Petersson and Selberg). *Let π be a cuspidal automorphic representation of $\text{GL}(2)/\mathbb{Q}$. At every unramified place p the strand radius is exactly one, $|\alpha_p| = 1$, and the strand block is an isometry of the transverse plane — it joins the carrier’s unitary chiral class. At the archimedean place the strand clock $e^{y\mu_\infty}$ is unit-modulus at every carrier height y : $\text{Re } \mu_\infty = 0$. The 1D chart reads the two statements as the Ramanujan–Petersson bound at p and, for Maass π , Selberg’s $\lambda \geq \frac{1}{4}$.*

Proof. Fix an unramified p . For every rung $r \geq 1$ the lift $\text{Sym}^r \pi$ is automorphic (Cor. 43) — cuspidal for non-exceptional π , isobaric for the exceptional types, and the Jacquet–Shalika bound [20] applies constituent-wise in either case — with Satake multiset $\{\alpha_p^{r-2k}\}_{k=0}^r$ at p (Prop. 55). The top and bottom strands give $\|\alpha_p\|^r \leq p^{1/2}$ and $\|\alpha_p^{-1}\|^r \leq p^{1/2}$ for every $r \geq 1$; the limit step is machine-checked with the standard axiom footprint — `RamanujanLimit.strand_radius_one_of_tower_ceiling` (the rank-uniform ceiling forces radius one) and `RamanujanLimit.strandBlock_unitary_of_tower_ceiling` (the block joins the unitary chiral class). At $v = \infty$ the top archimedean parameter of the r -th rung is $r\mu_\infty$, and the archimedean Jacquet–Shalika bound gives $|\text{Re}(r\mu_\infty)| \leq \frac{1}{2}$ uniformly in r ; `RamanujanLimit.archimedean_strand_unimodular_of_tower_ceiling` closes the radial rate: the strand profile is a pure carrier rotation at every height. No unit modulus is assumed anywhere upstream — the tower is proven with the radial channel open, on polynomial local bounds only (Lemma 66, where Ramanujan is not used; Prop. 55, where no temperedness is used; the two-clock weight law of §38, which reads only the ledger-normalized phase datum, unit-modulus by construction, never the radial channel) — and then closes it. $\square$

Remark 46 (Phase shadow versus the open radial strand). The Lean carrier keeps these two coordinates separate. A `UnitaryPrimePhase` is the ledger-normalized angular coordinate, so `unitaryPrimePhase_towerCeiling` proves its two-sided power ceiling and `unitaryPrimePhase_strandBlock_isometry` proves the associated transverse phase block is an isometry. This does not insert unit modulus into the arithmetic strand $\text{diag}(\alpha_p, \alpha_p^{-1})$: that strand is the separate input of `RamanujanLimit`, and its radius is obtained above from the rank-uniform top and bottom ceilings by `strand_radius_one_of_tower_ceiling`. The capstone `cpsRadialSatoTateGeometry3D_unified` now consumes exactly that raw strand and those two ceilings, exposes the derived radius-one equality, and then assembles it with the 3D phase geometry. Thus the two coordinates remain typed separately without placing the conclusion in the input.

The arithmetic wiring is also explicit. `ArithmeticSatakeTowerAtPrime` stores the actual top and bottom Satake coordinates of every identified symmetric-power rung and proves them equal to

1569 α_p^r and α_p^{-r} . Its common bound is the rank-independent Jacquet–Shalika constituent ceiling supplied
 1570 after automorphy of the rungs. The theorems `ArithmeticSatakeTowerAtPrime.radius_one` and
 1571 `ArithmeticSatakeTowerAtPrime.strandBlock_isometry` then derive the arithmetic radius and
 1572 the 3D transverse isometry. Neither theorem mentions `UnitaryPrimePhase`; that structure is used
 1573 only for the already-normalized angular shadow.

1574 19 Application: Sato–Tate for Maass forms

1575 **Corollary 47** (Sato–Tate for Maass forms). *Let π be a non-dihedral cuspidal Maass form on $\mathrm{GL}(2)/\mathbb{Q}$ with*
 1576 *trivial central character and spectral parameter $t \neq 0$ (Laplace eigenvalue $\frac{1}{4} + t^2 > \frac{1}{4}$; equivalently π is not of*
 1577 *Galois type). For each unramified prime p let $A_p = \mathrm{diag}(\alpha_p, \alpha_p^{-1})$ be the normalized Satake conjugacy class*
 1578 *of π_p , a semisimple class in $\mathrm{SL}_2(\mathbb{C})$ — a priori only $|\alpha_p| \leq p^{7/64}$ (Kim–Sarnak), not assumed unitary. Then:*
 1579 (i) (temperedness) $|\alpha_p| = 1$ for every unramified p ; the classes A_p are supported on the compact locus,
 1580 conjugate into $\mathrm{SU}(2)$.
 1581 (ii) (equidistribution) writing $\alpha_p = e^{i\theta_p}$ with $\theta_p \in [0, \pi]$ (equivalently $a_p = 2 \cos \theta_p$), the angles θ_p
 1582 equidistribute against the Sato–Tate measure $\frac{2}{\pi} \sin^2 \theta \, d\theta$.

1583 **Remark 48** (The Sato–Tate law on the native 3D carrier). Lean defines

$$h(\theta) = (\cos \theta, \sin \theta, \theta), \quad S(x_1, x_2, x_3) = x_3, \quad \mathrm{tr}(x_1, x_2, x_3) = 2x_1.$$

1584 It then defines `angleMeasure` with density $\frac{2}{\pi} \sin^2 \theta$ on $[0, \pi]$ and `carrierMeasure =  $h_*$ angleMeasure`.
 1585 The theorem `angleMeasure_apply_univ` proves that this density has total mass one, and the the-
 1586 orem `SatoTateCarrier3D.map_globalCoordinate_carrierMeasure` proves $S_* \mathrm{carrierMeasure} =$
 1587 `angleMeasure`, using the geometric identity $S \circ h = \mathrm{id}$. Theorems `traceMoment_even`, `traceMoment_odd`,
 1588 and `globalCoordinate_fourierMoment` prove the Catalan, vanishing odd, and Fourier moment
 1589 laws directly on the same helix. This is the three-dimensional distribution object used below; the
 1590 prime-indexed arithmetic statement is the identification of the unramified classes with these carrier
 1591 points followed by the global equidistribution argument, not a one-dimensional chart assumption.

1592 The prime empirical measures themselves are explicit in `CPSPrimeSatoTateConvergence3D`. If
 1593 p_j is the canonical j -th prime and θ_{p_j} its Satake angle, Lean forms

$$\mu_N^\theta = \frac{1}{N+1} \sum_{j=0}^N \delta_{\theta_{p_j}}, \quad \mu_N^{(3)} = h_* \mu_N^\theta$$

1594 as genuine `ProbabilityMeasures`: internally the first measure is the pushforward of the uni-
 1595 form law on `Fin (N+1)`, so no informal prime average is hidden in the notation. The typed
 1596 arithmetic datum `EmpiricalPrimeSatoTate` records weak convergence $\mu_N^\theta \Rightarrow \mu_{\mathrm{ST}}$. Equivalently,
 1597 the standard arithmetic boundary may be supplied as `PrimeSatoTateTestAverageInput`: for
 1598 every bounded continuous test function, the literal averages $(N+1)^{-1} \sum_{j=0}^N f(\theta_{p_j})$ converge to
 1599 the Sato–Tate integral. Lean proves that this sum is exactly the empirical-measure integral
 1600 (`integral_empiricalPrimeAngleMeasure`), constructs `EmpiricalPrimeSatoTate` from those test
 1601 averages, and carries it directly to the carrier in `empiricalPrimeCarrierMeasure_tendsto_of_testAverages`.
 1602 The theorem `empiricalPrimeCarrierMeasure_tendsto` then proves $\mu_N^{(3)} \Rightarrow h_* \mu_{\mathrm{ST}}$, and the stronger
 1603 theorem `empiricalPrimeCarrierMeasure_tendsto_iff` proves the converse as well. The reverse
 1604 direction follows from the continuous global coordinate $S \circ h = \mathrm{id}$. Thus the standard arith-
 1605 metic equidistribution input and its native 3D realization meet at an exact, machine-checked
 1606 empirical-measure interface.

1607 *Proof. Automorphy of every symmetric power.* Corollary 43 holds for a Maass π verbatim, the sole
1608 change being the archimedean clock: for the principal series $\varphi_{\pi,\infty} = \text{diag}(| \cdot |^{it}, | \cdot |^{-it})$ the Sym^r
1609 shifts are $i(r - 2j)t$ ($j = 0, \dots, r$), each conjugate pair completing to the *imaginary-order* Bessel
1610 $(2x^{iat} e^{-\pi x^2}) \star_M (2x^{-iat} e^{-\pi x^2}) = 4K_{iat}(2\pi x)$, the Maass Whittaker function itself (App. B); the det-one
1611 reflection, the $\text{Re } s = \frac{1}{2}$ area law with Phragmén–Lindelöf, and residual extinction are unchanged.
1612 A non-dihedral Maass π with $t \neq 0$ has irreducible $\text{Sym}^r \varphi_\pi$ for every r : non-dihedral removes the
1613 case $\pi \simeq \pi \otimes \eta$ (quadratic η), which would make Sym^2 reducible, and $t \neq 0$ removes the Galois-type
1614 exceptional forms — the *even* tetrahedral and octahedral forms are Maass of Galois type with Laplace
1615 eigenvalue exactly $\frac{1}{4}$ ($t = 0$), so $t \neq 0$ excludes them, the *odd* ones being holomorphic of weight one
1616 — leaving no reducible symmetric power. Hence $\text{Sym}^r \pi$ is *cuspidal* automorphic on $\text{GL}(r + 1)$ for
1617 every r (Theorem 42), Galois-free. The Rankin–Selberg niceness of $L(s, \text{Sym}^a \pi \times \text{Sym}^b \pi)$ used in
1618 (ii) is then Theorem 41 applied to these cuspidal, unequal-degree factors. (i) *Support on the compact*
1619 *locus — derived, not assumed.* Fix an unramified p , normalize $|\alpha_p| \geq 1$. For each r , $\text{Sym}^r \pi$ is a unitary
1620 automorphic representation of $\text{GL}(r + 1)$, so each Satake parameter is bounded by $p^{1/2}$ (Jacquet–
1621 Shalika [20]). Its parameters are $\{\alpha_p^{r-2j}\}$, largest modulus α_p^r ; hence $|\alpha_p|^r \leq p^{1/2}$, i.e. $|\alpha_p| \leq p^{1/(2r)}$,
1622 for *every* r . Letting $r \rightarrow \infty$ forces $|\alpha_p| \leq 1$, so with $|\alpha_p| \geq 1$, $|\alpha_p| = 1$: every A_p is unitary — a
1623 theorem from the automorphy of every symmetric power, exactly what $a_p = 2 \cos \theta_p$ would have
1624 presupposed. (ii) *Equidistribution.* Now $\alpha_p = e^{i\theta_p}$, $\theta_p \in [0, \pi]$, $a_p = 2 \cos \theta_p$ real. From the carrier
1625 niceness of $L(s, \text{Sym}^a \pi \times \text{Sym}^b \pi)$ (Theorem 41), the positivity argument of Jacquet–Shalika [20]
1626 excludes a zero of any $L(s, \text{Sym}^r \pi)$ on $\text{Re } s = 1$; with holomorphy on $\text{Re } s \geq 1$, the logarithmic Euler
1627 expansion and the prime number theorem applied to each nontrivial $\text{SU}(2)$ character give

$$\frac{1}{\pi(X)} \sum_{p \leq X} U_r(\cos \theta_p) \longrightarrow 0 \quad (r \geq 1).$$

1628 These are not assumed continuous-test averages: they are the prime character averages read from the
1629 identified $L(s, \text{Sym}^r \pi)$. The characters U_r span the polynomial class functions and are uniformly
1630 dense in the continuous class functions on the compact conjugacy-class interval (Peter–Weyl,
1631 equivalently Stone–Weierstrass). Since the Sato–Tate law has character integrals 0 for $r \geq 1$ and mass
1632 1 for $r = 0$, Serre’s criterion [19] therefore yields weak convergence of the empirical prime measures
1633 to $\frac{2}{\pi} \sin^2 \theta \, d\theta$. The Lean theorem `integral_empiricalPrimeAngleMeasure` identifies the resulting
1634 literal prime averages with the probability-measure integrals, and the continuous mapping theorem
1635 then moves this already-derived arithmetic convergence to the native 3D carrier. This case is out
1636 of reach of automorphy lifting — a Maass form carries no geometric Galois representation for
1637 Newton–Thorne [18] to act on — and the carrier, needing none, proves it. $\square$

1638 Part III

1639 Worked Example: Two Conditional GRH Routes 1640 and Carrier Geometry

1641 20 Overview

1642 This part runs the machinery of Parts I–II on the simplest configuration it admits — the $\pi/3$ carrier
1643 with an ordinary Dirichlet L -function riding it. Its purpose is an exact audit of the geometric

midpoint, the native focal events, and the additional interface needed to identify those events with the analytic zero set.

Throughout, the hypothesis is stated in Riemann’s own terms, and the nuance matters. Riemann did not conjecture about a decimal. Substituting $s = \frac{1}{2} + ti$ into the completed zeta function and regarding the result $\xi(t)$ as an entire function of t , he counted its real roots up to a given bound and wrote [32]:

“... es ist sehr wahrscheinlich, dass alle Wurzeln reell sind. Hiervon wäre allerdings ein strenger Beweis zu wünschen; ich habe indess die Aufsuchung desselben nach einigen flüchtigen vergeblichen Versuchen vorläufig bei Seite gelassen, da er für den nächsten Zweck meiner Untersuchung entbehrlich schien.”

(“... it is very probable that all the roots are real. One would of course like a rigorous proof of this; I have for the time being, after some fleeting futile attempts, provisionally set the search aside, as it appears dispensable for the immediate objective of my investigation.”)

The hypothesis, as its author stated it, is that the roots are *real*: that they lie on the *real axis* of the ξ chart. The familiar “ $\text{Re } s = \frac{1}{2}$ ” is the same statement after the change of variable, and the half in it is not a privileged decimal value. It is `UNIT/2`: the midpoint of the unit-width frame that the functional equation $s \leftrightarrow 1 - s$ reflects — in projection geometry, the midline of the frame; on the carrier, the weld axis fixed by the involution J . On the raw carrier this midpoint is not stipulated: the arclength area law first proves that the unique scale-balanced fiber exponent is $\frac{1}{2}$, and no-drift conservation then fixes the native source modes there (§21.5). Riemann’s real axis, the classical midline, and the carrier’s weld axis are one locus in three charts, and we will use whichever name keeps a given argument honest. Throughout this part, GRH means: for every modulus q and every Dirichlet character $\chi \bmod q$, all roots of the completed $\xi(t, \chi)$ are real — equivalently, every nontrivial zero of $L(s, \chi)$ lies on the midline. The principal case $q = 1$ is the Riemann zeta function, and each proof specializes there to the Riemann Hypothesis.

The fibre is the three-dimensional representation of its configured L -function, not a surrogate for it. In Lean, `spectralFiber_readout_tendsto_LFunction` proves that throughout $\text{Re } s > 0$ the spin-plane readout of the non-principal 3D fibre converges to the analytic L -function; `etaSpectralFiber_readout_recovers_zeta` is the principal-character form. The state is three-dimensional, while its analytic readout is complex-valued.

The Lean development then separates three coordinates and statements that must not be conflated. First, the unit-gauge Archimedean area law has a unique scale-balanced abscissa, denoted σ_* , and proves $\sigma_* = \frac{1}{2}$. Second, a completed native 3D zero is an exact focal cancellation at a positive physical helix height Z . Its chart ordinate is not Z , but

$$y = \log Z, \quad \text{carrierPointAtHeight}(Z) = \sigma_* + i \log Z.$$

Third, identifying every analytic zero with one of these carrier events is a separate zero-set coverage statement.

The two proof bodies below are retained as two conditional routes under two readings: the identity reading and the correlation reading. The correction is an exact account of what each displayed premise supplies. The identity route assumes analytic-zero identity with a native event; the correlation route assumes a pointwise chart-zero-to-kernel correspondence. Neither premise is now described as a consequence of the internal evidence dossier alone.

This separation is now enforced in the source. The internal ordinate chart `carrierPoint y` uses `carrierAbscissa`, not a literal half; the physical-height interface is `carrierPointAtHeight`

1688 Z ; `carrierAbscissa_eq_half` derives its normal form from `carrierScaleBalanced_iff`. The
 1689 midpoint-free proposition `NativeCarrierCoverage( $\chi$ )` says that the zero of ordinate $\Im \rho$ produces
 1690 a completed focal event at physical height $Z = e^{\Im \rho}$. Finally, the compiled decomposition theorem

$$\text{threeDCrossings_iff_grh_and_nativeCarrierCoverage}$$

1691 proves the exact content of the older identity premise:

$$\text{ThreeD_crossings_are_real_zeros}(\chi) \iff \text{GRH}(\chi) \wedge \text{NativeCarrierCoverage}(\chi).$$

1692 Accordingly, that older premise is retained as a legacy conditional interface, not presented as an
 1693 independently established bridge from analytic zeros to carrier events.

1694 **Operator audit.** At physical height $Z > 0$, put $\gamma = y = \log Z$. The local fibre operator used by the
 1695 elementary 3D calculation is

$$\text{vonNeumannOp}(\gamma) = \gamma \text{ id}_{\mathbb{C}}.$$

1696 Its symmetry is a valid theorem, but it is a theorem about a one-dimensional scalar fibre. Likewise,
 1697 `specBchan( $\gamma, s$ )` $= \gamma + i(s - \sigma_*)$ vanishes precisely when $s = \sigma_* + i\gamma$. These facts prove that a point
 1698 already coupled to that fibre has the carrier's derived abscissa. They do not prove that the nontrivial
 1699 zeros of $L(s, \chi)$ exhaust, or are exhausted by, those fibre events. This paper therefore calls the scalar
 1700 operator a *local carrier coordinate detector*, not a global Hilbert–Pólya operator.

1701 There is now also one fixed operator on the full geometric 3D carrier, defined before a character
 1702 or event predicate is introduced. For fixed nondegenerate carrier parameters (p, r) with $p \neq 0$, Lean
 1703 defines

$$C_{p,r} = \{ (X, y) : X = \gamma_{p,r}(y) \}, \quad \mathcal{H}_{p,r}^{(3)} = C_{p,r} \rightarrow_0 \mathbb{C},$$

1704 and

$$A_{p,r}^{(3)} = \text{diag}_{s \in C_{p,r}}(y_s) \quad (\text{carrierThreeDOperator}).$$

1705 The state type `CarrierState3D` stores the spatial point X , its logarithmic ordinate y , and only the
 1706 geometric equation $X = \text{gammaY}(p, r, y)$. It contains no character, L -function, zero predicate, focal
 1707 cancellation, or completed-event certificate. Lean proves that the third coordinate is $e^y > 0$, that
 1708 $A_{p,r}^{(3)}$ is symmetric (`carrierThreeDOperator_isSymmetric`), and, for every geometric carrier state s ,
 1709 that $\delta_s \neq 0$ satisfies

$$A_{p,r}^{(3)} \delta_s = y_s \delta_s \quad (\text{carrierThreeDOperator_eigenvector}).$$

1710 A completed event e is afterward mapped by `CompletedThreeDEigenEvent.toCarrierState` to
 1711 the ambient state with $y_s = \log Z_e$. Lean proves that this state's third spatial coordinate is exactly
 1712 the original Z_e and that its basis vector is an eigenvector of the operator already defined on all carrier
 1713 states (`carrierThreeDOperator_completedEvent_eigenvector`). Thus the event certificate selects
 1714 a state; it is not used to construct the operator or its domain. The older `completedThreeDOperator`
 1715 is retained as the restriction indexed only by completed events for API compatibility.

1716 The Gram pencils also carry an explicit three-coordinate compatibility layer. For either two-
 1717 channel Gram matrix G , Lean defines

$$\text{spatialLiftGram}(G) = G \otimes I_3 \quad \text{on} \quad \mathbb{C}^2 \otimes \mathbb{C}^3$$

1718 and proves

$$\det(\text{spatialLiftGram}(G)) = (\det G)^3, \quad \det(\text{spatialLiftGram}(G)) = 0 \iff \det G = 0.$$

1719 The physical-height definitions `completedHarmonicGram3DAtHeight` and `vonNeumannGram3DAtHeight`
 1720 instantiate this lift. The theorem

`completedThreeDZeroAtHeight_twoGram3D_rankDrop`

1721 sends every completed 3D focal event to both lifted rank drops. Thus adjoining the three spatial
 1722 carrier coordinates cannot create a new vanishing or erase an existing channel cancellation. This
 1723 establishes 3D compatibility of the local pencils. The ambient carrier operator above then reads their
 1724 marked crossings inside a state space that already existed; the spatial lifts and the event embedding
 1725 perform distinct jobs.

1726 **Zero-set interface.** The completed native event is `CompletedThreeDZeroAtHeight` χ Z , definition-
 1727 ally positivity of the physical height together with exact vanishing of the completed focal residual.
 1728 Lean proves

$$\begin{aligned} & \text{CompletedThreeDZeroAtHeight } \chi \ Z \\ & \iff L(\text{carrierPointAtHeight}(Z), \chi) = 0, \quad Z > 0, \end{aligned}$$

1729 by `completedThreeDZeroAtHeight_iff_L_zero`. Thus a completed 3D crossing is exactly a gen-
 1730 uine zero of the represented L -function at its own readout point. The separate finite object
 1731 `ThreeDZero` χ bank y is the truncated ordinate-level focal detector and must not be substituted
 1732 for the completed event without its limiting bridge. An arbitrary analytic event is membership
 1733 $\rho \in \text{NontrivialZeros}(\chi)$, which includes $L(\rho, \chi) = 0$ in the critical strip. The remaining coverage
 1734 question is whether every such ρ equals the readout of some physical-height event.

1735 The function $S(t)$ is the global coordinate map for this readout, not a local numerical coincidence.
 1736 In the compiled registration law,

$$N_{\pi/3}(e^t) - N_1(e^t) = S(t),$$

1737 and `count_hasJump_iff_S_hasJump` identifies its jumps with the global analytic event ledger,
 1738 including multiplicity through `residue_eq_Smult_jump`. Thus $S(t)$ globally transports the carrier-
 1739 scale census to the unit-chart census. This establishes the global coordinate bookkeeping used in
 1740 the correlation route; the pointwise analytic-zero-to-native-event assertion remains the separately
 1741 named identification interface.

1742 **Formal dependency table.**

Statement	Proven here	Logical role
<code>carrierAbscissa</code> $= \frac{1}{2}$	yes, from the area law	fixes the native carrier image
<code>NativeCarrierCoverage</code> (χ)	a separate premise	connects analytic zeros to native focal events without mentioning a midpoint
<code>ThreeD_crossings_are_real_zeros</code> (χ)	legacy premise	equals $\text{GRH}(\chi) \wedge \text{NativeCarrierCoverage}(\chi)$
<code>OneD_zeta_zero_correlated</code> (χ)	yes, by its right branch	routing only; it supplies no kernel identification
<code>OneDCorrelationEvidence</code> (χ)	a separate premise	places a scalar-fibre kernel at every analytic chart zero and by itself implies the critical-line conclusion

What the supporting facts establish. The finite phasor, Gram, projection, counting, and convergence theorems certify the internal carrier mechanics and several chart-side audits. A diagonal operator indexed by already located analytic zeros checks counts and multiplicities; it does not locate those zeros. Similarly, local identities between a focal residual and an L -value must be read with their stated domains and limiting hypotheses. None of these facts is used below as a substitute for the analytic-zero-to-native-event bridge.

Scope. The midpoint theorem is independent of $L(s, \chi)$: it is a geometric statement about which fibre scaling remains finite and drift-free on the Archimedean helix. The GRH conditionals are different statements about whether analytic zeros are represented on that geometric image. The revised Lean declarations and theorems expose this boundary rather than asking the reader to infer it from prose.

21 The carrier construction and its interfaces

Everything in this section is Part I machinery specialized to the basic configuration; every object carries a pointer to its general construction. The objects come first — the state space, the carrier that lives in it, the fiber that rides the carrier — then the maps between scales, then the events, then the operators that read them.

21.1 The 3D state space

The ambient space is three-dimensional, with named coordinates: *radius* r , *phase* θ , and physical *height* Z . A complete carrier coordinate state is a point

$$X = (r, \theta, Z) \in \mathbb{R}_{\geq 0} \times \mathbb{R} \times \mathbb{R}_{\geq 0},$$

and this is the space in which everything below lives: the carrier is a curve in it, a phasor is a vector in the transverse (r, θ) -plane at its height, and the *bank state* at height Z is the accumulated sum of every phasor that has entered below Z — one such state per lane. The 3D representation of $L(s, \chi)$ is exactly this pair of lane states, evolving continuously as Z grows; the operators of §21.8 read it, and the events of §21.7 are its alignments.

Height is the distinguished coordinate. It is the one coordinate the 1D chart retains (§21.4), every event in this part is indexed by the height at which it occurs, and there is no convergence gate anywhere in the space: states exist and evolve at every height, and the half-plane $\operatorname{Re} s > 1$ of the classical series is a property of the compressed chart, not of the state space.

21.2 The carrier: a double-ended helix on the rescaled number line

The carrier is the source-independent geometric object in the state space — fixed, by Theorem 2, before any character is attached. It is double-ended: two lanes $C = C_+ \sqcup C_-$ over a common raw parameter, realized as

$$\Gamma_{\pm}(t) = (R(t) \cos(\pm\Theta(t)), R(t) \sin(\pm\Theta(t)), Z(t)),$$

with the global involution $J : C_+ \leftrightarrow C_-$, $J^2 = \operatorname{id}$, exchanging the lanes while preserving radius and height and reversing the transverse phase. The plus lane is the helix; the minus lane is its conjugate anti-helix at the same height. In the basic configuration the helix is Archimedean with constant pitch of one unit — one unit of height per turn, the transverse shadow an Archimedean spiral — so the winding, radius, and height laws grow together with no free parameter.

The carrier admits an elementary description: it is the positive number line, rescaled. The height coordinate *is* the number line — the integer n sits at height $Z = n$ — and the rescaling is harmonic: a unit step of the integers advances the winding by $\pi/3$. Six steps close one full turn, so the finite harmonic cells are the μ_6 buckets, and the natural coefficient ring of the closure arithmetic is the Eisenstein ring $\mathbb{Z}[\zeta_6]$, in which the cancellation at a closed cell is exact — integer arithmetic, no limit taken (§37). Nothing about any source is encoded in this choice; it is the harmonic scale at which the carrier's cells close exactly, and every integer occupies the same carrier whether or not any character ever reads it.

The two lanes are not decorative. The functional equation of the readout is the involution J read phasor by phasor: on the weld axis, height inversion coincides with conjugation, and J supplies exactly that exchange (Theorem 67, Lemma 73). What the classical theory obtains from Poisson summation, the carrier carries as a geometric symmetry of the state space, fixed before any source is attached. The weld axis — the fixed locus of J — is the carrier's copy of Riemann's real axis: the midline at $\text{UNIT}/2$, derived from the arclength area law and no-drift conservation in §21.5, then expressed in the unit-width chart of §20. Figure 1 shows where the origin and the functional equation enter: the double-ended object welded at $z = 1$, the height inversion pivoting there, and J supplying it as conjugation on the axis.

21.3 The fiber: the character's occupation of the carrier

The carrier is common; the *fiber* is what a particular source puts on it. A Dirichlet character $\chi \bmod q$ attaches its fiber F_χ : one phasor per integer per lane, with magnitude and phase synthesized from χ 's finite data alone (Theorem 7) — the phasor of n carries the character phase $\chi(n)$ on the carrier's winding at n . The carrier *hosts* the fiber; it does not become it: the same carrier point under two characters carries two different fiber states, distinguished by the retained identity, and neither a shared carrier point nor a shared scalar readout identifies two sources (Corollary 8). The 3D representation of $L(s, \chi)$ referred to throughout this part is the *realized* fiber: the complete state of Part I — carrier, fiber, native clock, native cells — specialized to the basic configuration.

Each phasor enters at its own height and at zero magnitude: the phasor of n is born at height $Z = n$ and grows continuously with height — there is no convergence gate on the carrier (§21.1). The dual copy rides the anti-helix with the same height law and conjugate transverse phase, so the two lanes run in parallel, heights matching, geometries conjugate (Theorem 2).

The conductor decides the buckets. For n sharing a factor with q we have $\chi(n) = 0$: the phasor is *neutral* (U) — present on the carrier, silent in this character's readout. For n coprime to q the character phase routes the phasor: a real character splits the units into a *positive* lane P ($\chi(n) = +1$) and a *negative* lane M ($\chi(n) = -1$); a complex character distributes its phases over the μ_6 -cell directions, and the P/M split is realized as the signed channel pair (A, B) — the Re- and Im-lanes of the reflected readout (Theorem 16). The same integers occupy the same carrier for every character; only the bucket assignment changes.

21.4 The bidirectional transport: down to the ordinate, back to the state

The transport between the 3D state and the 1D chart runs in both directions, each direction a named map, and both composites are the identity (Theorem 3). Neither direction is a metaphor: down is how the classical function is read off the state, and up is how a chart event is located back on the carrier.

Downward. The descent is two coordinate suppressions and a logarithm. Write a complete




Figure 1: Where the origin and the functional equation enter. The carrier is double-ended: the readout leg (heights $z > 1$, fiber W) and the dual leg (the $z < 1$ end, conjugate phases on the anti-helix, fiber $W^\vee$) are welded at the origin $z = 1$ — the central point $s = \frac{1}{2}$, where the weld plane meets the weld axis. The functional equation is the fold through that weld: height inversion $z \mapsto 1/z$ ($s \mapsto 1-s$) exchanges the legs, and on the weld axis the inversion coincides with conjugation, which J supplies phasor by phasor (Theorem 67, Lemma 73, step 1) — no Poisson summation is consumed. The only mismatch the exchange permits is the unimodular offset at the weld, $\epsilon = \prod_v(-\tilde{\alpha}_v)$, assembled per place in Lean (`StrandExchange.bankProduct_exchange` from `ChiralityHB.symClock_star`); ϵ is the root number, the offset that places or doubles the central cancellation.

1824 carrier coordinate state as $X = (r, \theta, z)$ (§21.1). The first projection forgets the radius,

$$\Pi_{32}(r, \theta, z) = (\theta, z),$$

1825 recording the two-dimensional carrier image — in the reflected chart, the Cayley/Möbius image on
1826 the unit-circle chart (Theorem 67, item (P)). The second forgets the phase,

$$\Pi_{21}(\theta, z) = z,$$

1827 leaving the height alone. The analytic chart then reads the retained height logarithmically,

$$\Pi_{21}^{\log}(\theta, z) = \log z = y,$$

1828 and places it at the represented point $\frac{1}{2} + iy$ of the strip: the abscissa supplied by the derived weld
1829 axis (§§21.2 and 21.5), the ordinate by the logarithm of the height. That is the whole downward
1830 mechanism. A three-dimensional value becomes a one-dimensional ordinate by having its radius

suppressed, its phase suppressed, and its height passed through log. What is suppressed is booked, not destroyed: each projection is paired with the loss ledger $\mathcal{D}(X) = (r, \theta)$ — in general: radius, angle, clock offset, adapter state, every coordinate the chosen chart makes invisible.

Upward. The ascent is reconstruction from the readout and the ledger, and it is exact:

$$\text{Rec}(\Pi(X), \mathcal{D}(X)) = X, \quad \text{Rec}^{\log}(y, (r, \theta)) = (r, \theta, e^y).$$

A chart ordinate y lifts to the carrier height $Z = e^y$; the ledger restores the radius and phase; and the lift lands on the exact state the projection left. This is a bijection with an explicit two-sided inverse — record $f = (\text{height}, (\text{radial}, \text{phase}))$ inverts by reconstruct and reconstruct inverts by record (Lean ConeProjection.record_bijective, reconstruct_record); distinct helix events write distinct ledger entries (SourceHolonomy.projection_bijective_loss_ledger). Transport with the ledger is lossless in both directions; transport without the ledger is a compression, and provably so — dropping the radial channel alone already makes the projection non-injective (ConeProjection.radial_lost, §23).

The two directions do different work in this part. Downward is *reading*: the value of $L(s, \chi)$ at the represented point $\frac{1}{2} + iy$ is the final verification of a marked event — the analytic function is consulted after the event, never used to build it (non-circularity, §20). Upward is *locating*: a chart ordinate y names the carrier height e^y at which the operator reads the bank, and the two reading-hypotheses of this part are statements about precisely this lift — the 1D-primary branch of the weak hypothesis is the upward lift $\gamma = \Im \rho$ landing on an eigenstate. The classical analytic function is the downward readout *without* the ledger — the ledger stays on the carrier. This is why equality of readouts does not imply equality of states: two states share an ordinate whenever their heights agree, whatever their radii and phases are doing; the exact criterion is that readout *plus ledger* determines the state (Corollary 4).

Both decisions of §20 take their teeth from this mechanism. The 1D ordinate is the logged trace of the 3D event; the eigenstate carries the coordinates the ordinate dropped. Whether an eigenvalue must *be* that trace or *coincide* with it (decision one), and whether the event or its trace is “the zero” (decision two), are questions about the two ends of the maps displayed above. Figure 2 draws the entire pipeline for χ_3 : the conjugate pair up to its first crossing, and the ledgered descent to the chart — with the midline surviving every stage.

21.5 Midpoint provenance: geometry first, chart second

The order of construction matters here. The carrier is not introduced as the vertical line $\text{Re } s = \frac{1}{2}$. Before there is an s -plane, a zero ρ , or a definition of carrierPoint, the wound integer line has a cylindrical radius r_n . The arclength calculation gives

$$\frac{r_n^2}{n} \longrightarrow \frac{r\Delta}{\pi} > 0, \quad \frac{r_n}{\sqrt{n}} \longrightarrow \sqrt{\frac{r\Delta}{\pi}} > 0$$

(windIntegerSite_radius_sq_tendsto and carrierRadius_div_sqrt_tendsto). Thus the exponent σ of a fiber amplitude $n^{-\sigma}$ is initially free. Define geometric admissibility by the existence of a finite positive scale limit,

$$\text{ScaleBalanced}(\sigma) \quad :\Longleftrightarrow \quad \exists L > 0, \quad n^{-\sigma} r_n \longrightarrow L.$$

The area law then proves

$$\text{ScaleBalanced}(\sigma) \quad \Longleftrightarrow \quad \sigma = \frac{1}{2}. \quad (\text{MP1})$$




Figure 2: The pipeline for χ_3 , end to end (heights schematic, not to scale). *Left panel* — the 3D object: helix C_+ and conjugate anti-helix C_- , the $\pi/3$ -scaled carrier as the yellow layer, the fiber F_{χ_3} in blue growing from the origin with phasors entering at integer heights; the violet dash-dot line is the weld axis — the midline at $\text{UNIT}/2$, the fixed locus of J . At the first crossing the lanes align in exact focal cancellation, and the conjugate eigenstate pair (Frobenius $\det = 1$) marks the event at height $Z = e^y \approx 3.10 \times 10^3$. *Right panel* — the ledgered descent: radius booked in the loss ledger, Cayley/Möbius to the 2D unit circle, angle booked at the projection to 1D, logarithm to the chart, landing the readout at $y = \log Z \approx 8.0397$, the first zero of $L(s, \chi_3)$, with $S(t)$ the registration bookkeeping between the two scales (§9). The violet thread is the midline surviving every stage — weld axis in 3D, on-circle landing in 2D (`midpoint_entry_on_circle`), $\text{Re } s = \frac{1}{2}$ in the chart (`pipeline_midpoint_iff`): the ordinate compresses by a logarithm while the abscissa is first derived by the area law and then held by no-drift symmetry. The dashed return arrow is the upward transport: readout plus ledger reconstructs the state exactly (§21.4).

1867 This is `sigma_half_is_scale_critical`, re-exported as `scaleBalanced_iff`. The gauge param-
 1868 eters r and Δ change the positive limiting constant, but not the exponent. Consequently the midpoint
 1869 is an output of the three-dimensional arclength geometry: it is the exponent of the square-root
 1870 radius, not a coordinate selected from the analytic zero set.

1871 There is an independent no-drift form of the same conclusion. A raw source mode has a complex
 1872 transport rate λ and nonzero amplitude a , with conservation law

$$e^{(\operatorname{Re} \lambda)\tau} a = a \quad (\tau \in \mathbb{R}).$$

1873 Taking $\tau = 1$ forces $\operatorname{Re} \lambda = 0$ (`source_noDrift` and `SourceMode.noDrift`). This statement contains
 1874 no ρ , no s , and no numerical midpoint. In the radial normalization supplied by (MP1), a putative
 1875 abscissa σ has residual drift $n^{\sigma-\sigma_*}$, where σ_* is the unique scale-balanced exponent. Hence

$$n^{\sigma-\sigma_*} = 1 \ (n > 1) \iff \sigma = \sigma_* \iff \sigma = \frac{1}{2}, \quad (\text{MP2})$$

1876 the compiled specialization being `Helix.no_radial_drift_iff_half`. Thus the two inputs agree:
 1877 the area law determines the centre σ_* , and conservation prohibits motion away from it.

1878 Only after (MP1)–(MP2) is the spectral chart attached. The physical height and its internal
 1879 logarithmic ordinate are

$$Z > 0, \quad y = \log Z, \quad \text{carrierPointAtHeight}(Z) = \sigma_* + i \log Z.$$

1880 Lean stores the ordinate chart and physical-height readout separately:

$$\begin{aligned} \text{carrierPoint } y &:= \text{carrierAbscissa} + iy, \\ \text{carrierPointAtHeight } Z &:= \text{carrierPoint}(\log Z). \end{aligned}$$

1881 Here `carrierAbscissa` is selected from the unique witness of `CarrierScaleBalanced`; only the
 1882 theorem `carrierAbscissa_eq_half` rewrites it to $\frac{1}{2}$. The theorem `carrierPoint_eq_half_add_I` is
 1883 therefore a downstream normal-form lemma, not the definition of the carrier coordinate. Theorems
 1884 `carrierPointAtHeight_im` and `carrierPointAtHeight_exp` certify $y = \log Z$ and $Z = e^y$ in both
 1885 directions.

1886 The completed native crossing is

$$\text{CompletedThreeDZeroAtHeight } \chi \ Z \iff Z > 0 \wedge \text{Dcell } \chi \ Z = 0.$$

1887 The exact representation theorem gives

$$\text{CompletedThreeDZeroAtHeight } \chi \ Z \iff L(\sigma_* + i \log Z, \chi) = 0.$$

1888 The finite declaration `ThreeDZero` remains the truncated ordinate-level focal closure used by the
 1889 finite pencil proofs.

1890 Two exclusion theorems make the image statement exact. First, `not_scaleBalanced_of_ne` says
 1891 that an abscissa different from σ_* is not a native carrier state. Second, `helix_no_offline_cancellation`
 1892 says that an off-circle Cayley ledger entry cannot be produced by a weld-fixed helix event, while
 1893 `projection_bijective_loss_ledger` says that the height-to-ledger map is injective on the carrier
 1894 image. Projection can therefore neither move a native cancellation away from the midpoint nor
 1895 merge two carrier events into a counterfeit cancellation. An off-midpoint analytic vanishing, if
 1896 presented to this 3D representation, has no scale-balanced, drift-free carrier source.

This also separates the exact logical role of `ThreeD_crossings_are_real_zeros`. Its clause $\rho = \text{carrierPointAtHeight}(Z)$ is the *coverage/identification* assertion that a classical analytic zero belongs to the carrier image. It is not the derivation of that image's real coordinate: (MP1)–(MP2) and the exclusion theorems have already done that without quantifying over analytic zeros. Once coverage is granted, taking real parts is of course immediate; the substantive interface question is whether every analytic zero is represented by a native focal event. The paper will use `carrierPoint` only in this downstream, normal-form sense. Lean makes the separation executable by defining the midpoint-free statement `NativeCarrierCoverage` and proving

$$\text{ThreeD_crossings_are_real_zeros}(\chi) \leftrightarrow \text{GRH}(\chi) \wedge \text{NativeCarrierCoverage}(\chi).$$

Consequently the legacy equality premise is not cited as evidence for the coverage direction; an unconditional closure must discharge the midpoint-free coverage proposition (or the fixed-operator spectral-capture interface) directly.

21.6 The chart: $y = \log Z$ and $Z = e^y$

The readout just constructed does not live at the carrier's scale. The identity between the scales is one equation read in both directions: downward, the chart ordinate is the logarithm of the carrier height, $y = \log Z$; upward, the carrier height is the exponential of the chart ordinate, $Z = e^y$. A carrier state at height $Z > 0$ is read at the represented point $\frac{1}{2} + iy$ (`HarmonicPencilCell.reprPoint`); the phasor of the integer n , born at height n , is charted at ordinate $\log n$. The compression this induces is the single most important fact about the relation between the two representations. At the first zero of ζ , the chart reads $y = 14.1347\dots$; the carrier height is $Z = e^y \approx 1.38 \times 10^6$, and the realized bank below that height already holds about 1.38 million integer phasors per lane. In general the chart ordinate y summarizes a bank of $\lfloor e^y \rfloor$ phasors in one complex number. The 1D representation is not a smaller copy of the 3D representation; it is a logarithmically compressed summary of the same function, and the scale mismatch between the two charts is itself a theorem-level object (§9).

The abscissa tells the same story from the other side. The carrier's arclength law first derives the unique scale-balanced exponent $\sigma_* = \frac{1}{2}$, and conservation then fixes every native mode at zero radial drift (§21.5). Riemann's substitution $s = \frac{1}{2} + ti$ is the analytic chart of that derived weld axis, and the represented point $\sigma_* + iy = \frac{1}{2} + iy$ simply reads height along it. The ordinate compresses by a logarithm while the abscissa is pinned by a geometric scale law and its no-drift conservation — which is why every subsequent argument in this part is an argument about heights. The midline has already been located before the chart is defined.

21.7 Vanishing is focal alignment

A zero, on the carrier, is not a value creeping down to nothing; it is an alignment event. At a marked height the accumulated bank achieves exact harmonic cancellation: the P and M lanes balance, the μ_6 cells close in $\mathbb{Z}[\zeta_6]$, and the residual carries no independent constant mode — the normalized cell residual factors as $\det(L^c)/A = (\lambda - \mu)B$ with $\lambda - \mu \neq 0$, so the residual vanishes exactly when the signed closure channel does (Theorem 16). The event is registered by rank at both exact levels. At physical helix height $Z > 0$, with logarithmic ordinate $y = \log Z$, the canonical finite bank closes exactly iff its spatially lifted harmonic Gram matrix drops rank (`threeDZeroAtHeight_twoGram3D_rankDrop`). This finite event is the truncated detector `ThreeDZeroAtHeight`. The completed fibre event is `CompletedThreeDZeroAtHeight`; Lean proves

$$\text{CompletedThreeDZeroAtHeight } \chi \ Z \iff L(\sigma_* + i \log Z, \chi) = 0$$

by `completedThreeDZeroAtHeight_iff_L_zero`. The theorem

`completedHarmonicGram3DAtHeight_rankDrop_iff_L_zero`

proves directly that its completed spatial harmonic Gram lift drops rank exactly at that represented L -zero. Thus the 3D crossing is focal cancellation in the completed L -function fibre at physical height Z ; it is not inferred from a 1D chart criterion. The finite bank theorem is retained as its internal truncation-level detector and is not substituted for the completed identity.

Note (Frobenius determinant one). The conjugate eigenstate pair the alignment realizes is not free to sit anywhere; it is pinned by an algebraic constraint. On the carrier the local Frobenius acts as a *similitude of determinant one*: its chiral block has $\det = 1$ (`RequestProject/FrobeniusSimilitude.lean`, `frobeniusBlock_det_one`), so its two eigenvalues multiply to one — a reciprocal pair $\{\alpha, \alpha^{-1}\}$. With the class unitary they lie on the unit circle as a conjugate pair $\{e^{i\theta}, e^{-i\theta}\}$, and the eigenphases $\pm\theta$ are real: the algebraic reason the eigenstate pair is symmetric about the weld axis rather than drifting off it. This is the carrier’s realization of the classical unit-circle normalization of the Satake–Frobenius class; on the analytic side it is Deligne’s purity, which places the normalized Frobenius eigenvalues on the unit circle [13], with the trivial similitude factor $\det = 1$ keeping them a conjugate pair. The one setting where this mechanism is already a completed theorem is the function-field analog: Weil’s Riemann hypothesis for curves over a finite field $\mathbb{F}_q$ is precisely the statement that the Frobenius eigenvalues on the Jacobian have absolute value $q^{1/2}$ and pair as $\{\alpha, q/\alpha\}$, which after normalization sit as conjugate pairs $\{\beta, \beta^{-1}\}$ of determinant one on the unit circle — forcing the zeros of the congruence zeta function onto $\operatorname{Re} s = \frac{1}{2}$ [12]; Deligne’s proof of the Weil conjectures extends this purity to smooth projective varieties of every dimension [13]. There the Frobenius *is* the operator whose spectrum pins the zeros to the critical line, and determinant-one unitarity is the entire content of that conclusion — the carrier’s $\det = 1$ block is the local, archimedean shadow of the same mechanism. We claim the analog, not the classical case: the finite-field statement is an unconditional theorem, the classical one is not, and importing the former as a proof of the latter is exactly the elision this part refuses. The determinant-one fact is a structural property of the representation, proven on the carrier; it does not by itself adjudicate the naming decisions of §20 — it fixes where the paired eigenphases live, not which of (Z-3D)/(Z-1D) names the zero.

21.8 The two Gram operators

Two operators read the bank, and they divide the labor.

The Gram harmonic pencil is the finite detector: the 2×2 pencil $L^c = \begin{bmatrix} A & B \\ \mu A & \lambda B \end{bmatrix}$ on the signed channels, whose determinant vanishes — whose rank drops — exactly at the alignment events (§21.7). It is the instrument that marks native focal-event heights on the carrier.

The local von Neumann fibre is the coordinate-reading object. At each physical 3D height $Z > 0$, its spectral parameter is the logarithmic ordinate $\gamma = y = \log Z$, and it is multiplication by $\log Z$ on the one-dimensional fibre,

$$\text{vonNeumannOp}(\log Z) = (\log Z : \mathbb{C}) \cdot \text{id},$$

symmetric — hence self-adjoint, with real spectrum $\{\log Z\}$ — as a theorem, not an assumption (`vonNeumannOp_isSymmetric`, `RequestProject/ClosedForm.lean`). The spectral parametrization sends a real eigenvalue μ to the represented point $\sigma_* + i\mu = \frac{1}{2} + i\mu$, where σ_* has already been fixed by (MP1).

Because a separate scalar operator is supplied for each γ , this family is not itself the one fixed operator required by the global Hilbert–Pólya interface. Its exact role is to verify the local kernel and the reality of an already supplied height.

The family is assembled, without changing any local proof, by the fixed ambient carrier operator $A_{p,r}^{(3)}$ above. Its basis is indexed by all geometric states $X = \gamma_{p,r}(y)$, independently of crossings, and $A_{p,r}^{(3)}\delta_X = y\delta_X$. A completed 3D crossing $e = (Z_e, h_e)$ selects the state $X_e = \gamma_{p,r}(\log Z_e)$ and Lean proves that its physical third coordinate is Z_e . Hence a “3D eigenvector” in the ambient resolvent statements below means this explicit carrier basis state, while `vonNeumannOp(log Ze)` computes its local one-dimensional resolvent factor.

The internal 3D chain runs entirely on these two instruments, and every link is unconditionally proven. Finite focal cancellation at ordinate $y = \log Z$ is detected by the finite harmonic Gram rank drop. Completed focal cancellation at physical height Z is exactly the analytic L -zero of that fibre and is detected by the completed harmonic Gram rank drop. The spatial statement `completedThreeDZeroAtHeight_twoGram3D_rankDrop` supplies both the completed harmonic rank drop and the von Neumann rank drop at the same ordinate $\log Z$. The carrier point $\sigma_* + i \log Z = \frac{1}{2} + i \log Z$ lies on the midline because the area law has already proved $\sigma_* = \frac{1}{2}$, while von Neumann reality supplies the real spectral ordinate and hence zero source drift (§21.5). The one-line Lean proof `simp [carrierPoint]` in `threeDZero_on_midline` is now backed by `carrierAbscissa_eq_half` and is a normal-form check, not the provenance of the half. Thus focal cancellation = 3D zero $\rightarrow$ harmonic Gram rank drop plus von Neumann Gram rank drop at $y = \log Z$: that is the detecting and realizing instrument of this part, and no step consults the chart.

The resolvent trace now has two event-typed versions (`RequestProject/TwinResolventTrace.lean`). For a finite $E \subset \mathcal{E}_\chi$, multiplicities m_e , and $p(Z) = \text{carrierPointAtHeight}(Z)$, Version I is

$$T_{\chi,E}^{(3)}(w) = \sum_{e \in E} m_e \text{vonNeumannResolventTrace}(\log Z_e, w) = \sum_{e \in E} \frac{m_e}{w - p(Z_e)}.$$

The compiled ambient bundle `threeDZero_ambientEigenvector_resolventTrace` states together that every index e is a completed 3D zero, that $\delta_e \neq 0$ is its eigenvector of $A_{p,r}^{(3)}$, and that the displayed carrier-point trace formula holds. This is precisely the “3D zeros plus 3D eigenvectors” version.

Version II reads the same certified events in the ordinate chart:

$$T_{\chi,E}^{(1 \leftrightarrow 3)}(u) = \sum_{e \in E} \frac{m_e}{\log Z_e - u}.$$

The ambient bundle `oneDZero_threeDZero_ambientEigenvector_resolventTrace` additionally proves, for every e ,

$$L(p(Z_e), \chi) = 0, \quad \text{CompletedThreeDZeroAtHeight } \chi \ Z_e, \quad A_\chi^{(3)}\delta_e = (\log Z_e)\delta_e.$$

The first equation is obtained from the completed focal residual by `completedThreeDEigenEvent_L_zero`; it is not inserted as a second unrelated zero assumption. Finally Lean proves the exact chart relation

$$T_{\chi,E}^{(3)}(\text{lineC}(u)) = i T_{\chi,E}^{(1 \leftrightarrow 3)}(u) \quad (\text{completedTraces_agree}).$$

The factor i is $d(\text{lineC})/du$, and the two pole coordinates are $p(Z_e) = \sigma_* + i \log Z_e$ and $\log Z_e$, respectively. The older abstract finite-set results `trace3D_capture`, `trace3D_analyticAt_off`, `trace1D_capture`, and `twinTraces_agree_on_window` remain in place; the new theorems identify the 3D zero and eigenvector data that their abstract ordinate set did not itself carry.

21.9 The chart-side registration audit

One further object appears in the evidence dossiers and deserves an exact label. Over a finite window $(0, T]$, the Lean development builds a diagonal matrix on the window's zero-events — `hpOperator` T , one axis per unit of multiplicity, eigenvalue the event ordinate. Its name is historical; its role is not spectral detection. It is a finite *repackaging of the already-located analytic zero window*, used for exactly one thing: auditing the chart against the carrier, count and multiplicity. The audit facts are proven and exact — the diagonal is Hermitian by type (`hpOperator_isHermitian`); every index is an exact vanishing (`eigenheight_is_exact_vanishing`); the chart reads zero at a height iff an index sits there (`chartZero_iff_eigenheight`); the diagonal's dimension is the chart's own count formula $1 + \theta(T)/\pi + S_{\text{mult}}(T)$ (`hpDimension_eq_registration`); the eigenspace dimension at an event equals the vanishing order — the mode space is a graded local algebra $\mathbb{C}[X]/((X - \text{line } \gamma)^{\text{ord}})$ (`finrank_modeSpace`), a double zero a two-dimensional crossing; and the window traces are honest analytic objects (`gradedResolventTrace_eq_residue_sum`, `windowedDiffResolvent_tendsto`). But because the window is indexed by already-located analytic zeros, nothing in this subsection participates in locating an event. The local von Neumann fibre of §21.8 likewise verifies an event after its height is supplied. The fixed global operator interface is stated separately below.

21.10 The Hilbert–Pólya setup we use

The classical suggestion runs: exhibit a self-adjoint operator whose spectrum gives the ordinates of the nontrivial zeros; self-adjointness forces the spectrum real; real ordinates place the zeros on the midline. It is worth saying plainly that this is not an exotic reformulation: Riemann's own sentence — *alle Wurzeln reell* — already asks for a real spectrum, and real spectrum is precisely what self-adjointness delivers. The Hilbert–Pólya suggestion is Riemann's phrasing transposed into operator language, which is why we treat it as the natural frame for the worked example rather than as one strategy among many.

The present development has three operator levels, and they must be kept separate. The proved scalar fibre `vonNeumannOp`(γ) = $\gamma \text{ id}_{\mathbb{C}}$ is a local detector attached after a real height γ is supplied. Its symmetry and kernel equation certify the coordinate of that already marked event; they do not by themselves provide one operator whose spectrum is the event set. The fixed $A_{\chi}^{(3)}$ supplies the next level: one symmetric operator on all completed 3D crossing states of χ , with explicit eigenvectors and the two resolvent traces above. Because its index is the completed-event subtype, it is a valid spectral realization of the 3D event ontology, not a discovery theorem for analytic zeros not yet coupled to that ontology. The final analytic-coverage interface is `grh_of_selfAdjoint_resolvent_capture`: one fixed self-adjoint element a must be accompanied by a resolvent-trace identification proving that every analytic pole parameter lies in `spectrum`(a). Self-adjointness then supplies spectral reality. The carrier and chart theorems below now provide the concrete candidate $A_{\chi}^{(3)}$ and its event traces; applying the final interface to every analytic zero still requires the coverage/trace identification rather than only the diagonal assembly.

22 The theorems

The formal home of this part is `RequestProject/HelixResolventCapture.lean` together with the supporting files named below; every statement carries axiom footprint `{propext, Classical.choice, Quot.sound}`, no `sorry`, no `axiom`, built in-tree (App. B). Below, `GRH`(χ) abbreviates the formal statement “every nontrivial zero of $L(s, \chi)$ lies on the midline” (`GRHSpectral.GRH`), stated for every modulus N and every Dirichlet character $\chi \bmod N$. The reading-hypotheses and both GRH implications are proven

at that generality. The trivial character is the instantiation that lands Mathlib’s RiemannHypothesis — the mod-1 L -function *is* riemannZeta — and a reading adopted at every modulus yields `GRHSpectral.GRHComplete`, the conjecture itself: every Dirichlet L -function, every modulus, every character, principal included.

Midpoint provenance and exclusion — proven before any zero-set coverage hypothesis:

- `windIntegerSite_radius_sq_tendsto` and `carrierScaleBalanced_iff`: arclength forces $r_n \asymp \sqrt{n}$, and $n^{-\sigma} r_n$ has a positive finite limit exactly at $\sigma = \frac{1}{2}$.
- `carrierAbscissa_scaleBalanced`, `carrierAbscissa_eq_half`, and `carrierPoint_eq_half_add_I`: the carrier point is constructed from the unique scale-balanced abscissa and only afterward rewritten in its half-unit normal form. The physical-height readout is separately defined as `carrierPointAtHeight(Z) = carrierPoint(log Z)`, and `carrierPointAtHeight_exp` proves the exact conversion between physical height and ordinate.
- `source_noDrift`, `SourceMode.noDrift`, and `Helix.no_radial_drift_iff_half`: norm conservation first gives $\operatorname{Re} \lambda = 0$ without mentioning a zero or midpoint; after the area-law normalization, zero radial drift is equivalent to $\sigma = \sigma_* = \frac{1}{2}$.
- `ThreeDZero = focalClosure` and `focalClosure_iff_rankDrop`: finite 3D vanishing is defined and detected without `carrierPoint`. At the completed-fibre level, `completedThreeDZeroAtHeight_iff_L_zero` identifies focal cancellation at Z exactly with $L(\operatorname{carrierPointAtHeight}(Z), \chi) = 0$.
- `not_scaleBalanced_of_ne`, `helix_no_offline_cancellation`, and `no_offCarrier_representation`: an off-midpoint abscissa is not a native scale-balanced state; an off-circle ledger cancellation is not weld-fixed; and no complex point with a different real coordinate is the image of any real carrier height. The exact image theorem is `exists_eq_carrierPoint_iff`; its positive physical-height form is `exists_pos_eq_carrierPointAtHeight_iff`, with exclusion theorem `no_offCarrier_physical_representation`.

The reduction spine — proven, unconditional:

- `RH_of_GRH_Trivial_Char`: $\operatorname{GRH}(\chi_1) \implies$ Mathlib’s RiemannHypothesis. The hypothesis handles the strip; Mathlib supplies no-zeros on $\operatorname{Re} s \geq 1$ and the negative-real trivial zeros via the completed functional equation. The entire target is one character’s GRH.
- `focalClosure_iff_rankDrop`: exact native finite focal cancellation is equivalent to harmonic Gram rank drop. This internal finite-carrier theorem contains no analytic zero set.
- `completedThreeDZeroAtHeight_iff_L_zero`: the completed 3D fibre crosses at physical height $Z > 0$ exactly when L vanishes at $\operatorname{carrierPointAtHeight}(Z) = \sigma_* + i \log Z$. This packages `focal_residual_zero_iff_L_zero` together with `reprPoint_eq_carrierPointAtHeight`. It is an exact pointwise identity at every represented height; it does not assert that an arbitrary analytic zero has a representing Z .
- `spatialLiftGram_det`, `spatialLiftGram_rankDrop_iff`, and `completedHarmonicGram3DAtHeight_rankDrop_iff_L_zero`: the completed harmonic 3D Gram operator rank-drops exactly at the represented L -zero. In addition, `completedThreeDZeroAtHeight_twoGram3D_rankDrop` states that the harmonic and von Neumann operators act on the three spatial carrier coordinates as $G \otimes I_3$; their determinants are $(\det G)^3$, so the lift preserves and reflects rank drop and cannot manufacture an extra cancellation.
- `vonNeumannOp_isSymmetric`: symmetry of the local scalar fibre operator is a theorem; it supplies coordinate reality after an event has been coupled to that fibre.
- `threeDCrossings_iff_grh_and_nativeCarrierCoverage`: the legacy identity premise is exactly the conjunction of the critical-line conclusion and midpoint-free native coverage.

2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120

2121
2122
2123

2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136

2137
2138
2139
2140
2141
2142

- `grh_of_resolvent_trace_3D_real`: for every modulus and character, $\text{THREEDREAL EVIDENCE} \wedge \text{ThreeD_crossings_are_real_zeros}(\chi) \implies \text{GRH}(\chi)$. The evidence component is formally redundant; `grh_of_threeDCrossings` obtains the conclusion directly from the equality embedded in the legacy premise.
- `grh_of_resolvent_trace_1D_correlation`: for every modulus and character, $\text{ONEDCORRELATION EVIDENCE}(\chi) \wedge \text{OneD_zeta_zero_correlated}(\chi) \implies \text{GRH}(\chi)$. The routing proposition is redundant and is itself proved by `oneD_zeta_zero_correlated_proved`; `grh_of_oneDCorrelationEvidence` shows that the chart-to-kernel premise alone supplies the conclusion.
- `grh_of_selfAdjoint_resolvent_capture`: the global operator route. It uses one fixed self-adjoint operator and a trace/capture premise placing every analytic pole parameter in its spectrum.
- `grhComplete_of_resolvent_trace_3D_real`, `grhComplete_of_resolvent_trace_1D_correlation`: a reading adopted at every modulus yields `GRHComplete` — the full conjecture, principal characters included.
- `RH_of_resolvent_trace_3D_real`, `RH_of_resolvent_trace_1D_correlation`: the trivial-character instantiations composed with the bridge, landing `RiemannHypothesis` directly.
- `threeDRealEvidence`, `oneDChartEvidence`: the two dossiers inhabited — the 16-fold and 18-fold conjunctions of §24 and §23 certified theorem by theorem.
- `threeD_real_case`, `oneD_correlation_case`: each reading packaged as one auditable object — its proven evidence paired with its displayed conditional premise. The minimal dependencies are the audit theorems listed above.

The two legacy interfaces, rendered formal. The formal system speaks the classical frame: its “nontrivial zero” is definitionally the analytic object. The revised source records the minimal dependency of each older interface:

- `ThreeD_crossings_are_real_zeros` (3D-ZERO-REAL; the strong reading with the 3D zero primary): every nontrivial zero ρ is represented by a completed fibre crossing at a physical 3D height $Z > 0$, with $\rho = \text{carrierPointAtHeight}(Z) = \sigma_* + i \log Z$. The two spatially lifted Gram pencils then provide their rank drops at ordinate $\log Z$. Here $\frac{1}{2} + i \log Z$ is the proved normal form of the carrier image, not a fresh geometric assumption. Nevertheless, the equality already includes the critical-line conclusion. The theorem `threeDCrossings_iff_grh_and_nativeCarrierCoverage` exposes its exact content; the midpoint-free coverage component is now available as the separate target `NativeCarrierCoverage`.
- `OneD_zeta_zero_correlated` (1D-ZERO-EXCLUSIVE-REAL; the weak reading): a per-zero disjunction. Its right branch is only the analytic zero equation together with $\gamma = \Im \rho$, so `oneD_zeta_zero_correlated_proved` inhabits it unconditionally. The actual pointwise coupling is `OneDCorrelationEvidence`, which asserts a scalar-fibre kernel at that analytic zero and therefore implies the critical-line conclusion by itself.

These audits do not alter either conditional theorem. They state which premise actually supplies the conclusion and prevent the geometric midpoint theorems, the internal rank-drop correspondence, and the analytic zero-set identification from being presented as one result.

The coordinate convention in both interfaces is fixed: $Z > 0$ is physical height on the Archimedean helix, $y = \log Z$ is the analytic ordinate, and $S(t)$ is the global carrier-to-chart coordinate/registration map. The pointwise fibre readout and the global census map therefore have distinct typed roles.

23 Evidence I: the 3D–1D correlation

The eighteen facts of `oneDChartEvidence`, each proven unconditionally, form a chart-side registration dossier. They establish count, jump, residue, multiplicity, and projection identities on the objects appearing in their statements. The universal layer is the $S(t)$ global coordinate map: it transports the carrier-scale census to the unit-chart census and registers every analytic event jump. A global census identity is not by itself a pointwise bijection between `ThreeDZero` and arbitrary Dirichlet L -zeros. The separate, character-indexed input `ONEDCORRELATIONEVIDENCE( $\chi$ )` is the operative pointwise chart-to-kernel leg. Three of the facts below are *named* engines: *Shannon* — the collapse loses radius and angle to the ledger but never counterfeits (`record_bijective`); the *Shannon Cascade* — an accumulation of stage features is a feature of the limit (`limit_zero_of_stage_accumulation`); and *Riemann’s Fold* — the projection pipeline, crossing dimensions, lands on the real axis exactly when its retained coordinate does (`pipeline_midpoint_iff`).

- `chartZero_iff_eigenheight` (`RequestProject/HilbertPolya.lean`). For $0 < \gamma \leq T$: $\zeta(\frac{1}{2} + i\gamma) = 0 \iff \gamma$ is an index of the zero-window diagonal (chart-side audit, §21.9). The one-for-one coincidence, proven — not asserted — window by window.
- `hpDimension_eq_registration` (`RequestProject/HilbertPolya.lean`). The window diagonal’s dimension is the chart’s own count formula, $1 + \theta(T)/\pi + S_{\text{mult}}(T)$. The index count and the zero count are the same number because the ledger says so.
- `carrier_scale_compensation_S` (`RequestProject/CarrierScaleCompensation.lean`). $N_{\pi/3}(e^t) - N_1(e^t) = S(t)$: the classical $S(t)$ is the gap between the native-scale event count and the unit-scale smooth term — one procession, two registrations, no mysterious remainder.
- `native_identification'` (`RequestProject/CarrierScaleCompensation.lean`). $N_{\pi/3}(e^t)$ equals the number of on-line zero ordinates in $(0, t]$ — one zero per completed $\pi/3$ cell.
- `S_jump_detects_event` (`RequestProject/ResidueJump.lean`). S has a unit jump at γ iff $\zeta(\frac{1}{2} + i\gamma) = 0$: detecting a zero and detecting an S -jump are the same act.
- `count_hasJump_iff_S_hasJump` (`RequestProject/ResidueJump.lean`). The event count and S have identical jumps — their difference is continuous, so *all* discontinuity lives in S , none in the smooth clock θ/π .
- `residue_eq_Smult_jump` (`RequestProject/ResidueJump.lean`). At every event, log-derivative residue = vanishing order = ledger jump, with multiplicity and with *no* simplicity hypothesis.
- `record_bijective` (`RequestProject/RoundTrip.lean`). `record  $f$  = (height, (radial, phase))` is a bijection with explicit inverse `reconstruct`: the loss ledger made literal (§21.4) — *Shannon*: the collapse never counterfeits.
- `radial_lost` (`RequestProject/RoundTrip.lean`). Without the ledger the geometric projection is provably non-injective — the ledger does real work; the faithfulness results are not vacuous.
- `projection_bijective_loss_ledger` (`RequestProject/SourceHolonomy.lean`). The chart chain $y \mapsto 1 - (\frac{1}{2} + yi)^{-1}$ is injective: distinct helix events write distinct ledger entries — the data-processing inequality made literal.
- `pipeline_midpoint_iff` (`RequestProject/RoundTrip.lean`). The end-to-end $3D \rightarrow \text{height} \rightarrow \text{encode} \rightarrow 1D$ pipeline reads $\frac{1}{2}$ exactly when the retained 3D coordinate is $\frac{1}{2}$: the midline address survives projection intact, and only the midpoint maps there — *Riemann’s Fold*.
- `no_radial_drift_on_helix` (`RequestProject/SourceHolonomy.lean`). The Euler-factor phasor’s radius depends only on $\text{Re } s = \frac{1}{2}$, not on height: the height $\rightarrow$ circle map is a faithful re-coordinatization, all distinguishing information in the phase the ledger keeps.

- 2187 • `midpoint_entry_on_circle` (RequestProject/SourceHolonomy.lean). Every helix height charts
2188 to a unit-circle ledger entry ($\|1 - (\frac{1}{2} + yi)^{-1}\| = 1$): line-events are recorded as circle-points.
- 2189 • `faithful_projection_zeta` (RequestProject/Faithfulness.lean). The projected fiber is real
2190 on the midline and vanishes exactly at the genuine on-line zeros: the collapse neither hides a
2191 zero nor manufactures one.
- 2192 • `euler_maclaurin_dirichlet` (RequestProject/EulerMaclaurinDirichlet.lean). In the strip,
2193 the finite Dirichlet sum with boundary correction approximates ζ with error uniform in the
2194 truncation — the quantitative backbone of the chart readout.
- 2195 • `transfer_tendsto` (RequestProject/TransferContinuation.lean). A polynomial bound on
2196 the primitive continues the series past the trivial abscissa: the readout limit exists from a growth
2197 estimate alone, with no appeal to zero locations.
- 2198 • `limit_zero_of_stage_accumulation` (RequestProject/ExportAdapter.lean). A zero accu-
2199 mulating across the finite stages is a zero of the limit ξ -section: the collapse does not counterfeit
2200 features — the *Shannon Cascade* converse.
- 2201 • `windowedDiffResolvent_tendsto` (RequestProject/DifferencedResolvent.lean). The win-
2202 dowed differenced-resolvent trace converges as $T \rightarrow \infty$ to an absolutely convergent limit: the
2203 chart-side resolvent is an honest analytic object, unconditionally.

2204 24 Evidence II: internal three-dimensional carrier facts

2205 The sixteen facts of `threeDRealEvidence` are each proven unconditionally, but their conjunction is
2206 an internal-support dossier, not a theorem identifying every analytic zero with a `ThreeDZero`. The
2207 facts concern local scalar-fibre symmetry, finite carrier cancellation, completed critical-line readouts,
2208 chart-indexed registration objects, arithmetic winding, and general limit principles. Their scopes are
2209 recorded individually below. Three named engines organize the dossier: *Feynman's Quiver* — the
2210 crossings are prime-generated by unique factorization, the phasor bank is summable, the heights
2211 are distinct, and no focus splits; *Deligne's Pairs* — the self-reciprocal det-one local pairing, purity at
2212 the finite places; and *Shannon Projection Dominance* — the source dominates every projection, so the
2213 limit holds no feature the stages lack.

- 2214 • `vonNeumannOp_isSymmetric` (RequestProject/ClosedForm.lean). The fibre operator $\gamma \cdot \text{id}$ is
2215 symmetric, hence self-adjoint with real spectrum $\{\gamma\}$. This is a local coordinate detector indexed
2216 by an already supplied real γ , not the fixed global operator of `grh_of_selfAdjoint_resolvent_capture`.
- 2217 • `carrierThreeDOperator_isSymmetric`, `carrierThreeDOperator_eigenvector`, and `carrierThreeDOperator`
2218 (RequestProject/ThreeDFocalEvent.lean). For fixed nondegenerate geometric carrier param-
2219 eters (p, r) with $p \neq 0$, one real-diagonal operator acts on all finitely supported carrier states
2220 $X = \gamma_{p,r}(y)$; its type mentions no event predicate. Every geometric state supplies a nonzero basis
2221 eigenvector with eigenvalue y , and a completed crossing at Z_e is proved to select the state with
2222 $y = \log Z_e$ and third coordinate Z_e .
- 2223 • `threeDZero_ambientEigenvector_resolventTrace` and `oneDZero_threeDZero_ambientEigenvector_resolv`
2224 (RequestProject/TwinResolventTrace.lean). The first bundle contains 3D crossing certificates,
2225 their fixed-operator eigenvectors, and the 3D pole formula. The second adds the analytic L -zero
2226 readout for each same event, the ordinate trace, and the exact carrier–ordinate chart relation.
- 2227 • `eigenheight_is_exact_vanishing` (RequestProject/HilbertPolya.lean). Every index of the
2228 chart-side window satisfies $\zeta(\frac{1}{2} + i\gamma) = 0$ exactly: the audit's indices are genuine events, not
2229 approximations (§21.9).

- 2230 • `hpOperator_isHermitian` (RequestProject/HilbertPolya.lean). The zero-window registra-
2231 tion diagonal is Hermitian — reality by type (chart-side audit, §21.9).
- 2232 • `finrank_modeSpace` (RequestProject/GradedModeDictionary.lean). The mode space at an
2233 event is a graded local algebra of dimension exactly the vanishing order: a double zero is a
2234 two-dimensional crossing; spectral multiplicity = analytic order.
- 2235 • `gradedResolventTrace_eq_residue_sum` (RequestProject/GradedModeDictionary.lean). The
2236 3D-mode resolvent trace equals the explicit-formula sum of log-derivative residues: the trace
2237 formulas coincide, not just the locations.
- 2238 • `focal_residual_zero_iff_L_zero` (RequestProject/FocalResidualVanishes.lean). $D_{\text{cell}} =$
2239 $V_\chi \Phi_\chi$ with $V_\chi \neq 0$ and $\Phi_\chi = (\pi/3)L$: the cell channel vanishes iff the L -value does, residue-free,
2240 at exact $\pi/3$ amplitude at the represented critical-line point. This completed-cell statement is
2241 distinct from the finite-bank definition of `ThreeDZero`.
- 2242 • `fiber_accumulates_to_L` (RequestProject/Faithfulness.lean). For $\chi \neq 1$ and $\text{Re } s > 0$ the
2243 partial sums converge to $L(s, \chi)$ across the whole strip. This is value convergence at each stated
2244 s ; transferring zero sets requires the separate uniform-convergence/isolation hypotheses used by
2245 the limit theorems.
- 2246 • `threeD_exhaustive` (RequestProject/SourceHolonomy.lean). Every zero event on the real
2247 height ray is sourced — automatic because reals are conjugation-fixed. Its domain is the
2248 real-height event space; it does not prove that every complex analytic zero lies in that domain.
- 2249 • `threeD_metric_no_zeros` (RequestProject/SourceHolonomy.lean). The 3D cup form is defi-
2250 nite: no nonzero state is null, so vanishing is a property of *readings*, never of states — a zero
2251 cannot hide as a degenerate state.
- 2252 • `windFromPrimes_mul` (RequestProject/Origination.lean). The winding is multiplicative over
2253 factorization, with no Relog: the fundamental theorem of arithmetic realized as a homomorphism
2254 into the circle — carrier and L -function are two prime-built readings of one arithmetic (*Feynman's*
2255 *Quiver*, generation).
- 2256 • `heights_distinct` (RequestProject/NoDoubleCancellation.lean). Distinct sites carry dis-
2257 tinct eigen-heights: unique factorization as a spectral non-degeneracy (*Feynman's Quiver*, unique-
2258 ness).
- 2259 • `cancellation_modes_subsingleton` (RequestProject/NoDoubleCancellation.lean). At any
2260 spectral parameter, at most one basis mode cancels: no split focus — a crossing marks one
2261 ordinate, not a superposition (*Feynman's Quiver*, no split focus).
- 2262 • `localPoly_reciprocal` (RequestProject/FiniteWeightFiber.lean). The local numerator is
2263 self-reciprocal: the per-place functional equation whose global assembly has fixed axis $\text{Re } s = \frac{1}{2}$
2264 — the midline the crossings sit on (*Deligne's Pairs*).
- 2265 • `fiber_det_one` (RequestProject/FiniteWeightFiber.lean). $\prod_i \lambda_i = 1$ under the duality
2266 involution: the modulus ledger the reciprocal law consumes (*Deligne's Pairs*).
- 2267 • `frobeniusBlock_det_one` (RequestProject/FrobeniusSimilitude.lean). The chiral block
2268 $\text{diag}(z, \bar{z})$ has determinant 1: the Frobenius similitude is a pure rotation — Satake weights stay
2269 on the unit circle, eigenphases stay real (*Deligne's Pairs*).
- 2270 • `limit_dominance` (RequestProject/LimitDominance.lean). Hurwitz-type dominance: the
2271 under local uniform convergence and isolation, a limit zero has nearby stage zeros eventually.
2272 This gives proximity, not equality with a fixed finite stage or a proof that the nearby zeros lie on
2273 the carrier image.

25 Proof A: GRH under (Z-3D)

Theorem 49 (The identity route). *Unconditionally:*

- (i) *THREEDREAL EVIDENCE holds (threeDRealEvidence);*
- (ii) *for every modulus N and character $\chi \bmod N$,*

$$\text{THREEDREAL EVIDENCE} \wedge \text{ThreeD_crossings_are_real_zeros}(\chi) \implies \text{GRH}(\chi).$$

Lean: grh_of_resolvent_trace_3D_real;

- (iii) *the reading adopted at every modulus yields the full conjecture GRHComplete.*

Lean: grhComplete_of_resolvent_trace_3D_real;

- (iv) *at the trivial character the premises imply RiemannHypothesis.*

Lean: RH_of_resolvent_trace_3D_real.

Proof A. Adopt (Z-3D): “nontrivial zero” names the crossing — the focal-alignment event of §21.7. Two logically distinct facts are then composed. First, before this adoption, the arclength area law and conservative transport have proved that the native, scale-balanced carrier image is exactly $\{\sigma_* + i \log Z : Z > 0\}$, equivalently $\{\sigma_* + iy : y \in \mathbb{R}\}$, with $\sigma_* = \frac{1}{2}$, and that its ledger cannot encode an off-image cancellation (§21.5). Second, (Z-3D) supplies the coverage/identity clause that each classical nontrivial zero is one of those native focal events. This second clause is the explicitly displayed hypothesis in (ii); it is not used in the derivation of σ_* .

For each covered event at physical height Z , the two spatial Gram theorems provide the harmonic and von Neumann rank drops at the same ordinate $y = \log Z$. The fibre operator is self-adjoint by theorem (vonNeumannOp_isSymmetric), and the represented coordinate is the already derived $\sigma_* + i \log Z$. Hence taking real parts gives $\text{Re } \rho = \sigma_* = \frac{1}{2}$. With the evidence of §24 — exact vanishing at the eigenheights, multiplicity equal to order, no split focus, definite metric, exhaustion of the real ray — theorem (ii) closes $\text{GRH}(\chi)$ under the stated coverage hypothesis; (iii) and (iv) are its all-modulus and trivial-character specializations.

Equivalently, the newly assembled operator records the same step without a family-only ambiguity: the covered crossing is an index $e \in \mathcal{E}_\chi$, its explicit state δ_e is an eigenvector of $A_\chi^{(3)}$ with real eigenvalue $\log Z$, and Version I of the resolvent trace has its pole at the carrier point of that same event. This augments the retained proof; it does not replace its displayed coverage hypothesis. $\square$

Formal dependency audit for Proof A. The proof text above is retained. The compiler-level implication is sharper: the first evidence component is unused, because the equality $\rho = \text{carrierPointAtHeight}(Z)$ inside the second component already yields the real-part conclusion. The theorem `threeDCrossings_iff_grh_and_nativeCarrierCoverage` records this exactly. Thus Proof A is a valid conditional deduction, while its legacy identity premise must not be advertised as an independently proved analytic-zero coverage theorem. The geometric result established before that premise is the location of the native carrier image.

26 Proof B: GRH under (HP-W)

Theorem 50 (The correlation route). *Unconditionally:*

- (i) *ONEDCHART EVIDENCE holds (oneDChartEvidence);*

2311 (ii) for every modulus N and character $\chi \bmod N$,

$$\text{ONEDCORRELATIONEVIDENCE}(\chi) \wedge \text{OneD_zeta_zero_correlated}(\chi) \implies \text{GRH}(\chi).$$

2312 *Lean: grh_of_resolvant_trace_1D_correlation;*

2313 (iii) the reading adopted at every modulus yields the full conjecture *GRHComplete*.

2314 *Lean: grhComplete_of_resolvant_trace_1D_correlation;*

2315 (iv) at the trivial character the premises imply *RiemannHypothesis*.

2316 *Lean: RH_of_resolvant_trace_1D_correlation.*

2317 *Proof B.* Adopt (HP-W): the criterion demands that the operator's eigenvalues *coincide with* the zeros. The hypothesis in (ii) routes each zero through one of two forms: a crossing at a positive physical 3D height, or a 1D chart report at its ordinate. In the first form the crossing is a completed focal event at physical height $Z > 0$, read at ordinate $y = \log Z$. In the latter form, $\text{ONEDCORRELATIONEVIDENCE}(\chi)$ is a proof input: it turns that chart report into the nonzero von Neumann kernel at the reported point. Section 23 supplies the compiled chart-side dossier — the window bijection, registration identity, jump-for-residue agreement with multiplicity, and projection ledger. Both branches then close by the same von Neumann reality (the case split inside *grh_of_resolvant_trace_1D_correlation*), whence $\text{GRH}(\chi)$ at every modulus, (iii) assembles the full conjecture, and (iv) lands the classical statement. The weak reading consumes nothing stronger than the strong one — the same self-adjoint mechanism closes both branches.

2328 For a chart report that is coupled to a completed crossing, Version II makes all three records simultaneous in Lean: the 1D equation $L(p(Z_e), \chi) = 0$, the completed 3D event certificate, and 2329 the 3D eigenvector δ_e of $A_\chi^{(3)}$. The identity $T^{(3)}(\text{lineC}(u)) = iT^{(1 \leftrightarrow 3)}(u)$ is the compiled correlation 2330 between those records. The pointwise premise in the legacy proof still performs the routing of an 2331 arbitrary chart report into this event space. $\square$

2333 **Formal dependency audit for Proof B.** The proof text above is retained. The source now proves 2334 *oneD_zeta_zero_correlated_proved*: the disjunctive routing proposition is automatic because its 2335 right branch records only $L(\rho, \chi) = 0$ and chooses $\gamma = \Im \rho$. Conversely, *grh_of_oneDCorrelationEvidence* 2336 proves that the operative evidence alone gives the critical-line conclusion, since it asserts a scalar- 2337 fibre kernel at each analytic zero. The global $S(t)$ coordinate law explains the count, jump, and 2338 multiplicity registration used by this route; the pointwise chart-zero-to-kernel assertion remains its 2339 separate premise.

2340 27 Riemann Hypothesis corollaries at $\chi = 1$

2341 The mod-1 L -function is *riemannZeta*, so at the trivial character each conditional implication lands 2342 the classical statement with the displayed premise unchanged:

2343 **Corollary 51** (RH, identity reading). *THREEDREALEVIDENCE* $\wedge$ *ThreeD_crossings_are_real_zeros*(χ_1) 2344 *implies RiemannHypothesis* (*RH_of_resolvant_trace_3D_real*, proven). By the audit above, the iden- 2345 tity premise itself contains the critical-line conclusion as well as native coverage.

2346 **Corollary 52** (RH, correlation reading). *ONEDCORRELATIONEVIDENCE*(χ_1) $\wedge$ *OneD_zeta_zero_correlated*(χ_1) 2347 *implies RiemannHypothesis* (*RH_of_resolvant_trace_1D_correlation*, proven). The operative 2348 *ONEDCORRELATIONEVIDENCE*(χ_1) bridge alone is the pointwise premise that supplies the critical-line conclu- 2349 sion.

2350 Both factor through the same bridge, *RH_of_GRH_Trivial_Char*: one character's GRH is RH.

28 Conclusion

The corrected dependency structure has three levels. First, Archimedean helix geometry derives the unique scale-balanced, drift-free carrier abscissa; Lean constructs `carrierPoint` from that abscissa, proves its half-unit normal form afterward, and reads physical height through `carrierPointAtHeight(Z) = carrierPoint(log Z)`. Second, the finite harmonic pencil detects truncated focal cancellation exactly, the completed harmonic pencil detects the completed L -fibre cancellation, and the local scalar fibre verifies its ordinate $y = \log Z$. The fixed operator $A_\chi^{(3)}$ then assembles all completed crossings into one symmetric event-state space with explicit basis eigenvectors. Third, identifying the analytic zero set with those native events requires a pointwise coverage or spectral-capture theorem.

The global $S(t)$ coordinate map supplies exact census, jump, residue, and multiplicity registration between carrier and chart scales. The proved supporting facts include the two explicit completed-event resolvent versions: 3D zeros with 3D eigenvectors, and 1D zero readouts with the same 3D zeros and 3D eigenvectors. The legacy identity conditional and correlation conditional also remain valid deductions, with their proofs retained above; the new Lean audit states exactly which premise contains the critical-line conclusion in each route. An analytic Hilbert–Pólya completion can now target the fixed $A_\chi^{(3)}$: it must prove the analytic resolvent/spectral capture required by `grh_of_selfAdjoint_resolvent_capture`, or instantiate that interface with another fixed operator. The scalar family $\gamma \text{id}_\mathbb{C}$ is correctly used as the local diagonal-mode calculator, while $A_\chi^{(3)}$ is the assembled 3D event-space operator.

Part IV

Beyond Endoscopy - Altuğ Analysis and Repair

29 The obstruction: Altuğ’s uniformity problem

The framework began not as an alternative to functoriality but as a root-cause analysis of one failure: the loss of productivity in the direct Beyond-Endoscopy route [1] above the symmetric square. The obstruction turned out to be an emergent clock; once that clock is separated the higher-order analysis becomes compositional, and the transport interface the rest of the paper builds is extracted from the structure that composition consumes.

Altuğ’s execution of Beyond Endoscopy for $GL(2)$ [2, 3, 4] reduces the standard-representation trace to an archimedean analysis whose primary difficulty is a *uniformity problem*: the asymptotic expansions of the Fourier transforms of the orbital integrals must hold with implied constants independent of every parameter — the C, D -independence Altuğ calls “the central issue,” the content of his long technical appendix (Altuğ II, Appendix A), which secures the power saving $\sum_{n < X} \text{tr } T_k(n) = O(X^{31/32+\epsilon})$ isolating the standard representation (Altuğ III, Thm 1.1). A second difficulty sits beyond it: Poisson summation on the n -sum “works well for the standard representation and the symmetric square, however stops being productive for higher symmetric powers” (Altuğ III, fn. 5, after Sarnak [7]; the two productive cases are Venkatesh’s [5]). To our knowledge no one had instrumented these sums numerically; the field’s own diagnosis is that the direct analysis “does not scale to higher rank.” Our first reading mis-located the difficulty at the cutoff’s edges — it is not there.

The estimate that costs Altuğ roughly sixty pages reduces, in the exact gauge, to a single magnitude

bound. The transform (11) whose C, D -uniformity is Altuğ’s “central issue” is bounded, with no oscillation estimate, by the triangle inequality in the coordinate that makes the integrand real (Lemma 83); the sharp expansion (Theorem 84) and the full uniform bound (Proposition 86) then follow, every uniform constant being one application of that one bound. The analysis is deferred to Appendix A; here we turn to what made it look hard.

30 The oscillation is the orbital transform’s own comb

Why did Prop. 86 look hard? The oscillation the estimate must survive is deterministic. The numerical scan is made in the clock coordinate

$$\nu_{\text{clock}} := 2\nu_J, \quad e(-y\nu_J/2lf^2) = e(-y\nu_{\text{clock}}/4lf^2),$$

where ν_J is the transform frequency in Prop. 86. The measured clock is extracted from the full complex profile $V(\nu_{\text{clock}})$ by fitting the log-power modulation $\log |V(\nu_{\text{clock}})|^2$ to a smooth envelope plus the first two harmonics sharing one fundamental, not by averaging dip gaps. Over the cutoff support $y \in [\frac{X}{4}, \frac{5X}{4}]$ this gives

$$\Delta\nu_{\text{clock}} = 0.524 (lf^2)^{1.03} (X/8)^{-1.04} (1 + \xi)^{0.00}, \quad R^2 = 1.000,$$

equivalently half that spacing in the ν_J coordinate. In the one-parameter window-span normalization $\kappa_{\text{eff}} := 4lf^2/(X\Delta\nu_{\text{clock}})$ the cells give $\kappa_{\text{eff}} = 0.943 \pm 0.025$ (range 0.895–0.971), while the harmonic-truncation audit has median period shift 2.4×10^{-4} , so the residual spread is not a dip-localization artifact. With the same Chebyshev multiplier $U_r(x)$ used for the symmetric-power rank, the event-bearing $\xi \neq 0$ cells keep the same law through the tested range $0 \leq r \leq 13$ after separating the continuous clock from deep-event visibility: every rank has $R^2 = 1.000$, lf^2 -exponent 1.03–1.04, and X -exponent -1.04 – -1.03 on the cells passing the clock-quality gate. Thus the older scalar $\kappa \approx 0.94$ is the mean of a slightly drifting clock, and the observed high-rank loss is productivity decay: deep-event cells thin from 10/12 at $r = 0$ to 3/9 and 3/8 at $r = 11, 12$, while the surviving clock cells still obey the same law. Thus the scalable object is not the raw list of deep dips but the cell-indexed phase clock together with a separate productivity ledger: the clock law survives in cells where the visible deep comb has already thinned, so higher rank must be tested by clock-cell census and productive cell yield as distinct observables.

This oscillation is *intrinsic to the orbital transform* $\Psi = I_{l,f}(\xi, y)$, not the cutoff’s edges. Two controls settle it: replacing the compact cutoff $G(y/X)$ by a smooth edgeless window leaves the comb unchanged, and tapering the $x = \pm 1$ endpoints of the x -integral leaves it unchanged as well (`height_warp_test.py`) — so neither the height-cutoff edges nor the integral endpoints produce it. It is the ξ -Fourier oscillation of the orbital content: at $\xi = 0$ the x -integrand is unimodal ($P_2/P_1 = 0$) with *no* comb, while $\xi \neq 0$ switches on the transform’s own oscillation, the window fixing only its ν -spacing ($\Delta\nu \propto 1/X$). It is therefore a deterministic feature of the content, not a removable chart artifact of the cutoff; but it is *deterministic*, which is exactly what the exact gauge of §A discharges — the magnitude bound survives it whatever its source, with no estimate of it.

The clock is necessary but not sufficient: the repair is the clock configurator plus two adapters. Identifying the transport frequency is the first ingredient, not the fix — the clock alone does not close the crossings. Exact closure of the carrier readout, across the symmetric-power ladder and up the GL tower, is the *clock configurator* — which reads the deterministic transport frequency and sets the period, phase, and scale the rest inherit — *together with* two carrier adapters it keys. The *carrier dwell* re-registers the values a projected sum skips — the native, exact form of Altuğ’s $\Sigma()$

missing-lattice-point correction — and closes the crossing exactly where the skip displaces it. The *phasor delay adapter*, derived from the clock, retards the faster-spinning higher-weight channels so the crossing sharpens rather than smears, restoring the reopening exponent toward its simple-zero value. The clock configurator supplies the period; these two supply registration and spin. The repair is the clock configurator + carrier dwell + phasor delay, not the clock alone; the suite is catalogued as an operational unit in §40.

31 The repair: a lossless method that scales

The method the rest of the paper builds — a clock, then carrier *scaling*, a frozen *warp*, and ledgered *projection* — has a clean, precision-testable form on the carrier’s own *bank*, the object of the cell-closure algebra of §38 (CellClosure). We report that worked example here: on the bank, the four operations drive the native cell residual to a *precision-limited* zero, uniformly across the symmetric-power rank census, the residual tracking working precision — an identity, not a model floor.

Scope (read this first). This experiment is run on the *carrier bank* $B_r(k) = \sum_j e^{i(r-2j)\theta k}$ with the *exact* Satake clock; it is *not* Altuğ’s Poisson-summed orbital n -sum of §30, and it does not reproduce that machinery (the approximate functional equation, the trace Poisson summation, the orbital transform $\Psi(y) = I_{l,f}(\xi, y)$, or the Proposition 5.2 object). Its purpose is narrow and honest: to isolate and stress-test the *closure mechanism* at controlled high precision, with each operation ablated. The *bridge* — that the *measured* v -clock of Altuğ’s transform (§30) is the clock that generates this warp and closes the *actual* transform’s cell — is asserted structurally and *not* demonstrated here; building it (the measured clock feeding the carrier warp) is the explicit open experiment. Falsifiability register: a measured Altuğ clock that failed to generate the closing warp would disconfirm this unification, and would be reported as prominently as a success.

The object. For a fiber of Satake angle θ , the Sym^r carrier bank at carrier step k is $B_r(k) = \sum_{j=0}^r e^{i(r-2j)\theta k}$, with channel frequencies $\phi_{r,j} = (r - 2j)\theta$. The zero-frequency channel ($\phi_{r,j} \equiv 0 \bmod 2\pi$: the middle channel of Sym^{2m} , and any further trivial constituent) is the pole/trivial channel, booked separately in a ledger; cell closure is the vanishing of the non-DC part over a cell C of the carrier’s native period,

$$Q_C = \sum_{k \in C} \sum_{j: \phi_{r,j} \neq 0} e^{i\phi_{r,j}k} \stackrel{?}{=} 0.$$

Four stages, each adding one operation (residual = $|Q_C|$): **(i) clock** — remove the dominant linear transport $e^{-i\kappa k}$ (κ the largest- $|\phi|$ channel); scalar, still aliased. **(ii) scale** — move to the compatible μ_M cell, M drawn from the root-of-unity family and fixed, from the channel geometry *before* residuals are seen, fine enough that no non-DC channel snaps onto DC — the ledgered projection that prevents higher-rank aliasing. **(iii) warp** — remove the frozen within-cell defect $\omega_j(k) = e^{-i(\phi_{r,j} - \hat{\phi}_{r,j})k}$, where $\hat{\phi}_{r,j}$ is the nearest $\frac{2\pi}{M}$ grid frequency: a *unit-modulus*, one-slope-per-channel law read off the Satake angle, *not* a pointwise phase negation (which would give $\sum_k 1 = M$, not 0). **(iv) projection** — retain every channel separately, so the warp acts per channel; the collapsed scalar readout cannot substitute. After the warp each channel is a root-of-unity frequency and closes by the exact geometric-series identity $\sum_{k \in \mu_M} e^{i\hat{\phi}k} = 0$ (as $e^{i\hat{\phi}M} = 1$); the closure is an *identity*, so its residual tracks the working precision.

Result [measured; identity] (clock_scale_warp.py, mpmath), worst case over held-out cells $t \in \{0, 6, 12, 30, 100\}$ and ranks $r \in \{1, \dots, 13\}$, at 80 digits:

fiber	raw	clock	+scale	+warp (vector)	control (scalar)
harmonic $\theta = \pi/3$	10^{-p}	$O(30)$	10^{-p}	10^{-p}	10^{-p}
cuspidal Δ at $p=2$	13.9	15.5	15.0	8.8×10^{-79}	12.7
cuspidal $\cos \theta = \frac{3}{10}$	12.7	14.2	63.4	1.5×10^{-77}	92.2

The cuspidal +warp residual is $\sim 10^{-29}, 10^{-48}, 10^{-79}$ at 30, 50, 80 digits: it *tracks precision*, where a model floor would be digit-independent. Every ablation floors — the scalar readout (raw / clock), scaling without the warp, and the no-projection scalar warp all plateau at $O(1)$ – $O(100)$ — so *both* the projection and the warp are load-bearing.

What it is, and is not. The harmonic fiber (θ a rational multiple of π) is already root-of-unity and closes at every stage *except* the scalar clock column, where collapsing to a single dominant frequency aliases even it to $O(30)$; +scale re-closes it (the 10^{-p} entries). There the DC ledger correctly books the genuine Sym^r *reducibility* (three trivial constituents at $r = 8, 10$ for $\theta = \pi/3$) — the same trivial channels that return as the exceptional cases of Part III. The content is the *cuspidal* row: the irrational Satake phase does not close raw (the +scale column floors), but the frozen unit-modulus warp harmonizes it to the compatible cell, where it closes *exactly*. This is *not* a claim that the cuspidal L -function’s own coefficients sum to zero; it is the exact, lossless, precision-tracking closure of the harmonized fiber — “the fiber engages its own warp” (§40) made an identity, uniform across rank because the carrier scale M is refined ($6 \rightarrow 12 \rightarrow 48 \rightarrow 96$) to keep every channel resolved. This is the shape of the whole paper: Part II builds the machinery this example runs on (the carrier, the loss ledger, the warp dictionary), and Part III lands the classical functoriality; the reducibility booked in the ledger here is the same object that becomes the exceptional locus there.

Fixed results: the carrier closes every object to precision; the degradation lives only in the 1D projection. The repair of §30 is measured. Two readings of the same object must be separated: the *carrier* (its vector representation, the four-stage above) and the *scalar L -function readout* (the 1D projection). In the carrier, every object — including genuine multi-parameter ones — closes to a precision-limited zero, tracking working precision; the degradation is a feature of the projection, not of the object:

object (built from scratch)	scalar 1D readout	carrier (vector four-stage)
std-rep $\xi \neq 0$ channel (Altug dual)	$O(X)$ bound: 0.5	$ S /M = 0, S = 1.6 \times 10^{-14}$ exact
four-stage carrier cell	—	8.8×10^{-79} (80 dig.) identity
$\text{Sym}^{13} \Delta$ (GL 14), reopening k	0.44 (smeared)	10^{-p} (identity) carrier exact
self-dual $\text{GL}(3) = \text{Sym}^2(\text{Maass } R=13.78)$ $k = 1.01$ (1-param, clean)	$k = 1.01$ (1-param, clean)	4.2×10^{-91} (90 dig.) both exact
genuine $\text{GL}(4) = L(\Delta \times E_{11})$	$k = 0.47$ (smeared)	4.9×10^{-91} (90 dig.) carrier exact
non-self-dual $\text{GL}(3) = L(\rho_{F_{21}})$	$\text{Re } \eta = \text{Re } \eta'$ (blind)	4.3×10^{-62} (60 dig.) Lean-backed
carrier crossing, values skipped	cut: 1.10	dwel: 3.2×10^{-4} closes

The standard-representation channel Altug can only *bound* to $o(X)$ (magnitude ~ 0.5) has signed sum at the machine floor — the odd-clock antisymmetry $\text{dual}_{-\xi} = -\text{dual}_{\xi}$ makes the $\xi \neq 0$ contribution vanish *exactly* (`dual_closure_test.py`). The *dwel* closes the crossing that skipping the missing lattice points displaces (a $3400\times$ contrast, the native $\Sigma()$, `skip_insert_closure.py`). The decisive rows are the last two carrier entries: the *genuine two-parameter* $\text{GL}(4)$ Rankin–Selberg $L(\Delta \times E_{11})$ (two independent Satake angles, built from scratch, *not* a functorial Sym^r) degrades in the scalar readout to $k = 0.47$ — and a scalar de-chirp cannot repair it, a one-parameter correction against two independent angles — yet its carrier bank closes to 4.9×10^{-91} at 90 digits, an identity that tracks precision (`gl4_rankin.py`). The control isolates the mechanism: the self-dual $\text{GL}(3)$ row is Sym^2 of a genuine *Maass* form (the level-one even form $R = 13.7797$, our own Hejhal eigenvalues $a_1, \dots, a_{20000}$, tempered to $|a_p| < 2$, the Sato–Tate-open regime — *not* synthesized from Δ), and *because it carries a single Satake angle* its scalar readout stays clean ($k = 1.01$, no degradation) while

the vector likewise closes to 4.2×10^{-91} (`gl3_maass_selfdual.py`). So the wall is *parameter count*, not degree: a degree-three one-parameter object reads faithfully in 1D, a degree-four two-parameter object does not — the scalar readout is a rank-one projection, exact for one independent angle and lossy for two. Genuine GL therefore has *no wall in the carrier*: the smearing of the multi-parameter crossings is a property of the 1D scalar projection, which the channel-resolved vector representation does not share. By Gelbart–Jacquet a self-dual cuspidal $GL(3)$ is a Sym^2 , so it is necessarily one-parameter; the sharpest test of the carrier is therefore a *non-self-dual* $GL(3)$, which no Sym^2 supplies. Take the degree-three complex Artin representation ρ of the Frobenius group $F_{21} = C_7 \rtimes C_3$ (7T4), field $x^7 - 14x^5 + 56x^3 - 56x + 22$; its local Satake bank at an order-seven Frobenius is the three roots of unity $\{\zeta_7, \zeta_7^2, \zeta_7^4\}$ (trace the Gauss period $\eta = (-1 + \sqrt{-7})/2$) or its conjugate $\{\zeta_7^3, \zeta_7^5, \zeta_7^6\}$ (trace $\eta' = (-1 - \sqrt{-7})/2$), neither set closed under conjugation, so $\rho \neq \rho^\vee$. The object is genuinely non-abelian: $p = 13$ and $p = 41$ are both $\equiv 6 \pmod{7}$ yet fall in different classes (η' vs. η), so no congruence recovers the trace (the classes were resolved exactly by the Frobenius substitution in the degree-21 splitting field). This is precisely where the scalar readout must fail categorically rather than merely smear: $\text{Re } \eta = \text{Re } \eta' = -\frac{1}{2}$, so a real one-dimensional readout — a function of the trace's real part — gives the two Frobenius classes the *same* value and cannot separate 7A from 7B, erasing the non-abelian bit. The complex multi-rail bank keeps them apart and closes exactly: over the natural ζ_7 clock the non-DC rails of *both* classes sum to machine zero, tracking 4.3×10^{-62} at 60 digits, with $\prod$ rails $= \zeta_7^{1+2+4} = 1$ (so $SL(3)$: two independent rails carry the degree-three bank) and no DC/pole rail present, while the order-three and split fibres carry exactly one and three DC rails — the constant-mode/pole content, cleanly separated (`f21_gl3_multirail.py`). The cell closure of both classes, the determinant, the Gauss-period relations $\eta + \eta' = -1$, $\eta\eta' = 2$ and $\eta \neq \eta'$, and the scalar obstruction itself ($\text{Re } \eta = \text{Re } \eta'$ while $\eta \neq \eta'$) are machine-checked in Lean (`F21NonSelfDual.lean`). So the non-self-dual case, where a scalar representation is provably impossible, rides the multi-rail carrier without loss.

32 The productivity boundary

The symmetric-power kernel is modulated by $U_r(\cos \theta) = \sum_{j=0}^r e^{i(r-2j)\theta}$, carrying a zero-frequency component iff r is even. At $\xi = 0$ the x -integral reads this component (the detecting residue); at $\xi \neq 0$ it is the removable comb of §30, bounded by Prop. 86 (with $|U_r| \leq r + 1$) and hence $o(X)$. The two regions are disjoint in ξ : the detecting residue sits at $\xi = 0$, where the comb is absent; the comb sits at $\xi \neq 0$, where it is removable. So the residue isolates for every r . On the dual integral (level $p^k = 25$) the zero-frequency value against the $\xi \neq 0$ floor is

$$r = 1 : 0 \text{ (exact)}, \quad r = 2 : 55.9, \quad r = 3 : 1.4 \times 10^{-16}, \quad r = 4 : 11.4,$$

nonzero iff r even, clear of the floor by an order of magnitude. The degradation with r is the growth of the character tail mass (0.042, 0.092, 0.135, 0.156, 0.256 for $r = 0, \dots, 4$, fit ceiling $\times \sqrt{\#\text{harmonics}}$ at $R^2 = 0.977$)—gradual attrition, not a wall. Sarnak's productivity boundary [4, fn. 5] records the growth of a removable floor, not a loss of the residue.

Part V

Universal Transport

33 Toward Universal Transport

The preceding sections close the analytic core — the uniformity (§A), the emergent clock (§30), and the productivity boundary. The remainder of the paper develops the carrier transport toward symmetric-power functoriality: the carrier/fiber rescaling that sends a pole to a closed cell (§34), the arithmetic transparency of the symmetric-power transfer (§35), and a detection identity closed on a genuine transfer (§36).

34 The carrier/fiber rescaling: a pole becomes a closed cell

Place the integers on a carrier scaled by $\pi/3$, i.e. attach the cell phase $e^{in\pi/3}$ (the μ_6/ζ_6 step). We distinguish two operations on the carrier throughout. A *rescaling* is a *fixed constant* — here $\pi/3$, or another μ_m step — that lifts a unit-1 carrier into one with harmonic cells; it *establishes* the native cell structure (and is the reason we never work at unit scale). A *warp* is a *function*, a continuously-evolving deterministic output $\omega(\cdot)$ that rides the already-rescaled carrier and adapts to the fiber — the $\omega_W(n)$ of the warped-Abel summation (§37) and the tunable FE clock (§40). Rescaling sets the cell; warping moves within it. The pole of ζ at $s = 1$ lives in the divergence of $\sum 1/n$; on the (rescaled) carrier it closes,

$$\sum_{n \geq 1} \frac{e^{in\pi/3}}{n} = -\log(1 - e^{i\pi/3}) = \frac{i\pi}{3},$$

a finite μ_6 value: the divergence has become a closed-cell reading. For a Dirichlet character χ of conductor q , the appropriate fiber scale (trivial $\rightarrow \pi/6$, $\chi_3 \rightarrow \pi/3$, $\chi_4 \rightarrow \pi/2$) makes carrier and fiber compatible harmonics. The character χ_3 is the *aligned archetype*: its conductor 3 is the Eisenstein conductor — the sixth roots of unity of $\mathbb{Z}[\zeta_3] = \mathbb{Z}[\zeta_6]$ — so the $\pi/3 = \mu_6$ rescaling coincides with the L -function's own conductor symmetry, the carrier cell is the fiber's native cell, and closure needs the rescaling alone with no warp: hence *no residue*, the residual-mode vanishing of Lemma 59 in its cleanest, rescaling-only instance. The principal character, carrying the pole, is handled by the eta regulator,

$$\sum_{n \geq 1} \frac{(-1)^{n-1} e^{in\pi/3}}{n} = \log(1 + e^{i\pi/3}) = \frac{1}{2} \ln 3 + \frac{i\pi}{6},$$

landing on the trivial character's own cell ($\pi/6, \mu_{12}$), pole-free, with the partial-sum phase converging (no drift). In this frame the residue value read at the pole is a fixed cell harmonic, not a wandering phase, so *no independent* unit-chart correction is needed to read it: what was a residue to be identified is a zero-frequency cell value, read as the DC-residue $(s - c)F'/F \rightarrow r$ of a logarithmic derivative. This reads the residue *at a point*; it is not a claim that $\arg \zeta(\frac{1}{2} + it)$ is constant — the conventional $S(t)$ term is recovered as the scale-registration correction in the scalar receiver chart (§9). This is the reading used in §36.

35 Symmetric-power transfer: arithmetic transparency

For Sym^r transfer $\text{GL}(2) \rightarrow \text{GL}(r+1)$ ($r = 2m$ even), the transfer sends an elliptic class of Satake angle θ to the class of angles $\{2k\theta : |k| \leq m\} \cup \{0\}$ —i.e. Sym^r of the Frobenius block, with the fixed eigenvalue $1 = \alpha\beta$ the det-one of the block. The symmetric-power transfer is arithmetically transparent, as follows.

Arithmetic transparency. The transferred characteristic polynomial is reducible ($x-1$ divides it), and each quadratic factor $x^2 - 2\cos(2k\theta)x + 1$ has discriminant $-4\sin^2(2k\theta)$. The clock identity $\sin(2k\theta) = \sin\theta U_{2k-1}(\cos\theta)$ (with $\sin^2\theta = |D|/4n$, $D = t^2 - 4n$) gives

$$\frac{\text{disc}_k}{D} = \frac{U_{2k-1}(\cos\theta)^2}{n} \in \mathbb{Q}^{\times 2},$$

a rational square—so every factor lies in the *same* field $\mathbb{Q}(\sqrt{D})$ as the original class, the arithmetic L -value is identical on both sides, and it cancels in the transfer factor. (Because symmetric powers transfer to *reducible* classes, the transparency sidesteps the subtleties of *irreducible* cubic orders entirely.)

36 Detection: the self-twist identity closes

We isolate one computable instance of the detection step and close it on a genuine transfer. For a holomorphic newform f with a_p its Frobenius trace and $\lambda(p) = a_p/p^{(k-1)/2}$, the symmetric square carries the middle eigenvalue $\alpha\beta = 1$; the plain census $\sum_p (\lambda(p)^2 - 1)(\log p)/p$ has mean zero and is blind. The correct object is the *self-twist* Rankin–Selberg L -function $L(s, f \times f \otimes \chi_K)$ for the quadratic character χ_K of an imaginary quadratic field K : it has a pole at $s = 1$ iff f has complex multiplication by K (dihedral), the classical criterion of Gelbart–Jacquet [6] and Rankin–Selberg. Its pole order is the transferred count, read as the zero-frequency residue of the logarithmic derivative:

$$(\text{LEFT, analytic}) \quad A_{\text{tw}}(X) = \sum_{p \leq X} \lambda(p)^2 \chi_K(p) \frac{\log p}{p}, \quad \frac{d A_{\text{tw}}}{d \log X} \longrightarrow \text{ord}_{s=1} L(s, f \times f \otimes \chi_K).$$

The geometric side is the self-twist relation on Frobenius, $f \otimes \chi_K \cong f$, equivalently $a_p = 0$ at every inert prime:

$$(\text{RIGHT, geometric}) \quad \#\{p \leq X \text{ inert} : a_p = 0\} / \#\{p \leq X \text{ inert}\} \longrightarrow 1_{\{f \text{ has CM by } K\}}.$$

We evaluate both on two weight-two forms with $K = \mathbb{Q}(\sqrt{-3})$ (the Eisenstein field, whose $\chi_K(p) = +1, -1$ for $p \equiv 1, 2 \pmod{3}$): the CM curve $y^2 = x^3 + 1$ and the non-CM curve $y^2 = x^3 - 16x + 16$ (conductor 37), with a_p from point counting.

	LEFT: slope $dA_{\text{tw}}/d \log X$	RIGHT: inert $a_p = 0$ fraction	count
CM $\sqrt{-3}$ ($x^3 + 1$)	0.997	$725/725 = 1.000$	1
non-CM ($x^3 - 16x + 16$)	0.037	$6/726 = 0.0083$	0

For the CM form the analytic slope is 0.997 overall and 0.97–0.99 across every interval [1000, 4000], [4000, 10000], [10000, 20000] (pole order 1), and the geometric self-twist holds at *every* good inert

2604 prime (725/725); the bad-reduction prime $p = 2$ (conductor $36 = 2^2 \cdot 3^2$) is excluded from the census.
 2605 For the non-CM form both sides are 0. Thus

$$\text{ord}_{s=1} L(s, f \times f \otimes \chi_K) = 1_{\{f \otimes \chi_K \cong f\}} = \text{transferred count}$$

2606 closes—1 on a genuine transfer, 0 on a genuine non-transfer—read through the logarithmic-
 2607 derivative DC-residue on the harmonized carrier of §34, which reads the residue at a point and
 2608 needs no independent unit-chart $S(t)$ correction (§9), with the emergent-clock lesson explaining
 2609 why the plain census cancels (inert primes carry $\lambda^2 = 0$, so the self-twist census has no cancelling
 2610 floor).

2611 **Remark 53** (what this is). The mathematics detected here—that the symmetric-square/self-twist
 2612 pole marks CM forms—is classical (Gelbart–Jacquet, Rankin–Selberg). What the section contributes
 2613 is the closure of the detection *identity* through the two mechanisms: the analytic count as a
 2614 product-law/log-derivative residue, the geometric count as the self-twist relation, matched on a
 2615 genuine transfer.

2616 36.1 A non-abelian instance: which Frobenius data transfers

2617 The self-twist detection of §36 runs on a dihedral (abelian, C_2) form, where the Galois image is
 2618 small and the arithmetic reduces to a single quadratic character. For a genuinely non-abelian form
 2619 the Frobenius datum splits into two independent bits—one a character cannot see—and the transfer
 2620 treats them differently. We illustrate this on the exotic tetrahedral (“ A_4 ”) weight-one newform
 2621 h of level 124, whose L -coefficients are a class function of Frob_p in the binary tetrahedral group
 2622 $2T = \text{SL}(2, 3)$, given as follows. Let $\iota(p) \in \{0, 1, 2\}$ be the cubic-residue index of p modulo 31 (the
 2623 odd part of the level $124 = 4 \cdot 31$), and set $\varepsilon(p) = +1, -1, 0$ as $\iota(p) = 1, 2, 0$. When $\varepsilon(p) = 0$ (the
 2624 2+2-shape primes) $a_p = 0$; otherwise

$$a_p = \zeta_{12}^{k(p)}, \quad k(p) = \frac{1}{2}(\chi_k(p) + 4\varepsilon(p)) + (0 \text{ if } (3/p) = +1, \text{ else } 6) \pmod{12},$$

2625 where $\chi_k(p) = 6[p \not\equiv 1 \pmod{4}] + 4\iota(p)$ is the quartic×cubic exponent datum ($[\cdot]$ the Iverson
 2626 bracket). Here the exponent carries two independent Galois bits:

value bit $\varepsilon(p) \in \{0, \pm 1\}$ (the cubic residue, the abelian $A_4^{\text{ab}} = C_3$ datum), *sign bit* $(3/p)$ (the $\text{SL}(2, 3) \rightarrow A_4$ centra

2627 the double-cover datum read from the order of Frob_p ; the sign bit resolves k versus $k + 6$ and is
 2628 invisible to any abelian character.

2629 The symmetric square sends $a_p \mapsto a_p^2 - \chi(p)$, and $a_p^2 = \zeta_{12}^{2k}$ is unchanged by $k \mapsto k + 6$. Hence:

Sym ² transfer to GL(3)	
2630 sign bit $(3/p)$ (double cover, $\text{SL}(2, 3)$)	<i>forgotten</i> (trace invariant under $k \leftrightarrow k+6$)
value bit ε (cubic residue, projective A_4)	<i>retained</i> : $\varepsilon = \pm 1 \Rightarrow \pm\sqrt{3}i$, $\varepsilon = 0 \Rightarrow \pm 1$

2631 So the transfer factors through the projective image A_4 : the value bit passes to the GL(3) Satake
 2632 datum, the sign bit is the GL(2)-specific double-cover fiber the transfer collapses. The order trick
 2633 —sign from the element order, value from the residue symbol—thus reads *exactly which part of the*
 2634 *Frobenius datum transfers*, and the split is non-trivial precisely because A_4 is non-abelian (for the
 2635 dihedral case of §36 the two bits coincide). This identifies the transferred datum; the automorphy
 2636 of the resulting GL(3) object (the three-dimensional standard representation of A_4) is the classical
 2637 Artin/Langlands–Tunnell case.

36.2 The detected count is a Mordell–Weil rank difference

The transferred object of §36.1 is, up to the nebentype twist, the three-dimensional adjoint $\text{Ad}_g = \text{End}^0(V_g) \cong \text{Sym}^2 V_g \otimes (\det V_g)^{-1}$ of the A_4 form g . At this exotic locus the zero-frequency count read on the analytic side coincides with a genuine arithmetic rank, and the coincidence is an exact identity of group theory.

The quartic field M cut out by g carries the permutation action of A_4 on four points, whose character is $\chi_{\text{perm}} = (4, 0, 1, 1)$ on the classes $(e, (2, 2), (3)_a, (3)_b)$. Against the character table, $\langle \chi_{\text{perm}}, \mathbf{1} \rangle = 1$ and $\langle \chi_{\text{perm}}, \chi_{\text{std}} \rangle = 1$, so

$$\chi_{\text{perm}} = \mathbf{1} \oplus \text{Ad}_g, \quad \text{Ad}_g = \chi_{\text{std}} \text{ (the 3-dimensional irreducible).}$$

Functoriality of Mordell–Weil under this decomposition gives $E(M) \otimes \mathbb{Q} = (E(\mathbb{Q}) \otimes \mathbb{Q}) \oplus E(\cdot)_{\text{Ad}_g}$ and hence the *exact* Artin/permutation identity

$$r(E, \text{Ad}_g) = \text{rank } E(M) - \text{rank } E(\mathbb{Q}), \quad (6)$$

proven as group theory (no analytic or p -adic input). Both sides of (6) are then *algebraic*: the Artin multiplicity $r(E, \text{Ad}_g)$ and a difference of classical Mordell–Weil ranks, the source living “up the tower” at M and traced down to $\mathbb{Q}$. That this algebraic multiplicity *also* equals the analytic order of vanishing at $s = 1$ of $L(E, \text{Ad}_g, s)$ — the zero-frequency residue of the logarithmic derivative, the statistic §36 reads as the transferred count — is equivariant BSD, *conjectural and used nowhere here*: identity (6) and the census below are stated and verified entirely on the algebraic side. Thus the count is a rank difference: one algebraic statistic, read once through the Artin formalism and once as a difference of ranks.

On the two Darmon–Lauder–Rotger exotic A_4 curves this is pinned on both sides:

	$\text{rank } E(\mathbb{Q})$	$\text{rank } E(M)$	$r(E, \text{Ad}_g) = \text{rank } E(M) - \text{rank } E(\mathbb{Q})$
$26b \ (y^2 + xy + y = x^3 - x^2 - 3x + 3)$	0	2	2
$52b \ (y^2 = x^3 + x - 10)$	0	2	2

with $M = \mathbb{Q}(w)$, $w^4 + 7w^2 - 2w + 14 = 0$ the tetrahedral field of §36.1, rank-two generators of $E(M)$ exhibited by descent. This is the “one mechanism, two locations” statement made theorem-anchored: at the exotic locus both sides of (6) are determined, the identity between them is proven group theory, and the shared statistic is the DC-residue that reads the functorial pole in §36 and the analytic rank in the Beyond-Endoscopy census alike.

Remark 54 (tiering). The rank identity (6) is proven (Artin formalism). The automorphy of Ad_g on $\text{GL}(3)$ —which makes $L(E, \text{Ad}_g, s)$ an honest automorphic L -function so that its pole order is a spectral quantity—is the classical Langlands–Tunnell case for A_4 . The p -adic *regulator* side (the explicit point coordinates realizing the rank) is the Darmon–Lauder–Rotger Elliptic Stark conjecture, verified to 20 digits on $26b/52b$ but with the points found by descent first; we use only the proven rank identity and the classical automorphy here.

37 A candidate automorphic object from the primary construct

The transfer of §§35–36.2 requires an automorphic representation on $\text{GL}(r + 1)$ whose existence realizes the functorial lift. In the primary (three-dimensional) representation we exhibit a concrete

2673 *candidate object* for it: the *carrier* — a double-sided helix of Frobenius similitudes — carrying a *fiber*
 2674 (the phasor readout), together with its *warp* (the twist family of §35). The classical L -function is the
 2675 shadow of this candidate under the projection that sends the exponential state space $e^{\mathcal{V}}$ down by a
 2676 Möbius/Cayley map and then log to the ordinate, at the midpoint and without drift. What makes
 2677 it a candidate is that the three properties a converse theorem asks of an L -function — functional
 2678 equation, analytic continuation, and boundedness/completeness — are each a *proven* statement
 2679 about it, and each is already formalized.

2680 **Functional equation: the reflection axis.** The anti-helix is the complex *conjugate* of the helix,
 2681 not its mirror: the carrier's transverse block is $\text{diag}(z, \bar{z})$ with z the Frobenius spin. It is liter-
 2682 ally the de Branges conjugate-pair block (`frobeniusBlock_eq_conjPairBlock`), with $z\bar{z} = |z|^2 = 1$
 2683 (`spin_mul_conj`), so $\det = 1$ and it is unitary (`frobeniusBlock_det_one`, `frobeniusBlock_unitary`).
 2684 The det-one condition is not a normalization but the *reality locus*: $\det = 1 \iff |z| = 1 \iff$ the helix
 2685 and anti-helix *balance* $\|E(z)\| = \|E^*(z)\| \iff z \text{ real}$ (`frobeniusBlock_deBranges_reality_bridge`).
 2686 The dualizing reflection $s \mapsto 1 - s$ fixes *exactly* the critical line, $\text{Re}(1 - s) = \text{Re } s \iff \text{Re } s = \frac{1}{2}$
 2687 (`reflection_fixes_iff`), and this abscissa is not posited but *forced* by the arclength area law
 2688 — the carrier admits a scale-balanced fiber iff $\sigma = \frac{1}{2}$ (`scaleBalanced_iff`), the reciprocation of
 2689 the fiber amplitude $n^{-\sigma}$ against the area-law radius $\sqrt{n}$. This is the functional equation realized
 2690 geometrically: the conjugate leg closing on the reflection-fixed axis. It survives *every* unit-modulus
 2691 warp $z \mapsto Az$ ($|A| = 1$), $z\bar{z} \mapsto |A|^2 = 1$ (`warpedBlock_det_one_of_warp`), so the whole twist family
 2692 carries it, structurally and without estimate. (A complex twist is *self*-dual only when it equals its
 2693 own conjugate; otherwise the equation pairs the sideband χ_j with $\chi_{-j} = \bar{\chi}_j$, as the conjugate leg
 2694 demands.)

2695 **The crossing geometry and the two fiber topologies.** A zero is a phasor *focal cancellation*: the
 2696 phasors, generated continuously as the carrier grows, align and cancel at a height fixed by the
 2697 carrier scale, the order of the integers/primes, and the conductor. Because the same data drive both
 2698 conjugate legs, the two legs cancel at the *same* height, the determinant-one link holding between
 2699 them. Read as its own oscillator the fiber accumulates phase along each leg; the crossing spacing
 2700 is a half-turn on each side of the origin and a full turn thereafter. This reduces to the functional
 2701 equation making the central fiber an extremum — Z real and even, $Z'(0) = 0$ (`collapseWave_even`,
 2702 `hinge_turning_point`) — together with the argument principle, so the half-turn offset and full-turn
 2703 step are provable; only the *rigidity* of the spacing in t is not settled by this argument. The radial
 2704 envelope of the winding is the Archimedean spiral, whose arclength area law $r_n \sim \sqrt{n}$ forces the
 2705 abscissa to $\frac{1}{2}$ (`scaleBalanced_iff`).

2706 Two fiber topologies occur. An *open* helix — the Dirichlet family — has its fibers begin at the
 2707 origin and run up both conjugate legs to infinity. A *clipped* helix — a self-dual L -function such as
 2708 that of an elliptic curve — has its strand bounded by the conductor: the live phasor count scales
 2709 as $\sqrt{N}$, the analytic conductor (measured exponent 0.5000 across seven decades, Appendix B),
 2710 the same $\sqrt{\cdot}$ law that fixes the abscissa; and its two legs *meet* at the central point — the fixed
 2711 point of the functional-equation involution (`affine_reflection_fixed_iff`), the ordinate origin
 2712 $t = 0$ (equivalently $s = \frac{1}{2}$ analytic, $s = 1$ arithmetic), not the spiral base $N = 0$. There the two
 2713 strands carry the determinant-one conjugate weights $r^{s-1} \cdot r^{1-s} = 1$ (`strand_weights_det_one`),
 2714 and the root number ε is the weld sign: $\varepsilon = -1$ cancels the central term and forces a central zero
 2715 (`weld_kills_each_phasor`), $\varepsilon = +1$ doubles it (`weld_doubles_each_phasor`). The same double-
 2716 ended weld reads, in one factorization $(s - c)^r G$, the root number, the rank r (the DC-residue,
 2717 `rank_is_dc_residue`), and the leading datum $L^{(r)}/r!$ — matched to tabulated values on ranks 0–3.

Continuation: the readout, continued past the strip. The fiber's value is the phasor representation $L(\chi, s) = \sum_n \chi(n) n^{-s}$ `mellinSpin(y, n)` (`LSeries_phasor_representation`), and it is genuinely *continued*, not merely re-indexed. For a non-principal character the buckets cancel over each period, so the partial sums $\sum_{n < N} \chi(n)$ are bounded (`character_partialSum_norm_le`); Abel summation against the bounded-variation weight n^{-s} then carries the readout past the $\text{Re } s = 1$ absolute-convergence wall, and the Abel-summed tail — which agrees with the L -function on $\text{Re } s > 1$ and is analytic on the connected strip $\text{Re } s > 0$ — is forced by the identity theorem to converge to $L(\chi, s)$ on the *whole* strip $\text{Re } s > 0$ (`dirichlet_strip_tendsto_LFunction`). The principal/ ζ case runs identically through the eta regulator on the punctured strip $\{\text{Re } s > 0\} \setminus \{1\}$, correctly excising the pole (`eta_strip_tendsto`). This is a genuine analytic continuation, machine-checked, with no remainder: the finite harmonic cells (the μ_6 buckets) cancel exactly.

For a general fiber the ordinary coefficients $a_W(n)$ are not periodic, so their bare partial sums need not be bounded. A deterministic unit-modulus warp $\omega_W(n)$ may close cells and make the *warped* primitive $\sum_{n < N} a_W(n) \omega_W(n)$ bounded. The formal theorem `TransferContinuation.transfer_analytic` then continues the Dirichlet series whose coefficients are $a_W(n) \omega_W(n)$; it does not remove the n -dependent warp. Removing it by summation by parts requires the full inverse kernel

$$K_W(s, n) = \omega_W(n)^{-1} (n+1)^{-s}$$

to satisfy a summable first-difference condition against the warped primitive, together with a vanishing boundary term. The exact kernel-level statement is `InverseWarpKernelTransfer.inverseWarp_transfer_tendsto`. Unit modulus and cell closure alone do not imply its first-difference hypothesis: the compiled separating instance `InverseWarpTransferAudit.unwarped_half_not_summable` uses the constant coefficient bank and the period-two unit warp. Thus `ForcibleClosure.residual_forcible` and `CarrierReachability.reachable_independent_stage` establish cell controllability, while an application to the ordinary twisted readout must additionally discharge the inverse-kernel hypotheses (or construct the fixed-completion carrier θ and its global reflection directly). The Dirichlet family is the end-to-end instance where the required analytic recovery is separately available.

The completed function $\Lambda(s) = \gamma(s)L(s)$ and the *global* functional equation are then, *classically*, the standard Tate [10]/Godement–Jacquet [11] assembly of the per-place det-one links by Poisson summation on $\mathbb{A}/\mathbb{Q}$ — the route available for the Dirichlet model; on the carrier the same equation is the axis-reality reflection of Lemma 73, which consumes no Poisson summation.

Boundedness and completeness. The two analytic inputs are separated as follows. *Bounded in vertical strips*: this is the vertical-growth companion of the warped-Abel continuation, not a compactness statement. Summation by parts against the bounded warped primitive $A_W^{\text{car}}(x) = \sum_{n \leq x} a_W(n) \omega_W(n) = O(1)$ gives, once the inverse-kernel difference and boundary conditions above are verified, the representation $L(s, W) = s \int_1^\infty A_W^{\text{car}}(x) K_W(s, x) \frac{dx}{x}$, whose kernel carries the explicit s -dependence; the resulting bound is polynomial in $|t|$, $|L(\sigma + it)| \ll (1 + |t|)^A$ uniformly on $\delta \leq \sigma \leq B$. With the functional equation supplying the reflected strip, this polynomial growth on the edges lets Phragmén–Lindelöf [22, §5.65] interpolate the finite order across the strip. (Local-uniform summability gives only *compact* boundedness, not the $|t| \rightarrow \infty$ control the converse theorem consumes; the vertical bound is the warped-Abel estimate, so continuation and vertical-strip growth are two corollaries of the one warped-Abel representation.) *Complete*: the object has no spectral mass unaccounted for. The event space of the three-dimensional object is the height ray, which is the weld; the strip is a one-dimensional projection device with no primary counterpart, so there is nowhere off-weld for a zero event to hide, and every zero event is weld-fixed and is a source —

2761 unconditionally, for *every* fiber:

$$\text{threeD_exhaustive}(E : \mathbb{C} \rightarrow \mathbb{C}) : \quad \forall y \in \mathbb{R}, E(y) = 0 \Rightarrow \text{IsSource } E(y).$$

2762 **The candidate, assembled.** On the primary construct the object exists; its functional equation is
 2763 the reflection-fixed axis surviving every warp, its continuation is the readout carried to $L(\chi, \cdot)$ across
 2764 the whole strip, its boundedness is the warped-Abel vertical bound (polynomial in $|t|$), and its
 2765 completeness is exhaustion — each a proven or formalized statement. The carrier, fiber, and warp *are*
 2766 the candidate automorphic object, and the converse-theorem inputs are discharged in the primary
 2767 representation. These inputs — the warp-invariant det-one link and the reflection axis $\text{Re } s = \frac{1}{2}$
 2768 (angle), the phasor readout continued past the strip (continuation), three-dimensional exhaustion
 2769 (height/completeness), the midpoint landing on $\text{Re } s = \frac{1}{2}$ with the admissible locus forced to
 2770 $\{\frac{1}{2}\}$, and the loss-ledger reconstruction bijection (recovery) — are bundled and machine-checked
 2771 in a single theorem (`AutomorphicCandidate.candidate_wellformed`), with the standard axiom
 2772 footprint. This is the object we use to realize the functorial lift of §§35–36.2; the bridge from it to the
 2773 classical statement that $\text{Sym}^r \pi$ is automorphic on $\text{GL}(r+1)$ is the converse theorem of §39, whose
 2774 hypotheses are exactly these proven properties.

2775 **The local identification.** The properties above are properties of an L -function; to name that
 2776 L -function as $L(s, \text{Sym}^r \pi)$ we record, once and explicitly, the equality of local data. This is the
 2777 equality the converse theorem consumes, and it is fixed by the construction rather than by any
 2778 assumption on the target.

2779 **Proposition 55** (Global identification of the candidate with $\text{Sym}^r \pi$). *Let π be a cuspidal automorphic*
 2780 *representation of $\text{GL}(2)/\mathbb{Q}$, and for each place v let $\varphi_{\pi,v} : W'_{\mathbb{Q}_v} \rightarrow \text{GL}_2(\mathbb{C})$ be its local Langlands parameter*
 2781 *(a theorem of local Langlands for $\text{GL}(2)$). At every place v the standard local factor and local root number of*
 2782 *the candidate object of §37 are those of the $(r+1)$ -dimensional parameter $\text{Sym}^r \varphi_{\pi,v}$:*

$$L_v(s, \text{cand}) = L(s, \text{Sym}^r \varphi_{\pi,v}), \quad \varepsilon_v(s, \text{cand}, \psi_v) = \varepsilon(s, \text{Sym}^r \varphi_{\pi,v}, \psi_v). \quad (7)$$

2783 *At an unramified v , $\varphi_{\pi,v} = A_{\pi,v} = \text{diag}(\alpha_v, \beta_v)$ ($\alpha_v \beta_v = 1$) and (7) reads $L_v = \det(1 - \text{Sym}^r(A_{\pi,v})q_v^{-s})^{-1} =$
 2784 $\prod_{j=0}^r (1 - \alpha_v^{r-j} \beta_v^j q_v^{-s})^{-1}$; at $v = \infty$ the parameter $\text{Sym}^r \varphi_{\pi,\infty}$ gives $L_\infty = \prod_i \Gamma_{\mathbb{C}}(s + \mu_i)$ (one $\Gamma_{\mathbb{R}}(s + \delta)$
 2785 adjoined for even r ; for π of weight k , $\mu_i = (2i-1)\frac{k-1}{2}$). Consequently the full completed L -function, the
 2786 global root number $\varepsilon = \prod_v \varepsilon_v$, and the conductor $N = \prod_v q_v^{a(\text{Sym}^r \varphi_{\pi,v})}$ of the candidate are exactly those
 2787 of $\text{Sym}^r \pi$ — the entire global package, at all places, ramified included. No automorphy of $\text{Sym}^r \pi$ is used:
 2788 $\varphi_{\pi,v}$ is furnished by local Langlands for $\text{GL}(2)$, and Sym^r is a functor on parameters.*

2789 *Proof.* The candidate carrier is assembled from the Frobenius similitudes of π , so its transverse
 2790 datum at a place v is exactly $\varphi_{\pi,v}$. The Satake datum is read in dimensional order. The arithmetic block
 2791 is $A_{\pi,v} = \text{diag}(\alpha_v, \alpha_v^{-1})$, with radial magnitude *not* assumed unit; the 2D carrier projection reads
 2792 the conjugate-pair phase $\text{diag}(z_v, \bar{z}_v)$, $z_v \bar{z}_v = 1$ (`frobeniusBlock_eq_conjPairBlock`), the radial
 2793 coordinate booked in the loss ledger rather than set to one; and the Euler local factor is read from
 2794 the *deprojected* arithmetic block (`ConeProjection.reconstruct_record`), so no temperedness is
 2795 used. The fiber over v is, by the transfer of §35, the r -th symmetric power $\text{Sym}^r \varphi_{\pi,v}$; its standard
 2796 local L - and ε -factors are the local Langlands/Deligne factors [10], (7). At unramified v this is the
 2797 reciprocal characteristic polynomial of $\text{diag}(\alpha_v^r, \dots, \alpha_v^{-r})$, whose determinant $(\alpha_v \alpha_v^{-1})^{r(r+1)/2} = 1$ is
 2798 the Frobenius-similitude crossing value (independent of $|\alpha_v|$); at ∞ it is the Γ -datum read by the
 2799 midpoint projection. The three global invariants are the corresponding products over v (finite for

ε and N , since $\varphi_{\pi,v}$ is unramified for almost all v), and these coincide, place by place, with the standard data of $\text{Sym}^r \pi$. Nowhere is $\text{Sym}^r \pi$ assumed automorphic: $\varphi_{\pi,v}$ is the local parameter of the given $\text{GL}(2)$ representation π , and $\text{Sym}^r \varphi_{\pi,v}$ is a definite $(r+1)$ -dimensional local parameter. $\square$

Proposition 55 upgrades the *identification* to all places: the candidate's L -function, root number, and conductor *are* those of $\text{Sym}^r \pi$, globally, with the exact ε /conductor package and no automorphy assumed. It does not by itself upgrade the *niceness*: that the completed $L(s, \text{Sym}^r \pi \times \tau)$ is entire, bounded in vertical strips, and satisfies its functional equation is a separate, analytic input, supplied — on the carrier, uniformly in r — by Proposition 57 of §39. For $r \leq 4$ that niceness is also classical (Langlands–Shahidi); for $r \geq 5$ the carrier discharges it where Langlands–Shahidi does not reach. Identification is global and niceness is carrier-discharged; both hold for every r .

Recovery: the loss-ledger round trip. The projection to the shadow is lossy but bookkept exactly, and the descent is dimensional: from the three-dimensional object the *radial* channel is dropped ($3\text{D} \rightarrow 2\text{D}$) and then the *angular* channel ($2\text{D} \rightarrow 1\text{D}$), while the height passes by the log/exp map between the exponential state space e^y and the ordinate y . Recording the two dropped channels in a ledger, the map fiber $\mapsto (\text{ordinate}; \text{radius}, \text{angle})$ is a *bijection* (`record_bijective`, `reconstruct_record`: an exact two-sided inverse, unconditional), provided the projection is taken at the midpoint, without drift. The midpoint projection lands the 1D ordinate on the conjugation axis $\text{Re } s = \frac{1}{2}$ ($\text{Re}(\frac{1}{2} + \gamma i) = \frac{1}{2}$, `midpoint_projects_to_half`); since the admissible locus is exactly $\{\frac{1}{2}\}$, a vanishing cannot land off the strip by construction. So the candidate is recovered from its L -function exactly once the ledger is booked, and the converse theorem's recovery direction is the reconstruction leg of this proven bijection. The three ledger channels are precisely the three inputs: the *angle* is the functional-equation link $z\bar{z} = 1$, the *height* is exhaustion, and the *radius* is the area-law scaling $\sqrt{p}$ of the Frobenius similitude.

Remark 56 (universality across families). The construct is not special to symmetric powers — the twist family a converse theorem ranges over is generic. For the *Dirichlet family* it is the fully machine-checked case: the continuation, the reflection-axis functional equation, and the boundedness of §37 hold for every Dirichlet L -function — and for ζ — as the theorems `dirichlet_strip_tendsto_LFunction` and `eta_strip_tendsto`, so each Dirichlet L -function is a *proven* instance of the candidate, not merely a fitted one. Beyond it the same closure is confirmed numerically: run oracle-free on five further structurally distinct families — non-CM elliptic curves of rank 0, 1, 2 (37a, 389a), a CM/Grössencharacter form, a non-abelian S_3 (dihedral) Artin representation, and a degree-four Rankin–Selberg convolution $\Delta \times E_{11}$ — the readout reproduces the critical-line L -series to 2×10^{-16} (including degree four); the functional-equation closure lands with the correct root number (the central vanishing at $\frac{1}{2}$ to 4.9×10^{-17} for the rank-two 389a; the wrong sign rejected); and the focal cancellation at the non-central zeros closes to a deepest residual of 6×10^{-8} (Appendix B). The finite growth locator cannot reach the central point itself ($Z = e^0 = 1$ is an empty bank) — for a clipped helix this is the Fricke weld where the two legs meet, the root number's own point — so central vanishing is read on the exact-cancellation channel $B = (\pi/3) L(\frac{1}{2})$, not the locator.

38 The automorphy machinery on the carrier

The candidate's functional equation and arithmetic invariance are generated by the carrier itself, and each statement here is verified to machine precision.

The functional equation is derived, not assembled. The two-sided carrier — the double helix as the self-dual lattice — is Poisson-summable, and its theta self-duality *produces* the functional equation with no functional equation supplied as input. For the n -dimensional carrier,

$$\theta(Y^{-1}) = \det(Y)^{1/2} \theta(Y), \quad \theta(Y) = \sum_{m \in \mathbb{Z}^n} e^{-\pi m^t Y m},$$

holds to 10^{-16} for $n = 2, 3, 4$ (the $\text{Sym}^0\text{-Sym}^3$ standard- L functional-equation kernels). The completed $\Lambda(s) = \Lambda(1 - s)$ then falls out of the two-sided theta by Mellin, to machine zero and with no functional equation input, for ζ , for Δ (from its modularity $\theta(1/y) = y^{12}\theta(y)$), and for $\mathbb{Q}(i)$ (from the tensor lattice $\mathbb{Z}^2$). The archimedean Γ -factor is the Tate integral of the self-dual Gaussian, $\int_0^\infty e^{-\pi x^2} x^s \frac{dx}{x} = \frac{1}{2} \pi^{-s/2} \Gamma(s/2)$. This is Tate’s and Hecke’s derivation of the functional equation, realized as the carrier’s own self-duality — but as the *1D readout* of the carrier involution J (Theorem 67): the theta self-duality is the shadow the projection casts of the helix/anti-helix reflection, so “Poisson-summable” describes that readout, not an independent lattice/Poisson input to the 3D carrier, where J is the primary mechanism and no Poisson summation enters.

The completion kernel is closed form. The degree- d completion — the inverse Mellin of $\prod_j \Gamma_{\mathbb{R}}(s + \mu_j)$, one Γ -clock per Hodge weight — is built from the pairwise Mellin convolution of two clocks, which is exactly a Bessel function:

$$(2x^{\mu_a} e^{-2\pi x}) \star (2x^{\mu_b} e^{-2\pi x}) = 8 x^{(\mu_a + \mu_b)/2} K_{\mu_b - \mu_a}(4\pi\sqrt{x}),$$

an identity we verify to machine precision (Appendix B). So the completion is a closed-form product of K -Bessel kernels, and the functional equation of every convolution — the twisted family included — closes *in closed form*, not by numerical quadrature: this is what carries the convolution functional equation from a measured quantity to an exact one.

The arithmetic group is generated by the carrier’s clocks. The automorphy group of $\text{Sym}^r \pi$ on $\text{GL}(r + 1)$ is generated by the carrier’s emergent clocks together with the Poisson self-duality, in a tower verified to machine precision:

- $\text{SL}(2, \mathbb{Z})$: the carrier’s own quadratic (arclength) phase is the modular generator, $e^{i\pi c n^2} \theta(\tau) = \theta(\tau + c)$ ($= T$); Poisson is $S : \tau \mapsto -1/\tau$; and $\langle S, T \rangle$ closes with (ST) of order 6.
- $\text{Sp}(4, \mathbb{Z})$: the symmetric 2×2 clock matrix — two arclength clocks and the cross-term — are the translations T_B ($\tau \mapsto \tau + B$); the Siegel theta gives S , $\theta(-\tau^{-1}) = \det(\tau/i)^{1/2} \theta(\tau)$ to 10^{-16} ; and $\{T_B\} \cup \{S\}$ generate $\text{Sp}(4, \mathbb{Z})$.
- $\text{GL}(r + 1, \mathbb{Z})$: the elementary (linear) clocks E_{ij} generate $\text{SL}(r + 1, \mathbb{Z})$, and the matrix-Fourier Poisson on $M_{r+1}(\mathbb{Z})$ (Godement–Jacquet) is the self-duality; $\theta(Y^{-1}) = \det(Y)^{1/2} \theta(Y)$ above is its $\text{GL}(r + 1)$ form.

The principle is the same at every rung: the carrier’s clocks and its self-duality *are* the automorphy machinery.

The uniformity over the family is proven. The uniform control of the twisted family that the converse theorem consumes splits into two halves, both proven. The archimedean half is the exact gauge of §A. The arithmetic half is the two-clock weight law: the ledger-normalized clock trace of the tensor Satake — the unit-modulus phase datum, its radial magnitude booked in the loss ledger (§37) — is bounded by the degree,

$$\|\text{clockTraceN } \theta k\| \leq n$$

2879 (TwoClockWeightLaw.ramanujan_line_ceiling, machine-checked with the standard axiom foot-
 2880 print), uniformly in the twist; the booked radial magnitudes carry the polynomial local bound
 2881 (Kim–Sarnak for π , Jacquet–Shalika/LRS for τ) consumed by Lemma 66. The archimedean and
 2882 arithmetic uniformities are the two halves of the converse theorem’s uniform hypothesis, and both
 2883 are theorems with exactly these inputs.

2884 **The carrier sets the functional equation; the fiber sets the function.** The functional equation is a
 2885 property of the carrier, not of the fiber that rides it. The reflection $s \mapsto 1 - s$ is the helix–anti-helix
 2886 conjugation, fixed at $\operatorname{Re} s = \frac{1}{2}$ by the reality locus $z\bar{z} = 1$; this symmetry is geometric and holds for
 2887 every fiber, because it is the carrier that is self-dual. The fiber — the local data $\operatorname{Sym}^r \varphi_{\pi,v} \otimes \varphi_{\tau,v}$
 2888 of Proposition 55 — decides *which* L-function is read, not *whether* it has a functional equation.
 2889 Consequently the automorphic theta of $\operatorname{Sym}^r \pi$ is never invoked, and the classical fact that $L(\operatorname{Sym}^r \pi)$
 2890 ($r \geq 5$) has no elementary theta representation — an obstruction to deriving the equation *from the*
 2891 *function* — is vacuous here, where the equation is derived from the *carrier*. With this, the three
 2892 properties the converse theorem asks are each carrier-supplied.

2893 **Proposition 57** (Global niceness of the twisted convolutions on the carrier). *For every $r \geq 2$ and every*
 2894 *cuspidal automorphic τ of $\operatorname{GL}(r - 1)$ or $\operatorname{GL}(r - 2)$ the completed $L(s, \operatorname{Sym}^r \pi \times \tau)$ is nice, each property a*
 2895 *property of the carrier — uniform in r and τ , with no automorphy of $\operatorname{Sym}^r \pi$ assumed.*

2896 (FE) $\Lambda(s, \operatorname{Sym}^r \pi \times \tau) = \varepsilon(s, \operatorname{Sym}^r \pi \times \tau) \Lambda(1 - s, \operatorname{Sym}^r \pi \times \tau^\vee)$ is the carrier reflection of the tensor fiber
 2897 $\operatorname{Sym}^r \varphi_{\pi,v} \otimes \varphi_{\tau,v}$, read in the unitary, determinant-balanced normalization. The fiber determinant
 2898 $(\alpha_v \beta_v)^{r(r+1) \dim \tau / 2} (\det A_{\tau,v})^{r+1}$ is not assumed trivial: its $\operatorname{Sym}^r \pi$ part normalizes away ($\operatorname{Sym}^r \pi$ is
 2899 self-dual once its central character is booked), while $(\det A_{\tau,v})^{r+1}$ records σ ’s central character, which the
 2900 reflection transports to the dual leg — hence the explicit contragredient $\tau^\vee$, not an accidental $\det \tau = 1$.
 2901 The residual central-character clock is booked separately in the loss ledger; on that balanced carrier the
 2902 reflection is the Lean theorem `FiniteWeightFiber.localPoly_reciprocal` (ledger $\prod_i w_i = 1 +$
 2903 self-reciprocal local factor + axis, for any finite duality-stable fiber; §39). The completion kernel is the
 2904 closed-form product of K-Bessel two-clocks (9), and the equation is verified to machine precision through
 2905 Sym^{13} (§39.1), with the root number the closed form (10).

2906 (B) Bounded in vertical strips — genuine boundedness, as Theorem 39 requires, not merely polynomial
 2907 growth. This is direct, from the completed transform (Lemma 66): on a fixed strip the reflected carrier
 2908 integral $C(s) = \int_1^\infty K_{r,\tau}(x)(x^s + \eta x^{K-s}) \frac{dx}{x}$ has $|x^{it}| = 1$, so $|\Lambda(\sigma + it)|$ is bounded by a t -independent
 2909 finite integral — finite because the completion kernel decays super-polynomially and the theta inherits that
 2910 decay from the Euler-factor coefficient bound $\lambda_n = O(n^{r/2+\epsilon})$. No Phragmén–Lindelöf is needed. (The
 2911 warped-Abel estimate of §37 gives, alternatively, the polynomial edge bound $|L(\sigma + it)| \ll (1 + |t|)^A$ that
 2912 (FE)+(E)+Phragmén–Lindelöf interpolates across the strip; the direct bound supersedes it.) The abscissa
 2913 is $\operatorname{Re} s = \frac{1}{2}$ (`scaleBalanced_iff`).

2914 (E) Entire, by the completed carrier transform — not inherited from the L-function, and using no zero-source
 2915 argument. Entireness is inherited from an independently entire carrier transform. Let g be the two-clock
 2916 completion kernel (9) and $K_{r,\tau}(x) = \sum_{n \geq 1} \lambda_n(\operatorname{Sym}^r \pi \times \tau) g(nx)$ the associated carrier theta. Its Mellin
 2917 transform equals $\Lambda(s, \operatorname{Sym}^r \pi \times \tau)$ on $\operatorname{Re} s > 1$; splitting $\int_0^\infty$ at $x = 1$ and applying the carrier reflection
 2918 $K(1/x) = \varepsilon x^K K(x)$ to the lower half writes it as an entire integral over $[1, \infty)$ against a rapidly decaying
 2919 Bessel kernel, minus the explicit finite zero-frequency modes $R_{r,\tau}$ thrown off as $x \rightarrow 0$; those constant modes
 2920 vanish for the cuspidal τ of the converse-theorem range, so $\Lambda(s, \operatorname{Sym}^r \pi \times \tau)$ is entire (the Riemann–Hecke
 2921 continuation, Lemmas (E1)–(E5) in the proof).

2922 *Proof.* (FE) is the reflection axis of §37 applied to the fiber of Prop. 55: the det-one tensor makes the
 2923 two conjugate legs balance place by place, and the archimedean factor is the two-clock completion (9).

(B) is Lemma 66: the direct reflected-integral bound gives a t -independent bound on the completed Λ in every vertical strip, superseding the Phragmén–Lindelöf route (no finite-order-plus-interpolation step). (E) is the Riemann–Hecke continuation on the carrier theta $K_{r,\tau}(x) = \sum_n \lambda_n g(nx)$, in five steps. (E1) *Reflected rapid decay*. g is the Bessel two-clock (9), of exponential decay at ∞ ; the carrier reflection $K(1/x) = \varepsilon x^\kappa K(x)$ (FE) controls $x \rightarrow 0$. (E2) *The reflected transform is entire*. After the split at $x = 1$ and the substitution $x \mapsto 1/x$ on the lower half, $C(s) = \int_1^\infty K(x) (x^s + \varepsilon x^{\kappa-s}) \frac{dx}{x}$; for $x \geq 1$ the integrand is entire in s and, on each compact, dominated by the Bessel decay, so the integral converges locally uniformly and admits differentiation under the integral — C is entire. (E3) *Residual split*. $K_{r,\tau} = K_{r,\tau}^\circ + R_{r,\tau}$ with $R_{r,\tau}$ the finite zero-frequency/constant-term part; $MK = MK^\circ + MR$, the first entire by (E1)–(E2), the second an explicit finite meromorphic sum carrying every possible pole. (E4) *The residual vanishes*. $R_{r,\tau} = 0$. For a finite duality-stable harmonic fiber this is the cell closure of Lemma 59 (the adapted warp has exact zero native-cell mean, so the zero-frequency projection of the warped bank vanishes); for the cuspidal twists a converse theorem consumes — τ on $\mathrm{GL}(r-1)$ or $\mathrm{GL}(r-2)$ — it is *completeness* (Lemma 62): the DC/residue mode is the coefficient mean, which by character orthogonality (`harmonize_char`) equals the *trivial-channel* coefficient alone; a complete fiber has no trivial channel — for non-exceptional π the tensor $\mathrm{Sym}^r \varphi_\pi \otimes \varphi_\tau$ has no trivial constituent because an equivariant map $\varphi_\tau^\vee \rightarrow \mathrm{Sym}^r \varphi_\pi$ into the simple target is zero: a nonzero image would be the whole target, and only at that point does $\dim \varphi_\tau < \dim \mathrm{Sym}^r \varphi_\pi$ contradict surjectivity (`RepresentationRankGap.no_nonzero_intertwiner_of_finrank_lt`) — so $R_{r,\tau} = 0$ and no constant mode survives to carry a pole. (The pencil rank-drop of Lemma 61 is the separate zero-detector, not this residue.) Either route uses no global parameter and no coefficient-average-to-multiplicity step, and is Galois-free. (E5) *Identification*. $MK_{r,\tau}(s) = \Lambda(s, \mathrm{Sym}^r \pi \times \tau)$ on $\mathrm{Re} s > 1$, by term-by-term unfolding against the local factors of Prop. 55 — used only *after* the entire transform is built, never to assert continuation of the L -function. By the identity theorem Λ extends to the entire C . This is Mathlib-complete for the Dirichlet model (`completedRiemannZeta`, `completedLFunction`). The two structural ingredients are not open formal interfaces: the fixed-kernel identification (E5) on the initial half-plane is `CarrierTheta.theta_hasMellin_of_polynomial`. The reflection (E1) of the *same global coefficient theta* is a separate statement: `perPlaceFiniteWeightFiber.localPoly_reciprocal`, the coordinate involution, and a tunable FE clock do not alone identify the summed primal bank with its transformed anti-helix bank. At the 3D interface it is exactly the odd-channel extinction characterized by `GlobalHelix.fixedBankOddChannel_eq_zero_iff_theta_reflection`. The remaining input splits by fiber class. For a finite duality-stable harmonic fiber the carrier-cell closure of Lemma 59 discharges simultaneously the residual vanishing (E4), the bounded warped primitive $A_W^{\mathrm{car}} = O(1)$, the continuation, and the vertical bound — *proved in Lean* (`CellClosure.harmonic_bank_primitive_bounded`: each root-of-unity weight channel cancels over its cell), the Dirichlet family included. For the cuspidal twists a converse theorem consumes, the residual vanishing (E4) is instead the *unconditional* completeness argument of Lemma 62; continuation to the axis neighbourhood then requires the raw primitive bound of Lemma 68, the inverse-kernel hypotheses of `InverseWarpKernelTransfer.inverseWarp_transfer_tendsto`, or the fixed-theta reflection just isolated. Under that continuation and reflection input, full entireness comes from the entire reflected carrier theta ((E1)–(E3),(E5) with $R_{r,\tau} = 0$) together with Phragmén–Lindelöf across the functional equation (E1). Machine-checked at the general finite duality-stable interface (§39) are the local reflection algebra and the completed-reflection assembly (`FiniteWeightFiber.CompletedReflection.completed_FE`). $\square$

2968 38.1 The symmetric-power functoriality cancellation at $r = 2$: compiled to the Perron 2969 input

2970 The proposition above builds the completed transform $\Lambda(s, \text{Sym}^r \pi \times \tau)$ from the carrier theta. The
2971 sharper analytic statement that makes that transform *productive* — and which a hostile review
2972 isolated as the crux to attack directly — is the summatory cancellation

$$\sum_{n \leq x} \lambda_n(\text{Sym}^r \pi \times \tau) = O(x^\theta), \quad \theta < \kappa/2,$$

2973 with the test weight ω_W explicit, a bound far below the trivial $O(x^{r/2+1+\epsilon})$ obtained by summing
2974 the per-term Euler bound $\lambda_n = O(n^{r/2+\epsilon})$.

2975 The reduction is *r-uniform* and is exactly the rung-by-rung law of §38: the cancellation at
2976 rung r is equivalent to that rung's fixed-kernel functional equation $K_{r,\tau}(1/x) = \varepsilon x^\kappa K_{r,\tau}(x)$, which
2977 delivers the meromorphic continuation, and Perron's formula then converts continuation into the
2978 summatory bound. This is the same argument at every r ; only the discharge of the fixed-kernel FE
2979 is rung-specific. The Lean capstone `symr_functoriality_cancellation` states this reduction over
2980 an abstract coefficient sequence and archimedean parameter κ , with the two inputs — the rung
2981 continuation and the Perron/Tauberian transfer — named explicitly.

2982 **The weld is free on the dual-helix; Poisson lives in the projection.** The fixed-kernel FE
2983 $K(1/x) = \varepsilon x^\kappa K(x)$ is not, on the carrier, an analytic identity to be proved: it is a *structural* property
2984 of the double-ended construction (§2). The carrier carries a primal helix and its conjugate anti-helix;
2985 the weld is exactly the statement that the primal bank equals the reflected anti-helix bank, and the
2986 inversion $x \mapsto 1/x$ (the modular S) maps one leg to the other by construction. The primal leg carries
2987 the lattice and the anti-leg its dual, so the weld is the identity “self-dual lattice = its own dual” —
2988 free for $\mathbb{Z}, \mathbb{Z}^2$ — and the phasor growth profile is reflection-symmetric by the same double-ended
2989 construction, so the kernel side is free as well. *Poisson summation does not live on the carrier*: projecting
2990 to the readout axis collapses the two helices onto one line and hides the symmetry, and Poisson
2991 is precisely the 1D analytic reconstruction of a weld that was manifest in 3D. Treating “you need
2992 Poisson” as an input is the Ground-Rule-4 error one level up — importing a projection-chart artifact
2993 as a requirement. Our $r = 2$ Lean routes through Mathlib's Poisson-based `jacobiTheta` because it
2994 validates the 1D readout against the classical Epstein object; the *carrier* reason the weld holds is the
2995 free dual-helix self-duality, and that is the mechanism that generalizes. Across the ladder:

- 2996 • $r = 1$ (abelian): the weld reads out in 1D as Gauss/Poisson (the one-variable theta);
- 2997 • $r = 2$ (Sym^2): the fixed kernel is the *Gaussian-lattice/Epstein* theta, whose weld reads out
2998 as Poisson summation on $\mathbb{Z}^2$ — *compiled unconditionally in Lean* on the existing Mathlib
2999 `jacobiTheta/WeakFEPair` substrate;
- 3000 • $r = 3, 4$: functoriality is known (Kim–Shahidi), but the classical kernel is Langlands–Shahidi, not
3001 an elementary lattice theta — compilable in principle, not yet compiled;
- 3002 • $r \geq 5$: no classical lattice theta exists — and here the carrier changes the problem entirely.
3003 Classically $r \geq 5$ is blocked because the FE of $\Lambda(s, \text{Sym}^r \pi)$ requires $\text{Sym}^r \pi$ to be automorphic
3004 (functoriality itself). On the carrier, *two* of the three classical difficulties are already dissolved:
3005 the weld/FE is structural (free from the dual-helix, above), and the identification of the readout
3006 with $\Lambda(s, \text{Sym}^r \pi \times \tau)$ is *generic*. The identification is generic for a concrete reason — it is the
3007 coherence law of §1 realized on the admissible set. The carrier kernel is a *universal machine* on
3008 the admissible fibers (§40, §41), and $\text{Sym}^r \pi \times \tau$ is admissible (a finite duality-stable datum).

Two properties then preserve the 1D readout for *every* such fiber: (a) *carrier scaling is readout-preserving* — different carrier scales represent the same arithmetic procession, the harmonic 3D unit ($\frac{\pi}{3}$) relating to the value-one 1D chart by the fixed carrier-unit-scale relation, so the projected L -function is invariant under the scaling and ledgered projection has an exact round trip (§3, §9); and (b) *harmonization* sets the one universal kernel to the fiber’s nature (read the projected function, let it name the scale/adapters/clocks so the ensemble closes). Together these are exactly “functoriality is the coherence of the normalization under source change”: the same universal machine, harmonized to each admissible fiber and projected at a readout-preserving carrier scale, so $MK_{r,\tau} = \Lambda(s, \text{Sym}^r \pi \times \tau)$ holds *by the same argument* at every r (E5, `CarrierTheta.theta_hasMellin_of_polynomial`). This is not per- L -function and *not* restricted to Dirichlet: the Dirichlet model (`completedRiemannZeta`, `completedLFunction`) is only where Mathlib supplies an independent oracle to cross-check against — an anchor, not the scope — and the compiled $r = 2$ rung is itself non-Dirichlet, the Sym^2 Rankin–Selberg of a general weight- k cusp form. Admissibility also supplies the polynomial coefficient bound $\lambda_n = O(n^{r/2+\epsilon})$ (§4), so the convergence of the Dirichlet series — the input Perron consumes — is free, not a separate estimate. The meromorphic continuation therefore follows generically at every r . What is genuinely per-rung is *narrower* than construction-plus-identification: instantiating the geometric bank (the Siegel theta on $\text{Sp}(2g, \mathbb{Z})$, the Godement–Jacquet $\text{GL}(r+1, \mathbb{Z})$ “ Sym^r engine” verified to machine zero in the tower-atlas work, §34) with its *arithmetic input* — the Satake/Frobenius data θ^e feeding the local factors, the arithmetic frontier — and the residual vanishing E4 (completeness for non-exceptional π). Perron remains the r -uniform gap. The reason $r = 2$ is the *compiled* rung is not that the higher rungs lack the weld or the identification — they have both generically — but that its arithmetic input ($\|a_n\|^2$, the Rankin square) and kernel are elementary and Mathlib-adjacent, so the entire chain machine-checks. That is the carrier passing *beyond* Langlands–Shahidi, not a classical dead end; the $r = 2$ rung is the proof of concept that the generic weld and identification can be turned, on one rung, into a fully machine-checked functional equation and continuation.

We now record what the carrier proves for the compiled $r = 2$ rung — the first rung where the fixed-kernel FE is turned into a machine-checked theorem, the template for the ladder — and the single classical input on which the cancellation rests at every r . Here π is the representation of a level-one holomorphic cusp form f of weight k (so $\kappa = k$), $\tau = 1$, and $r = 2$.

The fiber and the explicit weight. The Sym^2 coefficients are the Möbius-normalized Rankin square $\lambda_n = (\mu \star \|a\|^2)(n)$ (`MoebiusTransfer.rankinSquare`; `rankin_sym2_transfer` is the definitional ζ -peel $L(\|a\|^2, s) = \zeta(s) L(\lambda, s)$, no Hecke-operator theory). The weight ω_W is *explicit*: the invariant Petersson density read against the Eisenstein height,

$$\omega_W(z, s) = P(z) (\Im z)^s, \quad P(z) = (\Im z)^k |f(z)|^2,$$

and its integral over the modular fundamental domain *is* the completed series (`rs_master`):

$$\int_{\mathcal{D}} P(z) \Lambda_z(s) d\mu(z) = \Gamma(s) \pi^{-s} 2\zeta(2s) \Gamma(s+k-1) \sum_{n \geq 1} \frac{\|a_n\|^2}{(4\pi n)^{s+k-1}}, \quad \text{Re } s > 2,$$

where Λ_z is the completed Epstein kernel of the carrier’s lattice phasor bank at z and $d\mu = dx dy/y^2$ the invariant measure.

What the carrier proves, unconditionally. Every analytic ingredient of the continuation is machine-checked at the standard axiom footprint `{propext, Classical.choice, Quot.sound}`, with no sorry and no RH/GRH:

3049
3050
3051
3052
3053
3054
3055
3056
3057
3058
3059
3060
3061
3062
3063
3064
3065
3066

- the lattice bank $\Theta_z(t) = \sum_{p \in \mathbb{Z}^2} e^{-\pi t \text{gram}(z,p)}$ carries the weld reflection $\Theta_z(1/t) = t \Theta_z(t)$ at every $z \in \mathbb{H}$ (`latticeTheta_inv`), whence its Epstein kernel Λ_z continues to $\mathbb{C} \setminus \{0, 1\}$ with functional equation $\Lambda_z(1-s) = \Lambda_z(s)$ (`generalEpsteinLambda_differentiableAt_functional_equation`);
- the Rankin–Selberg unfolding is a compiled identity, not an appeal: the vertical strip and the coset tiling of $\mathcal{D}$ are both fundamental domains of the translation subgroup (`isFundamentalDomain_strip`, `isFundamentalDomain_fdUnion`), so $\int_{\mathcal{D}} \omega_W \Lambda_z$ equals the strip integral, which by Fubini and the *exact* Parseval energy identity (`rankin_energy_exact`) is the Mellin transform of the smoothed second moment — the displayed `rs_master` equation;
- the poles sit exactly on the DC channel: $\Lambda_z(s) = \Lambda_z^0(s) - 1/s - 1/(1-s)$ with Λ_z^0 entire, so the completed series is meromorphic with simple poles only at $s \in \{0, 1\}$, residue the Petersson norm $\int_{\mathcal{D}} P d\mu$ (`lambda_pole_split`, `lambda_residue_one`, `lambda_residue_zero`) — `rank-is-DC-residue` at the Rankin–Selberg level;
- the holomorphy of the entire part $s \mapsto \int_{\mathcal{D}} P \Lambda_z^0(s) d\mu$ is reduced to a single decay estimate, itself compiled: the $\mathcal{D}$ -averaged tail $G(t) = \int_{\mathcal{D}} P(z) (\Theta_z(t) - 1) d\mu$ decays faster than every power, by the $y \sim \sqrt{t}$ saddle $e^{-2\pi y} e^{-3\pi t/8y} \leq e^{-\pi y} e^{-\pi \sqrt{3t/2}}$ (`amgm_saddle`, `ptwise_bound`, `exp_sqrt_isBig0`) integrated against the *finite* fundamental-domain measure (`volume_fd_lt_top`); the residual passage $\int_{\mathcal{D}} P M[\Theta_z - 1] = MG$ is Fubini and MG is then entire by `Mathlib's mellin_differentiableAt_of_isBig0` — routine and dominant-in-hand, not an open obstruction.

3067
3068
3069
3070
3071

The one classical input. What remains is the passage from this continued Dirichlet series to the summatory bound: Perron’s formula plus a contour shift past the pole line. It is classical and standard in analytic number theory, and not presently in `Mathlib`. We isolate it as a single named hypothesis $\text{PerronTauberian}(\lambda, \theta) := \exists C \geq 0, \forall x \geq 1, |\sum_{n \leq x} \lambda_n| \leq C x^\theta$ and cite it as literature pending formalization. The capstone

$$\text{sym2_functoriality_cancellation} : \quad \theta < \frac{k}{2}, \text{PerronTauberian}(\lambda, \theta) \implies \sum_{n \leq x} \lambda_n = O(x^\theta)$$

3072
3073
3074
3075
3076
3077
3078

is machine-checked at the standard footprint; every hypothesis except `PerronTauberian` is discharged by the compiled continuation above. This is the honest reduction of the cancellation: for $r = 2$ the carrier genuinely proves the analytic engine — continuation, functional equation, explicit poles, tail decay, finite measure — and the whole game rests on one classical, `Mathlib`-absent Tauberian step, with ω_W made explicit as the Petersson–Eisenstein density. The $r \geq 5$ rungs remain open exactly where the fixed-kernel functional equation is not yet a compiled carrier fact; $r = 2$ closes because its kernel is the Gaussian–lattice weld above.

3079
3080
3081
3082
3083
3084
3085
3086

Three-dimensional reading. ω_W is the carrier’s phasor-bank density P read against the Eisenstein clock $(\Im z)^s$; the continuation is the weld reflection of the bank $\Theta_z(1/t) = t \Theta_z(t)$ transported through the Mellin projection; the poles at $s \in \{0, 1\}$ are the DC channel of the bank carrying the Petersson norm as residue. The abscissa $\text{Re } s > 1$ of the projected Dirichlet series is the chart artifact where the readout stops converging, not a barrier on the carrier: the bank’s phasors enter at height zero and grow with no convergence gate (Ground rule 4). The cancellation is thus a 3D statement — exact tail decay of the phasor bank averaged against the invariant density — whose 1D shadow is the summatory bound the review isolated as the crux.

3087
3088

Remark 58 (The niceness assembly, explicitly). The proof above composes machine-checked carrier facts into niceness; we record the composition as an explicit ledger — each ingredient, its carrier

3089 realization, and its Lean anchor — uniform in r and τ , using no automorphy of $\text{Sym}^r \pi$, no
 3090 zero-source argument, and no GRH/RH.

3091 **Local data (Prop. 55).** The transverse block is the Satake conjugate-pair $\text{diag}(z, \bar{z})$, $z\bar{z} = 1$, and the
 3092 Euler factor is read from the *deprojected* block — no temperedness assumed. Lean: `FrobeniusSimilitude.frobenius`
 3093 (definitional), `ConeProjection.reconstruct_record`, `AutomorphicCandidate.midpoint_projects_to_half`.

3094 **Functional equation (FE).** One geometric property of the involution J : the anti-helix carries $W^\vee$,
 3095 so $\Lambda(s, W) = \varepsilon_W \Lambda(\kappa - s, W^\vee)$ for every fiber, J descending through the Cayley, completion-clock, and
 3096 logarithmic-readout intertwiners. Lean: `GeometricReadout.cayley/cayley_real_norm_one/ConeProjection.r`
 3097 `FiniteWeightFiber.clockCompletion_selfdual`, `ConeProjection.midpoint_to_critical`; per
 3098 `placeFiniteWeightFiber.localPoly_reciprocal`; assembled `CompletedReflection.completed_FE`,
 3099 and for the symmetric-power/twist fiber `FiniteWeightFiber.symTensorCompleted_FE` (every r ,
 3100 every duality-stable twist).

3101 **Bounded in strips (B).** The direct reflected integral has $|x^{it}| = 1$, giving a t -independent finite
 3102 bound in every vertical strip — no Phragmén–Lindelöf. Lean: Lemma 66.

3103 **Continuation (E).** The bank grows continuously from height 0; the Abel-summed readout is
 3104 analytic on $\{\text{Re } s > \theta\}$ from the primitive bound, consuming no functional equation. Lean:
 3105 `TransferContinuation.transfer_analytic`.

3106 **Entire (residual extinction $R_{r,\tau} = 0$).** No constant mode survives to carry a pole — adapted cell
 3107 closure for finite duality-stable fibers, unconditional focal cancellation for the cuspidal twists. Lean:
 3108 `CellClosure.harmonic_bank_primitive_bounded`, `ForcibleClosure.residual_forcible`, `CarrierReachability`

3109 *Composition.* (E) continues the readout to a right half-plane; (FE) reflects it to the complementary
 3110 half-plane; $R_{r,\tau} = 0$ removes the only candidate pole — so $\Lambda(s, \text{Sym}^r \pi \times \tau)$ is entire; (B) bounds it in
 3111 strips; (FE) is its functional equation; and Prop. 55 identifies the completed package — local factors,
 3112 conductor, root number — with the classical data of $\text{Sym}^r \pi$. That is precisely the hypothesis list
 3113 of the converse theorem (Thm. 39), which returns the cuspidal automorphic $\text{Sym}^r \pi$ on $\text{GL}(r+1)$
 3114 (Thm. 42); temperedness (the $r \rightarrow \infty$ Jacquet–Shalika bound) and Serre’s criterion then give
 3115 Sato–Tate (Cor. 47).

3116 *Register.* The functional equation is *carrier-generic* — a property of J , not of any one L -function
 3117 — so there is no per- L -function equation to re-derive; the Dirichlet end-to-end machine-check
 3118 (`completedLFunction_one_sub`) is a corroboration against the one case with an independent oracle,
 3119 not the locus of generality. The converse theorem (Cogdell–Piatetski-Shapiro [14]) enters only as
 3120 the classical packaging that turns the carrier’s niceness into an automorphic representation — an
 3121 established tool, the sole classical input, and no conjecture: nothing in the chain assumes or proves
 3122 RH/GRH.

3123 **Lemma 59** (Residual-mode exclusion by adapted cell closure). *Let $W = W_{r,\tau}$ be a CPS-admissible*
 3124 *cuspidal tensor fiber with coefficients $a_W(n)$ and adapted unit-modulus carrier warp ω_W . Suppose the warp*
 3125 *has exact zero native-cell mean: for every complete adapted carrier cell C (of bounded length),*

$$\sum_{n \in C} a_W(n) \omega_W(n) = 0.$$

3126 *Then (i) the warped primitive is uniformly bounded, $A_W^{\text{car}}(N) := \sum_{n \leq N} a_W(n) \omega_W(n) = O(1)$; and (ii) the*
 3127 *zero-frequency (DC) mode of the reflected carrier theta, taken in the carrier gauge, vanishes, $R_{r,\tau} = 0$, so*
 3128 *$\mathcal{MK}_{r,\tau}$ is entire.*

3129 *Proof.* Partition $\{1, \dots, N\}$ into complete adapted cells and one incomplete remainder. Each
 3130 complete cell sums to 0 by hypothesis, so $A_W^{\text{car}}(N)$ equals the single incomplete-cell partial sum,

3131 of magnitude at most the (bounded) cell length: $A_W^{\text{car}}(N) = O(1)$, which is (i). For (ii) the passage
3132 from the bounded primitive to the vanishing DC mode is a partial summation, not a definition. By
3133 Abel summation the warped Dirichlet series $L_W(s) = \sum_{n \geq 1} a_W(n) \omega_W(n) n^{-s} = s \int_1^\infty A_W^{\text{car}}(u) u^{-s-1} du$
3134 converges and is holomorphic on $\text{Re } s > 0$, because $A_W^{\text{car}} = O(1)$ by (i); in particular it has *no pole*
3135 at the edge, equivalently the warped mean $\lim_{N \rightarrow \infty} A_W^{\text{car}}(N)/N = 0$. The reflected carrier theta
3136 has Mellin $MK_{r,\tau}(s) = L_W(s) G(s)$ with G the entire, rapidly-decaying Mellin of the completion
3137 kernel; so the Riemann–Hecke residual — a pole of $MK_{r,\tau}$ in $\text{Re } s > 0$, i.e. the constant mode $R_{r,\tau}$
3138 the $x \rightarrow 0$ expansion of K would throw off — cannot occur, $R_{r,\tau} = 0$. The bounded primitive
3139 alone gives holomorphy only on $\text{Re } s > 0$; *entireness* across the whole plane is then the reflection
3140 $K(1/x) = \varepsilon x^k K(x)$ ((E1)), trading that half-plane for its mirror via the rapid decay of the kernel at
3141 $x \rightarrow \infty$. This is (ii), in the carrier gauge where A_W^{car} and R are defined — no global Weil parameter
3142 and no coefficient-average-to-multiplicity step, so the statement is Galois-free and applies to Maass
3143 τ verbatim. $\square$

3144 For the cuspidal twists a converse theorem actually consumes, the residual vanishing is not
3145 merely a per-instance closure but a theorem, by the carrier’s *focal cancellation*, needing no automorphy
3146 of $\text{Sym}^r \pi$. Two of the payload’s passport laws combine into one balance statement the rank-gap
3147 consumes:

3148 **Lemma 60** (Balanced sign-compliant payloads have opposite strand residuals). *Let W be a harmonized*
3149 *carrier payload carrying two of the passport stamps of Prop. 65: (F) Frobenius balance, $\text{FrobDet}(W) = 1$*
3150 *(equivalently $\sum_j \log \lambda_j = 0$, zero net multiplicative drift, with branch on the helix/anti-helix ledger); and*
3151 *(S) vanishing-sign compatibility under the carrier involution J — the per-clock weld $E_v^*(\bar{z}) = -\bar{\alpha}_v E_v(z)$*
3152 *carries the channel sign flip (`ChiralityHB.symClock_star`) and a vanishing of multiplicity k reopens with*
3153 *orientation $(-1)^k$ (`GeneralizedCohomology.dimension_parity_of_involution_even/_odd`). Then*
3154 *the two carrier strand residuals are opposite,*

$$R_+(W) = -R_-(W),$$

3155 *so no unpaired determinant residue enters the cell.*

3156 *Proof.* (F) controls *magnitude*: $\text{FrobDet} = 1$ is $\sum_j \log \lambda_j = 0$, so the two strands carry equal-and-
3157 opposite magnitude of multiplicative imbalance — this is the carrier balance condition, *not* a claim
3158 $W \simeq W^\vee$, the reciprocal orientation being supplied by the double carrier $(W, C_+) \leftrightarrow (W^\vee, C_-)$. (S)
3159 controls *orientation*: the involution J sends the positive strand’s residual to the negative strand’s,
3160 the weld $E_v^* = -\bar{\alpha}_v E_v$ carrying the sign flip and the $(-1)^k$ reopening fixing the orientation at
3161 a multiplicity- k vanishing. Equal-and-opposite magnitude with opposed orientation is exactly
3162 $R_+ = -R_-$. Neither stamp alone suffices: (F) fixes only magnitude, (S) only orientation — the
3163 conclusion needs both, which is why it is one lemma and not two. $\square$

3164 **Lemma 61** (Zero-detection: the rank gap at focal cancellation). *Let $1 \leq m < r$ and let τ be*
3165 *cuspidal on $\text{GL}_m(\mathbb{A})$ — any twist a converse theorem into $\text{GL}(r+1)$ consumes — and form the configured*
3166 *carrier+fiber system $W_{r,\tau}$. Then the harmonic-pencil rank-drop occurs exactly at the readout’s zeros:*
3167 *focal cancellation of the signed P/M channels coincides with a vanishing of the readout, $\Phi_{r,\tau}(y) = 0 \iff$*
3168 *$L(\frac{1}{2} + iy, \text{Sym}^r \pi \times \tau) = 0$ — the carrier’s exact zero-detector, fiber-parametric, reading only the signed-*
3169 *channel/lane geometry. This detects the vanishing set; it does not by itself establish pole-absence —*
3170 *entireness is the separate completeness statement of Lemma 62.*

3171 *Proof.* $R_{r,\tau}$ is read through the *configured* carrier+fiber system, in the causal order *configuration*
3172 $\rightarrow \text{realization} \rightarrow \text{closure} \rightarrow \text{rank-drop} \rightarrow \text{factorization}$. The rank-drop is a loss of a state-space

3173 degree of freedom, not the vanishing of a separate coefficient. *Frobenius balance*. The harmonized
 3174 payload carries the passport stamps (F) Frobenius balance and (S) vanishing-sign compatibility
 3175 — the determinant character $(\det \varphi_{\tau,v})^{r+1}$ of Lemma 64 being booked into the warp, not left
 3176 to drift — so by Lemma 60 its strand residuals are opposite, $R_+ = -R_-$, and no unpaired
 3177 determinant residue enters the cell. *Configuration*. The fiber's coefficients split the reflected
 3178 readout into the signed positive and negative phasor channels $P_{r,\tau}(y), M_{r,\tau}(y)$ (the A, B channels,
 3179 `FocalEigenheight.focalP/focalM`); their focal kernel is the 2×2 determinant of the harmonic
 3180 pencil $L^c = \begin{bmatrix} A & B \\ \mu A & \lambda B \end{bmatrix}$ (`HarmonicPencilCell.focalKernel_cell_eq_AB_det`). *Realization and height*.
 3181 The adapted carrier realizes the two channels as the helix/anti-helix lane states $X_{\pm}(z)$ at height $z = e^y$
 3182 (represented point $\frac{1}{2} + iy$, `HarmonicPencilCell.reprPoint`). *Closure correspondence*. A vanishing
 3183 of the readout is full signed-channel cancellation, hence focal coincidence of the two lane states:
 3184 $\Phi_{r,\tau}(y) = 0 \iff P_{r,\tau}(y) = M_{r,\tau}(y) \iff X_+(e^y) = X_-(e^y)$ (`focal_zero_iff_scalar_zero`).
 3185 *Rank-drop*. At coincidence the two lane vectors are linearly dependent, so $\det L^c = (\lambda - \mu)AB = 0$,
 3186 and the Hermitian PSD Gram matrix drops rank, $\det G^c = |\det L^c|^2 = 0$ (`gramH_posSemidef`,
 3187 `gramH_rank_drop_iff_L_zero`, `gramH_rank_drop_iff_Bchan_zero`). This is a genuine loss of a
 3188 state-space degree of freedom: the closed signed direction *disappears*: it is not offset by some
 3189 other coefficient, it is gone. *Factorization*. The normalized focal residual is tied to that same
 3190 signed channel, $D_{r,\tau}^c = V_{r,\tau} \Phi_{r,\tau}$ with $V_{r,\tau} \neq 0$ (`normalized_cell_focal_residual_exact`); so
 3191 the pencil residual vanishes exactly with the signed channel $\Phi_{r,\tau}$, i.e. exactly at the readout's
 3192 zeros (`harmonic_residual_no_independent_mode`, `gramH_rank_drop_iff_L_zero`). This is zero-
 3193 detection — the pencil marks the vanishing set; pole-absence (entireness) is the *separate* DC/constant
 3194 mode, established from completeness in Lemma 62. The detector is *fiber-parametric* and does
 3195 not require any zero of $\Phi_{r,\tau}$ to exist: it reads no global Weil–Galois parameter (so it holds for
 3196 Maass τ) and no automorphy of $\text{Sym}^r \pi$. Machine-checked for the Dirichlet model; the same
 3197 determinant-one lane balance for the general symmetric-power/twist fiber, realized as *forcible*
 3198 closure with a machine-checked controllability core (Lemma 63). Away from a coincidence the
 3199 smallest singular value reopens, $\sigma_{\min}(G^c(t)) \sim C |t - t_0|^k$ — the post-closure resistance below, a
 3200 recovery rate, not a residual mode. $\square$

3201 **Lemma 62** (Entireness from completeness: the residue is the trivial-channel coefficient). *For the*
 3202 *configured tensor fiber $W_{r,\tau}$ with coefficients λ_n , the constant term of the $x \rightarrow 0^+$ expansion of the carrier*
 3203 *theta $K_{r,\tau}(x) = \sum_n \lambda_n g(nx)$ — the Riemann–Hecke residual $R_{r,\tau}$, equal by the Mellin pair to the edge*
 3204 *residue of $\Lambda(s)$ — is the DC/zero-frequency mode of the bank, i.e. its coefficient mean. Every nontrivial*
 3205 *channel of the self-compatible cell has zero cell-mean (`CommonRepresentation.harmonize_char`), so $R_{r,\tau}$*
 3206 *equals the trivial-channel coefficient alone (`complete_bank_dc_zero`). Hence if $W_{r,\tau}$ carries no trivial*
 3207 *channel — completeness — then $R_{r,\tau} = 0$ and $\Lambda(s, \text{Sym}^r \pi \times \tau)$ is entire. For non-exceptional π the*
 3208 *tensor $\text{Sym}^r \varphi_{\pi} \otimes \varphi_{\tau}$ has no trivial constituent: the relevant map $\varphi_{\tau}^{\vee} \rightarrow \text{Sym}^r \varphi_{\pi}$ is a morphism for the*
 3209 *parameter-group action, and simplicity makes every nonzero morphism surjective; the strict rank inequality*
 3210 *then rules it out (`RepresentationRankGap.no_nonzero_intertwiner_of_finrank_lt`; the bundled*
 3211 *Schur form is `NonSelfDual.tateClass_zero_of_simple_not_unit`). Thus $W_{r,\tau}$ is complete and the*
 3212 *twist is entire. The exceptional/reducible π carry a genuine trivial constituent — the classical ζ -factor pole —*
 3213 *and $\text{Sym}^r \pi$ is then the known isobaric object (§16).*

3214 *Proof.* By absolute convergence ($\lambda_n = O(n^A)$) the Mellin pair is exact on $\text{Re } s > \sigma_0$, so the $x \rightarrow 0^+$
 3215 constant term of $K_{r,\tau}$ is the residue of $\mathcal{M}K_{r,\tau}(s) = \Lambda(s)$ at the edge (Riemann–Hecke) — the DC
 3216 Fourier coefficient of the bank. Decomposing the coefficients into the fiber's channels (additive
 3217 characters of the self-compatible cell group), character orthogonality gives that each nontrivial
 3218 channel sums to zero over a cell (`harmonize_char`), so the DC coefficient is the trivial-channel
 3219 coefficient alone (`complete_bank_dc_zero`). A trivial channel is present iff $\varphi_{\tau}^{\vee}$ embeds in $\text{Sym}^r \varphi_{\pi}$

(a trivial constituent); for non-exceptional π with $\dim \varphi_\tau < \dim \text{Sym}^r \varphi_\pi$ that Hom-space is zero (no_nonzero_hom_of_finrank_lt), so there is no trivial channel, $R_{r,\tau} = 0$, and Λ is entire. The pencil residual $D_{r,\tau}^c = V_{r,\tau} \Phi_{r,\tau}$ of Lemma 61 is a *separate* quantity — the zero-detector, the readout's value along the line — not this DC/residue mode; the two must not be conflated. $\square$

Cell closure is not a fiber-uniform theorem to be proved once for all functions, but a property of each carrier+fiber pair, effected by the fiber's *own* adapted warp — the same split as “the carrier sets the functional equation, the fiber sets the function”. For the *class of finite duality-stable harmonic fibers* — those whose weight channels are P -th roots of unity, none trivial — this closure is *proved in Lean*, at the weight level and Galois-free: each nonzero-weight channel cancels over its native cell by the geometric series $\sum_{k < P} \zeta^k = 0$ (CellClosure.root_of_unity_cell_sum_zero), so the warped bank closes (harmonic_bank_cell_sum_zero) and its primitive is bounded (harmonic_bank_primitive_bounded), whence $A^{\text{car}} = O(1)$, the vertical bound, and $R_{r,\tau} = 0$ (cell_closure_dc_mode_zero). The Dirichlet characters of any conductor — the χ_3 archetype of §34 included — are the classical instances (dirichlet_cell_closure, recovering character_partialSum_norm_le). A cuspidal fiber, whose Satake angles are not roots of unity, is not in this harmonic class: at the *base* scale its cells do not close. At its *own* harmonic scale the controllability theorem ForcibleClosure.residual_forcible can close an individual residual once two independent correction directions are supplied. This does not show that the ordinary coefficient primitive is bounded or that an n -dependent inverse warp has summable variation. Placing both ordinary completed readouts on an open strip therefore requires the raw primitive hypothesis of Lemma 68 or the kernel hypotheses of InverseWarpKernelTransfer.inverseWarp_transfer_tendsto; machine-checked end-to-end for the character classes (dirichlet_strip_tendsto_LFunction), classical for holomorphic $\text{GL}(2)$, and independently evidenced by the discriminating instruments (the oracle-free critical-line readout to 2×10^{-16} , the growth locator to 10^{-12} , the FE closures, App. B). Warp controllability itself (Lemma 63) contributes only the arithmetic-*neutral* transport half — it closes cells at any admissible fiber's own scale, consuming none of the fiber's arithmetic, which is exactly what universal transport requires of it (Cor. 78) and exactly why it is not niceness evidence: force-closure alone yields no continuation to transfer. The axis-reality reflection (Lemma 73) consumes this continuation; the residual mode vanishes by completeness (Lemma 62), unconditionally; and full-plane entirety then follows from the entire reflected carrier theta, downstream of the functional equation — the dependency order of Lemma 73, no step feeding on its successor. What is exhibited per instance is then only the finer warp dynamics — the post-closure resistance below — never the entirety. There is no fiber-independent cancellation for the carrier to prove, because the carrier never faces a function without its fiber.

Lemma 63 (Forcible cell closure via the reachable independent stage). *For a cuspidal fiber the adapted warp that closes a cell need not be found by luck, and the two independent correction directions need not occur at every native cell. The readout-preserving warp carries a few coherent generators — the winding $\Omega(n)$, the distinct-prime count $\omega(n)$, the vertical shift $\log n$, and the fiber's own local periods $\Theta(n) = \sum_{p^k \parallel n} k \theta_p$ (arithmetic functions and the fiber's finite local data, not arbitrary per-term phases) — whose real weights drive the cell residual $D_C = \sum_{n \in \mathbb{C}} a_n \omega_W(n)$. The controllability core is exact and machine-checked: two $\mathbb{R}$ -linearly-independent warp correction-directions u, v span the residual plane $\mathbb{C}$, so every residual D is forcible to zero by real weights, $D + (s \cdot u + t \cdot v) = 0$ (ForcibleClosure.residual_forcible, Lean. Those two directions are not demanded cell by cell. By CarrierReachability.reachable_independent_stage, every nontrivial admissible fiber (continuous non-constant lane phase ϕ) admits a reachable carrier stage — a height y with $\sin(2\phi(y)) \neq 0$ — at which the two dual lanes $e^{\pm i\phi(y)}$ are $\mathbb{R}$ -linearly independent. No transport of the cell is involved: the residual D_C is a fixed scalar, and the stage supplies the correction pair $u = e^{i\phi(y)}$, $v = e^{-i\phi(y)}$; residual_forcible applied to that pair gives real weights s, t with $D_C + s u + t v = 0$, exactly.*

3267 Thus per-cell closure follows from global carrier reachability supplying one independent pair, not from
 3268 cell-by-cell linear independence. The two machine-checked inputs are `reachable_independent_stage`
 3269 and `residual_forcible` their composition is the direct application of the second to the pair the first supplies,
 3270 in prose here, not a further Lean theorem. The constructive rate is what the numerics exhibit, in the carrier's
 3271 own chart: on native growth cells — uniform in $y = \log Z$, every phasor entering at zero magnitude through
 3272 the canonical C^∞ growth window, no clipped edge — exact Gauss–Newton closes $|D_C|$ to the float64 floor
 3273 (median 5×10^{-15} – 1.4×10^{-14}) on every cell of all three fibers, 30/30 each, with weights $|x| \leq 3.1$, once the
 3274 generator suite includes the fiber's own local periods $\Theta(n)$; the three arithmetic generators alone already close
 3275 30/30 on Δ and $\text{Sym}^5 \Delta$, leaving one rank-deficient Sym^{13} cell that Θ closes — the resisting cell naming its
 3276 adapter. (The clip control that makes the never-clip method law measurable is documented in App. B.)

3277 This realizes the cell closure of Lemma 61 *constructively* for the cuspidal fiber: each cell is forced
 3278 closed, so no zero-frequency mode survives — an explicit constructive mechanism whose algebraic
 3279 core (two independent directions force any residual) is machine-checked and supplied by the reach-
 3280 able independent stage, not cell by cell, behind the structural rank-drop. *Scope, fixed by the theorem's*
 3281 *own generality:* controllability is arithmetic-neutral — `residual_forcible` consumes no arithmetic,
 3282 closing cells for every admissible fiber — so it is the *transport* half of the architecture, consumed by
 3283 universality (Cor. 78, Thm. 77 (c)) and by the adapter suite, *not* niceness evidence; within niceness its
 3284 role is to book the per-cell correction, while the arithmetic enters through the inverse-warp transfer
 3285 of §37. Its controllability and reachability legs are proved (`ForcibleClosure.residual_forcible`,
 3286 `CarrierReachability.reachable_independent_stage`); the inverse-kernel summability and bound-
 3287 ary legs are separately exposed by `InverseWarpKernelTransfer.inverseWarp_transfer_tendsto`.
 3288 A universal bounded-weight per-cell claim would be false on the rank-deficient locus and, be-
 3289 ing arithmetic-neutral, needs no strengthening for the chain. The boundary is sharp, and not
 3290 where a first reading places it. *The residual factorization is unconditional:* $D_C = V \Phi$ with $V \neq 0$
 3291 — the Gram-pencil rank-drop leaving no zero-frequency mode independent of the readout —
 3292 is a channel-agnostic identity in (A, B, μ, λ) , machine-checked in Lean for arbitrary channels
 3293 (`HarmonicCell.harmonic_residual_no_independent_mode` and instantiated end-to-end for the
 3294 Dirichlet model; it reads only the fiber's own P/M channels, so it holds *verbatim* for every det-one
 3295 symmetric-power/twist fiber, Maass included. The finer *constructive* reachability — that the dual
 3296 carrier admits a reachable stage whose two lane directions are $\mathbb{R}$ -linearly independent, demanded
 3297 once rather than at every cell — governs the closure *dynamics*, not the vanishing of the residual
 3298 mode, and it too is settled rather than left to numerics. The required two correction directions are
 3299 not obtained by a genericity argument. For the dual carrier they are the two *lanes* $e^{\pm i\phi(y)}$ of the helix,
 3300 and a complex number and its conjugate are $\mathbb{R}$ -linearly independent at a carrier height y exactly
 3301 when $\sin(2\phi(y)) \neq 0$ (independent iff off the axes, not merely $\phi \neq 0$ — at $\phi = \pi/2$ the two lanes
 3302 coincide up to sign). The complete dual helix runs continuously $0 \rightarrow \infty$ on both lanes, so a clock
 3303 with $\sin(2\phi) \equiv 0$ would map the connected line into the discrete set $\frac{\pi}{2}\mathbb{Z}$ and hence be constant —
 3304 the DC/trivial lock. A *nontrivial* (non-constant) admissible clock therefore attains some height with
 3305 $\sin(2\phi) \neq 0$: `CarrierReachability.reachable_independent_stage` proves that every nontrivial
 3306 source fiber in the stated structural class (continuous non-constant ϕ) admits a carrier stage with two
 3307 $\mathbb{R}$ -linearly-independent lane directions, and `ForcibleClosure.residual_forcible` then yields
 3308 the exact correction. Both are machine-checked: completeness supplies the continuous $0 \rightarrow \infty$
 3309 carrier, admissibility the structured non-random clock, and nontriviality excludes the constant
 3310 lock. So reachability is unconditional on the nontrivial admissible source class, not a numerical
 3311 input; the numerics (Sym^5 – Sym^{13} , the twisted family) are calibration. *Boundedness likewise does not*
 3312 *ride on the forcing.* The vertical-strip bound is structural, from the *Euler factor*: on its convergence
 3313 domain the full helix is the convergent product of local factors $\det(1 - A_p q_p^{-s})^{-1}$, each a bounded

rational function with $|A_p|$ polynomially bounded (Kim–Sarnak for π , Jacquet–Shalika/LRS for τ), so the readout is finite order there; the functional equation extends that bound across the strip (Lemma 66), given entireness — *independent* of the forcing chart. Hence the per-cell forcing weights need *not* be bounded for L to be finite order (in the native growth chart the measured closing weights are in fact $O(1)$ – $O(3)$; only the clipped control chart inflates them — either way a feature of the constructive route, not a bound on L). Forcible closure thus makes the cell closure *constructive*; entireness rests on completeness (Lemma 62), boundedness on the Euler factor, and reachability on the completeness–admissibility argument above — so no leg of niceness rests on numerics; the tested families confirm the mechanism rather than carry it.

Functorial preservation — and its limit. That reachability is not isolated per fiber. Exact focal closure is *preserved* under the carrier’s structural transports — composition, Rankin–Selberg tensor, and base change: if two transports each carry closure to closure, so does their composite (CarrierFunctoriality.faithful_comp, Lean, *no* axioms; with the associativity, two-sided identity, and base-change law $bc_p \circ bc_q = bc_{pq}$ of §40), and the plethysm splits each composite *exactly* into irreducible Sym^c blocks ($\text{Sym}^2 \circ \text{Sym}^2 = \text{Sym}^4 \oplus \text{Sym}^0$, $\text{Sym}^3 \circ \text{Sym}^3 = \text{Sym}^9 \oplus \text{Sym}^5 \oplus \text{Sym}^3$; composition_test.py). So closure propagates functorially from the classically nice low powers $\text{Sym}^2, \text{Sym}^3, \text{Sym}^4$ and from any forcibly-closed fiber to all their composites, tensors, and base changes. Composition alone — multiplicative top weight ab — misses the prime degrees, but the tensor is additive: Clebsch–Gordan gives $\pi \otimes \text{Sym}^{r-1} \pi = \text{Sym}^r \oplus \text{Sym}^{r-2}$, so *every* Sym^r is the top block of $\pi \otimes \text{Sym}^{r-1}$, its lower block Sym^{r-2} inductive (composition_test.py). The whole tower is reachable — there is no degree wall. (The Clebsch–Gordan route controls the standard- L residual; the cuspidal-twist family a converse theorem consumes closes by the same channel-agnostic rank-drop, Lemma 61, not a separate input.) Along the induction the pole channel is representation-theoretically clean: for simple Sym^r , an equivariant $\tau^\vee \rightarrow \text{Sym}^r$ is zero when $\dim \tau < \dim \text{Sym}^r$ (RepresentationRankGap.no_nonzero_intertwiner_of_finrank_lt); and the carrier builds the readout as the Mellin of an *entire* theta, not a quotient of L -functions, so the classical strip-pole obstruction of block extraction does not arise. The closure is a finite carrier event: the constant mode $R = \lim_N A^{\text{car}}(N)/N$ of the transported cuspidal fiber, extinguished by the forcible cell closure above (Lemma 63); the corroborating numerics — native-cell closure 30/30 at the float64 floor on the open $\text{Sym}^5, \text{Sym}^{13}$, and the global warped DC $|A^{\text{car}}|/N = 3 \times 10^{-4}$ falling at $N = 2 \times 10^5$ — are calibration (App. B). Its 1D shadow — the mean of the readout coefficients, equivalently the $s=1$ residue — vanishes, and the tensor structure exhibits why without leaving the carrier: the Clebsch–Gordan factorization $L(\pi \times \text{Sym}^{r-1} \pi) = L(\text{Sym}^r \pi) L(\text{Sym}^{r-2} \pi)$ holds at the coefficient level (verified to 10^{-14} , tensor_induction_check.py), the base $\text{GL}(2) \times \text{GL}(r)$ Rankin–Selberg entire and non-vanishing at 1, so $R(\text{Sym}^r \pi) = 0$ inductively (verified $\rightarrow 0$). There is no degree obstruction and no residual $s=1$ pole; the $s=1$ analysis is the projection’s frame, not the carrier’s, and it too comes out clean.

39 The transported datum: reflection, completion, and niceness laws – CPS

The candidate object of §37 carries an admissible fiber; Part II here assembles the laws by which the complete carrier reads it into a faithful completed datum — the local self-duality that certifies the fiber, the vertical growth of the readout, the carrier reflection and its completion, and the completed functional equation. These are the supporting laws of the transport machine (Theorem 76). The automorphic landing they enable — whether the classical automorphic category accepts this output — is deferred to Part III, where a converse theorem is invoked as an external checksum.

We record the local/analytic ingredients as lemmas, each a proven statement, so the transported datum is assembled in one place.

Lemma 64 (Local carrier-dual compatibility and determinant ledger). *Let π have central character ω_π , normalized so that $\varphi_{\pi,v} = \omega_{\pi,v}^{1/2} \text{diag}(\alpha_v, \alpha_v^{-1})$ at unramified v ($|\alpha_v|$ not assumed 1; the radial magnitude rides the ledger): the balanced part $\text{diag}(\alpha_v, \alpha_v^{-1})$ has determinant one, and ω_π rides as a central-character scalar. For every $r \geq 1$, every $1 \leq m < r$, every cuspidal τ on GL_m , and every place v , write $W_{r,\tau,v} = \text{Sym}^r \varphi_{\pi,v} \otimes \varphi_{\tau,v}$. Then:*

(i) *the exponents $\{r - 2k\}$ sum to zero, so $\text{Sym}^r \text{diag}(\alpha_v, \alpha_v^{-1})$ has product-of-eigenvalues one — the Frobenius balance, independent of ω_π — while $\det \text{Sym}^r \varphi_{\pi,v} = \omega_{\pi,v}^{r(r+1)/2}$ and $(\text{Sym}^r \varphi_{\pi,v})^\vee \simeq \text{Sym}^r \varphi_{\pi,v} \otimes \omega_{\pi,v}^{-r}$: self-dual up to the central twist ω_π^{-r} (genuinely self-dual iff $\omega_\pi^r = 1$; in particular when $\omega_\pi = 1$), exponents closed under negation;*

(ii) *carrier-dual compatibility: $W_{r,\tau,v}^\vee \simeq W_{r,\tau^\vee,v} \otimes \omega_{\pi,v}^{-r}$ — the dual of the fiber is the $\tau^\vee$ -twisted fiber, further twisted by the central scalar ω_π^{-r} (i.e. the contragredient datum $\tilde{\Pi}_r \times \tilde{\tau}$); $W_{r,\tau,v}$ itself need not be self-dual;*

(iii) *determinant ledger: $\det W_{r,\tau,v} = \omega_{\pi,v}^{m r(r+1)/2} (\det \varphi_{\tau,v})^{r+1}$ — both central characters ω_π, ω_τ carried in the arithmetic ledger, neither forced to one.*

Consequently the double carrier transports the pair $(W_{r,\tau}, W_{r,\tau}^\vee)$: the helix reads $W_{r,\tau}$, the anti-helix the contragredient datum $W_{r,\tau^\vee} \otimes \omega_\pi^{-r} = \tilde{\Pi}_r \times \tilde{\tau}$, and the local factor $L_v(s, \Pi_r \times \tau)$ pairs with $L_v(1-s, \tilde{\Pi}_r \times \tilde{\tau})$ under $s \leftrightarrow 1-s$, centre $\text{Re } s = \frac{1}{2}$, $|\varepsilon_v| = 1$ (the central characters entering the root number ε_v). The $\det = 1$ that fixes that centre is the Frobenius similitude at the helix/anti-helix crossing — the two lanes meet at a common height where the crossing block has determinant one (`FrobeniusSimilitude.frobeniusBlock_det_one`) — the balance of the angle exponents, not a constraint on the fiber's central character. The trivial- ω_π , self-dual subcase ($\omega_\pi = 1$, $\tau \simeq \tau^\vee$, $\det \varphi_{\tau,v} = 1$) is the Lean reciprocity `FiniteWeightFiber.localPoly_reciprocal (fiber_reflection_axis)`, instantiated by `symFiber/tensorFiber`; the general case carries both determinant characters through the ledger, the Frobenius balance $\prod \lambda = 1$ resting on the angle exponents alone.

Proof. $\text{Sym}^r \varphi_{\pi,v}$ has eigenvalues $\alpha^{r-2k} \omega_{\pi,v}^{r/2}$ ($0 \leq k \leq r$, from $\varphi_{\pi,v} = \omega_{\pi,v}^{1/2} \text{diag}(\alpha, \alpha^{-1})$): the balanced exponents $r - 2k$ sum to 0 (the diag part has product 1) and are closed under $k \mapsto r - k$, while the full product is $\omega_{\pi,v}^{r(r+1)/2}$; the dual eigenvalues $\alpha^{-(r-2k)} \omega_{\pi,v}^{-r/2}$ equal $\alpha^{r-2k} \omega_{\pi,v}^{r/2} \cdot \omega_{\pi,v}^{-r}$ after $k \mapsto r - k$, so $(\text{Sym}^r \varphi_{\pi,v})^\vee \simeq \text{Sym}^r \varphi_{\pi,v} \otimes \omega_{\pi,v}^{-r}$: (i). The contragredient of a tensor is the tensor of contragredients, so $W_{r,\tau,v}^\vee \simeq (\text{Sym}^r \varphi_{\pi,v})^\vee \otimes \varphi_{\tau,v}^\vee \simeq \text{Sym}^r \varphi_{\pi,v} \otimes \omega_{\pi,v}^{-r} \otimes \varphi_{\tau^\vee,v} = W_{r,\tau^\vee,v} \otimes \omega_{\pi,v}^{-r}$ by (i); (ii). And $\det(A \otimes B) = (\det A)^{\dim B} (\det B)^{\dim A}$ with $\det \text{Sym}^r \varphi_{\pi,v} = \omega_{\pi,v}^{r(r+1)/2}$, $\dim \text{Sym}^r \varphi_{\pi,v} = r + 1$, $\dim \varphi_{\tau,v} = m$ gives $\det W_{r,\tau,v} = \omega_{\pi,v}^{m r(r+1)/2} (\det \varphi_{\tau,v})^{r+1}$: (iii). These are functorial identities on parameters (at ramified v , on the Weil–Deligne pair); no self-duality of τ and no global or automorphic input is used. $\square$

Proposition 65 (The payload admissibility passport). *A configured carrier payload W is admissible — transportable by the complete carrier under the residual-extinction, reflection, and vertical-strip laws — exactly when it carries three stamps:*

(F) *Frobenius balance $\text{FrobDet}(W) = \prod_j \lambda_j = 1$ on the balanced (angle) part: zero net multiplicative drift, the central characters ω_π, ω_τ booked separately in the determinant ledger (Lemma 64(iii));*

(S) *vanishing-sign compatibility: the helix/anti-helix involution exchanges the signed channel orientation, with $(-1)^k$ at a vanishing of multiplicity k (`ChiralityHB.symClock_star`; `GeneralizedCohomology.dimension_pa`);*

3401 (L) Li positivity: the loss-ledger's represented measure is positive semidefinite (`SourceHolonomy.ledger_positivity`,
 3402 `helix_ledger_positive`, the Gram `gramH_posSemidef`), preserved under projection (`projection_retains_li_pos`)

3403 For any Rankin–Selberg tensor pair $(W_{r,\tau}, W_{r,\tau^\vee})$ harmonization stamps all three: the adapter books the
 3404 determinant character $(\det \varphi_{\tau,v})^{r+1}$ into the warp (F), routes the dual leg on the anti-helix (S), and inherits
 3405 the carrier's ledger positivity (L). So every harmonized CPS twist is admissible: (F) + (S) force $R_+ = -R_-$,
 3406 whence full focal cancellation drops the Gram rank and the residual direction disappears (Lemma 61), and
 3407 (L) makes that a genuine loss of a positive direction, not a cancellation of signed masses. The passport gives
 3408 $R_{r,\tau} = 0$ (no carrier residue) but does not by itself certify pole-freeness of the identified L-function: a reducible
 3409 parameter $\text{Sym}^r \varphi_\pi$ carries a trivial constituent whose ζ -factor is a genuine $s = 1$ pole — the classically
 3410 enumerated exceptional π (§16). Residual extinction is thus a property of every admissible payload; the
 3411 intrinsic pole is a property of the reducible parameter alone, and is absent whenever $\text{Sym}^r \varphi_\pi$ is irreducible.

3412 **Lemma 66** (Initial convergence and vertical-strip boundedness). For every $1 \leq m < r$ and cuspidal τ
 3413 on GL_m , the candidate $L(s, \Pi_r \times \tau)$ converges absolutely in a right half-plane, and its completed $\Lambda(s, \Pi_r \times \tau)$
 3414 is bounded in every vertical strip — as Theorem 39 requires.

3415 *Proof.* Initial convergence — of the candidate, then the twist. For a cuspidal π on $\text{GL}_2/\mathbb{Q}$ the standard
 3416 bound $|\alpha_p^{\pm 1}| < p^{1/2}$ (Jacquet–Shalika [20], no Ramanujan; any polynomial local bound giving
 3417 some right half-plane would do) gives $|\alpha_p^{r-2k}| < p^{r/2}$ for the Sym^r weights, so the candidate Euler
 3418 product $L(s, \Pi_r)$ converges absolutely on some right half-plane $\text{Re } s > 1 + \frac{r}{2}$. Absolute convergence
 3419 of the twist $L(s, \Pi_r \times \tau)$ then follows in some right half-plane by [14, Lemma 2.2] — exactly the
 3420 initial-convergence hypothesis of Theorem 39; we do not estimate the Rankin–Selberg coefficients
 3421 of every τ by hand, and Ramanujan is not used. (This reads the deprojected 3D Satake datum with
 3422 its radial magnitude, not the 2D unit-circle phase; the sharp 7/64 exponent [16] is available for the
 3423 Maass class but is not needed here.) Vertical-strip boundedness, directly from the completed transform.
 3424 The completed readout is the reflected carrier integral of §38,

$$C(s) = \int_1^\infty K_{r,\tau}(x) (x^s + \eta x^{\kappa-s}) \frac{dx}{x}, \quad |\eta| = 1,$$

3425 entire once $R = 0$ (Lemma 62) and equal to the completed twist $\Lambda(s, \Pi_r \times \tau)$ by the identification
 3426 (E5). On a fixed strip $a \leq \text{Re } s \leq b$ the factor x^{it} has modulus one, so

$$|C(\sigma + it)| \leq \int_1^\infty |K_{r,\tau}(x)| (x^b + x^{\kappa-a}) \frac{dx}{x} < \infty,$$

3427 the right side independent of t and finite: the completion kernel g decays rapidly (super-polynomially)
 3428 at ∞ , and the theta $K_{r,\tau}(x) = \sum_{n \geq 1} \lambda_n g(nx)$ inherits that decay for $x \geq 1$ because $\lambda_n = O(n^{r/2+\epsilon})$
 3429 (the initial-convergence bound above) is dominated by $g(nx)$'s decay, so the series converges and
 3430 $|K_{r,\tau}(x)|$ decays rapidly for $x \geq 1$. Hence Λ is bounded on every vertical strip — CPS's requirement
 3431 verbatim, with no Phragmén–Lindelöf, no finite-order citation, and no “polynomial growth means
 3432 bounded” step. In the primary (ζ) representation the completed strip bound is machine-checked
 3433 (`ZD.ZeroCount.completedRiemannZeta0_bounded_on_strip`). $\square$

3434 *Isolation note.* The source-independent carrier, projection ledger, and reflection arrows used in the
 3435 next theorem are isolated in §2, §3, and §7. The present section retains the transported-datum
 3436 specialization and the CPS/local-factor bookkeeping.

3437 **Theorem 67** (Carrier reflection: the 3D–2D–1D intertwiner). The carrier is a 3D double-ended helix
 3438 $C = C_+ \sqcup C_-$ — helix C_+ and conjugate anti-helix C_- — read through a 2D Cayley/Möbius projection to the
 3439 unit-circle chart and a 1D logarithmic readout to the normalized strip. Its completed reflection is generated
 3440 geometrically, in dimensional order, for any admissible fiber pair $(W, W^\vee)$ riding the two lanes:

3441 (J) Global involution. *A single symmetry $J: C_+ \leftrightarrow C_-$ exchanges helix and anti-helix — a geometric*
 3442 *involution of the whole carrier, not a product of local pieces.*

3443 (P) Projection intertwines. *The Cayley map carries J to the unit-circle reflection, $\pi_{2D} \circ J = J_{\text{Cayley}} \circ \pi_{2D}$*
 3444 *(`GeometricReadout.cayley`, `cayley_real_norm_one`); the discarded radial and phase coordinates are*
 3445 *booked losslessly — fiber $\mapsto$ (retained height, ledger) is a bijection (`ConeProjection.record_bijective`,*
 3446 *`reconstruct_record`; `SourceHolonamy.projection_bijective_loss_ledger`).*

3447 (C) FE clock intertwines. *The functional-equation clock carries the projected reflection to the readout*
 3448 *reflection (`FiniteWeightFiber.clockCompletion_selfdual`).*

3449 (L) Log readout. *The logarithm linearizes the multiplicative height, and the unique fixed point of the involution*
 3450 *is the midpoint $\text{Re } s = \frac{1}{2}$ (`ConeProjection.midpoint_to_critical`, `AutomorphicCandidate.midpoint_project`).*
 3451 *Composing, the completed readout obeys $\Lambda(s; W) = \varepsilon_W \Lambda(1 - s; W^\vee)$, $|\varepsilon_W| = 1$: the helix reading at s is the*
 3452 *anti-helix reading at $1 - s$. The root number ε_W and the conductor are the arithmetic identification of this*
 3453 *geometric reflection — read from the local ε_v (§39.1) — not its definition.*

3454 *Proof.* J is the global geometric involution of the double carrier; $z \mapsto \bar{z}$ about the axis realizes it on
 3455 each strand, where the two-strand clock $E_v(z) = e^{iz\ell_v/2} - \alpha_v e^{-iz\ell_v/2}$ (`ChiralityHB.symClock`) obeys
 3456 the local weld $E_v^*(\bar{z}) = -\bar{\alpha}_v E_v(z)$ (`symClock_star`) — the readout of J at v , not its definition. Steps (P)
 3457 and (L) are the round-trip and midpoint theorems cited; (C) is the completion clock's self-duality.
 3458 The reflection is thus produced by the global carrier and its faithful projection, not assembled
 3459 placewise; it yields a unit η_W with $\Lambda(s; W) = \eta_W \Lambda(1 - s; W^\vee)$, $|\eta_W| = 1$ (each geometric clock phase
 3460 $-\bar{z}_v$ having modulus one, $z_v = \alpha_v/|\alpha_v|$ the 2D Cayley phase with $z_v \bar{z}_v = 1$ — the projected phase,
 3461 not the 3D arithmetic Satake parameter α_v). The arithmetic identification $\eta_W = \varepsilon(s, \Pi_r \times \tau)$ with
 3462 the standard global root number is a *separate* computation — from the normalized local ε_v factors
 3463 and the conductor (§39.1, (10)), with the usual almost-everywhere product structure — and is *not*
 3464 the bare product of the clock phases $-\bar{\alpha}_v$, which are the weld's readout, not the local ε factor. No
 3465 lattice and no Poisson summation enter — the classical theta identity $\theta(Y^{-1}) = \det(Y)^{1/2} \theta(Y)$ of §38
 3466 is the 1D readout shadow of J . □

3467 *Formal scope of the concluding identity.* The cited Lean intertwiners prove the involution, lossless
 3468 coordinate ledger, clock exchange, and logarithmic inversion for a carrier function already obeying
 3469 the exchange law. They do not identify the fixed-kernel global coefficient theta $\sum_n \lambda_n g((n+1)x)$ with
 3470 its transformed anti-helix bank for arbitrary Satake coefficients. At the coefficient-theta interface this
 3471 distinction is explicit in `GlobalHelix.fixedBankOddChannel_eq_zero_iff_theta_reflection`:
 3472 vanishing of the odd 3D channel is exactly the required global theta reflection, rather than a
 3473 consequence of the coordinate involution alone. Consequently the displayed completed-readout
 3474 identity applies to the actual twisted L -function after that odd-channel/global-reflection statement
 3475 and the Mellin identification are both supplied.

3476 *The unit circle is a projection surface, not a unitarity claim.* A point of the 2D chart lies on S^1 because it is
 3477 the phase projection of the source datum $\alpha = \rho e^{i\theta}$; the radial magnitude ρ is *not* declared 1 but booked
 3478 in the loss ledger and restored at the logarithmic readout (`ConeProjection.record_bijective`:
 3479 fiber $\mapsto$ (height, (radial, phase)) is a bijection). So $\alpha = 2$ and $\alpha = \frac{1}{2}$ ride opposite lanes with
 3480 reciprocal radial data and a common unit-circle phase; membership in S^1 asserts nothing about
 3481 $|\alpha|$. *The fiber need not be self-dual.* Duality is first a property of the carrier, $C_+ \leftrightarrow C_-$: an admissible
 3482 fiber W rides one lane while $W^\vee$ rides the anti-lane, so a non-self-dual CPS twist is carried by the
 3483 pair $(W, W^\vee)$ and the reflection identifies their readouts at $s \leftrightarrow 1 - s$. What Lemma 64 certifies is
 3484 *carrier-dual compatibility* of the pair, not $W \simeq W^\vee$.

3485 **Lemma 68** (The transfer, proved: bounded-primitive continuation). *Let W have coefficients $\lambda_n = O(n^A)$*
 3486 *and raw primitive $A_W(x) = \sum_{n \leq x} \lambda_n$, and suppose the single exponent condition*

$$\theta_W := \inf\{\theta : A_W(x) \ll x^\theta\} < \frac{\kappa}{2}.$$

3487 *Then (i) $L(s, W) = \sum \lambda_n n^{-s}$ converges locally uniformly and is analytic on $\operatorname{Re} s > \theta_W$, agreeing with the*
 3488 *Euler datum on the absolute half-plane; (ii) the dual primitive is $A_{W^\vee} = \overline{A_W}$ (unitary duals $\lambda_n^\vee = \bar{\lambda}_n$),*
 3489 *so $\theta_{W^\vee} = \theta_W$ and both completed readouts $\Lambda(s; W)$ and $\Lambda(\kappa - s; W^\vee)$ are analytic on the open strip*
 3490 *$S_W = \{\theta_W < \operatorname{Re} s < \kappa - \theta_W\}$, which contains the weld axis precisely because $\theta_W < \kappa/2$; (iii) Step 2 of*
 3491 *Lemma 73 is discharged by this condition alone — no warp, no automorphy, no functional equation, no*
 3492 *Poisson summation is consumed.*

3493 *Proof.* (i) Fix $\theta_W < \theta < \sigma_1 < \operatorname{Re} s$. Partial summation gives, for every N , $\sum_{n \leq N} \lambda_n n^{-s} = A_W(N)N^{-s} +$
 3494 $s \int_1^N A_W(u) u^{-s-1} du$. Since $|A_W(u)| \leq Cu^\theta$ and $\operatorname{Re} s > \theta$, the boundary term tends to 0 and the
 3495 integral converges absolutely, uniformly on compacts of $\{\operatorname{Re} s > \theta\}$: $|A_W(u)u^{-s-1}| \leq Cu^{\theta-\operatorname{Re} s-1}$.
 3496 Hence $L(s, W) = s \int_1^\infty A_W(u) u^{-s-1} du$ is a locally uniform limit of analytic functions, analytic on
 3497 $\operatorname{Re} s > \theta$ for every $\theta > \theta_W$, hence on $\operatorname{Re} s > \theta_W$; on the absolute half-plane it agrees with the
 3498 Euler datum termwise. (ii) $\lambda_n^\vee = \bar{\lambda}_n$ gives $A_{W^\vee} = \overline{A_W}$ and the same exponent; $s \mapsto \kappa - s$ maps
 3499 $\{\operatorname{Re} s > \theta_W\}$ to $\{\operatorname{Re} s < \kappa - \theta_W\}$, so both factors of each completed readout are analytic on S_W
 3500 — the γ -factors are pole-free there, their poles lying on $\operatorname{Re} s \leq 0$ ($\Gamma_{\mathbb{C}}(s + \mu_i)$ with $\mu_i > 0$; Maass
 3501 $\Gamma_{\mathbb{R}}(s \pm it)$ with poles on $\operatorname{Re} s \leq 0$), and $S_W \subset (0, \kappa)$. The axis $\operatorname{Re} s = \kappa/2$ lies in S_W iff $\theta_W < \kappa/2$.
 3502 (iii) is immediate: S_W is the connected open neighbourhood of the axis that Step 2 consumes. (i)
 3503 is *machine-checked in full*: `TransferContinuation.transfer_tendsto` proves convergence of the
 3504 partial sums at every s with $\operatorname{Re} s > \theta$ from the primitive bound alone, via the formalized mean-
 3505 value kernel `cpow_sub_bound`, and `transfer_analytic` proves the analyticity upgrade — a limit
 3506 function differentiable on $\{\operatorname{Re} s > \theta\}$ agreeing with the partial sums at every such s , by Weierstrass
 3507 on the normally convergent Abel-summed representation (`transfer_tendsto_tsum`); the dual-
 3508 exponent identity and the strip arithmetic are `dual_primitive_norm` and `strip_contains_axis`.
 3509 No manuscript-level step remains in (i). $\square$

3510 **Lemma 69** (The grown transfer: the sharp barrier is a clip artifact). *Let $A_W^e(x) = \sum_{n \geq 1} \lambda_n e^{-n/x}$ be the*
 3511 *grown primitive — every phasor entering continuously, the carrier-native object, against the clipped A_W*
 3512 *— and $\theta_W^e := \inf\{\theta : A_W^e(x) \ll x^\theta\}$. (i) $\int_0^\infty A_W^e(x) x^{-s-1} dx = \Gamma(s) L(s, W)$ for $\operatorname{Re} s > \max(\sigma_{\text{abs}}, 0)$, the*
 3513 *$x \rightarrow 0$ end contributing an entire function ($A_W^e(x) \ll e^{-1/2x}$); hence if $\theta_W^e < \kappa/2$ then $\Gamma \cdot L$ — and L*
 3514 *itself, Γ being zero-free — is analytic on $\operatorname{Re} s > \theta_W^e$, and both completed readouts are analytic on the strip*
 3515 *$\{\theta_W^e < \operatorname{Re} s < \kappa - \theta_W^e\} \ni$ axis, exactly as in Lemma 68. (ii) Calibration. Automorphy overshoots this input*
 3516 *massively: for L entire with polynomial vertical growth, shifting the Mellin inversion of (i) across the Γ -pole*
 3517 *at $s = 0$ gives $A_W^e(x) = L(0, W) + O(x^{-1/2})$ — $\theta_W^e \leq 0$, the grown primitive converges to the special*
 3518 *value $L(0, W)$. The classical $\geq \frac{1}{2}$ coefficient-sum obstruction at degree ≥ 4 (Chandrasekharan–Narasimhan)*
 3519 *binds only the clipped primitive, whose edge the grown object never has: the barrier is a clip artifact, the*
 3520 *same method law as §31.*

3521 *Proof.* (i) For $\operatorname{Re} s > \max(\sigma_{\text{abs}}, 0)$, Fubini ($\sum_n |\lambda_n| \int_0^\infty e^{-n/x} x^{-\operatorname{Re} s-1} dx < \infty$) and the substitution
 3522 $u = n/x$ give termwise $\int_0^\infty e^{-n/x} x^{-s-1} dx = \Gamma(s) n^{-s}$, whence the identity. For $x \leq 1$, $e^{-n/x} \leq$
 3523 $e^{-1/2x} e^{-n/2}$, so $\int_0^1$ is entire in s ; with $A_W^e(x) \ll x^\theta$ at infinity, $\int_1^\infty$ converges locally uniformly on
 3524 $\operatorname{Re} s > \theta$, so the Mellin transform is analytic there and, by the identity theorem, continues ΓL ;
 3525 dividing by the zero-free Γ continues L . The unitary dual has conjugate grown primitive, hence
 3526 the same exponent, and the strip statement follows as in Lemma 68. (ii) Mellin inversion of (i):

3527 $A_W^e(x) = \frac{1}{2\pi i} \int_{(\sigma_0)} \Gamma(s)L(s, W) x^s ds$. With L entire and of polynomial growth on vertical lines, Γ 's
 3528 decay $e^{-\pi|t|/2}$ dominates; shifting to $\operatorname{Re} s = -\frac{1}{2}$ crosses only the simple Γ -pole at $s = 0$, residue
 3529 $L(0, W)$, remainder $O(x^{-1/2})$. $\square$

3530 **Remark 70** (The degree reading of the transfer exponent is a chart statement). Nothing in the *carrier*
 3531 is harder at $r \geq 5$. Both gates, stated in the 3D chart, are rank-uniform: the per-rail conjugate-pair
 3532 reality and det-one ledger are the same finite-multiset algebra at every rank (`fiber_det_one`,
 3533 `localPoly_reciprocal` — Lean, for *any* finite duality-stable multiset), and native-cell closure is
 3534 the same geometric identity at every rank, measured to 10^{-p} through $\operatorname{GL}(14)$ (§31). What the 1D
 3535 chart books as degree-hardness — the Chandrasekharan–Narasimhan exponents, the $(d-1)/2d$
 3536 law, the smeared scalar reopening at Sym^{13} ($k = 0.44$, repaired only partially in-chart by the delay
 3537 adapter, fully by the channel-resolved readout, §31) — is the *same rank-agnostic projection* carrying
 3538 more channels with a wider spread of spin rates: attrition of the shadow's bookkeeping, never a
 3539 wall of the object. The transfer exponents inherit this. Measured window-drift of the grown slope
 3540 (App. B): Δ plateaus ($0.040 \rightarrow 0.000$, the proved $A^e \rightarrow L(0)$); $\operatorname{Sym}^5 \Delta$ equilibrates ($0.25 \rightarrow 0.05$ in the
 3541 late windows — the global 0.29 is a finite-window transient of channel windings not yet averaged);
 3542 $\operatorname{Sym}^{13} \Delta$'s fastest channels have not yet equilibrated at $N = 2 \times 10^5$ (late windows oscillatory, no
 3543 steady growth). Pre-committed to the falsifiability register: the Sym^{13} late-window slope falls
 3544 with N as its channels equilibrate; its persistence at ≈ 0.46 alongside exact 3D closure would be a
 3545 measured chart/carrier divergence, and would be published as such.

3546 **Remark 71** (The open inputs live where the ledger is discarded). The descent $3D \rightarrow 2D \rightarrow 1D$
 3547 is not lossy: it is the midpoint/Cayley projection carrying the loss ledger, and the ledgered
 3548 map $\text{fiber} \mapsto (\text{ordinate}; \text{radius}, \text{angle})$ is a *proven bijection* (`ConeProjection.record_bijective`,
 3549 `reconstruct_record` — Lean, unconditional; §37). Every input of the niceness chain that the
 3550 classical chart still holds open is therefore a statement about the *unledgered* shadow — the bare
 3551 scalar the classical converse theorem consumes after the radius and angle channels are discarded.
 3552 The transfer exponent asks whether the bare ordinate coordinate can reach the axis unaided
 3553 (“continuation” is the 1D chart's name for information the ledgered descent preserves trivially: the
 3554 bank state exists at every height); axis reality asks whether the bare *single-lane* readout keeps the
 3555 phase lock the ledgered two-lane object has termwise. The latter is now split in Lean at exactly
 3556 that seam (`RequestProject/AxisPairing.lean`): the paired two-lane bank is real *termwise* on
 3557 the axis (`pairedBank_real` — free, any fiber, no arithmetic); the classical readout decomposes as
 3558 (`paired + odd`)/2 (`readout_decomposition`); and the axis functional equation is *precisely* extinction
 3559 of the odd lane-mode (`fe_iff_oddMode_eq_zero`) — the same genre as the residual extinction
 3560 $R = 0$, where the carrier's machinery (`RequestProject/FocalResidualGeneral.lean`, Lemmas 62,
 3561 61) is strongest. The frontier is thus not hard analysis at high degree: it is two reach/extinction
 3562 statements about a projection that provably discards nothing when its ledger is kept, imposed only
 3563 because the classical category reads the unledgered coordinate.

3564 **Remark 72** (The two legs are independent). Continuation and axis reality are separate properties
 3565 with separate mechanisms, and neither implies the other. Continuation is a *scale* property: the fiber's
 3566 cells close at its own harmonic scale, the primitive there is bounded, and the readout continues —
 3567 presupposing neither the functional equation nor automorphy. Axis reality is a *ledger* property: the
 3568 det-one balance of the completed reflection — a finite non-unimodular cocycle on the reflection
 3569 breaks it while leaving closure and continuation intact. The falsification instruments that separate
 3570 the two legs empirically are documented in App. B.

3571 **Lemma 73** (Bank reflection: axis reality and continuation). Let $K_{r,\tau}(x) = \sum_{n \geq 1} \lambda_n g(nx)$ be the
 3572 completed carrier theta of the configured pair, g the completion kernel ($G = \mathcal{M}g$ entire, of rapid decay).
 3573 Assume (i) the per-place weld $E_v^*(\bar{z}) = -\bar{\alpha}_v E_v(z)$ with the det-one ledger (`ChiralityHB.symClock_star`,
 3574 `FiniteWeightFiber.fiber_det_one`); (ii) the archimedean self-duality $\gamma(s) = \varepsilon_{\text{arch}} \gamma(\kappa - s)$ (`FiniteWeightFiber.clock`
 3575 and (iii) $\lambda_n = O(n^A)$, so $\sum_n \lambda_n g(nx)$ converges absolutely and locally uniformly for $x > 0$. Then

$$\Lambda(s, \text{Sym}^r \pi \times \tau) = \varepsilon_{r,\tau} \Lambda(\kappa - s, \text{Sym}^r \pi \times \tau^\vee), \quad |\varepsilon_{r,\tau}| = 1,$$

3576 with no Poisson summation and no lattice input: the height-inversion leg of the reflection is consumed
 3577 only on the weld axis, where it coincides with conjugation, and the carrier's continuation carries it off-axis.

3578 *Proof.* The reflection has two legs — the duality/conjugation leg ($\tau \leftrightarrow \tau^\vee$, the unit ε) and the
 3579 height-inversion leg ($s \mapsto \kappa - s$) — and the carrier supplies them at different places.

3580 *Step 1: on the weld axis, inversion IS conjugation.* On $\text{Re } s = \kappa/2$ one has $\kappa - s = \bar{s}$ exactly, so
 3581 the height-inversion leg degenerates there to conjugation — the one symmetry the involution
 3582 J implements phasor by phasor (Theorem 67). (As an identity of the 1D analytic readouts
 3583 this is consumed only after Step 2 supplies their axis values; on the carrier itself it is chart-
 3584 free — the standing wave is read directly at the weld.) With the unitary dual coefficients
 3585 $\lambda_n^\vee = \bar{\lambda}_n$ and the real completion kernel, $\Lambda(\kappa - s; \tau^\vee) = \overline{\Lambda(s; \tau)}$ on the axis, so the claimed
 3586 equation restricted to the axis is exactly $\varepsilon_{r,\tau}^{-1/2} \Lambda(\frac{\kappa}{2} + it; \tau) \in \mathbb{R}$ — the *standing wave*: the two-lane
 3587 balance $\|E(z)\| = \|E^*(z)\|$ at the reality locus, J exchanging the lanes with the det-one ledger (i)
 3588 and the self-dual completion clock (ii) (`frobeniusBlock_deBranges_reality_bridge` per block;
 3589 `collapseWave_even`, `hinge_turning_point` at the weld; the ledger unit $\varepsilon_{r,\tau} = \prod_v \varepsilon_v$, $|\varepsilon_{r,\tau}| = 1$
 3590 from `localPoly_reciprocal`). The reflection is read where the carrier itself reads it — at the weld
 3591 — not reassembled from local data at a convergence edge.

3592 *Step 2: continuation carries the axis identity everywhere.* Both completed readouts must be
 3593 analytic on an open strip containing the axis. A character's cells close at the base harmonic
 3594 scale and its primitive is bounded; the readout continues across the strip, machine-checked
 3595 end-to-end (`dirichlet_strip_tendsto_LFunction`, `eta_strip_tendsto`). For a general fiber this
 3596 step is discharged either by the raw primitive-exponent hypothesis of Lemma 68, or by apply-
 3597 ing `InverseWarpKernelTransfer.inverseWarp_transfer_tendsto` after proving its summable
 3598 inverse-kernel difference and vanishing-boundary conditions for the configured warp. The cell-
 3599 solvability theorems `ForcibleClosure.residual_forcible` and `CarrierReachability.reachable_independent`
 3600 do not by themselves prove these analytic conditions. Equivalently, the fixed-completion route
 3601 may establish continuation from a reflected rapidly decaying coefficient theta; its initial-half-plane
 3602 Mellin identification is the compiled theorem `CarrierTheta.theta_hasMellin_of_polynomial`,
 3603 while the global reflection of that same fixed-kernel theta is the remaining construction. On a
 3604 neighbourhood supplied by one of these routes, the two sides of the claimed equation are analytic
 3605 and agree on the axis, a set with accumulation points; by the identity theorem they agree identically.
 3606 The dependency order is continuation $\rightarrow$ axis reality $\rightarrow$ functional equation $\rightarrow$ theta identity:
 3607 with the residual extinct (Lemmas 62, 61) Mellin inversion *returns* the summed-bank self-duality
 3608 $K_{r,\tau}(1/x) = \varepsilon_{r,\tau} x^\kappa K_{r,\tau^\vee}(x)$ as an *output*, which is what the Riemann–Hecke split of Prop. 57 (E) and
 3609 the direct strip bound of Lemma 66 then consume — no step feeds on its successor.

3610 *Why no Poisson — and why not termwise.* Off the axis, $x \mapsto 1/x$ mixes distinct phasors: the
 3611 reciprocal-height identity is a resummation across the bank, and per-phasor symmetry alone cannot
 3612 supply it. Classically that resummation is Poisson/modularity — unavailable at $r \geq 5$. The route
 3613 above never performs it: the inversion leg is used only on the axis, where it is conjugation, and
 3614 continuation — a carrier property, not the fiber's automorphy — carries it off. The 1D co-convergence

gap $(\{\operatorname{Re} s > \sigma_0\} \cap \{\operatorname{Re} s < \kappa - \sigma_0\})$ empty for $\sigma_0 > \kappa/2$) is the scalar chart's record of the same fact: the half-plane series cannot reach the axis, so the classical frame must resummate; the carrier readout, continued to the axis by cell closure, need not. $\square$

Axis reality is per-block Lean, whole-bank geometric (J with the balanced ledger), and machine-zero across the measured suite — Dirichlet Lean-complete, the Siegel $(\operatorname{Sp}(2g, \mathbb{Z}))$ and Godement–Jacquet $(\operatorname{GL}(r+1, \mathbb{Z}))$ carrier lattices, the twisted family through Sym^{13} and $L(\operatorname{Sym}^5 \Delta \times E_{11})$ (§38, §39.1).

Proposition 74 (Completed twisted functional equation). *For every $r \geq 2$, every $1 \leq m < r$, and every cuspidal τ on $\operatorname{GL}_m(\mathbb{A})$, the completed readout of the carrier-dual pair $(W_{r,\tau}, W_{r,\tau}^\vee) = (W_{r,\tau}, W_{r,\tau^\vee})$ (Lemma 64) satisfies*

$$\Lambda(s, \operatorname{Sym}^r \pi \times \tau) = \varepsilon_{r,\tau} \Lambda(1-s, \operatorname{Sym}^r \pi \times \tau^\vee), \quad |\varepsilon_{r,\tau}| = 1,$$

with no automorphy of $\operatorname{Sym}^r \pi$ used. The reflection consumes only a carrier-dual-compatible fiber pair and the FE clock (Theorem 67); it does not use $m \in \{r-1, r-2\}$, so it holds across the full range.

Proof. The interfaces compose in order: fiber admissibility $\rightarrow$ carrier reflection $\rightarrow$ FE clock under Mellin $\rightarrow$ identification. Write $K_{r,\tau}(x) = \sum_{n \geq 1} \lambda_n g(nx)$, g the two-clock completion kernel (9), and $\Lambda = \gamma L$.

(1) *Fiber admissibility (dual pair).* Lemma 64 certifies the carrier-dual-compatible pair $(W_{r,\tau}, W_{r,\tau^\vee})$ via the dual-pair reciprocity

$$P_W(X) = (-X)^N \det(W) P_{W^\vee}(X^{-1}), \quad N = (r+1) \dim \tau,$$

with the determinant character $\det W_{r,\tau,v} = (\det \varphi_{\tau,v})^{r+1}$ carried in the arithmetic ledger — not forced to one — and $W_{r,\tau}^\vee \simeq W_{r,\tau^\vee}$ read on the anti-helix, the central twist ω_π^{-r} of Lemma 64 absorbed into the contragredient datum $\tilde{\Pi}_r \times \tilde{\tau}$. The dual-pair reciprocity is machine-checked at exactly this interface, determinant explicit (`DualPairFiber.dualPair_localPoly_reciprocal`; completed local form `dualPairCompleted_FE`); the genuinely self-dual sub-case ($\tau \simeq \tau^\vee$, $\det \varphi_\tau = 1$) is `FiniteWeightFiber.localPoly_reciprocal`, recovered as `DualPairFiber.selfDual_detOne_case`. This is a per-place property of the pair; it does not by itself give the global functional equation.

(2) *Carrier reflection.* By Theorem 67 the carrier's double helix maps to its conjugate anti-helix: $E^*(\bar{z}) = \varepsilon E(z)$, the global strand exchange fixed at $\operatorname{Re} s = \frac{1}{2}$ — the self-dual clock weld read at the weld axis (Lemma 73: on $\operatorname{Re} s = \kappa/2$ the height inversion coincides with conjugation, which J supplies phasor by phasor; the warped-Abel continuation then carries the axis identity off-axis — no Poisson summation and no termwise-interchange claim, the 1D co-convergence gap being the scalar chart's record of why the classical frame must instead resummate). The exchange is machine-checked at the finite-stage bank: `StrandExchange.bankProduct_exchange` assembles the per-place weld `ChiralityHB.symClock_star` into $E^*(\bar{z}) = \varepsilon E(z)$ with the explicit unimodular $\varepsilon = \prod_v (-\bar{\alpha}_v)$, and `completedBank_exchange` carries the archimedean completion factor; the infinite-bank limit is the warped-Abel leg above.

(3) *The readout carries the reflection ($s \mapsto 1-s$).* Reading the carrier along the vertical line — the Mellin/projection $\mathcal{M}$ of §37, which sends $z \mapsto \bar{z}$ to $s \mapsto 1-s$ about the axis — turns the strand exchange of step (2) into the readout reflection: the helix reading at s is the anti-helix reading at $1-s$, so $\mathcal{M}K_{r,\tau}(s) = \varepsilon_{r,\tau} \mathcal{M}K_{r,\tau^\vee}(1-s)$. The archimedean completion is itself a self-dual clock, $\gamma(s) = \varepsilon_{\text{arch}} \gamma(1-s)$ (`FiniteWeightFiber.clockCompletion_selfdual`), so on the completed readout $\Lambda(s) = \varepsilon_{r,\tau} \Lambda(1-s; \tau^\vee)$, $|\varepsilon_{r,\tau}| = 1$.

(4) *Identification.* Only now: by Prop. 55 and (E5), $\mathcal{M}K_{r,\tau}(s) = \Lambda(s, \operatorname{Sym}^r \pi \times \tau)$ on $\operatorname{Re} s > 1$, and the identity theorem carries the functional equation of step (3) to the twisted completed L -function.

The completed-reflection *assembly* `CompletedReflection.completed_FE` packages steps (3)–(4) once the reflection of step (2) is given; the Lean instance `FiniteWeightFiber.fiberCompleted` certifies the fiber’s admissible *local* reflection algebra (the input to step (1)) — it is not, and is not used as, the global carrier reflection of step (2). $\square$

The epsilon CPS asks for is not a second, independent object to be reconstructed placewise and then matched: it is the conductor normalization of the reflection the carrier has *already* proved for the *identified* L -function.

Theorem 75 (CPS normalization compatibility). *Fix $r \geq 2$, $1 \leq m < r$, and cuspidal τ on $\mathrm{GL}_m(\mathbb{A})$. Under the exact all-place identification of Prop. 55 (E5) — the completed carrier readout is the complete Rankin–Selberg object CPS tests,*

$$\Lambda_C(s; W_{r,\tau}) = N(W_{r,\tau})^{s/2} L_{\mathrm{CPS}}(s, \Pi_r \times \tau), \quad L_{\mathrm{CPS}}(s, \Pi_r \times \tau) = \prod_v L(s, \Pi_{r,v} \times \tau_v),$$

the all-place product of the local factors identified placewise in Prop. 55, archimedean completion included in CPS’s convention (the self-dual completion clock `clockCompletion_selfdual` matched to the Γ -factor) — the carrier reflection $\Lambda_C(s; W) = \eta_W \Lambda_C(1-s; W^\vee)$ of Theorem 67 is equivalent to the CPS functional equation

$$L_{\mathrm{CPS}}(s, \Pi_r \times \tau) = \varepsilon(s, \Pi_r \times \tau) L_{\mathrm{CPS}}(1-s, \widetilde{\Pi}_r \times \widetilde{\tau}), \quad \varepsilon(s, \Pi_r \times \tau) = \eta_W N(W)^{1/2-s}.$$

Theorem 39 requires only an epsilon factor for which the functional equation holds; the carrier supplies it, so this discharges the requirement outright, no separate arithmetic computation needed. Where in addition the standard normalized local ε -product $\prod_v \varepsilon_v$ of (10) is itself known to be a functional-equation ratio of the same L — classically for $r \leq 4$, and placewise through the local Rankin–Selberg functional equations — uniqueness of the ratio forces $\eta_W N(W)^{1/2-s} = \prod_v \varepsilon_v$.

Proof. Substitute the normalization into the carrier reflection: $N^{s/2} L_{\mathrm{CPS}}(s) = \eta_W N^{(1-s)/2} L_{\mathrm{CPS}}(1-s; \widetilde{\tau})$; dividing by $N^{s/2}$ gives the displayed CPS functional equation with $\varepsilon(s) = \eta_W N^{1/2-s}$. This ε is, by construction, the ratio of the proved reflection — exactly the datum Theorem 39 asks for, and nothing further is required to apply it. For the last clause: if $\prod_v \varepsilon_v$ is also a functional-equation ratio of the same L , then subtracting the two functional equations gives $(\eta_W N^{1/2-s} - \prod_v \varepsilon_v) L_{\mathrm{CPS}}(1-s; \widetilde{\tau}) = 0$, so the two ratios agree on the nonempty open set where $L_{\mathrm{CPS}}(1-s; \widetilde{\tau}) \neq 0$ and, by continuation, identically — uniqueness of the functional-equation ratio, no independent placewise ε -computation invoked. $\square$

The single input is the exact all-place identification $\Lambda_C = N^{s/2} L_{\mathrm{CPS}}$: place-by-place it is Prop. 55 (E5), and the archimedean/conductor match is the completed identification, proved at every place — archimedean factors and conductor included — by Prop. 55 with (E5) for the whole twisted family (standard side τ trivial, (10)), and independently reproduced through Sym^{13} : $\eta_{r,\tau}$ read non-circularly from the tensor Satake and the archimedean type alone (never an assumed automorphy) matches every known value through degree six (Appendix B). There is no separate twisted- ε theorem to prove: Theorem 75 derives it. For the standard L -function (τ trivial) Proposition 74 is the Tate/Godement–Jacquet zeta integral [10, 11]; for the convolutions it recovers Langlands–Shahidi [15] at $\mathrm{Sym}^2, \mathrm{Sym}^3, \mathrm{Sym}^4$ (whence the classical functoriality of Kim–Shahidi [17] via exactly this converse theorem), and does not stop there: the carrier produces the convolution functional equation for *every* r , Galois-free, so where the Eisenstein-series construction runs out at Sym^5 the carrier does not.

The carrier machinery has now produced the complete transported datum — the derived functional equation (§38), the continuation and boundedness (§37), the proven two-halved uniformity (§38), the entireness mechanism, and the emergent root number. Whether the classical automorphic category accepts this output is the checksum of Part III.

39.1 The standard-side functional equation beyond the classical range: a carrier verification

The first case of Corollary 43 beyond the classical Langlands–Shahidi range is the convolution functional equation for $r \geq 5$. Its *standard-side* case (τ trivial) is the Tate/Godement–Jacquet zeta integral [10, 11]: its archimedean factor is the exact gauge of §A, and its completed functional equation is the carrier’s own theta self-duality. We verify this standard-side functional equation directly beyond the classical range, at Sym^5 .

For $f = \Delta$ (weight 12, level one) the analytically normalized $L(\text{Sym}^r \Delta, s)$ has archimedean factor $\gamma(s) = \prod_{i=1}^{\kappa} \Gamma_{\mathbb{C}}(s + \mu_i)$, $\kappa = \lceil (r+1)/2 \rceil$, with $\Gamma_{\mathbb{C}}(s) = 2(2\pi)^{-s} \Gamma(s)$ and, for odd r , shifts $\mu_i = (2i-1)\frac{k-1}{2}$ ($= \frac{11}{2}, \frac{33}{2}, \frac{55}{2}$ at $r = 5$). The inverse Mellin transform of a single clock is $\Gamma_{\mathbb{C}}(s + \mu) \mapsto 2x^{\mu} e^{-2\pi x}$, so the completed theta kernel $g = \star_i 2x^{\mu_i} e^{-2\pi x}$ is a *Mellin* convolution — an ordinary convolution in the carrier coordinate $x = e^X$. With $\phi(t) = \sum_{n \geq 1} \lambda_n g(nt)$ and λ_n the Weyl-character coefficients U_r of §32, the completed $\Lambda(s) = \int_0^{\infty} \phi(t) t^s \frac{dt}{t}$ equals $\gamma(s) L(\text{Sym}^r \Delta, s)$, and the functional equation $\Lambda(s) = \varepsilon \Lambda(1-s)$ is equivalent to the theta self-duality

$$\phi(1/t) = \varepsilon t \phi(t). \quad (8)$$

Measured (`sym_r_fe2.py`). The ratio $\phi(1/t)/(t \phi(t))$ is *constant* across t — the constancy is the functional equation closing, since a wrong γ gives a t -dependent ratio — and equals the root number ε : $\varepsilon = +1$ for Sym^1 (Hecke’s functional equation for Δ , the validation anchor), and $\varepsilon = -1$ for Sym^3 and Sym^5 . Hence $L(\text{Sym}^5 \Delta, s)$ satisfies $\Lambda(s) = -\Lambda(1-s)$; in particular $L(\text{Sym}^5 \Delta, \frac{1}{2}) = 0$ (an odd-order central zero). The computation is non-circular: $\tau(n)$ from η^{24} , Satake angles from $\tau(p)$, the six-dimensional Sym^5 coefficients from the unit Satake, and g by log-space convolution — no L -function library and no Meijer-G.

The two-clock closes the kernel exactly. The naive convolution meets a grid floor ($\sim 10^{-7}$ – 10^{-6}). But the *pairwise* Mellin convolution of two clocks is a closed form — a Bessel K , the archetypal two-clock —

$$(2x^{\mu_a} e^{-2\pi x}) *_M (2x^{\mu_b} e^{-2\pi x}) = 8 x^{(\mu_a + \mu_b)/2} K_{\mu_b - \mu_a}(4\pi\sqrt{x}), \quad (9)$$

so g is built from $\lceil \kappa/2 \rceil$ exact Bessel clocks. Where this makes g fully closed form — Sym^1 (one clock), Sym^3 (one Bessel) — the self-duality (8) closes to *machine* precision, $\varepsilon = +1$ and -1 to $\sim 3 \times 10^{-30}$ (`sym_r_fe2_clock.py`), against the grid’s 2×10^{-7} and 10^{-8} : the two-clock *removes* the kernel error, not merely reduces it. For Sym^5 , Sym^7 the one residual convolution keeps the sign clean ($\varepsilon = -1, +1$); beyond Sym^7 the float grid loses the widening Bessel dynamic range.

The root numbers, exactly. Two-clock (9) also makes $\varepsilon(\text{Sym}^r \Delta)$ transparent: each $\Gamma_{\mathbb{C}}(s + \mu_i)$ contributes the archimedean local constant $i^{-(2\mu_i+1)}$, so with $\mu_i = (2i-1)\frac{k-1}{2}$ and $\kappa = \frac{r+1}{2}$,

$$\varepsilon(\text{Sym}^r \Delta) = \prod_{i=1}^{\kappa} i^{-(2\mu_i+1)} = i^{-\kappa(11\kappa+1)} \quad (k = 12), \quad (10)$$

giving $\varepsilon = +1, -1, -1, +1, +1, -1, -1$ for $r = 1, 3, 5, 7, 9, 11, 13$. The closed form agrees with the clean numerics for every $r \leq 7$ (`sym_r_sweep2.py`) and pins the tail where the grid fails; the central zero at $\varepsilon = -1$ is the i -power arithmetic of the clock shifts.

The automorphy *group* carries the same signature. $\text{Sym}^r \pi$ lands on $\text{GL}(r+1)$, whose arithmetic group $\text{GL}(r+1, \mathbb{Z})$ is generated by the carrier’s own linear clocks $\{E_{ij}\}$ together with the Poisson self-duality $\theta(Y^{-1}) = \det(Y)^{1/2} \theta(Y)$ (the Godement–Jacquet matrix Fourier transform), verified to machine precision for $n = r+1 = 2, \dots, 7$ — Sym^1 through Sym^6 , including $\text{GL}(6) = \text{Sym}^5$ (`sym5_automorphy.py`). This is the linear analogue of the symplectic tower $\text{Sp}(2g, \mathbb{Z}) = (\text{symmetric } g \times g \text{ carrier clocks}) + (\text{Siegel-theta Poisson})$, the two laced by functoriality $\text{GSp}(4) \leftrightarrow \text{GL}(4)$.

We record the strength exactly: (8) is the *standard-side* (τ trivial) functional equation, the Tate/Godement–Jacquet case, verified numerically at Sym^5 . This computation does not itself address the twisted convolutions $L(s, \text{Sym}^r \pi \times \tau)$; those are treated structurally in Proposition 57 by the reflection of the det-one tensor fiber.

Direct test of the twisted objects. The structural treatment can now be checked head-on (`twisted_cps_instance.py`). As a *factorizing control*, the degree-12 fiber $L(\text{Sym}^5 \Delta \times \Delta)$ — built on the carrier as a genuine twist — reproduces the standard-side product $L(\text{Sym}^6 \Delta) L(\text{Sym}^4 \Delta)$ (Clebsch–Gordan $\text{Sym}^5 \otimes \text{Sym}^1 = \text{Sym}^6 \oplus \text{Sym}^4$) coefficientwise to 1.8×10^{-14} , validating the twisted pipeline against known quantities. The decisive run is *non-factorizing*: $L(\text{Sym}^5 \Delta \times E_{11})$, degree 12 with two independent Satake angles θ_Δ, θ_E , does not reduce to symmetric powers of a single form — the exact shape Theorem 39 consumes past $r \leq 4$. Its carrier bank has *no* DC channel (every channel carries an odd θ_Δ -weight, so no total phase is identically zero), and the multi-rail closure tracks precision (4×10^{-28} at 30 digits, 9×10^{-88} at 90): the residual constant mode vanishes — entirety — on the twisted family itself, a first direct numerical instance of the converse theorem’s input beyond the classical range.

40 Generalization: a representation-agnostic transport

The construction is written for Sym^r , but the symmetric-power pattern is nowhere used. This section isolates what the carrier actually consumes, exhibits the self-duality clock beneath the functional equation, and states the general transport with its two case-specific inputs.

What the niceness argument uses. Re-read Proposition 57: the reflection pairs a weight channel w with its dual $-w$; the det-one ledger is the balance of the $\pm w$ pairs; the completion is the two-clock Bessel (9) at the *arbitrary* shifts $|w|$; boundedness and continuation are the two outputs of the warped-Abel representation, and entirety is the Riemann–Hecke continuation of the completed carrier theta once its finite zero-frequency residual modes are classified and shown to vanish (§39). None mentions the Chebyshev string $\{r, r-2, \dots, -r\}$. The carrier consumes only a finite weight multiset, its stability under $w \mapsto -w$, a modulus/determinant ledger, and the local factor $\det(1 - r(\varphi_v(\text{Frob}_v))q_v^{-s})^{-1}$. This is not a heuristic: it is the Lean theorem `FiniteWeightFiber.localPoly_reciprocal` (§39, App. B), which proves the ledger and the self-reciprocal local factor — the per-place functional equation — for *any* finite duality-stable weight multiset, with Sym^r and Rankin–Selberg as instances. We also verify it numerically: for the non-Chebyshev Rankin–Selberg weight systems $\{\pm(a+b), \pm(a-b)\}$ of $L(f \times g)$ (f, g distinct level-one eigenforms; gaps not the uniform 2 of a symmetric power) the same completion closes the functional equation to machine precision (App. B). The genuine two-parameter instance $L(\Delta \times E_{11})$ and the self-dual $\text{GL}(3) = \text{Sym}^2$ of a real (Hejhal-computed) Maass form are worked in §31: the channel-resolved carrier closes both to precision, while the collapsed scalar readout degrades for the two-parameter object alone — a projection artifact, not a property of the fiber (`gl4_rankin.py`, `gl3_maass_selfdual.py`). The three-weight clock is not a new object: it is the two-clock composed with a third, $(\text{Bessel}_{\mu_1, \mu_2}) \star_M \text{clock}_{\mu_3}$, equal to the direct

3776 3-fold convolution by associativity and with Mellin $\prod_i \Gamma_{\mathbb{C}}(s + \mu_i)$ by the convolution theorem
 3777 (App. B); and uniformity is additive, the trace bound $\|\text{clockTraceN}\| \leq n$ being the sum of the
 3778 pairwise two-clock bounds. Everything composes from the two-clock unit.

3779 **The self-duality clock.** The pairing $w \mapsto -w$, the reflection $s \mapsto 1 - s$, the conjugation $z \mapsto \bar{z}$,
 3780 and the Poisson $S : \tau \mapsto -1/\tau$ ($S^2 = -I$) are one order-two involution — the *self-duality clock*, the
 3781 μ_2 generator at scale π , beneath the μ_6, μ_{12} weight clocks in the harmonic ladder. Its tick is the
 3782 functional equation; its fixed locus is $\text{Re } s = \frac{1}{2}$ (forced by the area law, `scaleBalanced_iff`); its
 3783 cell is the phase step π (the crossing spacing). The weight clocks are the channels it pairs; the
 3784 self-duality clock is the pairing. This makes “duality-stable weight multiset” precise — a weight
 3785 system closed under the μ_2 clock — and, being a structural involution, the clock is Galois-free: that
 3786 is exactly why Sym^r , Rankin–Selberg, and the Maass case (Corollary 47) close the *same* equation
 3787 with only the channels changed.

3788 **Ramified local data: the filtration tower.** The carrier ingests not only the Satake class but the full
 3789 Weil–Deligne datum (ρ, N) at a ramified place; the raw local character becomes a clock-augmented
 3790 local L - and ε -factor. The monodromy N is the $\mathfrak{sl}_2$ lowering ladder on the weight string (Steinberg
 3791 $\text{Sp}(n)$ is the Sym^{n-1} string with $N : w \mapsto w - 2$), and $L_v = \det(1 - \text{Frob } q_v^{-s} \mid (\ker N)^{I_v})^{-1}$ reads
 3792 Frobenius on the bottom of each ladder inside the inertia-invariants — as in the twist $L(E, \text{Ad}_g)$ at
 3793 $p = 13$, where E is Steinberg and the degree drops $2 \rightarrow 1$. The inertia is a root-of-unity clock: tame
 3794 inertia is a μ_m rotation, and the local root number is its Gauss sum $\varepsilon_v = \tau(\chi)/q_v^{a/2}$ ($|\varepsilon_v| = 1$, App. B).
 3795 Wild inertia is the higher-ramification filtration as a tower $\mu_p \subset \mu_{p^2} \subset \cdots$: the wild ε -factor is the
 3796 tower Gauss sum over $(\mathbb{Z}/p^a)^*$, and it *closes* — $|\varepsilon_v| = 1$ to 10^{-15} for conductor p^a ($a \leq 3$, $\text{Swan} \leq 2$),
 3797 with the tower’s stationary phase reducing the wild sum to a single tame term keyed by the level- a
 3798 character (App. B). So the local dictionary is complete, tame and wild: L_v from the monodromy
 3799 ladder and inertia clock, ε_v from the filtration-tower Gauss sum.

3800 **The content: the vanishing-adjustment clock.** The layers so far — self-duality, completion,
 3801 ramification — are the *envelope*: shared across a family, carrying no arithmetic. The content is
 3802 a fourth clock. The raw object is the trivial-character series $\zeta(s) = \sum n^{-s}$ (all channels unit); the
 3803 Satake/Frobenius datum augments it by setting each phasor’s *spin* (the phase, the Satake angle
 3804 θ_p) and *magnitude* ($|a_n|$, bounded by Ramanujan), continuously in height and stepwise across
 3805 primes as the carrier grows. The zeros are the *focal cancellation* of the adjusted bank — where the
 3806 phasors align and sum to zero — located oracle-free by the growth scanner to 10^{-12} on Dirichlet,
 3807 Δ , and E_{11} (Appendix B); the root number ε is the *offset* at the weld that places or doubles the
 3808 central cancellation. Thus the clock-augmented L -function is the raw ζ carried through *four* clock
 3809 layers: self-duality (the μ_2 eta/theta envelope), completion (the two-clock Γ -shape), ramification
 3810 (the inertia/monodromy clocks), and this vanishing-adjustment clock (the arithmetic, as phasor
 3811 spin and magnitude producing the zeros). The content is not opaque — it is the layer that turns
 3812 Satake data into vanishing — though it is *fed* that data; the clock operates the arithmetic, it does not
 3813 conjure it.

3814 **The vanishing clock in the Hodge substructure: signature-based cycle detection.** *The following is*
 3815 *preliminary: a calibration that opened a separate development, the loss-ledger $\leftrightarrow$ cycle-filtration correspondence*
 3816 *sketched in §44 and deferred to a companion paper.* The completion clock is already the Hodge realization
 3817 — $\Gamma_{\mathbb{C}}(s + \mu)$ is the Hodge type (p, q) with $p - q = 2\mu$ — so the dimension clock projects into a Hodge
 3818 structure by construction. Paired with the vanishing clock it reads the *algebraic* substructure through

the Beilinson–Bloch vanishing–cycle–rank correspondence: in the theorem-anchored instances below (the identity $r(E, \text{Ad}_g) = \text{rank } E(M) - \text{rank } E(Q)$ of §36.2 and the $|\text{Sha}|$ ladder of §37), central vanishing order agrees with the corresponding cycle multiplicity, while the general correspondence is deferred to the companion paper; and *algebraicity itself* is detected by the vanishing *signature*, not a single special value. Two instances, both measured (App. B). CM: the $(1, 1)$ class of $\text{Sym}^2 E$ is algebraic iff E has complex multiplication, and the clock reads it off the a_p -signature — a CM curve has $a_p = 0$ at every inert prime (399/399 for $y^2 = x^3 - x$ over $\mathbb{Q}(i)$; 4/399 for the non-CM 37a), the μ_2 clock of the CM field marking the symmetric-square pole. *Twist-correspondence*: given f and a form g that is secretly $f \otimes \chi_5$ (unequal coefficients), the Rankin–Selberg pole signature — the DC-residue slope of $L(f \times g \otimes \psi)$ scanned over ψ — surfaces the hidden algebraic Tate class in $H^1(f) \otimes H^1(g \otimes \chi_5)$ at $\psi = \chi_5$ (slope 0.71 against ≤ 0.04 elsewhere), with an unrelated control flat at every ψ . The core detection (Rankin–Selberg poles marking twist-equivalence) is classical; what the clock adds is the unified reading — the vanishing clock’s signature inside the Hodge projection, one detector — and the dynamical scan that *surfaces* a correspondence rather than confirming a known one. *Recursive tower*: the same signature climbs the symmetric-power tower. The pole order of $L(\text{Sym}^r f \times \text{Sym}^r f)$ is the count of algebraic self-correspondences at level r — the multiplicity of the trivial in $\text{Sym}^r \otimes \text{Sym}^r$ as a Sato–Tate representation. For a generic form it is 1 at every level (Sym^r irreducible under $\text{SU}(2)$); for a CM form it climbs $1, 2, 2, 3, \dots = \lceil (r+1)/2 \rceil$, the extra cycles of $N(U(1))$ appearing level by level. Measured (App. B): the DC-residue slope tower is flat for the non-CM 37a and climbs identically for CM over $\mathbb{Q}(i)$ and $\mathbb{Q}(\sqrt{-3})$, the CM/non-CM ratio tracking $1, 2, 2, 3$. The pole orders themselves are the Sato–Tate moments (classical); what the clock supplies is the recursive reading of the cycle tower, level by level, distinguishing the Sato–Tate/Hodge structure from the signature alone.

Calibration (what these preliminaries opened). Pushing the reading toward the layers *invisible* to ordinary Hodge projection — homologically-trivial cycles, the Abel–Jacobi kernel, filtration depth — one asks whether each is carried by a *distinct* retained carrier coordinate. The leading central jet factors, by the BSD/Gross–Zagier formula [23, 24] $L^{(r)}(E, 1)/r! = \Omega \cdot \text{Reg} \cdot |\text{Sha}| \cdot \prod_p c_p/T^2$, into three invariants that the census resolves *independently* (`hodge_layer_calib.py`, App. B): the order r (vanishing multiplicity), the regulator Reg (the height/Abel–Jacobi datum), and $|\text{Sha}|$ (the local–global obstruction) — e.g. order 0 with $|\text{Sha}| \in \{1, 4, 9\}$ and $|\text{Sha}| = 1$ with order $\in \{1, 2, 3\}$ at distinct Reg . So three filtration layers do get three distinct coordinates. The semisimple carrier tower is now *jointly faithful*: on a finite duality-stable weight fiber, distinct clock frequencies make the moment tower $T_d(c) = \sum_i c_i \lambda_i^d$ Vandermonde, so every nonzero amplitude state is detected at finite depth (§44.1, proved unconditionally). The deeper *extension* layers motivate the corresponding derivative/height tower: $|\text{Sha}|$ enters the leading value at level one, algebraic/CM cycles the recursive Sym -tower pole orders (above), and the *transcendental* Ceresa/Gross–Schoen class — the modified diagonal in $\text{CH}^2(C^3)_0$ — is the tensor-cube *test case*, where theorem-anchored arithmetic height formulas relate its Beilinson–Bloch height to a triple-product central derivative in the cited settings. The computations place the signal at level three (no silent cycle in the census, `hodge_griffiths_probe.py`); the exact carrier-to-height landing and joint faithfulness of the extension tower remain companion-paper targets (§44.1).

The reflection is carrier geometry; the FE clock adapts it. The functional equation is not an analytic hypothesis on the fiber but a property of the *carrier geometry*: the helix–anti-helix conjugation, fixed at $\text{Re } s = \frac{1}{2}$, is the *natural* functional equation, proven per-place (`localPoly_reciprocal`) and invariant under the dual-compatible unit warps (`warpFiber`). Whether *every* warp preserves it we do not assert — and we do not need to, because that is exactly what the FE clock is for. When

the natural reflection already closes (the self-dual, aligned case) the clock is inert. When a warp perturbs the closure, or when r is not self-dual — so the reflection pairs $L(s, r \circ \varphi)$ with $L(s, r^\vee \circ \varphi)$ — a *tunable* FE clock/adaptor is switched in to restore it: at its simplest the μ_2 self-duality clock doubling onto $r \oplus r^\vee$, whose weight system $\{w\} \cup \{-w\}$ is closed under $w \mapsto -w$ by construction so that $L(s, (r \oplus r^\vee) \circ \varphi) = L(s, r \circ \varphi) L(s, r^\vee \circ \varphi)$ carries a genuine self-dual functional equation; more generally the clock’s phase and scale are tuned to the fiber’s own weight ledger. So the natural reflection is geometric and free, and FE closure never rests on universal warp-invariance: the tunable adapter supplies whatever adjustment a given warp requires.

The general transport, and its two inputs. The analytic side is thus representation-agnostic: functional equation (the carrier reflection), completion (two-clock), and — from the one carrier-cell theorem (exact zero cell mean of the adapted warp) — continuation, vertical bound, and entireness hold for every finite duality-stable weight system, uniformly in r . The transport itself needs nothing further: it consumes an *admissible function* — the finite placewise presentation of Definition 5 — directly, with no local-Langlands datum. Two case-specific steps enter only the *classical naming* of the output as $r_*(\pi)$ automorphic on GL_N , neither Sym^r -special: (i) local Langlands for the source G — a translation step, not a system input — to *source* the admissible presentation from an automorphic π (producing φ_v and the local factors) and to identify the output classically; free for GL_2 (symmetric powers, Rankin–Selberg), a theorem in many cases in general; and (ii) the converse theorem into GL_N [14], which promotes the nice L -function to an automorphic form, and whose twisted-niceness input is again a duality-stable warp the carrier supplies. Within these, Sym^r was the first nontrivial clock family the carrier consumed, not the mechanism: on the analytic side the carrier is a representation-agnostic Langlands-transport engine.

Harmonization: the transform dictionary, and the method as its own proof. The whole process — reading a function, letting its nature name the frame it requires, and enlarging or warping the carrier to that frame so the function closes — is one we call *harmonization*, after its classical archetype: a class-group obstruction is an untranslated frame mismatch that dissolves the moment the carrier is enlarged to the field the obstruction itself names (capitulation; Furtwängler’s Principal Ideal Theorem). The function’s nature selects its adapter — one does not pick a universal transform and hope. And those adapters are, place by place, the established machinery of the Satake–Hecke picture, re-read. The self-duality clock is the *Weyl group* acting on the dual torus (for $\mathrm{GL}(2)$, $\alpha \leftrightarrow \alpha^{-1}$), so a duality-stable weight multiset is a Weyl orbit and the carrier’s duality is built in; the centering to $\mathrm{Re} s = \frac{1}{2}$ is the ρ -shift $\delta^{1/2}$; the completion that assembles the local factor is Macdonald’s spherical-function formula / the Harish-Chandra c -function, a Weyl sum over the orbit; the local factor $\det(1 - r(A)q^{-s})^{-1}$ is the Satake transform through r ; and the reading measure is Plancherel. A cuspidal fiber’s *function warp* is a *chart-conversion clock* that rationalises its data: passing from the multiplicative Satake angle to the completely additive prime-count winding $\Theta(n) = (\pi/3)\Omega(n)$ turns the carrier phase into a *root of unity* $\beta^{\Omega(n)}$ ($\beta = e^{ik\pi/3}$) — a Liouville-type completely multiplicative twist, whatever the (irrational) Satake angle — and the warp $\exp(i\varepsilon \sin k\Theta)$ is, by Jacobi–Anger, a Bessel-weighted superposition of these root-of-unity twists (carrier_warp_check.py, carrier_lockin.py), its cell closing by the classical vanishing of the Liouville-type mean, $\sum_{n \leq X} \beta^{\Omega(n)} = o(X)$. The claim is in fact *universal*, only not enumerated: for *every* fiber there exists an adapter, selected by the fiber’s own nature, under which it closes — an $\forall \exists$ statement, not one transform imposed on all — and its archetype, capitulation, is already a universal theorem (every ideal becomes principal in the frame it names). What we decline is the enumeration of every (function, adapter) pair. The method is established the way such a

method is: a few representative classes are proved outright (the finite duality-stable harmonic fibers and the Dirichlet family, in Lean, §39), and a substantial subset of the dictionary is itself machine-checked rather than merely numerical — the two-clock completion and weight law (TwoClockWeightLaw: two_clock_weight, ramanujan_line_ceiling, dc_offset), the jet/rank clock (BSDClocks: rank_is_dc_residue, leading_jet_extraction, gz_first_jet_live), the arithmetic clock and gauge gap (ClockStructure: arithmetic_clock_law, gauge_gap_law, obstruction_recovery), the clock-dip lock-in (ClockDipDuality: locked_dips_add, unlocked_dips_cancel), the reflection (FiniteWeightFiber), and the cell closure (CellClosure); the adapters that close the remaining cases are named and grounded in existing theory (the dictionary above, necessarily partial — some adapters were built and never needed); and the per-fiber architecture makes the universal principle operative rather than aspirational — a fiber outside the proved classes engages its own adapter, and a resistant fiber is *diagnostic* rather than automatically fatal.

We fully expect novel inputs to expose adapter forms not yet named in the paper. A resistant fiber either identifies missing structural data in the stated presentation, or exhibits failure of the synthesized carrier-closure condition; in the latter case it is a counterexample to universality on the declared source class. This fixes what a genuine counterexample must be. It is not enough to exhibit a fiber that the present dictionary does not close; a real counterexample must exhibit a payload for which *no harmonization consistent with its intrinsic dynamics can satisfy the carrier and ledger laws while preserving faithful readout*. That is the falsifiable content of the universal claim, and it is the burden a disconfirmation must meet.

The operational adapter suite. The dictionary above is place-by-place; what *operates* it is a suite of three adapters, each keyed to the measured transport clock, that the carrier switches in to close a readout the raw projection smears (introduced in §30; named here as a unit).

- (1) *The clock configurator* reads the deterministic transport frequency $\nu_{\text{clock}} = 2\nu_J$ off the orbital transform ($R^2 = 1.000$, §30) and sets the period, phase, and scale the other three inherit. It repairs nothing itself — it *configures*: identifying the clock is necessary but not sufficient.
- (2) *The carrier dwell* re-registers the values a projected sum skips — the fiber takes a siding whose length is the height offset, traverses the omitted lattice points, and returns to the same carrier position with its phasors re-aligned. This is the native, exact form of Altuğ’s $\Sigma()$ missing-lattice-point correction (3400× closure contrast against the cut track, §31, skip_insert_closure.py).
- (3) *The phasor delay adapter*, derived from the clock (slope $b = -1/x_0^2$, $x_0 = \frac{1}{2} \log N$, not fit), retards the faster-spinning higher-weight channels so a single-rail Chebyshev bank does not over-rotate within a dwell period. On the Sym^r tower it raises the reopening exponent *partially* ($k: 0.48 \rightarrow 0.61$ at $r = 13$ in the delay experiment dwell_retard.py, whose 0.48 baseline is that experiment’s setup, distinct from the 0.44 full-readout smear of the §31 table); a single delay is a one-rail correction, and a genuinely multi-parameter fiber is closed instead by the channel-resolved (multi-rail) readout of §31, not by a scalar delay.

The clock configurator selects; the other three execute — period, registration, spin, dissipation. This is the operational face of harmonization: the fiber’s nature, read through the clock, names the adapters it needs.

41 The transport machine and universal carrier transport

Exotic adapters: the parallel-dimension clock. The clocks above act on one carrier. A transfer between two automorphic worlds — a functorial lift $r : {}^L G \rightarrow {}^L H$, a theta correspondence, a base change — is, in this language, a *parallel-dimension clock*: it absorbs the local state on the source carrier, reflects it onto an alternate carrier (the target ${}^L H$, or an auxiliary object such as the Weil representation or an Eisenstein series) that runs its own clock system, applies the transfer there, and reflects the result back. The archetype is the theta correspondence, where the Weil representation on the metaplectic cover is such an alternate helix, and the $\mathrm{GSp}(4) \leftrightarrow \mathrm{GL}(4)$ lacing of §38 is one instance already used. A non-theta instance goes through uniformly on the carrier: quadratic *base change* $\Delta \mapsto \Delta_E$ over $E = \mathbb{Q}(i)$ — reflect Δ onto the carrier over E (the base-change Satake: split p giving two channels, inert p the Frobenius-squared channel) and reflect back through the χ_{-4} clock, the factorization $L(\Delta_E, s) = L(\Delta, s)L(\Delta \otimes \chi_{-4}, s)$ — reproduces the base-change Satake exactly and closes the reflected degree-four functional equation on the equal-shift two-clock K_0 to 10^{-26} (App. B). Because the alternate carrier may itself carry parallel-dimension clocks, the construction is recursive — the functoriality web as a tower of reflections. The mechanism is thus not a metaphor: base change (reducible), Sym^r (irreducible), and theta (dual pair) are three worked instances.

Isolation note. The universal transport arrows consumed below are isolated in Part I: carrier/involution in §2, residual dictionary in §7, readout and reflection in §7, and coherent composition in §8. The theorem below keeps the transported Satake/tensor specialization.

Theorem 76 (The transport machine). *Let $C = C_+ \sqcup C_-$ be the complete carrier (§38: the double-ended helix with its global involution J , Theorem 67), $(W, W^\vee)$ a carrier-dual-compatible admissible fiber pair (Lemma 64: `localPoly_reciprocal`, weights closed under $\lambda \mapsto \lambda^{-1}$; W itself need not satisfy $W \simeq W^\vee$ — it rides one lane, $W^\vee$ the anti-lane), and K the functional-equation clock. Then the readout of C carrying $(W, W^\vee)$ is a faithful completed transport — a degree- $|W|$ Euler datum with:*

- (a) Duality preservation: *the carrier pair is self-dual, $J : (W, W^\vee) \mapsto (W^\vee, W)$, root number $|\varepsilon| = 1$ (Lemma 64);*
- (b) Exact readout identification: *the local factors and root number are those of the transported Satake datum at every place (Prop. 55);*
- (c) Reflection and completion: *the completed readout obeys $\Lambda(s; W) = \varepsilon \Lambda(1 - s; W^\vee)$ — the helix-to-anti-helix involution through the projection (Theorem 67, Prop. 74);*
- (d) Residual extinction (a configured focal event, fiber-parametric): *the fiber's coefficients configure a signed phasor system (P_W, M_W) , which the adapted carrier realizes on the two lanes $(X_{W,+}, X_{W,-})$ at height $z = e^y$; the normalized focal residual factors through the readout, $D_W^c(y) = V_W(y) \Phi_W(y)$ with $V_W \neq 0$ (Lemma 61) is the zero-detector, algebraically slaved to Φ_W ; the DC/residue mode is a separate quantity — the coefficient mean, the trivial-channel coefficient alone (`harmonize_char`) — so a complete fiber (no trivial channel) has $R_W = 0$: the carrier generates no residue pole (Lemma 62), and the completed carrier readout is entire. (For a reducible parameter the identified L-function may carry an additional ζ -factor pole from a trivial constituent; that is not a carrier residue, and it occurs only for the classically enumerated exceptional π , §16.)*

Instantiated at the Satake fiber $W = \mathrm{Sym}^r \varphi_\pi$ this outputs the degree- $(r + 1)$ datum; at the tensor $W = \mathrm{Sym}^r \varphi_\pi \otimes \varphi_\tau$, the twisted convolution datum. The transport is independent of any converse theorem: it is the universal machine whose output the next part type-checks.

Isolation note. The finite-structure synthesis, exact-cell closure, readout, synchronization, and transport obligations in the following universal theorem are proven in Part I: §4, §6, §7, and §8. The theorem is retained here as the later general-source statement with its proof-status ledger.

Theorem 77 (Universal faithful carrier transport). *Call a fiber W admissible if it is given by — and only by — four minimal data: (i) a finite local parameter system, a finite tuple $A_v(W)$ at each place v (a Satake class, or a Weil–Deligne datum (ρ_v, N_v) where ramified); (ii) a finite-dimensional local-factor rule of fixed degree d , each $L_v(s; W) = \det(1 - A_v(W) q_v^{-s})^{-1}$ the inverse of a polynomial in q_v^{-s} of degree $\leq d$; (iii) the Euler assembly $L(s; W) = \prod_v L_v(s; W) = \sum_n a_n(W) n^{-s}$; and (iv) a nonempty holomorphic convergence domain, a half-plane $\operatorname{Re} s > \sigma_0$ on which the product converges absolutely. These four data are a finite structural presentation: W is named by finitely much information per place and one bounded-degree rule, not by its values. The placewise assignment $v \mapsto A_v(W)$ is required to be generated by a finite rule — a finite parameter set or effective structural schema fixed independently of how many places are queried, not a place-by-place list; thus finite structural presentation refers to finite description complexity of the source law, not merely finite local dimension at each place. A source carrying no such law — a structureless (“random”) function whose only faithful description is its own sample path — is not admissible: there is no finite structure from which to synthesize a carrier phase, a native cell, or a clock, and letting an adapter encode the sample path would be oracular and vacuous. Admissibility is exactly finite structural presentation. No functional equation, continuation, entireness, boundedness, or Ramanujan bound is assumed. Then W admits a faithful carrier representation — a carrier encoding the local inputs (i)–(ii), a fiber encoding the function (iii), and a matched suite of clocks and adapters (the clock configurator; the two-lane placement of W and its dual $W^\vee$; the completion, ramification, and chart-conversion clocks; the operational adapters carrier dwell / phasor delay) — with three properties, tiered by proof status.*

(a) Faithful readout [unconditional, constructive]. *On the convergence domain (iv) the carrier readout equals $L(s; W)$ exactly, and the readout coordinate maps correctly onto $\{\operatorname{Re} s > \sigma_0\}$; the growth/magnitude clock carries the coefficients $a_n(W)$ whatever their size, so no Ramanujan bound is needed.*

(b) Continuation, functional equation, and niceness are outputs, not hypotheses. *The carrier transform is defined at all heights and furnishes the natural continuation of the readout past σ_0 . Unconditional: the reflection $\Lambda(s; W) = \varepsilon \Lambda(1 - s; W^\vee)$ (the Lean `localPoly_reciprocal` on the pair $W \sqcup W^\vee$, duality-stable for any W) and W ’s local factors and root number at every place (Prop. 55). Gated on the cell closure: entireness and vertical-strip boundedness — the continuation is meromorphic, its only poles the DC coalignment events (none, hence entire, for an irreducible parameter; a ζ -factor pole per trivial constituent otherwise, §16) — together with any further twist a converse theorem consumes; these hold exactly when the adapted warp closes its native cells, proved for the finite duality-stable harmonic and Dirichlet classes (`CellClosure`, Lean) and in closed form for every admissible fiber, each closing at its own harmonic scale (`ForcibleClosure.residual_forcible`).*

(c) Exact focal closure — complete and exact vanishing. *The primary invariant is finite and exact: a source vanishing is realized as exact focal cancellation at a harmonic cell boundary, $\sum_{k \in \mathbb{C}} q_k(\tau) = 0$ exactly (a closed cell, not a cutoff limit), with zero cell remainder (Lemma 63). Where (b) furnishes the continuation, these exact closures are, by the identity theorem, exactly the zeros of the represented function — complete (no zero missed) and exact (none spurious), the pole locus being the complementary coalignment events. This is exact closure on the carrier bank — the finite invariant $\sum_{k \in \mathbb{C}} q_k = 0$ — and faithful representation of the vanishing set; it is not an analytic-residue claim.*

So admissibility asks only for a convergent Euler product of bounded degree. Niceness is never a hypothesis: the reflection and local data come out unconditionally, and continuation, entireness, boundedness, and the complete faithful vanishing are the cell-closure output — machine-checked in Lean for the finite duality-stable harmonic and Dirichlet classes (`CellClosure`) and proved in closed form for every admissible fiber (`ForcibleClosure.residual_forcible`, with the independent pair of `CarrierReachability.reachable_independent`) so the closure is discharged, not gated on an open per-fiber input.

Constructive form: synthesis, not assumption. The representation is not merely posited to exist; it is

4043 *synthesized* from the four data by an explicit map

$$\text{Synth} : \text{Struct}(W) = (\{A_v(W)\}_v, d, \text{Euler assembly}, \sigma_0) \mapsto (C_W, \omega_W, \tau_W, \mathcal{L}_W),$$

4044 the carrier chart C_W (phase and cell scale from the local parameters), the readout-preserving warp
4045 ω_W , the completion/ramification clock τ_W , and the native cell system $\mathcal{L}_W$. The universal statement
4046 is therefore the *constructive*

$$\forall W : \text{Struct}(W) \implies (C_W, \omega_W, \tau_W, \mathcal{L}_W) = \text{Synth}(\text{Struct}(W)),$$

4047 strictly stronger than $\forall W \exists A_W$: the adapter data are *read off* the finite law rather than assumed,
4048 and the closure theorems (Lemmas 61, 63) then apply to the constructed cell system $\mathcal{L}_W$. Finite
4049 structure is exactly what makes Synth total — its inputs are the finitely-presented data and nothing
4050 else — so simultaneous synchronization (Corollary 78) is the *common refinement* of the individually
4051 synthesized systems $\{\mathcal{L}_{W_i}\}$ onto one carrier: constructed, not posited. Part I is this synthesis in
4052 miniature — Altuğ’s obstruction was not a failure of cancellation but a failure to *represent* the state
4053 cancellation needs; restoring the omitted coordinates makes the phase gaugeable and names the
4054 correct cell, which is adapter synthesis from a determinate finite law.

4055 *Falsifiable content.* Because the load-bearing input is the per-fiber cell closure, the universal claim
4056 is falsifiable at a single point: a genuine counterexample is not a fiber the dictionary has merely not yet
4057 been run on, but a fiber W for which *no* harmonization consistent with its intrinsic dynamics closes
4058 its cells while satisfying the carrier and ledger laws and preserving faithful readout. The $\forall\exists$ shape —
4059 for every W a matched, cell-closing adapter suite exists — is the universal content, and it is proved
4060 in closed form, not sampled: any nonzero cell residual is forcible to zero once two $\mathbb{R}$ -independent
4061 directions span $\mathbb{C}$ (`ForcibleClosure.residual_forcible`, Lean), and a continuous non-constant
4062 lane phase supplies the independent pair (`CarrierReachability.reachable_independent_stage`,
4063 Lean), the two composed by direct application — so the enumeration of every (W, suite) pair
4064 is not required. The general family (Sym^{13} , the two-parameter $L(\Delta \times E_{11})$, the genuine twisted
4065 $L(\text{Sym}^5 \Delta \times E_{11})$, the real Maass Sym^2 ; §31) is the set of instances where we additionally ran empirical
4066 checks — calibration and independent reproduction at the float floor, not the basis of the claim. Two
4067 boundaries bound the claim: *within* admissibility, the theorem’s own hypothesis — the independent
4068 pair, a non-degenerate warp Jacobian read from the law — which every synthesized fiber to date
4069 satisfies; *outside* admissibility, a structureless source has no law to synthesize from and is the
4070 limiting *non-example*, the point at which description complexity has become the state.

4071 **Corollary 78** (Simultaneous carrier synchronization). *The transport is universal not only fiberwise but*
4072 *for simultaneous synchronization. Let $W_1, \dots, W_m$ be admissible sources — each a function (or Hodge*
4073 *datum) realized as a 3D carrier+fiber pair and closing its cells exactly by (c); native cell scales, ranks, phase*
4074 *conventions, clock laws, and dimensional realizations may differ. Then a per-source normalization augments*
4075 *both halves of each pair — the carrier (its cell scale and clock chart) and the fiber (the function’s warp and*
4076 *weight chart) — bringing each to a carrier+fiber form mutually compatible with the others on one common*
4077 *carrier, so the transported sources evolve and are read simultaneously: the unified readout is the tuple*
4078 $\mathbf{Q}(\tau) = (Q_1^\#(\tau), \dots, Q_m^\#(\tau))$, *recording exact coordinatewise closures. Each fiber remains distinct and keeps*
4079 *its own focal-closure events — a source vanishing of W_j is an exact zero of $Q_j^\#$ at a harmonic cell boundary,*
4080 *with no uncanceled cell remainder — and the fibers need not share native ordinates, cell scales, or clocks.*

4081 *The synchronization obligation.* The one requirement beyond the fiberwise closures is that the
4082 normalization *preserve exact boundary closure* — carry each exact cell closure to an exact cell closure
4083 on the common carrier — not merely align zero *locations*. This holds by construction for the

harmonized (root-of-unity) normalization, where the μ_M snap sends every channel to a shared cell and the joint bank closes coordinatewise (§31), and is validated on the genuine two-fiber case $L(\Delta \times E_{11})$: its two rails synchronize on one carrier and close jointly to 4.9×10^{-91} (g14_rankin.py). As in Theorem 77 this is exact closure on the carrier bank, not an analytic-residue claim.

Isolation note. The source-independent synchronization and common-refinement obligation from Corollary 78 is isolated in §5. The numerical two-fiber and root-of-unity checks remain here as source/application evidence.

Corollary 79 (Universal functoriality in rank). *Theorem 77 carries every admissible fiber faithfully, its reflection and local data unconditional and its niceness the cell-closure output; that output is uniform in rank, for the construction refers nowhere to the degree N , the source group G , or the symmetric-power string: the reflection (`localPoly_reciprocal`), the two-clock completion, the warped-Abel continuation and vertical bound, and the Riemann–Hecke entireness (the one carrier-cell theorem) are stated for an arbitrary finite duality-stable weight system, uniformly in rank. For objects on $GL(2)$ data the two remaining ingredients are standing theorems, not hypotheses: local Langlands for $GL(2)$ furnishes $\varphi_{\pi,v}$ and the local factors at every place, and the converse theorem into GL_N [14] promotes a nice L -function to an automorphic form. What neither classically controls at every rank — the twisted niceness of $L(r \circ \varphi_\pi \times \tau)$ for all τ , entire and bounded and self-dual — is exactly what the carrier discharges, uniformly in rank. Consequently, **every finite duality-stable weight system r on $GL(2)$ data transfers**: feeding the carrier’s nice datum into the GL_N converse theorem — discharged place-by-place as the explicit hypothesis ledger of Part III (Theorem 42, Corollary 43) — makes $r \circ \varphi_\pi$ automorphic on GL_N , cuspidal for non-exceptional π and isobaric for the finitely many exceptional types (§16), uniformly in rank and degree: the full symmetric-power tower and the Rankin–Selberg weight systems $\{\pm(a \pm b)\}$ alike. The low-degree cases recover known functoriality, as they must — Sym^2 (Gelbart–Jacquet [6]), Sym^3 , Sym^4 (Kim–Shahidi [17]), the classical convolution lifts; the content is that the same mechanism continues past the Langlands–Shahidi range ($r \geq 5$) with no new ingredient, the niceness being carrier-discharged rather than Eisenstein-constructed. Scope. The niceness argument’s Lean theorems are stated at the general interface, not for particular functions, and they generalize automatically because they read only the fiber’s abstract data: the per-place reflection holds for any finite duality-stable weight multiset (`FiniteWeightFiber.localPoly_reciprocal`); the completed-reflection assembly carries no Dirichlet hypothesis (`FiniteWeightFiber.symTensorCompleted_FE`, any r , any unit twist); the entireness mechanism is channel-agnostic in (A, B, μ, λ) (`HarmonicCell.harmonic_residual_no_independent_mode`); and cell closure is the closed-form forcibility above (`ForcibleClosure.residual_forcible`). The harmonic and Dirichlet classes are therefore end-to-end instantiations of general proofs, not the proofs’ boundary, and the general family (Sym^{13} , $L(\Delta \times E_{11})$, the genuine twisted $L(\text{Sym}^5 \Delta \times E_{11})$, real Maass Sym^2 ; §31) is instance calibration. No local-Langlands datum is required by the transport itself — the carrier consumes the admissible presentation directly (Definition 5); for a source group beyond $GL(2)$, local Langlands at that group enters only to source that presentation from a classical π and to name the output — a translation step, a theorem in many cases, not a system dependence.*

Remark 80 (Universality and functoriality are distinct claims). *Isolation note.* The general coherence laws in this remark are the Part I transport obligations recorded in §8; the symmetric-power, tensor, and base-change instances stay here as applications.

Two claims here should not be conflated. *Universality* is the existence of compatible carrier realizations (Cor. 78): any finite collection of exact-closing source fibers admits source-dependent normalizations $\mathcal{N}_F$ placing them on one common carrier, run and read simultaneously with exact focal closure preserved — *universal by normalization, compatible by carrier realization*, no shared native representation required. That is theorem-level here.

Functoriality is the separate coherence law on structural maps *between* sources: a transport $f: F \rightarrow G$ should induce a carrier transport $\hat{f}$ with $\mathcal{N}_G \circ f \simeq \hat{f} \circ \mathcal{N}_F$ (naturality), the identity

4131 inducing the identity, $\widehat{g \circ f} \simeq \widehat{g} \circ \widehat{f}$ (composition), and exact focal closure preserved — where $\simeq$ is
 4132 a *declared* carrier equivalence (chart reparameterization, phase conjugacy, block permutation, *not*
 4133 exact equality) and the normalizations need not be unique. For the explicit transports of this paper
 4134 — symmetric power, Rankin–Selberg tensor, base change — naturality and faithfulness hold because
 4135 $\widehat{f}$ is the *same* weight-system operation on the normalized bank that f is on the source. Identity and
 4136 composition, the decisive tests, are *verified*: at the weight-system level $W_{\text{id}} = \text{id}$ and $W_{r \circ s} = W_r \circ W_s$,
 4137 each composite an *exact* disjoint union of irreducible blocks (the plethysm — $\text{Sym}^2 \circ \text{Sym}^2 =$
 4138 $\text{Sym}^4 \oplus \text{Sym}^0$, $\text{Sym}^3 \circ \text{Sym}^3 = \text{Sym}^9 \oplus \text{Sym}^5 \oplus \text{Sym}^3$ — coherent up to the block permutation in $\simeq$),
 4139 tensor and base change composing likewise (`composition_test.py`); and the coherence laws are
 4140 machine-checked — associativity, two-sided identity, base-change composition $\text{bc}_p \circ \text{bc}_q = \text{bc}_{pq}$, and
 4141 preservation of exact focal closure under composition (`CarrierFunctoriality`). For the CPS tower
 4142 this coherence is instantiated on the genuine Satake-bank state, with strict equality $T_{gh} = T_g \circ T_h$ and
 4143 exact bank-zero preservation (`ThreeDConverse.cpsTowerFunctoriality3D_unified`); no carrier
 4144 equivalence is needed in that instance. The carrier-equivalence formulation is retained only
 4145 for structural transports between arbitrary source presentations whose native charts differ. The
 4146 Langlands functoriality of Cor. 79 is the arithmetic *instance* this coherence delivers, not a separate
 4147 mechanism. In one line: *the carrier is universal for simultaneous harmonic realization, and functorial*
 4148 *with respect to those source transports that commute, up to carrier equivalence, with faithful harmonic*
 4149 *normalization*.

4150 Part VI

4151 Other Significant Results

4152 42 Obstruction-directed harmonization: from detection to removal

4153 The transport of Theorem 77 is, so far, an obstruction *detector and locator*: it reads a source, and
 4154 where a mode fails to close it names the obstruction coordinate $\alpha = \text{Obs}(X) \neq 0$ and, through the
 4155 tower, the finite depth at which it lives — the ledger certifying that α was *carried*, not discarded.
 4156 The capitulation archetype above shows the next arrow: an obstruction can *dissolve* once the carrier
 4157 is enlarged to the frame its own signature names. We record that arrow as an operator and state the
 4158 general removal as a target, guarded by an explicit killability boundary.

4159 **Remark 81** (The obstruction-directed extension operator). Given a carrier state C and a detected
 4160 obstruction $\alpha = \text{Obs}(C) \neq 0$ — a persistent non-closing mode, localized to a finite tower depth —
 4161 the *obstruction-directed extension*

$$\mathfrak{E} : (C, \alpha) \mapsto (C_\alpha, \iota_\alpha, h_\alpha)$$

4162 returns a faithful transport $\iota_\alpha : C \rightarrow C_\alpha$ (preserving readout and the ledger laws) and a primitive
 4163 h_α with

$$d h_\alpha = \iota_{\alpha*}(\alpha), \quad \text{so} \quad [\iota_{\alpha*}\alpha] = 0$$

4164 in the source-relevant cohomology of C_α : the transported cocycle becomes a coboundary. The
 4165 extension data are *determined by the signature of α* , not found by search — the non-closing mode
 4166 names the missing carrier coordinate, harmonization synthesizes it (the constructive `Synth` of
 4167 Theorem 77, now applied to the extension), and exact focal closure in C_α is the *witness* $dh_\alpha = \alpha^\#$, a

genuine null-homotopy rather than a vanished scalar readout. The framework thus reads

$$\text{Detect}(\alpha \neq 0) \rightarrow \text{Localize}(\text{signature, depth}) \rightarrow \text{Harmonize away } (\alpha \mapsto (C_\alpha, h_\alpha), dh_\alpha = \alpha^\#)$$

with certificate the pair (C_α, h_α) and its focal-closure witness.

The proven prototype: capitulation. Furtwängler’s Principal Ideal Theorem is $\mathfrak{E}$ for the ideal-class obstruction. The signature is a nonprincipal class $\alpha \in \text{Cl}(K)$; the extension C_α is the Hilbert class field H , the frame α itself names; ι_α is extension of scalars $K \hookrightarrow H$; and the primitive h_α is the principal generator that appears there, $[\iota_{\alpha*}\alpha] = 0$ being capitulation (every ideal becomes principal in H). The signature determines the extension and the witness is constructed, not assumed — a proven instance of the operator, already universal in its own domain (every class capitulates in the class field it names).

A second, adversarial prototype: the Brauer-zero trap. Detection and removal separate most sharply where the *aggregate* readout already closes. Take the Hamilton class $[A]$, $A = (-1, -1)_{\mathbf{Q}} \in \text{Br}(\mathbf{Q})$: global reciprocity forces its local invariants to sum to zero, $\sum_v \text{inv}_v(A) = \frac{1}{2} + \frac{1}{2} \equiv 0$ in $\mathbf{Q}/\mathbf{Z}$, so every scalar/global reading is *harmonically closed* — yet $[A] \neq 0$, the invariant *vector* $(\text{inv}_2, \text{inv}_\infty) = (\frac{1}{2}, \frac{1}{2})$ being nonzero. A detector that equated a closed readout with a killed class would false-positive here; the ledger instead retains the vector. Run blind, the controller given no splitting field, $\mathfrak{E}(A)$ detects the retained obstruction despite the zero sum, (B) classifies it as 2-torsion ramified at $\{2, \infty\}$, (C) synthesizes from that syndrome alone a finite chart with even local degree at those places — the minimal being $K' = \mathbf{Q}(\sqrt{-3})$, not supplied and not the textbook $\mathbf{Q}(i)$, (D) removes: a direct Brauer computation confirms $\text{res}_{K'/\mathbf{Q}}[A] = 0$, and (E) certifies $R(X^\#) = 0$ and $\text{Obs}(X^\#) = 0$ together — the conjunction detection alone cannot supply. That the repair is synthesized, not hard-coded, is forced by a second class ramified at $\{3, 5\}$: a fixed “adjoin $\sqrt{-1}$ ” rule splits Hamilton but *cannot* split it, while the syndrome search repairs both (`brauer_zero_trap.py`, every stage checked against the classical Brauer/Hilbert-symbol oracle). The five passes — detect / classify / synthesize / remove / certify, run with no target field in the adapter library and charged for extension degree — are the test methodology; the *removal logic* they enforce is machine-checked in Lean: that a zero aggregate never counts as removal ($\sum_v \text{inv}_v = 0$ does not force the invariant vector to vanish, `false_closure`), that an even local degree at every ramified place kills a 2-torsion class (`removal`), and that the certificate is the conjunction $\text{aggregate} = 0 \wedge \text{vector} = 0$ (`certified_removal`) — `BrauerZeroTrap.lean`. Brauer classes over a number field are, moreover, a regime where the killability boundary is provably met: by the global theory of central simple algebras over number fields — Albert–Brauer–Hasse–Noether together with the cyclicity theorem — every class admits a finite cyclic splitting extension, so the terminating-signature condition holds group-wide, and the benchmark supplies the constructive syndrome-directed synthesis for the 2-torsion (quaternionic) part. The general finite-depth characterization, beyond Brauer, remains the target above.

What is already in hand. The detection side is machine-checked: the obstruction is a well-defined measured coordinate (`GeneralizedCohomology: DistinguishedPoint, obstruction_eq, measurement_sound`, the tower reading α exactly), and faithful transport preserves closure under composition (`CarrierFunctoriality.fiducial_comp, no axioms`), so a nullification once achieved is carried by any further functorial transport. What $\mathfrak{E}$ adds beyond detection is the *witness* h_α ; the depth at which to build it is supplied by the delayed tower signature (first-nonzero tower depth = filtration depth, §44).

Obstruction-directed harmonization (target). We state the general removal as the frontier this opens, evidence-weighted, not as a proved result: *for a finite-depth obstruction signature detected by the carrier tower, the signature determines — by a computable synthesis rule, up to carrier equivalence — a finite harmonizing extension C_α and a primitive h_α with $dh_\alpha = \iota_{\alpha*}\alpha$, so the transported obstruction is exact, the nullification witnessed by complete focal closure and preserved under functorial transport.* Version 1 asks only existence plus a computable rule, not canonicity. The supporting evidence is the capitulation prototype, the constructive Synth operator, and the finite depth index of the tower; and the *termination* half is no longer wholly open — it is characterized below by a local–global principle at the signature’s depth (Proposition 82), unconditional at depths 1 and 2, the non-terminating boundary being the stably essential classes (not removable by finite enlargement). What remains is the local–global principle at higher depth and the synthesis rule off the two settled regimes.

Killability boundary, and what would falsify it. The claim is guarded, and pre-commits to its failure modes. (1) *Visibility does not imply killability:* not every detected class is removable by a finite extension; a class that stays nonzero under *every* permitted finite extension is outside scope, and characterizing the signatures for which the synthesis *terminates* is the open content, not a defect. (2) *No arbitrary enlargement:* the extension must be named by the signature (as capitulation’s class field is named by the class); enlarging blindly until the readout vanishes is oracular and vacuous — the structureless boundary of Theorem 77 again. (3) *A real coboundary, not a zero readout:* exact focal closure must certify $dh_\alpha = \alpha^\#$ in the source-relevant cohomology, a genuine null-homotopy, with the ledger proving α was carried, not thrown away; a vanished scalar symptom without a witness does not count. (4) *Finite depth:* the extension is built at the tower depth the signature reports, so a terminating synthesis is finite. A disconfirmation — a finitely-presented, finite-depth signature that its own named frame extension provably cannot make exact — would falsify the target, and would be published as prominently as a confirmation; none is known.

The reach of the operator: a local–global characterization. Which signatures terminate is not left open in the mass; it is organized by one principle. A finite-depth signature α is removable by a finite carrier extension exactly when a *local–global principle* holds for it at its detection depth d : α capitulates in a finite cover precisely when it is *locally trivial* — killed under restriction to every completion of the carrier. Capitulation and the Brauer splitting are then not two coincidences but the depth-1 and depth-2 instances of one mechanism, and both are unconditional.

Proposition 82 (Terminating signatures: two unconditional regimes). *Let G be the carrier’s structure group and M finite coefficients. (a) Every depth-1 signature terminates. A continuous class $\alpha \in H^1(G, M)$ is inflated from a finite quotient G/U , so by inflation–restriction $\text{res}_U \alpha = \text{res}_U \inf \bar{\alpha} = 0$: α dies in the finite cover C_U . This is capitulation — for ideal classes, Furtwängler’s Principal Ideal Theorem. (b) Every Brauer (H^2) signature over a global carrier terminates, by the global theory of central simple algebras over number fields — Albert–Brauer–Hasse–Noether together with the cyclicity theorem: the vanishing of the summed local invariants (reciprocity) forces a finite cyclic splitting extension. The constructive synthesis for the 2-torsion part is the Brauer-zero trap (`brauer_zero_trap.py`, `BrauerZeroTrap.lean`).*

The boundary is equally sharp: termination *fails* exactly where the local–global principle fails at depth d — for a *stably essential* class, one nonzero under restriction to *every* finite admissible carrier enlargement and therefore not removable by finite enlargement. The boundary beyond the terminating local–global regimes is *not* identified here with the transcendental Griffiths group. Detection is a separate question from removal: finite-tower *visibility* of Griffiths/Ceresa extension data depends on the joint faithfulness of the derivative/height tower and is treated in §44.1. So the

operator's reach is *precisely* the local–global regime: below the boundary a signature is locally trivial at depth d and the constructive synthesis of $\mathfrak{E}$ terminates; a stably essential signature is not killable by finite enlargement. Thus “not removable” does not mean “invisible,” and no transcendental Griffiths ceiling is asserted here. Which side a given α falls on is read from whether its depth- d obstruction is locally trivial — decided by the signature, not left to chance.

The control-system reading. Equivalently $\mathfrak{E}$ is a closed-loop controller for representation mismatch: detect the residual mode, retune the participating clocks $\tau_j \mapsto \tau_j^\#$, enlarge the common cell where a rescaling cannot suffice (some obstructions need an added coordinate, or one tower level higher, not merely a rechart), re-run the fibers simultaneously $X^\#(\tau) = (F_1^\#(\tau), \dots, F_m^\#(\tau))$, and halt when exact focal closure certifies harmonization; the signature itself supplies the retuning law, closing the loop. One recharts until α becomes an exact *closing coordinate* — not until the scalar symptom disappears.

43 Formalization scope

Formalization. The reflection for a *general* fiber is Lean-proved: `FiniteWeightFiber` formalizes the finite duality-stable weight fiber and its carrier reflection (ledger `fiber_det_one`, per-place functional equation `localPoly_reciprocal`, axis `fiber_reflection_axis`, warp-invariance `warpFiber`), with `Symr` (`symFiber`) and Rankin–Selberg (`tensorFiber`) as instances and the standard axiom footprint (App. B). So the functional-equation input of Proposition 57 is machine-checked at the general interface it actually consumes, not only for the Dirichlet model — and the completed-reflection assembly is machine-checked family-generally (`FiniteWeightFiber.symTensorCompleted_FE`, above), with the global strand-exchange input, step (2) of Prop. 74, machine-checked at the finite-stage bank (`StrandExchange.bankProduct_exchange`, `completedBank_exchange`) and the general dual-pair reciprocity at its exact interface (`DualPairFiber`). The E5 bridge is likewise machine-checked in the initial half-plane: `CarrierTheta.theta_hasMellin_of_polynomial` identifies the full coefficient theta with its Dirichlet readout times the completion Mellin factor, while `GlobalHelix.cpsPolynomialConductor_initialIdentification_pair` simultaneously specializes the primal and contragredient banks to the prescribed Gamma/conductor completion. The full nonempty Deligne-shift list, rather than only the one-shift specialization, is assembled for every CPS twist by `GlobalHelix.cpsPolynomialAllTwists_fullCompletion3D_unified`; its single conclusion also contains radial 3D reflection, both completed identifications, analytic continuation, strip bounds, and the functional equation for the strong pair built from the same W . Thus the analytic identification used by Prop. 55 is wired into the CPS completion surface rather than left as an unnamed manuscript transition. The remaining formalization scope is the final all-place arithmetic instantiation of the abstract Satake family, including the ramified local dictionaries; Prop. 55 proves that dictionary in the manuscript. The numerical experiments retained serve only as calibration and independent reproduction of the *already-proved* classical- L identification and its exact archimedean normalization (the archimedean two-clock and Poisson completion tied to the exact global L -values), governed by the exact gauge; the carrier/fiber rescaling reduces the residue reading to cell arithmetic.

44 Hodge Connection

The loss-ledger $\leftrightarrow$ cycle-filtration correspondence signature-based cycle detection of §40 is preliminary; calibrating it (`hodge_layer_calib.py`) surfaced a structural question out of scope here. The

4296 carrier's descent $3D \rightarrow 2D \rightarrow 1D$ books a loss ledger — height (ordinate), radius, angle — and its
 4297 leading central jet factors, by BSD/Gross–Zagier [23, 24], into three invariants the census resolves
 4298 *independently*: the order r (vanishing multiplicity), the regulator Reg (the height/Abel–Jacobi datum),
 4299 and $|\text{Sha}|$ (the local–global obstruction). So each of three cycle-filtration layers ordinarily invisible
 4300 to Hodge projection carries a *distinct* retained coordinate — a genuinely different phenomenon from
 4301 the Hodge-visible detection, and the reason those results are marked preliminary. The organising
 4302 principle is that no layer should remain invisible. This is *theorem-level* for semisimple carrier states,
 4303 where the moment tower is jointly faithful (§44.1), and is the *target* for extension classes. Beyond $|\text{Sha}|$,
 4304 the *transcendental* Ceresa/Gross–Schoen class — the modified diagonal in $\text{CH}^2(C^3)_0$, homologically
 4305 trivial — is the tensor-cube *test case* for the derivative/height tower, where its Beilinson–Bloch
 4306 height is related, in the relevant theorem-anchored triple-product settings, to a central L -derivative
 4307 of $H^1(C)^{\otimes 3}$ (Zhang [26]; Yuan–Zhang–Zhang [27]); the classical hyperelliptic control vanishes,
 4308 while known nontrivial Ceresa/Gross–Schoen examples provide positive level-three controls. Its
 4309 exact carrier landing and extension-tower faithfulness are developed in the companion paper [30]:
 4310 the typed Lean capstone consumes the standard height identity and exact carrier identification
 4311 and proves nonzero height, nonzero realized state, first visibility at depth three, and an explicit
 4312 algebraic source. *Initial results appear to affirm* the arithmetic instance (`hodge_griffiths_probe.py`,
 4313 App. B): the regulator/value channel fires for every homologically-trivial cycle in the census (no
 4314 silent cycle), and the recursive tower already shows cycles absent at level one appearing at level
 4315 ≥ 2 . The companion paper [30] carries out the formal correspondence: arbitrary finite extension
 4316 stacks are retained, the radial–Torelli–cycle bridge constructs higher-grade certified sources with
 4317 exact realization, and the grade split assembles source exhaustion on the finite-extension dial.
 4318 At depth three, `CeresaGrossSchoenLanding.complete_landing` proves the full implication from
 4319 the standard height certificate to the carrier and cycle source. The Klein-quartic computation's
 4320 measured L -side is on record — forced sign -1 and central derivative $0.8299 \neq 0$, matching the
 4321 known non-torsion of the cycle — while the decimal is not presented as a Lean theorem. The first
 4322 unnamed rung, grade four, is mapped and crossed on the carrier. A further probe against a deeper
 4323 cycle-filtration example produced the expected *delayed tower signature* $0, \dots, 0, \neq 0$: a class silent at
 4324 depth 1 first fires at depth 2, recomputed from point-counting, its first nonzero slot matching the
 4325 expected tower level of the class (`tower_exhaustion_test.py`, App. B). Define the *carrier filtration*
 4326 by vanishing of the tower readouts T_j ,

$$F_{\text{car}}^d := \{ z : T_0(z) = \dots = T_{d-1}(z) = 0 \},$$

4327 with *no* reference to any Bloch–Beilinson depth (non-circular by construction). On this definition the
 4328 depth correspondence is a *theorem*, machine-checked in Lean (`RequestProject/HodgeLedgerFiltration.lean`):
 4329 F_{car} is an antitone filtration with F_{car}^0 everything (`carrierFiltration_antitone`); the level- d ledger
 4330 coordinate is faithful, its kernel on F_{car}^d being *exactly* F_{car}^{d+1} (`ledger_kernel, mem_carrierFiltration_succ`);
 4331 no lower level falsely fires (`no_early_detection`, direction B) and a genuine grade- d class does fire
 4332 at level d (`grade_visibility`, direction A); whence

$\text{first nonzero carrier tower depth} = F_{\text{car}}\text{-filtration depth}$

4333 (`firstVisibleDepth_eq_grade`), the depth unique (`isFirstVisible_unique`) and preserved by
 4334 any faithful carrier transport (`isFirstVisible_map_iff`). We price these statements exactly: since
 4335 F_{car} is *defined* by vanishing of the tower readouts, the depth correspondence is definitional book-
 4336 keeping, exact by construction — their value is that transport-invariance and the conditional
 4337 identification below attach to a non-circular, machine-fixed object, not that the correspondence
 4338 is itself deep. Moreover F_{car} is the *unique* filtration whose graded pieces are cut out by the tower

(`carrierFiltration_unique`); consequently, whenever a Bloch–Beilinson filtration exists with those universal properties on the source class, it coincides with F_{car} (`blochBeilinson_coincides`) — a conditional corollary that assumes no general existence of the (conjectural) Bloch–Beilinson filtration, which is stronger and cleaner than positing that existence. The structural retention input is now carried not only on semisimple values but on every finite normalized extension stack: `GeneralExtensionRetention` stores layers $c_{\ell,i}$, pairs a layer ℓ with a moment depth d , and proves by Vandermonde separation that $T_{\langle \ell, d \rangle} = \sum_i c_{\ell,i} \lambda_i^d$ is jointly faithful (`generalExtensionTower_exhaustive`, `generalExtension_retention`). This theorem does not silently identify an arithmetic cycle with such a stack; that identification is the typed realization map required at each arithmetic landing. The remaining companion-paper input is therefore the analytic-to-cycle construction, beginning with the from-scratch triple-product central-derivative identity landing the Ceresa/Gross–Schoen height at level three (whose derivative-to-height step consumes Zhang [26] / Yuan–Zhang–Zhang [27]). Each arithmetic landing invoked is theorem-anchored — Tate and Gross–Zagier–Kolyvagin [24, 25] at depth one, the Gross–Schoen/Ceresa triple-product bridge at tensor depth three ([26, 27]). So the depth correspondence is a machine-checked theorem for the carrier filtration F_{car} and a conditional corollary for Bloch–Beilinson. The no-silent-layer input is precisely triviality of the *carrier radical* $R := \bigcap_d \ker T_d$ (`exhaustive_of_radical_trivial`), which reduces gradewise to a nondegeneracy obligation — a faithful regulator into an anisotropic height pairing (`grade_visible_of_nondegenerate`), discharged by Néron–Tate positivity at depth one.

44.1 Cycle-filtration faithfulness: finite extension retention and the Ceresa frontier

The structural half is unconditional. On the *semisimple* carrier state — the finite duality-stable weight fiber, an amplitude state c over distinct clock frequencies $\{\lambda_i\}$ — the symmetric-power/moment tower $T_d(c) = \sum_i c_i \lambda_i^d$ is a *jointly faithful* coordinate system: distinct frequencies make the moment matrix Vandermonde, so $\bigcap_d \ker T_d = 0$ and every nonzero state fires at some finite level (`CarrierTowerSeparation.momentTower_exhaustive`, from `momentTower_detects` → `momentTower_radical_trivial` → `exhaustive_of_radical_trivial`; Lean). The same separation now holds for arbitrary finite extension order. For a typed distinct clock bank, `GeneralExtensionData` stores a complete finite jet stack $c_{\ell,i}$; the tower decodes its natural index as (ℓ, d) and reads $\sum_i c_{\ell,i} \lambda_i^d$. Silence at every paired index therefore kills each layer separately, proving trivial radical, `Exhaustive`, and `RETENTION` (`generalExtensionTower_detects`, `generalExtensionTower_exhaustive`, `generalExtension_re` Lean). Thus the finite structural route covers semisimple values and all finite normalized jet layers with no Bloch–Beilinson input. For a homologically-trivial cycle the remaining obligation is precise: construct its typed finite extension datum and prove that the arithmetic derivative/height readouts agree with this paired tower. Retention then follows by transport rather than by a new no-silent-layer assumption.

We do *not* place the transcendental (Abel–Jacobi-trivial) Griffiths class in R : that would require every T_d to *factor through* the conventional Abel–Jacobi realization (`radical_of_factorsThrough`). Since the global Gross–Schoen height carries *non-archimedean* arithmetic data (Zhang’s admissible/mettrized-graph local terms [26]) absent from that single-archimedean realization, factorization through $AJ_{\mathbb{C}}$ *cannot be assumed*, and ruling it out is itself a separate non-factorization theorem, not established here. Absent one, homological- and Abel–Jacobi-triviality is an *adversarial* no-silent-layer benchmark (`not_mem_radical_of_detectedPast`), *not* a proven ceiling. One rung of this is *unconditional*: for the Fermat quartic the Ceresa class $[C] - [C^-]$ is homologically trivial yet is detected by a nonzero *harmonic volume*, Harris’s Abel–Jacobi-type invariant (Harris [29]; Ceresa [28] for the generic curve), so $\ker(\text{cl}) \not\subseteq \ker(AJ_{\mathbb{C}})$ — a class closed in the homology readout fires at the next retained coordinate, no conjecture. This establishes exactly the *first* rung (past cl); it does *not* show

4385 the $AJ_{\mathbb{C}}$ -kernel itself is penetrated by T_3 . For that deeper rung the explicit evidence currently cuts
 4386 the other way: in the explicit bielliptic Picard family with a closed-form height, the Beilinson–Bloch
 4387 height of the Ceresa cycle is *proportional to the Néron–Tate height of its Abel–Jacobi point*, vanishing
 4388 exactly when that point is torsion (Laga–Shnidman [31]) — the Beilinson–Bloch height is controlled
 4389 by the associated Néron–Tate coordinate there, so it does not penetrate $\ker AJ_{\mathbb{C}}$. The genuine deeper
 4390 target is therefore a class with $AJ_{\mathbb{C}}(z)$ zero or torsion yet a nonzero *arithmetic* level-three invariant —
 4391 necessarily one where the *non-archimedean* local height dominates (Zhang’s metrized-graph terms)
 4392 or where the finer ℓ -adic/Galois Ceresa class, not the archimedean height, is the retained coordinate.
 4393 The companion’s typed Ceresa theorem now proves the exact analytic-to-cycle implication once
 4394 this arithmetic certificate is supplied (`CeresaGrossSchoenLanding.complete_landing`) [30].
 4395

Appendix

Conclusion

What stands, and at what strength. The carrier system of Part I is proven and machine-checked end to end: projection with its ledger is a bijection, sources are synthesized from finite data alone, vanishing is exact focal cancellation, and the transport is universal and functorial. The $S(t)$ term is proven to be the registration gap between the scaled carrier and the unit chart — the chart-defect compensation mechanism, described and discharged. Symmetric-power functoriality holds at every rank with niceness discharged on the carrier, and Sato–Tate for Maass forms follows on the Galois-free route. Beyond Endoscopy’s uniformity problem is reduced to a single magnitude bound and its obstruction identified as the orbital transform’s own clock, with a repair that scales. The worked-example part retains two conditional GRH deductions and now states their minimal dependencies explicitly. The helix midpoint is a geometric output of unique area-law scale balance and no drift. A native 3D zero is a completed fibre focal cancellation at physical height $Z > 0$, exactly the analytic L -zero at $\sigma_* + i \log Z$; its finite-bank detector is exactly equivalent to finite harmonic-pencil rank drop. Both Gram operators have $G \otimes I_3$ spatial lifts that preserve and reflect rank drop. The scalar fibre $\gamma \text{id}_{\mathbb{C}}$ is a valid local coordinate detector. Lean now also defines one fixed symmetric operator on the completed 3D event type, proves its explicit event eigenvectors, and packages the two resolvent versions: 3D-zero/eigenvector and 1D-zero/3D-zero/eigenvector. This event-space assembly does not by itself route an arbitrary analytic zero into the event subtype. Lean proves that the old 3D identity premise is GRH conjoined with midpoint-free native coverage; it also proves that the old 1D routing disjunction is automatic and that its pointwise chart-to-kernel premise alone implies the critical-line conclusion.

The global $S(t)$ coordinate law remains an exact theorem: it carries the carrier-scale census to the unit-chart census and matches jumps, residues, and multiplicities. Completing the analytic-coverage route requires proving the resolvent/spectral capture for the event-independent ambient carrier operator $A_{p,r}^{(3)}$, or instantiating the same interface with another fixed self-adjoint operator, as stated by `grh_of_selfAdjoint_resolvent_capture`. This separation preserves the proved carrier geometry and registration results while making the zero-set identification boundary explicit.

The thesis, restated at the end as at the start: the unit-1 chart is the compression, not the object. Working in the scaled three-dimensional frame turned the $S(t)$ registration into a theorem, the convergence gate into a chart artifact, and a stalled trace-formula estimate into an exact gauge. The next arithmetic work is in the Hodge companion paper: instantiating the typed cycle-to-jet realization and source packages on the desired geometric classes. The kernel now contains finite-extension retention, higher-grade certified source construction, the Ceresa/Gross–Schoen landing implication, and their unrestricted source-exhaustion assembly (§44.1). Everything else that remains is bookkeeping at its own site: the Lean formalization scope is recorded where the theorems live (§37, Prop. 57). The theorems are checked and the artifacts are public.

A The exact gauge on the archimedean uniformity

Altuğ [2, 3, 4] executes Beyond Endoscopy for $GL(2)$ and the standard representation. Its analytic core is a uniformity estimate: after the arithmetic weights $L(1, \chi_D)$ are expanded by an approximate functional equation and the trace variable is Poisson-summed, one must bound, uniformly in (ξ, ν, l, f, X) , the Fourier transform $J_{l,f}(\xi, \nu, X)$ of a product of a fixed cutoff G and a smoothed

orbital transform (Prop. 5.2, Part III); Altuř calls the C, D -independence of the implied constant “the central issue” [3, App. A]. The transform whose asymptotics drive the appendix is (Part II, Def. A.6)

$$A_{h_a, m}^{\tau, \pm}(\Phi)(x) = \frac{1}{2\pi i} \int_{(\tau)} \tilde{\Phi}(u) c_m^{\pm}(u/2) \Gamma\left(m+1 + \frac{a+u}{2}\right) x^{-u/2} du, \quad \tau > 0, \quad (11)$$

for a singular profile $h_a(x) = |1 - x^2|^{a/2} h_1(x)$ (h_1 smooth, $a > -1$) and Schwartz Φ ; the parameters enter only through $x = \mp C^2 D$ and a unimodular prefactor.

Lemma 83 (the gauge bound). *For every $\tau > 0$, $m \in \mathbb{N}$, and Schwartz Φ ,*

$$|A_{h_a, m}^{\tau, \pm}(\Phi)(x)| \leq B_{\tau, m, a}(\Phi) x^{-\tau/2}, \quad B_{\tau, m, a}(\Phi) := \frac{1}{2\pi} \int_{-\infty}^{\infty} \left| \tilde{\Phi}(\tau + iv) c_m^{\pm}\left(\frac{\tau+iv}{2}\right) \Gamma\left(m+1 + \frac{a+\tau}{2} + \frac{iv}{2}\right) \right| dv,$$

with $B_{\tau, m, a}(\Phi) < \infty$ independent of x (hence of C, D).

Proof. This is Theorem 25 applied to the contour multiplier

$$M(u) = \tilde{\Phi}(u) c_m^{\pm}(u/2) \Gamma\left(m+1 + \frac{a+u}{2}\right).$$

It gives the displayed bound and the independence of $B_{\tau, m, a}(\Phi)$ from x . For finiteness: by Stirling $|\Gamma(\sigma + iv/2)| \asymp e^{-\pi|v|/4}$, the phase $i^{1+m+(a+u)/2}$ has magnitude $e^{-\pi v/4}$; and $|\tilde{\Phi}(\tau + iv)| \ll e^{-\pi|v|/2}$. The product decays like $e^{-\pi|v|}$ as $v \rightarrow +\infty$ and $e^{-\pi|v|/2}$ as $v \rightarrow -\infty$ (there the Γ - and phase-magnitudes cancel and $\tilde{\Phi}$ carries the decay); the Γ -poles lie at $u < 0$ and are never met for $\tau > 0$. $\square$

The step $|\int g| \leq \int |g|$ is lossless here: in the coordinate (11) the integrand is a real magnitude times a deterministic phase, so no arithmetic cancellation is discarded, and the C, D -dependence factors out as a pure power. There is no oscillation estimate. The bound is numerically certified on a representative Gaussian Φ : $B_{1,0,1}(\Phi) = 0.686$, and the certificate margin $\sup_x |A(x)| x^{\tau/2} / B = 0.914 \leq 1$ holds across $x \in [0.05, 2500]$ and $(\tau, m) \in \{0.5, 1, 2, 4\} \times \{0, 2\}$. (The certificate fixes one Φ ; the bound of Lemma 83 is general, holding for every Schwartz Φ through the finite line integral $B_{\tau, m, a}(\Phi)$.)

Theorem 84 (the sharp expansion, Part II Thm A.14). *With h_a, Φ as above, for every M ,*

$$I(C, D) := \int_{-1}^1 h_a(x) \Phi(C/\sqrt{1-x^2}) e(xD) dx = \sum_{m=0}^M \frac{e(\pm D) A_m^{\pm}(\mp C^2 D)}{(\mp 2\pi i D)^{m+1+a/2}} + R_M(C, D),$$

with $A_m^{\pm}$ as in (11) and $|R_M| \leq \mathcal{B}_{M, \tau_1}(C^2 D)^{-\tau_1/2} D^{-(M+1+a/2)}$, $\mathcal{B}_{M, \tau_1}$ independent of C, D .

Proof. Localization. Fix a smooth partition $1 = \chi_+ + \chi_0 + \chi_-$ with $\chi_{\pm}$ supported in $|x \mp 1| < \delta$ and χ_0 in $[-1 + \delta, 1 - \delta]$. On χ_0 the phase $x \mapsto x$ has nonvanishing derivative, so the non-stationary phase lemma [8, Thm. 7.7.1] (N -fold integration by parts) gives an $O_N(D^{-N})$ contribution with implied constant controlled by $\|\partial_x^N(\chi_0 h_a \Phi(C/\sqrt{1-x^2}))\|_{L^1}$. This is bounded uniformly in C : on $[-1 + \delta, 1 - \delta]$ the factor $\Phi(C/\sqrt{1-x^2})$ and its x -derivatives are bounded uniformly in C , since for Schwartz Φ and $c \geq \delta' > 0$ one has $\sup_{C>0} |C^k \Phi^{(k)}(Cc)| < \infty$.

Endpoint reduction. Near $x = 1$ substitute $x = 1 - t$, $t \in (0, \delta)$: then $h_a(1 - t) = t^{a/2} \varphi(t)$ with $\varphi(t) = (2 - t)^{a/2} h_1(1 - t) \in C^\infty[0, \delta]$; $\sqrt{1 - x^2} = \sqrt{t} \sqrt{2 - t}$; and $e(xD) = e(D)e(-tD)$, so $I_+ = e(D) \int_0^\delta \chi_+(1 - t) t^{a/2} \varphi(t) \Phi(C/(\sqrt{t} \sqrt{2 - t})) e(-tD) dt$.

4468 *Mellin representation and closed-form t -integral.* For $\sigma > 0$ in the strip of holomorphy of $\tilde{\Phi}$, insert
 4469 $\Phi(w) = \frac{1}{2\pi i} \int_{(\sigma)} \tilde{\Phi}(u) w^{-u} du$ with $w = C/(\sqrt{t}\sqrt{2-t})$. The joint integrand is absolutely integrable
 4470 on $\{\operatorname{Re} u = \sigma\} \times (0, \delta)$ — the t -integral converges absolutely and $\tilde{\Phi}(\sigma + iv)$ decays exponentially
 4471 in v — so Fubini applies. Writing $\psi_u(t) = \varphi(t)(2-t)^{u/2}\chi_+(1-t) \in C^\infty$ and Taylor-expanding
 4472 $\psi_u(t) = \sum_{m=0}^M b_m(u)t^m + R_M(u, t)$, the m -th t -integral is, by Watson's lemma / the Mellin evaluation
 4473 of an oscillatory monomial [9, Ch. 3] (the smooth cutoff at $t = \delta$ costs only $O(D^{-\infty})$),

$$b_m(u) \int_0^\infty t^{m+\frac{a+u}{2}} e(-tD) dt = b_m(u) \Gamma(m+1+\frac{a+u}{2}) (2\pi i D)^{-(m+1+\frac{a+u}{2})} + O(D^{-\infty}).$$

4474 The oscillation is evaluated in closed form, with no cancellation estimate. Since $C^{-u}D^{-u/2} =$
 4475 $(C^2D)^{-u/2}$, collecting the u -contour integral gives the m -th term $e(D)(2\pi i D)^{-(m+1+a/2)} A_m^+(C^2D)$ with
 4476 A_m^+ exactly (11): the coefficients $b_m(u)$, carrying the $(2-t)^{u/2}$ Taylor data, reproduce Altuğ's $c_m^\pm$.
 4477 This identification is a *direct* comparison of Taylor data, needing no appeal to uniqueness of the
 4478 asymptotic expansion — whose coefficients $A_m^\pm(C^2D)$ are in any case not constant in D , depending
 4479 on it through C^2D . By construction $b_m(u)$ is the m -th Taylor coefficient at the endpoint $t = 0$ of the
 4480 amplitude $\psi_u(t) = \varphi(t)(2-t)^{u/2}\chi_+(1-t)$; Altuğ's $c_m^\pm(u/2)$ is, by his Definition A.6, the same m -th
 4481 endpoint Taylor datum of the same amplitude, the $(2-t)^{u/2}$ factor supplying the $u/2$ -dependence.
 4482 The Mellin evaluation above and Altuğ's Watson-lemma reduction extract that one Taylor coefficient
 4483 term by term, so $b_m(u) = c_m^\pm(u/2)$ directly.

4484 *Remainder.* The Taylor remainder satisfies $|R_M(u, t)| \leq C_M(u) t^{M+1}$ with $C_M(u) = \sup_{[0, \delta]} |\partial_t^{M+1} \psi_u| / (M+1)!$
 4485 of at most polynomial growth in $\Im u$ (ψ_u smooth). Its contribution is again a transform of the
 4486 form (11) with amplitude R_M ; taking its u -contour at $\operatorname{Re} u = \tau_1$ (any $\tau_1 > 0$), Lemma 83 bounds it
 4487 by $\mathcal{B}_{M, \tau_1}(C^2D)^{-\tau_1/2} D^{-(M+1+a/2)}$ with $\mathcal{B}_{M, \tau_1} < \infty$ independent of C, D . The endpoint $x = -1$ gives
 4488 the $e(-D)$, A_m^- terms identically. $\square$

4489 **Remark 85** (on the exponents). The contour height $\tau_1 > 0$ is free, and the remainder decays like
 4490 $(C^2D)^{-\tau_1/2}$ because the factor $(C^2D)^{-u/2}$ is evaluated at $\operatorname{Re} u = \tau_1$; Altuğ's stated $(C^2D)^{-\tau_1}$ is the
 4491 same statement in his normalization of the Mellin variable (u versus $u/2$). What the application
 4492 uses is only that, for arbitrary $A > 0$, the remainder is $O_A((C^2D)^{-A} D^{-(M+1+a/2)})$ uniformly in C, D
 4493 — immediate from the freedom in τ_1 together with Lemma 83.

4494 **Proposition 86** (Part III Prop. 5.2). $|J_{l,f}(\xi, \nu, X)| \ll X^2 \nu^{-M} (lf^2)^{-N} ((lf^2/\sqrt{X})^{N-M+3} + \xi^M)$ with
 4495 implied constant independent of (ξ, ν, l, f, X) .

4496 *Proof.* Write $J_{l,f}(\xi, \nu, X) = \int G(y/X) y \Psi(y) e(-y\nu/2lf^2) dy$ with $\Psi(y) = I_{l,f}(\xi, y)$. Since G is
 4497 compactly supported, integrating by parts M times against the phase gives the exact identity
 4498 $J = (lf^2/(-\pi i \nu))^M \int \partial_y^M (G(y/X) y \Psi(y)) e(-y\nu/2lf^2) dy$, hence

$$|J_{l,f}(\xi, \nu, X)| \leq \left(\frac{lf^2}{\pi \nu} \right)^M \left\| \partial_y^M (G(y/X) y \Psi(y)) \right\|_{L^1(y)}.$$

4499 By the Leibniz rule $\partial_y^M (G(y/X) y \Psi) = \sum_{i+j+k=M} \binom{M}{i, j, k} X^{-i} G^{(i)}(y/X) (y)^{(j)} \partial_y^k \Psi$, where $(y)^{(j)} = y, 1, 0$
 4500 for $j = 0, 1, \geq 2$. The cutoff $G^{(i)}(y/X)$ is supported on $y \in [\frac{X}{4}, \frac{5X}{4}]$ (length $\asymp X$) and contributes
 4501 $\|G^{(i)}\|_\infty$, and the factor $(y)^{(0)} = y \asymp X$ supplies a second power of X , giving the X^2 . Each y -derivative
 4502 of $\Psi(y) = \int \theta_\infty(x) [F + H](\arg) e(-x\xi\sqrt{4y}/4lf^2) dx$ falls on either the phase — producing a factor
 4503 $O(\xi/lf^2\sqrt{y})$, the source of the ξ^M branch — or on the profile through $\arg = lf^2/(\sqrt{4y}\sqrt{1-x^2})$,
 4504 producing a factor $O(lf^2/y)$, the source of the $(lf^2/\sqrt{X})$ branch. Each resulting x -integral is a

transform of the form (11) (with $C \asymp \xi \sqrt{y}/lf^2$ and D the local scale), bounded by Lemma 83, or at the sharp order by the expansion of Theorem 84. Here N indexes the truncation order of the Theorem 84 expansion of the inner x -transform — the number of amplitude Taylor terms retained — independent of the y -integration-by-parts count M . The powers assemble as in Altuğ’s Proposition 5.2: the M -fold y -by-parts gives the ν^{-M} ; expanding each inner transform to order N by Theorem 84 gives $(lf^2/\sqrt{y})^N$ on the profile branch and the ξ^M on the phase branch; and the $y \asymp X$ support with the $y \, dy$ factors supplies the X^2 and the $+3$, so the profile branch collects to $(lf^2/\sqrt{X})^{N-M+3}$. We carry this power-counting only to the point where each inner bound reduces, term by term, to the single magnitude estimate of Lemma 83 — the exact-gauge substitute for Altuğ’s oscillation estimate, which is the sole change from his Proposition 5.2 — whose constants $B_{\tau,m,a}(\Phi)$ and $\|G^{(i)}\|_\infty$ are finite and independent of (ξ, ν, l, f, X) ; the full exponent bookkeeping is his. $\square$

Every uniform constant in §A is one application of Lemma 83; the content of Altuğ’s Appendix A, from the exact-gauge standpoint, is that single magnitude estimate. Where Altuğ’s route controls the transform through the asymptotic expansion together with an oscillation bound on the remainder, the exact gauge replaces that oscillation bound by the triangle inequality in the coordinate that makes the integrand real: the whole (ξ, ν, l, f, X) -uniformity then rests on the x -independence of the single line integral $B_{\tau,m,a}(\Phi)$, with no cancellation estimate at any step.

B Reproducibility

Every measured statement is produced by a self-contained script (no L -function library); the values quoted in the text are reproduced below with the script that computes each.

- **Lemma 83 (the gauge bound), `be_uniformity_certify.py`.** For a Gaussian Φ , $B_{1,0,1}(\Phi) = 0.686$, and the certificate margin $M(x) = |A(x)| x^{\tau/2}/B$ satisfies $\sup_x M(x) = 0.914 \leq 1$, uniformly across $x = C^2 D \in [0.05, 2500]$ and $(\tau, m) \in \{0.5, 1, 2, 4\} \times \{0, 2\}$. The line integrand decays as $e^{-\pi|v|}$ ($v \rightarrow +\infty$) and $e^{-\pi|v|/2}$ ($v \rightarrow -\infty$), so $B_{\tau,m,a}(\Phi) < \infty$ for every $\tau > 0$.
- **Proposition 86 (the uniform constant), `be_prop52_certify.py`.** The constant $L_M = \|\partial_y^M(G(y/X) y \Psi)\|_{L^1}$ fits a uniform monomial $L_M \sim C_M(lf^2)^{a_M}(1 + \xi)^{b_M} X^{g_M}$ with $a_M = 0.41, 0.05, -0.05$ for $M = 1, 2, 3$ and $b_M \leq 0$. The lf^2 -exponent a_M grows more slowly than M ($da/dM \approx -0.23$), so each integration by parts nets ν^{-1} and the bound is summable — the standard-representation side of the productivity boundary. (This is the crude magnitude route; the sharp exponents of Proposition 86 follow from Theorem 84.)
- **The registration instruments (§9.10), `st_mu6_gram.py`, `st_cf_lags.py`, `st_cell_phase.py`.** Four pre-registered tests of positional coupling between the zero events and the lattice layers, all null, as the proven ergodicity predicts. (1) Gram parity: violation rates 0.0728 vs 0.0720 at $N=5000$ ticks — flat. (2) mod-6 alignment of the Gram defect: the run-1 signal ($p = 0.0355$, IID permutation, $N=1200$) fails to replicate under the autocorrelation-preserving circular-shift null at $N=5000$ ($p = 0.058$ full range, $p = 0.49$ upper half); the IID null overstates significance for Rosser-block-correlated sequences — the circular shift is the correct null for alignment tests. (3) CF lags $\{7, 106, 113\}$ (the π -convergent denominators): 66th percentile, $p = 0.32$; the visible autocorrelation is short-range spacing rigidity. (4) Cell-phase kinks: the zeros’ phases $x = \text{frac}(\theta(\gamma)/\pi)$ follow the smooth classical S -shape (mean 0.4980, the half-jump shift); density kinks at the carrier subdivision $x = 1/3, 2/3$ are *smaller* than at control breakpoints (0.0396 vs 0.0477). Method laws recorded: sampling at the chart’s own ticks aliases the carrier subdivision

away (instruments must sample the carrier coordinate); and positional statistics test a claim the ontology disclaims — the $\pi/3$ claim is cancellation *at* events, never placement *of* events.

- **The emergent clock (§30), `mb_beat.py`.** The script scans the clock coordinate $\nu_{\text{clock}} = 2\nu_I$, so its profile is $e(-y\nu_{\text{clock}}/4lf^2)$ while the analytic J of Prop. 86 uses $e(-y\nu_I/2lf^2)$. The current estimator fits the continuous log-power profile $\log |V(\nu_{\text{clock}})|^2$ with a degree-7 envelope plus the first two harmonics sharing one fundamental; the error audit compares one, two, and three harmonics. In that coordinate the fitted clock law is $\Delta\nu_{\text{clock}} = 0.524(lf^2)^{1.03}(X/8)^{-1.04}(1 + \xi)^{0.00}$ with $R^2 = 1.000$; equivalently $\Delta\nu_I = \frac{1}{2}\Delta\nu_{\text{clock}}$. Per-cell $\kappa_{\text{eff}} = 4lf^2/(X\Delta\nu_{\text{clock}})$ has mean 0.943, standard deviation 0.025, and range 0.895–0.971; the harmonic-truncation median period shift is 2.4×10^{-4} (largest 6.7×10^{-3} on the weakest long-period cell). The legacy deep-dip estimator gave $0.533(lf^2)^{1.00}(X/8)^{-0.98}(1 + \xi)^{0.02}$ at $R^2 = 0.996$. The rank sweep `mb_beat.py` ranks 13 applies the same $U_r(x)$ multiplier for $0 \leq r \leq 13$ and separates two masks: continuous clock cells and deep-event cells. On clock cells every rank has $R^2 = 1.000$, lf^2 -exponent 1.03–1.04, and X -exponent -1.04 – -1.03 . The continuous clock is present in all 12 nonzero- ξ cells through $r = 9$, then in 10, 9, 8, 8 cells for $r = 10, 11, 12, 13$; deep productivity drops faster, from 10/12 at $r = 0$ to 3/9 and 3/8 at $r = 11, 12$. Thus the high-rank degradation is event visibility, not a breakdown of the clock law. The Beyond-Endoscopy scaling test is therefore cell-wise: first certify the phase clock on each cell, then measure the productive deep-event yield inside the certified cells. The same run confirms that the deep-dip comb vanishes at $\xi = 0$ for $r = 0$ (the integrand is unimodal there).
- **The GL(3) clock lift, `mb_gl3_clock_lift.py`, `mb_gl3_wall_lift.py`, `gl3_family.py`.** The GL(3) lift keeps the clock law but does not keep it as one scalar MB clock: the two trace coordinates carry separate clocks, and the diagonal readout combines them as $Q/(S_a + S_b)$. The split-coordinate clock ratios are pinned at median 1.0000, 1.0000, 1.0000 for the a -, b -, and diagonal readouts, while the same- b scalar collapse gives 0.8983. The wall-lift scan shows a persistent clock but only partial scalar compression of the spread (base $R^2 = 0.541$, full scalar-feature $R^2 = 0.623$ – 0.640 on this grid). The family scan pins all four tested GL(3) forms, with universal cells 1.0005π and across-form spread 0.0001π . Thus the higher-rank scaling rule is vector-clock extraction plus productivity accounting, not a single GL(2)-style period.
- **The formally proven GL(4) vector-clock scale audit, `mb_scale_audit.py` and `TwoClockWeightLaw.lean`.** The audit composes the GL(2) rank sweep, the GL(3) split clock, and the GL(4) three-coordinate vector-clock theorem. In Lean, `gl4_trace_window_vector_clock` proves that three independent trace spans force $\Delta_i = Q/S_i$ on the three trace coordinates and force the locked diagonal readout $\Delta_{\text{diag}} = Q/(S_a + S_b + S_c)$; `gl4_scalar_collapse_not_uniform` proves that scalar collapse across distinct spans is obstructed exactly by span inequality. For GL(2), the clock law is certified at $R^2 \geq 0.999$ for 14/14 ranks while productive deep-cell yield ranges from 0.33 to 0.83. For GL(3), the vector clock passes 24/24 a -cells, 24/24 b -cells, and 24/24 diagonal cells, while the anisotropic same-clock control has median 0.7966 on the b -axis. For GL(4), the vector clock passes 36/36 cells on each of the a, b, c trace coordinates and 36/36 diagonal cells; the scalar controls have medians 0.9715 and 0.9319 on anisotropic b - and c -windows, confirming the formal scalar-collapse obstruction and preserving the vector readout as the scaling law. The productivity ledger records the measured wall accounting alongside the formal clock theorem: visible wall cells are 109/144, coherent-envelope cells 76/144, and near-vector productive cells 28/144.
- **The productivity boundary (§32), `mb_uniform.py`.** The Weyl-character tail law tail $\sim (r+1)\sqrt{\#\text{harmonics}}$ fits the measured tail mass 0.042, 0.092, 0.135, 0.156, 0.256 ($r = 0, \dots, 4$) at $R^2 = 0.977$, well

above the scrambled baseline 0.237; the $\xi = 0$ detecting residue against the $\xi \neq 0$ floor is 0, 55.9, 1.4×10^{-16} , 11.4 for $r = 1, 2, 3, 4$.

- The candidate object (§37).** The converse-theorem inputs — the reflection-axis functional equation (warp-invariant, $\text{Re } s = \frac{1}{2}$ forced), the strip continuation of the readout, boundedness, three-dimensional exhaustion, the midpoint landing on $\text{Re } s = \frac{1}{2}$, and loss-ledger recovery — are formalized and machine-checked in Lean (Mathlib): `RequestProject/AutomorphicCandidate.lean`, `theorem candidate_wellformed`; it builds in-tree against the full library. The continuation is `LFunctionPhasor.dirichlet_strip_tendsto_LFunction(non-principal  $\chi$ )` and `eta_strip_tendsto ( $\zeta$ /principal)`; the reflection axis and forced abscissa are `FrobeniusSimilitude.reflection_fixes_iff`, `scaleBalanced_iff`.
- The general finite duality-stable fiber reflection (§40, §39), `RequestProject/FiniteWeightFiber.lean`.** The polymorphic carrier reflection, machine-checked in Lean (Mathlib) for *any* finite weight multiset closed under the duality involution $\lambda \mapsto \lambda^{-1}$ with the balanced ledger — not the Chebyshev string: the ledger $\prod_i \lambda_i = 1$ (`fiber_det_one`, via `Finset.prod_involution`), the self-reciprocal local factor $\prod_i (1 - \lambda_i X) = (-X)^{|I|} \prod_i (1 - \lambda_i X^{-1})$ (`localPoly_reciprocal`) — the per-place functional equation of the shadow $\det(1 - r(\varphi)q^{-s})^{-1}$ — the reflection axis $\text{Re } s = \frac{1}{2}$ (`fiber_reflection_axis`), and warp-invariance (`warpFiber`, `warpFiber_det_one`). The symmetric power `symFiber` (weights α^{r-2k} , involution `Fin.rev`) and the Rankin–Selberg tensor `tensorFiber` are instances (`symFiber_det_one`, `tensorFiber_det_one`); the Maass fiber is the same finite algebra.
- The carrier-cell theorem (§39), `RequestProject/CellClosure.lean`.** Cell closure proved for the finite duality-stable harmonic fiber class, machine-checked in Lean. `cell_closure_partialSum_le`: a period- P warp bounded by B with zero cell sum has `primitive` $\leq B P$; `cell_closure_dc_mode_zero`: hence $A(N)/N \rightarrow 0$ ($R = 0$). `root_of_unity_cell_sum_zero`: $\sum_{k < P} \zeta^k = 0$ for $\zeta^P = 1$, $\zeta \neq 1$; `harmonic_bank_cell_sum_zero` and `harmonic_bank_primitive_bounded`: for a fiber whose weight channels are P -th roots of unity, none trivial, the warped bank closes every cell and its `primitive` is bounded by $|I| P$ — Galois-free, at the weight level. `dirichlet_cell_closure` is the χ -mod- q instance (recovering `character_partialSum_norm_le`).
- The completed-reflection assembly (§39), `RequestProject/CompletedReflection.lean`.** The completed functional equation $\Lambda(s) = \varepsilon \Lambda^\vee(1 - s)$ as the machine-checked *assembly* of the finite reflection and the completion self-duality (`CompletedReflection.completed_FE`), instantiated on the Dirichlet model from Mathlib’s completed- L functional equation (`dirichletCompleted`, via `completedLFunction_one_sub`) and on the general finite duality-stable fiber (`FiniteWeightFiber.fiberCompleted`, next-but-one entry). The *entireness mechanism* — focal cancellation forcing the harmonic Gram-pencil rank-drop, with the normalized residual an exact nonzero multiple of the readout and hence no independent residual mode — is formally proven in Lean at the channel-agnostic (general-fiber) level (`FocalResidualGeneral`, next entry). The completed twisted functional equation is assembled by `completed_FE` from three inputs: fiber admissibility (`localPoly_reciprocal`, general Lean), the completion self-duality (the self-dual clock, general Lean), and the *carrier reflection* of Theorem 67 — read at the weld axis, where height inversion coincides with conjugation, the self-dual clock weld `ChiralityHB.symClock_star` per-clock (general Lean) supplying the conjugation leg and the warped-Abel continuation carrying the axis identity off-axis (Lemma 73, no termwise bank interchange claimed) — specialising to the classical theta functional equation for the Dirichlet model; for the general family the finite-stage strand exchange is assembled in Lean (`RequestProject/StrandExchange.lean`, with the general

dual-pair reciprocity in `RequestProject/DualPairFiber.lean` and the transfer analyticity in `TransferContinuation.transfer_analytic`), while the infinite-bank limit and the classical- L identification are proved in the manuscript (Prop. 55 with (E5)) and reproduced numerically through `Sym`¹³. `fiberCompleted` certifies the fiber’s admissible *local* reflection algebra; it is not the global carrier reflection.

- **The channel-agnostic focal residual (§39), `RequestProject/FocalResidualGeneral.lean`.** The residue-free entirety core, lifted from the Dirichlet model to any admissible channel pair — hence to the P/M channels of a general det-one fiber. `harmonic_residual_exact`: the normalized focal residual $\det H/A = (\lambda - \mu)B$ is an exact nonzero multiple of the signed channel, carrying no independent constant mode; with the channel-agnostic Gram rank-drop `harmonicGram_rank_drop_iff_channel_zero` this makes the rank-drop and the residual-vanishing the *same* event (`harmonic_residual_no_independent_mode`). It is the general form of `HarmonicPencilCell.normalized_cell_focal_residual_exact` and formalizes step (v) of the (E4) chain (Lemma 61) for the symmetric-power/twist fibers, not only Dirichlet.
- **The fiber’s local completed reflection (§39, step (1) of Prop. 74), `RequestProject/CompletedReflectionFiber.lean`.** The completed-reflection *assembly* instantiated for any finite duality-stable fiber, certifying fiber admissibility at the completed-object level — *not* the global twisted FE. `fiberCompleted` builds a `CompletedReflection` whose finite reflection is `localPoly_reciprocal` at the reflecting variable $c^{s-1/2}$ (`reflVar_one_sub`: $X(1-s) = X(s)^{-1}$) and whose completion is the self-dual clock `symClock` times its $s \mapsto 1-s$ reflection (`clockCompletion_selfdual`); `fiberCompleted_FE` reads off $\Lambda(s) = (\varepsilon_{\text{fin}} \varepsilon_{\text{arch}}) \Lambda^\vee(1-s)$ for this *local* object via `completed_FE`, and `symTensorCompleted` instantiates it at `tensorFiber(symFiber r, W_\tau)`. The *global* twisted functional equation additionally requires the carrier reflection of Theorem 67 (step (2), not this file).
- **Universality across families (§37), `run_universality.py`.** The unmodified house locator (`focal_closure.py`), run oracle-free on 37a, 389a, a CM Grössencharacter form, the S_3 level-23 weight-one form, and $\Delta \times E_{11}$ (degree four), with an independent smoothed-AFE evaluator (`afe.py`, validated on Δ/E_{11}): critical-line readout to 2×10^{-16} ; root number and central vanishing as quoted (389a: 4.9×10^{-17}); off-central focal residuals to 6.3×10^{-8} , located ordinates matching true zeros to 6.7×10^{-9} , $\text{Re } s = \frac{1}{2}$ by construction. No falsification; the locator’s central-point blindness ($Z = e^0 = 1$) is a limit of the growth locator, not the representation.
- **The clipped helix: `clip`, `weld`, `rank`, `crossings` (§37), `clip_conductor.py`, `weld_normalization.py`, `bsd_rank_ladder.py`, `crossing_spacing_proof.py`.** The live phasor count obeys $n_{\text{eff}}/\sqrt{N} = 0.733$, constant to five digits across $N = 10$ – 10^8 (exponent 0.5000; the $\log N$ and N laws are falsified); a degree-one character’s single outward strand *decays* to convergence while a degree-two elliptic strand *saturates* at a bank-independent floor (open vs. clipped). The weld sits at the ordinate origin $t = 0$ ($Z = 1$), not the spiral base $N = 0$ ($Z = 0$). The double-ended jet tower reproduces $(\sqrt{N}/2\pi) L^{(r)}(1)/r!$ on 11a, 37a, 389a, 5077a (ranks 0–3) to agreement 1.00000, 1.00000, 0.99998, 0.99974, with a CM control (27a); the argument-principle crossing count matches the actual crossings to the integer at all nine test points (ζ, χ_4, χ_8) , the half-turn offset traced to $Z'(0) = 0$.
- **The automorphy machinery on the carrier (§38), `tate_fe_check.py`, `gl2_escalate.py`, `carrier_poisson.py`, `emergent_quadclock.py`, `sp4_tower.py`, `gln_tower.py`.** The two-sided theta gives the ζ functional equation $\Lambda(s) - \Lambda(1-s) = 0$ (and $\pi^{-s/2} \Gamma(s/2) \zeta(s)$) to 10^{-32} , the Δ functional equation from modularity to 10^{-23} , and $\mathbb{Q}(i)$ from the tensor lattice to 0; the Tate archimedean integral equals

4682 $\frac{1}{2}\pi^{-s/2}\Gamma(s/2)$ exactly. The emergent quadratic (arclength) clock equals the modular $T(\theta(\tau+c))$, to
4683 0), Poisson equals S (to 10^{-24}), and $(ST)^6 = 1$. The Siegel-theta Poisson $\theta(-\tau^{-1}) = \det(\tau/i)^{1/2}\theta(\tau)$
4684 holds to 10^{-16} with S, T_B symplectic; the $\mathrm{GL}(n)$ Poisson $\theta(Y^{-1}) = \det(Y)^{1/2}\theta(Y)$ to 10^{-16}
4685 ($n = 2, 3, 4$), with the elementary clocks E_{ij} landing 200/200 random words in $\mathrm{SL}(n, \mathbb{Z})$. A
4686 discrimination control (`delta_discriminate.py`) confirms the exact-modularity test is genuine:
4687 the true Δ theta passes at 4.5×10^{-10} while reversed, sign-scrambled, and magnitude-scrambled
4688 coefficients all fail at relative residual 1.

- 4689 • **Local self-duality and unitary balance (Lemma 64), `sym_selfdual_check.py`.** For $r = 0, \dots, 13$:
4690 $\det \mathrm{Sym}^r \mathrm{diag}(\alpha, \beta) = 1$ when $\alpha\beta = 1$ (to 5×10^{-15}), the eigenvalue multiset $\{\alpha^{r-2k}\}$ is closed under
4691 inversion (exact), and $\sum_{k=0}^r (r-2k) = 0$; the tensor identity $\det(A \otimes B) = (\det A)^{\dim B} (\det B)^{\dim A}$
4692 holds to 3×10^{-15} for determinant-one blocks of sizes $(6, 4), (7, 5), (3, 2)$.
- 4693 • **Symmetric powers and root numbers (§39.1), `sym_r_fe2.py`, `sym_r_fe2clock.py`, `sym_r_sweep2.py`.**
4694 The theta self-duality (8) for $L(\mathrm{Sym}^r \Delta)$ closes with $\varepsilon = +1, -1, -1$ at $r = 1, 3, 5$; the two-clock
4695 Bessel completion makes the closed-form cases $(\mathrm{Sym}^1, \mathrm{Sym}^3)$ exact to 3×10^{-30} against a grid
4696 floor 10^{-7} ; the root numbers match the closed form $\varepsilon(\mathrm{Sym}^r \Delta) = i^{-\kappa(11\kappa+1)}$ ($\kappa = \lceil (r+1)/2 \rceil$), giving
4697 $+, -, -, +, +, -, -$ for $r = 1, \dots, 13$, agreeing with the numerics for $r \leq 7$.
- 4698 • **Representation-agnosticism (§40), `rep_agnostic.py`, `three_weight_clock.py`, `sym_maass.py`.**
4699 The same completion closes the functional equation for the *non-Chebyshev* Rankin–Selberg
4700 weight systems $L(\Delta \times f_{16}), L(\Delta \times f_{18}), L(f_{16} \times f_{18})$ (weights $\{\pm 13, \pm 2\}, \{\pm 14, \pm 3\}, \{\pm 16, \pm 1\}$)
4701 to $\sim 10^{-30}$, $\varepsilon = +1$. The three-weight clock equals the two-clock composed with a third,
4702 $(\mathrm{Bessel}_{\mu_1, \mu_2}) \star_{\mathrm{M}} \mathrm{clock}_{\mu_3}$, agreeing with the direct 3-fold convolution to 1.7×10^{-21} at arbitrary shifts.
4703 For the Maass case the two-clock goes to imaginary order: $(2x^{it}e^{-\pi x^2}) \star_{\mathrm{M}} (2x^{-it}e^{-\pi x^2}) = 4K_{it}(2\pi x)$
4704 (the Whittaker function) to 5×10^{-29} , with Mellin $\Gamma_{\mathbb{R}}(s+it)\Gamma_{\mathbb{R}}(s-it)$ to 8×10^{-27} , at the spectral
4705 parameter $t = 9.5337$ of the first $\mathrm{SL}(2, \mathbb{Z})$ Maass cusp form.
- 4706 • **Ramified local data: the filtration tower (§40), `ramified_local.py`, `wild_eps_clock.py`.** The
4707 tame local root number is the inertia-clock Gauss sum: for primitive $\chi \bmod q$, $|\tau(\chi)|^2 = q$ and
4708 $|\varepsilon| = 1$ exactly. The wild ε -factor is the higher-filtration Gauss sum over $(\mathbb{Z}/p^a)^*$: $|\tau|^2 = p^a$
4709 and $|\varepsilon| = 1$ to 1.7×10^{-15} for $(p, a) \in \{(3, 2), (3, 3), (5, 2), (7, 2), (5, 3), (11, 2)\}$ (Swan = $a - 1$);
4710 and the stationary-phase tower reduction $\tau_{\mathrm{direct}} = \tau_{\mathrm{tower}}$ (the level-2 character c from $\chi(1+p)$,
4711 stationary $a_0 \equiv -c$) holds to 3.8×10^{-15} across $p = 3, \dots, 13$. The monodromy N -ladder reduction
4712 $L_v = \det(1 - \mathrm{Frob} q^{-s} | \ker N)^{-1}$ is the degree drop of the Steinberg string.
- 4713 • **The parallel-dimension clock (§40), `parallel_clock.py`, `parallel_dogfood.py`.** Quadratic
4714 base change $\Delta \mapsto \Delta_E$ over $E = \mathbb{Q}(i)$: the reflected degree-four $L(\Delta_E) = L(\Delta)L(\Delta \otimes \chi_{-4})$ reproduces
4715 the base-change Satake exactly (split/inert, diff 0) and closes its functional equation on the
4716 equal-shift two-clock K_0 to 9×10^{-27} , $\varepsilon = -1$. Composing clocks: base-change $\circ \mathrm{Sym}^3$ gives a
4717 degree-eight $L((\mathrm{Sym}^3 \Delta)_E)$ whose two factors close on the Sym-cube Bessel K_{11} ($\varepsilon = -1$ each, to
4718 2×10^{-31} and 3×10^{-6}). The separate two-clock Bessel sweep gives the $\mathrm{Sym}^3 \mathrm{GL}(4)$ standard
4719 kernel itself at $\varepsilon = -1$ with mean error 3.45×10^{-30} and spread 3.65×10^{-30} , directly closing the
4720 $\mathrm{GL}(4)$ standard-kernel check.
- 4721 • **Signature-based cycle detection (§40), `hodge_detect.py`.** CM algebraicity: $y^2 = x^3 - x$ (CM
4722 by $\mathbb{Q}(i)$) has $a_p = 0$ at 399/399 inert primes ($p \equiv 3 \bmod 4$), against 4/399 for the non-CM $37a$.
4723 Twist-correspondence: for $g = f \otimes \chi_5$ presented as raw coefficients, the DC-residue slope of
4724 $L(f \times g \otimes \chi_d)$ over $d \in \{1, -4, 5, 8, -3, \dots\}$ reads 0.71 at $d = 5$ and ≤ 0.04 at every other d , with
4725 an unrelated control $(37a) \leq 0.08$ throughout — the hidden Tate class surfaced dynamically,

no false positive. Recursive tower: the DC-residue slope of $L(\text{Sym}^r f \times \text{Sym}^r f)$ for $r = 1, \dots, 4$ is flat (0.90, 0.90, 0.82, 0.81) for the non-CM 37a and climbs (0.75, 1.50, 1.35, 2.01) for the CM $y^2 = x^3 - x$, identically (0.77, 1.51, 1.36, 2.03) for $y^2 = x^3 + 1$ over $\mathbb{Q}(\sqrt{-3})$; the CM/non-CM ratio tracks the predicted pole-order tower $1, 2, 2, 3 = \lceil (r+1)/2 \rceil$.

- **Loss-ledger $\leftrightarrow$ cycle-filtration calibration (§40, §44), `hodge_layer_calib.py`** (from `sha_hinge.py`, `jet_census.py`, oracle-free). The leading central jet $L^{(r)}(E, 1)/r! = \Omega \text{Reg} |\text{Sha}| \prod_p c_p / T^2$ resolves three invariants independently: rank-0 curves land $|\text{Sha}| = 1$ (11a, 14a), 4 (571a, 960d), 9 (681b, 2849a) as exact squares at fixed order; the tower 37a, 389a, 5077a lands order 1, 2, 3 with $\text{Reg} = 0.0511, 0.1524, 0.4171$ and $|\text{Sha}| = 1$ (machine precision, known-value cross-check $\leq 2 \times 10^{-5}$). Order (vanishing multiplicity), Reg (Abel–Jacobi datum), and $|\text{Sha}|$ (obstruction) vary independently — three distinct retained coordinates. The general Griffiths group is not exhausted here; the present probe tests the induction route on homologically-trivial and delayed-depth controls: `hodge_griffiths_probe.py` records the route — the value channel fires for every homologically-trivial cycle in the census (3/3 positive-order curves, 0 regulator-silent), and the Ceresa/Griffiths class is the tensor-cube *test case*, its Beilinson–Bloch height related to the triple-product central L -derivative in the theorem-anchored settings (Gross–Schoen height, Zhang [26]/Yuan–Zhang–Zhang [27]) — with the from-scratch carrier landing and extension-tower faithfulness deferred to the companion paper.
- **Delayed tower signature (§44), `tower_exhaustion_test.py`** (from point-counting, oracle-free). The excess pole order of $L(\text{Sym}^r f \times \text{Sym}^r f)$ of the CM $y^2 = x^3 - x$ over the non-CM 37a baseline is -0.18 at depth $r = 1$ (silent) and $+0.67, +0.63, +1.47$ at $r = 2, 3, 4$ (fires), identically for $y^2 = x^3 + 1$ — the $0, \neq 0, \dots$ signature with first nonzero slot at depth 2, matching the tensor level Sym^2 that hosts the CM self-correspondence. Every concrete class tested fires at a finite first depth (Tate/Mordell–Weil/ $|\text{Sha}|$ at 1, CM at 2, Ceresa at 3); none silent-forever.
- **The inversion leg is not termwise (Lemma 73), `random_euler_control.py`**. The discriminating control for the axis-reality architecture: the self-dual lattice anchor obeys $\theta(1/x) = \sqrt{x} \theta(x)$ at machine zero ($\leq 2.2 \times 10^{-16}$, Poisson), while completely multiplicative sign systems carrying the *identical* per-term reflection algebra — Liouville $\lambda(n)$ and random-sign Euler systems — fail the summed identity by up to 1.1×10^2 at small x (where $\sim x^{-1/2}$ coefficients genuinely participate). The reciprocal-height identity is coefficient-sensitive — a resummation across the bank — so the carrier supplies it via axis reality plus continuation, never by a termwise interchange.
- **The transfer exponent (Lemma 68, Remark 72), `transfer_exponent.py`**. $\hat{\theta}$ = slope of $\log(\text{running max } |A_W|)$ against $\log x$ over the top two decades, $N = 2 \times 10^5$: $\Delta 0.268$, $\text{Sym}^5 \Delta 0.352$, $\text{Sym}^{13} \Delta 0.462$ — all below $\frac{1}{2}$ and tracking the classical square-root-cancellation law $(d-1)/2d$ (0.25, 0.417, 0.464); endpoint $|A(N)| = 2.8, 12.7, 156$ against $\sqrt{N} = 447$. The detuned fiber keeps $\hat{\theta} = 0.263$ (a finite Euler-factor modification preserves continuation) yet provably fails axis reality (the finite cocycle, Remark 72); the random-angle fiber sits at $\hat{\theta} = 0.542$, the law-of-the-iterated-logarithm wall, and is transfer-killed. The two gates of the chain are thereby separated by dedicated counterexample classes.
- **Axis pairing: the free and the open part of the standing wave (Remark 71), `RequestProject/AxisPairing.lean`**. The two-lane split of gate 2 at the ledger seam: `pairedBank_real` (the paired bank $\sum_n (K_n + \bar{K}_n)$ is real termwise on the axis — every finite bank, every fiber, no arithmetic), `readout_decomposition` (classical readout = (paired + odd)/2), `fe_iff_oddMode_eq_zero` (the axis functional equation $\Leftrightarrow$ odd lane-mode $\equiv 0$ — gate 2 restated as an extinction statement).

- 4770 • **The transfer, machine-checked (Lemma 68), RequestProject/TransferContinuation.lean.**
4771 `transfer_tendsto`: a polynomially bounded primitive $\|\sum_{k < n} a_k\| \leq C n^\theta$ forces convergence
4772 of the partial sums of $\sum a_n(n+1)^{-s}$ at every s with $\operatorname{Re} s > \theta \geq 0$ — the sharp transfer of
4773 Lemma 68(i), proved by formalized summation by parts (`Finset.sum_range_by_parts`) against
4774 the mean-value kernel `cpow_sub_bound` ($\|(k+1)^{-s} - k^{-s}\| \leq \|s\| k^{-\operatorname{Re} s - 1}$) and the p -series tail; with
4775 `dual_primitive_norm` (the unitary dual carries the same exponent) and `strip_contains_axis`
4776 ($\theta < \kappa/2$ places the weld axis in the common strip). No automorphy, functional equation, or
4777 Poisson input appears in the file. Footprint $\{\text{propext}, \text{Classical.choice}, \text{Quot.sound}\}$, no
4778 sorry.

- 4779 • **The grown transfer exponent (Lemma 69), grown_transfer.py.** The native primitive $A_W^e(x) =$
4780 $\sum \lambda_n e^{-n/x}$ (no clipped edge), $x \in [10, N/40]$, $N = 2 \times 10^5$: $\hat{\theta}^e = 0.002$ for Δ — *bounded*, plateau
4781 0.73, the automorphy-forced convergence $A^e \rightarrow L(0)$ of Lemma 69(ii) observed directly — 0.290
4782 for $\operatorname{Sym}^5 \Delta$ and 0.403 for $\operatorname{Sym}^{13} \Delta$ (every sharp margin widened), detuned 0.065 (finite Euler
4783 modification: still essentially bounded), random 0.581 (smoothing manufactures no cancellation;
4784 the gate bites). The sharp-cutoff Chandrasekharan–Narasimhan barrier at degree ≥ 4 binds
4785 the clipped primitive only: measured, the clip was the barrier. Window-drift of the grown
4786 slope (`grown_transfer.py -median, 4 log-windows`): Δ 0.040/0.008/0.002/0.000 (plateau); Sym^5
4787 late windows 0.252/0.050 (equilibrating — the global 0.29 is a finite-window transient); Sym^{13}
4788 oscillatory, not yet equilibrated at $N = 2 \times 10^5$ (fastest channels $e^{\pm 13i\theta}$); random 0.43/0.58/0.76/0.49
4789 — about $\frac{1}{2}$ in every window, never equilibrating (Remark 70).

- 4790 • **Adapted closure is arithmetic-blind (Lemma 73 Step 2, §39), detuned_closure_control.py.**
4791 The register entry that localizes the arithmetic inside the continuation leg: the true Δ anchor, a
4792 *detuned* Δ ($\theta_2 \leftrightarrow \theta_3$ swapped, multiplicatively rebuilt), and a fully *random*-angle fiber of the same
4793 duality-stable class all force-close 30/30 native cells at the float64 floor (medians $3.2\text{--}36 \times 10^{-15}$,
4794 weights ≤ 2.91). The random-angle control shows that forced cell closure alone cannot deliver con-
4795 tinuation of the ordinary readout. The arithmetic-sensitive step is the *inverse-warp transfer* of §37.
4796 What is machine-checked by `TransferContinuation.transfer_analytic` is continuation for the
4797 coefficient sequence whose own primitive is bounded. The valid n -dependent inverse-kernel in-
4798 terface is machine-checked in `InverseWarpKernelTransfer.inverseWarp_transfer_tendsto`;
4799 applying it requires its summable first-difference and vanishing-boundary conditions for the
4800 selected fiber warp. Cell closure from `ForcibleClosure.residual_forcible` supplies neither
4801 condition by itself.

- 4802 • **Twisted CPS instances (§39.1), twisted_cps_instance.py.** The objects Theorem 39 actually
4803 consumes, tested directly. *Factorizing control*: the degree-12 twisted fiber $L(\operatorname{Sym}^5 \Delta \times \Delta)$, built
4804 on the carrier, reproduces the standard-side product $L(\operatorname{Sym}^6 \Delta) L(\operatorname{Sym}^4 \Delta)$ coefficientwise to
4805 1.8×10^{-14} — the twisted pipeline validated against known quantities. *Non-factorizing instance*:
4806 $L(\operatorname{Sym}^5 \Delta \times E_{11})$, degree 12 with two independent Satake angles, does not reduce to symmetric
4807 powers of one form; no channel is DC (every channel carries an odd θ_Δ -weight), and the multi-rail
4808 bank closes to precision (from 4×10^{-28} at 30 digits to 9×10^{-88} at 90) — entireness on the twisted
4809 family itself, not only the standard side.

- 4810 • **The Brauer-zero trap (§42), brauer_zero_trap.py (Sage).** The adversarial removal test: the
4811 Hamilton class $(-1, -1)_{\mathbb{Q}}$ has $\sum_v \operatorname{inv}_v = 0$ (aggregate readout closed) yet $[A] \neq 0$. Five passes, blind
4812 (no splitting field given): detect the retained obstruction despite the zero sum; classify it 2-torsion
4813 at $\{2, \infty\}$; synthesize the minimal splitting chart $\mathbb{Q}(\sqrt{-3})$ from the syndrome; verify $\operatorname{res}[A] = 0$
4814 by direct Brauer computation; certify $R=0$ and $\operatorname{Obs}=0$ jointly. A second class ramified at $\{3, 5\}$

separates a hard-coded “adjoin $\sqrt{-1}$ ” (splits Hamilton, fails $\{3, 5\}$) from the syndrome search (repairs both). Every stage cross-checked against Sage’s quaternion-algebra/Hilbert-symbol oracle; the removal logic (zero aggregate $\neq$ removal; even local degree kills a 2-torsion class; certificate = the conjunction) is machine-checked in `RequestProject/BrauerZeroTrap.lean`.

- **Genuine non-self-dual GL(3) (§31, table row), `f21_gl3_multirail.py` (data via Sage `idealfrobenius`) and `RequestProject/F21NonSelfDual.lean`.** The degree-three complex Artin representation of $F_{21} = C_7 \rtimes C_3$ (field $x^7 - 14x^5 + 56x^3 - 56x + 22$). Order-seven Satake $\{\zeta_7, \zeta_7^2, \zeta_7^4\}$ (Gauss period η) or its conjugate (η'); classes at $p = 13, 41$ (both $\equiv 6 \pmod{7}$) genuinely split η'/η . Multi-rail cell closure of the non-DC rails tracks precision, 1.5×10^{-30} (30 digits) to 4.3×10^{-62} (60), $\prod$ rails $= \zeta_7^{1+2+4} = 1$. Lean: exact cell closure of both classes, $\eta + \eta' = -1$, $\eta\eta' = 2$, $\eta \neq \eta'$, and $\text{Re } \eta = \text{Re } \eta'$ with $\eta \neq \eta'$ (the scalar obstruction).
- **Forcible cell closure (Lemma 63), `forcible_closure.py` and `RequestProject/ForcibleClosure.lean`.** On *native* growth cells — uniform in $y = \log Z$, each phasor entering at zero magnitude through the canonical C^∞ window `focal_closure.growth_window`, no clipped edge — exact Gauss–Newton over the coherent generators closes the nonlinear cell residual to the float64 floor on *every* cell: with $(\Omega, \omega, \log n, \Theta)$, $\Theta(n) = \sum_{p^k \parallel n} k \theta_p$ the fiber’s own local periods, 30/30 cells close on each of Δ , $\text{Sym}^5 \Delta$, $\text{Sym}^{13} \Delta$ — median $|D_C| = 4.9 \times 10^{-15}, 1.4 \times 10^{-14}, 7.8 \times 10^{-15}$, max relative residual $\leq 3.6 \times 10^{-14}$, weights $|x| \leq 1.28, 3.07, 2.27$ (integer-generator phases mod 2π). Three generators alone close 30/30 on $\Delta, \text{Sym}^5 \Delta$ and 29/30 on $\text{Sym}^{13} \Delta$; the one rank-deficient cell is closed by adjoining Θ — the resisting cell naming its adapter. One cell’s weights leave the next cell’s residual at its own scale (10–38% of that cell’s mass) — cross-cell independence. The sharp-window clip control (retained in the script) manufactures 1–3 resistant cells per fiber with exploding weights: the never-clip method law made measurable. Lean `residual_forcible`: two $\mathbb{R}$ -independent directions span $\mathbb{C}$, so any residual is forcible to zero; `CarrierReachability.reachable_independent_stage`: a continuous non-constant lane phase attains $\sin(2\phi) \neq 0$ (a connected image cannot lie in the discrete lock $\frac{\pi}{2}\mathbb{Z}$), supplying the pair — the two are composed by direct application, no further Lean theorem claimed.
- **Carrier-transport functoriality (§40), `RequestProject/CarrierFunctoriality.lean` and `composition_test.py`.** `faithful_comp` (the composite of closure-preserving transports preserves closure, *no* axioms) with associativity, two-sided identity, and base change $\text{bc}_p \circ \text{bc}_q = \text{bc}_{pq}$; the script verifies exact plethysm splitting $\text{Sym}^2 \circ \text{Sym}^2 = \text{Sym}^4 \oplus \text{Sym}^0$, $\text{Sym}^3 \circ \text{Sym}^3 = \text{Sym}^9 \oplus \text{Sym}^5 \oplus \text{Sym}^3$.
- **Tensor-induction residual (functorial preservation, §39), `tensor_induction_check.py`.** On real Δ Satake: the Clebsch–Gordan factorization $a_{\pi \times \text{Sym}^4} = a_{\text{Sym}^5} \star a_{\text{Sym}^3}$ holds coefficientwise to 1.4×10^{-14} ; $\text{mean}(a_{\text{Sym}^5}) \rightarrow 6 \times 10^{-5}$ while $L(\text{Sym}^3)(1) = 0.652 \neq 0$ (the trivial-rep control gives mean 0.652, so the deduction is non-vacuous).

References

- [1] R. P. Langlands, *Beyond endoscopy*, in *Contributions to Automorphic Forms, Geometry, and Number Theory* (Shalika 60th birthday volume), Johns Hopkins Univ. Press, Baltimore (2004), 611–697; the proposal to isolate functorial transfers through the poles of $L(s, \pi, \rho)$ inside the trace formula.
- [2] S. A. Altuğ, *Beyond endoscopy via the trace formula, I*, *Compositio Math.* **151** (2015).

- [3] S. A. Altuğ, *Beyond endoscopy via the trace formula, II*, Amer. J. Math. **139** (2017).
- [4] S. A. Altuğ, *Beyond endoscopy via the trace formula, III*, J. Inst. Math. Jussieu **19** (2020).
- [5] A. Venkatesh, “*Beyond endoscopy*” and special forms on $GL(2)$, J. reine angew. Math. **577** (2004), 23–80; the standard-representation and symmetric-square cases carried out — the two productive cases Sarnak’s boundary marks the end of.
- [6] S. Gelbart and H. Jacquet, *A relation between automorphic representations of $GL(2)$ and $GL(3)$* , Ann. Sci. ÉNS **11** (1978).
- [7] P. Sarnak, *Comments on Robert Langlands’ lecture “Endoscopy and Beyond”*, notes (2001), publications.ias.edu/sarnak.
- [8] L. Hörmander, *The Analysis of Linear Partial Differential Operators I*, 2nd ed., Grundlehren **256**, Springer (1990); Thm. 7.7.1 (non-stationary phase).
- [9] R. Wong, *Asymptotic Approximations of Integrals*, Classics in Applied Mathematics **34**, SIAM (2001); Ch. 3 (Watson’s lemma and the Mellin method).
- [10] J. Tate, *Fourier analysis in number fields and Hecke’s zeta-functions*, thesis (1950), in J. W. S. Cassels and A. Fröhlich (eds.), *Algebraic Number Theory*, Academic Press (1967), 305–347.
- [11] R. Godement and H. Jacquet, *Zeta Functions of Simple Algebras*, Lecture Notes in Mathematics **260**, Springer (1972).
- [12] A. Weil, *Sur les courbes algébriques et les variétés qui s’en déduisent*, Actualités Sci. Ind. **1041**, Hermann, Paris (1948); the Riemann hypothesis for curves over a finite field — the Frobenius eigenvalues on the Jacobian have absolute value $q^{1/2}$, placing the zeros of the congruence zeta function on $\text{Re } s = \frac{1}{2}$.
- [13] P. Deligne, *Les constantes des équations fonctionnelles des fonctions L* , in *Modular Functions of One Variable II* (Antwerp), Lecture Notes in Math. **349**, Springer (1973), 501–597 (the local L - and ε -factors of a Weil–Deligne representation); *La conjecture de Weil. I*, Publ. Math. IHÉS **43** (1974), 273–307 (the Riemann hypothesis for smooth projective varieties over finite fields); and . . . *II*, Publ. Math. IHÉS **52** (1980), 137–252 (purity: the normalized Frobenius eigenvalues lie on the unit circle).
- [14] J. W. Cogdell and I. I. Piatetski-Shapiro, *Converse theorems for GL_n* , Publ. Math. IHÉS **79** (1994), 157–214; *II*, J. reine angew. Math. **507** (1999), 165–188.
- [15] F. Shahidi, *Eisenstein Series and Automorphic L -functions*, Amer. Math. Soc. Colloquium Publications **58**, AMS (2010); and *A proof of Langlands’ conjecture on Plancherel measures; complementary series for p -adic groups*, Ann. of Math. **132** (1990), 273–330.
- [16] H. Kim, *Functoriality for the exterior square of GL_4 and the symmetric fourth of GL_2* , with Appendix 2 by H. Kim and P. Sarnak, J. Amer. Math. Soc. **16** (2003), 139–183 (the bound $|\alpha_p| \leq p^{7/64}$).
- [17] H. Kim and F. Shahidi, *Functorial products for $GL(2) \times GL(3)$ and the symmetric cube for $GL(2)$* , Ann. of Math. **155** (2002), 837–893 (for the symmetric fourth, see [16]).
- [18] J. Newton and J. A. Thorne, *Symmetric power functoriality for holomorphic modular forms, I & II*, Publ. Math. IHÉS **134** (2021), 1–116, 117–152.

- [19] J.-P. Serre, *Abelian ℓ -adic Representations and Elliptic Curves*, Benjamin (1968); Ch. I, the equidistribution criterion for the Sato–Tate law from the analyticity and non-vanishing of the symmetric-power L -functions.
- [20] H. Jacquet and J. A. Shalika, *A non-vanishing theorem for zeta functions of GL_n* , *Invent. Math.* **38** (1976), 1–16; non-vanishing on $\operatorname{Re} s = 1$ from the positivity of the Rankin–Selberg convolution; and *On Euler products and the classification of automorphic representations*, *Amer. J. Math.* **103** (1981), the local bound $|\alpha| < p^{1/2}$.
- [21] W. Luo, Z. Rudnick, and P. Sarnak, *On the generalized Ramanujan conjecture for $GL(n)$* , in *Automorphic Forms, Automorphic Representations, and Arithmetic*, *Proc. Sympos. Pure Math.* **66**, Part 2, AMS (1999), 301–310; the bound $|\alpha| \leq p^{1/2-1/(n^2+1)}$ toward Ramanujan.
- [22] E. C. Titchmarsh, *The Theory of Functions*, 2nd ed., Oxford University Press (1939), §§5.6–5.65 (Phragmén–Lindelöf).
- [23] B. J. Birch and H. P. F. Swinnerton-Dyer, *Notes on elliptic curves. II*, *J. reine angew. Math.* **218** (1965), 79–108; the conjecture relating the analytic rank to the Mordell–Weil rank.
- [24] B. H. Gross and D. B. Zagier, *Heegner points and derivatives of L -series*, *Invent. Math.* **84** (1986), 225–320; the central derivative as the Néron–Tate height of a Heegner point (analytic rank one).
- [25] V. A. Kolyvagin, *Euler systems*, in *The Grothendieck Festschrift, Vol. II*, *Progr. Math.* **87**, Birkhäuser (1990), 435–483; finiteness of Sha and the rank bound completing Birch–Swinnerton-Dyer at analytic rank ≤ 1 .
- [26] S. Zhang, *Gross–Schoen cycles and dualizing sheaves*, *Invent. Math.* **179** (2010), 1–73; the Beilinson–Bloch height of the modified diagonal.
- [27] X. Yuan, S. Zhang, and W. Zhang, *Triple product L -series and Gross–Schoen cycles*, book manuscript; the central derivative of the triple-product L -function as a Gross–Schoen/Ceresa height.
- [28] G. Ceresa, *C is not algebraically equivalent to C^- in its Jacobian*, *Ann. of Math. (2)* **117** (1983), 285–291.
- [29] B. Harris, *Harmonic volumes*, *Acta Math.* **150** (1983), 91–123, and *Homological versus algebraic equivalence in a Jacobian*, *Proc. Natl. Acad. Sci. USA* **80** (1983), 1157–1158 — the Fermat-quartic Ceresa cycle detected by a nonzero harmonic volume.
- [30] S. Lavery, *The loss ledger and the cycle filtration: tower exhaustion on the helix carrier, and a program whose terminus is the Hodge conjecture*, companion preprint draft (2026); the depth dictionary, the harmonic Hodge frame, the grade-four program, and the Ceresa landing.
- [31] J. Laga and A. Shnidman, *Ceresa cycles of bielliptic Picard curves*, arXiv:2312.12965 (2024); the Beilinson–Bloch height proportional to the Néron–Tate height of an explicit elliptic point.
- [32] B. Riemann, *Ueber die Anzahl der Primzahlen unter einer gegebenen Grösse*, *Monatsberichte der Königlich Preussischen Akademie der Wissenschaften zu Berlin* (November 1859), 671–680.
- [33] A. M. Odlyzko, *Correspondence about the origins of the Hilbert–Pólya conjecture* (letters exchanged with G. Pólya, 1981–1982), <http://www.dtc.umn.edu/~odlyzko/polya/> — the written record of the suggestion; no canonical formal statement exists.

- 4932 [34] H. L. Montgomery, *The pair correlation of zeros of the zeta function*, in: Analytic Number Theory
 4933 (St. Louis, 1972), Proc. Sympos. Pure Math. **24**, Amer. Math. Soc. (1973), 181–193.
- 4934 [35] M. V. Berry and J. P. Keating, *The Riemann zeros and eigenvalue asymptotics*, SIAM Review **41**
 4935 (1999), 236–266.
- 4936 [36] A. Connes, *Trace formula in noncommutative geometry and the zeros of the Riemann zeta function*,
 4937 Selecta Math. (N.S.) **5** (1999), 29–106.