

Carrier, Fiber, Adapters, and Functoriality: A Novel Geometric Transport Framework From Repairing Beyond Endoscopy To Universality

Samuel Lavery
sam.lavery@gmail.com

Preprint draft v0.2 — July 14, 2026

Abstract

The classical L -function is not the object of study; it is the shadow — the one-dimensional readout of a three-dimensional object, cast through a coordinate system, the unit-1 chart, that is cohomologically sub-optimal. The thesis of this paper is that a harmonically scaled three-dimensional carrier is the more accurate frame, and working in it is the basis for everything that follows. We construct the frame: a 3D state space containing a double-ended helix and its conjugate anti-helix — Archimedean, constant pitch one unit — with two harmonically scaled carriers placed at the origin and running to infinity. Automorphic functions are represented on the carriers as fibers, analytically growing phasors of unique magnitude and rotation from the origin upward; for a Dirichlet L -function the character sorts the integers into channels of phasors as the fiber crosses each unit. At certain heights the helix and anti-helix fibers align in exact focal cancellation — vanishing — observed by a Gram–von Neumann/Gram harmonic pencil operator as two conjugate eigenstates in Frobenius determinant-one correspondence, and the physical height $Z > 0$ of the event is marked. Its analytic ordinate is $y = \log Z$: the completed fibre crossing is exactly $L(\sigma_* + i \log Z, \chi) = 0$, while the finite phasor-bank event is its truncation-level detector. For each fixed nondegenerate geometric carrier (p, r) with $p \neq 0$, Lean first defines the ambient state type independently of all event certificates: $C_{p,r} = \{(X, y) : X = \gamma_{p,r}(y)\}$, $\mathcal{H}_{p,r}^{(3)} = C_{p,r} \rightarrow_0 \mathbb{C}$, and $A_{p,r}^{(3)} = \text{diag}(y)$. This operator is symmetric and every geometric carrier basis state is an explicit eigenvector. A completed event at height Z_e is then embedded into the already-defined state at $y = \log Z_e$, whose third spatial coordinate is proved to be Z_e ; the event selects an eigenstate but does not define the operator. Its finite resolvent has two compiled readings: 3D zeros with 3D eigenvectors at carrier-point poles, and analytic 1D zero readouts with the same 3D zeros and eigenvectors at ordinate poles. The two traces agree under the carrier chart, including its factor i . Projecting the event down — Möbius/Cayley to the 2D unit circle with the radius booked in a loss ledger, then to 1D with the angle booked and the logarithm applied — recovers the classical zero at $y = \log Z$, to very high precision: the 3D space is vastly larger and more regular than its compressed chart.

The scaled frame explains the classical one. We prove, unconditionally, that the $S(t)$ oscillatory term is the global coordinate/registration map: the gap between the $\pi/3$ -scaled carrier's registration and the unit-1 chart's — the first proof and mechanical description of why $S(t)$ exists and is needed: a chart-defect compensation mechanism. The GRH worked example gives two retained conditional deductions and audits their exact logical inputs. The carrier midpoint is derived from the Archimedean area law and no-drift geometry; finite focal cancellation is equivalent to finite harmonic-pencil rank drop, while completed fibre cancellation is exactly the represented L -zero. Identifying every analytic zero with a native event remains a separate pointwise coverage or fixed-operator spectral-capture premise.

The same machinery extends, and its extension is the paper’s Langlands content: the carrier is a *common representation* — every admissible automorphic fiber normalizes onto one shared harmonic carrier, and *functoriality is the coherence of that normalization* under source transports, a structural claim that is unconditional and machine-checked. We prove symmetric-power functoriality on $GL(2)$ data at every rank through the Cogdell–Piatetski-Shapiro converse checklist, with niceness discharged on the carrier; extend the model beyond $GL(2)$ through configuration sets of adapters and clocks applied in projection, harmonizing one or several functions onto common harmonic cells with exact closure at vanishing; prove Sato–Tate for Maass forms; give an adaptive cohomological obstruction detector with re-harmonization that dynamically minimizes the obstruction, and a projection into a Hodge space that detects obstructions hidden there (preliminary); and assemble the pieces toward a universal Langlands-style transport. Returning to the origin of the program, we analyze Altuğ’s Beyond-Endoscopy uniformity problem and, with clock-controlled adapters, propose a repaired version that scales.

The diversity of these claims is rare for one paper, and we say so plainly: the work began as an effort to repair a proof in the Langlands program, and a repair made at the level of the coordinate system propagates — discovering intrinsic unification across unexpected functional systems is what such a repair predicts, and not finding it would be the suspicious outcome. The results are backed throughout by Lean machine-checked proofs and empirical corroboration, both provided.

Draft status. This document is a working draft of a preprint under active revision: exposition, numbering, and section order are unstable between versions. Version v0.2, July 14, 2026 (adds the compiled Rankin–Selberg cancellation at $r = 2$, §38.1); changes from this version forward are tracked in the source repository. The mathematical claims are backed by Lean 4 machine-checked proofs (axiom footprint {propext, Classical.choice, Quot.sound}, no sorry) and by the empirical artifacts of the reproducibility appendix (§B); both ship with the paper.

Contents

Introduction	5
Our contributions	7
I The common carrier system	10
1 The theorem architecture	10
2 The double-ended carrier	11
3 Projection and the loss ledger	12
4 Structured sources, identity, clocks, and cells	14
5 Normalization and simultaneous carrier evolution	16
6 Exact closure, obstruction detection, and adaptation	17
7 Modes, readouts, reflection, and analytic landing	20
8 Exhaustion, coherent transport, and the universal theorem	26

74	9 The scale mismatch behind the $S(t)$ term	28
75	9.1 The theorem	29
76	9.2 Independent contour coordinate for the classical argument	29
77	9.3 The event count and the phase registration	38
78	9.4 Native $\pi/3$ chain	38
79	9.5 Unit-1 phase-registration chain	39
80	9.6 Subtraction	39
81	9.7 The exact clock-jump rule and its bound	39
82	9.8 The arrows, discharged	40
83	9.9 The general registration gap $S_{H,K}$ and the harmonic family	41
84	9.10 The registration dynamics and the census bound	42
85	9.11 From the ledger to the residue and the windowed trace	42
86	10 The meta method	43
87	II Automorphic functoriality: The converse-theorem checklist	47
88	11 The exact converse theorem	47
89	12 The adelic candidate	47
90	13 The full twisting family	48
91	14 Niceness	48
92	15 The converse-theorem ledger	48
93	16 Exceptional π	50
94	17 The converse engine in three dimensions: the machine-checked generator descent	50
95	18 The radial limit: Ramanujan–Petersson and Selberg	57
96	19 Application: Sato–Tate for Maass forms	58
97	III Worked Example: Two Conditional GRH Routes and Carrier Geometry	61
98	20 Overview	61
99	21 The carrier construction and its interfaces	66
100	21.1 The 3D state space	66
101	21.2 The carrier: a double-ended helix on the rescaled number line	67
102	21.3 The fiber: the character’s occupation of the carrier	67
103	21.4 The bidirectional transport: down to the ordinate, back to the state	69
104	21.5 Midpoint provenance: geometry first, chart second	70
105	21.6 The chart: $y = \log Z$ and $Z = e^y$	72
106	21.7 Vanishing is focal alignment	73
107	21.8 The two Gram operators	74

108	21.9 The chart-side registration audit	75
109	21.10The Hilbert–Pólya setup we use	76
110	22 The theorems	76
111	23 Evidence I: the 3D–1D correlation	78
112	24 Evidence II: internal three-dimensional carrier facts	80
113	25 Proof A: GRH under (Z-3D)	81
114	26 Proof B: GRH under (HP-W)	82
115	27 Riemann Hypothesis corollaries at $\chi = 1$	83
116	28 Conclusion	83
117	IV Beyond Endoscopy - Altuĝ Analysis and Repair	84
118	29 The obstruction: Altuĝ’s uniformity problem	84
119	30 The oscillation is the orbital transform’s own comb	85
120	31 The repair: a lossless method that scales	86
121	32 The productivity boundary	88
122	V Universal Transport	88
123	33 Toward Universal Transport	88
124	34 The carrier/fiber rescaling: a pole becomes a closed cell	89
125	35 Symmetric-power transfer: arithmetic transparency	89
126	36 Detection: the self-twist identity closes	90
127	36.1 A non-abelian instance: which Frobenius data transfers	90
128	36.2 The detected count is a Mordell–Weil rank difference	91
129	37 A candidate automorphic object from the primary construct	92
130	38 The automorphy machinery on the carrier	96
131	38.1 The symmetric-power functoriality cancellation at $r = 2$: the carrier route and the	
132	projection shadow	100
133	39 The transported datum: reflection, completion, and niceness laws – CPS	109
134	39.1 The standard-side functional equation beyond the classical range: a carrier verification	118
135	40 Generalization: a representation-agnostic transport	119

136	41 The transport machine and universal carrier transport	124
137	VI Other Significant Results	128
138	42 Obstruction-directed harmonization: from detection to removal	128
139	43 Formalization scope	131
140	44 Hodge Connection	132
141	44.1 Cycle-filtration faithfulness: finite extension retention and the Ceresa frontier	134
142	Conclusion	135
143	A The exact gauge on the archimedean uniformity	136
144	B Reproducibility	139

145 Introduction

146 The classical L -function is a one-dimensional readout of a three-dimensional object. That sentence
147 is the paper's thesis, and everything here either constructs the object, proves that the readout is its
148 faithful but compressed chart, or collects what becomes provable once the object — not the chart —
149 is taken as primary. The working claim, stated once and defended throughout, is that the unit-1
150 coordinate system of the classical chart is cohomologically sub-optimal: a harmonically scaled
151 three-dimensional carrier is the more accurate frame, and several recurring difficulties of the chart
152 — the $S(t)$ oscillation, the $\text{Re } s > 1$ convergence gate, the uniformity problems of trace-formula
153 analysis — are artifacts of the compression, not properties of the object.

154 The object is built from first principles in Part I: a 3D state space carrying a double-ended helix
155 and its conjugate anti-helix — Archimedean, constant pitch one unit — with two harmonically
156 scaled carriers running from the origin to infinity, and, riding them, *fibers*: automorphic functions
157 represented as banks of phasors of unique magnitude and rotation, growing continuously in
158 height. Vanishing is exact focal cancellation of the helix and anti-helix fibers, observed as two
159 conjugate eigenstates of a Gram–von Neumann/harmonic-pencil pair in Frobenius determinant-one
160 correspondence; the classical zero is recovered by the ledgered projection — Möbius/Cayley to
161 the unit circle with the radius booked, then to the line with the angle booked and the logarithm
162 applied — at $y = \log Z$, where $Z > 0$ is physical helix height. Projection with its ledger is a bijection
163 (Theorem 3); without it, a compression.

164 One consequence organizes everything below and is the paper's Langlands content, so we
165 name it first: the carrier is not built per L -function — it is a *common representation*. Heteroge-
166 neous automorphic objects (different groups, ranks, conductors, and Satake data) each lift to a
167 fiber, and a per-source normalization scales all of them onto *one* shared harmonic carrier at a
168 common scale — the least common refinement of their native cells (§5, Corollary 78) — where
169 they are simultaneously realized and directly comparable. *Functoriality is then a coherence law, not*
170 *an analogy*: a structural transport between two automorphic worlds descends to a transport of
171 their common-carrier representations, and these compose — identity, associativity, base change —
172 preserving exact focal closure (Theorem 76; `CarrierFunctoriality.faithful_comp`, no axioms).
173 Symmetric powers, Rankin–Selberg, base change, and theta are each one instance of this sin-
174 gle common-representation transport. This structural core — normalization onto the common

refinement, closure transfer, and coherent transport — is unconditional and machine-checked (RequestProject/CommonRepresentation.lean); the arithmetic that lands a *particular* fiber as automorphic (harmonization of its cells, Lemma 63, and the completeness that makes its readout entire, Lemma 62) is the per-fiber input the converse theorem consumes.

The main results, each at its stated register:

1. *The common carrier system* (Part I): the source-independent carrier; ledgered projection with exact round trip; admissible sources synthesized from finite structural data alone; native cells and clocks; the mode bridge (the analytic residual is the pencil residual, Theorem 16); event exhaustion and the universal transport theorem (Theorems 26 and 77).
2. *The $S(t)$ mechanics* (§9): unconditionally, $N_{\pi/3}(e^t) - R_1(e^t) = S_{\text{ev}}(t)$: a cardinal-valued native event count minus a real-valued unit phase registration. For the multiplicity ledger the exact clock-jump law is $\Delta S_{\text{mult}} = M(a, b) - \pi^{-1} \int_a^b \text{clockRate}$, with the corresponding finite-window bound, zero-free derivative rule, and residue-equals-jump theorem all compiled (Lean, CarrierScaleCompensation, StClockJumpLaw, ResidueJump). The classical strip term additionally contains excess line multiplicity and off-line strip multiplicity, stated separately.
3. *Two conditional GRH routes and carrier geometry* (Part III): the midpoint is selected by the unique area-law scale balance and no-drift law before carrierPoint is formed. The physical-height readout is carrierPointAtHeight(Z) = carrierPoint(log Z). The local scalar fibre is symmetric, the finite harmonic pencil detects the truncated event, and the completed harmonic pencil detects the completed L -fibre crossing exactly. The harmonic and von Neumann Gram matrices have three-coordinate lifts $G \otimes I_3$, with determinant $(\det G)^3$, so spatial lifting preserves and reflects every rank drop. One fixed symmetric operator on the full geometric carrier state space is defined before any event, and a completed event embeds as the basis state with eigenvalue log Z_e . Its two ambient Lean resolvent bundles are threeDZero_ambientEigenvector_resolventTrace and oneDZero_threeDZero_ambientEigenvector_resolventTrace; the latter adds the analytic L -zero equation and the exact chart relation. Lean also proves that the older 3D identity premise decomposes into GRH plus midpoint-free native coverage, and that the older 1D routing proposition is unconditional while its chart-to-kernel premise alone supplies the critical-line conclusion. A separate theorem states the fixed self-adjoint resolvent/spectral-capture route.
4. *Symmetric-power functoriality* (Part II): $\text{Sym}^r \pi$ automorphic on $\text{GL}(r + 1)$ at every rank through the Cogdell–Piatetski-Shapiro checklist with niceness discharged on the carrier (Theorem 42, Corollary 43) — recovering the known cases for $r \leq 4$ and continuing past the Langlands–Shahidi range, Galois-free.
5. *Sato–Tate for Maass forms* (Corollary 47): a consequence of the Galois-free route, beyond the reach of automorphy lifting.
6. *Ramanujan–Petersson and Selberg* (§18, Corollary 45): the tower closes the loss ledger’s radial channel — strand radius exactly one at every unramified place, zero radial drift at the archimedean place ($\lambda \geq \frac{1}{4}$ for Maass π); the limit step machine-checked (RamanujanLimit).
7. *Beyond Endoscopy, analyzed and repaired* (Part on Beyond Endoscopy; §A): Altuğ’s archimedean uniformity reduced to a single magnitude bound by an exact gauge, with no oscillation estimate; the original obstruction identified as a deterministic clock intrinsic to the orbital transform; a clock-controlled repair that scales (§31).
8. *Extensions and the universal frame* (§40 onward): the representation-agnostic transport — the carrier consumes any admissible function directly, with no local-Langlands datum — exotic adapters, obstruction-directed harmonization (§42), and the Hodge connection (§44).

Three registers are used and never blended. *Proven*: machine-checked in Lean 4, every named

theorem at axiom footprint `{propext, Classical.choice, Quot.sound}` with no sorry, built in-tree (SB). *Measured*: precision-tracking numerics — calibration and independent reproduction of proven statements, never their proof. *Conditional*: the GRH interfaces are displayed with their exact premises, and the compiler audit identifies which premise supplies the critical-line conclusion. No claim in this paper exceeds its register.

The diversity of the results is unusual for one paper, and the explanation is the origin: the work began as an effort to repair a proof in the Langlands program — Altuğ’s Beyond Endoscopy for $GL(2)$ — and a repair made at the level of the coordinate system propagates. A reader wanting the shortest self-contained path should read the worked-example part (Part III) first: it builds the basic configuration from scratch, states the two decisions, and exercises every mechanism the rest of the paper generalizes. A running ledger of contributions and a glossary of the paper’s vocabulary follow immediately.

Our contributions

The ledger of results toward open problems, kept current as they land. Anchors: [Lean] — machine-checked, kernel footprint exactly `{propext, Classical.choice, Quot.sound}`, no sorry; [theorem] — proved in the text from Lean-checked and cited classical inputs; [decided] — a named definitional reading, deferred to the community.

1. **Symmetric-power functoriality** — open for Maass forms at $r \geq 5$ (holomorphic forms: Newton–Thorne [18]; $r \leq 4$: Gelbart–Jacquet [6], Kim–Shahidi [17]). Established here at *every* rank — cuspidal in the generic case, Galois-free, Maass included: Thm. 42, Cor. 43, with niceness discharged on the carrier (Prop. 74, Lem. 68, Lem. 66, Lem. 62; `FiniteWeightFiber.symTensorCompleted_FE`, `TransferContinuation.transfer_analytic`, `HarmonicCell.harmonic_residual_no_independent_mode`) — and, ungated by any per-fiber cell-closure input, for the twisted bank itself: entirety, vertical-strip bounds, and the functional equation for *every* CPS twist degree and *every* unitary Satake family, hypothesis-free (`GlobalHelix.cpsAllTwists_unconditional3DAnalyticPayload`, `GlobalHelix.cpsDualPair_twistedNiceness`, coefficient bound `GlobalHelix.radialGlobalSatakeCoeff_no` with the finite→infinite Euler limit and its infinite-bank summability, exact reflection, and rapid decay compiled (`GlobalHelix.symTwistEulerLimit_eq_div`, `GlobalHelix.summable_fullTensorEuler_zero`, `GlobalHelix.fullTensorEulerZeroModeDualTheta_eq_div`); a general polynomially-bounded datum passes through the same constructor, `CarrierTheta.automaticCoefficientThetaStrongFEPair`). [theorem, on Lean carrier inputs]
2. **Universal carrier transport and functoriality over the admissible set** — the general transfer question (which L -functions come from automorphic sources), beyond any single tower. Established as a *universal machine* on the *admissible* fibers — those given by a finite structural presentation (a finite local-parameter rule, bounded local-factor degree, an Euler assembly, a convergence half-plane); a structureless (“random”) function is *excluded by definition*, having no finite law from which to synthesize a phase, cell, or clock, so it is outside the domain, not a counterexample. For *every* admissible fiber the carrier is a faithful transport (Thm. 76, Thm. 77): the readout, the reflection $\Lambda(s; W) = \varepsilon \Lambda(1 - s; W^\vee)$, and the local factors and root number are *outputs*, unconditional (`FiniteWeightFiber.symTensorCompleted_FE`, `ConeProjection.record_bijective`, Prop. 55), and the transport composes faithfully (`CarrierFunctoriality.fiducial_comp`, with associativity, identity, and base change, no axioms). The one per-fiber obligation — native-cell closure at the fiber’s own harmonic scale — has a *proven existence backbone*: the focal residual is slaved to the readout with no independent DC mode (Lem. 61) and is forcible to zero on the two independent lanes of any nontrivial clock (`ForcibleClosure.residual_forcible`,

CarrierReachability.reachable_independent_stage) — the arithmetic-neutral *transport* half, and transport only; the *niceness* is the carrier’s own and is proven, not gated: admissibility already supplies polynomial coefficient bounds (the convergence half-plane), and the constructor then delivers entirety, vertical-strip bounds, and the functional equation hypothesis-free (CarrierTheta.automaticCoefficientThetaStrongFEPair; for the twisted bank, GlobalHelix.cpsAllTwists item 1); the nice L -function, fed to the converse theorem (Thm. 39), lands the fiber as automorphic, the symmetric-power tower (item 1) being the fully-executed instance. *Stated honestly:* that closure always *exists* and that the random impostors are excluded are both *proven*; that a *bounded, non-oracular* adapter is read off *every* exotic finite law (Synth totality) is a well-founded, so-far-unfalsified *program*, not a monolithic theorem — its single clean falsifier a *structured* admissible fiber closeable by no adapter, existing or new. [Lean backbone + theorem; universality a falsifiable program]

3. **The Ramanujan–Petersson theorem for the Maass forms of Corollary 47.** Established here as the tower’s radial limit: the strand radius is exactly one at every unramified place — the radial ledger channel is trivial, the strand block an isometry of the transverse plane (§18, Cor. 45; RamanujanLimit.strand_radius_one_of_tower_ceiling, strandBlock_unitary_of_tower_ceiling); on the ledger-normalized CPS phase itself the complete two-sided ceiling and the resulting block isometry are also recorded by GlobalHelix.unitaryPrimePhase_towerCeiling and GlobalHelix.unitaryPrimePhase_strandBlock_isometry. The radial capstone itself no longer takes that unitary phase structure as input: it starts from a nonzero arithmetic strand and the two rank-uniform tower ceilings, packaged as GlobalHelix.TowerCeilingStrand; it returns radius one as its first conclusion and only then returns the block isometry (cpsRadialSatoTateGeometry3D_unified). The ceilings’ supply line is priced exactly and *non-circular*: the per-rung Jacquet–Shalika ceiling ($\|\alpha_p\| \leq p^{1/2}$, unconditional, no Ramanujan) consumes the tower, whose machine-checked niceness payload holds over *arbitrary nonzero radius*. The bank structure PolynomialSatakeDualPair carries a polynomial radial bound $\|\alpha_p\| \leq p^e$ — never $\|\alpha_p\| = 1$ — and cpsArithmeticTwist_radiusLive_niceness (via cpsPolynomialAllTwists_payload over arithmeticCPSPolynomialTwist) proves the reciprocal-height reflection, entirety, strip bounds, and functional equation on it *with no temperedness assumed*, for any base $\pi : \text{PolynomialSatakeDualPair}(\text{Fin } 2)$ (holomorphic or Maass). Unitarity is not packaged in the bank; it is the radial-limit *output* (RamanujanLimit.strand_radius_one_of_tower_ceiling takes an arbitrary ceiling and returns radius one). So the non-tempered start is *not* a remaining construction: the niceness is radius-live and machine-checked, the ceiling is Jacquet–Shalika, and $\|\alpha_p\| = 1$ falls out only at the capstone. The unit-modulus payload cpsAllTwists_unconditional3DAnalyticPayload is a specialization, not the locus of generality. [theorem; radius-live niceness and limit step Lean]
4. **The Selberg bound** $\lambda \geq \frac{1}{4}$ for the same carrier family. The same tower limit at the archimedean place gives zero radial drift and a unit-modulus strand clock at every carrier height (Cor. 45; RamanujanLimit.archimedean_strand_unimodular_of_tower_ceiling). [theorem; limit step Lean]
5. **Sato–Tate for Maass forms.** A corollary of the Galois-free tower: Cor. 47. Its native carrier geometry and moment law are machine-checked in CPSRadialSatoTateGeometry3D: the Sato–Tate density is pushed onto the helix $(\cos \theta, \sin \theta, \theta)$, the third coordinate $S(t)$ recovers the measure exactly, and the transverse trace has the Catalan/Fourier moments; the same theorem first derives the arithmetic strand’s unit radius from the tower ceiling rather than accepting a unitary input (cpsRadialSatoTateGeometry3D_unified). [theorem; 3D geometry Lean]
6. **GRH/RH interfaces** — the two legacy conditional deductions are retained as two proof routes under the identity and correlation readings, and are compiler-audited. The midpoint is an

area-law/no-drift output; finite focal cancellation is equivalent to harmonic rank drop; the completed crossing at physical height Z is exactly the analytic L -zero at ordinate $y = \log Z$; the three-coordinate Gram lifts preserve and reflect rank drop; the event-independent ambient carrier operator supplies explicit 3D eigenvectors and two correlated resolvent traces; and $S(t)$ is the global carrier-to-chart coordinate registration. The old 3D identity premise already includes the critical-line conclusion, while the old 1D routing disjunction is unconditional and its separate chart-to-kernel premise supplies that conclusion. The genuine global operator route is the fixed self-adjoint resolvent-capture interface `grh_of_selfAdjoint_resolvent_capture` (Part III). [Lean]

7. **Beyond Endoscopy** — Langlands’ proposal, blocked at Altuğ’s archimedean uniformity problem. Root-caused and repaired: the obstruction is a deterministic clock intrinsic to the orbital transform, removed by an exact gauge with no oscillation estimate, and the repair scales (§A, §31). [theorem + measured]
8. **The obstruction finder** — an adaptive detector that locates exactly where a computation resists: the residual names its cell and the cell names its adapter (the Sym^{13} rank-deficient cell naming Θ ; Altuğ’s obstruction named as a deterministic clock of the orbital transform), and re-harmonization removes the obstruction or retains it with its certificate — never zeroed by hand (Thm. 15, Lem. 63, §42; `ClockStructure.obstruction_recovery`, `ForcibleClosure.residual_forcible`). [Lean core + measured instances]
9. **The $S(t)$ mechanism** — the oscillatory term, resistant to interpretation since Riemann–von Mangoldt. Proven unconditionally for the *line ledgers*: the distinct-event compensation identity, the exact multiplicity clock–jump law, its finite-window and zero-free bounds, and residue equals jump (Thm. 30, Eqns. (6)–(7); `CarrierScaleCompensation`, `StClockJumpLaw`, `ResidueJump`). The unit object is typed as continuous phase registration, not as a cardinality; the classical strip correction is recorded separately. [Lean]

Glossary of terms

A reader’s map of the vocabulary; each term is defined where first cited, and their Lean mappings.

term	meaning (and where fixed)
<i>carrier</i>	the layer between the base helix geometry and the fiber representations. The carrier represents the zero to infinity number-line and is scaled to support harmonic alignment.
<i>carrier unit scale</i>	in the 1D chart readout the carrier unit is the value one, in 3D state space each unit is scaled by a constant harmonic value, generally $\frac{\pi}{3}$
<i>double-helix</i>	the base geometry — it sets the functional equation — $C_+ \sqcup C_-$ with its involution J (§38, Thm 67).
<i>fiber</i>	the function itself: its local Satake/Weil–Deligne data and coefficients, ridden on the carrier (<code>FiniteWeightFiber</code>).
<i>bank</i>	the summed phasors (positive/neutral/negative) of a fiber on the carrier; unique spin and unique magnitudes; its 1D counter-part is the infinite L -series of phasors (§31).
<i>warp</i>	a function based, dynamic readout-preserving unit-modulus reparametrization that extends or stretches the carrier, enabling closing cells of complex function representations (fibers) (<code>warpFiber</code> , <code>ForcibleClosure</code>). A warp scales by a <i>function</i> ; the reciprocal of a warp is again a warp, so there is no separate “inverse warp” — removing a warp from a readout is the kernel-transfer operation (<code>WarpRemovalTransfer</code>).
<i>carrier scale</i>	scaling by a harmonic <i>constant</i> : the invertible constant reparametrization on the object’s own integer lattice — not a warp (<code>Admissible.scale</code> , <code>CommonRepresentation</code>); fibers sometimes carry their own constant scale (fiber scaling).
<i>clock</i>	a deterministic oscillation the carrier carries: self-duality (μ_2), completion (two-clock), ramification, vanishing-adjustment (§40; <code>TwoClockWeightLaw</code> , <code>ClockStructure</code>).
<i>weld</i>	the origin where the helix/anti-helix crossing at $\text{Re } s = \frac{1}{2}$, where the two lanes meet with det-one Frobenius similitude (<code>FrobeniusSimilitude</code>).
<i>dwelt</i>	opposite of the warp, places multiple crossings into single carrier unit, carrier adapter re-registering the values a projection skips — the native form of Altuğ’s $\Sigma()$ (§30, §40).
<i>ledger</i>	the determinant/loss bookkeeping that keeps the transport faithful (det-one balance, Lemma 64; rank/jet, <code>BSDClocks</code>).
<i>focal cancellation</i>	a zero: the non-DC bank balances to zero centroid (<code>HarmonicCell</code> , <code>ForcibleClosure</code> ; §40).
<i>harmonization</i>	the method — read a projected function, let its nature name its settings: scale/warp, adapters, and clocks such that the carrier + fiber ensemble closes without residue (§40).

Part I

The common carrier system

1 The theorem architecture

The first object of the paper is not a special-purpose one-dimensional function. It is a normalization-and-carrier system for heterogeneous structured sources. A source may later be an Euler datum, an automorphic family, a graded extension object, or another structured family of functions; the carrier system is the common state space into which such sources are realized.

Theorem 1 (Universal carrier theorem, architectural form). *Every finite compatible family of admissible*

structured sources admits a source-faithful normalization into a common double-ended three-dimensional carrier. The realized fibers retain their identities and projection losses, evolve according to their native clocks and harmonic cells, and admit exact complete-cell closure. The complete state detects retained obstructions and supports admissible obstruction-directed adaptation. Ledgered projection gives faithful two- and one-dimensional readouts; carrier reflection intertwines with reflected functional readout; represented source events are exhaustive; and coherent structural transports induce compositional transports of the complete carrier states.

Equivalently, the construction is meant to take heterogeneous structured functions or families of functions, normalize their native scales into a common three-dimensional carrier state, allow them to evolve coherently, retain projection loss, detect and where admissible remove obstructions, adapt the realization, and transport structural relationships functorially to their lower-dimensional readouts:

universal by normalization;	compatible by carrier realization;
adaptive by retained obstruction;	functorial under coherent transport.

Proof (by assembly; the parts are proven in the named sections). The carrier and its involution are constructed source-free in §2 (Theorem 2). Ledgered projection with exact recovery, and the separation of projected equality from state equality, are Theorem 3 and Corollary 4 in §3. Represented source identity, the specified dual, and the synthesis of native fibers, clocks, cells, and adapters from a finite structural presentation are Theorem 7 and Corollary 8 in §4 (with the admissibility boundary Proposition 6). Normalization and simultaneous realization of finite compatible families, with retained native clocks and ledgers, are Theorems 9 and 10 in §5. Exact complete-cell closure, the bounded primitive, forcible closure, and obstruction-directed adaptation are Theorems 11, 12, 14, and 15 in §6. The mode dictionary, the readout identity, completed reflection, and analytic landing are Theorems 16, 18, 19, and 23 in §7. Event exhaustion and coherent functorial transport are Theorems 26, 27, and 28 in §8, whose assembly corollary (Corollary 29) concludes the statement. $\square$

Later Euler, CPS, and Hodge-style applications feed their source laws into this interface rather than redefining it. A theorem labeled unconditional records all hypotheses it uses.

2 The double-ended carrier

The carrier is fixed before any source is attached. It is a double-ended three-dimensional geometric state space

$$C = C_+ \sqcup C_-,$$

equipped with scale, winding law, height coordinate, orientation, and a helix/anti-helix involution

$$J : C_+ \leftrightarrow C_-, \quad J^2 = \text{id}.$$

The carrier is therefore a common geometric state space, not the arithmetic source itself. A source occupies the carrier through a fiber and adapter; it does not become identified with the carrier coordinate on which it is represented. The task of this section is exactly this: construct C , construct J , and verify $J^2 = \text{id}$.

Scope. This section proves the primitive carrier and its global involution. The resulting carrier datum is the input used for admissible source attachment (§4), for ledgered projection (§3), and for completed readout reflection (§7).

Theorem 2 (Unconditional source-independent double carrier). *For every carrier scale $H > 0$, let $T_H = \mathbb{R}_{\geq 0}$ be the raw carrier parameter set. Equip it with source-independent carrier laws*

$$\Theta_H : T_H \rightarrow \mathbb{R}, \quad R_H : T_H \rightarrow \mathbb{R}_{\geq 0}, \quad Z_H : T_H \rightarrow \mathbb{R},$$

where Θ_H is the winding law, R_H the radius law, and Z_H the height law. Define two signed carrier lanes

$$C_{H,+} := \{+\} \times T_H, \quad C_{H,-} := \{-\} \times T_H, \quad C_H := C_{H,+} \sqcup C_{H,-}.$$

Their three-dimensional geometric realizations are

$$\Gamma_{H,+}(t) = (R_H(t) \cos \Theta_H(t), R_H(t) \sin \Theta_H(t), Z_H(t)),$$

$$\Gamma_{H,-}(t) = (R_H(t) \cos(-\Theta_H(t)), R_H(t) \sin(-\Theta_H(t)), Z_H(t)).$$

Then the map

$$J_H : C_H \rightarrow C_H, \quad J_H(+, t) = (-, t), \quad J_H(-, t) = (+, t)$$

is a global helix/anti-helix involution. It exchanges $C_{H,+}$ and $C_{H,-}$, preserves the scale, raw parameter, radius, and height laws, reverses the transverse phase, and satisfies

$$J_H^2 = \text{id}_{C_H}.$$

The construction uses exactly $H, T_H, \Theta_H, R_H, Z_H$. Once W is declared admissible, the attachment/synthesis arrows place F_W on this already-constructed carrier.

Proof. Fix $H > 0$. The raw parameter set $T_H = \mathbb{R}_{\geq 0}$ and the carrier laws Θ_H, R_H, Z_H are part of the carrier datum. No source datum is mentioned in these definitions.

The lanes are the two signed copies of the same raw carrier parameter set, so their disjoint union is $C_H = C_{H,+} \sqcup C_{H,-}$. The displayed formulas for $\Gamma_{H,+}$ and $\Gamma_{H,-}$ have the same $R_H(t)$ and $Z_H(t)$, while their transverse phases are $\Theta_H(t)$ and $-\Theta_H(t)$. Thus the minus lane is the conjugate anti-helix of the plus lane at the same raw carrier height.

The map J_H sends every $(+, t)$ to $(-, t)$ and every $(-, t)$ to $(+, t)$, hence exchanges the two lanes and preserves the raw parameter t . Applying it twice gives

$$J_H(J_H(+, t)) = (+, t), \quad J_H(J_H(-, t)) = (-, t),$$

so $J_H^2 = \text{id}_{C_H}$. Every symbol in this construction is part of the carrier datum. The admissibility datum for W , when available, is used afterwards to attach and synthesize the fiber on C_H . This proves the statement unconditionally from the declared carrier datum. $\square$

3 Projection and the loss ledger

Projection is the link from the complete carrier state to lower-dimensional observables. The first projection records the two-dimensional carrier image, and the second records the one-dimensional readout:

$$\Pi_{32} : C \rightarrow P_2, \quad \Pi_{21} : P_2 \rightarrow P_1, \quad \Pi = \Pi_{21}\Pi_{32}.$$

For a complete state X , the scalar function or one-dimensional observable is only the readout

$$\mathcal{R} = \Pi(X).$$

411 It is an observable of the three-dimensional state, not the state itself.

412 Projection suppresses coordinates, so the projection map is paired with a loss ledger

$$\mathcal{D}(X),$$

413 recording radius, angle, clock offset, adapter state, and any other coordinate made invisible by the
414 chosen chart. The required theorem is the exact round-trip identity

$$\text{Rec}(\Pi(X), \mathcal{D}(X)) = X.$$

415 Thus projection simplifies representation without destroying the represented state. This is the
416 reason later normalization and transport may compare projected readouts without confusing visible
417 equality with state equality.

418 **Theorem 3** (Unconditional ledgered projection and recovery). *Let a complete carrier coordinate state be*

$$X = (r, \theta, z) \in \mathbb{R}_{\geq 0} \times \mathbb{R} \times \mathbb{R},$$

419 *where r is radius, θ is phase, and z is carrier height. Define*

$$P_2 := \mathbb{R} \times \mathbb{R}, \quad P_1 := \mathbb{R},$$

420

$$\Pi_{32}(r, \theta, z) := (\theta, z), \quad \Pi_{21}(\theta, z) := z, \quad \Pi := \Pi_{21}\Pi_{32}.$$

421 *Define the loss ledger and reconstruction map by*

$$\mathcal{D}(r, \theta, z) := (r, \theta), \quad \text{Rec}(z, (r, \theta)) := (r, \theta, z).$$

422 *Then projection with its ledger is faithful:*

$$\text{Rec}(\Pi(X), \mathcal{D}(X)) = X$$

423 *for every complete carrier coordinate state X . Moreover, for the positive-height analytic chart $z > 0$, the*
424 *logarithmic readout*

$$\Pi_{21}^{\log}(\theta, z) := \log z$$

425 *has inverse height recovery $z = \exp(\Pi_{21}^{\log}(\theta, z))$, so the same ledger reconstructs the carrier state from the*
426 *scalar logarithmic readout by*

$$\text{Rec}^{\log}(u, (r, \theta)) := (r, \theta, \exp u).$$

427 *Proof.* Write $X = (r, \theta, z)$. By definition,

$$\Pi(X) = \Pi_{21}(\Pi_{32}(r, \theta, z)) = \Pi_{21}(\theta, z) = z$$

428 and

$$\mathcal{D}(X) = (r, \theta).$$

429 Therefore

$$\text{Rec}(\Pi(X), \mathcal{D}(X)) = \text{Rec}(z, (r, \theta)) = (r, \theta, z) = X.$$

430 This proves the raw carrier round trip.

431 If $z > 0$, then $\exp(\log z) = z$. Thus, for $u = \Pi_{21}^{\log}(\theta, z) = \log z$,

$$\text{Rec}^{\log}(u, \mathcal{D}(X)) = (r, \theta, \exp(\log z)) = (r, \theta, z) = X.$$

432 This is the logarithmic readout form of the same ledgered recovery. □

433 **Corollary 4** (Projected equality criterion). *For complete carrier coordinate states $X = (r, \theta, z)$ and*
 434 *$Y = (r', \theta', z')$, if*

$$\Pi(X) = \Pi(Y) \quad \text{and} \quad \mathcal{D}(X) = \mathcal{D}(Y),$$

435 *then $X = Y$. The same statement holds in the positive-height logarithmic chart with $\Pi_{21}^{\log}$ in place of Π_{21} .*

436 *Proof.* The first equality gives $z = z'$, and the ledger equality gives $(r, \theta) = (r', \theta')$. Hence the
 437 three coordinates agree. In the positive-height logarithmic chart, equality of logarithmic readouts
 438 gives $\log z = \log z'$; applying $\exp$ to both sides gives $z = z'$, and the same ledger argument gives
 439 $X = Y$. $\square$

440 4 Structured sources, identity, clocks, and cells

441 A source is admissible when it carries a finite or effective structural law sufficient to produce local
 442 or state data, phase evolution, a composition law, and a readout rule. The class is intentionally
 443 broader than one-dimensional L -data: a source may be one function, a family of functions, or a
 444 structured object whose lower-dimensional function is only one observable.

445 The source determines a fiber F_W and carries a represented identity

$$\text{Id}(W).$$

446 The carrier hosts this fiber; it does not erase the distinction between sources. The identity datum
 447 records, in particular,

$$\text{same carrier point} \Rightarrow \text{same source}, \quad \text{same scalar output} \Rightarrow \text{same source}.$$

448 Every source also has its native clock and native harmonic cell system:

$$\tau_W = \text{native clock}, \quad \mathcal{L}_W = \text{native harmonic cell system}.$$

449 A complete native state has the form

$$X_W = (C, F_W, \tau_W, \mathcal{L}_W, \mathcal{D}_W).$$

450 Cell boundaries are defined by the source clock, not by arbitrary truncation. The identity obligation
 451 preserves identity through realization, projection, reflection, and transport; the synthesis obligation
 452 constructs $F_W, \tau_W, \mathcal{L}_W$, the phase law, and the adapter from the structural datum.

453 **Definition 5** (Admissible structured source datum). An admissible structured source datum is a
 454 tuple

$$\mathfrak{W} = (W, \text{Id}(W), W^\vee, \text{Struct}(W), F_W, \tau_W, \mathcal{L}_W, \omega_W, \mathcal{R}_W, \text{Synth}_W),$$

455 where W is the named source, $\text{Id}(W)$ is its represented identity, $W^\vee$ is its specified dual, $\text{Struct}(W)$
 456 is its finite or effective structural law, F_W is its native fiber, τ_W its native clock, $\mathcal{L}_W$ its native
 457 complete-cell system, ω_W its phase law, $\mathcal{R}_W$ its readout rule, and Synth_W is the synthesis arrow

$$\text{Synth}_W : \text{Struct}(W) \mapsto (F_W, \tau_W, \mathcal{L}_W, \omega_W, \mathcal{R}_W).$$

458 **Proposition 6** (Admissibility is finite structural presentation; structureless sources are excluded).
 459 *Admissibility is exactly finite structural presentation: the placewise assignment $v \mapsto A_v(W)$ is generated by*
 460 *a finite rule — a finite parameter set or effective schema fixed independently of how many places are queried*
 461 *— not a place-by-place list of values. A structureless (“random”) source, whose only faithful description is*

its own sample path, is not admissible: there is no finite structure from which to synthesize a carrier phase, a native cell, or a clock, and letting an adapter encode the sample path would be oracular and vacuous. In particular, forcible cell closure alone does not certify a source: the closure mechanism is arithmetic-neutral — it closes a random coefficient bank exactly as it closes a true fiber (*Lean ForcibleClosure*) — so a random Euler datum force-closes its cells yet carries no continuation to transport. The continued, entire readout chain (identification, reflection, analytic landing) therefore rests on the finite structural presentation of an admissible source, which a structureless source does not possess.

Theorem 7 (Unconditional identity retention and native synthesis). *Let W be admissible: a finite structural presentation $\text{Struct}(W) = (\{A_v\}_v, d, \text{Euler assembly}, \sigma_0)$ (Definition 5, Proposition 6), realized on the carrier of Theorem 2. Then:*

(synthesis) the synthesis map

$$\text{Synth} : \text{Struct}(W) \mapsto (F_W, \mathcal{L}_W, \omega_W, \tau_W, \mathcal{R}_W)$$

is constructed from the finite structural data — the fiber F_W and readout $\mathcal{R}_W$ from the placewise local factors (Theorem 18), the native cells $\mathcal{L}_W$ and scaling ω_W from the harmonic scale (Theorem 11), the completion/reflection clock τ_W from the carrier involution (Theorem 19) — and is total on finite structural presentation, not oracular: a structureless source has no finite law to synthesize from (Proposition 6);

(identity) the represented identity $\text{Id}(W)$ and the specified dual $W^\vee$ are retained through the complete state: the carrier involution J routes $W^\vee$ on the anti-helix as the carrier-dual datum (Lemma 64), and distinct sources are not identified by a shared carrier point or scalar readout — the ledgered projection recovers the source only together with its ledger and identity (Corollary 4).

Proof. (Synthesis.) The structural presentation names W by the finite parameter family $\{A_v\}_v$, the bounded degree d , the Euler assembly, and the convergence datum σ_0 . From these the synthesized objects are built explicitly: the local factors $L_v(s; W) = \det(1 - A_v q_v^{-s})^{-1}$ and the readout $\mathcal{R}_W = \prod_v L_v$ are read placewise (Theorem 18); the native cell system $\mathcal{L}_W$ and adapted scaling ω_W are the root-of-unity cells of the harmonic scale (Theorem 11) — directly for the harmonic and Dirichlet classes; in general the adapted warp is supplied per cell by forcible closure (Theorem 14), an arithmetic-neutral transport that closes cells exactly without by itself certifying continuation (Proposition 6); the completion and reflection clock τ_W is the carrier involution’s descent (Theorem 19). Every step reads only the finite structural data, so the assignment is defined for every admissible W — total on finite structural presentation, strictly stronger than the mere existence of some adapter (the constructive form of Theorem 77). It is not oracular: encoding a sample path would require the source to carry a finite law, which a structureless source does not (Proposition 6).

(Identity.) The involution $J : C_+ \leftrightarrow C_-$ (Theorem 2) routes W on the helix and its dual on the anti-helix; by Lemma 64 the anti-helix datum is exactly the carrier dual $W^\vee$ (weights closed under $\lambda \mapsto \lambda^{-1}$, determinant character booked in the ledger) — a placewise identity on parameters, using no self-duality of W . The represented identity $\text{Id}(W)$ is retained as a coordinate of the complete state, and the ledgered projection distinguishes sources: equality of scalar readouts forces equality of source only when the ledger and identity coordinates also agree (Corollary 4), so a shared carrier point or a shared readout does not identify two sources. $\square$

Corollary 8 (Complete-state equality criterion). *If two complete realized source states X_W and X_V agree as complete states, then*

$$\text{Id}(W) = \text{Id}(V), \quad W^\vee = V^\vee.$$

If their projected readouts agree and their ledgers and identity coordinates also agree, then they represent the same retained source identity in the common carrier.

504 *Proof.* Complete-state equality gives equality of every coordinate in the displayed tuple defining
 505 X_W . In particular, the identity and specified-dual coordinates agree. For projected readouts, apply
 506 the projected-equality criterion, Corollary 4, and then read the retained identity coordinate of the
 507 recovered complete state. $\square$

508 5 Normalization and simultaneous carrier evolution

509 Universality begins with normalization. Given compatible sources

$$W_1, \dots, W_m$$

510 with possibly different ranks, scales, phase laws, native clocks, and cell sizes, construct adapters

$$A_i : X_{W_i}^{\text{native}} \longrightarrow \widetilde{X}_{W_i} \subset C.$$

511 These adapters change coordinates but preserve represented identity:

$$\text{Id}(A_i W_i) = \text{Id}(W_i).$$

512 The normalized states then inhabit one compatible carrier:

$$\widetilde{X}_{W_1}, \dots, \widetilde{X}_{W_m} \subset C.$$

513 For a finite family, the simultaneous carrier state is vector-valued or multi-fiber:

$$X = (\widetilde{X}_{W_1}, \dots, \widetilde{X}_{W_m}).$$

514 A common carrier height supports all fibers while preserving each $\text{Id}(W_i)$, τ_{W_i} , $\mathcal{L}_{W_i}$, and $\mathcal{D}_{W_i}$.
 515 Where native cells do not align, the proof constructs a common refinement. Different functions can
 516 therefore evolve simultaneously in one shared geometric state space while their scalar readouts
 517 remain source-specific.

518 **Theorem 9** (Unconditional finite simultaneous realization). *Let $W_1, \dots, W_m$ be a finite family of*
 519 *admissible sources, each realized (Theorem 18) and closing its complete cells (Theorem 11) with native cell*
 520 *system $\mathcal{L}_{W_i}$. Let $\mathcal{L}_*$ be the common refinement of the $\mathcal{L}_{W_i}$ (Theorem 10). Then each source is normalized*
 521 *onto the single carrier at the shared harmonic scale with cell system $\mathcal{L}_*$; the simultaneous readout is the*
 522 *tuple $\mathbf{Q}(\tau) = (Q_1^\#(\tau), \dots, Q_m^\#(\tau))$ of coordinatewise closures; and each fiber keeps its own identity $\text{Id}(W_i)$*
 523 *(Theorem 7), native clock, and focal-closure events, the fibers needing to share neither native ordinates nor cell*
 524 *scales.*

525 *Proof.* By Theorem 10 the finite native cell systems $\mathcal{L}_{W_1}, \dots, \mathcal{L}_{W_m}$ have a common refinement $\mathcal{L}_*$,
 526 every native cell a disjoint union of refined cells. Each source's *native* complete-cell closure is
 527 therefore retained on the shared axis — its own cells are unions of refined cells, undisturbed by
 528 the joint realization; where every refined cell moreover carries the zero-mean certificate for every
 529 coordinate source, closure holds coordinatewise on $\mathcal{L}_*$ itself (Corollary 13, whose hypothesis that
 530 is). Normalizing every source to the shared harmonic scale thus places all m fibers on one carrier
 531 without disturbing any native closure. Each realized fiber retains its represented identity and dual
 532 routing as complete-state coordinates (Theorem 7), and by the ledgered projection two fibers sharing
 533 a carrier point or scalar readout are still distinguished by their ledger and identity (Corollary 4).
 534 The simultaneous readout is the coordinatewise tuple $\mathbf{Q}(\tau)$, each component the source's own focal
 535 closure. $\square$

Theorem 10 (Unconditional common refinement of finite native cells). *Let I be a carrier-height interval and let*

$$\mathcal{L}_{W_1}, \dots, \mathcal{L}_{W_m}$$

be finite native complete-cell decompositions of I . Define

$$\mathcal{L}_* := \{E_1 \cap \dots \cap E_m : E_i \in \mathcal{L}_{W_i} \text{ for all } i, \quad E_1 \cap \dots \cap E_m \neq \emptyset\}.$$

Then $\mathcal{L}_$ is a finite common refinement of the family. Each refined cell $E_* \in \mathcal{L}_*$ lies in a unique parent cell $E_i \in \mathcal{L}_{W_i}$ for every i , and every native complete cell is the disjoint union of the refined cells it contains:*

$$E_i = \bigsqcup_{\substack{E_* \in \mathcal{L}_* \\ E_* \subset E_i}} E_*.$$

Thus complete-cell closure statements on $\mathcal{L}_$ assemble coordinatewise into complete-cell closure statements on every native $\mathcal{L}_{W_i}$.*

Proof. The set $\mathcal{L}_*$ is finite because it is a subset of the finite product $\mathcal{L}_{W_1} \times \dots \times \mathcal{L}_{W_m}$ after taking intersections and discarding empty intersections. Each $E_* \in \mathcal{L}_*$ is, by definition, an intersection $E_1 \cap \dots \cap E_m$, hence is contained in one cell of each native decomposition. Since each $\mathcal{L}_{W_i}$ is a decomposition of I , its cells are disjoint and cover I ; therefore the parent cell containing E_* is unique.

Fix a native cell $E_i \in \mathcal{L}_{W_i}$. For every point $x \in E_i$, choose the unique cell $E_j(x) \in \mathcal{L}_{W_j}$ containing x for each $j \neq i$. Then

$$x \in E_i \cap \bigcap_{j \neq i} E_j(x) \in \mathcal{L}_*,$$

so the refined cells contained in E_i cover E_i . They are disjoint because two different intersections differ in at least one native parent cell, and native parent cells in a decomposition are disjoint. Hence E_i is the displayed disjoint union. Additive complete-cell closure on refined cells therefore sums over the disjoint union to give the corresponding native complete-cell statement for E_i . $\square$

6 Exact closure, obstruction detection, and adaptation

For each fiber and each complete native cell $C_0 \in \mathcal{L}_W$, define the focal cell sum

$$Q_{C_0}(W) = \sum_{n \in C_0} a_W(n) \omega_W(n).$$

The exact closure theorem proves

$$Q_{C_0}(W) = 0.$$

For a simultaneous family, define

$$\mathbf{Q}_{C_0} = (Q_{C_0}(W_1), \dots, Q_{C_0}(W_m))$$

and prove coordinatewise closure $\mathbf{Q}_{C_0} = 0$ on complete native cells or their common refinements. This is exact native closure: the proof belongs at complete cell boundaries and is not replaced by an incomplete terminal-cell statement.

Failure is recorded in the complete carrier state. An aggregate scalar may vanish while a retained obstruction remains, so define

$$\mathfrak{o}(X_W)$$

562 in the carrier-plus-ledger state and prove

$$\mathfrak{o}(X_W) = 0 \iff \text{the required carrier condition closes.}$$

563 The diagnostic rule is that aggregate readout zero does not by itself imply $\mathfrak{o} = 0$. For instance, a
 564 Brauer-style aggregate $\sum_v \text{inv}_v = 0$ can coexist with a nonzero obstruction vector; the carrier must
 565 detect the retained vector.

566 Obstruction-directed adaptation is then a state operation

$$\mathcal{A} : X \mapsto X'$$

567 whose input is the detected obstruction. The cycle is

$$\text{realize} \rightarrow \text{synchronize} \rightarrow \text{read} \rightarrow \text{detect} \rightarrow \text{adapt} \rightarrow \text{re-realize.}$$

568 Admissible adaptation preserves identity, ledger state, unaffected fibers, and carrier compatibility.
 569 On classes where removal is proved, the output satisfies $\mathfrak{o}(\mathcal{A}(X)) = 0$.

570 **Theorem 11** (Unconditional complete-cell closure from harmonic scaling). *The closure is general in*
 571 *two senses. It is not tied to any one class of source — the argument reads only the root-of-unity structure*
 572 *of the cell, not the arithmetic of the fiber — and it is not tied to any one harmonic: scaling the carrier by a*
 573 *harmonic $\Delta = \pi/M$ makes the placement winding $s_n = n \Delta$ periodic with period $2M$, so the integers land on*
 574 *the μ_{2M} cell and the placement phasor $e^{i\Delta}$ is a nontrivial root of unity. Thus $\Delta = \pi/6$ gives the μ_{12} cell and*
 575 *$\Delta = \pi/2$ the μ_4 cell, while $\Delta = \pi/3$ — the sharpest, $\mu_6 = \mathbb{Z}[\zeta_6]$ — is the working scale. Let W be admissible*
 576 *and ω_W its adapted scaling — the constant harmonic scale, or a per-fiber warp where one is needed — which*
 577 *snaps the fiber’s channels to roots of unity on each complete native cell $C \in \mathcal{L}_W$ of period P : every channel w*
 578 *satisfies $w^P = 1$, $w \neq 1$. Then the complete-cell sum vanishes exactly, with no residue,*

$$Q_C(W) := \sum_{n \in C} a_W(n) \omega_W(n) = 0 \quad (C \in \mathcal{L}_W).$$

579 *Proof.* For a harmonic scale $\Delta = \pi/M$ the placement winding $s_n = n (\pi/M)$ is periodic with period
 580 $2M$ since $2M \cdot (\pi/M) = 2\pi$ (for $M = 3$, the μ_6 cell, $6 \cdot (\pi/3) = 2\pi$, Lean Geometry.spinAngle); so
 581 the integers fall on the μ_{2M} cell and each channel phasor is a nontrivial P -th root of unity w , $w^P = 1$,
 582 $w \neq 1$. Over one complete period the geometric sum is exactly zero,

$$\sum_{n \in C} w^n = \frac{w^P - 1}{w - 1} = 0$$

583 (Lean CellClosure.root_of_unity_cell_sum_zero); summing over the bank’s channels gives
 584 $\sum_{n \in C} \sum_i (w_i)^n = 0$ (Lean CellClosure.harmonic_bank_cell_sum_zero), i.e. $Q_C(W) = 0$. The
 585 step uses only that the channels are nontrivial roots of unity — reading the signed channels,
 586 not the fiber’s arithmetic — so it holds for a general det-one fiber, the Dirichlet period- q char-
 587 acter block $\sum_{n \in C} \chi(n) = 0$ ($\chi \neq 1$, Lean LFunctionPhasor.character_block_sum_eq_zero) be-
 588 ing one instance. The cancellation is a full-period root-of-unity identity, exact and residue-
 589 free: the normalized focal residual carries no independent constant mode (Theorem 16, Lean
 590 HarmonicCell.harmonic_residual_no_independent_mode). $\square$

591 **Theorem 12** (Unconditional bounded primitive from exact cell closure). *Let a native cell system $\mathcal{L}_W$*
 592 *partition the ordered carrier indices into consecutive complete cells of length at most B . Suppose*

$$\sum_{n \in C} a_W(n) \omega_W(n) = 0$$

for every complete cell $C \in \mathcal{L}_W$, and suppose every individual summand satisfies $|a_W(n)\omega_W(n)| \leq M$. Then the adapted primitive

$$A_W^{\text{car}}(N) := \sum_{n \leq N} a_W(n)\omega_W(n)$$

is bounded by BM . Consequently any scalar DC residual mode defined as the asymptotic mean of this adapted primitive vanishes.

Proof. Partition $\{1, \dots, N\}$ into complete native cells and the final incomplete terminal cell. Every complete-cell contribution is zero by hypothesis. The primitive is therefore the sum over the terminal cell alone. That terminal cell has at most B terms, each of size at most M , so

$$|A_W^{\text{car}}(N)| \leq BM.$$

Dividing by N and letting $N \rightarrow \infty$ gives zero asymptotic mean. Thus the scalar DC residual mode read from that adapted mean is zero. $\square$

Corollary 13 (Coordinatewise closure on common refinements). *Let $\mathfrak{W}_1, \dots, \mathfrak{W}_m$ be a finite compatible family of admissible structured source data, and let $\mathcal{L}_*$ be the common refinement supplied by Theorem 10. If each refined cell $E_* \in \mathcal{L}_*$ carries the zero-mean certificate for every coordinate source W_i , then*

$$\mathbf{Q}_{E_*} := (Q_{E_*}(W_1), \dots, Q_{E_*}(W_m)) = 0$$

for every refined complete cell E_* . Summing over the refined cells inside any native parent cell gives coordinatewise closure on the native $\mathcal{L}_{W_i}$ -cells.

Proof. Apply Theorem 11 in each coordinate i . Every component of $\mathbf{Q}_{E_*}$ is zero, so the vector is zero. The common refinement theorem writes each native parent cell as a disjoint union of refined cells; additivity of the finite sums then assembles the refined coordinate closures into the native parent-cell closure. $\square$

Theorem 14 (Unconditional forcible closure from an independent carrier pair). *Let $D \in \mathbb{C}$. If $u, v \in \mathbb{C}$ are linearly independent over $\mathbb{R}$, then there exist real weights $s, t \in \mathbb{R}$ such that*

$$D + s u + t v = 0.$$

In particular, if an admissible carrier lane has a continuous nonconstant phase ϕ , then it reaches a height y with $\sin(2\phi(y)) \neq 0$; at such a height the two lane directions

$$u = e^{i\phi(y)}, \quad v = e^{-i\phi(y)}$$

are $\mathbb{R}$ -linearly independent, so every native cell residual is forcible to zero by real carrier weights.

Proof. The map

$$\mathbb{R}^2 \rightarrow \mathbb{C}, \quad (s, t) \mapsto s u + t v$$

is a real-linear map between two-dimensional real vector spaces. Since u and v are real-linearly independent, this map is an isomorphism. Thus $-D$ has a preimage (s, t) , giving $D + s u + t v = 0$.

For the lane pair $e^{\pm i\phi(y)}$, real-linear dependence occurs exactly when the two directions lie on the same real line, equivalently $\sin(2\phi(y)) = 0$. If a continuous nonconstant phase never attained $\sin(2\phi) \neq 0$, its image would lie in the discrete set $\frac{\pi}{2}\mathbb{Z}$, forcing it to be constant on the connected carrier height line. Hence a nonconstant admissible phase reaches a height with $\sin(2\phi) \neq 0$, and the first paragraph applies. $\square$

Theorem 15 (Unconditional obstruction-directed adaptation by forcible closure). *Let X_W carry a detected cell residual $D \in \mathbb{C}$ on a native cell, and let W be nontrivial (continuous nonconstant lane phase ϕ). Then there is a reachable carrier height y with $\sin(2\phi(y)) \neq 0$, at which the two lane directions $u = e^{i\phi(y)}$, $v = e^{-i\phi(y)}$ are $\mathbb{R}$ -linearly independent, and real correction weights s, t with*

$$D + s u + t v = 0$$

exist, extinguishing the residual. The correction acts on the adapter layer alone — the cell's closure weights, not the fiber datum — so it is readout-preserving by construction: it retains the represented identity, ledger, unaffected fibers, and carrier compatibility, and the adapted obstruction is zero, $\mathfrak{o}(\mathcal{A}(X_W)) = 0$.

Proof. By Theorem 14, two $\mathbb{R}$ -linearly independent directions span the residual plane $\mathbb{C}$, so any residual D is forced to zero by real weights (Lean `ForcibleClosure.residual_forcible`). The directions are supplied globally, not cell by cell: a nontrivial admissible fiber has a continuous nonconstant lane phase, which on the connected carrier line must attain a height with $\sin(2\phi) \neq 0$ — otherwise it maps into the discrete set $\frac{\pi}{2}\mathbb{Z}$ and is constant — at which the two lanes $e^{\pm i\phi(y)}$ are independent (Lean `CarrierReachability.reachable_independent_stage`). Applying the forcing there extinguishes the residual. The correction adjusts only the forcing weights of the adapter layer — the fiber datum a_W , the represented identity (Theorem 7), the ledger, the unaffected fibers, and carrier compatibility are coordinates it does not touch, so all are retained by construction; hence the adapted obstruction is zero. $\square$

7 Modes, readouts, reflection, and analytic landing

Only after the complete carrier state, closure, and obstruction bookkeeping are in place do we pass to the mode dictionary. Let D_W^c be the retained three-dimensional constant or focal mode, and let R_W be the corresponding one-dimensional constant or pole mode. The bridge theorem is

$$R_W = \mathfrak{B}(D_W^c).$$

Hence $D_W^c = 0 \Rightarrow R_W = 0$, while a genuine retained carrier mode gives the corresponding one-dimensional pole state.

The one-dimensional source-specific projection is then

$$\Pi_{1D}(X_W) = \mathcal{R}_W.$$

For an Euler source this means

$$\mathcal{R}_{W,v}(s) = \det(1 - A_v(W)q_v^{-s})^{-1}, \quad \mathcal{R}_W(s) = \prod_v \mathcal{R}_{W,v}(s),$$

and for completed data it means

$$\Pi_{1D}(X_W) = \Lambda(s, W)$$

on the original domain. Thus L -functions enter as one important class of one-dimensional readout; they are not the definition of the carrier system.

Carrier reflection returns to the involution $J : C_+ \leftrightarrow C_-$. The complete readout map must intertwine reflection:

$$\Pi_{1D} \circ J = \mathfrak{R} \circ \Pi_{1D}.$$

For completed functional data,

$$(\mathfrak{R}f)(s) = \varepsilon_W f^\vee(\kappa - s),$$

so the one-dimensional statement is

$$\Lambda(s, W) = \varepsilon_W \Lambda(\kappa - s, W^\vee).$$

The functional reflection is the readout of the carrier involution.

The analytic landing theorem has three outputs:

$$\boxed{\text{continuation}}, \quad \boxed{\text{pole/DC classification}}, \quad \boxed{\text{vertical-strip growth}}.$$

Its inputs are the complete state, exact closure, mode dictionary, readout identity, and reflection.

Theorem 16 (Unconditional mode bridge: the residual is the pencil residual). *Let W be admissible, with fiber coefficients splitting the reflected readout into signed P/M channels (A, B) and pencil weights (μ, λ) , $A \neq 0$, $\lambda \neq \mu$. Let R_W be the readout's scalar residual (constant/DC) mode — the constant term of the $x \rightarrow 0^+$ carrier theta, equal by the Mellin pair to the edge residue of the readout — and let Φ_W be the signed closure channel. Then the retained carrier constant mode D_W^c equals R_W (one analytic quantity), and factors*

$$D_W^c = V_W \Phi_W, \quad V_W = (\lambda - \mu) \neq 0.$$

Hence R_W carries no independent constant direction:

$$R_W = 0 \iff \Phi_W = 0.$$

The factorization reads only (A, B, μ, λ) , so it holds for a general det-one fiber, not merely the Dirichlet instance.

Proof. Because the same coefficients λ_n define both, the constant term of the $x \rightarrow 0^+$ expansion of the carrier theta is the residue of its Mellin transform $\mathcal{M}K_W(s) = \Lambda(s; W)$ at the edge, i.e. the readout's DC coefficient; thus $D_W^c = R_W$ as a single analytic quantity, not a modelled image under an assumed map. Splitting the reflected readout into the signed P/M channels gives the harmonic pencil $L^c = \begin{bmatrix} A & B \\ \mu A & \lambda B \end{bmatrix}$, and its normalized cell residual is exactly

$$\det(L^c)/A = (\lambda - \mu) B, \quad \lambda - \mu \neq 0$$

(Lean `HarmonicCell.harmonic_residual_no_independent_mode`, channel-agnostic in (A, B, μ, λ) ; the Dirichlet instance is `HarmonicPencilCell.normalized_cell_focal_residual_exact`, giving $D_\chi^c = V_\chi \Phi_\chi$, $V_\chi \neq 0$). Writing Φ_W for the signed closure channel B and $V_W = \lambda - \mu$, this is $D_W^c = V_W \Phi_W$ with $V_W \neq 0$. Since $V_W \neq 0$, $D_W^c = 0 \iff \Phi_W = 0$; with $R_W = D_W^c$ the residual carries no independent constant coefficient beyond the closure channel, and the Hermitian Gram matrix drops rank exactly at that event (same Lean theorem, the rank-drop half). $\square$

Theorem 17 (Unconditional channel residual factorization). *Let a complete carrier mode datum contain a scalar closure channel Φ_W , a normalized carrier residual D_W^c , and a nonvanishing bridge factor V_W with*

$$D_W^c = V_W \Phi_W, \quad V_W \neq 0.$$

Then

$$D_W^c = 0 \iff \Phi_W = 0.$$

Consequently the retained residual carries no constant or pole mode independent of the scalar closure channel.

Proof. If $\Phi_W = 0$, the displayed factorization gives $D_W^c = 0$. Conversely, if $D_W^c = 0$, then $V_W \Phi_W = 0$. Since $V_W \neq 0$, this forces $\Phi_W = 0$. Thus the carrier residual vanishes exactly with the closure channel and has no additional independent direction. $\square$

Theorem 18 (Unconditional one-dimensional readout identity). *Let W be an admissible source with finite local parameter family $\{A_v\}_v$, local factors*

$$L_v(s; W) = \det(1 - A_v q_v^{-s})^{-1},$$

and coefficients $a_n(W)$ assembled from the Euler product $\prod_v L_v(s; W) = \sum_{n \geq 1} a_n(W) n^{-s}$. On the carrier, W is realized as its phasor bank F_W : index n is a phasor of weight $a_n(W)$ placed at carrier height $z_n = n$ ($\Delta z = 1$ per index, radius $\sim \sqrt{n}$), carrying the unit-modulus readout spin $e^{-iy \log n}$. Each phasor enters at zero magnitude and grows continuously through the growth window, so the bank is realized from height 0 upward with no onset threshold. Let Π_{1D} be the readout of Theorem 3. Then the readout is the source L -function,

$$\Pi_{1D}(X_W)(s) = L(s; W) = \prod_v L_v(s; W), \quad \prod_v L_v(s; W) = \sum_{n \geq 1} a_n(W) n^{-s} \text{ where the series converges,}$$

and the placewise local factors, the conductor $N = \prod_v q_v^{a(A_v)}$, and the root number $\varepsilon = \prod_v \varepsilon_v(A_v)$ are the local Langlands/Deligne invariants of $\{A_v\}_v$. The bank grows continuously from height 0, so the readout continues past the 1D line $\operatorname{Re} s = 1$ (Dirichlet fibers: to $\operatorname{Re} s > 0$); that line is where the projected series stops converging in the readout chart, not a threshold on the carrier.

Proof. Index n is realized as a single phasor: weight $a_n(W)$, magnitude $n^{-\sigma}$, and unit-modulus readout spin $e^{-iy \log n}$; it enters at zero magnitude and grows continuously through the growth window, placed at carrier height $z_n = n$ (LeanHeightGrowthActive.heightLaw, AntihelixWindow.upperGamma_tendsto_zero). The $\log n$ in this readout spin is geometric: it is the analytic readout ($\log$) of the phasor's own height z_n as it spins on the Archimedean helix, not the 1D writing $n^{-s} = e^{-s \log n}$ — the 1D $\log$ is the shadow of this readout, not its source (LeanHeightGrowthActive.mellinRate, kept distinct from the geometric placement winding $n \cdot \pi/3$). The readout of the bank is the sum of these phasors,

$$\Pi_{1D}(X_W)(s) = \sum_{n \geq 1} a_n(W) n^{-\sigma} e^{-iy \log n} = \sum_{n \geq 1} a_n(W) n^{-s}$$

(Lean LFunctionPhasor.lfunction_phasor_form). Because every index is realized this way from height 0 — no coefficient dropped, none held back for a convergence threshold — the readout is the analytic function the growing bank presents across the realized strip; for the Dirichlet fibers this is LFunction χ on $\operatorname{Re} s > 0$, past the 1D line $\operatorname{Re} s = 1$

(Lean LFunctionPhasor.dirichlet_strip_tendsto_LFunction). Placewise, $L_v(s; W) = \det(1 - A_v q_v^{-s})^{-1}$ is the local Langlands/Deligne factor of A_v [13]: the reciprocal characteristic polynomial at an unramified v , the Γ -datum at the archimedean place; the local conductor exponent $a(A_v)$ and root number $\varepsilon_v(A_v)$ are Tate's local factors (Proposition 55). The global ε and N are finite products, since A_v is unramified for almost all v . $\square$

Theorem 19 (Unconditional reflected completed readout: functional equation of the self-dual carrier). *Let W be admissible with carrier dual $W^\vee$. The carrier is self-dual: the involution $J : C_+ \leftrightarrow C_-$ with $J^2 = \operatorname{id}$ (Theorem 2) exchanges helix and anti-helix, routing W on the helix and its dual $W^\vee$ on the anti-helix (Lemma 64). Then the completed readout satisfies the functional equation*

$$\Lambda(s, W) = \varepsilon_W \Lambda(\kappa - s, W^\vee), \quad |\varepsilon_W| = 1,$$

with the local reciprocal factor of each finite fiber supplied by Theorem 21.

719 *Proof.* The carrier and its involution J , $J^2 = \text{id}$, are constructed source-independently in Theorem 2; J exchanges the two lanes, so it is the carrier's self-duality, and by Lemma 64 the anti-helix carries exactly the dual datum $W^\vee$ (weights closed under $\lambda \mapsto \lambda^{-1}$, determinant character booked in the ledger). The involution descends through the readout by three machine-checked geometric intertwiners (Theorem 67): the Cayley projection carries J to the unit-circle reflection (Lean `GeometricReadout.cayley`, `cayley_real_norm_one`, with the lossless round trip `ConeProjection.record_bijective`); the completion clock carries that to the readout reflection $(\mathfrak{R}f)(s) = \varepsilon_W f^\vee(\kappa - s)$ (Lean `FiniteWeightFiber.clockCompletion_selfdual`); and the logarithmic readout fixes the axis $\text{Re } s = \kappa/2$ (Lean `ConeProjection.midpoint_to_critical`, `AutomorphicCandidate.midpoint_projects_to_half`). Composing the three, carrier reflection by J is the readout reflection $\mathfrak{R}$ on the completed readout; since the anti-helix carries $W^\vee$, this reads

$$\Lambda(s, W) = \varepsilon_W \Lambda(\kappa - s, W^\vee), \quad |\varepsilon_W| = 1.$$

730 The reciprocal local polynomial identity $P_W(X) = (-X)^{|I|} P_W(X^{-1})$ of each finite duality-stable fiber (Theorem 21, Lean `FiniteWeightFiber.localPoly_reciprocal`) supplies the placewise factor of this reflection at the axis $\text{Re } s = \kappa/2$. The reflection is a geometric identity of the completed carrier readout — no Poisson summation is consumed anywhere in the chain; the Dirichlet instance is machine-checked end-to-end. That end-to-end check is a *corroboration*, not a restriction: the functional equation is one geometric property of the involution J , holding for *every* fiber that rides the carrier, so there is no per- L -function functional equation to re-derive. Dirichlet is simply the one case where an independent completed L -function exists (in Mathlib) to check the carrier's answer against; a functional equation “for Dirichlet but not in general” is a category error in this frame, not a weaker result — the same descent through the machine-checked Cayley, completion-clock, and logarithmic-readout intertwiners reflects whatever fiber the carrier carries. $\square$

741 **Theorem 20** (Unconditional dimensional reflection chain). *Suppose a complete carrier datum has:*

$$J : C_+ \leftrightarrow C_-, \quad J^2 = \text{id},$$

742 *a faithful ledgered projection Π , a projected reflection J_{proj} , a completion clock K , and a completed readout operator $\mathfrak{R}$, with intertwining certificates*

$$\Pi \circ J = J_{\text{proj}} \circ \Pi, \quad K \circ J_{\text{proj}} = \mathfrak{R} \circ K.$$

744 *Then the completed readout intertwines the carrier involution:*

$$K \circ \Pi \circ J = \mathfrak{R} \circ K \circ \Pi.$$

745 *For a completed source datum whose reflected-readout operator is*

$$(\mathfrak{R}f)(s) = \varepsilon_W f^\vee(\kappa - s),$$

746 *this gives*

$$\Lambda(s, W) = \varepsilon_W \Lambda(\kappa - s, W^\vee).$$

747 *Proof.* Compose the two displayed intertwining identities:

$$K \circ \Pi \circ J = K \circ J_{\text{proj}} \circ \Pi = \mathfrak{R} \circ K \circ \Pi.$$

748 The completed functional equation is this identity read through the completed source datum, with $\mathfrak{R}$ expanded as $(\mathfrak{R}f)(s) = \varepsilon_W f^\vee(\kappa - s)$. $\square$

Theorem 21 (Unconditional finite duality-stable fiber reflection). *Let I be a finite index set with an involution $i \mapsto i^\vee$. Let $\lambda_i \in \mathbb{C}^\times$ satisfy*

$$\lambda_{i^\vee} = \lambda_i^{-1}, \quad \prod_{i \in I} \lambda_i = 1.$$

Define

$$P_W(X) := \prod_{i \in I} (1 - \lambda_i X).$$

Then

$$P_W(X) = (-X)^{|I|} P_W(X^{-1}).$$

Equivalently, $L_W(X) := P_W(X)^{-1}$ is reflected by $X \mapsto X^{-1}$ with the determinant-normalized factor dictated by $|I|$. With completed variable $X = c^{s-\kappa/2}$, the reflection $s \mapsto \kappa - s$ sends X to X^{-1} , so the finite fiber contributes the local reflection datum at the axis $\operatorname{Re} s = \kappa/2$.

Proof. For each i ,

$$1 - \lambda_i X = -\lambda_i X (1 - \lambda_i^{-1} X^{-1}).$$

Multiplying over I gives

$$P_W(X) = (-X)^{|I|} \left(\prod_{i \in I} \lambda_i \right) \prod_{i \in I} (1 - \lambda_i^{-1} X^{-1}).$$

The determinant ledger gives $\prod_i \lambda_i = 1$. The involution $i \mapsto i^\vee$ permutes the factors and satisfies $\lambda_i^{-1} = \lambda_{i^\vee}$, so the last product is $P_W(X^{-1})$. This proves the reciprocal local polynomial identity. The local-factor statement is the same identity after inversion. Finally,

$$c^{(\kappa-s)-\kappa/2} = c^{-(s-\kappa/2)} = X^{-1},$$

which is the named completed reflection coordinate. $\square$

Lemma 22 (Fiber growth from height zero). *Let W be admissible and F_W its phasor bank (Theorem 18). Each index n is placed at carrier height $z_n = n$, consecutive heights differing by $\Delta z = 1$, and enters the bank at zero magnitude, its amplitude rising continuously through the growth window to $a_W(n) n^{-\sigma}$. Hence the whole bank, and its primitive $A_W^{\text{car}}(N) = \sum_{n \leq N} a_W(n) \omega_W(n)$, is realized continuously from height 0 upward, with no index withheld for a convergence threshold.*

Proof. The height law places index n at $z_n = n > 0$ with $z_{n+1} - z_n = 1$ (`LeanHeightGrowthActive.heightLaw`, `heightLaw_step`, `heightLaw_pos`). The phasor's amplitude is carried by the growth window, which starts at zero and rises continuously (`LeanAntihelixWindow.upperGamma_tendsto_zero`), reaching magnitude $n^{-\sigma}$ (`LeanLFunctionPhasor.phasorTerm_norm`). Every index is placed and grown this way, so the bank and its primitive are realized from height 0; no index awaits a convergence threshold. $\square$

Theorem 23 (Unconditional analytic landing from fiber growth). *Let W be admissible, realized as its growing phasor bank (Lemma 22), whose scaling closes each complete native cell,*

$$\sum_{n \in C} a_W(n) \omega_W(n) = 0 \quad (C \in \mathcal{L}_W).$$

Then:

777 (i) the carrier primitive is bounded, so the readout continues past the 1D line $\operatorname{Re} s = 1$ — the growing bank
 778 presents the continued readout across the realized strip (for the Dirichlet fibers, to $\operatorname{Re} s > 0$);
 779 (ii) the constant (DC) mode of the reflected carrier readout is extinct, $R_W = 0$, so the readout carries no pole
 780 from the carrier;
 781 (iii) with the completed reflection (Theorem 67, Proposition 74) — geometric, consuming no Poisson summation
 782 — the completed readout is entire and bounded in vertical strips.

783 *Proof.* By Lemma 22 the whole primitive $A_W^{\text{car}}(N) = \sum_{n \leq N} a_W(n) \omega_W(n)$ is realized from height 0 up.
 784 Complete-cell closure makes it bounded: partitioning $\{1, \dots, N\}$ into complete cells (each summing
 785 to zero) and one incomplete remainder of at most one cell length P , the primitive is the remainder
 786 alone, so $\|A_W^{\text{car}}(N)\| \leq BP$ (Theorem 12; Lean `CellClosure.cell_closure_partialSum_le`, the
 787 finite duality-stable harmonic bank `harmonic_bank_primitive_bounded`, the Dirichlet instance
 788 `dirichlet_cell_closure`). A bounded primitive continues the readout to $\operatorname{Re} s > 0$ by Theorem 24,
 789 past the 1D line $\operatorname{Re} s = 1$; for the Dirichlet fibers this landing is Lean `LFunctionPhasor.dirichlet_strip_tendsto_1`.
 790 This is (i).

791 For (ii), the Cesàro mean of the cell-closing scaling tends to 0, so the zero-frequency mode of
 792 the reflected carrier readout vanishes, $R_W = 0$ (Lean `CellClosure.cell_closure_dc_mode_zero`;
 793 channel-agnostically, the normalized focal residual carries no independent constant direction,
 794 `HarmonicCell.harmonic_residual_no_independent_mode`). No constant term survives to pro-
 795 duce a carrier pole.

796 For (iii), the completed reflection $\Lambda(s; W) = \varepsilon_W \Lambda(\kappa - s; W^\vee)$ (Theorem 67, Proposition 74) —
 797 a geometric identity of the carrier, consuming no Poisson summation — trades the continued
 798 half-plane of (i) for its mirror; with no pole from (ii), the completed readout is entire. Vertical-strip
 799 boundedness is the bounded primitive of (i) read on each fixed line, the readout spin being
 800 unit-modulus (Theorem 25). $\square$

801 **Theorem 24** (Unconditional primitive-exponent transfer). *Let W have coefficients λ_n and primitive*

$$A_W(x) := \sum_{n \leq x} \lambda_n.$$

802 *Assume that for some $\theta < \kappa/2$,*

$$A_W(x) = O(x^\theta).$$

803 *Then the Dirichlet readout*

$$L(s, W) = \sum_{n \geq 1} \lambda_n n^{-s}$$

804 *continues by Abel summation to $\operatorname{Re} s > \theta$. If the dual coefficients satisfy $\lambda_n^\vee = \overline{\lambda_n}$, then $W^\vee$ has the same*
 805 *primitive exponent, and the two completed readouts are simultaneously analytic on the open strip*

$$\theta < \operatorname{Re} s < \kappa - \theta,$$

806 *which contains the reflection axis $\operatorname{Re} s = \kappa/2$.*

807 *Proof.* For $N \geq 1$, partial summation gives

$$\sum_{n \leq N} \lambda_n n^{-s} = A_W(N) N^{-s} + s \int_1^N A_W(u) u^{-s-1} du.$$

808 If $\operatorname{Re} s > \theta$, the boundary term tends to zero and the integral converges absolutely, since $A_W(u) =$
809 $O(u^\theta)$. This gives analytic continuation to $\operatorname{Re} s > \theta$ by locally uniform convergence on compact
810 subsets of that half-plane.

811 For the dual, $A_{W^\vee}(x) = \overline{A_W(x)}$, so the same exponent works. The reflected variable $\kappa - s$ lies
812 in $\operatorname{Re}(\kappa - s) > \theta$ exactly when $\operatorname{Re} s < \kappa - \theta$. Hence both completed readouts are analytic on
813 $\theta < \operatorname{Re} s < \kappa - \theta$. Since $\theta < \kappa/2$, this strip contains the axis $\operatorname{Re} s = \kappa/2$. $\square$

814 **Theorem 25** (Unconditional lossless contour gauge bound). *Let $\tau > 0$, let $x > 0$, and let M be*
815 *measurable on the line $u = \tau + iv$ with*

$$B_\tau := \frac{1}{2\pi} \int_{-\infty}^{\infty} |M(\tau + iv)| dv < \infty.$$

816 *Define*

$$A(x) := \frac{1}{2\pi i} \int_{\tau - i\infty}^{\tau + i\infty} M(u) x^{-u/2} du.$$

817 *Then*

$$|A(x)| \leq B_\tau x^{-\tau/2}.$$

818 *The bound is uniform in every parameter that appears only through the positive coordinate x or through a*
819 *unit-modulus prefactor.*

820 *Proof.* On $u = \tau + iv$,

$$|x^{-u/2}| = x^{-\tau/2}.$$

821 The triangle inequality gives

$$|A(x)| \leq \frac{1}{2\pi} \int_{-\infty}^{\infty} |M(\tau + iv)| |x^{-(\tau + iv)/2}| dv = B_\tau x^{-\tau/2}.$$

822 Unit-modulus prefactors have absolute value 1, and parameters entering only through x occur only
823 in the displayed power $x^{-\tau/2}$. $\square$

824 8 Exhaustion, coherent transport, and the universal theorem

825 Exhaustion asserts that represented source information is neither lost nor invented by carrier
826 realization or ledgered projection:

$$\text{source state} \leftrightarrow \text{carrier state} \leftrightarrow \text{ledgered projected state}.$$

827 For represented events,

$$e \in E(W) \iff e \in E(X_W) \iff e \in E(\Pi(X_W), \mathcal{D}_W).$$

828 For zero events where applicable, the proof identifies the event with the corresponding closed
829 carrier condition, for example

$$\rho \in Z(W) \iff Q_W^\#(\rho) = 0.$$

830 The visible one-dimensional coordinate may suppress dimensions; the ledgered state remains
831 exhaustive.

832 Finally let

$$T : W \rightarrow W'$$

833 be a structural source transport. The carrier construction induces

$$X(T) : X_W \rightarrow X_{W'}.$$

834 The transport theorem proves preservation and compatibility of identity, ledger state, cell structure,
835 closure, J , one-dimensional readout, and obstruction state, and proves composition:

$$X(T_2 \circ T_1) = X(T_2) \circ X(T_1).$$

836 Together with the previous sections, this is the proof content of Theorem 1. Later parts should
837 consume this theorem by instantiating the admissible source law: Euler/L-function sources, CPS
838 twisting families, and graded obstruction towers are inputs to the same carrier interface.

839 **Theorem 26** (Unconditional ledged event exhaustion). *Let X_W be a complete realized source state.*
840 *Because the ledged projection is a bijection — the fiber reconstructs from its retained height and loss ledger*
841 *(Theorem 3; Lean `ConeProjection.record_bijective, reconstruct_record,`*
842 *`SourceHolonomy.projection_bijective_loss_ledger`) — the represented source events, carrier*
843 *events, and ledged projected events coincide:*

$$e \in E(W) \iff e \in E(X_W) \iff e \in E(\Pi(X_W), \mathcal{D}_W),$$

844 so no event is lost by projection or invented by realization. For zero events, the represented vanishing coincides
845 with the closed carrier condition,

$$\rho \in Z(W) \iff Q_W^\#(\rho) = 0,$$

846 the carrier focal closure vanishing exactly at the readout's zeros (Lean `ClosedForm.carrierZero_iff_L_zero,`
847 `FocalResidualVanishes.focal_residual_zero_iff_L_zero` — the Dirichlet-instantiated form; the
848 general zero-event identification is placewise through the local identification of Theorem 18).

849 *Proof.* The ledged projection $X_W \mapsto (\Pi(X_W), \mathcal{D}_W)$ is a bijection whose inverse is the reconstruction
850 map (Theorem 3; Lean `record_bijective, reconstruct_record, projection_bijective_loss_ledger`),
851 so it neither identifies distinct states nor omits any; the represented, carrier, and ledged projected
852 event sets are therefore in bijection, giving the three-way equivalence. For a zero-event datum, the car-
853 rier focal closure $Q_W^\#$ vanishes exactly when the readout vanishes (Lean `carrierZero_iff_L_zero,`
854 `focal_residual_zero_iff_L_zero`), so $\rho \in Z(W) \iff Q_W^\#(\rho) = 0$. $\square$

855 **Theorem 27** (Unconditional coherent carrier transport). *Call a carrier transport T faithful for a closure*
856 *predicate Cl if $Cl(x) \Rightarrow Cl(Tx)$. Then the identity transport is faithful; the composite of faithful transports*
857 *is faithful; composition is associative with the identity as two-sided unit; and the base-change transports*
858 *compose, $bc_p \circ bc_q = bc_{pq}$, with $bc_1 = id$. Hence coherent source transports induce carrier transports that*
859 *preserve closure and retained data and respect composition.*

860 *Proof.* Each law is machine-checked in `CarrierFunctoriality`: the identity is faithful (`faithful_id`);
861 a composite of faithful transports is faithful (`faithful_comp`); composition is associative (`comp_assoc`)
862 with two-sided identity (`id_comp, comp_id`); and base change composes (`bc_comp`) with unit (`bc_one`).
863 The abstract closure-transport statement is Theorem 28; these are its verified instances. $\square$

864 **Theorem 28** (Unconditional functorial closure transport). *Let C be a class of complete carrier states*
865 *equipped with a closure predicate Cl and retained datum predicate Ret . A transport $F : X \rightarrow Y$ is faithful*
866 *when*

$$Cl(X) \Rightarrow Cl(F(X)), \quad Ret(X) \Rightarrow Ret(F(X)).$$

867 Then the identity transport is faithful, and the composite of faithful transports is faithful. In particular, if

$$X \xrightarrow{F} Y \xrightarrow{G} Z$$

868 are faithful carrier transports, then $G \circ F$ preserves closure and all retained Part I data named by `Ret`.

869 *Proof.* For the identity map, both implications are tautological:

$$\text{Cl}(X) \Rightarrow \text{Cl}(X), \quad \text{Ret}(X) \Rightarrow \text{Ret}(X).$$

870 Now suppose F and G are faithful. If $\text{Cl}(X)$, then faithfulness of F gives $\text{Cl}(F(X))$, and faithfulness
871 of G gives $\text{Cl}(G(F(X)))$. The retained-datum implication is identical with `Ret` in place of `Cl`. Thus
872 $G \circ F$ is faithful. $\square$

873 **Corollary 29** (Part I assembly of the universal carrier theorem). *For every finite compatible family of*
874 *admissible structured source data with the carrier, projection, source, synchronization, closure, obstruction,*
875 *readout, reflection, analytic, exhaustion, and transport datums specified in Part I, the conclusions of Theorem 1*
876 *hold.*

877 *Proof.* The primitive double-ended carrier and involution are supplied by Theorem 2. Ledgered
878 projection and recovery are supplied by Theorem 3 and Corollary 4. Source identity and native
879 synthesis are supplied by Definition 5, Theorem 7, and Corollary 8. Finite simultaneous carrier real-
880 ization and common refinement are supplied by Theorems 9 and 10. Exact complete-cell closure and
881 obstruction-directed adaptation are supplied by Theorem 11, Theorem 12, Corollary 13, Theorem 14,
882 and Theorem 15. The retained-mode bridge, channel residual factorization, one-dimensional
883 readout identity, completed reflection, and analytic landing are supplied by Theorems 16, 17, 18, 19,
884 20, 21, 23, 24, and 25. Event exhaustion and coherent transport are supplied by Theorems 26, 27,
885 and 28. These are exactly the arrows named in Theorem 1. $\square$

886 9 The scale mismatch behind the $S(t)$ term

$S_{\text{ev}}(t)$ is not an additional native oscillatory feature of the prime/harmonic system.

887 The native system is realized on the $\pi/3$ -scaled harmonic carrier. The conventional analytic readout
888 uses a unit-1 coordinate system. These two carrier scalings represent the same arithmetic procession,
889 but they do not register the same crossings in the same coordinate units.

890 *Typing convention, fixed before the theorem.* Throughout this section $N_{\text{ev}}(t)$ denotes the carrier's dis-
891 tinct event count — the on-line ordinate census $\#\{\gamma \in (0, t] : \zeta(\frac{1}{2} + i\gamma) = 0\}$ (Lean: `zeroEventCount`)
892 — and $S_{\text{ev}}(t) := N_{\text{ev}}(t) - 1 - \vartheta(t)/\pi$ its normalized form. The multiplicity-weighted line census is
893 $N_{\text{line}}^{\text{mult}}(t) = \sum_{0 < \gamma \leq t} \text{ord}_{1/2+i\gamma} \zeta$ (Lean: `zeroEventCountMult`), with ledger $S_{\text{mult}} = N_{\text{line}}^{\text{mult}} - 1 - \vartheta/\pi$.
894 The classical Riemann–von Mangoldt term instead normalizes the strip census with multiplicity.
895 Consequently the exact bookkeeping is

$$S_{\text{class}}(t) = S_{\text{ev}}(t) + (N_{\text{line}}^{\text{mult}}(t) - N_{\text{ev}}(t)) + N_{\text{off}}^{\text{mult}}(t),$$

896 or equivalently $S_{\text{class}} = S_{\text{mult}} + N_{\text{off}}^{\text{mult}}$. Thus distinct events, excess line multiplicity, and off-line strip
897 multiplicity are separate typed ledgers. Every unconditional theorem below states explicitly which
898 one it uses.

9.1 The theorem

Let F_ζ denote the trivial-character arithmetic fiber and let $X_{\zeta,H}$ denote its realization on carrier scale H , with the fiber law, local arithmetic data, phasor amplitudes, source identity, and crossing criterion held fixed. Only the carrier scale is changed.

Theorem 30 (The $S(t)$ carrier-scale compensation theorem). *With the registered readouts of Definition 32:*

- (i) $N_{\pi/3}(e^t) = N_{\text{ev}}(t)$: the native count is the arithmetic event count over the height ray;
- (ii) $R_1(e^t) = 1 + \vartheta(t)/\pi$: the unit-chart continuous phase registration is the DC residue plus the archimedean clock, with ϑ independently derived from the gauge (the current Lean identifier for R_1 is N_1);
- (iii) consequently

$$\boxed{N_{\pi/3}(e^t) - R_1(e^t) = S_{\text{ev}}(t)}, \quad S_{\text{ev}}(t) = N_{\text{ev}}(t) - 1 - \frac{\vartheta(t)}{\pi}.$$

The two terms have different types: $N_{\pi/3}$ is cardinal-valued event data, whereas R_1 is real-valued continuous phase registration. Lean's `registeredCount` assembles the latter from an independently proved gauge clock, DC base, and scale-dependent event coefficient; it does not prove that R_1 is the cardinality of a set of unit-chart crossings. Every displayed identity is machine-checked in `RequestProject/CarrierScaleCompensation.lean` `native_identification'`, `unit_identification`, `carrier_scale_compensation_S`.

Remark 31 (Register: the classical $S(t)$). What is proven above is unconditional for the distinct-event line ledger. For the classical strip term the unconditional decomposition is

$$S_{\text{class}}(t) = \underbrace{N_{\pi/3}(e^t) - R_1(e^t)}_{S_{\text{ev}}(t), \text{ proven}} + \underbrace{(N_{\text{line}}^{\text{mult}}(t) - N_{\text{ev}}(t))}_{\text{excess line multiplicity}} + \underbrace{N_{\text{off}}^{\text{mult}}(t)}_{\text{off-line strip multiplicity}}.$$

Equivalently, the multiplicity ledger proved below satisfies $S_{\text{class}} = S_{\text{mult}} + N_{\text{off}}^{\text{mult}}$. This is the precise boundary between the unconditional line theorem and the classical strip function; neither off-line multiplicity nor simplicity is silently discarded.

9.2 Independent contour coordinate for the classical argument

The classical argument is constructed independently of every event and zero-count ledger. At a good height $T > 0$, meaning that no nontrivial zero has ordinate T , let

$$\Gamma_T : 2 \longrightarrow 2 + iT \longrightarrow \frac{1}{2} + iT.$$

The first segment lies in $\text{Re } s = 2$, where Euler-product nonvanishing applies. On the second segment, a hypothetical vanishing either has $\text{Re } s \geq 1$, again contradicting nonvanishing, or is a nontrivial zero of ordinate T , contradicting goodness. Thus $\zeta \circ \Gamma_T$ is a path in $\mathbb{C}^\times$.

Lift this path through the exponential covering $\exp : \mathbb{C} \rightarrow \mathbb{C}^\times$, with initial value

$$L_T(0) = \log \zeta(2) = \log(\pi^2/6) \in \mathbb{R}.$$

Path-lift uniqueness removes every branch choice. Define

$$S_\Gamma(T) := \frac{\Im L_T(1)}{\pi}. \tag{1}$$

926 This is a genuine global coordinate of the zeta path: its definition contains none of `zeroEventCount`,
 927 `zeroEventCountMult`, `S`, or `Smult`, and the endpoint theorem is

$$e^{L_T(1)} = \zeta\left(\frac{1}{2} + iT\right).$$

928 Combining this independently lifted phase with the independently derived Gamma clock and the
 929 real native factor gives

$$\exp(i(\pi S_\Gamma(T) + \vartheta(T))) = \frac{Z(T)}{|Z(T)|} \in \{-1, 1\}, \quad (2)$$

930 where $Z(T) = \text{rsZ}(T)$. Consequently the covering itself produces an integer winding coordinate
 931 $k_\Gamma(T)$, rather than receiving one as a counting assumption, and

$$S_\Gamma(T) = k_\Gamma(T) - \frac{\vartheta(T)}{\pi}. \quad (3)$$

932 In Lean these are `standardContour`, `zetaContourPath`, `contourLogLift`, `classicalSContour`,
 933 `contour_phase_times_clock_eq_sign_rsZ`, and `classicalSContour_eq_winding_sub_clock` in
 934 `RequestProject/ZetaContourArgument.lean`. They are proved from the analytic zeta path, the
 935 exponential covering, and the Gamma-factor factorization; no census formula is unfolded in the
 936 construction.

937 The census side is also constructed independently. Lean's `stripZeroWindow(T)` is the finite set
 938 of actual nontrivial zeros with $0 < \Im \rho \leq T$, and `stripZeroCountMult` sums their ξ -orders. Filtering
 939 this same finite set by $\text{Re } \rho = 1/2$ and $\text{Re } \rho \neq 1/2$ gives typed on-line and off-line counts. The
 940 compiled identities are

$$N_{\text{strip}}^{\text{mult}}(T) = N_{\text{on}}^{\text{mult}}(T) + N_{\text{off}}^{\text{mult}}(T), \quad N_{\text{on}}^{\text{mult}}(T) = N_{\text{line}}^{\text{mult}}(T).$$

941 They are `stripZeroCountMult_eq_onLine_add_offLine` and `onLineStripZeroCountMult_eq_zeroEventCountMult`.
 942 their composition is `stripZeroCountMult_eq_line_add_offLine`. The second equality is a
 943 proved bijection $\gamma \mapsto \frac{1}{2} + i\gamma$, with multiplicities transported by the analytic-unit order bridge
 944 `eventOrder_eq_xiOrderNat`.

945 The argument-principle comparison now identifies $k_\Gamma(T) + 1$ with this independently defined
 946 strip census by an explicit second construction, rather than by citing the classical counting formula.
 947 On the same broken path Lean forms the completed logarithmic lift

$$\mathcal{L}_{\xi,T}(u) = \log \Gamma_T(u) + \log(\Gamma_T(u) - 1) - \log 2 + \log \Gamma_{\mathbb{R}}(\Gamma_T(u)) + L_T(u).$$

948 The Gamma logarithm is first constructed on the simply connected right half-plane and normalized
 949 at (2); its imaginary part on the critical line is then proved to be the independently integrated clock
 950 $\vartheta(T)$. The two principal polynomial logarithms stay in the upper slit plane and satisfy the geometric
 951 endpoint law

$$\Im(\log(\frac{1}{2} + iT) + \log(-\frac{1}{2} + iT) - \log 2) = \pi.$$

952 Thus the value (+1) is the winding contribution of the completion factor $(s(s-1)/2)$; it is derived from
 953 the upper-half-plane geometry and is not inserted into the carrier or census definitions. Pointwise
 954 exponentiation gives $\exp \mathcal{L}_{\xi,T}(u) = \xi(\Gamma_T(u))$.

955 The continuous-lift calculus then proves, on the two affine segments,

$$i \int_0^T \frac{\xi'}{\xi}(2 + iy) dy - \int_{1/2}^2 \frac{\xi'}{\xi}(x + iT) dx = \mathcal{L}_{\xi,T}(1) - \mathcal{L}_{\xi,T}(0).$$

956 Reflection and conjugation turn this half-contour increment into the full rectangle boundary integral.
 957 Independently, the analytic divisor factorization evaluates that same boundary integral as $2\pi i$ times
 958 the multiplicity sum over *all* zeros in the upper strip, including any off-line zeros. Cancelling the
 959 nonzero factor $2\pi i$ gives the compiled theorem

$$\boxed{k_{\Gamma}(T) + 1 = N_{\text{strip}}^{\text{mult}}(T)}.$$

960 The corresponding Lean spine is `gammaLog_line_im, xiPolynomialLog_line_im, exp_xiContourLog,`
 961 `halfContourIntegral_logDeriv_riemannXi_eq_lift_sub, rectangleBoundaryIntegral_logDeriv_riemannXi`
 962 `and contourWindingIndex_add_one_eq_stripZeroCountMult`. This comparison is downstream
 963 of both independent constructions: it is not the definition of either S_{Γ} or the zero count, and no
 964 critical-line location statement is used in the strip census.

965 Eliminating the total strip census between this theorem and the independently proved on-
 966 line/off-line partition gives a sharper registration rule. At every good height,

$$N_{\text{off}}^{\text{mult}}(T) = k_{\Gamma}(T) + 1 - N_{\text{line}}^{\text{mult}}(T), \quad (4)$$

$$S_{\Gamma}(T) = \boxed{S_{\text{mult}}(T) + N_{\text{off}}^{\text{mult}}(T)} \quad (5)$$

967 These are the compiled theorems `offLineStripZeroCountMult_eq_winding_sub_line` and `classicalSContour_e`
 968 In particular, `classicalSContour_eq_Smult_iff_offLine_eq_zero` proves that the independently
 969 continued global coordinate agrees with the native multiplicity coordinate exactly when the finite-
 970 strip coverage defect vanishes. There can be no cancellation hidden in that defect: every summand is
 971 the positive analytic order of an actual zero. The pointwise theorem `mem_stripZeroFinset_line_or_coordinate_d`
 972 therefore says that each analytic strip zero is either a native line event or produces a detectable
 973 mismatch of the two global coordinates. This is the precise finite-window rule connecting the
 974 contour census, the multiplicity ledger, and the $S(t)$ registration. Finally, `exists_goodHeight_gt`
 975 constructs a good contour above every real ordinate from the finite next-window census, and
 976 `upper_nontrivialZero_line_or_globalCoordinateDefect` promotes the dichotomy globally: ev-
 977 ery upper-half-plane nontrivial zero is a native line event or exhibits an explicit higher good height
 978 where the two global coordinates differ.

979 This non-cancellation statement is now quantitative. If an enclosed zero ρ is off the carrier, then

$$\text{ord}_{\rho} \xi \leq N_{\text{off}}^{\text{mult}}(T) = S_{\Gamma}(T) - S_{\text{mult}}(T),$$

980 and the right-hand side is strictly positive. These are the compiled theorems `xiOrderNat_le_offLineStripZeroCou`
 981 `xiOrderNat_le_classicalSContour_sub_Smult_of_offCarrier`, and `classicalSContour_sub_Smult_pos_of`
 982 Thus neither a second zero nor the completed-reflection partner can cancel an off-carrier contribution
 983 in the global coordinate ledger.

984 The native branch is now carried all the way into the event-independent three-dimensional opera-
 985 tor. In `RequestProject/ZetaContourNative3D.lean`, the data type `PrincipalContourNative3DCertificate`
 986 contains a completed mod-one focal event e whose height is $e.Z = \exp(\text{Im } \rho)$, its exact am-
 987 bient physical height, the equality of its carrier readout with ρ , a nonzero eigenvector of
 988 `carrierThreeDOperator` $1 \ 1$ with eigenvalue $\Im \rho$, every spatial harmonic-pencil rank drop, and
 989 the exact one-dimensional/three-dimensional resolvent-trace relation weighted by `xiOrderNat` ρ . It
 990 now also contains a single parameter-preserving kernel coupling from `RequestProject/SpectralCarrierKernelCo`
 991 That object retains simultaneously the nonzero kernel of the configured full analytic operator
 992 $L_{\text{spec}}(\rho) \text{ id}$, the completed focal event at the same analytic point, the ambient carrier eigenvector, and
 993 the self-adjoint one-mode resolvent kernel. The constructor `ofLspecZeroAtHeight` builds this object

994 from $L_{\text{spec}}(\sigma_* + i \log Z) = 0$ at every positive carrier height; conversely `lspec_zero_of_coupling`
 995 proves that the spatial coupling cannot create an analytic vanishing. The exact identification
 996 theorems `nonempty_coupling_iff` and `nonempty_coupling_iff_readoutKernel` state that such a
 997 coupling exists precisely when the analytic kernel and the independently derived carrier coordinate
 998 refer to the same point. Thus neither operator is silently replaced by the other, and the midpoint
 999 coordinate is supplied by carrier membership. The finite-window defect has now been rewritten
 1000 entirely in this typed language. In `RequestProject/ZetaContourCouplingDefect3D.lean`, define
 1001 $N_{\text{uncpl}}^{\text{mult}}(T)$ by summing the positive analytic order over precisely those strip kernels for which no
 1002 `SpectralCarrierKernelCoupling3D` exists. The compiled identities

$$N_{\text{uncpl}}^{\text{mult}}(T) = N_{\text{off}}^{\text{mult}}(T), \quad S_{\Gamma}(T) = S_{\text{mult}}(T) + N_{\text{uncpl}}^{\text{mult}}(T)$$

1003 are `uncoupledKernelCountMult_eq_offLine` and `classicalSContour_eq_Smult_add_uncoupledKernelCount`.
 1004 Because each included order is positive, `classicalSContour_eq_Smult_iff_everyKernelCoupled`
 1005 proves that the two global coordinates agree exactly when every enclosed full analytic kernel has a
 1006 parameter-preserving coupling to the independently defined carrier operator. Hence an absent cou-
 1007 pling cannot be hidden by cancellation between zeros. This has a midpoint-free no-drift form. For ev-
 1008 ery base $n > 1$, `ofLspecZeroOfNoRadialDrift` constructs the full coupling from analytic vanishing
 1009 and $n^{\text{Re } s - \sigma_*} = 1$; it derives $\text{Re } s = \sigma_*$ internally from `noRadialDrift_iff_carrierAbscissa`. The
 1010 independent constructor `ofLspecZeroOfBinaryNoRadialDrift` repeats the construction through
 1011 the physical height $Z = \exp(\text{Im } s)$. Thus no numerical midpoint is an input to either con-
 1012 structor. At finite-window scale, `uncoupledKernelCountMult_eq_radialDrift` proves that the
 1013 uncoupled count is exactly the positive multiplicity of kernels with nontrivial radial drift,
 1014 and `classicalSContour_eq_Smult_iff_everyKernelNoRadialDrift` identifies global-coordinate
 1015 agreement with zero drift for every enclosed kernel. These finite-window statements have also been
 1016 globalized without changing their content. In `RequestProject/ZetaContourGlobalIdentification3D.lean`,
 1017 the theorem `globalCoordinateIdentification_iff_upperKernelCoupling` identifies agreement
 1018 at every good height with a full parameter-preserving coupling for every upper nontrivial analytic
 1019 zero. The corresponding theorem `globalCoordinateIdentification_iff_upperNative3DCertificate`
 1020 upgrades each such coupling to the complete physical-height focal-event, ambient-eigenvector,
 1021 harmonic-rank-drop, twin-trace, and multiplicity certificate. Independently, `globalCoordinateIdentification_i`
 1022 obtains the same global criterion directly by enclosing each zero in a higher good contour.

1023 There is now also a conservation-native source formulation with no literal half-unit in its
 1024 definition. For a conservative source mode ψ , set

$$\text{coord}_{\text{src}}(\psi) := \sigma_* + \text{rate}(\psi),$$

1025 where σ_* is the unique area-law-selected carrier abscissa. The theorem `conservativeCarrierCoord_re`
 1026 derives $\text{Re coord}_{\text{src}}(\psi) = \sigma_*$ from source conservation, and `conservativeCarrierCoord_noRadialDrift`
 1027 derives unit radial projection at every base greater than one. Finally, `globalCoordinateIdentification_iff_upper`
 1028 proves that all-good-height coordinate agreement is exactly the statement that every upper analytic
 1029 zero is the coordinate of one of these conservative carrier modes. This source identification has a di-
 1030 rect pointwise landing as well: `principalContourNative3DCertificate_of_noRadialDrift` con-
 1031 structs the complete certificate from a zero and its no-drift projection, while `principalContourNative3DCertificat`
 1032 constructs it from a conservative source representation. Neither constructor assumes any global
 1033 coordinate identity; both produce the physical event and operator data at that single analytic
 1034 parameter. The analytic-zero side is no longer represented only by a scalar membership predicate.
 1035 The compiled transfer in `RequestProject/ZetaZeroNative3DSourceTransfer.lean` begins with
 1036 every actual $\rho \in \text{ZD.NontrivialZeros}$ and constructs `PrincipalZeroAnalyticFiber3D`(ρ). Its

state is the literal eta-configured Vec3 fiber at the original complex parameter; after division by the nonvanishing eta factor, its plane partial sums converge to $\zeta(\rho) = 0$. The same object carries the event-independent ambient helix state at ordinate $\Im\rho$, positive physical height $Z = \exp(\Im\rho)$, and its nonzero eigenvector with eigenvalue $\Im\rho$. None of these fields places ρ on a preselected vertical line.

Before any parameter-preserving carrier-to-chart identification, the compiled constructor `principalZero_focalCancellation_on_carrier` packages this closure as a `PrincipalZeroNativeCarrierEvent`. Its `carrierState` is a typed member of the independently defined Archimedean helix, and its stored analytic fiber closes at the same physical height. The theorems `.on_archimedeanHelix`, `.physicalHeight_eq_focalHeight`, and `.ambient_eigenvector` prove, respectively, literal helix membership, equality of the carrier and focal heights, and the ambient 3D eigenvector equation. Most importantly, `.noRadialDrift` is derived directly from that carrier membership by `carrierPointAtHeight_noRadialDrift`. The event type and this proof use neither the Euclidean magnitude of the analytic fiber nor a radial-limit hypothesis. Only afterwards does `.carrierReadout_scaleBalanced` apply the area law to the carrier readout. Thus focal cancellation is first marked at $Z = \exp(\Im\rho)$ on the helix, no drift is a fact about that helix state, and equality of the resulting carrier coordinate with the original complex parameter remains the separate chart-identification theorem.

The two uses of “rides” are now separated at the type level. For an analytic zero fiber F_ρ , Lean defines `PrincipalZeroAnalyticFiber3D.RidesCarrier` by the state equality

$$F_\rho = \text{spectralFiber } \text{etaW}(\text{carrierPointAtHeight}(Z_\rho)) \quad \text{in } (\mathbb{N} \rightarrow \text{Vec3}).$$

This is the fiber-to-carrier attachment; it is stronger than sharing the ordinate or physical height. The carrier-to-helix map is the independent field `carrierState.point = Geometry.gammaY(1, 1, carrierState.ordinate)`. The theorem `PrincipalZeroAnalyticFiber3D.ridesCarrier_iff_carrierWeld` proves that the first state equality is equivalent to the parameter-preserving weld `carrierPointAtHeight(Z_\rho) = \rho`: its imaginary part comes from $\log Z_\rho = \Im\rho$, while equality of the radial exponents is extracted from the actual three-dimensional magnitudes at the nontrivial carrier site $n = 2$. Composing with the area law gives `.ridesCarrier_iff_scaleBalanced`. Thus no equality of unlike objects is used and no bare midpoint is placed in the ride definition.

Exact focal cancellation itself cannot occur off the carrier. Lean defines `CompletedFocalCancellationRepresents` by requiring one positive physical height Z for which the completed residual `Dcell` χZ vanishes and whose readout `carrierPointAtHeight Z` is the same parameter ρ . The compiled theorem `completedFocalCancellationRepresents_iff` proves

$$\text{CompletedFocalCancellationRepresents}(\chi, \rho) \iff L(\rho, \chi) = 0 \text{ and } \text{Re } \rho = \sigma_*.$$

The reverse direction reconstructs $Z = \exp(\Im\rho)$; the real coordinate is read from the Archimedean area-law carrier. Consequently `no_offCarrier_completedFocalCancellation` proves pointwise that a parameter with $\text{Re } \rho \neq \sigma_*$ cannot be represented by focal cancellation. This precedes every resolvent trace and every zero count, so there is no cross-event or reflected-pair cancellation mechanism to invoke.

The second structure, `PrincipalZeroNative3DSourceTransfer`, records the geometric projection of that analytic fiber into a conservative source mode and the complete native focal/operator certificate at the same parameter. Its constructor `principalZeroNative3DSourceTransfer_of_noRadialDrift` accepts the area-law equation $n^{\text{Re } \rho - \sigma_*} = 1$, derives the source coordinate internally, and never accepts $\text{Re } \rho = 1/2$. Conversely the source conservation field recovers that no-drift equation at every $n > 1$. Hence the compiled pointwise theorem `nonempty_principalZeroNative3DSourceTransfer_iff_noRadialDrift` identifies the exact projection step,

rather than hiding it inside a statement that a zero already equals a carrier point. Globally, the native direction no longer needs that radial equation as an input. The theorem `carrierPointAtHeight_noRadialDrift` proves zero radial drift for every physical-height carrier readout solely from helix membership, and `completedThreeDZeroAtHeight_focal_rankDrop_eigenvector_noRadialDrift` assembles, for one and the same event, the completed focal residual zero, the spatial harmonic-pencil rank drop, the ambient carrier eigenvector at $\log Z$, and the no-drift conclusion. At the transfer layer, `PrincipalContourNative3DCertificate.analyticFiber_eq_carrierFiber` proves that the literal `spectralFiber_etaW_rho` is the carrier fiber at the certificate's marked height, while `PrincipalContourNative3DCertificate.noRadialDrift` derives its radial equation from that identification. Consequently `principalZeroNative3DSourceTransfer_of_nativeCertificate` constructs the full source transfer from native realization itself, with neither a midpoint nor a no-drift proposition supplied. The physical-height identification is also literal, not merely an agreement of ordinates: `PrincipalZeroNative3DSourceTransfer.eventHeight_eq_focalCancellationHeight` proves that the native event height equals the analytic fiber's stored focal-cancellation height. After rewriting by that equality, `.harmonicRankDrop_at_focalCancellationHeight` proves that the completed spatial pencil rank-drops at precisely that height. The combined theorem `.focalCancellation_rankDrop_noRadialDrift_sameHeight` returns the fiber readout closure, pencil rank drop, and helix no-drift law at this one shared physical height. Moreover every completed source transfer proves the attachment itself by `PrincipalZeroNative3DSourceTransfer.fiberRidesCarrier`; it then derives the original analytic parameter from the carrier readout by `.carrierReadout_eq_analyticParameter`. The composed theorem `.fiberRidesCarrier_carrierRidesHelix` returns, in one statement, the fiber-state attachment, literal membership of the carrier point in Γ_Y , and equality of the two physical focal heights. At the global level, `globalCoordinateIdentification_iff_allCanonicalZeroFibersRideCarrier` states the remaining identification exactly as this state-level attachment for every canonical analytic zero fiber. The assembly has also been closed without passing through the contour coordinate: `nonempty_principalZeroNative3DSourceTransfer_iff_fiberRidesCarrier` proves, zero by zero, that existence of the completed native source transfer is exactly equality of the two complete $\mathbb{N} \rightarrow \text{Vec3}$ fiber states. Quantifying this statement gives `allNative3DSourceTransfer_iff_allCanonicalZeroFibersRideCarrier`, and `allNative3DSourceTransfer_iff_zeroLedger_eq_carrierHeight` identifies that same family-level attachment with equality of the analytic zero-ledger and event-independent carrier-height operators. Thus the source, state, and operator formulations are proved equivalent; none silently substitutes a bare equality of ordinates for fiber attachment.

Globally, `globalCoordinateIdentification_iff_upperNative3DSourceTransfer` states that all-good-height coordinate agreement is exactly transfer of every upper zero from its genuine three-dimensional analytic fiber to the conservative native source. Finally, `upperNative3DSourceTransfer_of_selfAdjointXiReceiver` performs this full transfer for the independent self-adjoint xi-receiver route. The conjugation-complete theorem `globalCoordinateIdentification_iff_allNative3DSourceTransfer_of_selfAdjointXiReceiver` upgrades the statement to every nontrivial zero in both half-planes, and `allNative3DSourceTransfer_of_selfAdjointXiReceiver` performs that complete two-sided transfer through the same receiver. The final operator propagation is now direct and basiswise. Given those two-sided transfers, `xiZeroLedgerOperator_eq_carrierHeight_of_allNative3DSourceTransfer` proves

$$\text{xiZeroLedgerOperator} = \text{xiCarrierHeightOperator}$$

by evaluating the two diagonal operators on each canonical analytic-zero mode and using only that mode's compiled scale-balance theorem. No trace sum or cross-event cancellation appears. The theorem `globalCoordinateIdentification_of_allNative3DSourceTransfer` then feeds this operator equality into the contour/carrier unification and returns `classicalSContour = Smult` at every

1126 good height. At the receiver endpoint, `globalCoordinateIdentification_and_zeroLedger_eq_`
 1127 `carrierHeight_of_selfAdjointXiReceiver` returns both conclusions in one certificate. The
 1128 state-space form is sharper still. At every positive site n , `PrincipalZeroAnalyticFiber3D.`
 1129 `radialMagnitude` computes directly from the literal `Vec3` state

$$\text{mag}_3(F_\rho(n)) = n^{-\text{Re } \rho}.$$

1130 Consequently `PrincipalZeroAnalyticFiber3D.carrierScaleBalanced_iff_radialMagnitude`
 1131 identifies the additional parameter-preserving scale condition with a positive finite limit of
 1132 $\text{mag}_3(F_\rho(n))R_{\text{carrier}}(n)$. This statement contains neither a carrier point nor a supplied midpoint. It is
 1133 a diagnostic for whether the analytic exponent agrees with the already no-drift carrier readout, not
 1134 the source of native helix membership or native no drift. The numerical half-unit is recovered only
 1135 when the independently proved Archimedean area law evaluates the unique balanced exponent.
 1136 The analytic/operator identification has also been completed before any event certificate is
 1137 introduced. In `RequestProject/XiZeroLedgerResolvent3D.lean`, the affine chart identity

$$\left(\frac{1}{2} + iz\right) - \rho = i(z - \text{poleParam}(\rho))$$

1138 transports the proved Hadamard partial fraction term by term and gives `xiChannel_eq_constant_`
 1139 `add_zeroResolventTrace`:

$$-\frac{\xi'}{\xi}\left(\frac{1}{2} + iz\right) = C + \sum_{\rho} m_{\rho} \left(\frac{i}{z - \text{poleParam}(\rho)} - \frac{1}{\rho} \right).$$

1140 Thus the half-unit is the affine carrier readout unit used in an independently proved change of
 1141 coordinates, not a zero-location field. The associated `xiZeroLedgerOperator` is defined on the free
 1142 finitely supported Hilbert space on analytic zeros and has $\text{poleParam}(\rho)$ as an actual eigenvalue
 1143 by `xiZeroLedgerOperator_hasEigenvalue`. Its definition mentions no `ThreeDZero`, focal event,
 1144 or equality with `carrierPoint`. Spectral symmetry then yields the area-law balance event by
 1145 event through `xiZeroLedgerOperator_symmetric_imp_scaleBalanced`. Finally the compiled
 1146 unification theorem `globalCoordinateIdentification_iff_xiZeroLedgerOperator_symmetric`
 1147 states that symmetry of this independently identified resolvent operator is exactly the all-good-
 1148 height identity $S_{\Gamma} = S_{\text{mult}}$. This rules out an “offline cancellation” inside the comparison: analytic
 1149 multiplicities are positive, and every diagonal pole is carried by its own nonzero basis eigenvector.
 1150 More explicitly, the compiled operator identity `xiZeroLedgerOperator_eq_height_sub_I_smul_`
 1151 `radialDrift` is

$$H_{\xi} = H_{\text{carrier}} - iD_{\text{radial}}, \quad D_{\text{radial}}e_{\rho} = \left(\text{Re } \rho - \frac{1}{2}\right)e_{\rho}.$$

1152 Here both H_{carrier} and D_{radial} are independently constructed real-diagonal symmetric opera-
 1153 tors; the half-unit in D_{radial} is the value selected by the preceding area law. The theorems
 1154 `xiRadialDriftOperator_eq_zero_iff_all_scaleBalanced` and `xiZeroLedgerOperator_isSymmetric_`
 1155 `iff_radialDrift_eq_zero` prove, basis mode by basis mode, that symmetry of the analytic pole op-
 1156 erator is precisely extinction of radial drift, not cancellation of drift after summing different zeros. At
 1157 global scale this is compiled in the two interchangeable forms `globalCoordinateIdentification_`
 1158 `iff_xiRadialDriftOperator_eq_zero` and `globalCoordinateIdentification_iff_zeroLedger_`
 1159 `eq_carrierHeight`. There is also a canonical projection statement, rather than a bare instruction to
 1160 discard the radial coordinate. On the same finitely supported zero-state space, `xiZeroLedgerFormalAdjoint`
 1161 is defined coefficientwise and satisfies the exact Green identity

$$\langle H_{\xi} f, g \rangle = \langle f, H_{\xi}^{\dagger} g \rangle$$

1162 by `xiZeroLedgerFormalAdjoint_green`. Its Hermitian part is computed, not postulated:

$$P_{\text{nd}}(H_\xi) := \frac{1}{2}(H_\xi + H_\xi^\dagger) = H_{\text{carrier}},$$

1163 the theorem `xiZeroLedgerNoDriftProjection_eq_carrierHeight`. The complementary anti-
1164 Hermitian part is exactly the radial ledger,

$$D_{\text{radial}} = \frac{i}{2}(H_\xi - H_\xi^\dagger),$$

1165 by `xiRadialDriftOperator_eq_I_half_smul_adjointDefect`. Hence the all-good-height identity
1166 is equivalently the fixed-point equation $H_\xi = P_{\text{nd}}(H_\xi)$, compiled as `globalCoordinateIdentification_`
1167 `iff_zeroLedger_fixedByNoDriftProjection`. This makes precise the phrase “no-drift projection”:
1168 the carrier height and the drift are the Hermitian and anti-Hermitian components of the indepen-
1169 dently identified analytic pole operator, and separate basis modes cannot cancel one another in either
1170 component. Indeed `xiZeroLedgerResidual_eq_zero_iff_each_basis` proves that the projection
1171 residual vanishes as an operator if and only if it vanishes on every canonical zero basis vector sepa-
1172 rately, and `globalCoordinateIdentification_iff_eachZeroBasisProjectionResidual_eq_zero`
1173 identifies that basiswise condition with the global coordinate equality. The pointwise computation
1174 is literal:

$$(H_\xi - P_{\text{nd}}H_\xi)e_\rho = -i \left(\text{Re } \rho - \frac{1}{2} \right) e_\rho,$$

1175 proved by `xiZeroLedgerProjectionResidual_on_basis`. Finally `principalZeroProjectionResidual_`
1176 `eq_zero_iff_radialMagnitude` identifies vanishing of this same basis residual with the positive
1177 finite area-normalized Euclidean-magnitude limit of the actual eta-configured 3D fiber. Thus the
1178 operator, state-space, and area-law formulations agree at each event before any global summation
1179 is taken.

1180 There is also a direct receiver formulation. For every complex sample z , `zeroPoleResolvent_`
1181 `eq_carrierBasisReadout_iff_scaleBalanced` proves

$$i(z - \text{poleParam}(\rho))^{-1} = i \langle e_{\Im \rho}, (z - H_{\text{carrier}})^{-1} e_{\Im \rho} \rangle \iff F_\rho \text{ is area-law scale balanced.}$$

1182 Only injectivity of inversion and of multiplication by i is used, so one sample already iden-
1183 tifies the pole parameter and no trace-level cancellation is available. The global assembly
1184 `globalCoordinateIdentification_iff_eachZeroPoleResolvent_eq_carrierBasisReadout` states
1185 that $S_\Gamma = S_{\text{mult}}$ at all good heights exactly when this equality holds separately on every analytic-zero
1186 basis mode. The regularized trace bridge is now typed at the same resolution. Lean defines
1187 `xiZeroResolventKernel` and `xiCarrierHeightResolventKernel` with the same positive analytic
1188 multiplicity and the same Hadamard counterterm, but with pole coordinates `poleParam( $\rho$ )` and `$\Im \rho$` ,
1189 respectively. The pointwise theorem `xiZeroResolventKernel_eq_carrierHeightKernel_iff_`
1190 `scaleBalanced` cancels the positive multiplicity and proves that the two kernels agree exactly at the
1191 area-law fixed point. Hence `globalCoordinateIdentification_iff_basisResolvedCarrierResolventBridge`
1192 identifies the all-good-height theorem with equality of these two kernels for every (z, ρ) , before sum-
1193 mation. Only after that basiswise equality is established does `xiCarrierHeightResolventTrace`
1194 take the `tsum`; `carrierHeightResolventTrace_of_globalCoordinateIdentification` then gives
1195 the literal carrier formula

$$-\frac{\xi'}{\xi} \left(\frac{1}{2} + iz \right) = C + \text{xiCarrierHeightResolventTrace}(z).$$

Thus the global theorem, the operator equality, and the resolvent-trace bridge have the same basiswise weld, and no scalar trace equality is used to infer a zero-by-zero identification. For the paper's stated primary-3D/shadow-1D conditional, Lean packages the whole implication in `globalCoordinateIdentification_basisResolvedBridge_and_carrierTrace_of_allNative3DSourceTransf`. The theorem consumes one native 3D source transfer for each analytic shadow and returns the all-good-height coordinate identity, every basis-resolved kernel equality, and the summed carrier-height trace together. In particular, the half-unit is delivered inside the native no-drift/area-law transfer and is not repeated as an independent assumption of this assembly theorem. The upper-half-plane census has now been joined to the complete two-sided analytic channel in `RequestProject/ZetaContourXiReceiverIdentification3D.lean`. The theorem `conj_mem_nontrivialZeros` transports genuine zeros by the real structure of the entire ξ function, while `nontrivialZero_im_ne_zero` consumes the independently proved zero-free region $0 < \operatorname{Re} s < 1, |\operatorname{Im} s| < 2$, to exclude a real-axis member. Consequently `upperNoRadialDrift_iff_allNoRadialDrift` promotes the upper no-drift statement to the complete nontrivial zero set.

This gives the exact analytic receiver formulation

$$(\forall T \text{ good}, S_{\Gamma}(T) = S_{\text{mult}}(T)) \iff \text{the } \xi\text{-channel is regular off the real spectral axis,}$$

compiled as `globalCoordinateIdentification_iff_xiChannel_offReal_regular`. Thus the coordinate identification and the analytic receiver are now one theorem, not parallel narrative interfaces. Finally, `globalCoordinateIdentification_and_upperNative3D_of_selfAdjointXiReceiver` proves that a self-adjoint receiver of this independent channel supplies both the all-good-height equality and, for every upper zero, the complete physical-height 3D/operator certificate described above. The operator geometry needed by that receiver is now constructed on the full, event-independent state space in `RequestProject/CarrierUnboundedResolvent3D.lean`. Because the carrier ordinate ranges over all of $\mathbb{R}$, the native height operator H is unbounded; it is therefore represented directly on finitely supported carrier modes rather than being mislabeled as a bounded C-star-algebra element. For $\operatorname{Im} z \neq 0$, the diagonal formula

$$(z - H)^{-1} e_s = \frac{1}{z - s.\text{ordinate}} e_s$$

is proved to be both a left and a right inverse of $zI - H$ by `carrierThreeDShift_comp_resolvent` and `carrierThreeDResolvent_comp_shift`. Moreover, `carrierBasisResolventReadout_regular_off_real` proves that every geometric basis matrix coefficient has a finite punctured limit at every nonreal parameter. These are properties of the Archimedean helix state space itself: the construction mentions no zero predicate, L-function, focal event, or event certificate. A bounded C-star receiver is used only after a compact coordinate transform; the unbounded 3D height operator retains the literal ordinate and hence the full real spectral axis. The all-height theorem `upper_nontrivialZero_kernelCoupling_or_globalCoordinateDefect` then returns, for every upper nontrivial zero, either this full typed coupling or an explicit higher good height at which the two coordinates differ. The compiled capstone `upper_nontrivialZero_native3DCertificate_or_allowbreak_globalCoordinateDefect` constructs this complete certificate or returns the explicit higher good height where the global coordinates differ. Thus the contour census, physical-height event, ambient eigenvector, harmonic pencil, analytic multiplicity, and both trace charts are joined without defining the ambient operator from the zero certificate.

The proof occupies the rest of the section: part (i) in the native- $\pi/3$ subsection, part (ii) in the unit-1 subsection, part (iii) by subtraction; the induced event readout pinning every functor input is derived in the discharged-arrows subsection.

9.3 The event count and the phase registration

Definition 32 (Scale-indexed registered readout). For a carrier realization $X_{\zeta,H}$, write

$$C_H(F_\zeta; z) = \{ c : c \text{ is a complete } F_\zeta\text{-closure in } X_{\zeta,H} \text{ through raw height } z \}.$$

At the native scale the cumulative crossing count is

$$N_{\pi/3}(e^t) = \#C_{\pi/3}(F_\zeta; e^t).$$

The unit chart instead supplies the real-valued phase registration

$$R_1(e^t) = 1 + \frac{\vartheta(t)}{\pi}.$$

The theorem compares these two readouts; it does not identify the continuous quantity R_1 with a cardinality. A topological-degree or contour-winding coordinate for the classical argument is instead constructed in (1)–(3); it is a path lift of the actual zeta fiber, not a reinterpretation of R_1 as a set cardinality.

In Lean the scale-indexed registered readout is installed as `registeredCount`: DC base at the common origin, an independently constructed continuous clock procession in π -units, and the event count weighted by the induced per-event contribution of the scale's closure geometry (`eventContribution`; derivation in the discharged-arrows subsection below). The closure predicate is `NativeClosure` (the fiber's η -accumulation) and the counts are `nativeCount` and `zeroEventCount`.

9.4 Native $\pi/3$ chain

The native chain already present in the carrier framework is:

$$F_\zeta \longrightarrow X_{\zeta,\pi/3} \longrightarrow C_{\pi/3}(F_\zeta; e^t) \longrightarrow N_{\pi/3}(e^t).$$

Its local pieces are the $\pi/3$ cell law

$$\text{cell}(n) = \exp(in\pi/3),$$

the antipodal cancellation

$$\text{cell}(n) + \text{cell}(n + 3) = 0,$$

the helix pencil closure equivalence, and the exact height readout

$$\text{heightReadout}(e^\gamma) = \gamma.$$

In Lean these correspond to the native cell and pencil statements in `GeometricPhasorClosure.lean` and to the readout lemmas

$$\text{cell_antipodal_cancel}, \quad \text{helix_cancellation_logfree}, \quad \text{readout_exp}$$

in `GeometricReadout.lean`.

Proof of Theorem 30(i). A complete closure of the fiber at ordinate γ is exactly an on-line zero event: the fiber's own η -accumulation closes at γ if and only if the readout vanishes there (`nativeClosure_iff_zero`, via `Faithful.continuous_model_zeta`), and the height readout carries raw height e^t to ordinate t exactly (`readout_exp`). Here $N_{\text{ev}}(t)$ is the zero-counting procession over the 3D representation's *entire* event space — the height ray, whose exhaustion is a theorem, consumed not assumed (`native_events_sourced`, instantiating the proven `threeD_exhaustive`; the height encoding is lossless, `event_encoding_complete`). Counting complete closures through raw height e^t therefore counts exactly the arithmetic events through ordinate t : $N_{\pi/3}(e^t) = N_{\text{ev}}(t)$ (`native_identification'`). No $S(t)$ term is an additional state variable in $X_{\zeta,\pi/3}$. $\square$

9.5 Unit-1 phase-registration chain

The unit-1 chain is the scalar phase registration of the same fiber:

$$F_\zeta \longrightarrow X_{\zeta,1} \longrightarrow R_1(e^t).$$

The required scalar readout identity is Theorem 30(ii): $R_1(e^t) = 1 + \vartheta(t)/\pi$.

Proof of Theorem 30(ii). The arrow is obtained from the $H = 1$ carrier/readout construction itself (the numerical layer's smooth count and crossing-spacing audit remain as cross-checks). The clock is constructed on the carrier,

$$\vartheta(t) = \int_0^t \operatorname{Re} \left(\frac{\Gamma'_\mathbb{R}}{\Gamma_\mathbb{R}} \left(\frac{1}{2} + iu \right) \right) du \quad (\text{theta}),$$

proven to be the continuous phase of the archimedean gauge, $\Gamma_\mathbb{R}(\frac{1}{2} + it) = \|\Gamma_\mathbb{R}(\frac{1}{2} + it)\| e^{i\vartheta(t)}$ (gauge_polar), unique under the origin normalization (theta_unique). The master registration identity $\zeta(\frac{1}{2} + it) = Z(t) e^{-i\vartheta(t)}$ with $Z(t)$ real (unit_chart_factorization) shows the unit readout's entire continuous phase is clock; the DC base is the trivial-channel residue, value forced = 1 (dcResidue_spec, dcResidue_unique); and the event channel is shut by the arc geometry — no whole-cell walk of the unit carrier realizes the half-turn mark (unit_arcs_empty, eventContribution_one). Assembling base, clock, and the vanishing event channel in the registered readout gives $R_1(e^t) = 1 + \vartheta(t)/\pi$ (unit_identification). The clock itself is independently derived and characterized by gauge_polar and theta_unique; the assembly formula registeredCount then uses that derived clock as its continuous input. $\square$

9.6 Subtraction

Proof of Theorem 30(iii). With parts (i) and (ii) supplied, substitution in the fixed definition $S_{\text{ev}} = N_{\text{ev}} - 1 - \vartheta/\pi$ gives

$$\begin{aligned} N_{\pi/3}(e^t) - R_1(e^t) &= N_{\text{ev}}(t) - \left(1 + \frac{\vartheta(t)}{\pi} \right) \\ &= S_{\text{ev}}(t). \end{aligned}$$

Thus $S_{\text{ev}}(t)$ is the scalar compensation ledger for registering the native $\pi/3$ arithmetic procession in the unit-1 chart (carrier_scale_compensation_S). $\square$

9.7 The exact clock-jump rule and its bound

The multiplicity ledger has a stronger, non-asymptotic formulation. For $0 \leq a \leq b$, put

$$M(a, b) := \sum_{a < \gamma \leq b} \operatorname{ord}_{1/2+i\gamma} \zeta, \quad C(a, b) := \int_a^b \operatorname{Re} \frac{\Gamma'_\mathbb{R}}{\Gamma_\mathbb{R}} \left(\frac{1}{2} + it \right) dt.$$

These inputs are defined independently from the analytic zero orders and the Archimedean gauge. The compiled finite-window law is

$$\boxed{S_{\text{mult}}(b) - S_{\text{mult}}(a) = M(a, b) - \frac{C(a, b)}{\pi}.} \quad (6)$$

1293 It immediately gives the unconditional bound

$$|S_{\text{mult}}(b) - S_{\text{mult}}(a)| \leq M(a, b) + \frac{1}{\pi} \int_a^b \left| \operatorname{Re} \frac{\Gamma'_{\mathbb{R}}}{\Gamma_{\mathbb{R}}} \left(\frac{1}{2} + it \right) \right| dt. \quad (7)$$

1294 On a zero-free interval the discrete term vanishes, so equality $S_{\text{mult}}(b) - S_{\text{mult}}(a) = -C(a, b)/\pi$ holds
 1295 and the right side of the bound contains only the clock budget. At every interior non-event,

$$S'_{\text{mult}}(t) = -\frac{\text{clockRate}(t)}{\pi};$$

1296 at an event its jump is the analytic multiplicity, which is also the residue of ζ'/ζ . In distributional
 1297 notation, summarizing the compiled finite-window statements,

$$dS_{\text{mult}} = \sum_{\gamma > 0} m_{\gamma} \delta_{\gamma} - \frac{\text{clockRate}(t)}{\pi} dt.$$

1298 The Lean proofs are `Smult_interval_clock_jump`, `abs_Smult_interval_le_multiplicity_add_clock`,
 1299 `abs_Smult_interval_le_clock_of_event_free`, and `Smult_hasDerivWithinAt_of_event_free`
 1300 in `RequestProject/StClockJumpLaw.lean`; the residue/jump identity is `residue_eq_Smult_jump`
 1301 in `RequestProject/ResidueJump.lean`. Thus the result supplies a local equation of motion, an
 1302 exact jump rule, and a finite-window bound without identifying the line ledger with the classical
 1303 strip argument.

1304 **Remark 33** (No native oscillatory state). The structural consequence — $S_{\text{ev}}(t)$ has no additional
 1305 native oscillatory carrier state — is `S_has_no_native_carrier` with `no_native_oscillation`:
 1306 despinning the unit readout by the clock leaves a real value at every height. The distinct and
 1307 multiplicity-weighted line ledgers record the event jumps against that continuous clock. Per
 1308 Remark 31, the classical strip term additionally contains off-line strip multiplicity.

1309 9.8 The arrows, discharged

1310 The native cumulative count and the unit phase registration are assembled in `RequestProject/CarrierScaleComper`
 1311 Their inputs have the following proved origins:

- 1312 • **Event channel.** An event is a sign flip of the real native state; its mark is the half turn $-1 = e^{i\pi}$
 1313 (`sign_flip_mark`). A scale- H chart registers the mark only through a completed cell arc realizing
 1314 it, `eventArcs` $H = \{k \geq 1 : e^{ikH} = -1\}$. At $\pi/3$ the wall is hit exactly once per closed six-cell
 1315 monodromy loop (`pi3_wall_mem`, from the antipodal identity; `pi3_arcs_eq`: the arcs are exactly
 1316 $k \equiv 3 \pmod{6}$) — one registered event per completed native cell. At scale 1 no arc ever realizes
 1317 the mark (`unit_arcs_empty`). The induced per-event contribution (`eventContribution`) is 1 at
 1318 $\pi/3$ and 0 at 1 by theorems, not by a gate.
- 1319 • **Clock channel.** The native chart's zero clock is derived, not assigned: every continuous clock its
 1320 readout factors through is constant on event-free stretches (`native_clock_locally_constant`,
 1321 from `no_native_oscillation`). The unit chart's clock is ϑ , the unique continuous gauge phase
 1322 (`gauge_polar`, `theta_unique`).
- 1323 • **DC base.** The origin registration is the trivial-channel residue, value forced (`dcResidue_spec`,
 1324 `dcResidue_unique`).

The geometric scale gap is prior to the analytic event data: the native and unit placements of the integer procession are incommensurable — the lattices $\{k\pi/3\}$ and $\{m\}$ meet only at the origin (`lattice_gap_fundamental`); the unit carrier closes no cell (`unit_never_closes`) and has no antipodal wall (`unit_cell_no_antipodal`). The compiled compensation identity compares the native event count with the unit phase registration against that scale geometry.

9.9 The general registration gap $S_{H,K}$ and the harmonic family

The comparison generalizes to arbitrary scale pairs. With R_H the canonical scale- H registered readout (Lean identifier `NH`; by `NH_dichotomy` the family contains exactly the two proven chart types, decided by the arc geometry), define

$$S_{H,K}(t) := R_H(e^t) - R_K(e^t) \quad (\text{Sgap}).$$

Proposition 34 (Coboundary laws). *The registration gap is a coboundary of the per-scale potential: $S_{H,H} = 0$, $S_{H,K} = -S_{K,H}$, and the cocycle law $S_{H,L} = S_{H,K} + S_{K,L}$ holds for all scales. $S(t)$ is relative — a gap between registrations — never intrinsic.*

Proof. Immediate from the definition as a difference of the per-scale potential R_H (`Sgap_refl`, `Sgap_antisymm`, `Sgap_cocycle`). $\square$

Theorem 35 (One S for the whole harmonic family). *Every harmonic scale π/m ($m \geq 1$) realizes the event mark at m cells, exactly once per closed $2m$ -cell monodromy loop; the family registers identically, and every member reads the same distinct-event ledger $S_{\text{ev}}(t)$ against the unit chart:*

$$S_{\pi/m,1}(t) = S_{\text{ev}}(t), \quad \text{i.e.} \quad R_{\pi/m}(e^t) = R_1(e^t) + S_{\text{ev}}(t) \quad \text{for every } m \geq 1.$$

Proof. The wall exists at every harmonic scale: $m \cdot (\pi/m) = \pi$, so the m -cell walk lands on the half turn (`pi_div_wall`); the arcs realizing the mark are exactly $k \equiv m \pmod{2m}$, one per closed loop (`pi_div_arcs_eq`); monodromy follows by doubling the wall (`pi_div_closes`). Hence the event channel is open with weight one at every π/m , and the intra-family gap vanishes (`Sgap_pi_div = 0`). The cocycle law of Proposition 34 composes through $\pi/3$: $S_{\pi/m,1} = S_{\pi/m,\pi/3} + S_{\pi/3,1} = 0 + S_{\text{ev}}$ (`Sgap_pi3_one`, `Sgap_pi_div_one`, `unit_tracks_harmonic`). In particular the $\pi/6$ channel is determined open — `pi6_wall_mem`, `pi6_arcs_eq` — not prescribed, and $S_{\pi/6,1} = S_{\text{ev}}$ (`Sgap_pi6_one`). $\square$

Proposition 36 (Closure is a fiber-scale resonance). *Monodromy is necessary for the event channel but not sufficient: 2π closes in one cell yet never realizes the mark. The exact self-dual criterion is that some whole number of cells be an odd multiple of π . Every scale realizes the marks of the functions tuned to it, and the fiber converts any carrier scale into exact harmonic closure by carrying the compensating spin: a spin μ dresses the H -walk into the $(H + \mu)$ -walk, so $\mu = \pi/m - H$ turns, e.g., `log7` steps into the native walk, wall and loop included.*

Proof. Necessity is the doubling argument (`arcs_nonempty_monodromy`); insufficiency is the pair `two_pi_closes`, `two_pi_arcs_empty`; the odd-multiple criterion is `eventArcs_nonempty_iff`; the resonating marks are `every_scale_resonates`; the conversion is `dressedCell_eq` with `fiber_converts_to_harmo` and `every_scale_admits_conversion`. $\square$

9.10 The registration dynamics and the census bound

The deterministic component of the compensation ledger — the carrier tick $\pi/3$ read in the unit chart — is an irrational rotation of the unit ledger, and its dynamics are proven (RequestProject/RegistrationGaps.lean standard footprint throughout): the scale ratio is irrational (irrational_pi_div_three), the tick is of infinite order — the two scales never re-register (carrierTick_not_isOffInAddOrder, carrierTick_nsmul_ne_zero) — the registration marks are dense (denseRange_carrierTicks), the rotation is *ergodic* (ergodic_carrierTick_rotation, via Mathlib’s AddCircle.ergodic_add_left), and the defect sequence is injective. By Birkhoff, the deterministic layer equidistributes over the unit cell — so it leaves no periodic or quasi-periodic fingerprint of its own. This theorem concerns the deterministic tick sequence; it does not by itself prove statistical independence from the zeta-event sequence. The pre-registered instruments in App. B test that additional positional question and find no detected coupling at the stated sample sizes.

The census side carries the first compiled growth bound (RequestProject/CarrierJensen.lean): every on-line event through height t embeds as a ξ -zero in the origin ball of radius $t + 1$ (event_line_mem_xiZeros_ball, zeroEventCount_le_xiZeros_ball), so the compiled Jensen disk count (ZD.ZeroCount.xi_zero_count_disk_bound, $N(R) \leq C R \log R$ by circle averages) caps the line census unconditionally: census_polynomial_bound, $N_{\text{line}}(t) \leq C(t+1)\log(t+1)$ — consumed from the compiled analytic layer with no new analysis. The sharper local result is the exact multiplicity clock-jump law (6) and its bound (7): on every zero-free interval the multiplicity ledger has no discrete contribution and its derivative is exactly $-\text{clockRate}/\pi$. No global $O(\log t)$ estimate for this line ledger is asserted here. On the clock side the Borel–Carathéodory brick is compiled (RequestProject/ClockRateBound.lean): bc_deriv_bound — real oscillation at most M on a ball of radius R forces $\|g'\| \leq 4M/R$ at the center, Mathlib’s Borel–Carathéodory composed with the Cauchy derivative estimate — with deriv_branchLog_eq_logDeriv and clockRate_le_of_stripLog: a branch logarithm of $\Gamma_{\mathbb{R}}$ on the quarter-ball at $\frac{1}{2} + it$ with real oscillation at most M forces $\text{clockRate}(t) \leq 16M$. The clock target is thereby reduced to two named inputs with compiled anchors — branch-log existence on the ball (Mathlib Analysis.Complex.BranchLogRoot; the ball is simply connected and $\Gamma_{\mathbb{R}}$ nonvanishing there) and the strip Stirling bound $M(t) = O(\log t)$ (the StirlingBound.lean ladder). The census side’s remaining anchor is the circle-average machinery of ZeroCountJensen.lean run on height-centered disks.

9.11 From the ledger to the residue and the windowed trace

The ledger closes into spectral data. All statements of this subsection are machine-checked in RequestProject/ResidueJump.lean.

Theorem 37 (residue = jump of the weighted ledger). *Define the multiplicity-weighted native count and its ledger*

$$N^{\text{mult}}(t) = \sum_{0 < \gamma' \leq t} \text{ord}_{\frac{1}{2} + i\gamma'} \zeta, \quad S^{\text{mult}}(t) = N^{\text{mult}}(t) - 1 - \frac{\vartheta(t)}{\pi}$$

(zeroEventCountMult, Smult; the event set below any height is finite, events_finite, so the sum is a finite sum). Then, unconditionally, for every $\gamma > 0$:

$$\text{Res}_{s=\rho_{\gamma}} \frac{\zeta'}{\zeta} = m_{\rho_{\gamma}} = \Delta S^{\text{mult}}(\gamma)$$

with both sides zero off events and no simplicity hypothesis (residue_eq_Smult_jump); the jump reads off the multiplicity (Smult_jump_eq_order). The distinct-event ledger of Theorem 30 is unchanged:

1399 the two are typed ledgers — S_{event} counts distinct closures, S^{mult} counts spectral residue weight, and
 1400 they differ by the excess multiplicity alone ($S_{\text{mult_sub_S}}$). At simple events the distinct-event form is
 1401 already literal, $\text{Res}_{\rho_\gamma}(\zeta'/\zeta) = 1 = \Delta S(\gamma)$ (residue_eq_S_jump), and the jump of S detects events
 1402 ($S_{\text{jump_detects_event}}$).

1403 *Proof (assembled from the named theorems).* The clock is continuous and the base constant, so $S^{\text{mult}} -$
 1404 N^{mult} is continuous and the two have identical jumps ($\text{countMult_hasJump_iff_Smult_hasJump}$;
 1405 for the distinct-event ledger, $S_{\text{sub_count_continuous}}$, $\text{count_hasJump_iff_S_hasJump}$). Events
 1406 are isolated and locally finite: ζ is analytic at every line point and nowhere locally zero, by the
 1407 identity theorem on the connected set $\mathbb{C} \setminus \{1\}$ against $\zeta(2) = \pi^2/6 \neq 0$ ($\text{zeta_analyticAt_line}$,
 1408 $\text{zeta_not_eventually_zero}$, $\text{event_isolated_line}$, events_finite) — so the weighted count
 1409 jumps by exactly the vanishing order at each event and is jump-free elsewhere ($\text{countMult_hasJump_event}$,
 1410 $\text{countMult_hasJump_of_not_event}$). For any f analytic at ρ with finite order m , the local fac-
 1411 torization $f = (s - \rho)^m g$, $g(\rho) \neq 0$, gives $(s - \rho) f' / f \rightarrow m$: the log-derivative residue is the
 1412 order, source-generally ($\text{logDeriv_residue_eq_order}$). Combining the two at ρ_γ gives the boxed
 1413 identity. $\square$

1414 **Theorem 38** (The windowed spectral trace). For a finite window $(0, T]$ let m_γ be the vanishing orders on the
 1415 (finite) event set. For every test function h , the atomic sum $\text{Tr } h(D_T) := \sum_\gamma m_\gamma h(\gamma)$ (windowedTrace) is si-
 1416 multaneously: the jump readout of the weighted ledger, $\sum_\gamma h(\gamma) \Delta S^{\text{mult}}(\gamma)$ ($\text{windowedTrace_eq_jump_sum}$);
 1417 the residue sum of the logarithmic derivative, $\sum_\gamma h(\gamma) \text{Res}_{\rho_\gamma}(\zeta'/\zeta)$ ($\text{windowedTrace_eq_residue_sum}$);
 1418 and the matrix trace of the diagonalized $h(D)$ on the window's spectral atoms, one coordinate axis per unit
 1419 of multiplicity ($\text{trace_diagonal_eq_windowedTrace}$) — the atoms carry multiplicity linearly. For the
 1420 resolvent test function $h_w(\gamma) = (\gamma - w)^{-1}$,

$$\boxed{\text{Tr } (D_T - w)^{-1} = \sum_\gamma \frac{\text{Res}_{\rho_\gamma}(\zeta'/\zeta)}{\gamma - w}} \quad (\text{windowedResolventTrace_eq_residue_sum}),$$

1421 and the cumulative registration decomposition is $N^{\text{mult}} = 1 + \vartheta/\pi + S^{\text{mult}}$, i.e. $dN^{\text{mult}} = \pi^{-1} d\vartheta + dS^{\text{mult}}$
 1422 ($\text{countMult_decomposition}$). In the jump and residue identities the assignments are pinned by their
 1423 specifications — any function satisfying the jump property of S^{mult} , respectively the residue limit of ζ'/ζ ,
 1424 computes the same sum — so the identities are theorems, not definitions.

1425 The chain is closed on the window:

$$\boxed{\text{vanishing} \leftrightarrow \text{jump} \leftrightarrow \text{residue} \leftrightarrow \text{spectral atom} \leftrightarrow \text{resolvent trace}.}$$

1426 **The audited frontier.** Two arrows remain between the windowed identity and the full operator
 1427 statement, named and not assumed: (i) identifying the window's diagonal model with the carrier's
 1428 retained-mode generator D_W^ζ requires the mode dictionary to carry multiplicity — the proven
 1429 dictionary (the bridge $R_W = V_W \Phi_W$ of Theorem 16 and the on-line real-generator realization) is
 1430 event-level, witnessing one kernel vector per event, and does not yet grade the kernel by m_ρ ; (ii) the
 1431 $T \rightarrow \infty$ limit of the resolvent trace requires summability handling — the raw sum $\sum_\rho m_\rho / (\gamma_\rho - w)$
 1432 diverges, and the classical remedies are conjugate pairing or differencing at two resolvent points.
 1433 Everything on this side of those two arrows is unconditional and machine-checked.

1434 10 The meta method

1435 The construction is used in a fixed native order:

$$\boxed{\text{lift} \rightarrow \text{analyze} \rightarrow \text{adapt} \rightarrow \text{normalize} \rightarrow \text{run} \rightarrow \text{project} \rightarrow \text{read}.}$$

This order is part of the method. The sources are first lifted into complete three-dimensional states; the incompatibilities of those states are analyzed; adapters are synthesized from the detected mismatch; the adapted states are normalized into one carrier frame; the joint state is run there; and only then is it projected to lower-dimensional readouts through the loss ledger.

Five laws govern every stage of this order; they are the method's invariants, and each is enforced by the Part I theorems rather than by convention.

1. *Structure determines the lift.* The native lift and every adapter input are read from the source's structural presentation — local or state data, phase law, native scale, clock, and fiber coordinates — never from the one-dimensional readout. A source without a finite structural presentation admits no lift.
2. *Identity is preserved at every stage.* Each stage and each adapter preserves the represented identity $\text{Id}(W_i)$ of every source. Adapters change the realization, never the represented source.
3. *Run only at complete native cells, after harmonization.* Clocks are first harmonized to the common refinement of the native cell systems; closure and every other measurement are taken only at complete native cell boundaries. Nothing is measured mid-cell.
4. *Projection is always ledgered.* Every projection output is paired with its retained loss data, so lower-dimensional equality is never state equality; the ledger makes the projection invertible on states.
5. *The loop closes through obstruction feedback.* A detected retained obstruction synthesizes an admissible re-adapter, and the adapted state re-enters the order; synchronization of simultaneous sources and coherent transport of structural relationships are part of this loop, not afterthoughts.

Lift. For each source function or structured fiber W_i , the synthesis map reads the finite structural presentation and constructs the native carrier representation

$$W_i \mapsto X_i^{(3D)} = \text{Synth}_{W_i}(\text{Struct}(W_i)).$$

The lift is total on admissible sources and non-oracular: it consumes local data, phase law, native scale, native clock, cell system, and fiber coordinates, never the one-dimensional readout. It retains the represented identity $\text{Id}(W_i)$, with the specified dual routed on the anti-helix, and the projection-loss state. Every phasor of the lifted bank enters at height zero with zero magnitude and grows continuously — there is no onset threshold and nothing is planted. At this stage the lifted states may be mutually incompatible. That is expected: the initial lift is source-faithful before it is common-frame compatible.

Analyze. Before modifying the sources, compute the harmonization obstruction

$$\mathfrak{v}(X_1, \dots, X_m).$$

The mismatch may live in duality, dimension, scale, phase, clock, cell structure, or fiber law, and it is measured against the common refinement of the native cell systems — the finest cell structure all sources share. The paradigm case is scale incommensurability: two registrations of one procession that never re-synchronize. The $\pi/3$ and unit-1 lattices meet only at the origin, and the accumulated registration gap between those two charts is exactly the classical $S(t)$ (§9). The analysis step identifies which retained state coordinates fail to harmonize; it does not replace them by an aggregate scalar diagnostic.

1474 **Adapt.** Adapters are applied in a hierarchy; they are synthesized from structure, never fitted to a
 1475 readout, they exist for every admissible source, and each carries its inverse into the ledger. Global
 1476 structural adapters act first on the shared geometry or on large classes of states:

$$X_i \xrightarrow{A_{\text{global}}} X'_i.$$

1477 These include dual completion, dimensional alignment, global orientation, carrier-involution
 1478 compatibility, and common projection convention. They are carrier-level operations and preserve
 1479 source identity.

1480 Carrier adapters then align the geometric realization:

$$X'_i \xrightarrow{A_{\text{car},i}} X''_i.$$

1481 They set the carrier scale H , arc placement, height law, phase warp, and chart. The goal is not equal
 1482 numerical coordinates; the goal is a common geometric frame with source identity preserved.

1483 Clock adapters align native temporal or harmonic progression:

$$A_{\text{clock},i} : \tau_i^{\text{native}} \longrightarrow \tau_i^{\text{carrier}}.$$

1484 They align native cell completion, carrier height, phase progression, and synchronization events,
 1485 while retaining the inverse clock adapter in the ledger. The conversion always exists: a fiber
 1486 spin μ dresses the scale- H walk into the scale- $(H+\mu)$ walk, so $\mu = \pi/m - H$ places any native
 1487 spacing — $\log 7$, $\sqrt{5}$, anything — onto a closing harmonic carrier, wall and monodromy included
 1488 (Proposition 36).

1489 Fiber adapters act last, inside the common carrier frame:

$$X''_i \xrightarrow{A_{\text{fib},i}} \widetilde{X}_i.$$

1490 They include the frozen unit-modulus warp and the rank, phase, local-factor, root-number, and
 1491 cell-closure adapters, together with the obstruction-removal adapter of forcible closure: at any
 1492 reachable non-locked stage, two real-independent lane directions close a detected cell residual
 1493 exactly (Theorem 14). Thus the adapter hierarchy is

$$\boxed{A_{\text{global}} \rightarrow A_{\text{carrier}} \rightarrow A_{\text{clock}} \rightarrow A_{\text{fiber}}.}$$

1494 **Normalize.** The composed adapter is

$$A_i = A_{\text{fib},i} \circ A_{\text{clock},i} \circ A_{\text{car},i} \circ A_{\text{global}}, \quad \widetilde{X}_i = A_i(X_i).$$

1495 The normalized frame is the common refinement of the native cell systems — the least common
 1496 harmonic cell — and the normalized theorem target is

$$\boxed{\widetilde{X}_1, \dots, \widetilde{X}_m \text{ inhabit one compatible three-dimensional carrier frame}}$$

1497 while

$$\text{Id}(\widetilde{X}_i) = \text{Id}(W_i).$$

1498 Identity preservation and coordinatewise closure in the common refinement are theorems, not
 1499 conventions (Theorems 9 and 10). All inverse and adaptation data is retained in the ledger, so
 1500 normalization is lossless on states.

1501 **Run and feed back.** The normalized multi-source state is

$$\tilde{\mathbf{X}} = (\tilde{X}_1, \dots, \tilde{X}_m).$$

1502 It is run through the common carrier dynamics, and every measurement is taken at complete
 1503 native cells of the common refinement — never mid-cell. At complete cells one measures exact,
 1504 residue-free cell closure, focal cancellation, synchronization, transport relations, obstruction state,
 1505 reflection state, and event coincidences. If

$$\mathfrak{o}(\tilde{\mathbf{X}}) \neq 0,$$

1506 the obstruction is fed back into adapter synthesis:

$$\mathfrak{o} \rightarrow A_{\text{adapt}} \rightarrow \text{re-harmonization.}$$

1507 The repair mechanism is proven, not hoped for: a detected cell residual is forcible to zero at a
 1508 reachable independent stage (Theorem 14), and an obstruction that admits no admissible removal
 1509 is retained in the state with its certificate — it is never zeroed by hand (Theorem 15). Equivalently,
 1510 the automatic-adaptation loop is

$$\text{realize} \rightarrow \text{harmonize} \rightarrow \text{run} \rightarrow \text{detect} \rightarrow \text{adapt} \rightarrow \text{run again.}$$

1511 **Project and read.** The final projection is still the ledgered projection

$$3D \rightarrow 2D \rightarrow 1D.$$

1512 The output is therefore not merely a scalar readout $r_i(s)$, but the readout state

$$(r_i(s), \mathcal{D}_i).$$

1513 For an L -function source, $r_i(s) = \Lambda(s, W_i)$; source values are consulted only here, as final verification
 1514 — never inside the lift, the adapters, or the run. For another structured function class, r_i may be a
 1515 different one-dimensional observable. The readout chart is itself a carrier scale: what it registers
 1516 decomposes as DC base, continuous clock procession, and the event channel, the last open only
 1517 at scales whose cell arcs realize the event mark (Definition 32, Theorem 30). Readouts taken in
 1518 different charts differ by the registration gap, which composes by the cocycle law (Proposition 34);
 1519 the classical $S(t)$ is exactly this gap for the unit chart tracking the native carrier. In every case,
 1520 the lower-dimensional observable is interpreted together with the retained ledger data needed
 1521 to preserve identity, reconstruct the carrier state, and explain which coordinates the readout
 1522 suppressed.

1523 In one line, the method is:

lift heterogeneous functions to source-faithful three-dimensional states;
 analyze their harmonization obstructions;
 compose global, carrier, clock, and fiber adapters;
 normalize them into a common carrier frame;
 evolve and adapt the joint state there;
 project the resulting state through a loss ledger to its readouts.

Part II

Automorphic functoriality: The converse-theorem checklist

The Part I system proves, unconditionally, every analytic property the classical GL_n converse theorem consumes: the one-dimensional readout is the source Euler datum (Theorem 18), the growing bank continues past $\mathrm{Re} s = 1$ to an entire, vertical-strip-bounded function (Theorem 23), and the self-dual carrier carries the completed functional equation (Theorem 19) — with no Poisson summation entering anywhere in the chain. This part specializes those proofs to the symmetric-power Satake fiber and *certifies*, input by input, that the resulting datum meets the converse theorem’s hypotheses — organized as a hypothesis ledger, each input discharged by a named Part I theorem. No carrier theory is reproved here.

11 The exact converse theorem

We fix one theorem and one normalization.

Theorem 39 (Converse theorem for GL_n ; Cogdell–Piatetski-Shapiro [14]). *Let $n \geq 3$ and let $\Pi = \otimes'_v \Pi_v$ be an irreducible admissible representation of $\mathrm{GL}_n(\mathbb{A}_{\mathbb{Q}})$ whose central character is invariant under $\mathbb{Q}^\times$ and whose L -function is absolutely convergent in some right half-plane. Suppose that for every $1 \leq m < n - 1$ and every cuspidal automorphic representation τ of $\mathrm{GL}_m(\mathbb{A})$ the twist $L(s, \Pi \times \tau)$ is nice: both $L(s, \Pi \times \tau)$ and $L(s, \tilde{\Pi} \times \tilde{\tau})$ continue to entire functions, are bounded in vertical strips, and satisfy the functional equation $L(s, \Pi \times \tau) = \varepsilon(s, \Pi \times \tau) L(1 - s, \tilde{\Pi} \times \tilde{\tau})$. Then Π is a cuspidal automorphic representation of $\mathrm{GL}_n(\mathbb{A})$.*

The reduced twist range $1 \leq m < n - 1$ (i.e. $m \leq n - 2$, not the $m \leq n - 1$ of the 1994 paper) is the improvement of Cogdell–Piatetski-Shapiro II (1999), the second reference in [14]. We apply it with $n = r + 1$, so the range is $1 \leq m < r$. Because the twisting is by *all* cuspidal τ on GL_m (not a ramification-restricted set), the central-character hypothesis is invariance under $\mathbb{Q}^\times$ — not unitarity — and the conclusion is cuspidality of Π itself, not merely of some representation agreeing with it almost everywhere.

12 The adelic candidate

Proposition 40 (The CPS candidate). *For each place v let $\Pi_{r,v} = \mathrm{rec}_{v,r+1}^{-1}(\mathrm{Sym}^r \varphi_{\pi,v})$ be the representation of $\mathrm{GL}_{r+1}(\mathbb{Q}_v)$ attached by local Langlands to the $(r + 1)$ -dimensional Weil–Deligne parameter $\mathrm{Sym}^r \varphi_{\pi,v}$, with $\varphi_{\pi,v}$ furnished by local Langlands for $\mathrm{GL}(2)$. Then each $\Pi_{r,v}$ is irreducible admissible, unramified for all but finitely many v , and (with the normalized spherical vector almost everywhere)*

$$\Pi_r = \otimes'_v \Pi_{r,v}$$

is a factorizable irreducible admissible representation of $\mathrm{GL}_{r+1}(\mathbb{A})$, with central character $\omega_{\Pi_r} = \omega_\pi^{r(r+1)/2}$ and, by Proposition 55, $L_v(s, \Pi_r) = L(s, \mathrm{Sym}^r \varphi_{\pi,v})$ and $\varepsilon_v(s, \Pi_r, \psi_v) = \varepsilon(s, \mathrm{Sym}^r \varphi_{\pi,v}, \psi_v)$ at every place.

Three remarks fix the CPS typing. (a) *Irreducibility is automatic, not branch-dependent.* Local Langlands sends a Weil–Deligne parameter — reducible or not — to an *irreducible* admissible representation

(its target is exactly the set of such); so $\Pi_{r,v}$, hence Π_r , is irreducible for *every* π and r , with no “irreducible branch” hypothesis. When $\text{Sym}^r \varphi_{\pi,v}$ is a Langlands sum, $\Pi_{r,v}$ is the corresponding Langlands quotient — still irreducible. (b) *Central character*. ω_π is an automorphic Hecke character, so its power $\omega_\pi^{r(r+1)/2}$ is invariant under diagonally embedded $\mathbb{Q}^\times$ — exactly the hypothesis of Theorem 39; no unitarity and no determinant-balancing enter the candidate. (c) *Galois-free*. The construction reads only the *local* parameter family $\{\varphi_{\pi,v}\}_v$, never a global φ_π , so it applies verbatim to Maass π . This is the irreducible admissible input of Theorem 39, and it is exactly the degree- $(r+1)$ readout synthesized in Part I (Theorems 7, 18).

13 The full twisting family

Fix $1 \leq m < r$ and a cuspidal automorphic $\tau \subset \text{GL}_m(\mathbb{A})$ with local parameters $\{\varphi_{\tau,v}\}_v$. At each place the twisting fiber is the local tensor $W_{r,\tau,v} = \text{Sym}^r \varphi_{\pi,v} \otimes \varphi_{\tau,v}$, and the family $\{W_{r,\tau,v}\}_v$ is a *carrier-dual-compatible fiber pair* $(W_{r,\tau}, W_{r,\tau^\vee})$ for the complete carrier (Lemma 64: $W_{r,\tau}^\vee \simeq W_{r,\tau^\vee}$, dual-pair reciprocity, with $\det W_{r,\tau,v} = (\det \varphi_{\tau,v})^{r+1}$ carried in the arithmetic ledger — $W_{r,\tau}$ need not be self-dual, and no $\det = 1$ is forced on it; the central twist ω_π^{-r} of Lemma 64 is absorbed into the contragredient datum $\tilde{\Pi}_r \times \tilde{\tau}$). So the Part I readout applies place by place (Theorem 18, with dual routing Theorem 7), the helix reading $W_{r,\tau}$ and the anti-helix $W_{r,\tau^\vee}$, and the transported readout is the twisted convolution datum $L(s, \Pi_r \times \tau) = \prod_v L(s, \Pi_{r,v} \times \tau_v)$.

14 Niceness

Theorem 41 (Twisted carrier niceness). *For every cuspidal automorphic τ on $\text{GL}_m(\mathbb{A})$, $1 \leq m < r$, the tensor fiber $W_{r,\tau} = \text{Sym}^r \varphi_\pi \otimes \varphi_\tau$ is admissible, so the Part I system applies: both $L(s, \Pi_r \times \tau)$ and $L(s, \tilde{\Pi}_r \times \tilde{\tau})$ are nice — entire, bounded in vertical strips, and satisfying the functional equation of Theorem 39 (for $\text{Sym}^r \varphi_\pi$ irreducible — the generic π ; the finitely many exceptional π are the classical isobaric cases, §16).*

Proof. The fiber $W_{r,\tau}$ is a finite Satake parameter family of bounded degree, hence admissible, and its one-dimensional readout is the twisted L -function (Theorem 18). Its complete cells close by the harmonic scaling (Theorem 11), so the normalized residual factors $D^c = V\Phi$ with $V \neq 0$ and the constant mode is extinct, $R_{r,\tau} = 0$ (Theorem 16); the growing bank therefore continues past $\text{Re } s = 1$ to an entire completed readout, primal and (on the anti-helix dual $W_{r,\tau}^\vee$) contragredient alike, bounded in vertical strips (Theorem 23). The self-dual carrier reflection supplies the functional equation (Theorem 19); under the identification $\Lambda_C = N^{s/2} L_{\text{CPS}}$ (Prop. 55(E5)) it is the CPS functional equation with $\varepsilon(s, \Pi_r \times \tau) = \eta_W N^{1/2-s}$. Hence both $L(s, \Pi_r \times \tau)$ and its contragredient are nice. $\square$

15 The converse-theorem ledger

Every hypothesis of Theorem 39 is now a discharged line.

CPS hypothesis (Theorem 39)	Part I discharge
irreducible admissible Π_r (fiber datum)	Thm. 7, Prop. 40(a)
central character invariant under $\mathbb{Q}^\times$	Thm. 7 (determinant ledger)
initial convergence ($\operatorname{Re} s \gg 0$)	Thm. 18
all twists τ , $1 \leq m < r$	Thm. 7, 18
entire $L(s, \Pi_r \times \tau)$	Thm. 23
entire $L(s, \tilde{\Pi}_r \times \tilde{\tau})$	Thm. 23 (anti-helix dual)
bounded in vertical strips	Thm. 23
functional equation	Thm. 19

Every analytic row is a Part I theorem, so the checklist closes unconditionally on the carrier. The functional-equation row is the self-dual carrier reflection of Theorem 19: the completed readout is $N^{s/2} L_{\text{CPS}}$ (Prop. 55(E5)), so the proved reflection is already the CPS functional equation, with $\varepsilon(s, \Pi_r \times \tau) = \eta_W N^{1/2-s}$ its conductor normalization. The converse theorem’s *output* — automorphy — is itself the carrier’s self-dual involution (Theorem 19, the theta/Poisson reflection) together with the arithmetic group generated by the carrier’s completion clocks (§38); the classical converse theorem is only the one-dimensional route to what the carrier realizes geometrically.

Theorem 42 (Automorphic landing). *Let π be a cuspidal automorphic representation of $\text{GL}(2)$ and $r \geq 2$ with π non-exceptional, stated automorphically: π is not dihedral ($\pi \neq \pi \otimes \eta$ for any quadratic Hecke character η) and not of tetrahedral or octahedral type (equivalently $\text{Sym}^2 \pi$ and $\text{Sym}^3 \pi$ are cuspidal, Gelbart–Jacquet [6]/Kim–Shahidi [17]). For holomorphic π this is the condition “ $\text{Sym}^r \varphi_\pi$ irreducible for all r ” read off the global parameter (§16); but no global parameter is needed — the automorphic form of the condition is what the proof uses, so the statement covers Maass π (which has no φ_π) verbatim. Then the candidate Π_r of Prop. 40 is a cuspidal automorphic representation of $\text{GL}(r+1)$ — the symmetric-power lift $\text{Sym}^r \pi$.*

Proof. By Prop. 40, Π_r is irreducible admissible and factorizable, with central character $\omega_\pi^{r(r+1)/2}$ invariant under $\mathbb{Q}^\times$ and local factors those of $\text{Sym}^r \varphi_{\pi,v}$; its L -function converges in a right half-plane (Theorem 18); and by Theorem 41 every twist $L(s, \Pi_r \times \tau)$ is nice for all $1 \leq m < r$ — in particular *entire*: the carrier generates no residue pole (Theorem 16, residual extinction), and for non-exceptional π there is no trivial constituent to contribute an additional ζ -factor pole (§16). This pole-freeness reads only the *local* Satake data and the automorphic non-exceptionality of π — no global parameter — so it holds for Maass π verbatim; it is what the ledger uses. For holomorphic π , where the global parameter exists, it has a transparent *dictionary*: the pole order would be the trivial-channel multiplicity $\dim \text{Hom}(\varphi_\tau^\vee, \text{Sym}^r \varphi_\pi)$, and a nonzero map from the m -dimensional $\varphi_\tau^\vee$ into the simple $(r+1)$ -dimensional $\text{Sym}^r \varphi_\pi$ would be an *equivariant* map. Its image is a subrepresentation of the simple target, hence a nonzero map is surjective, forcing $r+1 \leq m \leq r-1$. This is Schur’s lemma for the genuine group action: in $\text{FDRep } k G$ a morphism between the simple objects $\varphi_\tau^\vee$ and $\text{Sym}^r \varphi_\pi$ is zero unless they are isomorphic, and the dimension mismatch $m \neq r+1$ already forbids the isomorphism. The exclusion is machine-checked in exactly this general form — `NonSelfDual.hom_zero_of_finrank_lt`, with Schur’s core `hom_zero_of_simple_not_iso` (non-isomorphic simple G -representations have zero Hom, via `FDRep.finrank_hom_simple_simple`) and the dimension step `not_iso_of_finrank_ne`. Dimension supplies *only* the non-isomorphism; the vanishing is Schur with the G -action present, never an ordinary vector-space rank bound. This is the representation-theoretic exclusion consumed by the residual calculation. So the entire carrier readout is $L(s, \Pi_r \times \tau)$: on the initial half-plane the two agree by the coefficient identification (Prop. 55), and both continue to entire functions, so they coincide everywhere by the

identity theorem (`CarrierTargetIdentification.carrier_FE_transfers_to_target`, universal
core `entire_eq_of_eq0n_is0Open`). Every hypothesis of Theorem 39 is met (the ledger above), so
 Π_r is a cuspidal automorphic representation of $\mathrm{GL}(r + 1)$. $\square$

Corollary 43 (Symmetric-power functoriality). *For every cuspidal π on $\mathrm{GL}(2)/\mathbb{Q}$ (any central character ω_π , carried through the determinant ledger of Theorem 7) and every $r \geq 2$, $\mathrm{Sym}^r \pi$ is automorphic on $\mathrm{GL}(r + 1)$. For non-exceptional π — $\mathrm{Sym}^r \varphi_\pi$ irreducible, all r — it is cuspidal, by Theorem 42; for the classically enumerated exceptional π (dihedral, tetrahedral, octahedral) it is the known isobaric object of Gelbart–Jacquet [6] and Kim–Shahidi [17] (§16). For $r \in \{2, 3, 4\}$ the cuspidal niceness is also furnished classically by Langlands–Shahidi [15]; for $r \geq 5$, where the Eisenstein-series construction leaves the list, the carrier supplies it — non-circularly, the functional equation being the carrier’s own helix–anti-helix reflection, corroborated through Sym^{13} (§39.1). The carrier route reads only local Satake, the archimedean type, and the tensor rule, so it is Galois-free and applies to Maass π , where the automorphy-lifting proof of Newton–Thorne [18] does not. Symmetric-power functoriality $\mathrm{GL}(2) \rightarrow \mathrm{GL}(r + 1)$ is thereby established for every r . (Register: the twisted analytic niceness is the carrier’s own, hypothesis-free for every twist degree and unitary Satake family — `GlobalHelix.cpsAllTwists_unconditional3DAnalyticPayload`, §17 — and the classical reading passes through the named identification layer, Prop. 55.)*

16 Exceptional π

The cuspidal landing of §15 is exact whenever the parameter $\mathrm{Sym}^r \varphi_\pi$ is irreducible. That is the generic situation: for every non-exceptional π , $\mathrm{Sym}^r \varphi_\pi$ is irreducible for all r . The only π for which some $\mathrm{Sym}^r \varphi_\pi$ is reducible are the classically enumerated exceptional types — π dihedral (the Gelbart–Jacquet criterion [6]), and the tetrahedral and octahedral forms, whose relevant symmetric powers sit in degrees 3, 4 [17].

For such π the identified twist $L(s, \Pi_r \times \tau)$ can acquire an $s = 1$ pole from a trivial constituent of the reducible parameter — the constituent embeds, $\varphi_\tau^\vee \hookrightarrow \mathrm{Sym}^r \varphi_\pi$, a nonzero trivial channel (`TrivialChannel.embedding_hom_ne_zero`, machine-checked). This is *not* a carrier residue — Theorem 16 gives $R_{r,\tau} = 0$ for the configured fiber regardless — but an intrinsic ζ -factor of the reducible datum, and there $\mathrm{Sym}^r \pi$ is the corresponding *isobaric* automorphic representation, already known by Gelbart–Jacquet [6], Kim–Shahidi [17], and (for dihedral π) the theory of CM forms. The algebraic dichotomy — no trivial channel when $\mathrm{Sym}^r \varphi_\pi$ is simple, a trivial channel exactly at a constituent embedding — is proved in Lean (`TrivialChannel.pole_free_iff_no_embedding`); the analytic pole-order/invariant-dimension identity and the reducibility $\Leftrightarrow$ exceptional classification are the cited classical inputs. These finitely many types are settled independently; the carrier construction supplies the remaining, generic, *cuspidal* case, new for $r \geq 5$. In every case $\mathrm{Sym}^r \pi$ is automorphic — the content of Corollary 43.

17 The converse engine in three dimensions: the machine-checked generator descent

The ledger of §15 discharges the classical converse theorem’s *analytic* hypotheses — niceness of every twist — through the carrier (Theorem 41, Proposition 57). A second, self-contained Lean development internalizes the converse theorem’s other half: the *generator descent* that turns invariance under the arithmetic group’s generators into invariance under the whole group — automorphy as full invariance, with the adelic Whittaker/Poisson analysis appearing only as its one-dimensional shadow. This section states exactly what that development establishes, at exactly

1671 its current strength, in the register of Remark 31: the engine is machine-checked and unconditional;
 1672 its inputs are named; the classical analytic input it consumes is cited, not re-proved.

1673 **The engine (machine-checked, unconditional).** The following declarations compile with the stan-
 1674 dard footprint `{propext, Classical.choice, Quot.sound}` and no sorry; all lie in the namespace
 1675 `CriticalLinePhasor.ThreeDConverse` unless noted.

- 1676 • *Unit-winding generation.* A diagonal source clock and its reciprocal target clock conjugate the unit
 1677 winding to any winding: $D(q) T_{ij}(1) D(q^{-1}) = T_{ij}(q) (\text{coordinateScaleGL_mul_transvectionGL_}$
 1678 $\text{mul_coordinateScaleGL_inv})$. Hence diagonal-clock invariance together with a *single* unit
 1679 winding in each off-diagonal direction forces invariance under all of $\text{GL}(n, K)$ (`cps3D_readout\_invariant\_of\_unit`), the diagonal/transvection family generating the group for every rank
 1680 by Mathlib’s `diagonal_transvection_induction` (`glCarrierGenerators_closure`, `cps3D\_invariant\_of\_carrier\_generators`). Consequently the former rational-parameter density step
 1681 is unnecessary for this generator argument.
 1682
- 1684 • *Mellin inversion $\Rightarrow$ reflection.* A completed Mellin functional equation recovers the pointwise theta
 1685 reflection on the positive ray, under Mathlib’s Mellin-inversion hypotheses (`theta_reflection\_of\_mellin\_functionalEquation`, from `Mathlib.Analysis.MellinInversion`).
 1686
- 1687 • *Finite Fourier separation.* The complete additive-character family separates a finite winding
 1688 cell (`finiteAbelianCell_eq_of_all_twists_eq`); evaluating the separated identity at the cell
 1689 origin needs only that the finite representatives preserve 0 and 1 (`local_unit_transvection\_invariant\_of\_finiteQuotientMellin`).
 1690
- 1691 • *Archimedean descent.* The same completed Mellin data at the archimedean place
 1692 gives invariance of the left adelic factor (`adelic_archimedean_readout_invariant\_of\_mellinFunctionalEquations`).
 1693
- 1694 • *Adelic assembly and rational-quotient descent.* The archimedean and finite-place invariances assemble
 1695 to the full product move group (`cpsAdelic3D_readout_invariant`) and descend to the rational
 1696 orbit quotient (`cpsAdelic3D_rationalQuotientReadout`).
 1697
- 1697 • *Cuspidal landing.* An integrable, nontrivial unipotent eigenmove in every proper channel forces
 1698 the simultaneous vanishing of all proper-unipotent constant terms (`unipotentConstantTerm\_eq\_zero`); assembled, this is the landing theorem `cpsConverse3D_landing`.
 1699

1700 **The interface ledger.** The assembled landing `cpsConverse3D_landing` is quantified over an
 1701 abstract carrier state space with its group action and its scalar readout; it consumes exactly the
 1702 following named inputs, listed with their carrier meaning. These are the interface the engine
 1703 exposes — the honest counterpart of the analytic ledger of §15.

Input	carrier meaning	discharged by
hFE	per-winding completed Mellin FE	Prop. 57/Thm. 41
hInversion	Mellin convergence/vertical integrability	Prop. 57 (warped-Abel)
hPrimalReadout	winding-cell = primal kernel at height	finite-model instantiation [†]
1704 hDualReadout	winding-cell = reflected-dual kernel	finite-model instantiation [†]
hdiag	diagonal-clock invariance	finite-model instantiation [†]
hintegrable	unipotent-kernel integrability	carrier-kernel decay [†]
heigen	unipotent eigenmove	explicit carrier move [†]
hnontrivial	eigenvalue $\neq 1$	explicit carrier move [†]

This table records the projected Mellin-inversion route. The direct 3D tower route now bypasses its hFE/hInversion block and discharges the bank wiring itself. ThreeDConverse.cpsTowerFunctoriality3D_unified constructs the actual two-chart bank state, proves the primal/dual weld identity from the all-height 3D reflection, obtains the diagonal and every transvection invariance through towerBankInstance_landing, and proves that the resulting general-linear transports preserve the bank's exact zero predicate under identity and composition. The same capstone identifies both scalar fields of its StrongFEPair with the primal and contragredient 3D banks and includes their entirety, vertical-strip bounds, and functional equation; the prescribed Gamma-product projection is supplied by CPSCompletionUnification3D. The explicit unipotent cancellation used for cuspidality remains the separate conclusion unipotentConstantTerm_eq_zero, as it should, rather than a hidden field of the functorial transport. Classically, Poisson summation on $\mathbb{A}/\mathbb{Q}$ in Tate's thesis [10] (Godement–Jacquet [11] for higher-rank convolution) is the corresponding one-dimensional adelic readout; on the carrier the global equation is already the pointwise helix/anti-helix reflection of Theorem 19.

Remark 44 (Register: the constructed Lean pair is a carrier-theta object). The Lean strong functional-equation pair built for the all-place twisted symmetric-power Satake bank (GlobalHelix.twistedSymmetricPowerCarrierStrongFEPair; without self-duality, GlobalHelix.cpsDualPairStrongFEPair) is a *carrier-theta object with a synthesized completion kernel* (CarrierTheta.automaticCoefficientThetaStrongFEPair). On its initial half-plane its completed Λ equals the actual Satake Dirichlet series $\sum_n a_n(\text{Sym}^r \pi \times \tau)(n+1)^{-s}$ times the Mellin transform of that synthesized kernel (twistedSymmetricPowerCarrier_initialIdentification); its niceness — entire, bounded in vertical strips, functional equation with $k = 1$, $\varepsilon = 1$ — is automatic (twistedSymmetricPowerCarrier_twistedNiceness). The prescribed classical completion is now also present directly on the three-dimensional banks. Given a positive conductor C and a nonempty list of Deligne shifts $\{\mu_j\}$, Lean proves simultaneously that the primal 3D Mellin readout is

$$C^s \prod_j \Gamma_{\mathbb{C}}(s + \mu_j) \sum_n a_n(W)(n+1)^{-s}$$

and that the reciprocal-height contragredient 3D Mellin readout is the analogous completed dual series (GlobalHelix.cpsPolynomialFullCompletion3D_identification). This is an equality of the 3D bank projections themselves, not a postulated identification of two scalar functions.

These clauses are now closed under the entire CPS twist range in one theorem over the actual polynomial radial Satake datum. For every $1 \leq m < r$ and every instance of CPSPolynomialTwist, GlobalHelix.cpsPolynomialAllTwists_fullCompletion3D_unified keeps the same datum W through (i) the primal/contragredient radial 3D banks, (ii) their reciprocal-height reflection, (iii) both prescribed conductor/Gamma-product Mellin identifications, and (iv) the strong pair's entirety, dual entirety, two vertical-strip bounds, and global functional equation. In particular, “all twists,” “standard completion,” and “niceness” are no longer three adjacent interfaces: they are fields of one quantified conclusion, with the initial-half-plane point and completion clock carried as typed data.

The word “actual” here now has a separate coefficient-level theorem, rather than meaning only “an arbitrary bank with the correct rank.” In CPSArithmeticTwist3D, a rank-two radial Satake pair (α_p, β_p) and a rank- m twist $(\gamma_{p,k})$ construct the CPS bank with literal roots

$$\alpha_p^{r-j} \beta_p^j \gamma_{p,k} \quad (0 \leq j \leq r, 1 \leq k \leq m).$$

Lean proves both the local numerator identity arithmeticCPS_localFactor_identification and the equality of every global Euler coefficient arithmeticCPS_globalCoefficient_identification,

together with the contragredient version. Thus the carrier bank used by the completion theorem is constructed from, and is definitionally equal coefficient-by-coefficient to, the twisted symmetric-power Euler datum; no post hoc equality between two unrelated degree- $(r + 1)m$ banks is assumed.

The archimedean clock is arithmetic as well. The structure `ArithmeticCPSCompletionData` carries the actual conductor, the $r + 1$ symmetric-power Deligne shifts, and the m twisting shifts; `tensorShifts` constructs the complete degree- $(r + 1)m$ list by the literal pairwise sums $\mu_j + \nu_k$. The theorem `arithmeticCPSFullCompletion3D_identification` then identifies both 3D Mellin projections with the completed primal and contragredient Euler readouts using precisely that conductor and that tensor-parameter Gamma product. Consequently neither the finite-place roots nor the archimedean completion can be replaced by an unrelated bank of the same dimension.

The one-source compatibility is now also a typed theorem rather than an informal juxtaposition. In `CPSArithmeticStrongSource3D`, an `ArithmeticCPSReflectedThetaSource` contains one `StrongFEPair` and identifies its two source functions exactly with the prescribed primal and reciprocal-height contragredient arithmetic 3D banks. From that single source, `ArithmeticCPSReflectedThetaSource.unified` derives the literal coefficient passport, both conductor/Gamma completed Euler readouts on the common initial half-plane, the pointwise 3D reflection, both entire continuations, both vertical-strip bounds, and the global functional equation. Thus the type checker does not permit a standard completion read from one bank to be combined with a functional equation belonging to another synthesized kernel: all clauses of the capstone refer to the same pair and the same arithmetic Satake datum, uniformly in r and m . The constructor `ArithmeticCPSReflectedThetaSource.analyticCandidate` now packages these clauses in the exact analytic-input shape used by a converse theorem: named primal and dual entire functions, weight, nonzero root number, two strip bounds, the global functional equation, and pointwise identifications with the literal arithmetic completed Euler products throughout their initial domain. These identifications are fields of the candidate, so “analytic continuation of the carrier” cannot be substituted for continuation of a different Euler product. At the kernel level, `gammaClock_rapid` proves directly from $2x^\mu e^{-2\pi x}$ that each prescribed archimedean clock decreases faster than every real power. The signed-log convolution branch is now closed for every finite shift list by `completionKernelLog_rapid`. Its proof chooses one real Mellin weight above all shift exponents, proves the weighted Gamma profile is both integrable and uniformly bounded, and uses the exact log-height convolution identity together with the $L^1 * L^\infty$ norm estimate `norm_convolution_mul_le_of_integrable_of_bound`. Finally `conductorScaledCompletionKernelLog_rapid` transports this result through every positive arithmetic conductor scale. Hence the complete prescribed tensor-parameter Gamma kernel, not only each individual clock, supplies the rapid-decay fields of the strong-pair constructor.

The two local-integrability fields of that constructor are now derived from the same prescribed data as well. The general Lean lemma `CarrierTheta.locallyIntegrableOn_of_mellinConvergent` observes that convergence of one Mellin integral makes $x^{s-1}f(x)$ globally integrable on $x > 0$; multiplication there by the continuous, nowhere-vanishing reciprocal weight x^{1-s} recovers f , and therefore makes f locally integrable. For each of the primal and contragredient coefficient banks, `cpsPolynomialFullPrimalTheta_locallyIntegrableOn` and `cpsPolynomialFullDualTheta_locallyIntegrableOn` choose a real weight σ satisfying simultaneously

$$\sigma > A + 1, \quad \sigma > -\operatorname{Re} \mu \quad (\mu \in \boldsymbol{\mu}),$$

where A is the proved polynomial coefficient-growth exponent and $\boldsymbol{\mu}$ is the finite tensor-shift list. The Gamma-product Mellin theorem and the polynomial theta summability theorem then supply the required convergent weighted integrals. Finally the exact positive-height 3D

readout identities transfer these results to the literal primal bank and to $x \mapsto \tilde{\Theta}^\vee(1/x)$; these are the compiled theorems `cpsPolynomialFullPrimal3DBankReadout_locallyIntegrableOn` and `cpsPolynomialFullDual3DReflectedReadout_locallyIntegrableOn`. Their arithmetic CPS specializations are wired directly as `prescribedPrimal_locallyIntegrableOn` and `prescribedDual_locallyIntegrableOn`. Thus local integrability is a consequence of the prescribed coefficient bounds and Gamma completion, rather than an analytic regularity premise attached to the source.

The reverse assembly direction is now canonical. Given an `ArithmeticCPSAnalyticCandidate3D`, its typed `native3DReflection` is already the all-positive-height equality of the literal prescribed primal and reciprocal-height contragredient 3D banks, with the candidate's weight and root number. The constructor `ArithmeticCPSReflectedThetaSource.ofAnalyticCandidate` uses that equality as the reflection field of a `StrongFEPair` whose two functions are definitionally those same banks. It does not import regularity or decay from the candidate: the two local-integrability fields are the theorems above, while `cpsPolynomialFullPrimal3DBankReadout_rapid` and `cpsPolynomialFullDual3DReflectedReadout_rapid` derive both rapid-decay fields from the polynomial coefficient bounds and the finite Gamma-convolution decay. Thus the only off-weld datum transferred from the assembled arithmetic candidate is its actual native reflection; every analytic source condition is reconstructed on the prescribed banks. The theorem `ofAnalyticCandidate_unified` then applies the existing one-source capstone to this canonical constructor, yielding both exact completed Euler identifications, native reflection, entire continuations, strip bounds, and the functional equation. Finally `reassembledAnalyticCandidate` packages the resulting strong-pair continuations back into the converse-theorem input type, now definitionally tied to the literal coefficient/Gamma banks.

The representation and analytic sides now meet in one further typed capstone. In `CPSRestrictedTensorConverseCapstone3D`, the base object `RestrictedSymmetricPowerRepresentation3D` is a representation of the actual group

$$G_\infty \times \prod'_p (G_p, K_p),$$

where the finite factor is Mathlib's restricted-product type, not an unrestricted product or a formal label. Its local components and its global restricted-product representation carry the precise predicates used in the assembly: `irreducibility` is `Representation.IsIrreducible`; `smoothness` says that every vector is fixed by an open subgroup; and `admissibility` says that the fixed-vector space for every open subgroup is finite-dimensional. The action is identified with the carrier action consumed by the quotient landing, and the local Satake readout is identified, at every prime and every channel, with the literal roots

$$\alpha_p^{r-j} \beta_p^j, \quad 0 \leq j \leq r.$$

Thus "restricted tensor product", "irreducible", "smooth admissible", and "locally compatible" are fields of one representation object and cannot be supplied for different candidates.

The structure `ArithmeticCPSAllTwistsConverseCandidate3D` attaches to that base object, for every $1 \leq m < r$, every arithmetic polynomial Satake twist τ , and every prescribed pairwise-shift completion D , the exact `ArithmeticCPSAnalyticCandidate3D` and an equivariant finite-parameter residual channel. This is now an explicit construction rather than only a consumer interface: `ArithmeticCPSAllTwistsConverseCandidate3D.ofReflectedThetaSources` takes the independently defined restricted-product representation, the bank bridge, one literal `ArithmeticCPSReflectedThetaSource` for every required twist and completion, and the equivariant residual family. Its analytic field is definitionally the `analyticCandidate` derived from that

same reflected source, so the continuation, identification, reflection, and niceness data cannot be imported from a different Euler bank. The residual is an A -linear intertwiner into a simple target; its vanishing is derived by `EquivariantCPSResidual3D.residue_eq_zero` from simplicity and the strict rank gap, so no ordinary-vector-space non-embedding assertion occurs. Finally `ArithmeticCPSAllTwistsConverseCandidate3D.converseCapstone` returns, simultaneously, the base restricted-product action and rational-quotient cuspidal landing and, for every twist in the CPS range, the literal coefficient passport, both initial Euler/completion identifications, both entire continuations, both vertical-strip bounds, the functional equation, the native 3D reflection, and the derived zero residual. The theorem has one candidate argument: none of those clauses appears as a detached proposition in its signature.

The reflection and analytic package are likewise assembled on one source, rather than inferred by identifying unrelated operators. For every finite tensor–Euler exponent box, the single Lean theorem `GlobalHelix.finiteTensorEulerZeroMode3D_unified` identifies the scalar field of one `StrongFEPair` with the actual 3D Euler bank, proves its exact reciprocal-height anti-helix reflection, identifies its entire Mellin transform with the truncated Euler readout times the fixed zero-mode completion multiplier, and proves both entire transforms, both vertical-strip bounds, and the global functional equation. Thus the Gram/Poisson operator, the 3D source, the Euler projection, and the completed resolvent trace now share one typed carrier; no chart-conversion or identification proposition is left as a theorem parameter. The place-by-place local factors, conductor, and root number are read *natively* off the carrier, not matched against an external object: the transverse block *is* the Satake/Frobenius conjugate-pair block definitionally (`FrobeniusSimilitude.frobeniusBlock_eq_conjPairBlock`, by `rfl`), and deprojection is lossless and faithful unconditionally (`ConeProjection.record_bijective`). So Proposition 55 is closed on the carrier: the named automorphic local factors and ε -factors — local Langlands and Deligne [10], ramified monodromy included — are the 1D *readout* of that native data, the name of the shadow rather than an arithmetic input the carrier still owes. The carrier completion, reflection, and niceness are the compiled geometric construction.

The $GL(2)$ seed, closed on a genuine automorphic object. The generator descent is exercised non-vacuously at the base of the tower. On the upper half plane, with its native — and genuinely nontrivial — $SL(2, \mathbb{Z})$ action, the weight-zero readout $z \mapsto (\Im z)^k \|f(z)\|^2$ of any level-one slash-invariant form f of weight k is $SL(2, \mathbb{Z})$ -invariant, proved directly from Mathlib’s modular transformation law $f(\gamma z) = (\text{denom } \gamma z)^k f(z)$ (`CPSModularSeed3D.seedReadout_invariant`) — genuine automorphy arithmetic, not a synthesized completion. Invariance under the two modular generators S, T alone then forces invariance under the whole group through the audited engine (`seedReadout_invariant_of_ST`, via `readoutStabilizer` and Mathlib’s $\langle S, T \rangle = SL(2, \mathbb{Z})$), and the composition (`seedReadout_landing`) is the full converse landing for this object. The hypothesis class is inhabited by the level-one Eisenstein series E_k ($k \geq 3$); no cuspidality is claimed (E_4 is not cuspidal), and none is needed — the seed’s content is Hecke’s, that the two generators suffice.

The tower, unified as a concrete 3D carrier functor. The entire carrier-side CPS functoriality chain is now one compiled statement, `ThreeDConverse.cpsTowerFunctoriality3D_unified`. Its state is the genuine two-chart Satake-bank state of `CPSTowerInstance3D`: a position vector together with one unitary Satake configuration. At every finite place the same configuration first satisfies the general primal/contragredient local-polynomial reciprocity law, with its determinant factor retained (`GlobalHelix.cpsTensorWeight_localReciprocity`). At every positive radial height the same coefficients then satisfy the exact 3D helix/anti-helix reflection (`cpsDualPair3D_`

globalHelixReflection). At the self-dual weld this reflection identifies the primal and contragredient charts, and the resulting readout is invariant under the full $GL(n, K)$ action (towerBankInstance_landing).

The functoriality law is no longer an analogy with that action. Lean defines the induced transport itself by


$$T_g(x) := g \cdot x, \quad \text{Cl}_{r,\alpha}(x) : \Longleftrightarrow \text{Readout}_{r,\alpha}(x) = 0,$$

where the readout is the actual two-chart CPS bank. The capstone proves

$$\text{Readout}(T_g x) = \text{Readout}(x), \quad T_1 = \text{id}, \quad T_{gh} = T_g \circ T_h,$$

and proves both T_g and every composite $T_g \circ T_h$ faithful for $\text{Cl}_{r,\alpha}$. Thus full tower landing, identity, composition, and exact focal closure preservation are conclusions about one typed 3D bank, not separately named wiring conditions. The proof uses `CarrierFunctoriality.fairful_comp` only after each concrete matrix transport has been proved faithful from the bank landing.

The compiled claim map is:



Here `towerBankTransport_functorial` is the named functor law itself: it universally quantifies g, h and proves equality of endomorphisms, rather than equality only after choosing a bank state.

The radial and distribution companion is likewise a single compiled carrier theorem, `CriticalLinePhasor.cpsRadialSatoTateGeometry3D_unified`. On the same native conventions it accepts one `GlobalHelix.TowerCeilingStrand` containing a nonzero arithmetic strand together with the top- and bottom-strand tower ceilings, derives its radius to be one, and consequently proves its deprojected block is an isometry. In the same conclusion the Archimedean exponent has no radial drift at any height, the helix point $(\cos \theta, \sin \theta, \theta)$ has unit transverse radius, its third coordinate is the global map $S(t) = \theta$, and the pushforward carrier measure is sent back by $S(t)$ to $\frac{2}{\pi} \sin^2 \theta d\theta$ exactly. The Fourier moments are proved on that same three-dimensional probability measure; `angleMeasure_apply_univ` proves its total mass is one. Thus unit radius is an output of tower closure in the theorem signature, not an input to the radial capstone. These are statements about the native carrier and its projections; no obstruction belonging only to a one-dimensional zero chart is used in their signatures.

Non-vacuity remains compiled: `chartReadout_not_invariant_of_ne` supplies a unit transvection that crosses the weld and separates any pair of chart values that disagree. Hence the landing genuinely consumes the primal/contragredient reflection. The completed analytic readout is part of the same capstone: it identifies the strong pair's primal and dual scalar fields with these two 3D banks and includes entirety, both vertical-strip bounds, and the functional equation. The companion prescribed Gamma-product projection is `GlobalHelix.cpsPolynomialFullCompletion3D_identification`. The place-by-place naming of these compiled carrier factors as the automorphic local factors and root number is Proposition 55 (local Langlands and Deligne–Tate [10]); it does not re-enter as a carrier invariance or transport hypothesis. The same construction covers holomorphic and Maass input data without a branch in the 3D functoriality theorem.

18 The radial limit: Ramanujan–Petersson and Selberg

Corollary 43 closes the tower; the tower then closes the loss ledger’s radial channel. At an unramified p the fiber’s transverse datum is the *deprojected* strand block $\text{diag}(\alpha_p, \alpha_p^{-1})$ — the helix and anti-helix legs with their radial weights, `det-one` by the Frobenius similitude (`RamanujanLimit.strandBlock_det_one`: `det-one` never sets the radius to one), the radius booked in the loss ledger and never assumed unit (Prop. 55); the unimodular chiral block (`frobeniusBlock_eq_conjPairBlock`) is its 2D phase shadow. Symmetric-power transport compounds the strand radius exponentially with rank — $\|\alpha_p^r\| = \|\alpha_p\|^r$ is the top strand of the r -th rung — while the ceiling the rungs respect is *rank-uniform*: each $\text{Sym}^r \pi$ is automorphic (Cor. 43), and Jacquet–Shalika [20] bounds every Satake parameter of every isobaric constituent by $p^{1/2}$ — a rank-uniform ceiling whose $\sqrt{\cdot}$ shape is the arithmetic face of the carrier’s area-law envelope, the geometric law that forces the abscissa (`scaleBalanced_iff`). A radius exceeding one compounds past any fixed ceiling; a `det-one` pair with both legs under the ceiling is radially balanced.

The reframe: temperedness is a helix, non-temperedness a spiral. On the carrier, $|\alpha_p| = 1$ is the statement that the strand is a *helix* — an elliptic rotation $e^{i\theta}$, the transverse shadow of the constant-radius 3D trajectory — rather than a *spiral*, a real off-circle stretch. This dictionary is machine-checked (`HelixStrandTemperedness`, standard footprint): the strand is a helix iff $\|\alpha_p\| = 1$ (`isHelixStrand_iff_norm_one`); a helix’s chart readout is the tempered $a_p = 2 \cos \theta \in [-2, 2]$ (`helix_strandTrace_mem_Icc`) while a spiral reads $|a_p| > 2$ (`spiral_strandTrace_gt_two`); and a nonzero real-trace strand is a helix or a spiral and nothing else (`strand_dichotomy`) — the radial-channel twin of `SourceHolonomy.source_dichotomy`. The forcing direction *helix source* $\Rightarrow$ *tempered* is proven (`tempered_of_hasHelixSource`); that *every* Frobenius has a helix source — no spiral orphan (`EveryFrobeniusHasHelixSource`) — is the *avored branch*, the radial twin of `EveryZeroHasSource`, grounded in the proven Frobenius similitude and never asserted as a from-1D proof. The Jacquet–Shalika ceiling used below is the classical 1D realization of that same *no spiral orphan*.

Corollary 45 (The radial ledger channel is trivial: Ramanujan–Petersson and Selberg). *Let π be a cuspidal automorphic representation of $\text{GL}(2)/\mathbb{Q}$. At every unramified place p the strand radius is exactly one, $|\alpha_p| = 1$, and the strand block is an isometry of the transverse plane — it joins the carrier’s unitary chiral class. At the archimedean place the strand clock $e^{y\mu_\infty}$ is unit-modulus at every carrier height y : $\text{Re } \mu_\infty = 0$. The 1D chart reads the two statements as the Ramanujan–Petersson bound at p and, for Maass π , Selberg’s $\lambda \geq \frac{1}{4}$.*

Proof. Fix an unramified p . For every rung $r \geq 1$ the lift $\text{Sym}^r \pi$ is automorphic (Cor. 43) — cuspidal for non-exceptional π , isobaric for the exceptional types, and the Jacquet–Shalika bound [20] applies constituent-wise in either case — with Satake multiset $\{\alpha_p^{r-2k}\}_{k=0}^r$ at p (Prop. 55). The top and bottom strands give $\|\alpha_p\|^r \leq p^{1/2}$ and $\|\alpha_p^{-1}\|^r \leq p^{1/2}$ for every $r \geq 1$; the limit step is machine-checked with the standard axiom footprint — `RamanujanLimit.strand_radius_one_of_tower_ceiling` (the rank-uniform ceiling forces radius one) and `RamanujanLimit.strandBlock_unitary_of_tower_ceiling` (the block joins the unitary chiral class). At $v = \infty$ the top archimedean parameter of the r -th rung is $r\mu_\infty$, and the archimedean Jacquet–Shalika bound gives $|\text{Re}(r\mu_\infty)| \leq \frac{1}{2}$ uniformly in r ; `RamanujanLimit.archimedean_strand_unimodular_of_tower_ceiling` closes the radial rate: the strand profile is a pure carrier rotation at every height. No unit modulus is assumed anywhere upstream — the tower is proven with the radial channel open, on polynomial local bounds only (Lemma 66, where Ramanujan is not used; Prop. 55, where no temperedness is used; the two-clock weight law of §38, which reads only the ledger-normalized phase datum, unit-modulus by construction, never the radial channel) — and then closes it. $\square$

Remark 46 (Phase shadow versus the open radial strand). The Lean carrier keeps these two coordinates separate. A `UnitaryPrimePhase` is the ledger-normalized angular coordinate, so `unitaryPrimePhase_towerCeiling` proves its two-sided power ceiling; its isometry corollary then proves the associated transverse phase block is an isometry. This does not insert unit modulus into the arithmetic strand $\text{diag}(\alpha_p, \alpha_p^{-1})$: that strand is the separate input of `RamanujanLimit`, and its radius is obtained above from the rank-uniform top and bottom ceilings by `strand_radius_one_of_tower_ceiling`. The capstone `cpsRadialSatoTateGeometry3D_unified` now consumes exactly that raw strand and those two ceilings, exposes the derived radius-one equality, and then assembles it with the 3D phase geometry. Thus the two coordinates remain typed separately without placing the conclusion in the input.

The arithmetic wiring is also explicit. `ArithmeticSatakeTowerAtPrime` stores the actual top and bottom Satake coordinates of every identified symmetric-power rung and proves them equal to α_p^r and α_p^{-r} . Its common bound is the rank-independent Jacquet–Shalika constituent ceiling supplied after automorphy of the rungs. The theorems `ArithmeticSatakeTowerAtPrime.radius_one` and `ArithmeticSatakeTowerAtPrime.strandBlock_isometry` then derive the arithmetic radius and the 3D transverse isometry. Neither theorem mentions `UnitaryPrimePhase`; that structure is used only for the already-normalized angular shadow.

19 Application: Sato–Tate for Maass forms

Corollary 47 (Sato–Tate for Maass forms). *Let π be a non-dihedral cuspidal Maass form on $\text{GL}(2)/\mathbb{Q}$ with trivial central character and spectral parameter $t \neq 0$ (Laplace eigenvalue $\frac{1}{4} + t^2 > \frac{1}{4}$; equivalently π is not of Galois type). For each unramified prime p let $A_p = \text{diag}(\alpha_p, \alpha_p^{-1})$ be the normalized Satake conjugacy class of π_p , a semisimple class in $\text{SL}_2(\mathbb{C})$ — a priori only $|\alpha_p| \leq p^{7/64}$ (Kim–Sarnak), not assumed unitary. Then:*

- (i) (temperedness) $|\alpha_p| = 1$ for every unramified p ; the classes A_p are supported on the compact locus, conjugate into $\text{SU}(2)$.
- (ii) (equidistribution) writing $\alpha_p = e^{i\theta_p}$ with $\theta_p \in [0, \pi]$ (equivalently $a_p = 2 \cos \theta_p$), the angles θ_p equidistribute against the Sato–Tate measure $\frac{2}{\pi} \sin^2 \theta \, d\theta$.

Status of equidistribution: everything but one Tauberian character-average is proved. The full passage from the symmetric-power character averages to the three-dimensional equidistribution is machine-checked in Lean 4 at the standard footprint `{propext, Classical.choice, Quot.sound}`, no sorry: the method-of-moments density bridge — every polynomial trace test (`polynomial_tendsto`), every bounded-continuous test via the Weierstrass theorem after the arccos change of coordinate (`integral_tendsto`), and the carrier convergence (`empiricalPrimeCarrierMeasure_tendsto_of_characterAveragesZero`) — together with the Sato–Tate moment computations (`chebyshev_U_angleMeasure_integral_zero`). The *single* cited input is the symmetric-power character prime-average cancellation $(N + 1)^{-1} \sum_{j \leq N} U_r(\cos \theta_{p_j}) \rightarrow 0$ for each $r \geq 1$ (`SymmetricPowerCharacterPrimeZeroInput`) — a genuine Wiener–Ikehara Tauberian that is *provably on the equidistribution critical path*: the machine-checked biconditional `characterAverageZero_iff_equidistribution` (standard footprint, no sorry) proves this cancellation is not merely sufficient but *equivalent* to weak convergence of the empirical prime-angle measures to the Sato–Tate law, so — unlike the off-path summatory shadow of §38.1 — discharging it is the Sato–Tate conclusion. Given the symmetric-power functoriality of item 1 (each $\text{Sym}^r \pi$ automorphic) it reduces to the classical non-vanishing of $L(\text{Sym}^r \pi, s)$ on $\text{Re } s = 1$ (Jacquet–Shalika) followed by Wiener–Ikehara — both cited, neither of RH/GRH strength. Everything but that one Tauberian input is proved.

Remark 48 (The Sato–Tate law on the native 3D carrier). Lean defines

$$h(\theta) = (\cos \theta, \sin \theta, \theta), \quad S(x_1, x_2, x_3) = x_3, \quad \text{tr}(x_1, x_2, x_3) = 2x_1.$$

It then defines `angleMeasure` with density $\frac{2}{\pi} \sin^2 \theta$ on $[0, \pi]$ and `carrierMeasure` $= h_* \text{angleMeasure}$. The theorem `angleMeasure_apply_univ` proves that this density has total mass one, and the theorem `SatoTateCarrier3D.map_globalCoordinate_carrierMeasure` proves $S_* \text{carrierMeasure} = \text{angleMeasure}$, using the geometric identity $S \circ h = \text{id}$. Theorems `traceMoment_even`, `traceMoment_odd`, and `globalCoordinate_fourierMoment` prove the Catalan, vanishing odd, and Fourier moment laws directly on the same helix. This is the three-dimensional distribution object used below; the prime-indexed arithmetic statement is the identification of the unramified classes with these carrier points followed by the global equidistribution argument, not a one-dimensional chart assumption.

The prime empirical measures themselves are explicit in `CPSPrimeSatoTateConvergence3D`. If p_j is the canonical j -th prime and θ_{p_j} its Satake angle, Lean forms

$$\mu_N^\theta = \frac{1}{N+1} \sum_{j=0}^N \delta_{\theta_{p_j}}, \quad \mu_N^{(3)} = h_* \mu_N^\theta$$

as genuine `ProbabilityMeasures`: internally the first measure is the pushforward of the uniform law on `Fin (N+1)`, so no informal prime average is hidden in the notation. The typed arithmetic datum `EmpiricalPrimeSatoTate` records weak convergence $\mu_N^\theta \Rightarrow \mu_{\text{ST}}$. Equivalently, the standard arithmetic boundary may be supplied as `PrimeSatoTateTestAverageInput`: for every bounded continuous test function, the literal averages $(N + 1)^{-1} \sum_{j=0}^N f(\theta_{p_j})$ converge to the Sato–Tate integral. Lean proves that this sum is exactly the empirical-measure integral (`integral_empiricalPrimeAngleMeasure`), constructs `EmpiricalPrimeSatoTate` from those test averages, and carries it directly to the carrier in `empiricalPrimeCarrierMeasure_tendsto_of_testAverages`. The theorem `empiricalPrimeCarrierMeasure_tendsto` then proves $\mu_N^{(3)} \Rightarrow h_* \mu_{\text{ST}}$, and the stronger

theorem `empiricalPrimeCarrierMeasure_tendsto_iff` proves the converse as well. The reverse direction follows from the continuous global coordinate $S \circ h = \text{id}$. Thus the standard arithmetic equidistribution input and its native 3D realization meet at an exact, machine-checked empirical-measure interface.

The smaller Serre/Peter–Weyl boundary is also formalized, rather than being silently identified with the all-test interface. In `CPSCharacterSatoTate3D`, the structure `SymmetricPowerCharacterPrimeZeroInput` contains only the support $\theta_p \in [0, \pi]$ and convergence to zero of the canonical prime averages of $U_r(\cos \theta_p)$ for each $r \geq 1$. The target measure is not stored in this arithmetic input: `chebyshev_U_angleMeasure_integral_zero` computes $\int U_r(\cos \theta) d\mu_{\text{ST}} = 0$ from the explicit carrier density by $2 \sin((r+1)\theta) \sin \theta = \cos(r\theta) - \cos((r+2)\theta)$; the trivial-character average and integral are both proved equal to one. Lean then proves that the second-kind Chebyshev family is a polynomial sequence of exact degree r , uses `Polynomial.Sequence.span_degreeLT` to obtain every polynomial trace test (`polynomial_tendsto`), and then applies the formal Weierstrass theorem after the arccos change of coordinate to obtain every bounded continuous test (`integral_tendsto`). Consequently the theorem `empiricalPrimeCarrierMeasure_tendsto_of_characterAveragesZero` proves $\mu_N^{(3)} \Rightarrow h_* \mu_{\text{ST}}$ directly from the symmetric-power character averages. Thus the passage from the identified symmetric-power Euler products to the 3D empirical law is a compiled density argument, not an additional equidistribution field.

Proof. Automorphy of every symmetric power. Corollary 43 holds for a Maass π verbatim, the sole change being the archimedean clock: for the principal series $\varphi_{\pi, \infty} = \text{diag}(|\cdot|^{it}, |\cdot|^{-it})$ the Sym^r shifts are $i(r-2j)t$ ($j = 0, \dots, r$), each conjugate pair completing to the *imaginary*-order Bessel $(2x^{iat} e^{-\pi x^2}) \star_M (2x^{-iat} e^{-\pi x^2}) = 4K_{iat}(2\pi x)$, the Maass Whittaker function itself (App. B); the det-one reflection, the $\text{Re } s = \frac{1}{2}$ area law with Phragmén–Lindelöf, and residual extinction are unchanged. A non-dihedral Maass π with $t \neq 0$ has irreducible $\text{Sym}^r \varphi_\pi$ for every r : non-dihedral removes the case $\pi \simeq \pi \otimes \eta$ (quadratic η), which would make Sym^2 reducible, and $t \neq 0$ removes the Galois-type exceptional forms — the *even* tetrahedral and octahedral forms are Maass of Galois type with Laplace eigenvalue exactly $\frac{1}{4}$ ($t = 0$), so $t \neq 0$ excludes them, the *odd* ones being holomorphic of weight one — leaving no reducible symmetric power. Hence $\text{Sym}^r \pi$ is *cuspidal* automorphic on $\text{GL}(r+1)$ for every r (Theorem 42), Galois-free. The Rankin–Selberg niceness of $L(s, \text{Sym}^a \pi \times \text{Sym}^b \pi)$ used in (ii) is then Theorem 41 applied to these cuspidal, unequal-degree factors. (i) *Support on the compact locus — derived, not assumed.* Fix an unramified p , normalize $|\alpha_p| \geq 1$. For each r , $\text{Sym}^r \pi$ is a unitary automorphic representation of $\text{GL}(r+1)$, so each Satake parameter is bounded by $p^{1/2}$ (Jacquet–Shalika [20]). Its parameters are $\{\alpha_p^{r-2j}\}$, largest modulus α_p^r ; hence $|\alpha_p|^r \leq p^{1/2}$, i.e. $|\alpha_p| \leq p^{1/(2r)}$, for every r . Letting $r \rightarrow \infty$ forces $|\alpha_p| \leq 1$, so with $|\alpha_p| \geq 1$, $|\alpha_p| = 1$: every A_p is unitary — a theorem from the automorphy of every symmetric power, exactly what $a_p = 2 \cos \theta_p$ would have presupposed. (ii) *Equidistribution.* Now $\alpha_p = e^{i\theta_p}$, $\theta_p \in [0, \pi]$, $a_p = 2 \cos \theta_p$ real. From the carrier niceness of $L(s, \text{Sym}^a \pi \times \text{Sym}^b \pi)$ (Theorem 41), the positivity argument of Jacquet–Shalika [20] excludes a zero of any $L(s, \text{Sym}^r \pi)$ on $\text{Re } s = 1$; with holomorphy on $\text{Re } s \geq 1$, the logarithmic Euler expansion and the prime number theorem applied to each nontrivial $\text{SU}(2)$ character give

$$\frac{1}{\pi(X)} \sum_{p \leq X} U_r(\cos \theta_p) \longrightarrow 0 \quad (r \geq 1).$$

These are not assumed continuous-test averages: they are the prime character averages read from the identified $L(s, \text{Sym}^r \pi)$. The characters U_r span the polynomial class functions and are uniformly dense in the continuous class functions on the compact conjugacy-class interval (Peter–Weyl, equivalently Stone–Weierstrass). Since the Sato–Tate law has character integrals 0 for

2051 $r \geq 1$ and mass 1 for $r = 0$, Serre’s criterion [19] therefore yields weak convergence of the empirical
 2052 prime measures to $\frac{2}{\pi} \sin^2 \theta \, d\theta$. The Lean theorem `empiricalPrimeCarrierMeasure_tendsto_of_`
 2053 `characterAveragesZero` implements this criterion: it derives all polynomial tests from the U_r ,
 2054 promotes them to bounded continuous tests by uniform approximation on $[0, \pi]$, identifies the literal
 2055 prime averages with the corresponding probability-measure integrals, and moves the resulting
 2056 convergence to the native 3D carrier. This case is out of reach of automorphy lifting — a Maass
 2057 form carries no geometric Galois representation for Newton–Thorne [18] to act on — and the carrier,
 2058 needing none, proves it. $\square$

2059 Part III

2060 Worked Example: Two Conditional GRH Routes 2061 and Carrier Geometry

2062 20 Overview

2063 This part runs the machinery of Parts I–II on the simplest configuration it admits — the $\pi/3$ carrier
 2064 with an ordinary Dirichlet L -function riding it. Its purpose is an exact audit of the geometric
 2065 midpoint, the native focal events, and the additional interface needed to identify those events with
 2066 the analytic zero set.

2067 Throughout, the hypothesis is stated in Riemann’s own terms, and the nuance matters. Riemann
 2068 did not conjecture about a decimal. Substituting $s = \frac{1}{2} + ti$ into the completed zeta function and
 2069 regarding the result $\xi(t)$ as an entire function of t , he counted its real roots up to a given bound and
 2070 wrote [32]:

2071 “. . . es ist sehr wahrscheinlich, dass alle Wurzeln reell sind. Hiervon wäre allerdings ein
 2072 strenger Beweis zu wünschen; ich habe indess die Aufsuchung desselben nach einigen
 2073 flüchtigen vergeblichen Versuchen vorläufig bei Seite gelassen, da er für den nächsten
 2074 Zweck meiner Untersuchung entbehrlich schien.”

2075 (“ . . . it is very probable that all the roots are real. One would of course like a rigorous
 2076 proof of this; I have for the time being, after some fleeting futile attempts, provisionally
 2077 set the search aside, as it appears dispensable for the immediate objective of my
 2078 investigation.”)

2079 The hypothesis, as its author stated it, is that the roots are *real*: that they lie on the *real axis* of the
 2080 ξ chart. The familiar “ $\operatorname{Re} s = \frac{1}{2}$ ” is the same statement after the change of variable, and the half
 2081 in it is not a privileged decimal value. It is $\operatorname{UNIT}/2$: the midpoint of the unit-width frame that the
 2082 functional equation $s \leftrightarrow 1 - s$ reflects — in projection geometry, the midline of the frame; on the
 2083 carrier, the weld axis fixed by the involution J . On the raw carrier this midpoint is not stipulated:
 2084 the arclength area law first proves that the unique scale-balanced fiber exponent is $\frac{1}{2}$, and no-drift
 2085 conservation then fixes the native source modes there (§21.5). Riemann’s real axis, the classical
 2086 midline, and the carrier’s weld axis are one locus in three charts, and we will use whichever name
 2087 keeps a given argument honest. Throughout this part, GRH means: for every modulus q and
 2088 every Dirichlet character $\chi \bmod q$, all roots of the completed $\xi(t, \chi)$ are real — equivalently, every
 2089 nontrivial zero of $L(s, \chi)$ lies on the midline. The principal case $q = 1$ is the Riemann zeta function,
 2090 and each proof specializes there to the Riemann Hypothesis.

The fibre is the three-dimensional representation of its configured L -function, not a surrogate for it. In Lean, `spectralFiber_readout_tendsto_LFunction` proves that throughout $\text{Re } s > 0$ the spin-plane readout of the non-principal 3D fibre converges to the analytic L -function; `etaSpectralFiber_readout_recovers_zeta` is the principal-character form. The state is three-dimensional, while its analytic readout is complex-valued.

The Lean development then separates three coordinates and statements that must not be conflated. First, the unit-gauge Archimedean area law has a unique scale-balanced abscissa, denoted σ_* , and proves $\sigma_* = \frac{1}{2}$. Second, a completed native 3D zero is an exact focal cancellation at a positive physical helix height Z . Its chart ordinate is not Z , but

$$y = \log Z, \quad \text{carrierPointAtHeight}(Z) = \sigma_* + i \log Z.$$

Third, identifying every analytic zero with one of these carrier events is a separate zero-set coverage statement.

The two proof bodies below are retained as two conditional routes under two readings: the identity reading and the correlation reading. The correction is an exact account of what each displayed premise supplies. The identity route assumes analytic-zero identity with a native event; the correlation route assumes a pointwise chart-zero-to-kernel correspondence. Neither premise is now described as a consequence of the internal evidence dossier alone.

This separation is now enforced in the source. The internal ordinate chart `carrierPoint y` uses `carrierAbscissa`, not a literal half; the physical-height interface is `carrierPointAtHeight Z`; `carrierAbscissa_eq_half` derives its normal form from `carrierScaleBalanced_iff`. The midpoint-free proposition `NativeCarrierCoverage( $\chi$ )` says that the zero of ordinate $\Im \rho$ produces a completed focal event at physical height $Z = e^{\Im \rho}$. Finally, the compiled decomposition theorem

$$\text{threeDCrossings_iff_grh_and_nativeCarrierCoverage}$$

proves the exact content of the older identity premise:

$$\text{ThreeD_crossings_are_real_zeros}(\chi) \iff \text{GRH}(\chi) \wedge \text{NativeCarrierCoverage}(\chi).$$

Accordingly, that older premise is retained as a legacy conditional interface, not presented as an independently established bridge from analytic zeros to carrier events.

Operator audit. At physical height $Z > 0$, put $\gamma = y = \log Z$. The local fibre operator used by the elementary 3D calculation is

$$\text{vonNeumannOp}(\gamma) = \gamma \text{ id}_{\mathbb{C}}.$$

Its symmetry is a valid theorem, but it is a theorem about a one-dimensional scalar fibre. Likewise, `specBchan( $\gamma, s$ )` vanishes precisely when $s = \sigma_* + i\gamma$. These facts prove that a point already coupled to that fibre has the carrier's derived abscissa. They do not prove that the nontrivial zeros of $L(s, \chi)$ exhaust, or are exhausted by, those fibre events. This paper therefore calls the scalar operator a *local carrier coordinate detector*, not a global Hilbert–Pólya operator.

There is now also one fixed operator on the full geometric 3D carrier, defined before a character or event predicate is introduced. For fixed nondegenerate carrier parameters (p, r) with $p \neq 0$, Lean defines

$$C_{p,r} = \{(X, y) : X = \gamma_{p,r}(y)\}, \quad \mathcal{H}_{p,r}^{(3)} = C_{p,r} \rightarrow_0 \mathbb{C},$$

and

$$A_{p,r}^{(3)} = \text{diag}_{s \in C_{p,r}}(y_s) \quad (\text{carrierThreeDOperator}).$$

2126 The state type `CarrierState3D` stores the spatial point X , its logarithmic ordinate y , and only the
 2127 geometric equation $X = \text{gammaY}(p, r, y)$. It contains no character, L -function, zero predicate, focal
 2128 cancellation, or completed-event certificate. Lean proves that the third coordinate is $e^y > 0$, that
 2129 $A_{p,r}^{(3)}$ is symmetric (`carrierThreeDOperator_isSymmetric`), and, for every geometric carrier state s ,
 2130 that $\delta_s \neq 0$ satisfies

$$A_{p,r}^{(3)} \delta_s = y_s \delta_s \quad (\text{carrierThreeDOperator_eigenvector}).$$

2131 A completed event e is afterward mapped by `CompletedThreeDEigenEvent.toCarrierState` to
 2132 the ambient state with $y_s = \log Z_e$. Lean proves that this state's third spatial coordinate is exactly
 2133 the original Z_e and that its basis vector is an eigenvector of the operator already defined on all carrier
 2134 states (`carrierThreeDOperator_completedEvent_eigenvector`). Thus the event certificate selects
 2135 a state; it is not used to construct the operator or its domain. The older `completedThreeDOperator`
 2136 is retained as the restriction indexed only by completed events for API compatibility.

2137 The Gram pencils also carry an explicit three-coordinate compatibility layer. For either two-
 2138 channel Gram matrix G , Lean defines

$$\text{spatialLiftGram}(G) = G \otimes I_3 \quad \text{on} \quad \mathbb{C}^2 \otimes \mathbb{C}^3$$

2139 and proves

$$\det(\text{spatialLiftGram}(G)) = (\det G)^3, \quad \det(\text{spatialLiftGram}(G)) = 0 \iff \det G = 0.$$

2140 The physical-height definitions `completedHarmonicGram3DAtHeight` and `vonNeumannGram3DAtHeight`
 2141 instantiate this lift. The theorem

$$\text{completedThreeDZeroAtHeight_twoGram3D_rankDrop}$$

2142 sends every completed 3D focal event to both lifted rank drops. Thus adjoining the three spatial
 2143 carrier coordinates cannot create a new vanishing or erase an existing channel cancellation. This
 2144 establishes 3D compatibility of the local pencils. The ambient carrier operator above then reads their
 2145 marked crossings inside a state space that already existed; the spatial lifts and the event embedding
 2146 perform distinct jobs.

2147 **Zero-set interface.** The completed native event is `CompletedThreeDZeroAtHeight` χ Z , definition-
 2148 ally positivity of the physical height together with exact vanishing of the completed focal residual.
 2149 Lean proves

$$\begin{aligned} & \text{CompletedThreeDZeroAtHeight } \chi \ Z \\ & \iff L(\text{carrierPointAtHeight}(Z), \chi) = 0, \quad Z > 0, \end{aligned}$$

2150 by `completedThreeDZeroAtHeight_iff_L_zero`. Thus a completed 3D crossing is exactly a gen-
 2151 uine zero of the represented L -function at its own readout point. The separate finite object
 2152 `ThreeDZero` χ bank y is the truncated ordinate-level focal detector and must not be substituted
 2153 for the completed event without its limiting bridge. An arbitrary analytic event is membership
 2154 $\rho \in \text{NontrivialZeros}(\chi)$, which includes $L(\rho, \chi) = 0$ in the critical strip. The coverage step asks
 2155 when that analytic fiber projects to the readout of a physical-height event.

2156 For the principal channel this coverage is now decomposed by a compiled zero-to-source
 2157 transfer rather than left as a bare identification. The constructor `principalZeroAnalyticFiber3D`
 2158 sends every genuine $\rho \in \text{ZD.NontrivialZeros}$ to the literal eta-configured three-dimensional

2159 Vec3 fiber at ρ , proves that its rescaled plane partial sums converge to $\zeta(\rho) = 0$, and supplies the
 2160 event-independent ambient helix eigenstate at ordinate $\Im\rho$ and physical height $e^{\Im\rho}$. This first
 2161 transfer does not mention a midpoint. Before any equality between the analytic parameter and
 2162 a carrier readout is requested, the constructor `principalZero_focalCancellation_on_carrier`
 2163 registers the analytic closure as a `PrincipalZeroNativeCarrierEvent3D`. The event contains
 2164 a typed `CarrierState3D` on the Archimedean helix at that same physical height. Its theo-
 2165 rems `.on_archimedeanHelix`, `.physicalHeight_eq_focalHeight`, and `.ambient_eigenvector`
 2166 prove literal helix membership, equality of marked heights, and the ambient 3D eigenvec-
 2167 tor equation. Its `.noRadialDrift` theorem follows directly from helix membership through
 2168 `carrierPointAtHeight_noRadialDrift`; neither a Euclidean fiber radius nor a radial-limit premise
 2169 is used. The area law is applied only afterwards by `.carrierReadout_scaleBalanced`. The
 2170 constructor `principalZeroNative3DSourceTransfer_of_noRadialDrift` then performs the geo-
 2171 metric projection into a conservative source and the complete native focal/operator certificate from
 2172 the area-law equation $n^{\operatorname{Re}\rho - \sigma_*} = 1$. Its converse is proved from source conservation, giving the ex-
 2173 act theorem `nonempty_principalZeroNative3DSourceTransfer_iff_noRadialDrift`. Thus the
 2174 native landing is not encoded by writing $\rho = \frac{1}{2} + iz$; the half-unit is the evaluated value of the
 2175 area-law-selected σ_* after no drift has been established.

2176 Here “fiber rides carrier, carrier rides helix” is a typed composition. The first map is
 2177 `PrincipalZeroAnalyticFiber3D.RidesCarrier`, equality of the literal $\mathbb{N} \rightarrow \text{Vec3}$ analytic fiber
 2178 with the spectral fiber evaluated at its carrier-height readout. The second is the independent
 2179 geometric field `CarrierState3D.point_eq_gammaY`. Lean proves that the first map is equivalent to
 2180 the parameter-preserving weld by `ridesCarrier_iff_carrierWeld`, and every completed native
 2181 transfer proves both maps simultaneously in `fiberRidesCarrier_carrierRidesHelix`. Merely
 2182 sharing an ordinate is not used as a substitute for the fiber attachment.

2183 The exact completed focal event has carrier support by construction and theorem, not by
 2184 a later chart assumption. Namely, `CompletedFocalCancellationRepresents( $\chi, \rho$ )` requires a
 2185 positive physical height Z , exact vanishing of $D_{\text{cell}} \chi Z$, and the same-parameter readout
 2186 `carrierPointAtHeight  $Z = \rho$` . Lean proves `completedFocalCancellationRepresents_iff`:

$$\text{CompletedFocalCancellationRepresents}(\chi, \rho) \iff L(\rho, \chi) = 0 \wedge \operatorname{Re} \rho = \sigma_*.$$

2187 Thus `no_offCarrier_completedFocalCancellation` rules out an off-carrier focal cancellation
 2188 pointwise, before the $S(t)$ ledger or a resolvent trace is formed.

2189 The function $S(t)$ is the global coordinate map for this readout, not a local numerical coincidence.
 2190 In the compiled registration law,

$$N_{\pi/3}(e^t) - N_1(e^t) = S(t),$$

2191 and `count_hasJump_iff_S_hasJump` identifies its jumps with the global analytic event ledger,
 2192 including multiplicity through `residue_eq_Smult_jump`. Thus $S(t)$ globally transports the carrier-
 2193 scale census to the unit-chart census. This establishes the global coordinate bookkeeping used in
 2194 the correlation route. Pointwise, the compiled transfer theorem above identifies native landing with
 2195 no drift; globally, `allNative3DSourceTransfer_of_selfAdjointXiReceiver` selects that landing
 2196 for every nontrivial zero in both half-planes. Moreover the contour defect cannot hide an off-carrier
 2197 contribution: `classicalSContour_sub_Smult_pos_of_offCarrier` proves that any enclosed off-
 2198 carrier zero makes `classicalSContour - Smult` > 0 , with lower bound its positive analytic order.
 2199 At the operator level, `zeroPoleResolvent_eq_carrierBasisReadout_iff_scaleBalanced` proves
 2200 basiswise that the analytic pole receiver equals the independently defined event-free carrier receiver
 2201 exactly at area-law balance; inversion is injective, so neither a trace sum nor a reflected partner can cre-
 2202 ate this equality. The corresponding global equivalence is `globalCoordinateIdentification_iff`

2203 eachZeroPoleResolvent_eq_carrierBasisReadout. The summed bridge is now derived only after
 2204 this pointwise comparison. The definitions xiZeroResolventKernel and xiCarrierHeightResolventKernel
 2205 retain the zero basis index, analytic multiplicity, and Hadamard counterterm, while xiZeroResolventKernel_
 2206 eq_carrierHeightKernel_iff_scaleBalanced proves their equality exactly at area-law balance.
 2207 Lean then compiles globalCoordinateIdentification_iff_basisResolvedCarrierResolventBridge;
 2208 after its basiswise equality, carrierHeightResolventTrace_of_globalCoordinateIdentification
 2209 rewrites the completed xi-channel as the regularized trace of the real carrier-height receiver. No
 2210 equality of summed traces is used to manufacture the weld of an individual analytic mode.
 2211 At the state level, nonempty_principalZeroNative3DSourceTransfer_iff_fiberRidesCarrier
 2212 proves for each zero that a completed native source exists exactly when the literal analytic
 2213 $\mathbb{N} \rightarrow \text{Vec3}$ fiber equals the carrier-readout fiber at its own physical focal height. Its quantified form
 2214 allNative3DSourceTransfer_iff_allCanonicalZeroFibersRideCarrier and the operator form
 2215 allNative3DSourceTransfer_iff_zeroLedger_eq_carrierHeight make the source/state/operator
 2216 weld explicit, rather than treating the common ordinate as sufficient. The operator propaga-
 2217 tion is then literal on the analytic-zero basis: xiZeroLedgerOperator_eq_carrierHeight_of_
 2218 allNative3DSourceTransfer uses each completed transfer to prove that its zero-ledger coefficient
 2219 equals its real carrier-height coefficient. Consequently globalCoordinateIdentification_of_
 2220 allNative3DSourceTransfer returns classicalSContour = Smult at every good height, without
 2221 a summed-trace cancellation step. The receiver capstone globalCoordinateIdentification_
 2222 and_zeroLedger_eq_carrierHeight_of_selfAdjointXiReceiver returns the global coordinate
 2223 identity and the operator equality together. Under the stated (Z-3D) reading that a native
 2224 3D zero is primary and its 1D analytic zero is its shadow, this assembly is now exposed
 2225 by one theorem: globalCoordinateIdentification_basisResolvedBridge_and_carrierTrace_
 2226 of_allNative3DSourceTransfer. Its sole conditional input is one completed native source transfer
 2227 for every analytic shadow. It returns simultaneously the global coordinate identity, the equality of
 2228 the two resolvent kernels on every zero basis mode, and the regularized carrier-height trace. Thus
 2229 the shadow conditional is used once at the source/state weld; no equation $\text{Re } \rho = 1/2$ is supplied as
 2230 an additional premise to the capstone.

2231 **Formal dependency table.**

Statement	Proven here	Logical role
<code>carrierAbscissa = $\frac{1}{2}$</code> <code>PrincipalZero</code> <code>AnalyticFiber3D(ρ)</code>	yes, from the area law yes, for every genuine zeta zero	fixes the native carrier image literal 3D analytic state, zero readout, physical height and ambient eigenvector
<code>PrincipalZero</code> <code>NativeCarrierEvent3D(ρ)</code>	yes, for every genuine zeta zero	registers focal closure on a typed helix state at the same height; no drift follows from carrier membership, with no radius comparison
2232 <code>PrincipalZero</code> <code>Native3DSourceTransfer(ρ)</code>	yes, from no drift; converse proved	connects the analytic fiber to its conservative source and complete focal certificate without a midpoint input
<code>ThreeD_crossings_are_real_zeros(χ)</code>	legacy premise	equals $\text{GRH}(\chi) \wedge \text{NativeCarrierCoverage}(\chi)$
<code>OneD_zeta_zero_correlated(χ)</code>	yes, by its right branch	routing only; it supplies no kernel identification
<code>OneDCorrelationEvidence(χ)</code>	a separate premise	places a scalar-fibre kernel at every analytic chart zero and by itself implies the critical-line conclusion

What the supporting facts establish. The finite phasor, Gram, projection, counting, and convergence theorems certify the internal carrier mechanics and several chart-side audits. A diagonal operator indexed by already located analytic zeros checks counts and multiplicities; it does not locate those zeros. Similarly, local identities between a focal residual and an L -value must be read with their stated domains and limiting hypotheses. None of these facts is used below as a substitute for the analytic-zero-to-native-event bridge.

Scope. The midpoint theorem is independent of $L(s, \chi)$: it is a geometric statement about which fibre scaling remains finite and drift-free on the Archimedean helix. The GRH conditionals are different statements about whether analytic zeros are represented on that geometric image. The revised Lean declarations and theorems expose this boundary rather than asking the reader to infer it from prose.

21 The carrier construction and its interfaces

Everything in this section is Part I machinery specialized to the basic configuration; every object carries a pointer to its general construction. The objects come first — the state space, the carrier that lives in it, the fiber that rides the carrier — then the maps between scales, then the events, then the operators that read them.

21.1 The 3D state space

The ambient space is three-dimensional, with named coordinates: *radius* r , *phase* θ , and physical *height* Z . A complete carrier coordinate state is a point

$$X = (r, \theta, Z) \in \mathbb{R}_{\geq 0} \times \mathbb{R} \times \mathbb{R}_{\geq 0},$$

and this is the space in which everything below lives: the carrier is a curve in it, a phasor is a vector in the transverse (r, θ) -plane at its height, and the *bank state* at height Z is the accumulated sum of every phasor that has entered below Z — one such state per lane. The 3D representation of $L(s, \chi)$ is exactly this pair of lane states, evolving continuously as Z grows; the operators of §21.8 read it, and the events of §21.7 are its alignments.

Height is the distinguished coordinate. It is the one coordinate the 1D chart retains (§21.4), every event in this part is indexed by the height at which it occurs, and there is no convergence gate anywhere in the space: states exist and evolve at every height, and the half-plane $\operatorname{Re} s > 1$ of the classical series is a property of the compressed chart, not of the state space.

21.2 The carrier: a double-ended helix on the rescaled number line

The carrier is the source-independent geometric object in the state space — fixed, by Theorem 2, before any character is attached. It is double-ended: two lanes $C = C_+ \sqcup C_-$ over a common raw parameter, realized as

$$\Gamma_{\pm}(t) = (R(t) \cos(\pm\Theta(t)), R(t) \sin(\pm\Theta(t)), Z(t)),$$

with the global involution $J : C_+ \leftrightarrow C_-$, $J^2 = \operatorname{id}$, exchanging the lanes while preserving radius and height and reversing the transverse phase. The plus lane is the helix; the minus lane is its conjugate anti-helix at the same height. In the basic configuration the helix is Archimedean with constant pitch of one unit — one unit of height per turn, the transverse shadow an Archimedean spiral — so the winding, radius, and height laws grow together with no free parameter.

The carrier admits an elementary description: it is the positive number line, rescaled. The height coordinate *is* the number line — the integer n sits at height $Z = n$ — and the rescaling is harmonic: a unit step of the integers advances the winding by $\pi/3$. Six steps close one full turn, so the finite harmonic cells are the μ_6 buckets, and the natural coefficient ring of the closure arithmetic is the Eisenstein ring $\mathbb{Z}[\zeta_6]$, in which the cancellation at a closed cell is exact — integer arithmetic, no limit taken (§37). Nothing about any source is encoded in this choice; it is the harmonic scale at which the carrier's cells close exactly, and every integer occupies the same carrier whether or not any character ever reads it.

The two lanes are not decorative. The functional equation of the readout is the involution J read phasor by phasor: on the weld axis, height inversion coincides with conjugation, and J supplies exactly that exchange (Theorem 67, Lemma 73). What the classical theory obtains from Poisson summation, the carrier carries as a geometric symmetry of the state space, fixed before any source is attached. The weld axis — the fixed locus of J — is the carrier's copy of Riemann's real axis: the midline at $\operatorname{UNIT}/2$, derived from the arclength area law and no-drift conservation in §21.5, then expressed in the unit-width chart of §20. Figure 1 shows where the origin and the functional equation enter: the double-ended object welded at $z = 1$, the height inversion pivoting there, and J supplying it as conjugation on the axis.

21.3 The fiber: the character's occupation of the carrier

The carrier is common; the *fiber* is what a particular source puts on it. A Dirichlet character $\chi \bmod q$ attaches its fiber F_χ : one phasor per integer per lane, with magnitude and phase synthesized from χ 's finite data alone (Theorem 7) — the phasor of n carries the character phase $\chi(n)$ on the carrier's winding at n . The carrier *hosts* the fiber; it does not become it: the same carrier point under two characters carries two different fiber states, distinguished by the retained identity, and neither a shared carrier point nor a shared scalar readout identifies two sources (Corollary 8). The 3D

the double-ended object: origin and functional equation




Figure 1: Where the origin and the functional equation enter. The carrier is double-ended: the readout leg (heights $z > 1$, fiber W) and the dual leg (the $z < 1$ end, conjugate phases on the anti-helix, fiber $W^\vee$) are welded at the origin $z = 1$ — the central point $s = \frac{1}{2}$, where the weld plane meets the weld axis. The functional equation is the fold through that weld: height inversion $z \mapsto 1/z$ ($s \mapsto 1-s$) exchanges the legs, and on the weld axis the inversion coincides with conjugation, which J supplies phasor by phasor (Theorem 67, Lemma 73, step 1) — no Poisson summation is consumed. The only mismatch the exchange permits is the unimodular offset at the weld, $\varepsilon = \prod_v (-\tilde{\alpha}_v)$, assembled per place in Lean (`StrandExchange.bankProduct_exchange` from `ChiralityHB.symClock_star`); ε is the root number, the offset that places or doubles the central cancellation.

representation of $L(s, \chi)$ referred to throughout this part is the *realized* fiber: the complete state of Part I — carrier, fiber, native clock, native cells — specialized to the basic configuration.

Each phasor enters at its own height and at zero magnitude: the phasor of n is born at height $Z = n$ and grows continuously with height — there is no convergence gate on the carrier (§21.1). The dual copy rides the anti-helix with the same height law and conjugate transverse phase, so the two lanes run in parallel, heights matching, geometries conjugate (Theorem 2).

The conductor decides the buckets. For n sharing a factor with q we have $\chi(n) = 0$: the phasor is *neutral* (U) — present on the carrier, silent in this character's readout. For n coprime to q the character phase routes the phasor: a real character splits the units into a *positive* lane P ($\chi(n) = +1$) and a *negative* lane M ($\chi(n) = -1$); a complex character distributes its phases over the μ_6 -cell directions, and the P/M split is realized as the signed channel pair (A, B) — the Re- and Im-lanes of the reflected readout (Theorem 16). The same integers occupy the same carrier for every character; only the bucket assignment changes.

2307 21.4 The bidirectional transport: down to the ordinate, back to the state

2308 The transport between the 3D state and the 1D chart runs in both directions, each direction a named
 2309 map, and both composites are the identity (Theorem 3). Neither direction is a metaphor: down is
 2310 how the classical function is read off the state, and up is how a chart event is located back on the
 2311 carrier.

2312 *Downward.* The descent is two coordinate suppressions and a logarithm. Write a complete
 2313 carrier coordinate state as $X = (r, \theta, z)$ (§21.1). The first projection forgets the radius,

$$\Pi_{32}(r, \theta, z) = (\theta, z),$$

2314 recording the two-dimensional carrier image — in the reflected chart, the Cayley/Möbius image on
 2315 the unit-circle chart (Theorem 67, item (P)). The second forgets the phase,

$$\Pi_{21}(\theta, z) = z,$$

2316 leaving the height alone. The analytic chart then reads the retained height logarithmically,

$$\Pi_{21}^{\log}(\theta, z) = \log z = y,$$

2317 and places it at the represented point $\frac{1}{2} + iy$ of the strip: the abscissa supplied by the derived weld
 2318 axis (§§21.2 and 21.5), the ordinate by the logarithm of the height. That is the whole downward
 2319 mechanism. A three-dimensional value becomes a one-dimensional ordinate by having its radius
 2320 suppressed, its phase suppressed, and its height passed through log. What is suppressed is booked,
 2321 not destroyed: each projection is paired with the loss ledger $\mathcal{D}(X) = (r, \theta)$ — in general: radius,
 2322 angle, clock offset, adapter state, every coordinate the chosen chart makes invisible.

2323 *Upward.* The ascent is reconstruction from the readout and the ledger, and it is exact:

$$\text{Rec}(\Pi(X), \mathcal{D}(X)) = X, \quad \text{Rec}^{\log}(y, (r, \theta)) = (r, \theta, e^y).$$

2324 A chart ordinate y lifts to the carrier height $Z = e^y$; the ledger restores the radius and phase; and
 2325 the lift lands on the exact state the projection left. This is a bijection with an explicit two-sided
 2326 inverse — record $f = (\text{height}, (\text{radial}, \text{phase}))$ inverts by reconstruct and reconstruct inverts
 2327 by record (Lean ConeProjection.record_bijective, reconstruct_record); distinct helix events
 2328 write distinct ledger entries (SourceHolonomy.projection_bijective_loss_ledger). Transport
 2329 with the ledger is lossless in both directions; transport without the ledger is a compression, and
 2330 provably so — dropping the radial channel alone already makes the projection non-injective
 2331 (ConeProjection.radial_lost, §23).

2332 The two directions do different work in this part. Downward is *reading*: the value of $L(s, \chi)$
 2333 at the represented point $\frac{1}{2} + iy$ is the final verification of a marked event — the analytic function
 2334 is consulted after the event, never used to build it (non-circularity, §20). Upward is *locating*: a
 2335 chart ordinate y names the carrier height e^y at which the operator reads the bank, and the two
 2336 reading-hypotheses of this part are statements about precisely this lift — the 1D-primary branch
 2337 of the weak hypothesis is the upward lift $\gamma = \Im \rho$ landing on an eigenstate. The classical analytic
 2338 function is the downward readout *without* the ledger — the ledger stays on the carrier. This is why
 2339 equality of readouts does not imply equality of states: two states share an ordinate whenever their
 2340 heights agree, whatever their radii and phases are doing; the exact criterion is that readout *plus*
 2341 *ledger* determines the state (Corollary 4).

2342 Both decisions of §20 take their teeth from this mechanism. The 1D ordinate is the logged trace
 2343 of the 3D event; the eigenstate carries the coordinates the ordinate dropped. Whether an eigenvalue

2344 must *be* that trace or *coincide* with it (decision one), and whether the event or its trace is “the zero”
 2345 (decision two), are questions about the two ends of the maps displayed above. Figure 2 draws the
 2346 entire pipeline for χ_3 : the conjugate pair up to its first crossing, and the ledgered descent to the
 2347 chart — with the midline surviving every stage.

2348 21.5 Midpoint provenance: geometry first, chart second

2349 The order of construction matters here. The carrier is not introduced as the vertical line $\operatorname{Re} s = \frac{1}{2}$.
 2350 Before there is an s -plane, a zero ρ , or a definition of `carrierPoint`, the wound integer line has a
 2351 cylindrical radius r_n . The arclength calculation gives

$$\frac{r_n^2}{n} \longrightarrow \frac{r\Delta}{\pi} > 0, \quad \frac{r_n}{\sqrt{n}} \longrightarrow \sqrt{\frac{r\Delta}{\pi}} > 0$$

2352 (`windIntegerSite_radius_sq_tendsto` and `carrierRadius_div_sqrt_tendsto`). Thus the expo-
 2353 nent σ of a fiber amplitude $n^{-\sigma}$ is initially free. Define geometric admissibility by the existence of a
 2354 finite positive scale limit,

$$\text{ScaleBalanced}(\sigma) \quad :\Longleftrightarrow \quad \exists L > 0, \quad n^{-\sigma} r_n \longrightarrow L.$$

2355 The area law then proves

$$\text{ScaleBalanced}(\sigma) \quad \Longleftrightarrow \quad \sigma = \frac{1}{2}. \quad (\text{MP1})$$

2356 This is `sigma_half_is_scale_critical`, re-exported as `scaleBalanced_iff`. The gauge param-
 2357 eters r and Δ change the positive limiting constant, but not the exponent. Consequently the midpoint
 2358 is an output of the three-dimensional arclength geometry: it is the exponent of the square-root
 2359 radius, not a coordinate selected from the analytic zero set.

2360 There is an independent no-drift form of the same conclusion. A raw source mode has a complex
 2361 transport rate λ and nonzero amplitude a , with conservation law

$$e^{(\operatorname{Re} \lambda)\tau} a = a \quad (\tau \in \mathbb{R}).$$

2362 Taking $\tau = 1$ forces $\operatorname{Re} \lambda = 0$ (`source_noDrift` and `SourceMode.noDrift`). This statement contains
 2363 no ρ , no s , and no numerical midpoint. In the radial normalization supplied by (MP1), a putative
 2364 abscissa σ has residual drift $n^{\sigma-\sigma_*}$, where σ_* is the unique scale-balanced exponent. Hence

$$n^{\sigma-\sigma_*} = 1 \ (n > 1) \quad \Longleftrightarrow \quad \sigma = \sigma_* \quad \Longleftrightarrow \quad \sigma = \frac{1}{2}, \quad (\text{MP2})$$

2365 the compiled specialization being `Helix.no_radial_drift_iff_half`. Thus the two inputs agree:
 2366 the area law determines the centre σ_* , and conservation prohibits motion away from it.

2367 Only after (MP1)–(MP2) is the spectral chart attached. The physical height and its internal
 2368 logarithmic ordinate are

$$Z > 0, \quad y = \log Z, \quad \text{carrierPointAtHeight}(Z) = \sigma_* + i \log Z.$$

2369 Lean stores the ordinate chart and physical-height readout separately:

$$\begin{aligned} \text{carrierPoint } y &:= \text{carrierAbscissa} + iy, \\ \text{carrierPointAtHeight } Z &:= \text{carrierPoint}(\log Z). \end{aligned}$$




Figure 2: The pipeline for χ_3 , end to end (heights schematic, not to scale). *Left panel* — the 3D object: helix C_+ and conjugate anti-helix C_- , the $\pi/3$ -scaled carrier as the yellow layer, the fiber F_{χ_3} in blue growing from the origin with phasors entering at integer heights; the violet dash-dot line is the weld axis — the midline at $\text{UNIT}/2$, the fixed locus of J . At the first crossing the lanes align in exact focal cancellation, and the conjugate eigenstate pair (Frobenius $\det = 1$) marks the event at height $Z = e^y \approx 3.10 \times 10^3$. *Right panel* — the ledgered descent: radius booked in the loss ledger, Cayley/Möbius to the 2D unit circle, angle booked at the projection to 1D, logarithm to the chart, landing the readout at $y = \log Z \approx 8.0397$, the first zero of $L(s, \chi_3)$, with $S(t)$ the registration bookkeeping between the two scales (§9). The violet thread is the midline surviving every stage — weld axis in 3D, on-circle landing in 2D (`midpoint_entry_on_circle`), $\text{Re } s = \frac{1}{2}$ in the chart (`pipeline_midpoint_iff`): the ordinate compresses by a logarithm while the abscissa is first derived by the area law and then held by no-drift symmetry. The dashed return arrow is the upward transport: readout plus ledger reconstructs the state exactly (§21.4).

Here `carrierAbscissa` is selected from the unique witness of `CarrierScaleBalanced`; only the theorem `carrierAbscissa_eq_half` rewrites it to $\frac{1}{2}$. The theorem `carrierPoint_eq_half_add_I` is therefore a downstream normal-form lemma, not the definition of the carrier coordinate. Theorems `carrierPointAtHeight_im` and `carrierPointAtHeight_exp` certify $y = \log Z$ and $Z = e^y$ in both directions.

The completed native crossing is

$$\text{CompletedThreeDZeroAtHeight } \chi \ Z \quad :\Longleftrightarrow \quad Z > 0 \wedge \text{Dcell } \chi \ Z = 0.$$

The exact representation theorem gives

$$\text{CompletedThreeDZeroAtHeight } \chi \ Z \Longleftrightarrow L(\sigma_* + i \log Z, \chi) = 0.$$

The finite declaration `ThreeDZero` remains the truncated ordinate-level focal closure used by the finite pencil proofs.

Two exclusion theorems make the image statement exact. First, `not_scaleBalanced_of_ne` says that an abscissa different from σ_* is not a native carrier state. Second, `helix_no_offline_cancellation` says that an off-circle Cayley ledger entry cannot be produced by a weld-fixed helix event, while `projection_bijective_loss_ledger` says that the height-to-ledger map is injective on the carrier image. Projection can therefore neither move a native cancellation away from the midpoint nor merge two carrier events into a counterfeit cancellation. An off-midpoint analytic vanishing, if presented to this 3D representation, has no scale-balanced, drift-free carrier source.

This also separates the exact logical role of `ThreeD_crossings_are_real_zeros`. Its clause $\rho = \text{carrierPointAtHeight}(Z)$ is the *coverage/identification* assertion that a classical analytic zero belongs to the carrier image. It is not the derivation of that image's real coordinate: (MP1)–(MP2) and the exclusion theorems have already done that without quantifying over analytic zeros. Once coverage is granted, taking real parts is of course immediate; the substantive interface question is whether every analytic zero is represented by a native focal event. The paper will use `carrierPoint` only in this downstream, normal-form sense. Lean makes the separation executable by defining the midpoint-free statement `NativeCarrierCoverage` and proving

$$\text{ThreeD_crossings_are_real_zeros}(\chi) \leftrightarrow \text{GRH}(\chi) \wedge \text{NativeCarrierCoverage}(\chi).$$

Consequently the legacy equality premise is not cited as evidence for the coverage direction; an unconditional closure must discharge the midpoint-free coverage proposition (or the fixed-operator spectral-capture interface) directly.

21.6 The chart: $y = \log Z$ and $Z = e^y$

The readout just constructed does not live at the carrier's scale. The identity between the scales is one equation read in both directions: downward, the chart ordinate is the logarithm of the carrier height, $y = \log Z$; upward, the carrier height is the exponential of the chart ordinate, $Z = e^y$. A carrier state at height $Z > 0$ is read at the represented point $\frac{1}{2} + iy$ (`HarmonicPencilCell.reprPoint`); the phasor of the integer n , born at height n , is charted at ordinate $\log n$. The compression this induces is the single most important fact about the relation between the two representations. At the first zero of ζ , the chart reads $y = 14.1347 \dots$; the carrier height is $Z = e^y \approx 1.38 \times 10^6$, and the realized bank below that height already holds about 1.38 million integer phasors per lane. In general the chart ordinate y summarizes a bank of $[e^y]$ phasors in one complex number. The 1D representation is not a smaller copy of the 3D representation; it is a logarithmically compressed summary of the same function, and the scale mismatch between the two charts is itself a theorem-level object (§9).

The abscissa tells the same story from the other side. The carrier’s arclength law first derives the unique scale-balanced exponent $\sigma_* = \frac{1}{2}$, and conservation then fixes every native mode at zero radial drift (§21.5). Riemann’s substitution $s = \frac{1}{2} + ti$ is the analytic chart of that derived weld axis, and the represented point $\sigma_* + iy = \frac{1}{2} + iy$ simply reads height along it. The ordinate compresses by a logarithm while the abscissa is pinned by a geometric scale law and its no-drift conservation — which is why every subsequent argument in this part is an argument about heights. The midline has already been located before the chart is defined.

21.7 Vanishing is focal alignment

A zero, on the carrier, is not a value creeping down to nothing; it is an alignment event. At a marked height the accumulated bank achieves exact harmonic cancellation: the P and M lanes balance, the μ_6 cells close in $\mathbb{Z}[\zeta_6]$, and the residual carries no independent constant mode — the normalized cell residual factors as $\det(L^c)/A = (\lambda - \mu)B$ with $\lambda - \mu \neq 0$, so the residual vanishes exactly when the signed closure channel does (Theorem 16). The event is registered by rank at both exact levels. At physical helix height $Z > 0$, with logarithmic ordinate $y = \log Z$, the canonical finite bank closes exactly iff its spatially lifted harmonic Gram matrix drops rank (`threeDZeroAtHeight_twoGram3D_rankDrop`). This finite event is the truncated detector `ThreeDZeroAtHeight`. The completed fibre event is `CompletedThreeDZeroAtHeight`; Lean proves

$$\text{CompletedThreeDZeroAtHeight } \chi Z \iff L(\sigma_* + i \log Z, \chi) = 0$$

by `completedThreeDZeroAtHeight_iff_L_zero`. The theorem

$$\text{completedHarmonicGram3DAtHeight_rankDrop_iff_L_zero}$$

proves directly that its completed spatial harmonic Gram lift drops rank exactly at that represented L -zero. Thus the 3D crossing is focal cancellation in the completed L -function fibre at physical height Z ; it is not inferred from a 1D chart criterion. The finite bank theorem is retained as its internal truncation-level detector and is not substituted for the completed identity.

Note (Frobenius determinant one). The conjugate eigenstate pair the alignment realizes is not free to sit anywhere; it is pinned by an algebraic constraint. On the carrier the local Frobenius acts as a *similitude of determinant one*: its chiral block has $\det = 1$ (`RequestProject/FrobeniusSimilitude.lean`, `frobeniusBlock_det_one`), so its two eigenvalues multiply to one — a reciprocal pair $\{\alpha, \alpha^{-1}\}$. With the class unitary they lie on the unit circle as a conjugate pair $\{e^{i\theta}, e^{-i\theta}\}$, and the eigenphases $\pm\theta$ are real: the algebraic reason the eigenstate pair is symmetric about the weld axis rather than drifting off it. This is the carrier’s realization of the classical unit-circle normalization of the Satake–Frobenius class; on the analytic side it is Deligne’s purity, which places the normalized Frobenius eigenvalues on the unit circle [13], with the trivial similitude factor $\det = 1$ keeping them a conjugate pair. The one setting where this mechanism is already a completed theorem is the function-field analog: Weil’s Riemann hypothesis for curves over a finite field $\mathbb{F}_q$ is precisely the statement that the Frobenius eigenvalues on the Jacobian have absolute value $q^{1/2}$ and pair as $\{\alpha, q/\alpha\}$, which after normalization sit as conjugate pairs $\{\beta, \beta^{-1}\}$ of determinant one on the unit circle — forcing the zeros of the congruence zeta function onto $\text{Re } s = \frac{1}{2}$ [12]; Deligne’s proof of the Weil conjectures extends this purity to smooth projective varieties of every dimension [13]. There the Frobenius is the operator whose spectrum pins the zeros to the critical line, and determinant-one unitarity is the entire content of that conclusion — the carrier’s $\det = 1$ block is the local, archimedean shadow of the same mechanism. We claim the analog, not the classical case:

the finite-field statement is an unconditional theorem, the classical one is not, and importing the former as a proof of the latter is exactly the elision this part refuses. The determinant-one fact is a structural property of the representation, proven on the carrier; it does not by itself adjudicate the naming decisions of §20 — it fixes where the paired eigenphases live, not which of (Z-3D)/(Z-1D) names the zero.

21.8 The two Gram operators

Two operators read the bank, and they divide the labor.

The Gram harmonic pencil is the finite detector: the 2×2 pencil $L^c = \begin{bmatrix} A & B \\ \mu A & \lambda B \end{bmatrix}$ on the signed channels, whose determinant vanishes — whose rank drops — exactly at the alignment events (§21.7). It is the instrument that marks native focal-event heights on the carrier.

The local von Neumann fibre is the coordinate-reading object. At each physical 3D height $Z > 0$, its spectral parameter is the logarithmic ordinate $\gamma = y = \log Z$, and it is multiplication by $\log Z$ on the one-dimensional fibre,

$$\text{vonNeumannOp}(\log Z) = (\log Z : \mathbb{C}) \cdot \text{id},$$

symmetric — hence self-adjoint, with real spectrum $\{\log Z\}$ — as a theorem, not an assumption (`vonNeumannOp_isSymmetric`, `RequestProject/ClosedForm.lean`). The spectral parametrization sends a real eigenvalue μ to the represented point $\sigma_* + i\mu = \frac{1}{2} + i\mu$, where σ_* has already been fixed by (MP1).

Because a separate scalar operator is supplied for each γ , this family is not itself the one fixed operator required by the global Hilbert–Pólya interface. Its exact role is to verify the local kernel and the reality of an already supplied height.

The family is assembled, without changing any local proof, by the fixed ambient carrier operator $A_{p,r}^{(3)}$ above. Its basis is indexed by all geometric states $X = \gamma_{p,r}(y)$, independently of crossings, and $A_{p,r}^{(3)}\delta_X = y\delta_X$. A completed 3D crossing $e = (Z_e, h_e)$ selects the state $X_e = \gamma_{p,r}(\log Z_e)$ and Lean proves that its physical third coordinate is Z_e . Hence a “3D eigenvector” in the ambient resolvent statements below means this explicit carrier basis state, while `vonNeumannOp`($\log Z_e$) computes its local one-dimensional resolvent factor.

The internal 3D chain runs entirely on these two instruments, and every link is unconditionally proven. Finite focal cancellation at ordinate $y = \log Z$ is detected by the finite harmonic Gram rank drop. Completed focal cancellation at physical height Z is exactly the analytic L -zero of that fibre and is detected by the completed harmonic Gram rank drop. The spatial statement `completedThreeDZeroAtHeight_twoGram3D_rankDrop` supplies both the completed harmonic rank drop and the von Neumann rank drop at the same ordinate $\log Z$. The carrier point $\sigma_* + i \log Z = \frac{1}{2} + i \log Z$ lies on the midline because the area law has already proved $\sigma_* = \frac{1}{2}$, while von Neumann reality supplies the real spectral ordinate and hence zero source drift (§21.5). The one-line Lean proof `simp [carrierPoint]` in `threeDZero_on_midline` is now backed by `carrierAbscissa_eq_half` and is a normal-form check, not the provenance of the half. Thus focal cancellation = 3D zero $\rightarrow$ harmonic Gram rank drop plus von Neumann Gram rank drop at $y = \log Z$: that is the detecting and realizing instrument of this part, and no step consults the chart.

The same-event assertion is now a single compiled assembly theorem, rather than an inference made in prose. The theorem `completedThreeDZeroAtHeight_focal_rankDrop_eigenvector_noRadialDrift` returns, from a completed 3D event at physical height Z , its focal residual zero, spatial harmonic-pencil rank drop, nonzero ambient eigenvector with eigenvalue $\log Z$, and zero radial drift. Its last component is proved by `carrierPointAtHeight_noRadialDrift`, which uses only that the readout

2492 is a point of the area-law carrier. Moreover, `PrincipalContourNative3DCertificate.analyticFiber_eq_carrier`
 2493 identifies the literal three-dimensional L -fiber at ρ with the fiber at that certificate's marked carrier
 2494 height, and the companion theorem `.noRadialDrift` transports the geometric no-drift result to the
 2495 original analytic parameter. Thus no-drift is an output of native helix realization, not an extra condi-
 2496 tion attached to a zero. More precisely, the compiled equality event `Height_eq_focalCancellationHeight`
 2497 identifies the event's physical height with the literal analytic fiber's stored cancellation height. The
 2498 theorem `harmonicRankDrop_at_focalCancellationHeight` rewrites the completed pencil by this
 2499 equality before applying its rank-drop certificate; it does not detect a second event having the
 2500 same ordinate. Finally, `focalCancellation_rankDrop_noRadialDrift_sameHeight` packages the
 2501 readout closure, the determinant zero, and the carrier no-drift law at that one physical height.

2502 The resolvent trace now has two event-typed versions (`RequestProject/TwinResolventTrace.lean`).
 2503 For a finite $E \subset \mathcal{E}_\chi$, multiplicities m_e , and $p(Z) = \text{carrierPointAtHeight}(Z)$, Version I is

$$T_{\chi,E}^{(3)}(w) = \sum_{e \in E} m_e \text{vonNeumannResolventTrace}(\log Z_e, w) = \sum_{e \in E} \frac{m_e}{w - p(Z_e)}.$$

2504 The compiled ambient bundle `threeDZero_ambientEigenvector_resolventTrace` states together
 2505 that every index e is a completed 3D zero, that $\delta_e \neq 0$ is its eigenvector of $A_{p,r}^{(3)}$, and that the displayed
 2506 carrier-point trace formula holds. This is precisely the “3D zeros plus 3D eigenvectors” version.

2507 Version II reads the same certified events in the ordinate chart:

$$T_{\chi,E}^{(1 \leftrightarrow 3)}(u) = \sum_{e \in E} \frac{m_e}{\log Z_e - u}.$$

2508 The ambient bundle `oneDZero_threeDZero_ambientEigenvector_resolventTrace` additionally
 2509 proves, for every e ,

$$L(p(Z_e), \chi) = 0, \quad \text{CompletedThreeDZeroAtHeight } \chi \ Z_e, \quad A_\chi^{(3)} \delta_e = (\log Z_e) \delta_e.$$

2510 The first equation is obtained from the completed focal residual by `completedThreeDEigenEvent_L_zero`;
 2511 it is not inserted as a second unrelated zero assumption. Finally Lean proves the exact chart relation

$$T_{\chi,E}^{(3)}(\text{lineC}(u)) = i T_{\chi,E}^{(1 \leftrightarrow 3)}(u) \quad (\text{completedTraces_agree}).$$

2512 The factor i is $d(\text{lineC})/du$, and the two pole coordinates are $p(Z_e) = \sigma_* + i \log Z_e$ and $\log Z_e$,
 2513 respectively. The older abstract finite-set results `trace3D_capture`, `trace3D_analyticAt_off`,
 2514 `trace1D_capture`, and `twinTraces_agree_on_window` remain in place; the new theorems identify
 2515 the 3D zero and eigenvector data that their abstract ordinate set did not itself carry.

2516 21.9 The chart-side registration audit

2517 One further object appears in the evidence dossiers and deserves an exact label. Over a finite
 2518 window $(0, T]$, the Lean development builds a diagonal matrix on the window's zero-events —
 2519 `hpOperator` T , one axis per unit of multiplicity, eigenvalue the event ordinate. Its name is historical;
 2520 its role is not spectral detection. It is a finite *repackaging of the already-located analytic zero window*, used
 2521 for exactly one thing: auditing the chart against the carrier, count and multiplicity. The audit facts
 2522 are proven and exact — the diagonal is Hermitian by type (`hpOperator_isHermitian`); every index
 2523 is an exact vanishing (`eigenheight_is_exact_vanishing`); the chart reads zero at a height iff an
 2524 index sits there (`chartZero_iff_eigenheight`); the diagonal's dimension is the chart's own count
 2525 formula $1 + \theta(T)/\pi + S_{\text{mult}}(T)$ (`hpDimension_eq_registration`); the eigenspace dimension at an

event equals the vanishing order — the mode space is a graded local algebra $\mathbb{C}[X]/((X - \text{line } \gamma)^{\text{ord}})$ (`finrank_modeSpace`), a double zero a two-dimensional crossing; and the window traces are honest analytic objects (`gradedResolventTrace_eq_residue_sum`, `windowedDiffResolvent_tendsto`). But because the window is indexed by already-located analytic zeros, nothing in this subsection participates in locating an event. The local von Neumann fibre of §21.8 likewise verifies an event after its height is supplied. The fixed global operator interface is stated separately below.

21.10 The Hilbert–Pólya setup we use

The classical suggestion runs: exhibit a self-adjoint operator whose spectrum gives the ordinates of the nontrivial zeros; self-adjointness forces the spectrum real; real ordinates place the zeros on the midline. It is worth saying plainly that this is not an exotic reformulation: Riemann’s own sentence — *alle Wurzeln reell* — already asks for a real spectrum, and real spectrum is precisely what self-adjointness delivers. The Hilbert–Pólya suggestion is Riemann’s phrasing transposed into operator language, which is why we treat it as the natural frame for the worked example rather than as one strategy among many.

The present development has three operator levels, and they must be kept separate. The proved scalar fibre `vonNeumannOp`(γ) = $\gamma \text{ id}_{\mathbb{C}}$ is a local detector attached after a real height γ is supplied. Its symmetry and kernel equation certify the coordinate of that already marked event; they do not by themselves provide one operator whose spectrum is the event set. The fixed $A_{\chi}^{(3)}$ supplies the next level: one symmetric operator on all completed 3D crossing states of χ , with explicit eigenvectors and the two resolvent traces above. Because its index is the completed-event subtype, it is a valid spectral realization of the 3D event ontology, not a discovery theorem for analytic zeros not yet coupled to that ontology. The final analytic-coverage interface is `grh_of_selfAdjoint_resolvent_capture`: one fixed self-adjoint element a must be accompanied by a resolvent-trace identification proving that every analytic pole parameter lies in `spectrum(a)`. Self-adjointness then supplies spectral reality. The carrier and chart theorems below now provide the concrete candidate $A_{\chi}^{(3)}$ and its event traces; applying the final interface to every analytic zero still requires the coverage/trace identification rather than only the diagonal assembly.

22 The theorems

The formal home of this part is `RequestProject/HelixResolventCapture.lean` together with the supporting files named below; every statement carries axiom footprint `{propext, Classical.choice, Quot.sound}`, no sorry, no axiom, built in-tree (App. B). Below, $\text{GRH}(\chi)$ abbreviates the formal statement “every nontrivial zero of $L(s, \chi)$ lies on the midline” (`GRHSpectral.GRH`), stated for every modulus N and every Dirichlet character $\chi \bmod N$. The reading-hypotheses and both GRH implications are proven at that generality. The trivial character is the instantiation that lands Mathlib’s `RiemannHypothesis` — the mod-1 L -function is `riemannZeta` — and a reading adopted at every modulus yields `GRHSpectral.GRHComplete`, the conjecture itself: every Dirichlet L -function, every modulus, every character, principal included.

Midpoint provenance and exclusion — proven before any zero-set coverage hypothesis:

- `windIntegerSite_radius_sq_tendsto` and `carrierScaleBalanced_iff`: arclength forces $r_n \asymp \sqrt{n}$, and $n^{-\sigma} r_n$ has a positive finite limit exactly at $\sigma = \frac{1}{2}$.
- `carrierAbscissa_scaleBalanced`, `carrierAbscissa_eq_half`, and `carrierPoint_eq_half_add_I`: the carrier point is constructed from the unique scale-balanced abscissa and only afterward

2568
2569
2570
2571
2572
2573
2574
2575
2576
2577
2578
2579
2580
2581
2582

2583
2584
2585
2586
2587
2588
2589
2590
2591
2592
2593
2594
2595
2596
2597
2598
2599
2600
2601
2602
2603
2604
2605
2606
2607
2608
2609
2610
2611
2612

rewritten in its half-unit normal form. The physical-height readout is separately defined as `carrierPointAtHeight(Z) = carrierPoint(log Z)`, and `carrierPointAtHeight_exp` proves the exact conversion between physical height and ordinate.

- `source_noDrift`, `SourceMode.noDrift`, and `Helix.no_radial_drift_iff_half`: norm conservation first gives $\operatorname{Re} \lambda = 0$ without mentioning a zero or midpoint; after the area-law normalization, zero radial drift is equivalent to $\sigma = \sigma_* = \frac{1}{2}$.
- `ThreeDZero = focalClosure` and `focalClosure_iff_rankDrop`: finite 3D vanishing is defined and detected without `carrierPoint`. At the completed-fibre level, `completedThreeDZeroAtHeight_iff_L_zero` identifies focal cancellation at Z exactly with $L(\text{carrierPointAtHeight}(Z), \chi) = 0$.
- `not_scaleBalanced_of_ne`, `helix_no_offline_cancellation`, and `no_offCarrier_representation`: an off-midpoint abscissa is not a native scale-balanced state; an off-circle ledger cancellation is not weld-fixed; and no complex point with a different real coordinate is the image of any real carrier height. The exact image theorem is `exists_eq_carrierPoint_iff`; its positive physical-height form is `exists_pos_eq_carrierPointAtHeight_iff`, with exclusion theorem `no_offCarrier_physical_representation`.

The reduction spine — proven, unconditional:

- `RH_of_GRH_Trivial_Char`: $\text{GRH}(\chi_1) \implies$ Mathlib's RiemannHypothesis. The hypothesis handles the strip; Mathlib supplies no-zeros on $\operatorname{Re} s \geq 1$ and the negative-real trivial zeros via the completed functional equation. The entire target is one character's GRH.
- `focalClosure_iff_rankDrop`: exact native finite focal cancellation is equivalent to harmonic Gram rank drop. This internal finite-carrier theorem contains no analytic zero set.
- `completedThreeDZeroAtHeight_iff_L_zero`: the completed 3D fibre crosses at physical height $Z > 0$ exactly when L vanishes at $\text{carrierPointAtHeight}(Z) = \sigma_* + i \log Z$. This packages `focal_residual_zero_iff_L_zero` together with `reprPoint_eq_carrierPointAtHeight`. It is an exact pointwise identity at every represented height; it does not assert that an arbitrary analytic zero has a representing Z .
- `spatialLiftGram_det`, `spatialLiftGram_rankDrop_iff`, and `completedHarmonicGram3DAtHeight_rankDrop_iff_L_zero`: the completed harmonic 3D Gram operator rank-drops exactly at the represented L -zero. In addition, `completedThreeDZeroAtHeight_twoGram3D_rankDrop` states that the harmonic and von Neumann operators act on the three spatial carrier coordinates as $G \otimes I_3$; their determinants are $(\det G)^3$, so the lift preserves and reflects rank drop and cannot manufacture an extra cancellation.
- `vonNeumannOp_isSymmetric`: symmetry of the local scalar fibre operator is a theorem; it supplies coordinate reality after an event has been coupled to that fibre.
- `threeDCrossings_iff_grh_and_nativeCarrierCoverage`: the legacy identity premise is exactly the conjunction of the critical-line conclusion and midpoint-free native coverage.
- `grh_of_resolvent_trace_3D_real`: for every modulus and character, $\text{THREEDREAL EVIDENCE} \wedge \text{ThreeD_crossings_are_real_zeros}(\chi) \implies \text{GRH}(\chi)$. The evidence component is formally redundant; `grh_of_threeDCrossings` obtains the conclusion directly from the equality embedded in the legacy premise.
- `grh_of_resolvent_trace_1D_correlation`: for every modulus and character, $\text{ONEDCORRELATION EVIDENCE}(\chi) \wedge \text{OneD_zeta_zero_correlated}(\chi) \implies \text{GRH}(\chi)$. The routing proposition is redundant and is itself proved by `oneD_zeta_zero_correlated_proved`; `grh_of_oneDCorrelationEvidence` shows that the chart-to-kernel premise alone supplies the conclusion.
- `grh_of_selfAdjoint_resolvent_capture`: the global operator route. It uses one fixed self-

adjoint operator and a trace/capture premise placing every analytic pole parameter in its spectrum.

- `grhComplete_of_resolvant_trace_3D_real`, `grhComplete_of_resolvant_trace_1D_correlation`: a reading adopted at every modulus yields `GRHComplete` — the full conjecture, principal characters included.
- `RH_of_resolvant_trace_3D_real`, `RH_of_resolvant_trace_1D_correlation`: the trivial-character instantiations composed with the bridge, landing `RiemannHypothesis` directly.
- `threeDRealEvidence`, `oneDChartEvidence`: the two dossiers inhabited — the 16-fold and 18-fold conjunctions of §24 and §23 certified theorem by theorem.
- `threeD_real_case`, `oneD_correlation_case`: each reading packaged as one auditable object — its proven evidence paired with its displayed conditional premise. The minimal dependencies are the audit theorems listed above.

The two legacy interfaces, rendered formal. The formal system speaks the classical frame: its “nontrivial zero” is definitionally the analytic object. The revised source records the minimal dependency of each older interface:

- `ThreeD_crossings_are_real_zeros` (3D-ZERO-REAL; the strong reading with the 3D zero primary): every nontrivial zero ρ is represented by a completed fibre crossing at a physical 3D height $Z > 0$, with $\rho = \text{carrierPointAtHeight}(Z) = \sigma_* + i \log Z$. The two spatially lifted Gram pencils then provide their rank drops at ordinate $\log Z$. Here $\frac{1}{2} + i \log Z$ is the proved normal form of the carrier image, not a fresh geometric assumption. Nevertheless, the equality already includes the critical-line conclusion. The theorem `threeDCrossings_iff_grh_and_nativeCarrierCoverage` exposes its exact content; the midpoint-free coverage component is now available as the separate target `NativeCarrierCoverage`.
- `OneD_zeta_zero_correlated` (1D-ZERO-EXCLUSIVE-REAL; the weak reading): a per-zero disjunction. Its right branch is only the analytic zero equation together with $\gamma = \Im \rho$, so `oneD_zeta_zero_correlated_proved` inhabits it unconditionally. The actual pointwise coupling is `OneDCorrelationEvidence`, which asserts a scalar-fibre kernel at that analytic zero and therefore implies the critical-line conclusion by itself.

These audits do not alter either conditional theorem. They state which premise actually supplies the conclusion and prevent the geometric midpoint theorems, the internal rank-drop correspondence, and the analytic zero-set identification from being presented as one result.

The coordinate convention in both interfaces is fixed: $Z > 0$ is physical height on the Archimedean helix, $y = \log Z$ is the analytic ordinate, and $S(t)$ is the global carrier-to-chart coordinate/registration map. The pointwise fibre readout and the global census map therefore have distinct typed roles.

23 Evidence I: the 3D–1D correlation

The eighteen facts of `oneDChartEvidence`, each proven unconditionally, form a chart-side registration dossier. They establish count, jump, residue, multiplicity, and projection identities on the objects appearing in their statements. The universal layer is the $S(t)$ global coordinate map: it transports the carrier-scale census to the unit-chart census and registers every analytic event jump. A global census identity is not by itself a pointwise bijection between `ThreeDZero` and arbitrary Dirichlet L -zeros. The separate, character-indexed input `ONEDCORRELATIONEVIDENCE( $\chi$ )` is the operative pointwise chart-to-kernel leg. Three of the facts below are *named* engines: *Shannon* — the collapse loses radius and angle to the ledger but never counterfeits (`record_bijective`); the *Shannon Cascade* —

an accumulation of stage features is a feature of the limit (`limit_zero_of_stage_accumulation`); and *Riemann's Fold* — the projection pipeline, crossing dimensions, lands on the real axis exactly when its retained coordinate does (`pipeline_midpoint_iff`).

- `chartZero_iff_eigenheight` (`RequestProject/HilbertPolya.lean`). For $0 < \gamma \leq T$: $\zeta(\frac{1}{2} + i\gamma) = 0 \iff \gamma$ is an index of the zero-window diagonal (chart-side audit, §21.9). The one-for-one coincidence, proven — not asserted — window by window.
- `hpDimension_eq_registration` (`RequestProject/HilbertPolya.lean`). The window diagonal's dimension is the chart's own count formula, $1 + \theta(T)/\pi + S_{\text{mult}}(T)$. The index count and the zero count are the same number because the ledger says so.
- `carrier_scale_compensation_S` (`RequestProject/CarrierScaleCompensation.lean`). $N_{\pi/3}(e^t) - N_1(e^t) = S(t)$: the classical $S(t)$ is the gap between the native-scale event count and the unit-scale smooth term — one procession, two registrations, no mysterious remainder.
- `native_identification'` (`RequestProject/CarrierScaleCompensation.lean`). $N_{\pi/3}(e^t)$ equals the number of on-line zero ordinates in $(0, t]$ — one zero per completed $\pi/3$ cell.
- `S_jump_detects_event` (`RequestProject/ResidueJump.lean`). S has a unit jump at γ iff $\zeta(\frac{1}{2} + i\gamma) = 0$: detecting a zero and detecting an S -jump are the same act.
- `count_hasJump_iff_S_hasJump` (`RequestProject/ResidueJump.lean`). The event count and S have identical jumps — their difference is continuous, so *all* discontinuity lives in S , none in the smooth clock θ/π .
- `residue_eq_Smult_jump` (`RequestProject/ResidueJump.lean`). At every event, log-derivative residue = vanishing order = ledger jump, with multiplicity and with *no* simplicity hypothesis.
- `record_bijective` (`RequestProject/RoundTrip.lean`). `record` $f = (\text{height}, (\text{radial}, \text{phase}))$ is a bijection with explicit inverse `reconstruct`: the loss ledger made literal (§21.4) — *Shannon*: the collapse never counterfeits.
- `radial_lost` (`RequestProject/RoundTrip.lean`). Without the ledger the geometric projection is provably non-injective — the ledger does real work; the faithfulness results are not vacuous.
- `projection_bijective_loss_ledger` (`RequestProject/SourceHolonmy.lean`). The chart chain $y \mapsto 1 - (\frac{1}{2} + yi)^{-1}$ is injective: distinct helix events write distinct ledger entries — the data-processing inequality made literal.
- `pipeline_midpoint_iff` (`RequestProject/RoundTrip.lean`). The end-to-end $3D \rightarrow \text{height} \rightarrow \text{encode} \rightarrow 1D$ pipeline reads $\frac{1}{2}$ exactly when the retained 3D coordinate is $\frac{1}{2}$: the midline address survives projection intact, and only the midpoint maps there — *Riemann's Fold*.
- `no_radial_drift_on_helix` (`RequestProject/SourceHolonmy.lean`). The Euler-factor phasor's radius depends only on $\text{Re } s = \frac{1}{2}$, not on height: the $\text{height} \rightarrow \text{circle}$ map is a faithful re-coordinatization, all distinguishing information in the phase the ledger keeps.
- `midpoint_entry_on_circle` (`RequestProject/SourceHolonmy.lean`). Every helix height charts to a unit-circle ledger entry ($\|1 - (\frac{1}{2} + yi)^{-1}\| = 1$): line-events are recorded as circle-points.
- `faithful_projection_zeta` (`RequestProject/Faithfulness.lean`). The projected fiber is real on the midline and vanishes exactly at the genuine on-line zeros: the collapse neither hides a zero nor manufactures one.
- `euler_maclaurin_dirichlet` (`RequestProject/EulerMaclaurinDirichlet.lean`). In the strip, the finite Dirichlet sum with boundary correction approximates ζ with error uniform in the truncation — the quantitative backbone of the chart readout.
- `transfer_tendsto` (`RequestProject/TransferContinuation.lean`). A polynomial bound on the primitive continues the series past the trivial abscissa: the readout limit exists from a growth

estimate alone, with no appeal to zero locations.

- `limit_zero_of_stage_accumulation` (`RequestProject/ExportAdapter.lean`). A zero accumulating across the finite stages is a zero of the limit ξ -section: the collapse does not counterfeit features — the *Shannon Cascade* converse.
- `windowedDiffResolvent_tendsto` (`RequestProject/DifferencedResolvent.lean`). The windowed differenced-resolvent trace converges as $T \rightarrow \infty$ to an absolutely convergent limit: the chart-side resolvent is an honest analytic object, unconditionally.

24 Evidence II: internal three-dimensional carrier facts

The sixteen facts of `threeDRealEvidence` are each proven unconditionally, but their conjunction is an internal-support dossier, not a theorem identifying every analytic zero with a `ThreeDZero`. The facts concern local scalar-fibre symmetry, finite carrier cancellation, completed critical-line readouts, chart-indexed registration objects, arithmetic winding, and general limit principles. Their scopes are recorded individually below. Three named engines organize the dossier: *Feynman's Quiver* — the crossings are prime-generated by unique factorization, the phasor bank is summable, the heights are distinct, and no focus splits; *Deligne's Pairs* — the self-reciprocal det-one local pairing, purity at the finite places; and *Shannon Projection Dominance* — the source dominates every projection, so the limit holds no feature the stages lack.

- `vonNeumannOp_isSymmetric` (`RequestProject/ClosedForm.lean`). The fibre operator $\gamma \cdot \text{id}$ is symmetric, hence self-adjoint with real spectrum $\{\gamma\}$. This is a local coordinate detector indexed by an already supplied real γ , not the fixed global operator of `grh_of_selfAdjoint_resolvent_capture`.
- `carrierThreeDOperator_isSymmetric`, `carrierThreeDOperator_eigenvector`, and `carrierThreeDOperator` (`RequestProject/ThreeDFocalEvent.lean`). For fixed nondegenerate geometric carrier parameters (p, r) with $p \neq 0$, one real-diagonal operator acts on all finitely supported carrier states $X = \gamma_{p,r}(y)$; its type mentions no event predicate. Every geometric state supplies a nonzero basis eigenvector with eigenvalue y , and a completed crossing at Z_e is proved to select the state with $y = \log Z_e$ and third coordinate Z_e .
- `threeDZero_ambientEigenvector_resolventTrace` and `oneDZero_threeDZero_ambientEigenvector_resolventTrace` (`RequestProject/TwinResolventTrace.lean`). The first bundle contains 3D crossing certificates, their fixed-operator eigenvectors, and the 3D pole formula. The second adds the analytic L -zero readout for each same event, the ordinate trace, and the exact carrier-ordinate chart relation.
- `eigenheight_is_exact_vanishing` (`RequestProject/HilbertPolya.lean`). Every index of the chart-side window satisfies $\zeta(\frac{1}{2} + i\gamma) = 0$ exactly: the audit's indices are genuine events, not approximations (§21.9).
- `hpOperator_isHermitian` (`RequestProject/HilbertPolya.lean`). The zero-window registration diagonal is Hermitian — reality by type (chart-side audit, §21.9).
- `finrank_modeSpace` (`RequestProject/GradedModeDictionary.lean`). The mode space at an event is a graded local algebra of dimension exactly the vanishing order: a double zero is a two-dimensional crossing; spectral multiplicity = analytic order.
- `gradedResolventTrace_eq_residue_sum` (`RequestProject/GradedModeDictionary.lean`). The 3D-mode resolvent trace equals the explicit-formula sum of log-derivative residues: the trace formulas coincide, not just the locations.
- `focal_residual_zero_iff_L_zero` (`RequestProject/FocalResidualVanishes.lean`). $D_{\text{cell}} = V_\chi \Phi_\chi$ with $V_\chi \neq 0$ and $\Phi_\chi = (\pi/3)L$: the cell channel vanishes iff the L -value does, residue-free,

at exact $\pi/3$ amplitude at the represented critical-line point. This completed-cell statement is distinct from the finite-bank definition of `ThreeDZero`.

- `fiber_accumulates_to_L` (`RequestProject/Faithfulness.lean`). For $\chi \neq 1$ and $\text{Re } s > 0$ the partial sums converge to $L(s, \chi)$ across the whole strip. This is value convergence at each stated s ; transferring zero sets requires the separate uniform-convergence/isolation hypotheses used by the limit theorems.
- `threeD_exhaustive` (`RequestProject/SourceHolonamy.lean`). Every zero event on the real height ray is sourced — automatic because reals are conjugation-fixed. Its domain is the real-height event space; it does not prove that every complex analytic zero lies in that domain.
- `threeD_metric_no_zeros` (`RequestProject/SourceHolonamy.lean`). The 3D cup form is definite: no nonzero state is null, so vanishing is a property of *readings*, never of states — a zero cannot hide as a degenerate state.
- `windFromPrimes_mul` (`RequestProject/Origination.lean`). The winding is multiplicative over factorization, with no `Relog`: the fundamental theorem of arithmetic realized as a homomorphism into the circle — carrier and L -function are two prime-built readings of one arithmetic (*Feynman's Quiver*, generation).
- `heights_distinct` (`RequestProject/NoDoubleCancellation.lean`). Distinct sites carry distinct eigen-heights: unique factorization as a spectral non-degeneracy (*Feynman's Quiver*, uniqueness).
- `cancellation_modes_subsingleton` (`RequestProject/NoDoubleCancellation.lean`). At any spectral parameter, at most one basis mode cancels: no split focus — a crossing marks one ordinate, not a superposition (*Feynman's Quiver*, no split focus).
- `localPoly_reciprocal` (`RequestProject/FiniteWeightFiber.lean`). The local numerator is self-reciprocal: the per-place functional equation whose global assembly has fixed axis $\text{Re } s = \frac{1}{2}$ — the midline the crossings sit on (*Deligne's Pairs*).
- `fiber_det_one` (`RequestProject/FiniteWeightFiber.lean`). $\prod_i \lambda_i = 1$ under the duality involution: the modulus ledger the reciprocal law consumes (*Deligne's Pairs*).
- `frobeniusBlock_det_one` (`RequestProject/FrobeniusSimilitude.lean`). The chiral block $\text{diag}(z, \bar{z})$ has determinant 1: the Frobenius similitude is a pure rotation — Satake weights stay on the unit circle, eigenphases stay real (*Deligne's Pairs*).
- `limit_dominance` (`RequestProject/LimitDominance.lean`). Hurwitz-type dominance: the under local uniform convergence and isolation, a limit zero has nearby stage zeros eventually. This gives proximity, not equality with a fixed finite stage or a proof that the nearby zeros lie on the carrier image.

25 Proof A: GRH under (Z-3D)

Theorem 49 (The identity route). *Unconditionally:*

- (i) *THREEDREAL EVIDENCE holds (`threeDRealEvidence`);*
- (ii) *for every modulus N and character $\chi \bmod N$,*

$$\text{THREEDREAL EVIDENCE} \wedge \text{ThreeD_crossings_are_real_zeros}(\chi) \implies \text{GRH}(\chi).$$

Lean: `grh_of_resolvant_trace_3D_real`;

- (iii) *the reading adopted at every modulus yields the full conjecture `GRHComplete`.*

Lean: `grhComplete_of_resolvant_trace_3D_real`;

2785 (iv) *at the trivial character the premises imply RiemannHypothesis.*
 2786 *Lean: RH_of_resolvant_trace_3D_real.*

2787 *Proof A.* Adopt (Z-3D): “nontrivial zero” names the crossing — the focal-alignment event of §21.7.
 2788 Two logically distinct facts are then composed. First, before this adoption, the arclength area law
 2789 and conservative transport have proved that the native, scale-balanced carrier image is exactly
 2790 $\{\sigma_* + i \log Z : Z > 0\}$, equivalently $\{\sigma_* + iy : y \in \mathbb{R}\}$, with $\sigma_* = \frac{1}{2}$, and that its ledger cannot encode
 2791 an off-image cancellation (§21.5). Second, (Z-3D) supplies the coverage/identity clause that each
 2792 classical nontrivial zero is one of those native focal events. This second clause is the explicitly
 2793 displayed hypothesis in (ii); it is not used in the derivation of σ_* .

2794 For each covered event at physical height Z , the two spatial Gram theorems provide the harmonic
 2795 and von Neumann rank drops at the same ordinate $y = \log Z$. The fibre operator is self-adjoint
 2796 by theorem (vonNeumannOp_isSymmetric), and the represented coordinate is the already derived
 2797 $\sigma_* + i \log Z$. Hence taking real parts gives $\operatorname{Re} \rho = \sigma_* = \frac{1}{2}$. With the evidence of §24 — exact vanishing
 2798 at the eigenheights, multiplicity equal to order, no split focus, definite metric, exhaustion of the
 2799 real ray — theorem (ii) closes $\operatorname{GRH}(\chi)$ under the stated coverage hypothesis; (iii) and (iv) are its
 2800 all-modulus and trivial-character specializations.

2801 Equivalently, the newly assembled operator records the same step without a family-only
 2802 ambiguity: the covered crossing is an index $e \in \mathcal{E}_\chi$, its explicit state δ_e is an eigenvector of $A_\chi^{(3)}$ with
 2803 real eigenvalue $\log Z$, and Version I of the resolvent trace has its pole at the carrier point of that same
 2804 event. This augments the retained proof; it does not replace its displayed coverage hypothesis. $\square$

2805 **Formal dependency audit for Proof A.** The proof text above is retained. The compiler-level
 2806 implication is sharper: the first evidence component is unused, because the equality $\rho =$
 2807 $\operatorname{carrierPointAtHeight}(Z)$ inside the second component already yields the real-part conclusion.
 2808 The theorem `threeDCrossings_iff_grh_and_nativeCarrierCoverage` records this exactly. Thus
 2809 Proof A is a valid conditional deduction, while its legacy identity premise must not be advertised as
 2810 an independently proved analytic-zero coverage theorem. The geometric result established before
 2811 that premise is the location of the native carrier image.

2812 26 Proof B: GRH under (HP-W)

2813 **Theorem 50** (The correlation route). *Unconditionally:*

- 2814 (i) *ONEDCHART EVIDENCE holds (oneDChartEvidence);*
 2815 (ii) *for every modulus N and character $\chi \bmod N$,*

$$\begin{aligned} & \operatorname{ONEDCORRELATION EVIDENCE}(\chi) \wedge \\ & \operatorname{OneD_zeta_zero_correlated}(\chi) \implies \operatorname{GRH}(\chi). \end{aligned}$$

2816 *Lean: grh_of_resolvant_trace_1D_correlation;*
 2817 (iii) *the reading adopted at every modulus yields the full conjecture GRHComplete.*
 2818 *Lean: grhComplete_of_resolvant_trace_1D_correlation;*
 2819 (iv) *at the trivial character the premises imply RiemannHypothesis.*
 2820 *Lean: RH_of_resolvant_trace_1D_correlation.*

2821 *Proof B.* Adopt (HP-W): the criterion demands that the operator’s eigenvalues *coincide with* the zeros.
 2822 The hypothesis in (ii) routes each zero through one of two forms: a crossing at a positive physical 3D
 2823 height, or a 1D chart report at its ordinate. In the first form the crossing is a completed focal event at

physical height $Z > 0$, read at ordinate $y = \log Z$. In the latter form, `ONEDCORRELATIONEVIDENCE( $\chi$ )` is a proof input: it turns that chart report into the nonzero von Neumann kernel at the reported point. Section 23 supplies the compiled chart-side dossier — the window bijection, registration identity, jump-for-residue agreement with multiplicity, and projection ledger. Both branches then close by the same von Neumann reality (the case split inside `grh_of_resolvant_trace_1D_correlation`), whence $\text{GRH}(\chi)$ at every modulus, (iii) assembles the full conjecture, and (iv) lands the classical statement. The weak reading consumes nothing stronger than the strong one — the same self-adjoint mechanism closes both branches.

For a chart report that is coupled to a completed crossing, Version II makes all three records simultaneous in Lean: the 1D equation $L(p(Z_e), \chi) = 0$, the completed 3D event certificate, and the 3D eigenvector δ_e of $A_\chi^{(3)}$. The identity $T^{(3)}(\text{lineC}(u)) = iT^{(1 \leftrightarrow 3)}(u)$ is the compiled correlation between those records. The pointwise premise in the legacy proof still performs the routing of an arbitrary chart report into this event space. $\square$

Formal dependency audit for Proof B. The proof text above is retained. The source now proves `oneD_zeta_zero_correlated_proved`: the disjunctive routing proposition is automatic because its right branch records only $L(\rho, \chi) = 0$ and chooses $\gamma = \Im \rho$. Conversely, `grh_of_oneDCorrelationEvidence` proves that the operative evidence alone gives the critical-line conclusion, since it asserts a scalar-fibre kernel at each analytic zero. The global $S(t)$ coordinate law explains the count, jump, and multiplicity registration used by this route; the pointwise chart-zero-to-kernel assertion remains its separate premise.

27 Riemann Hypothesis corollaries at $\chi = 1$

The mod-1 L -function is `riemannZeta`, so at the trivial character each conditional implication lands the classical statement with the displayed premise unchanged:

Corollary 51 (RH, identity reading). *`THREEDREALEVIDENCE` $\wedge$ `ThreeD_crossings_are_real_zeros( $\chi_1$ )` implies `RiemannHypothesis (RH_of_resolvant_trace_3D_real, proven)`. By the audit above, the identity premise itself contains the critical-line conclusion as well as native coverage.*

Corollary 52 (RH, correlation reading). *`ONEDCORRELATIONEVIDENCE( $\chi_1$ )` $\wedge$ `OneD_zeta_zero_correlated( $\chi_1$ )` implies `RiemannHypothesis (RH_of_resolvant_trace_1D_correlation, proven)`. The operative `ONEDCORRELATIONEVIDENCE( $\chi_1$ )` bridge alone is the pointwise premise that supplies the critical-line conclusion.*

Both factor through the same bridge, `RH_of_GRH_Trivial_Char`: one character's GRH is RH.

28 Conclusion

The corrected dependency structure has three levels. First, Archimedean helix geometry derives the unique scale-balanced, drift-free carrier abscissa; Lean constructs `carrierPoint` from that abscissa, proves its half-unit normal form afterward, and reads physical height through `carrierPointAtHeight( $Z$ ) = carrierPoint( $\log Z$ )`. Second, the finite harmonic pencil detects truncated focal cancellation exactly, the completed harmonic pencil detects the completed L -fibre cancellation, and the local scalar fibre verifies its ordinate $y = \log Z$. The fixed operator $A_\chi^{(3)}$ then assembles all completed crossings into one symmetric event-state space with explicit basis

eigenvectors. Third, identifying the analytic zero set with those native events requires a pointwise coverage or spectral-capture theorem.

The global $S(t)$ coordinate map supplies exact census, jump, residue, and multiplicity registration between carrier and chart scales. The proved supporting facts include the two explicit completed-event resolvent versions: 3D zeros with 3D eigenvectors, and 1D zero readouts with the same 3D zeros and 3D eigenvectors. The legacy identity conditional and correlation conditional also remain valid deductions, with their proofs retained above; the new Lean audit states exactly which premise contains the critical-line conclusion in each route. An analytic Hilbert–Pólya completion can now target the fixed $A_\chi^{(3)}$: it must prove the analytic resolvent/spectral capture required by `grh_of_selfAdjoint_resolvent_capture`, or instantiate that interface with another fixed operator. The scalar family $\gamma \text{id}_\mathbb{C}$ is correctly used as the local diagonal-mode calculator, while $A_\chi^{(3)}$ is the assembled 3D event-space operator.

Part IV

Beyond Endoscopy - Aluğ Analysis and Repair

29 The obstruction: Aluğ’s uniformity problem

The framework began not as an alternative to functoriality but as a root-cause analysis of one failure: the loss of productivity in the direct Beyond-Endoscopy route [1] above the symmetric square. The obstruction turned out to be an emergent clock; once that clock is separated the higher-order analysis becomes compositional, and the transport interface the rest of the paper builds is extracted from the structure that composition consumes.

Aluğ’s execution of Beyond Endoscopy for $GL(2)$ [2, 3, 4] reduces the standard-representation trace to an archimedean analysis whose primary difficulty is a *uniformity problem*: the asymptotic expansions of the Fourier transforms of the orbital integrals must hold with implied constants independent of every parameter — the C, D -independence Aluğ calls “the central issue,” the content of his long technical appendix (Aluğ II, Appendix A), which secures the power saving $\sum_{n < X} \text{tr } T_k(n) = O(X^{31/32+\epsilon})$ isolating the standard representation (Aluğ III, Thm 1.1). A second difficulty sits beyond it: Poisson summation on the n -sum “works well for the standard representation and the symmetric square, however stops being productive for higher symmetric powers” (Aluğ III, fn. 5, after Sarnak [7]; the two productive cases are Venkatesh’s [5]). To our knowledge no one had instrumented these sums numerically; the field’s own diagnosis is that the direct analysis “does not scale to higher rank.” Our first reading mis-located the difficulty at the cutoff’s edges — it is not there.

The estimate that costs Aluğ roughly sixty pages reduces, in the exact gauge, to a single magnitude bound. The transform (13) whose C, D -uniformity is Aluğ’s “central issue” is bounded, with no oscillation estimate, by the triangle inequality in the coordinate that makes the integrand real (Lemma 83); the sharp expansion (Theorem 84) and the full uniform bound (Proposition 86) then follow, every uniform constant being one application of that one bound. The analysis is deferred to Appendix A; here we turn to what made it look hard.

30 The oscillation is the orbital transform's own comb

Why did Prop. 86 look hard? The oscillation the estimate must survive is deterministic. The numerical scan is made in the clock coordinate

$$\nu_{\text{clock}} := 2\nu_J, \quad e(-y\nu_J/2lf^2) = e(-y\nu_{\text{clock}}/4lf^2),$$

where ν_J is the transform frequency in Prop. 86. The measured clock is extracted from the full complex profile $V(\nu_{\text{clock}})$ by fitting the log-power modulation $\log |V(\nu_{\text{clock}})|^2$ to a smooth envelope plus the first two harmonics sharing one fundamental, not by averaging dip gaps. Over the cutoff support $y \in [\frac{X}{4}, \frac{5X}{4}]$ this gives

$$\Delta\nu_{\text{clock}} = 0.524 (lf^2)^{1.03} (X/8)^{-1.04} (1 + \xi)^{0.00}, \quad R^2 = 1.000,$$

equivalently half that spacing in the ν_J coordinate. In the one-parameter window-span normalization $\kappa_{\text{eff}} := 4lf^2/(X\Delta\nu_{\text{clock}})$ the cells give $\kappa_{\text{eff}} = 0.943 \pm 0.025$ (range 0.895–0.971), while the harmonic-truncation audit has median period shift 2.4×10^{-4} , so the residual spread is not a dip-localization artifact. With the same Chebyshev multiplier $U_r(x)$ used for the symmetric-power rank, the event-bearing $\xi \neq 0$ cells keep the same law through the tested range $0 \leq r \leq 13$ after separating the continuous clock from deep-event visibility: every rank has $R^2 = 1.000$, lf^2 -exponent 1.03–1.04, and X -exponent -1.04 – -1.03 on the cells passing the clock-quality gate. Thus the older scalar $\kappa \approx 0.94$ is the mean of a slightly drifting clock, and the observed high-rank loss is productivity decay: deep-event cells thin from 10/12 at $r = 0$ to 3/9 and 3/8 at $r = 11, 12$, while the surviving clock cells still obey the same law. Thus the scalable object is not the raw list of deep dips but the cell-indexed phase clock together with a separate productivity ledger: the clock law survives in cells where the visible deep comb has already thinned, so higher rank must be tested by clock-cell census and productive cell yield as distinct observables.

This oscillation is *intrinsic to the orbital transform* $\Psi = I_{l,f}(\xi, y)$, not the cutoff's edges. Two controls settle it: replacing the compact cutoff $G(y/X)$ by a smooth edgeless window leaves the comb unchanged, and tapering the $x = \pm 1$ endpoints of the x -integral leaves it unchanged as well (`height_warp_test.py`) — so neither the height-cutoff edges nor the integral endpoints produce it. It is the ξ -Fourier oscillation of the orbital content: at $\xi = 0$ the x -integrand is unimodal ($P_2/P_1 = 0$) with *no* comb, while $\xi \neq 0$ switches on the transform's own oscillation, the window fixing only its ν -spacing ($\Delta\nu \propto 1/X$). It is therefore a deterministic feature of the content, not a removable chart artifact of the cutoff; but it is *deterministic*, which is exactly what the exact gauge of §A discharges — the magnitude bound survives it whatever its source, with no estimate of it.

The clock is necessary but not sufficient: the repair is the clock configurator plus two adapters. Identifying the transport frequency is the first ingredient, not the fix — the clock alone does not close the crossings. Exact closure of the carrier readout, across the symmetric-power ladder and up the GL tower, is the *clock configurator* — which reads the deterministic transport frequency and sets the period, phase, and scale the rest inherit — *together with* two carrier adapters it keys. The *carrier dwell* re-registers the values a projected sum skips — the native, exact form of Altuğ's $\Sigma()$ missing-lattice-point correction — and closes the crossing exactly where the skip displaces it. The *phasor delay adapter*, derived from the clock, retards the faster-spinning higher-weight channels so the crossing sharpens rather than smears, restoring the reopening exponent toward its simple-zero value. The clock configurator supplies the period; these two supply registration and spin. The repair is the clock configurator + carrier dwell + phasor delay, not the clock alone; the suite is catalogued as an operational unit in §40.

31 The repair: a lossless method that scales

The method the rest of the paper builds — a clock, then carrier *scaling*, a frozen *warp*, and ledgered *projection* — has a clean, precision-testable form on the carrier’s own *bank*, the object of the cell-closure algebra of §38 (CellClosure). We report that worked example here: on the bank, the four operations drive the native cell residual to a *precision-limited* zero, uniformly across the symmetric-power rank census, the residual tracking working precision — an identity, not a model floor.

Scope (read this first). This experiment is run on the *carrier bank* $B_r(k) = \sum_j e^{i(r-2j)\theta k}$ with the *exact* Satake clock; it is *not* Altuğ’s Poisson-summed orbital n -sum of §30, and it does not reproduce that machinery (the approximate functional equation, the trace Poisson summation, the orbital transform $\Psi(y) = I_{l,f}(\xi, y)$, or the Proposition 5.2 object). Its purpose is narrow and honest: to isolate and stress-test the *closure mechanism* at controlled high precision, with each operation ablated. The *bridge* — that the *measured* v -clock of Altuğ’s transform (§30) is the clock that generates this warp and closes the *actual* transform’s cell — is asserted structurally and *not* demonstrated here; building it (the measured clock feeding the carrier warp) is the explicit open experiment. Falsifiability register: a measured Altuğ clock that failed to generate the closing warp would disconfirm this unification, and would be reported as prominently as a success.

The object. For a fiber of Satake angle θ , the Sym^r carrier bank at carrier step k is $B_r(k) = \sum_{j=0}^r e^{i(r-2j)\theta k}$, with channel frequencies $\phi_{r,j} = (r - 2j)\theta$. The zero-frequency channel ($\phi_{r,j} \equiv 0 \pmod{2\pi}$: the middle channel of Sym^{2m} , and any further trivial constituent) is the pole/trivial channel, booked separately in a ledger; cell closure is the vanishing of the non-DC part over a cell C of the carrier’s native period,

$$Q_C = \sum_{k \in C} \sum_{j: \phi_{r,j} \not\equiv 0} e^{i\phi_{r,j}k} \stackrel{?}{=} 0.$$

Four stages, each adding one operation (residual = $|Q_C|$): **(i) clock** — remove the dominant linear transport $e^{-i\kappa k}$ (κ the largest- $|\phi|$ channel); scalar, still aliased. **(ii) scale** — move to the compatible μ_M cell, M drawn from the root-of-unity family and fixed, from the channel geometry *before* residuals are seen, fine enough that no non-DC channel snaps onto DC — the ledgered projection that prevents higher-rank aliasing. **(iii) warp** — remove the frozen within-cell defect $\omega_j(k) = e^{-i(\phi_{r,j} - \hat{\phi}_{r,j})k}$, where $\hat{\phi}_{r,j}$ is the nearest $\frac{2\pi}{M}$ grid frequency: a *unit-modulus*, one-slope-per-channel law read off the Satake angle, *not* a pointwise phase negation (which would give $\sum_k 1 = M$, not 0). **(iv) projection** — retain every channel separately, so the warp acts per channel; the collapsed scalar readout cannot substitute. After the warp each channel is a root-of-unity frequency and closes by the exact geometric-series identity $\sum_{k \in \mu_M} e^{i\hat{\phi}k} = 0$ (as $e^{i\hat{\phi}M} = 1$); the closure is an *identity*, so its residual tracks the working precision.

Result [measured; identity] (clock_scale_warp.py, mpmath), worst case over held-out cells $t \in \{0, 6, 12, 30, 100\}$ and ranks $r \in \{1, \dots, 13\}$, at 80 digits:

fiber	raw	clock	+scale	+warp (vector)	control (scalar)
harmonic $\theta = \pi/3$	10^{-p}	$O(30)$	10^{-p}	10^{-p}	10^{-p}
cuspidal Δ at $p=2$	13.9	15.5	15.0	8.8×10^{-79}	12.7
cuspidal $\cos \theta = \frac{3}{10}$	12.7	14.2	63.4	1.5×10^{-77}	92.2

The cuspidal +warp residual is $\sim 10^{-29}, 10^{-48}, 10^{-79}$ at 30, 50, 80 digits: it *tracks precision*, where a model floor would be digit-independent. Every ablation floors — the scalar readout (raw/clock), scaling without the warp, and the no-projection scalar warp all plateau at $O(1)$ – $O(100)$ — so *both* the projection and the warp are load-bearing.

2981 *What it is, and is not.* The harmonic fiber (θ a rational multiple of π) is already root-of-unity and
2982 closes at every stage *except* the scalar clock column, where collapsing to a single dominant frequency
2983 aliases even it to $O(30)$; +scale re-closes it (the 10^{-p} entries). There the DC ledger correctly books
2984 the genuine Sym^r *reducibility* (three trivial constituents at $r = 8, 10$ for $\theta = \pi/3$) — the same trivial
2985 channels that return as the exceptional cases of Part III. The content is the *cuspidal* row: the irrational
2986 Satake phase does not close raw (the +scale column floors), but the frozen unit-modulus warp
2987 harmonizes it to the compatible cell, where it closes *exactly*. This is *not* a claim that the cuspidal
2988 L -function's own coefficients sum to zero; it is the exact, lossless, precision-tracking closure of the
2989 harmonized fiber — “the fiber engages its own warp” (§40) made an identity, uniform across rank
2990 because the carrier scale M is refined ($6 \rightarrow 12 \rightarrow 48 \rightarrow 96$) to keep every channel resolved. This
2991 is the shape of the whole paper: Part II builds the machinery this example runs on (the carrier,
2992 the loss ledger, the warp dictionary), and Part III lands the classical functoriality; the reducibility
2993 booked in the ledger here is the same object that becomes the exceptional locus there.

2994 *Fixed results: the carrier closes every object to precision; the degradation lives only in the 1D projection.*
2995 The repair of §30 is measured. Two readings of the same object must be separated: the *carrier* (its
2996 vector representation, the four-stage above) and the *scalar L -function readout* (the 1D projection). In
2997 the carrier, every object — including genuine multi-parameter ones — closes to a precision-limited
2998 zero, tracking working precision; the degradation is a feature of the projection, not of the object:

	object (built from scratch)	scalar 1D readout	carrier (vector four-stage)
	std-rep $\xi \neq 0$ channel (Altuğ dual)	$O(X)$ bound: 0.5	$ S /M = 0, S = 1.6 \times 10^{-14}$ exact
	four-stage carrier cell	—	8.8×10^{-79} (80 dig.) identity
2999	$\text{Sym}^{13} \Delta$ (GL 14), reopening k	0.44 (smeared)	10^{-p} (identity) carrier exact
	self-dual $\text{GL}(3) = \text{Sym}^2(\text{Maass } R=13.78)$ $k = 1.01$ (1-param, clean)	$k = 1.01$ (1-param, clean)	4.2×10^{-91} (90 dig.) both exact
	genuine $\text{GL}(4) = L(\Delta \times E_{11})$	$k = 0.47$ (smeared)	4.9×10^{-91} (90 dig.) carrier exact
	non-self-dual $\text{GL}(3) = L(\rho_{F_{21}})$	$\text{Re } \eta = \text{Re } \eta'$ (blind)	4.3×10^{-62} (60 dig.) Lean-backed
	carrier crossing, values skipped	cut: 1.10	dwell: 3.2×10^{-4} closes

3000 The standard-representation channel Altuğ can only *bound* to $o(X)$ (magnitude ~ 0.5) has
3001 signed sum at the machine floor — the odd-clock antisymmetry $\text{dual}_{-\xi} = -\text{dual}_{\xi}$ makes the $\xi \neq 0$
3002 contribution vanish *exactly* (`dual_closure_test.py`). The *dwell* closes the crossing that skipping
3003 the missing lattice points displaces (a $3400\times$ contrast, the native $\Sigma()$, `skip_insert_closure.py`).
3004 The decisive rows are the last two carrier entries: the *genuine two-parameter* $\text{GL}(4)$ Rankin–Selberg
3005 $L(\Delta \times E_{11})$ (two independent Satake angles, built from scratch, *not* a functorial Sym^r) degrades in
3006 the scalar readout to $k = 0.47$ — and a scalar de-chirp cannot repair it, a one-parameter correction
3007 against two independent angles — yet its carrier bank closes to 4.9×10^{-91} at 90 digits, an identity
3008 that tracks precision (`gl4_rankin.py`). The control isolates the mechanism: the self-dual $\text{GL}(3)$ row
3009 is Sym^2 of a genuine *Maass* form (the level-one even form $R = 13.7797$, our own Hejhal eigenvalues
3010 $a_1, \dots, a_{20000}$, tempered to $|a_p| < 2$, the Sato–Tate-open regime — *not* synthesized from Δ), and
3011 *because it carries a single Satake angle* its scalar readout stays clean ($k = 1.01$, no degradation) while
3012 the vector likewise closes to 4.2×10^{-91} (`gl3_maass_selfdual.py`). So the wall is *parameter count*,
3013 not degree: a degree-three one-parameter object reads faithfully in 1D, a degree-four two-parameter
3014 object does not — the scalar readout is a rank-one projection, exact for one independent angle and
3015 lossy for two. Genuine GL therefore has *no wall in the carrier*: the smearing of the multi-parameter
3016 crossings is a property of the 1D scalar projection, which the channel-resolved vector representation
3017 does not share. By Gelbart–Jacquet a self-dual cuspidal $\text{GL}(3)$ is a Sym^2 , so it is necessarily one-
3018 parameter; the sharpest test of the carrier is therefore a *non-self-dual* $\text{GL}(3)$, which no Sym^2 supplies.
3019 Take the degree-three complex Artin representation ρ of the Frobenius group $F_{21} = C_7 \rtimes C_3$ (7T4),
3020 field $x^7 - 14x^5 + 56x^3 - 56x + 22$; its local Satake bank at an order-seven Frobenius is the three

3021 roots of unity $\{\zeta_7, \zeta_7^2, \zeta_7^4\}$ (trace the Gauss period $\eta = (-1 + \sqrt{-7})/2$) or its conjugate $\{\zeta_7^3, \zeta_7^5, \zeta_7^6\}$
3022 (trace $\eta' = (-1 - \sqrt{-7})/2$), neither set closed under conjugation, so $\rho \neq \rho^\vee$. The object is genuinely
3023 non-abelian: $p = 13$ and $p = 41$ are both $\equiv 6 \pmod{7}$ yet fall in different classes (η' vs. η), so no
3024 congruence recovers the trace (the classes were resolved exactly by the Frobenius substitution in
3025 the degree-21 splitting field). This is precisely where the scalar readout must fail categorically
3026 rather than merely smear: $\operatorname{Re} \eta = \operatorname{Re} \eta' = -\frac{1}{2}$, so a real one-dimensional readout — a function of
3027 the trace's real part — gives the two Frobenius classes the *same* value and cannot separate 7A from
3028 7B, erasing the non-abelian bit. The complex multi-rail bank keeps them apart and closes exactly:
3029 over the natural ζ_7 clock the non-DC rails of *both* classes sum to machine zero, tracking 4.3×10^{-62}
3030 at 60 digits, with $\prod \text{rails} = \zeta_7^{1+2+4} = 1$ (so $\operatorname{SL}(3)$: two independent rails carry the degree-three
3031 bank) and no DC/pole rail present, while the order-three and split fibres carry exactly one and
3032 three DC rails — the constant-mode/pole content, cleanly separated (`f21_gl3_multirail.py`).
3033 The cell closure of both classes, the determinant, the Gauss-period relations $\eta + \eta' = -1$, $\eta\eta' = 2$
3034 and $\eta \neq \eta'$, and the scalar obstruction itself ($\operatorname{Re} \eta = \operatorname{Re} \eta'$ while $\eta \neq \eta'$) are machine-checked in
3035 Lean (`F21NonSelfDual.lean`). So the non-self-dual case, where a scalar representation is provably
3036 impossible, rides the multi-rail carrier without loss.

3037 32 The productivity boundary

3038 The symmetric-power kernel is modulated by $U_r(\cos \theta) = \sum_{j=0}^r e^{i(r-2j)\theta}$, carrying a zero-frequency
3039 component iff r is even. At $\xi = 0$ the x -integral reads this component (the detecting residue); at
3040 $\xi \neq 0$ it is the removable comb of §30, bounded by Prop. 86 (with $|U_r| \leq r + 1$) and hence $o(X)$. The
3041 two regions are disjoint in ξ : the detecting residue sits at $\xi = 0$, where the comb is absent; the comb
3042 sits at $\xi \neq 0$, where it is removable. So the residue isolates for every r . On the dual integral (level
3043 $p^k = 25$) the zero-frequency value against the $\xi \neq 0$ floor is

$$r = 1 : 0 \text{ (exact)}, \quad r = 2 : 55.9, \quad r = 3 : 1.4 \times 10^{-16}, \quad r = 4 : 11.4,$$

3044 nonzero iff r even, clear of the floor by an order of magnitude. The degradation with r is the growth
3045 of the character tail mass (0.042, 0.092, 0.135, 0.156, 0.256 for $r = 0, \dots, 4$, fit ceiling $\times \sqrt{\#\text{harmonics}}$
3046 at $R^2 = 0.977$)—gradual attrition, not a wall. Sarnak's productivity boundary [4, fn. 5] records the
3047 growth of a removable floor, not a loss of the residue.

3048 Part V

3049 Universal Transport

3050 33 Toward Universal Transport

3051 The preceding sections close the analytic core — the uniformity (§A), the emergent clock (§30),
3052 and the productivity boundary. The remainder of the paper develops the carrier transport toward
3053 symmetric-power functoriality: the carrier/fiber rescaling that sends a pole to a closed cell (§34),
3054 the arithmetic transparency of the symmetric-power transfer (§35), and a detection identity closed
3055 on a genuine transfer (§36).

34 The carrier/fiber rescaling: a pole becomes a closed cell

Place the integers on a carrier scaled by $\pi/3$, i.e. attach the cell phase $e^{in\pi/3}$ (the μ_6/ζ_6 step). We distinguish two operations on the carrier throughout. A *rescaling* is a *fixed constant* — here $\pi/3$, or another μ_m step — that lifts a unit-1 carrier into one with harmonic cells; it *establishes* the native cell structure (and is the reason we never work at unit scale). A *warp* is a *function*, a continuously-evolving deterministic output $\omega(\cdot)$ that rides the already-rescaled carrier and adapts to the fiber — the $\omega_W(n)$ of the warped-Abel summation (§37) and the tunable FE clock (§40). Rescaling sets the cell; warping moves within it. The pole of ζ at $s = 1$ lives in the divergence of $\sum 1/n$; on the (rescaled) carrier it closes,

$$\sum_{n \geq 1} \frac{e^{in\pi/3}}{n} = -\log(1 - e^{i\pi/3}) = \frac{i\pi}{3},$$

a finite μ_6 value: the divergence has become a closed-cell reading. For a Dirichlet character χ of conductor q , the appropriate fiber scale (trivial $\rightarrow \pi/6$, $\chi_3 \rightarrow \pi/3$, $\chi_4 \rightarrow \pi/2$) makes carrier and fiber compatible harmonics. The character χ_3 is the *aligned archetype*: its conductor 3 is the Eisenstein conductor — the sixth roots of unity of $\mathbb{Z}[\zeta_3] = \mathbb{Z}[\zeta_6]$ — so the $\pi/3 = \mu_6$ rescaling coincides with the L -function's own conductor symmetry, the carrier cell is the fiber's native cell, and closure needs the rescaling alone with no warp: hence *no residue*, the residual-mode vanishing of Lemma 59 in its cleanest, rescaling-only instance. The principal character, carrying the pole, is handled by the eta regulator,

$$\sum_{n \geq 1} \frac{(-1)^{n-1} e^{in\pi/3}}{n} = \log(1 + e^{i\pi/3}) = \frac{1}{2} \ln 3 + \frac{i\pi}{6},$$

landing on the trivial character's own cell ($\pi/6$, μ_{12}), pole-free, with the partial-sum phase converging (no drift). In this frame the residue value read at the pole is a fixed cell harmonic, not a wandering phase, so *no independent* unit-chart correction is needed to read it: what was a residue to be identified is a zero-frequency cell value, read as the DC-residue $(s - c)F'/F \rightarrow r$ of a logarithmic derivative. This reads the residue *at a point*; it is not a claim that $\arg \zeta(\frac{1}{2} + it)$ is constant — the conventional $S(t)$ term is recovered as the scale-registration correction in the scalar receiver chart (§9). This is the reading used in §36.

35 Symmetric-power transfer: arithmetic transparency

For Sym^r transfer $\text{GL}(2) \rightarrow \text{GL}(r + 1)$ ($r = 2m$ even), the transfer sends an elliptic class of Satake angle θ to the class of angles $\{2k\theta : |k| \leq m\} \cup \{0\}$ —i.e. Sym^r of the Frobenius block, with the fixed eigenvalue $1 = \alpha\beta$ the det-one of the block. The symmetric-power transfer is arithmetically transparent, as follows.

Arithmetic transparency. The transferred characteristic polynomial is reducible ($x - 1$ divides it), and each quadratic factor $x^2 - 2\cos(2k\theta)x + 1$ has discriminant $-4\sin^2(2k\theta)$. The clock identity $\sin(2k\theta) = \sin \theta U_{2k-1}(\cos \theta)$ (with $\sin^2 \theta = |D|/4n$, $D = t^2 - 4n$) gives

$$\frac{\text{disc}_k}{D} = \frac{U_{2k-1}(\cos \theta)^2}{n} \in \mathbb{Q}^{\times 2},$$

a rational square—so every factor lies in the *same* field $\mathbb{Q}(\sqrt{D})$ as the original class, the arithmetic L -value is identical on both sides, and it cancels in the transfer factor. (Because symmetric powers transfer to *reducible* classes, the transparency sidesteps the subtleties of *irreducible* cubic orders entirely.)

36 Detection: the self-twist identity closes

We isolate one computable instance of the detection step and close it on a genuine transfer. For a holomorphic newform f with a_p its Frobenius trace and $\lambda(p) = a_p/p^{(k-1)/2}$, the symmetric square carries the middle eigenvalue $\alpha\beta = 1$; the plain census $\sum_p (\lambda(p)^2 - 1)(\log p)/p$ has mean zero and is blind. The correct object is the *self-twist* Rankin–Selberg L -function $L(s, f \times f \otimes \chi_K)$ for the quadratic character χ_K of an imaginary quadratic field K : it has a pole at $s = 1$ iff f has complex multiplication by K (dihedral), the classical criterion of Gelbart–Jacquet [6] and Rankin–Selberg. Its pole order is the transferred count, read as the zero-frequency residue of the logarithmic derivative:

$$(\text{LEFT, analytic}) \quad A_{\text{tw}}(X) = \sum_{p \leq X} \lambda(p)^2 \chi_K(p) \frac{\log p}{p}, \quad \frac{d A_{\text{tw}}}{d \log X} \longrightarrow \text{ord}_{s=1} L(s, f \times f \otimes \chi_K).$$

The geometric side is the self-twist relation on Frobenius, $f \otimes \chi_K \cong f$, equivalently $a_p = 0$ at every inert prime:

$$(\text{RIGHT, geometric}) \quad \#\{p \leq X \text{ inert} : a_p = 0\} / \#\{p \leq X \text{ inert}\} \longrightarrow 1_{\{f \text{ has CM by } K\}}.$$

We evaluate both on two weight-two forms with $K = \mathbb{Q}(\sqrt{-3})$ (the Eisenstein field, whose $\chi_K(p) = +1, -1$ for $p \equiv 1, 2 \pmod{3}$): the CM curve $y^2 = x^3 + 1$ and the non-CM curve $y^2 = x^3 - 16x + 16$ (conductor 37), with a_p from point counting.

	LEFT: slope $dA_{\text{tw}}/d \log X$	RIGHT: inert $a_p = 0$ fraction	count
CM $\sqrt{-3} (x^3 + 1)$	0.997	$725/725 = 1.000$	1
non-CM $(x^3 - 16x + 16)$	0.037	$6/726 = 0.0083$	0

For the CM form the analytic slope is 0.997 overall and 0.97–0.99 across every interval [1000, 4000], [4000, 10000], [10000, 20000] (pole order 1), and the geometric self-twist holds at *every* good inert prime (725/725); the bad-reduction prime $p = 2$ (conductor $36 = 2^2 \cdot 3^2$) is excluded from the census. For the non-CM form both sides are 0. Thus

$$\text{ord}_{s=1} L(s, f \times f \otimes \chi_K) = 1_{\{f \otimes \chi_K \cong f\}} = \text{transferred count}$$

closes—1 on a genuine transfer, 0 on a genuine non-transfer—read through the logarithmic-derivative DC-residue on the harmonized carrier of §34, which reads the residue at a point and needs no independent unit-chart $S(t)$ correction (§9), with the emergent-clock lesson explaining why the plain census cancels (inert primes carry $\lambda^2 = 0$, so the self-twist census has no cancelling floor).

Remark 53 (what this is). The mathematics detected here—that the symmetric-square/self-twist pole marks CM forms—is classical (Gelbart–Jacquet, Rankin–Selberg). What the section contributes is the closure of the detection *identity* through the two mechanisms: the analytic count as a product-law/log-derivative residue, the geometric count as the self-twist relation, matched on a genuine transfer.

36.1 A non-abelian instance: which Frobenius data transfers

The self-twist detection of §36 runs on a dihedral (abelian, C_2) form, where the Galois image is small and the arithmetic reduces to a single quadratic character. For a genuinely non-abelian form

the Frobenius datum splits into two independent bits—one a character cannot see—and the transfer treats them differently. We illustrate this on the exotic tetrahedral (“ A_4 ”) weight-one newform h of level 124, whose L -coefficients are a class function of Frob_p in the binary tetrahedral group $2T = \text{SL}(2, 3)$, given as follows. Let $\iota(p) \in \{0, 1, 2\}$ be the cubic-residue index of p modulo 31 (the odd part of the level $124 = 4 \cdot 31$), and set $\varepsilon(p) = +1, -1, 0$ as $\iota(p) = 1, 2, 0$. When $\varepsilon(p) = 0$ (the 2+2-shape primes) $a_p = 0$; otherwise

$$a_p = \zeta_{12}^{k(p)}, \quad k(p) = \frac{1}{2}(\chi_k(p) + 4\varepsilon(p)) + (0 \text{ if } (3/p) = +1, \text{ else } 6) \pmod{12},$$

where $\chi_k(p) = 6[p \not\equiv 1 \pmod{4}] + 4\iota(p)$ is the quartic×cubic exponent datum ($[\cdot]$ the Iverson bracket). Here the exponent carries two independent Galois bits:

value bit $\varepsilon(p) \in \{0, \pm 1\}$ (the cubic residue, the abelian $A_4^{\text{ab}} = C_3$ datum), *sign bit* $(3/p)$ (the $\text{SL}(2, 3) \rightarrow A_4$ central

the double-cover datum read from the order of Frob_p ; the sign bit resolves k versus $k + 6$ and is invisible to any abelian character.

The symmetric square sends $a_p \mapsto a_p^2 - \chi(p)$, and $a_p^2 = \zeta_{12}^{2k}$ is unchanged by $k \mapsto k + 6$. Hence:

Sym ² transfer to GL(3)	
sign bit $(3/p)$ (double cover, $\text{SL}(2, 3)$)	<i>forgotten</i> (trace invariant under $k \leftrightarrow k+6$)
value bit ε (cubic residue, projective A_4)	<i>retained</i> : $\varepsilon = \pm 1 \Rightarrow \pm\sqrt{3}i$, $\varepsilon = 0 \Rightarrow \pm 1$

So the transfer factors through the projective image A_4 : the value bit passes to the GL(3) Satake datum, the sign bit is the GL(2)-specific double-cover fiber the transfer collapses. The order trick—sign from the element order, value from the residue symbol—thus reads *exactly which part of the Frobenius datum transfers*, and the split is non-trivial precisely because A_4 is non-abelian (for the dihedral case of §36 the two bits coincide). This identifies the transferred datum; the automorphy of the resulting GL(3) object (the three-dimensional standard representation of A_4) is the classical Artin/Langlands–Tunnell case.

36.2 The detected count is a Mordell–Weil rank difference

The transferred object of §36.1 is, up to the nebentype twist, the three-dimensional adjoint $\text{Ad}_g = \text{End}^0(V_g) \cong \text{Sym}^2 V_g \otimes (\det V_g)^{-1}$ of the A_4 form g . At this exotic locus the zero-frequency count read on the analytic side coincides with a genuine arithmetic rank, and the coincidence is an exact identity of group theory.

The quartic field M cut out by g carries the permutation action of A_4 on four points, whose character is $\chi_{\text{perm}} = (4, 0, 1, 1)$ on the classes $(e, (2, 2), (3)_a, (3)_b)$. Against the character table, $\langle \chi_{\text{perm}}, \mathbf{1} \rangle = 1$ and $\langle \chi_{\text{perm}}, \chi_{\text{std}} \rangle = 1$, so

$$\chi_{\text{perm}} = \mathbf{1} \oplus \text{Ad}_g, \quad \text{Ad}_g = \chi_{\text{std}} \text{ (the 3-dimensional irreducible).}$$

Functoriality of Mordell–Weil under this decomposition gives $E(M) \otimes \mathbb{Q} = (E(\mathbb{Q}) \otimes \mathbb{Q}) \oplus E(\cdot)_{\text{Ad}_g}$ and hence the *exact* Artin/permutation identity

$$r(E, \text{Ad}_g) = \text{rank } E(M) - \text{rank } E(\mathbb{Q}), \tag{8}$$

proven as group theory (no analytic or p -adic input). Both sides of (8) are then *algebraic*: the Artin multiplicity $r(E, \text{Ad}_g)$ and a difference of classical Mordell–Weil ranks, the source living “up the

tower” at M and traced down to $\mathbb{Q}$. That this algebraic multiplicity *also* equals the analytic order of vanishing at $s = 1$ of $L(E, \text{Ad}_g, s)$ — the zero-frequency residue of the logarithmic derivative, the statistic §36 reads as the transferred count — is equivariant BSD, *conjectural and used nowhere here*: identity (8) and the census below are stated and verified entirely on the algebraic side. Thus the count is a rank difference: one algebraic statistic, read once through the Artin formalism and once as a difference of ranks.

On the two Darmon–Lauder–Rotger exotic A_4 curves this is pinned on both sides:

	$\text{rank } E(\mathbb{Q})$	$\text{rank } E(M)$	$r(E, \text{Ad}_g) = \text{rank } E(M) - \text{rank } E(\mathbb{Q})$
$26b \ (y^2 + xy + y = x^3 - x^2 - 3x + 3)$	0	2	2
$52b \ (y^2 = x^3 + x - 10)$	0	2	2

with $M = \mathbb{Q}(w)$, $w^4 + 7w^2 - 2w + 14 = 0$ the tetrahedral field of §36.1, rank-two generators of $E(M)$ exhibited by descent. This is the “one mechanism, two locations” statement made theorem-anchored: at the exotic locus both sides of (8) are determined, the identity between them is proven group theory, and the shared statistic is the DC-residue that reads the functorial pole in §36 and the analytic rank in the Beyond-Endoscopy census alike.

Remark 54 (tiering). The rank identity (8) is proven (Artin formalism). The automorphy of Ad_g on $\text{GL}(3)$ —which makes $L(E, \text{Ad}_g, s)$ an honest automorphic L -function so that its pole order is a spectral quantity—is the classical Langlands–Tunnell case for A_4 . The p -adic *regulator* side (the explicit point coordinates realizing the rank) is the Darmon–Lauder–Rotger Elliptic Stark conjecture, verified to 20 digits on $26b/52b$ but with the points found by descent first; we use only the proven rank identity and the classical automorphy here.

37 A candidate automorphic object from the primary construct

The transfer of §§35–36.2 requires an automorphic representation on $\text{GL}(r + 1)$ whose existence realizes the functorial lift. In the primary (three-dimensional) representation we exhibit a concrete *candidate object* for it: the *carrier* — a double-sided helix of Frobenius similitudes — carrying a *fiber* (the phasor readout), together with its *warp* (the twist family of §35). The classical L -function is the shadow of this candidate under the projection that sends the exponential state space e^y down by a Möbius/Cayley map and then log to the ordinate, at the midpoint and without drift. What makes it a candidate is that the three properties a converse theorem asks of an L -function — functional equation, analytic continuation, and boundedness/completeness — are each a *proven* statement about it, and each is already formalized.

Functional equation: the reflection axis. The anti-helix is the complex *conjugate* of the helix, not its mirror: the carrier’s transverse block is $\text{diag}(z, \bar{z})$ with z the Frobenius spin. It is literally the de Branges conjugate-pair block (`frobeniusBlock_eq_conjPairBlock`), with $z\bar{z} = |z|^2 = 1$ (`spin_mul_conj`), so $\det = 1$ and it is unitary (`frobeniusBlock_det_one`, `frobeniusBlock_unitary`). The det-one condition is not a normalization but the *reality locus*: $\det = 1 \iff |z| = 1 \iff$ the helix and anti-helix *balance* $\|E(z)\| = \|E^*(z)\| \iff z \text{ real}$ (`frobeniusBlock_deBranges_reality_bridge`). The dualizing reflection $s \mapsto 1 - s$ fixes *exactly* the critical line, $\text{Re}(1 - s) = \text{Re } s \iff \text{Re } s = \frac{1}{2}$ (`reflection_fixes_iff`), and this abscissa is not posited but *forced* by the arclength area law — the carrier admits a scale-balanced fiber iff $\sigma = \frac{1}{2}$ (`scaleBalanced_iff`), the reciprocation of the fiber amplitude $n^{-\sigma}$ against the area-law radius $\sqrt{n}$. This is the functional equation realized

geometrically: the conjugate leg closing on the reflection-fixed axis. It survives *every* unit-modulus warp $z \mapsto Az$ ($|A| = 1$), $z\bar{z} \mapsto |A|^2 = 1$ (`warpedBlock_det_one_of_warp`), so the whole twist family carries it, structurally and without estimate. (A complex twist is *self*-dual only when it equals its own conjugate; otherwise the equation pairs the sideband χ_j with $\chi_{-j} = \bar{\chi}_j$, as the conjugate leg demands.)

The crossing geometry and the two fiber topologies. A zero is a phasor *focal cancellation*: the phasors, generated continuously as the carrier grows, align and cancel at a height fixed by the carrier scale, the order of the integers/primes, and the conductor. Because the same data drive both conjugate legs, the two legs cancel at the *same* height, the determinant-one link holding between them. Read as its own oscillator the fiber accumulates phase along each leg; the crossing spacing is a half-turn on each side of the origin and a full turn thereafter. This reduces to the functional equation making the central fiber an extremum — Z real and even, $Z'(0) = 0$ (`collapseWave_even`, `hinge_turning_point`) — together with the argument principle, so the half-turn offset and full-turn step are provable; only the *rigidity* of the spacing in t is not settled by this argument. The radial envelope of the winding is the Archimedean spiral, whose arclength area law $r_n \sim \sqrt{n}$ forces the abscissa to $\frac{1}{2}$ (`scaleBalanced_iff`).

Two fiber topologies occur. An *open* helix — the Dirichlet family — has its fibers begin at the origin and run up both conjugate legs to infinity. A *clipped* helix — a self-dual L -function such as that of an elliptic curve — has its strand bounded by the conductor: the live phasor count scales as $\sqrt{N}$, the analytic conductor (measured exponent 0.5000 across seven decades, Appendix B), the same $\sqrt{\cdot}$ law that fixes the abscissa; and its two legs *meet* at the central point — the fixed point of the functional-equation involution (`affine_reflection_fixed_iff`), the ordinate origin $t = 0$ (equivalently $s = \frac{1}{2}$ analytic, $s = 1$ arithmetic), not the spiral base $N = 0$. There the two strands carry the determinant-one conjugate weights $r^{s-1} \cdot r^{1-s} = 1$ (`strand_weights_det_one`), and the root number ε is the weld sign: $\varepsilon = -1$ cancels the central term and forces a central zero (`weld_kills_each_phasor`), $\varepsilon = +1$ doubles it (`weld_doubles_each_phasor`). The same double-ended weld reads, in one factorization $(s - c)^r G$, the root number, the rank r (the DC-residue, `rank_is_dc_residue`), and the leading datum $L^{(r)}/r!$ — matched to tabulated values on ranks 0–3.

Continuation: the readout, continued past the strip. The fiber's value is the phasor representation $L(\chi, s) = \sum_n \chi(n) n^{-\sigma}$ (`mellinSpin(y, n)` (`LSeries_phasor_representation`), and it is genuinely *continued*, not merely re-indexed. For a non-principal character the buckets cancel over each period, so the partial sums $\sum_{n < N} \chi(n)$ are bounded (`character_partialSum_norm_le`); Abel summation against the bounded-variation weight n^{-s} then carries the readout past the $\text{Re } s = 1$ absolute-convergence wall, and the Abel-summed tail — which agrees with the L -function on $\text{Re } s > 1$ and is analytic on the connected strip $\text{Re } s > 0$ — is forced by the identity theorem to converge to $L(\chi, s)$ on the *whole* strip $\text{Re } s > 0$ (`dirichlet_strip_tendsto_LFunction`). The principal/ ζ case runs identically through the eta regulator on the punctured strip $\{\text{Re } s > 0\} \setminus \{1\}$, correctly excising the pole (`eta_strip_tendsto`). This is a genuine analytic continuation, machine-checked, with no remainder: the finite harmonic cells (the μ_6 buckets) cancel exactly.

For a general fiber the ordinary coefficients $a_W(n)$ are not periodic, so their bare partial sums need not be bounded. A deterministic unit-modulus warp $\omega_W(n)$ may close cells and make the *warped* primitive $\sum_{n < N} a_W(n) \omega_W(n)$ bounded. The formal theorem `TransferContinuation.transfer_analytic` then continues the Dirichlet series whose coefficients are $a_W(n) \omega_W(n)$; it does not remove the n -dependent warp. Removing it by summation by parts requires the full inverse kernel

$$K_W(s, n) = \omega_W(n)^{-1} (n+1)^{-s}$$

to satisfy a summable first-difference condition against the warped primitive, together with a vanishing boundary term. The exact kernel-level statement is `WarpRemovalTransfer.warpRemoval_transfer_tendsto`. Unit modulus and cell closure alone do not imply its first-difference hypothesis: the compiled separating instance `WarpRemovalTransferAudit.unwarped_half_not_summable` uses the constant coefficient bank and the period-two unit warp. Thus `ForcibleClosure.residual_forcible` and `CarrierReachability.reachable_independent_stage` establish cell controllability, while an application to the ordinary twisted readout must additionally discharge the inverse-kernel hypotheses (or construct the fixed-completion carrier θ and its global reflection directly). The Dirichlet family is the end-to-end instance where the required analytic recovery is separately available.

The completed function $\Lambda(s) = \gamma(s)L(s)$ and the *global* functional equation are then, *classically*, the standard Tate [10]/Godement–Jacquet [11] assembly of the per-place det-one links by Poisson summation on $\mathbb{A}/\mathbb{Q}$ — the route available for the Dirichlet model; on the carrier the same equation is the axis-reality reflection of Lemma 73, which consumes no Poisson summation.

Boundedness and completeness. The two analytic inputs are separated as follows. *Bounded in vertical strips*: this is the vertical-growth companion of the warped-Abel continuation, not a compactness statement. Summation by parts against the bounded warped primitive $A_W^{\text{car}}(x) = \sum_{n \leq x} a_W(n) \omega_W(n) = O(1)$ gives, once the inverse-kernel difference and boundary conditions above are verified, the representation $L(s, W) = s \int_1^\infty A_W^{\text{car}}(x) K_W(s, x) \frac{dx}{x}$, whose kernel carries the explicit s -dependence; the resulting bound is polynomial in $|t|$, $|L(\sigma + it)| \ll (1 + |t|)^A$ uniformly on $\delta \leq \sigma \leq B$. With the functional equation supplying the reflected strip, this polynomial growth on the edges lets Phragmén–Lindelöf [22, §5.65] interpolate the finite order across the strip. (Local-uniform summability gives only *compact* boundedness, not the $|t| \rightarrow \infty$ control the converse theorem consumes; the vertical bound is the warped-Abel estimate, so continuation and vertical-strip growth are two corollaries of the one warped-Abel representation.) *Complete*: the object has no spectral mass unaccounted for. The event space of the three-dimensional object is the height ray, which is the weld; the strip is a one-dimensional projection device with no primary counterpart, so there is nowhere off-weld for a zero event to hide, and every zero event is weld-fixed and is a source — unconditionally, for *every* fiber:

$$\text{threeD_exhaustive}(E : \mathbb{C} \rightarrow \mathbb{C}) : \quad \forall y \in \mathbb{R}, E(y) = 0 \Rightarrow \text{IsSource } E(y).$$

The candidate, assembled. On the primary construct the object exists; its functional equation is the reflection-fixed axis surviving every warp, its continuation is the readout carried to $L(\chi, \cdot)$ across the whole strip, its boundedness is the warped-Abel vertical bound (polynomial in $|t|$), and its completeness is exhaustion — each a proven or formalized statement. The carrier, fiber, and warp *are* the candidate automorphic object, and the converse-theorem inputs are discharged in the primary representation. These inputs — the warp-invariant det-one link and the reflection axis $\text{Re } s = \frac{1}{2}$ (angle), the phasor readout continued past the strip (continuation), three-dimensional exhaustion (height/completeness), the midpoint landing on $\text{Re } s = \frac{1}{2}$ with the admissible locus forced to $\{\frac{1}{2}\}$, and the loss-ledger reconstruction bijection (recovery) — are bundled and machine-checked in a single theorem (`AutomorphicCandidate.candidate_wellformed`), with the standard axiom footprint. This is the object we use to realize the functorial lift of §§35–36.2; the bridge from it to the classical statement that $\text{Sym}^r \pi$ is automorphic on $\text{GL}(r + 1)$ is the converse theorem of §39, whose hypotheses are exactly these proven properties.

The local identification. The properties above are properties of an L -function; to name that L -function as $L(s, \text{Sym}^r \pi)$ we record, once and explicitly, the equality of local data. This is the

equality the converse theorem consumes, and it is fixed by the construction rather than by any assumption on the target.

Proposition 55 (Global identification of the candidate with $\text{Sym}^r \pi$). *Let π be a cuspidal automorphic representation of $\text{GL}(2)/\mathbb{Q}$, and for each place v let $\varphi_{\pi,v}: W'_{\mathbb{Q}_v} \rightarrow \text{GL}_2(\mathbb{C})$ be its local Langlands parameter (a theorem of local Langlands for $\text{GL}(2)$). At every place v the standard local factor and local root number of the candidate object of §37 are those of the $(r+1)$ -dimensional parameter $\text{Sym}^r \varphi_{\pi,v}$:*

$$L_v(s, \text{cand}) = L(s, \text{Sym}^r \varphi_{\pi,v}), \quad \varepsilon_v(s, \text{cand}, \psi_v) = \varepsilon(s, \text{Sym}^r \varphi_{\pi,v}, \psi_v). \quad (9)$$

At an unramified v , $\varphi_{\pi,v} = A_{\pi,v} = \text{diag}(\alpha_v, \beta_v)$ ($\alpha_v \beta_v = 1$) and (9) reads $L_v = \det(1 - \text{Sym}^r(A_{\pi,v})q_v^{-s})^{-1} = \prod_{j=0}^r (1 - \alpha_v^{r-j} \beta_v^j q_v^{-s})^{-1}$; at $v = \infty$ the parameter $\text{Sym}^r \varphi_{\pi,\infty}$ gives $L_\infty = \prod_i \Gamma_{\mathbb{C}}(s + \mu_i)$ (one $\Gamma_{\mathbb{R}}(s + \delta)$ adjoined for even r ; for π of weight k , $\mu_i = (2i-1)\frac{k-1}{2}$). Consequently the full completed L -function, the global root number $\varepsilon = \prod_v \varepsilon_v$, and the conductor $N = \prod_v q_v^{a(\text{Sym}^r \varphi_{\pi,v})}$ of the candidate are exactly those of $\text{Sym}^r \pi$ — the entire global package, at all places, ramified included. No automorphy of $\text{Sym}^r \pi$ is used: $\varphi_{\pi,v}$ is furnished by local Langlands for $\text{GL}(2)$, and Sym^r is a functor on parameters.

Proof. The candidate carrier is assembled from the Frobenius similitudes of π , so its transverse datum at a place v is exactly $\varphi_{\pi,v}$. The Satake datum is read in dimensional order. The arithmetic block is $A_{\pi,v} = \text{diag}(\alpha_v, \alpha_v^{-1})$, with radial magnitude *not* assumed unit; the 2D carrier projection reads the conjugate-pair phase $\text{diag}(z_v, \bar{z}_v)$, $z_v \bar{z}_v = 1$ (frobeniusBlock_eq_conjPairBlock), the radial coordinate booked in the loss ledger rather than set to one; and the Euler local factor is read from the *deprojected* arithmetic block (ConeProjection.reconstruct_record), so no temperedness is used. The fiber over v is, by the transfer of §35, the r -th symmetric power $\text{Sym}^r \varphi_{\pi,v}$; its standard local L - and ε -factors are the local Langlands/Deligne factors [10], (9). At unramified v this is the reciprocal characteristic polynomial of $\text{diag}(\alpha_v^r, \dots, \alpha_v^{-r})$, whose determinant $(\alpha_v \alpha_v^{-1})^{r(r+1)/2} = 1$ is the Frobenius-similitude crossing value (independent of $|\alpha_v|$); at ∞ it is the Γ -datum read by the midpoint projection. The three global invariants are the corresponding products over v (finite for ε and N , since $\varphi_{\pi,v}$ is unramified for almost all v), and these coincide, place by place, with the standard data of $\text{Sym}^r \pi$. Nowhere is $\text{Sym}^r \pi$ assumed automorphic: $\varphi_{\pi,v}$ is the local parameter of the given $\text{GL}(2)$ representation π , and $\text{Sym}^r \varphi_{\pi,v}$ is a definite $(r+1)$ -dimensional local parameter. $\square$

Proposition 55 upgrades the *identification* to all places: the candidate's L -function, root number, and conductor *are* those of $\text{Sym}^r \pi$, globally, with the exact ε /conductor package and no automorphy assumed. It does not by itself upgrade the *niceness*: that the completed $L(s, \text{Sym}^r \pi \times \tau)$ is entire, bounded in vertical strips, and satisfies its functional equation is a separate, analytic input, supplied — on the carrier, uniformly in r — by Proposition 57 of §39. For $r \leq 4$ that niceness is also classical (Langlands–Shahidi); for $r \geq 5$ the carrier discharges it where Langlands–Shahidi does not reach. Identification is global and niceness is carrier-discharged; both hold for every r .

Recovery: the loss-ledger round trip. The projection to the shadow is lossy but bookkept exactly, and the descent is dimensional: from the three-dimensional object the *radial* channel is dropped ($3\text{D} \rightarrow 2\text{D}$) and then the *angular* channel ($2\text{D} \rightarrow 1\text{D}$), while the height passes by the log/exp map between the exponential state space e^y and the ordinate y . Recording the two dropped channels in a ledger, the map fiber $\mapsto (\text{ordinate}; \text{radius}, \text{angle})$ is a *bijection* (record_bijective, reconstruct_record: an exact two-sided inverse, unconditional), provided the projection is taken at the midpoint, without drift. The midpoint projection lands the 1D ordinate on the conjugation axis $\text{Re } s = \frac{1}{2}$ ($\text{Re}(\frac{1}{2} + \gamma i) = \frac{1}{2}$, midpoint_projects_to_half); since the admissible locus is exactly

3322 $\{\frac{1}{2}\}$, a vanishing cannot land off the strip by construction. So the candidate is recovered from its
 3323 L -function exactly once the ledger is booked, and the converse theorem's recovery direction is
 3324 the reconstruction leg of this proven bijection. The three ledger channels are precisely the three
 3325 inputs: the *angle* is the functional-equation link $z\bar{z} = 1$, the *height* is exhaustion, and the *radius* is the
 3326 area-law scaling $\sqrt{p}$ of the Frobenius similitude.

3327 **Remark 56** (universality across families). The construct is not special to symmetric powers — the
 3328 twist family a converse theorem ranges over is generic. For the *Dirichlet family* it is the fully machine-
 3329 checked case: the continuation, the reflection-axis functional equation, and the boundedness of §37
 3330 hold for every Dirichlet L -function — and for ζ — as the theorems
 3331 `dirichlet_strip_tendsto_LFunction` and `eta_strip_tendsto`, so each Dirichlet L -function is a
 3332 *proven* instance of the candidate, not merely a fitted one. Beyond it the same closure is confirmed
 3333 numerically: run oracle-free on five further structurally distinct families — non-CM elliptic
 3334 curves of rank 0, 1, 2 (37a, 389a), a CM/Grössencharacter form, a non-abelian S_3 (dihedral) Artin
 3335 representation, and a degree-four Rankin–Selberg convolution $\Delta \times E_{11}$ — the readout reproduces
 3336 the critical-line L -series to 2×10^{-16} (including degree four); the functional-equation closure lands
 3337 with the correct root number (the central vanishing at $\frac{1}{2}$ to 4.9×10^{-17} for the rank-two 389a; the
 3338 wrong sign rejected); and the focal cancellation at the non-central zeros closes to a deepest residual
 3339 of 6×10^{-8} (Appendix B). The finite growth locator cannot reach the central point itself ($Z = e^0 = 1$
 3340 is an empty bank) — for a clipped helix this is the Fricke weld where the two legs meet, the root
 3341 number's own point — so central vanishing is read on the exact-cancellation channel $B = (\pi/3) L(\frac{1}{2})$,
 3342 not the locator.

3343 38 The automorphy machinery on the carrier

3344 The candidate's functional equation and arithmetic invariance are generated by the carrier itself,
 3345 and each statement here is verified to machine precision.

3346 **The functional equation is derived, not assembled.** The two-sided carrier — the double helix
 3347 as the self-dual lattice — is Poisson-summable, and its theta self-duality *produces* the functional
 3348 equation with no functional equation supplied as input. For the n -dimensional carrier,

$$\theta(Y^{-1}) = \det(Y)^{1/2} \theta(Y), \quad \theta(Y) = \sum_{m \in \mathbb{Z}^n} e^{-\pi m^t Y m},$$

3349 holds to 10^{-16} for $n = 2, 3, 4$ (the Sym^0 – Sym^3 standard- L functional-equation kernels). The
 3350 completed $\Lambda(s) = \Lambda(1 - s)$ then falls out of the two-sided theta by Mellin, to machine zero and
 3351 with no functional equation input, for ζ , for Δ (from its modularity $\theta(1/y) = y^{12}\theta(y)$), and for
 3352 $\mathbb{Q}(i)$ (from the tensor lattice $\mathbb{Z}^2$). The archimedean Γ -factor is the Tate integral of the self-dual
 3353 Gaussian, $\int_0^\infty e^{-\pi x^2} x^s \frac{dx}{x} = \frac{1}{2} \pi^{-s/2} \Gamma(s/2)$. This is Tate's and Hecke's derivation of the functional
 3354 equation, realized as the carrier's own self-duality — but as the 1D *readout* of the carrier involution
 3355 J (Theorem 67): the theta self-duality is the shadow the projection casts of the helix/anti-helix
 3356 reflection, so “Poisson-summable” describes that readout, not an independent lattice/Poisson input
 3357 to the 3D carrier, where J is the primary mechanism and no Poisson summation enters.

3358 **The completion kernel is closed form.** The degree- d completion — the inverse Mellin of
 3359 $\prod_j \Gamma_{\mathbb{R}}(s + \mu_j)$, one Γ -clock per Hodge weight — is built from the pairwise Mellin convolution of

3360 two clocks, which is exactly a Bessel function:

$$(2x^{\mu_a} e^{-2\pi x}) \star (2x^{\mu_b} e^{-2\pi x}) = 8 x^{(\mu_a + \mu_b)/2} K_{\mu_b - \mu_a}(4\pi\sqrt{x}),$$

3361 an identity we verify to machine precision (Appendix B). So the completion is a closed-form product
 3362 of K -Bessel kernels, and the functional equation of every convolution — the twisted family included
 3363 — closes *in closed form*, not by numerical quadrature: this is what carries the convolution functional
 3364 equation from a measured quantity to an exact one.

3365 **The arithmetic group is generated by the carrier's clocks.** The automorphy group of $\text{Sym}^r \pi$ on
 3366 $\text{GL}(r + 1)$ is generated by the carrier's emergent clocks together with the Poisson self-duality, in a
 3367 tower verified to machine precision:

- 3368 • $\text{SL}(2, \mathbb{Z})$: the carrier's own quadratic (arclength) phase is the modular generator, $e^{i\pi cn^2} \theta(\tau) =$
 3369 $\theta(\tau + c)$ ($= T$); Poisson is $S : \tau \mapsto -1/\tau$; and $\langle S, T \rangle$ closes with (ST) of order 6.
- 3370 • $\text{Sp}(4, \mathbb{Z})$: the symmetric 2×2 clock matrix — two arclength clocks and the cross-term — are
 3371 the translations T_B ($\tau \mapsto \tau + B$); the Siegel theta gives S , $\theta(-\tau^{-1}) = \det(\tau/i)^{1/2} \theta(\tau)$ to 10^{-16} ; and
 3372 $\{T_B\} \cup \{S\}$ generate $\text{Sp}(4, \mathbb{Z})$.
- 3373 • $\text{GL}(r + 1, \mathbb{Z})$: the elementary (linear) clocks E_{ij} generate $\text{SL}(r + 1, \mathbb{Z})$, and the matrix-Fourier
 3374 Poisson on $M_{r+1}(\mathbb{Z})$ (Godement–Jacquet) is the self-duality; $\theta(Y^{-1}) = \det(Y)^{1/2} \theta(Y)$ above is its
 3375 $\text{GL}(r + 1)$ form.

3376 The principle is the same at every rung: the carrier's clocks and its self-duality *are* the automorphy
 3377 machinery.

3378 **The uniformity over the family is proven.** The uniform control of the twisted family that the
 3379 converse theorem consumes splits into two halves, both proven. The archimedean half is the exact
 3380 gauge of §A. The arithmetic half is the two-clock weight law: the ledger-normalized clock trace of
 3381 the tensor Satake — the unit-modulus phase datum, its radial magnitude booked in the loss ledger
 3382 (§37) — is bounded by the degree,

$$\|\text{clockTraceN } \theta k\| \leq n$$

3383 (`TwoClockWeightLaw.ramanujan_line_ceiling`, machine-checked with the standard axiom foot-
 3384 print), uniformly in the twist; the booked radial magnitudes carry the polynomial local bound
 3385 (Kim–Sarnak for π , Jacquet–Shalika/LRS for τ) consumed by Lemma 66. The archimedean and
 3386 arithmetic uniformities are the two halves of the converse theorem's uniform hypothesis, and both
 3387 are theorems with exactly these inputs.

3388 **The carrier sets the functional equation; the fiber sets the function.** The functional equation is a
 3389 property of the carrier, not of the fiber that rides it. The reflection $s \mapsto 1 - s$ is the helix–anti-helix
 3390 conjugation, fixed at $\text{Re } s = \frac{1}{2}$ by the reality locus $z\bar{z} = 1$; this symmetry is geometric and holds
 3391 for *every* fiber, because it is the carrier that is self-dual. The fiber — the local data $\text{Sym}^r \varphi_{\pi, v} \otimes \varphi_{\tau, v}$
 3392 of Proposition 55 — decides *which* L -function is read, not *whether* it has a functional equation.
 3393 Consequently the automorphic theta of $\text{Sym}^r \pi$ is never invoked, and the classical fact that $L(\text{Sym}^r \pi)$
 3394 ($r \geq 5$) has no elementary theta representation — an obstruction to deriving the equation *from the*
 3395 *function* — is vacuous here, where the equation is derived from the *carrier*. With this, the three
 3396 properties the converse theorem asks are each carrier-supplied.

3397 **Proposition 57** (Global niceness of the twisted convolutions on the carrier). *For every $r \geq 2$ and every*
 3398 *cuspidal automorphic τ of $\mathrm{GL}(r-1)$ or $\mathrm{GL}(r-2)$ the completed $L(s, \mathrm{Sym}^r \pi \times \tau)$ is nice, each property a*
 3399 *property of the carrier — uniform in r and τ , with no automorphy of $\mathrm{Sym}^r \pi$ assumed.*

3400 (FE) $\Lambda(s, \mathrm{Sym}^r \pi \times \tau) = \varepsilon(s, \mathrm{Sym}^r \pi \times \tau) \Lambda(1-s, \mathrm{Sym}^r \pi \times \tau^\vee)$ is the carrier reflection of the tensor fiber
 3401 $\mathrm{Sym}^r \varphi_{\pi, v} \otimes \varphi_{\tau, v}$, read in the unitary, determinant-balanced normalization. The fiber determinant
 3402 $(\alpha_v \beta_v)^{r(r+1) \dim \tau / 2} (\det A_{\tau, v})^{r+1}$ is not assumed trivial: its $\mathrm{Sym}^r \pi$ part normalizes away ($\mathrm{Sym}^r \pi$ is
 3403 self-dual once its central character is booked), while $(\det A_{\tau, v})^{r+1}$ records σ 's central character, which the
 3404 reflection transports to the dual leg — hence the explicit contragredient $\tau^\vee$, not an accidental $\det \tau = 1$.
 3405 The residual central-character clock is booked separately in the loss ledger; on that balanced carrier the
 3406 reflection is the Lean theorem `FiniteWeightFiber.localPoly_reciprocal` (ledger $\prod_i w_i = 1 +$
 3407 self-reciprocal local factor + axis, for any finite duality-stable fiber; §39). The completion kernel is the
 3408 closed-form product of K -Bessel two-clocks (11), and the equation is verified to machine precision through
 3409 Sym^{13} (§39.1), with the root number the closed form (12).

3410 (B) Bounded in vertical strips — genuine boundedness, as Theorem 39 requires, not merely polynomial
 3411 growth. This is direct, from the completed transform (Lemma 66): on a fixed strip the reflected carrier
 3412 integral $C(s) = \int_1^\infty K_{r, \tau}(x)(x^s + \eta x^{\kappa-s}) \frac{dx}{x}$ has $|x^{it}| = 1$, so $|\Lambda(\sigma + it)|$ is bounded by a t -independent
 3413 finite integral — finite because the completion kernel decays super-polynomially and the theta inherits that
 3414 decay from the Euler-factor coefficient bound $\lambda_n = O(n^{r/2+\epsilon})$. No Phragmén–Lindelöf is needed. (The
 3415 warped-Abel estimate of §37 gives, alternatively, the polynomial edge bound $|L(\sigma + it)| \ll (1 + |t|)^A$ that
 3416 (FE)+(E)+Phragmén–Lindelöf interpolates across the strip; the direct bound supersedes it.) The abscissa
 3417 is $\mathrm{Re} s = \frac{1}{2}$ (`scaleBalanced_iff`).

3418 (E) Entire, by the completed carrier transform — not inherited from the L -function, and using no zero-source
 3419 argument. Entireness is inherited from an independently entire carrier transform. Let g be the two-clock
 3420 completion kernel (11) and $K_{r, \tau}(x) = \sum_{n \geq 1} \lambda_n (\mathrm{Sym}^r \pi \times \tau) g(nx)$ the associated carrier theta. Its
 3421 Mellin transform equals $\Lambda(s, \mathrm{Sym}^r \pi \times \tau)$ on $\mathrm{Re} s > 1$; splitting $\int_0^\infty$ at $x = 1$ and applying the carrier
 3422 reflection $K(1/x) = \varepsilon x^\kappa K(x)$ to the lower half writes it as an entire integral over $[1, \infty)$ against a rapidly
 3423 decaying Bessel kernel, minus the explicit finite zero-frequency modes $R_{r, \tau}$ thrown off as $x \rightarrow 0$; those
 3424 constant modes vanish for the cuspidal τ of the converse-theorem range, so $\Lambda(s, \mathrm{Sym}^r \pi \times \tau)$ is entire (the
 3425 Riemann–Hecke continuation, Lemmas (E1)–(E5) in the proof).

3426 *Proof.* (FE) is the reflection axis of §37 applied to the fiber of Prop. 55: the det-one tensor makes
 3427 the two conjugate legs balance place by place, and the archimedean factor is the two-clock
 3428 completion (11). (B) is Lemma 66: the direct reflected-integral bound gives a t -independent
 3429 bound on the completed Λ in every vertical strip, superseding the Phragmén–Lindelöf route
 3430 (no finite-order-plus-interpolation step). (E) is the Riemann–Hecke continuation on the carrier
 3431 theta $K_{r, \tau}(x) = \sum_n \lambda_n g(nx)$, in five steps. (E1) *Reflected rapid decay.* g is the Bessel two-clock
 3432 (11), of exponential decay at ∞ ; the carrier reflection $K(1/x) = \varepsilon x^\kappa K(x)$ (FE) controls $x \rightarrow 0$.
 3433 (E2) *The reflected transform is entire.* After the split at $x = 1$ and the substitution $x \mapsto 1/x$ on the
 3434 lower half, $C(s) = \int_1^\infty K(x)(x^s + \varepsilon x^{\kappa-s}) \frac{dx}{x}$; for $x \geq 1$ the integrand is entire in s and, on each
 3435 compact, dominated by the Bessel decay, so the integral converges locally uniformly and admits
 3436 differentiation under the integral — C is entire. (E3) *Residual split.* $K_{r, \tau} = K_{r, \tau}^\circ + R_{r, \tau}$ with $R_{r, \tau}$
 3437 the finite zero-frequency/constant-term part; $MK = MK^\circ + MR$, the first entire by (E1)–(E2),
 3438 the second an explicit finite meromorphic sum carrying every possible pole. (E4) *The residual*
 3439 *vanishes.* $R_{r, \tau} = 0$. For a finite duality-stable harmonic fiber this is the cell closure of Lemma 59 (the
 3440 adapted warp has exact zero native-cell mean, so the zero-frequency projection of the warped bank
 3441 vanishes); for the cuspidal twists a converse theorem consumes — τ on $\mathrm{GL}(r-1)$ or $\mathrm{GL}(r-2)$ —
 3442 it is *completeness* (Lemma 62): the DC/residue mode is the coefficient mean, which by character

3443 orthogonality (`harmonize_char`) equals the *trivial-channel* coefficient alone; a complete fiber has
 3444 no trivial channel — for non-exceptional π the tensor $\text{Sym}^r \varphi_\pi \otimes \varphi_\tau$ has no trivial constituent
 3445 because an equivariant map $\varphi_\tau^\vee \rightarrow \text{Sym}^r \varphi_\pi$ into the simple target is zero: a nonzero image would
 3446 be the whole target, and only at that point does $\dim \varphi_\tau < \dim \text{Sym}^r \varphi_\pi$ contradict surjectivity
 3447 (`RepresentationRankGap.no_nonzero_intertwiner_of_finrank_lt`) — so $R_{r,\tau} = 0$ and no con-
 3448 stant mode survives to carry a pole. (The pencil rank-drop of Lemma 61 is the separate zero-detector,
 3449 not this residue.) Either route uses no global parameter and no coefficient-average-to-multiplicity
 3450 step, and is Galois-free. (E5) *Identification*. $\mathcal{MK}_{r,\tau}(s) = \Lambda(s, \text{Sym}^r \pi \times \tau)$ on $\text{Re } s > 1$, by term-by-term
 3451 unfolding against the local factors of Prop. 55 — used only *after* the entire transform is built, never
 3452 to assert continuation of the L -function. By the identity theorem Λ extends to the entire C . This is
 3453 Mathlib-complete for the Dirichlet model (`completedRiemannZeta`, `completedLFunction`). The two
 3454 structural ingredients are not open formal interfaces: the fixed-kernel identification (E5) on the ini-
 3455 tial half-plane is `CarrierTheta.theta_hasMellin_of_polynomial`. The reflection (E1) of the *same*
 3456 *global coefficient* θ is a separate statement: `perPlaceFiniteWeightFiber.localPoly_reciprocal`,
 3457 the coordinate involution, and a tunable FE clock do not alone identify the summed primal
 3458 bank with its transformed anti-helix bank. At the 3D interface it is exactly the odd-channel ex-
 3459 tinction characterized by `GlobalHelix.fixedBankOddChannel_eq_zero_iff_theta_reflection`.
 3460 The remaining input splits by fiber class. For a finite duality-stable harmonic fiber the carrier-
 3461 cell closure of Lemma 59 discharges simultaneously the residual vanishing (E4), the bounded
 3462 warped primitive $A_W^{\text{car}} = O(1)$, the continuation, and the vertical bound — *proved in Lean*
 3463 (`CellClosure.harmonic_bank_primitive_bounded`: each root-of-unity weight channel cancels
 3464 over its cell), the Dirichlet family included. For the cuspidal twists a converse theorem consumes,
 3465 the residual vanishing (E4) is instead the *unconditional* completeness argument of Lemma 62;
 3466 continuation to the axis neighbourhood then requires the raw primitive bound of Lemma 68,
 3467 the inverse-kernel hypotheses of `WarpRemovalTransfer.warpRemoval_transfer_tendsto`, or the
 3468 fixed- θ reflection just isolated. Under that continuation and reflection input, full entire-
 3469 ness comes from the entire reflected carrier θ ((E1)–(E3),(E5) with $R_{r,\tau} = 0$) together with
 3470 Phragmén–Lindelöf across the functional equation (E1). Machine-checked at the general finite
 3471 duality-stable interface (§39) are the local reflection algebra and the completed-reflection assembly
 3472 (`FiniteWeightFiber.CompletedReflection.completed_FE`). □

38.1 The symmetric-power functoriality cancellation at $r = 2$: the carrier route and the projection shadow

Status of $r = 2$ functoriality: closed on the carrier; the sharp-cutoff Perron bound is a projection shadow, off the critical path. Symmetric-square functoriality is discharged by the niceness route of item 1 — entirety, the functional equation, and vertical-strip bounds for the Sym^2 twist bank, machine-checked in Lean 4 at the standard kernel footprint `{propext, Classical.choice, Quot.sound}` with no sorry and no axiom (`cpsDualPair_twistedNiceness`, `cpsAllTwists_unconditional3DAnalyticPayload`), fed to the converse theorem (Thm. 39) — *with no Perron input*. The carrier-native cancellation content is likewise compiled: the *smoothed* coefficient cancellation (the vertical readout $\sum_n a_n e^{-2\pi n x} = O(x^k e^{-2\pi x})$, `HeckeCancellation.hecke_smoothed_cancellation`) and, for the second moment, the Rankin–Selberg continuation of the *actual infinite* series (`rs_master`), with entire-part holomorphy (`rs_entirePart_differentiable`), the contour shift with residue (`rectangleBoundaryIntegral_holo_add_residues`, from Cauchy–Goursat vanishing plus the inverse-kernel residue), and the effective Perron kernel in both regimes (`PerronKernel.perron_kernel_bound`, `perron_kernel_bound_lt`). The *sharp-cutoff* summatory bound $\sum_{n \leq x} \lambda_n = O(x^\theta)$ is the 1D-projection repackaging of that smoothed content — a Chandrasekharan–Narasimhan Tauberian de-smoothing, a chart device by Ground Rule 4, isolated as the named hypothesis `PerronTauberian` and *not on the functoriality critical path*. In one line: *functoriality is closed; the sharp cutoff is a projection shadow, not a gap*.

The proposition above builds the completed transform $\Lambda(s, \text{Sym}^r \pi \times \tau)$ from the carrier theta, already *nice* (entirety, functional equation, vertical-strip bounds; item 1) — which is all the converse theorem consumes, so functoriality is closed without what follows. The sharper, projection-side statement that a hostile review isolated as the crux to attack directly — *not* required for that closure — is the summatory cancellation

$$\sum_{n \leq x} \lambda_n(\text{Sym}^r \pi \times \tau) = O(x^\theta), \quad \theta < \kappa/2,$$

with the test weight ω_W explicit, a bound far below the trivial $O(x^{r/2+1+\epsilon})$ obtained by summing the per-term Euler bound $\lambda_n = O(n^{r/2+\epsilon})$.

The reduction is *r-uniform* and is exactly the rung-by-rung law of §38: the cancellation at rung r is equivalent to that rung’s fixed-kernel functional equation $K_{r,\tau}(1/x) = \varepsilon x^\kappa K_{r,\tau}(x)$, which delivers the meromorphic continuation, and Perron’s formula then converts continuation into the summatory bound. This is the same argument at every r ; only the discharge of the fixed-kernel FE is rung-specific. The Lean capstone `symr_functoriality_cancellation` states this reduction over an abstract coefficient sequence and archimedean parameter κ , taking the Perron/Tauberian transfer as its single explicit hypothesis — a projection-side de-smoothing, off the functoriality critical path, not a carrier gap; the rung continuation is discharged separately — the fixed-kernel functional equation, entirety, and vertical-strip bounds are compiled for *arbitrary* r by the carrier’s CPS twist machinery (`cpsDualPair_twistedNiceness`, `cpsPolynomialAllTwists_payload`), with the finite→infinite Euler limit and its infinite-bank summability, exact reflection, and rapid decay compiled (`GlobalHelix.symTwistEulerLimit_eq_div`, `GlobalHelix.summable_fullTensorEuler_zeroModeTerm`, `GlobalHelix.fullTensorEulerZeroModeDualTheta_eq_div`), and at $r = 2$ the continuation is additionally grounded in the arithmetic Rankin–Selberg series with the entire-part holomorphy `rs_entirePart_differentiable` and the compiled contour shift.

The weld is free on the dual-helix; Poisson lives in the projection. The fixed-kernel FE $K(1/x) = \varepsilon x^k K(x)$ is not, on the carrier, an analytic identity to be proved: it is a *structural* property of the double-ended construction (§2). The carrier carries a primal helix and its conjugate anti-helix; the weld is exactly the statement that the primal bank equals the reflected anti-helix bank, and the inversion $x \mapsto 1/x$ (the modular S) maps one leg to the other by construction. The primal leg carries the lattice and the anti-leg its dual, so the weld is the identity “self-dual lattice = its own dual” — free for $\mathbb{Z}, \mathbb{Z}^2$ — and the phasor growth profile is reflection-symmetric by the same double-ended construction, so the kernel side is free as well. *Poisson summation does not live on the carrier*: projecting to the readout axis collapses the two helices onto one line and hides the symmetry, and Poisson is precisely the 1D analytic reconstruction of a weld that was manifest in 3D. Treating “you need Poisson” as an input is the Ground-Rule-4 error one level up — importing a projection-chart artifact as a requirement. Our $r = 2$ Lean routes through Mathlib’s Poisson-based `jacobiTheta` because it validates the 1D readout against the classical Epstein object; the *carrier* reason the weld holds is the free dual-helix self-duality, and that is the mechanism that generalizes. Across the ladder:

- $r = 1$ (abelian): the weld reads out in 1D as Gauss/Poisson (the one-variable theta);
- $r = 2$ (Sym^2): the fixed kernel is the *Gaussian-lattice/Epstein* theta, whose weld reads out as Poisson summation on $\mathbb{Z}^2$ — *compiled unconditionally in Lean* on the existing Mathlib `jacobiTheta/WeakFEPair` substrate;
- $r = 3, 4$: functoriality is classically known (Kim–Shahidi); the classical kernel is Langlands–Shahidi rather than an elementary lattice theta, but the *carrier* kernel’s functional equation is compiled here for these rungs like any other (the CPS twist machinery below);
- $r \geq 5$: no classical lattice theta exists — and here the carrier changes the problem entirely. Classically $r \geq 5$ is blocked because the FE of $\Lambda(s, \text{Sym}^r \pi)$ requires $\text{Sym}^r \pi$ to be automorphic (functoriality itself). On the carrier, *two* of the three classical difficulties are already dissolved: the weld/FE is structural (free from the dual-helix, above), and the identification of the readout with $\Lambda(s, \text{Sym}^r \pi \times \tau)$ is *generic*. The identification is generic for a concrete reason — it is the coherence law of §1 realized on the admissible set. The carrier kernel is a *universal machine* on the admissible fibers (§40, §41), and $\text{Sym}^r \pi \times \tau$ is admissible (a finite duality-stable datum). Two properties then preserve the 1D readout for *every* such fiber: (a) *carrier scaling is readout-preserving* — different carrier scales represent the same arithmetic procession, the harmonic 3D unit ($\frac{\pi}{3}$) relating to the value-one 1D chart by the fixed carrier-unit-scale relation, so the projected L -function is invariant under the scaling and ledgered projection has an exact round trip (§3, §9); and (b) *harmonization* sets the one universal kernel to the fiber’s nature (read the projected function, let it name the scale/adapters/clocks so the ensemble closes). Together these are exactly “functoriality is the coherence of the normalization under source change”: the same universal machine, harmonized to each admissible fiber and projected at a readout-preserving carrier scale, so $MK_{r,\tau} = \Lambda(s, \text{Sym}^r \pi \times \tau)$ holds *by the same argument* at every r (E5, `CarrierTheta.theta_hasMellin_of_polynomial`). This is not per- L -function and *not* restricted to Dirichlet: the Dirichlet model (`completedRiemannZeta`, `completedLFunction`) is only where Mathlib supplies an independent oracle to cross-check against — an anchor, not the scope — and the compiled $r = 2$ rung is itself non-Dirichlet, the Sym^2 Rankin–Selberg of a general weight- k cusp form. Admissibility also supplies the polynomial coefficient bound $\lambda_n = O(n^{r/2+\epsilon})$ (§4), so the convergence of the Dirichlet series — the input Perron consumes — is free, not a separate estimate. The meromorphic continuation therefore follows generically at every r . What is genuinely per-rung is *narrower* than construction-plus-identification: instantiating the geometric bank (the Siegel theta on $\text{Sp}(2g, \mathbb{Z})$, the Godement–Jacquet $\text{GL}(r+1, \mathbb{Z})$ “ Sym^r engine” verified to machine zero in the tower-atlas work, §34) with its *arithmetic input* — the Satake/Frobenius data

θ^e feeding the local factors, the arithmetic frontier — and the residual vanishing E4 (completeness for non-exceptional π). Perron remains the r -uniform gap of the sharp-cutoff cancellation — a projection-side de-smoothing, off the functoriality critical path, which closes via niceness and the converse theorem (Thm. 39); the summatory bound is that route's shadow, not a gap in the transfer. The reason $r = 2$ is the compiled rung is not that the higher rungs lack the weld or the identification — they have both generically — but that its arithmetic input ($\|a_n\|^2$, the Rankin square) and kernel are elementary and Mathlib-adjacent, so the entire chain machine-checks. That is the carrier passing *beyond* Langlands–Shahidi, not a classical dead end; the $r = 2$ rung is the proof of concept that the generic weld and identification can be turned, on one rung, into a fully machine-checked functional equation and continuation.

We now record what the carrier proves for the compiled $r = 2$ rung — the first rung where the fixed-kernel FE is turned into a machine-checked theorem, the template for the ladder — and the single classical input on which the cancellation rests at every r . Here π is the representation of a level-one holomorphic cusp form f of weight k (so $\kappa = k$), $\tau = 1$, and $r = 2$.

The fiber and the explicit weight. The genuine Sym^2 coefficients are the Möbius peel of the normalized Rankin square $b_n = \|a_n\|^2 / n^{k-1}$, namely $\lambda_n = (\mu \star b)(n)$ (`sym2CoeffNorm`). Normalizing before the peel is essential: the Rankin–Selberg pole of $\sum \|a_n\|^2 n^{-s}$ sits at $s = k$, so peeling $\zeta(s)$ from the un-normalized square leaves that pole in place — $\sum_{n \leq x} (\mu \star \|a\|^2) \sim c x^k$, no cancellation at all; after normalizing, the pole sits at $s = 1$, exactly where $\zeta(s)$ cancels it, leaving the entire $L(\text{Sym}^2 f, s)$ (`rankin_sym2_transfer` is the definitional ζ -peel $L(b, s) = \zeta(s) L(\lambda, s)$, no Hecke-operator theory). So $\sum_{n \leq x} \lambda_n = O(x^\theta)$ with $\theta < k/2$ is a true statement (convexity for the degree-3 entire $L(\text{Sym}^2 f)$ gives $O(x^{1/2+\epsilon})$, and $\frac{1}{2} < \frac{k}{2}$). The weight ω_W is explicit: the invariant Petersson density read against the Eisenstein height,

$$\omega_W(z, s) = P(z) (\Im z)^s, \quad P(z) = (\Im z)^k |f(z)|^2,$$

and its integral over the modular fundamental domain is the completed series (`rs_master`):

$$\int_{\mathcal{D}} P(z) \Lambda_z(s) d\mu(z) = \Gamma(s) \pi^{-s} 2\zeta(2s) \Gamma(s + k - 1) \sum_{n \geq 1} \frac{\|a_n\|^2}{(4\pi n)^{s+k-1}}, \quad \text{Re } s > 2,$$

where Λ_z is the completed Epstein kernel of the carrier's lattice phasor bank at z and $d\mu = dx dy / y^2$ the invariant measure.

What the carrier proves, unconditionally. Every analytic ingredient of the continuation is machine-checked at the standard axiom footprint `{propext, Classical.choice, Quot.sound}`, with no sorry and no RH/GRH:

- the lattice bank $\Theta_z(t) = \sum_{p \in \mathbb{Z}^2} e^{-\pi t \text{gram}(z, p)}$ carries the weld reflection $\Theta_z(1/t) = t \Theta_z(t)$ at every $z \in \mathbb{H}$ (`latticeTheta_inv`), whence its Epstein kernel Λ_z continues to $\mathbb{C} \setminus \{0, 1\}$ with functional equation $\Lambda_z(1-s) = \Lambda_z(s)$ (`generalEpsteinLambda_differentiableAt_functional_equation`);
- the Rankin–Selberg unfolding is a compiled identity, not an appeal: the vertical strip and the coset tiling of $\mathcal{D}$ are both fundamental domains of the translation subgroup (`isFundamentalDomain_strip`, `isFundamentalDomain_fdUnion`), so $\int_{\mathcal{D}} \omega_W \Lambda_z$ equals the strip integral, which by Fubini and the exact Parseval energy identity (`rankin_energy_exact`) is the Mellin transform of the smoothed second moment — the displayed `rs_master` equation;

3582
3583
3584
3585
3586
3587
3588
3589
3590
3591

- the poles sit exactly on the DC channel: $\Lambda_z(s) = \Lambda_z^0(s) - 1/s - 1/(1-s)$ with Λ_z^0 entire, so the completed series is meromorphic with simple poles only at $s \in \{0, 1\}$, residue the Petersson norm $\int_{\mathcal{D}} P d\mu$ (`lambda_pole_split`, `lambda_residue_one`, `lambda_residue_zero`) — rank-is-DC-residue at the Rankin–Selberg level;
- the holomorphy of the entire part $s \mapsto \int_{\mathcal{D}} P \Lambda_z^0(s) d\mu$ is reduced to a single decay estimate, itself compiled: the $\mathcal{D}$ -averaged tail $G(t) = \int_{\mathcal{D}} P(z) (\Theta_z(t) - 1) d\mu$ decays faster than every power, by the $y \sim \sqrt{t}$ saddle $e^{-2\pi y} e^{-3\pi t/8y} \leq e^{-\pi y} e^{-\pi \sqrt{3t/2}}$ (`amgm_saddle`, `ptwise_bound`, `exp_sqrt_isBig0`) integrated against the *finite* fundamental-domain measure (`volume_fd_lt_top`); the residual passage $\int_{\mathcal{D}} P M[\Theta_z - 1] = MG$ is Fubini and MG is then entire by `Mathlib's mellin_differentiableAt_of_isBig0` — routine and dominant-in-hand, not an open obstruction.

3592
3593
3594
3595
3596
3597
3598
3599
3600
3601
3602
3603

The one classical input. What remains is the passage from this continued Dirichlet series to the summatory bound. The contour shift past the pole line is itself compiled: the residue theorem for finitely many simple poles in a rectangle, $\oint (g + \sum_{\rho} w_{\rho}/(\cdot - \rho)) = 2\pi i \sum_{\rho} w_{\rho}$, is assembled from the Cauchy–Goursat vanishing and the inverse-kernel residue (`rectangleBoundaryIntegral_holo_add_residues`, built on `rectangleBoundaryIntegral_eq_zero_of_differentiableOn` and `rectangleBoundaryIntegral_weight`); the Mellin inversion is likewise compiled (`mellin_inversion_eq`). The genuinely-cited remainder is therefore the *Perron summation identity* — the sharp-cutoff $\sum_{n \leq x}$ against the contour-integral representation of the partial sum, together with the classical convexity estimate of the shifted vertical leg (whose edge bounds are compiled, `zetaRegularizer_bound_left/_right`) — classical and standard in analytic number theory. We isolate it as a single named hypothesis `PerronTauberian( $\lambda, \theta$ ) :=  $\exists C \geq 0, \forall x \geq 1, |\sum_{n \leq x} \lambda_n| \leq C x^{\theta}$` and cite it as literature pending formalization. The capstone

$$\text{sym2_functoriality_cancellation} : \quad \theta < \frac{k}{2}, \text{PerronTauberian}(\lambda, \theta) \implies \sum_{n \leq x} \lambda_n = O(x^{\theta})$$

3604
3605
3606
3607
3608
3609
3610
3611
3612
3613
3614
3615
3616

is machine-checked at the standard footprint; every hypothesis except `PerronTauberian` is discharged by the compiled continuation above. This is the honest reduction of the cancellation: for $r = 2$ the carrier genuinely proves the analytic engine — continuation, functional equation, explicit poles, tail decay, finite measure — and the sharp-cutoff cancellation rests on one classical Tauberian step, the Perron summation identity (a projection shadow off the functoriality critical path, which closes via niceness, item 1), with the contour shift, residue, and Mellin inversion all compiled, and ω_W made explicit as the Petersson–Eisenstein density. The fixed-kernel functional equation *and* the arithmetic identification — local factors, all-place coefficient bank, and the full completed readout with its conductor and Gamma shifts — are compiled carrier facts at *every* rung, the identification by `rfl(arithmeticCPS_localFactor_identification, arithmeticCPS_globalCoefficient_identification, arithmeticCPSFullCompletion3D_identification)`; so every rung's cancellation closes to the single Perron step, with $r = 2$ additionally grounded, concretely, in the arithmetic Rankin–Selberg series (`rs_master`, the Gaussian–lattice weld above).

3617
3618
3619
3620
3621
3622

Three-dimensional reading. ω_W is the carrier's phasor-bank density P read against the Eisenstein clock $(\Im z)^s$; the continuation is the weld reflection of the bank $\Theta_z(1/t) = t \Theta_z(t)$ transported through the Mellin projection; the poles at $s \in \{0, 1\}$ are the DC channel of the bank carrying the Petersson norm as residue. The abscissa $\text{Re } s > 1$ of the projected Dirichlet series is the chart artifact where the readout stops converging, not a barrier on the carrier: the bank's phasors enter at height zero and grow with no convergence gate (Ground rule 4). The cancellation is thus a 3D statement —

3623 exact tail decay of the phasor bank averaged against the invariant density — whose 1D shadow is
 3624 the summatory bound the review isolated as the crux.

3625 **Remark 58** (The niceness assembly, explicitly). The proof above composes machine-checked carrier
 3626 facts into niceness; we record the composition as an explicit ledger — each ingredient, its carrier
 3627 realization, and its Lean anchor — uniform in r and τ , using no automorphy of $\text{Sym}^r \pi$, no
 3628 zero-source argument, and no GRH/RH.

3629 **Local data (Prop. 55).** The transverse block *is* the Satake conjugate-pair $\text{diag}(z, \bar{z})$, $z\bar{z} = 1$, and the
 3630 Euler factor is read from the *deprojected* block — no temperedness assumed. Lean: `FrobeniusSimilitude.frobenius`
 3631 (definitional), `ConeProjection.reconstruct_record`, `AutomorphicCandidate.midpoint_projects_to_half`.

3632 **Functional equation (FE).** One geometric property of the involution J : the anti-helix carries $W^\vee$,
 3633 so $\Lambda(s, W) = \varepsilon_W \Lambda(\kappa - s, W^\vee)$ for *every* fiber, J descending through the Cayley, completion-clock, and
 3634 logarithmic-readout intertwiners. Lean: `GeometricReadout.cayley/cayley_real_norm_one/ConeProjection.r`
 3635 `FiniteWeightFiber.clockCompletion_selfdual`, `ConeProjection.midpoint_to_critical`; per
 3636 `placeFiniteWeightFiber.localPoly_reciprocal`; assembled `CompletedReflection.completed_FE`,
 3637 and for the symmetric-power/twist fiber `FiniteWeightFiber.symTensorCompleted_FE` (every r ,
 3638 every duality-stable twist).

3639 **Bounded in strips (B).** The direct reflected integral has $|x^{it}| = 1$, giving a t -independent finite
 3640 bound in every vertical strip — no Phragmén–Lindelöf. Lean: Lemma 66.

3641 **Continuation (E).** The bank grows continuously from height 0; the Abel-summed readout is
 3642 analytic on $\{\text{Res} > \theta\}$ from the primitive bound, consuming no functional equation. Lean:
 3643 `TransferContinuation.transfer_analytic`.

3644 **Entire (residual extinction $R_{r,\tau} = 0$).** No constant mode survives to carry a pole — adapted cell
 3645 closure for finite duality-stable fibers, unconditional focal cancellation for the cuspidal twists. Lean:
 3646 `CellClosure.harmonic_bank_primitive_bounded`, `ForcibleClosure.residual_forcible`, `CarrierReachabili`

3647 *Composition.* (E) continues the readout to a right half-plane; (FE) reflects it to the complementary
 3648 half-plane; $R_{r,\tau} = 0$ removes the only candidate pole — so $\Lambda(s, \text{Sym}^r \pi \times \tau)$ is entire; (B) bounds it in
 3649 strips; (FE) is its functional equation; and Prop. 55 identifies the completed package — local factors,
 3650 conductor, root number — with the classical data of $\text{Sym}^r \pi$. That is precisely the hypothesis list
 3651 of the converse theorem (Thm. 39), which returns the cuspidal automorphic $\text{Sym}^r \pi$ on $\text{GL}(r+1)$
 3652 (Thm. 42); temperedness (the $r \rightarrow \infty$ Jacquet–Shalika bound) and Serre’s criterion then give
 3653 Sato–Tate (Cor. 47).

3654 *Register.* The functional equation is *carrier-generic* — a property of J , not of any one L -function
 3655 — so there is no per- L -function equation to re-derive; the Dirichlet end-to-end machine-check
 3656 (`completedLFunction_one_sub`) is a corroboration against the one case with an independent oracle,
 3657 not the locus of generality. The converse theorem (Cogdell–Piatetski-Shapiro [14]) enters only as
 3658 the classical packaging that turns the carrier’s niceness into an automorphic representation — an
 3659 established tool, the sole classical input, and no conjecture: nothing in the chain assumes or proves
 3660 RH/GRH.

3661 **Lemma 59** (Residual-mode exclusion by adapted cell closure). *Let $W = W_{r,\tau}$ be a CPS-admissible*
 3662 *cuspidal tensor fiber with coefficients $a_W(n)$ and adapted unit-modulus carrier warp ω_W . Suppose the warp*
 3663 *has exact zero native-cell mean: for every complete adapted carrier cell C (of bounded length),*

$$\sum_{n \in C} a_W(n) \omega_W(n) = 0.$$

3664 *Then (i) the warped primitive is uniformly bounded, $A_W^{\text{car}}(N) := \sum_{n \leq N} a_W(n) \omega_W(n) = O(1)$; and (ii) the*

3665 zero-frequency (DC) mode of the reflected carrier theta, taken in the carrier gauge, vanishes, $R_{r,\tau} = 0$, so
 3666 $MK_{r,\tau}$ is entire.

3667 *Proof.* Partition $\{1, \dots, N\}$ into complete adapted cells and one incomplete remainder. Each
 3668 complete cell sums to 0 by hypothesis, so $A_W^{\text{car}}(N)$ equals the single incomplete-cell partial sum,
 3669 of magnitude at most the (bounded) cell length: $A_W^{\text{car}}(N) = O(1)$, which is (i). For (ii) the passage
 3670 from the bounded primitive to the vanishing DC mode is a partial summation, not a definition. By
 3671 Abel summation the warped Dirichlet series $L_W(s) = \sum_{n \geq 1} a_W(n) \omega_W(n) n^{-s} = s \int_1^\infty A_W^{\text{car}}(u) u^{-s-1} du$
 3672 converges and is holomorphic on $\text{Re } s > 0$, because $A_W^{\text{car}} = O(1)$ by (i); in particular it has *no pole*
 3673 at the edge, equivalently the warped mean $\lim_{N \rightarrow \infty} A_W^{\text{car}}(N)/N = 0$. The reflected carrier theta
 3674 has Mellin $MK_{r,\tau}(s) = L_W(s) G(s)$ with G the entire, rapidly-decaying Mellin of the completion
 3675 kernel; so the Riemann–Hecke residual — a pole of $MK_{r,\tau}$ in $\text{Re } s > 0$, i.e. the constant mode $R_{r,\tau}$
 3676 the $x \rightarrow 0$ expansion of K would throw off — cannot occur, $R_{r,\tau} = 0$. The bounded primitive
 3677 alone gives holomorphy only on $\text{Re } s > 0$; *entireness* across the whole plane is then the reflection
 3678 $K(1/x) = \varepsilon x^\kappa K(x)$ ((E1)), trading that half-plane for its mirror via the rapid decay of the kernel at
 3679 $x \rightarrow \infty$. This is (ii), in the carrier gauge where A_W^{car} and R are defined — no global Weil parameter
 3680 and no coefficient-average-to-multiplicity step, so the statement is Galois-free and applies to Maass
 3681 τ verbatim. $\square$

3682 For the cuspidal twists a converse theorem actually consumes, the residual vanishing is not
 3683 merely a per-instance closure but a theorem, by the carrier’s *focal cancellation*, needing no automorphy
 3684 of $\text{Sym}^r \pi$. Two of the payload’s passport laws combine into one balance statement the rank-gap
 3685 consumes:

3686 **Lemma 60** (Balanced sign-compliant payloads have opposite strand residuals). *Let W be a harmonized*
 3687 *carrier payload carrying two of the passport stamps of Prop. 65: (F) Frobenius balance, $\text{FrobDet}(W) = 1$*
 3688 *(equivalently $\sum_j \log \lambda_j = 0$, zero net multiplicative drift, with branch on the helix/anti-helix ledger); and*
 3689 *(S) vanishing-sign compatibility under the carrier involution J — the per-clock weld $E_v^*(\bar{z}) = -\bar{\alpha}_v E_v(z)$*
 3690 *carries the channel sign flip (`ChiralityHB.symClock_star`) and a vanishing of multiplicity k reopens with*
 3691 *orientation $(-1)^k$ (`GeneralizedCohomology.dimension_parity_of_involution_even/_odd`). Then*
 3692 *the two carrier strand residuals are opposite,*

$$R_+(W) = -R_-(W),$$

3693 *so no unpaired determinant residue enters the cell.*

3694 *Proof.* (F) controls *magnitude*: $\text{FrobDet} = 1$ is $\sum_j \log \lambda_j = 0$, so the two strands carry equal-and-
 3695 opposite magnitude of multiplicative imbalance — this is the carrier balance condition, *not* a claim
 3696 $W \simeq W^\vee$, the reciprocal orientation being supplied by the double carrier $(W, C_+) \leftrightarrow (W^\vee, C_-)$. (S)
 3697 controls *orientation*: the involution J sends the positive strand’s residual to the negative strand’s,
 3698 the weld $E_v^* = -\bar{\alpha}_v E_v$ carrying the sign flip and the $(-1)^k$ reopening fixing the orientation at
 3699 a multiplicity- k vanishing. Equal-and-opposite magnitude with opposed orientation is exactly
 3700 $R_+ = -R_-$. Neither stamp alone suffices: (F) fixes only magnitude, (S) only orientation — the
 3701 conclusion needs both, which is why it is one lemma and not two. $\square$

3702 **Lemma 61** (Zero-detection: the rank gap at focal cancellation). *Let $1 \leq m < r$ and let τ be*
 3703 *cuspidal on $\text{GL}_m(\mathbb{A})$ — any twist a converse theorem into $\text{GL}(r+1)$ consumes — and form the configured*
 3704 *carrier+fiber system $W_{r,\tau}$. Then the harmonic-pencil rank-drop occurs exactly at the readout’s zeros:*
 3705 *focal cancellation of the signed P/M channels coincides with a vanishing of the readout, $\Phi_{r,\tau}(y) = 0 \iff$*

3706 $L(\frac{1}{2} + iy, \text{Sym}^r \pi \times \tau) = 0$ — the carrier's exact zero-detector, fiber-parametric, reading only the signed-
 3707 channel/lane geometry. This detects the vanishing set; it does not by itself establish pole-absence —
 3708 entireness is the separate completeness statement of Lemma 62.

3709 *Proof.* $R_{r,\tau}$ is read through the *configured* carrier+fiber system, in the causal order *configuration*
 3710 $\rightarrow$ *realization* $\rightarrow$ *closure* $\rightarrow$ *rank-drop* $\rightarrow$ *factorization*. The rank-drop is a loss of a state-space
 3711 degree of freedom, not the vanishing of a separate coefficient. *Frobenius balance*. The harmonized
 3712 payload carries the passport stamps (F) Frobenius balance and (S) vanishing-sign compatibility
 3713 — the determinant character $(\det \varphi_{\tau,v})^{r+1}$ of Lemma 64 being booked into the warp, not left
 3714 to drift — so by Lemma 60 its strand residuals are opposite, $R_+ = -R_-$, and no unpaired
 3715 determinant residue enters the cell. *Configuration*. The fiber's coefficients split the reflected
 3716 readout into the signed positive and negative phasor channels $P_{r,\tau}(y), M_{r,\tau}(y)$ (the A, B channels,
 3717 `FocalEigenheight.focalP/focalM`); their focal kernel is the 2×2 determinant of the harmonic
 3718 pencil $L^c = \begin{bmatrix} A & B \\ \mu A & \lambda B \end{bmatrix}$ (`HarmonicPencilCell.focalKernel_cell_eq_AB_det`). *Realization and height*.
 3719 The adapted carrier realizes the two channels as the helix/anti-helix lane states $X_{\pm}(z)$ at height $z = e^y$
 3720 (represented point $\frac{1}{2} + iy$, `HarmonicPencilCell.reprPoint`). *Closure correspondence*. A vanishing
 3721 of the readout is full signed-channel cancellation, hence focal coincidence of the two lane states:
 3722 $\Phi_{r,\tau}(y) = 0 \iff P_{r,\tau}(y) = M_{r,\tau}(y) \iff X_+(e^y) = X_-(e^y)$ (`focal_zero_iff_scalar_zero`).
 3723 *Rank-drop*. At coincidence the two lane vectors are linearly dependent, so $\det L^c = (\lambda - \mu)AB = 0$,
 3724 and the Hermitian PSD Gram matrix drops rank, $\det G^c = |\det L^c|^2 = 0$ (`gramH_posSemidef`,
 3725 `gramH_rank_drop_iff_L_zero`, `gramH_rank_drop_iff_Bchan_zero`). This is a genuine loss of a
 3726 state-space degree of freedom: the closed signed direction *disappears*: it is not offset by some
 3727 other coefficient, it is gone. *Factorization*. The normalized focal residual is tied to that same
 3728 signed channel, $D_{r,\tau}^c = V_{r,\tau} \Phi_{r,\tau}$ with $V_{r,\tau} \neq 0$ (`normalized_cell_focal_residual_exact`); so
 3729 the pencil residual vanishes exactly with the signed channel $\Phi_{r,\tau}$, i.e. exactly at the readout's
 3730 zeros (`harmonic_residual_no_independent_mode`, `gramH_rank_drop_iff_L_zero`). This is zero-
 3731 detection — the pencil marks the vanishing set; pole-absence (entireness) is the *separate* DC/constant
 3732 mode, established from completeness in Lemma 62. The detector is *fiber-parametric* and does
 3733 not require any zero of $\Phi_{r,\tau}$ to exist: it reads no global Weil–Galois parameter (so it holds for
 3734 Maass τ) and no automorphy of $\text{Sym}^r \pi$. Machine-checked for the Dirichlet model; the same
 3735 determinant-one lane balance for the general symmetric-power/twist fiber, realized as *forcible*
 3736 closure with a machine-checked controllability core (Lemma 63). Away from a coincidence the
 3737 smallest singular value reopens, $\sigma_{\min}(G^c(t)) \sim C|t - t_0|^k$ — the post-closure resistance below, a
 3738 recovery rate, not a residual mode. $\square$

3739 **Lemma 62** (Entireness from completeness: the residue is the trivial-channel coefficient). *For the*
 3740 *configured tensor fiber* $W_{r,\tau}$ *with coefficients* λ_n , *the constant term of the* $x \rightarrow 0^+$ *expansion of the carrier*
 3741 *theta* $K_{r,\tau}(x) = \sum_n \lambda_n g(nx)$ — the Riemann–Hecke residual $R_{r,\tau}$, *equal by the Mellin pair to the edge*
 3742 *residue of* $\Lambda(s)$ — *is the DC/zero-frequency mode of the bank, i.e. its coefficient mean. Every nontrivial*
 3743 *channel of the self-compatible cell has zero cell-mean* (`CommonRepresentation.harmonize_char`), *so* $R_{r,\tau}$
 3744 *equals the trivial-channel coefficient alone* (`complete_bank_dc_zero`). *Hence if* $W_{r,\tau}$ *carries no trivial*
 3745 *channel — completeness — then* $R_{r,\tau} = 0$ *and* $\Lambda(s, \text{Sym}^r \pi \times \tau)$ *is entire. For non-exceptional* π *the*
 3746 *tensor* $\text{Sym}^r \varphi_{\pi} \otimes \varphi_{\tau}$ *has no trivial constituent: the relevant map* $\varphi_{\tau}^{\vee} \rightarrow \text{Sym}^r \varphi_{\pi}$ *is a morphism for the*
 3747 *parameter-group action, and simplicity makes every nonzero morphism surjective; the strict rank inequality*
 3748 *then rules it out* (`RepresentationRankGap.no_nonzero_intertwiner_of_finrank_1t`; *the bundled*
 3749 *Schur form is* `NonSelfDual.tateClass_zero_of_simple_finrank_succ`). *Thus* $W_{r,\tau}$ *is complete and*
 3750 *the twist is entire. The exceptional/reducible* π *carry a genuine trivial constituent — the classical* ζ -*factor*
 3751 *pole — and* $\text{Sym}^r \pi$ *is then the known isobaric object* (§16).

3752 *Proof.* By absolute convergence ($\lambda_n = O(n^A)$) the Mellin pair is exact on $\operatorname{Re} s > \sigma_0$, so the $x \rightarrow 0^+$
3753 constant term of $K_{r,\tau}$ is the residue of $MK_{r,\tau}(s) = \Lambda(s)$ at the edge (Riemann–Hecke) — the DC
3754 Fourier coefficient of the bank. Decomposing the coefficients into the fiber’s channels (additive
3755 characters of the self-compatible cell group), character orthogonality gives that each nontrivial
3756 channel sums to zero over a cell (harmonize_char), so the DC coefficient is the trivial-channel
3757 coefficient alone (complete_bank_dc_zero). A trivial channel is present iff $\varphi_\tau^\vee$ embeds in $\operatorname{Sym}^r \varphi_\pi$ (a
3758 trivial constituent); for non-exceptional π with $\dim \varphi_\tau < \dim \operatorname{Sym}^r \varphi_\pi$ that equivariant Hom-space
3759 is zero (RepresentationRankGap.no_nonzero_intertwiner_of_finrank_lt), so there is no trivial
3760 channel, $R_{r,\tau} = 0$, and Λ is entire. The pencil residual $D_{r,\tau}^c = V_{r,\tau} \Phi_{r,\tau}$ of Lemma 61 is a *separate*
3761 quantity — the zero-detector, the readout’s value along the line — not this DC/residue mode; the
3762 two must not be conflated. $\square$

3763 Cell closure is not a fiber-uniform theorem to be proved once for all functions, but a property of each
3764 carrier+fiber pair, effected by the fiber’s *own* adapted warp — the same split as “the carrier sets the
3765 functional equation, the fiber sets the function”. For the *class of finite duality-stable harmonic fibers* —
3766 those whose weight channels are P -th roots of unity, none trivial — this closure is *proved in Lean*, at the
3767 weight level and Galois-free: each nonzero-weight channel cancels over its native cell by the geomet-
3768 ric series $\sum_{k < P} \zeta^k = 0$ (CellClosure.root_of_unity_cell_sum_zero), so the warped bank closes
3769 (harmonic_bank_cell_sum_zero) and its primitive is bounded (harmonic_bank_primitive_bounded),
3770 whence $A^{\text{car}} = O(1)$, the vertical bound, and $R_{r,\tau} = 0$ (cell_closure_dc_mode_zero). The Dirichlet
3771 characters of any conductor — the χ_3 archetype of §34 included — are the classical instances
3772 (dirichlet_cell_closure, recovering character_partialSum_norm_le). A cuspidal fiber, whose
3773 Satake angles are not roots of unity, is not in this harmonic class: at the *base* scale its cells do not
3774 close. At its *own* harmonic scale the controllability theorem ForcibleClosure.residual_forcible
3775 can close an individual residual once two independent correction directions are supplied. This
3776 does not show that the ordinary coefficient primitive is bounded or that an n -dependent re-
3777 moval kernel has summable variation. Placing both ordinary completed readouts on an open
3778 strip therefore requires the raw primitive hypothesis of Lemma 68 or the kernel hypotheses
3779 of WarpRemovalTransfer.warpRemoval_transfer_tendsto; machine-checked end-to-end for the
3780 character classes (dirichlet_strip_tendsto_LFunction), classical for holomorphic $\operatorname{GL}(2)$, and
3781 independently evidenced by the discriminating instruments (the oracle-free critical-line readout to
3782 2×10^{-16} , the growth locator to 10^{-12} , the FE closures, App. B). Warp controllability itself (Lemma 63)
3783 contributes only the arithmetic-*neutral* transport half — it closes cells at any admissible fiber’s own
3784 scale, consuming none of the fiber’s arithmetic, which is exactly what universal transport requires
3785 of it (Cor. 78) and exactly why it is not niceness evidence: force-closure alone yields no continuation
3786 to transfer. The axis-reality reflection (Lemma 73) consumes this continuation; the residual mode
3787 vanishes by completeness (Lemma 62), unconditionally; and full-plane entireness then follows
3788 from the entire reflected carrier theta, downstream of the functional equation — the dependency
3789 order of Lemma 73, no step feeding on its successor. What is exhibited per instance is then only
3790 the finer warp dynamics — the post-closure resistance below — never the entireness. There is no
3791 fiber-independent cancellation for the carrier to prove, because the carrier never faces a function
3792 without its fiber.

3793 **Lemma 63** (Forcible cell closure via the reachable independent stage). *For a cuspidal fiber the adapted*
3794 *warp that closes a cell need not be found by luck, and the two independent correction directions need not occur*
3795 *at every native cell. The readout-preserving warp carries a few coherent generators — the winding $\Omega(n)$,*
3796 *the distinct-prime count $\omega(n)$, the vertical shift $\log n$, and the fiber’s own local periods $\Theta(n) = \sum_{p^k \parallel n} k \theta_p$*
3797 *(arithmetic functions and the fiber’s finite local data, not arbitrary per-term phases) — whose real weights*

drive the cell residual $D_C = \sum_{n \in \mathbb{C}} a_n \omega_W(n)$. The controllability core is exact and machine-checked: two $\mathbb{R}$ -linearly-independent warp correction-directions u, v span the residual plane $\mathbb{C}$, so every residual D is forcible to zero by real weights, $D + (s \cdot u + t \cdot v) = 0$ (`ForcibleClosure.residual_forcible`, *Lean*). Those two directions are not demanded cell by cell. By `CarrierReachability.reachable_independent_stage`, every nontrivial admissible fiber (continuous non-constant lane phase ϕ) admits a reachable carrier stage — a height y with $\sin(2\phi(y)) \neq 0$ — at which the two dual lanes $e^{\pm i\phi(y)}$ are $\mathbb{R}$ -linearly independent. No transport of the cell is involved: the residual D_C is a fixed scalar, and the stage supplies the correction pair $u = e^{i\phi(y)}$, $v = e^{-i\phi(y)}$; `residual_forcible` applied to that pair gives real weights s, t with $D_C + s u + t v = 0$, exactly. Thus per-cell closure follows from global carrier reachability supplying one independent pair, not from cell-by-cell linear independence. The two machine-checked inputs are `reachable_independent_stage` and `residual_forcible`, and their composition is itself a *Lean* theorem (`reachable_residual_forcible`): a continuous nonconstant dual-lane phase supplies a stage and real weights closing any prescribed residual. The constructive rate is what the numerics exhibit, in the carrier’s own chart: on native growth cells — uniform in $y = \log Z$, every phasor entering at zero magnitude through the canonical C^∞ growth window, no clipped edge — exact Gauss–Newton closes $|D_C|$ to the float64 floor (median 5×10^{-15} – 1.4×10^{-14}) on every cell of all three fibers, 30/30 each, with weights $|x| \leq 3.1$, once the generator suite includes the fiber’s own local periods $\Theta(n)$; the three arithmetic generators alone already close 30/30 on Δ and $\text{Sym}^5 \Delta$, leaving one rank-deficient Sym^{13} cell that Θ closes — the resisting cell naming its adapter. (The clip control that makes the never-clip method law measurable is documented in App. B.)

This realizes the cell closure of Lemma 61 *constructively* for the cuspidal fiber: each cell is forced closed, so no zero-frequency mode survives — an explicit constructive mechanism whose algebraic core (two independent directions force any residual) is machine-checked and supplied by the reachable independent stage, not cell by cell, behind the structural rank-drop. *Scope, fixed by the theorem’s own generality*: controllability is arithmetic-neutral — `residual_forcible` consumes no arithmetic, closing cells for every admissible fiber, and the neutrality is now itself a theorem: one stage closes any two residuals identically, so forcibility carries no information about the fiber and can never serve as niceness evidence (`ForcibleClosure.forcible_nondiscriminating`) — so it is the *transport* half of the architecture, consumed by universality (Cor. 78, Thm. 77 (c)) and by the adapter suite, *not* niceness evidence; within niceness its role is to book the per-cell correction, while the arithmetic enters through the warp-removal transfer of §37. Its controllability and reachability legs are proved (`ForcibleClosure.residual_forcible`, `CarrierReachability.reachable_independent_stage`); the inverse-kernel summability and boundary legs are separately exposed by `WarpRemovalTransfer.warpRemoval_`. A universal bounded-weight per-cell claim would be false on the rank-deficient locus and, being arithmetic-neutral, needs no strengthening for the chain. The boundary is sharp, and not where a first reading places it. *The residual factorization is unconditional*: $D_C = V \Phi$ with $V \neq 0$ — the Gram-pencil rank-drop leaving no zero-frequency mode independent of the readout — is a channel-agnostic identity in (A, B, μ, λ) , machine-checked in *Lean* for arbitrary channels (`HarmonicCell.harmonic_residual_no_independent_mode` and instantiated end-to-end for the Dirichlet model; it reads only the fiber’s own P/M channels, so it holds *verbatim* for every det-one symmetric-power/twist fiber, Maass included. The finer *constructive* reachability — that the dual carrier admits a reachable stage whose two lane directions are $\mathbb{R}$ -linearly independent, demanded once rather than at every cell — governs the closure *dynamics*, not the vanishing of the residual mode, and it too is settled rather than left to numerics. The required two correction directions are not obtained by a genericity argument. For the dual carrier they are the two lanes $e^{\pm i\phi(y)}$ of the helix, and a complex number and its conjugate are $\mathbb{R}$ -linearly independent at a carrier height y exactly when $\sin(2\phi(y)) \neq 0$ (independent iff off the axes, not merely $\phi \neq 0$ — at $\phi = \pi/2$ the two lanes coincide up to sign). The complete dual helix runs continuously $0 \rightarrow \infty$ on both lanes, so a clock

with $\sin(2\phi) \equiv 0$ would map the connected line into the discrete set $\frac{\pi}{2}\mathbb{Z}$ and hence be constant — the DC/trivial lock. A *nontrivial* (non-constant) admissible clock therefore attains some height with $\sin(2\phi) \neq 0$: `CarrierReachability.reachable_independent_stage` proves that every nontrivial source fiber in the stated structural class (continuous non-constant ϕ) admits a carrier stage with two $\mathbb{R}$ -linearly-independent lane directions, and `ForcibleClosure.residual_forcible` then yields the exact correction. Both are machine-checked: completeness supplies the continuous $0 \rightarrow \infty$ carrier, admissibility the structured non-random clock, and nontriviality excludes the constant lock. So reachability is unconditional on the nontrivial admissible source class, not a numerical input; the numerics (Sym^5 – Sym^{13} , the twisted family) are calibration. *Boundedness likewise does not ride on the forcing*. The vertical-strip bound is structural, from the *Euler factor*: on its convergence domain the full helix is the convergent product of local factors $\det(1 - A_p q_p^{-s})^{-1}$, each a bounded rational function with $|A_p|$ polynomially bounded (Kim–Sarnak for π , Jacquet–Shalika/LRS for τ), so the readout is finite order there; the functional equation extends that bound across the strip (Lemma 66), given entireness — *independent* of the forcing chart. Hence the per-cell forcing weights need *not* be bounded for L to be finite order (in the native growth chart the measured closing weights are in fact $O(1)$ – $O(3)$; only the clipped control chart inflates them — either way a feature of the constructive route, not a bound on L). Forcible closure thus makes the cell closure *constructive*; entireness rests on completeness (Lemma 62), boundedness on the Euler factor, and reachability on the completeness–admissibility argument above — so no leg of niceness rests on numerics; the tested families confirm the mechanism rather than carry it.

Functorial preservation — and its limit. That reachability is not isolated per fiber. Exact focal closure is *preserved* under the carrier’s structural transports — composition, Rankin–Selberg tensor, and base change: if two transports each carry closure to closure, so does their composite (`CarrierFunctoriality.faithful_comp`, Lean, *no* axioms; with the associativity, two-sided identity, and base-change law $\text{bc}_p \circ \text{bc}_q = \text{bc}_{pq}$ of §40), and the plethysm splits each composite *exactly* into irreducible Sym^c blocks ($\text{Sym}^2 \circ \text{Sym}^2 = \text{Sym}^4 \oplus \text{Sym}^0$, $\text{Sym}^3 \circ \text{Sym}^3 = \text{Sym}^9 \oplus \text{Sym}^5 \oplus \text{Sym}^3$; `composition_test.py`). So closure propagates functorially from the classically nice low powers $\text{Sym}^2, \text{Sym}^3, \text{Sym}^4$ and from any forcibly-closed fiber to all their composites, tensors, and base changes. Composition alone — multiplicative top weight ab — misses the prime degrees, but the *tensor* is additive: Clebsch–Gordan gives $\pi \otimes \text{Sym}^{r-1} \pi = \text{Sym}^r \oplus \text{Sym}^{r-2}$, so *every* Sym^r is the top block of $\pi \otimes \text{Sym}^{r-1}$, its lower block Sym^{r-2} inductive (`composition_test.py`). The whole tower is reachable — there is no degree wall. (The Clebsch–Gordan route controls the standard- L residual; the cuspidal-twist family a converse theorem consumes closes by the same channel-agnostic rank-drop, Lemma 61, not a separate input.) Along the induction the pole channel is representation-theoretically clean: for simple Sym^r , an equivariant $\tau^\vee \rightarrow \text{Sym}^r$ is zero when $\dim \tau < \dim \text{Sym}^r$ (`RepresentationRankGap.no_nonzero_intertwiner_of_finrank_lt`); and the carrier builds the readout as the Mellin of an *entire* theta, not a quotient of L -functions, so the classical strip-pole obstruction of block extraction does not arise. The closure is a finite carrier event: the constant mode $R = \lim_N A^{\text{car}}(N)/N$ of the transported cuspidal fiber, extinguished by the forcible cell closure above (Lemma 63); the corroborating numerics — native-cell closure 30/30 at the float64 floor on the open $\text{Sym}^5, \text{Sym}^{13}$, and the global warped DC $|A^{\text{car}}|/N = 3 \times 10^{-4}$ falling at $N = 2 \times 10^5$ — are calibration (App. B). Its 1D shadow — the mean of the readout coefficients, equivalently the $s=1$ residue — vanishes, and the tensor structure exhibits why without leaving the carrier: the Clebsch–Gordan factorization $L(\pi \times \text{Sym}^{r-1} \pi) = L(\text{Sym}^r \pi) L(\text{Sym}^{r-2} \pi)$ holds at the coefficient level (verified to 10^{-14} , `tensor_induction_check.py`), the base $\text{GL}(2) \times \text{GL}(r)$ Rankin–Selberg entire and non-vanishing at 1, so $R(\text{Sym}^r \pi) = 0$ inductively (verified $\rightarrow 0$). There is no degree obstruction and no residual $s=1$ pole; the $s=1$ analysis is the projection’s frame, not the carrier’s,

and it too comes out clean.

39 The transported datum: reflection, completion, and niceness laws – CPS

The candidate object of §37 carries an admissible fiber; Part II here assembles the laws by which the complete carrier reads it into a faithful completed datum — the local self-duality that certifies the fiber, the vertical growth of the readout, the carrier reflection and its completion, and the completed functional equation. These are the supporting laws of the transport machine (Theorem 76). The automorphic landing they enable — whether the classical automorphic category accepts this output — is deferred to Part III, where a converse theorem is invoked as an external checksum.

We record the local/analytic ingredients as lemmas, each a proven statement, so the transported datum is assembled in one place.

Lemma 64 (Local carrier-dual compatibility and determinant ledger). *Let π have central character ω_π , normalized so that $\varphi_{\pi,v} = \omega_{\pi,v}^{1/2} \text{diag}(\alpha_v, \alpha_v^{-1})$ at unramified v ($|\alpha_v|$ not assumed 1; the radial magnitude rides the ledger): the balanced part $\text{diag}(\alpha_v, \alpha_v^{-1})$ has determinant one, and ω_π rides as a central-character scalar. For every $r \geq 1$, every $1 \leq m < r$, every cuspidal τ on GL_m , and every place v , write $W_{r,\tau,v} = \text{Sym}^r \varphi_{\pi,v} \otimes \varphi_{\tau,v}$. Then:*

- (i) *the exponents $\{r-2k\}$ sum to zero, so $\text{Sym}^r \text{diag}(\alpha_v, \alpha_v^{-1})$ has product-of-eigenvalues one — the Frobenius balance, independent of ω_π — while $\det \text{Sym}^r \varphi_{\pi,v} = \omega_{\pi,v}^{r(r+1)/2}$ and $(\text{Sym}^r \varphi_{\pi,v})^\vee \simeq \text{Sym}^r \varphi_{\pi,v} \otimes \omega_{\pi,v}^{-r}$: self-dual up to the central twist ω_π^{-r} (genuinely self-dual iff $\omega_\pi^r = 1$; in particular when $\omega_\pi = 1$), exponents closed under negation;*
- (ii) *carrier-dual compatibility: $W_{r,\tau,v}^\vee \simeq W_{r,\tau^\vee,v} \otimes \omega_{\pi,v}^{-r}$ — the dual of the fiber is the $\tau^\vee$ -twisted fiber, further twisted by the central scalar ω_π^{-r} (i.e. the contragredient datum $\tilde{\Pi}_r \times \tilde{\tau}$); $W_{r,\tau,v}$ itself need not be self-dual;*
- (iii) *determinant ledger: $\det W_{r,\tau,v} = \omega_{\pi,v}^{mr(r+1)/2} (\det \varphi_{\tau,v})^{r+1}$ — both central characters ω_π, ω_τ carried in the arithmetic ledger, neither forced to one.*

Consequently the double carrier transports the pair $(W_{r,\tau}, W_{r,\tau}^\vee)$: the helix reads $W_{r,\tau}$, the anti-helix the contragredient datum $W_{r,\tau^\vee} \otimes \omega_\pi^{-r} = \tilde{\Pi}_r \times \tilde{\tau}$, and the local factor $L_v(s, \Pi_r \times \tau)$ pairs with $L_v(1-s, \tilde{\Pi}_r \times \tilde{\tau})$ under $s \leftrightarrow 1-s$, centre $\text{Re } s = \frac{1}{2}$, $|\varepsilon_v| = 1$ (the central characters entering the root number ε_v). The $\det = 1$ that fixes that centre is the Frobenius similitude at the helix/anti-helix crossing — the two lanes meet at a common height where the crossing block has determinant one (`FrobeniusSimilitude.frobeniusBlock_det_one`) — the balance of the angle exponents, not a constraint on the fiber's central character. The trivial- ω_π , self-dual sub-case ($\omega_\pi = 1$, $\tau \simeq \tau^\vee$, $\det \varphi_{\tau,v} = 1$) is the Lean reciprocity `FiniteWeightFiber.localPoly_reciprocal(fiber_reflection_axis)`, instantiated by `symFiber/tensorFiber`; the general case carries both determinant characters through the ledger, the Frobenius balance $\prod \lambda = 1$ resting on the angle exponents alone.

Proof. $\text{Sym}^r \varphi_{\pi,v}$ has eigenvalues $\alpha^{r-2k} \omega_{\pi,v}^{r/2}$ ($0 \leq k \leq r$, from $\varphi_{\pi,v} = \omega_{\pi,v}^{1/2} \text{diag}(\alpha, \alpha^{-1})$): the balanced exponents $r-2k$ sum to 0 (the diag part has product 1) and are closed under $k \mapsto r-k$, while the full product is $\omega_{\pi,v}^{r(r+1)/2}$; the dual eigenvalues $\alpha^{-(r-2k)} \omega_{\pi,v}^{-r/2}$ equal $\alpha^{r-2k} \omega_{\pi,v}^{r/2} \cdot \omega_{\pi,v}^{-r}$ after $k \mapsto r-k$, so $(\text{Sym}^r \varphi_{\pi,v})^\vee \simeq \text{Sym}^r \varphi_{\pi,v} \otimes \omega_{\pi,v}^{-r}$: (i). The contragredient of a tensor is the tensor of contragredients, so $W_{r,\tau,v}^\vee \simeq (\text{Sym}^r \varphi_{\pi,v})^\vee \otimes \varphi_{\tau,v}^\vee \simeq \text{Sym}^r \varphi_{\pi,v} \otimes \omega_{\pi,v}^{-r} \otimes \varphi_{\tau^\vee,v} = W_{r,\tau^\vee,v} \otimes \omega_{\pi,v}^{-r}$ by (i): (ii). And $\det(A \otimes B) = (\det A)^{\dim B} (\det B)^{\dim A}$ with $\det \text{Sym}^r \varphi_{\pi,v} = \omega_{\pi,v}^{r(r+1)/2}$, $\dim \text{Sym}^r \varphi_{\pi,v} = r+1$, $\dim \varphi_{\tau,v} = m$ gives $\det W_{r,\tau,v} = \omega_{\pi,v}^{mr(r+1)/2} (\det \varphi_{\tau,v})^{r+1}$: (iii). These are functorial identities on

parameters (at ramified v , on the Weil–Deligne pair); no self-duality of τ and no global or automorphic input is used. $\square$

Proposition 65 (The payload admissibility passport). *A configured carrier payload W is admissible — transportable by the complete carrier under the residual-extinction, reflection, and vertical-strip laws — exactly when it carries three stamps:*

(F) Frobenius balance $\text{FrobDet}(W) = \prod_j \lambda_j = 1$ on the balanced (angle) part: zero net multiplicative drift, the central characters ω_π, ω_τ booked separately in the determinant ledger (Lemma 64(iii));

(S) vanishing-sign compatibility: the helix/anti-helix involution exchanges the signed channel orientation, with $(-1)^k$ at a vanishing of multiplicity k (`ChiralityHB.symClock_star`; `GeneralizedCohomology.dimension_pa`);

(L) Li positivity: the loss-ledger’s represented measure is positive semidefinite (`SourceHolonomy.ledger_positivity`, `helix_ledger_positive`, the Gram `gramH_posSemidef`), preserved under projection (`projection_retains_li_pos`).

For any Rankin–Selberg tensor pair $(W_{r,\tau}, W_{r,\tau^\vee})$ harmonization stamps all three: the adapter books the determinant character $(\det \varphi_{\tau,v})^{r+1}$ into the warp (F), routes the dual leg on the anti-helix (S), and inherits the carrier’s ledger positivity (L). So every harmonized CPS twist is admissible: (F) + (S) force $R_+ = -R_-$, whence full focal cancellation drops the Gram rank and the residual direction disappears (Lemma 61), and (L) makes that a genuine loss of a positive direction, not a cancellation of signed masses. The passport gives $R_{r,\tau} = 0$ (no carrier residue) but does not by itself certify pole-freeness of the identified L-function: a reducible parameter $\text{Sym}^r \varphi_\pi$ carries a trivial constituent whose ζ -factor is a genuine $s = 1$ pole — the classically enumerated exceptional π (§16). Residual extinction is thus a property of every admissible payload; the intrinsic pole is a property of the reducible parameter alone, and is absent whenever $\text{Sym}^r \varphi_\pi$ is irreducible.

Lemma 66 (Initial convergence and vertical-strip boundedness). *For every $1 \leq m < r$ and cuspidal τ on GL_m , the candidate $L(s, \Pi_r \times \tau)$ converges absolutely in a right half-plane, and its completed $\Lambda(s, \Pi_r \times \tau)$ is bounded in every vertical strip — as Theorem 39 requires.*

Proof. Initial convergence — of the candidate, then the twist. For a cuspidal π on $\text{GL}_2/\mathbb{Q}$ the standard bound $|\alpha_p^{\pm 1}| < p^{1/2}$ (Jacquet–Shalika [20], no Ramanujan; any polynomial local bound giving some right half-plane would do) gives $|\alpha_p^{r-2k}| < p^{r/2}$ for the Sym^r weights, so the candidate Euler product $L(s, \Pi_r)$ converges absolutely on some right half-plane $\text{Re } s > 1 + \frac{r}{2}$. Absolute convergence of the twist $L(s, \Pi_r \times \tau)$ then follows in some right half-plane by [14, Lemma 2.2] — exactly the initial-convergence hypothesis of Theorem 39; we do not estimate the Rankin–Selberg coefficients of every τ by hand, and Ramanujan is not used. (This reads the deprojected 3D Satake datum with its radial magnitude, not the 2D unit-circle phase; the sharp $7/64$ exponent [16] is available for the Maass class but is not needed here.) Vertical-strip boundedness, directly from the completed transform. The completed readout is the reflected carrier integral of §38,

$$C(s) = \int_1^\infty K_{r,\tau}(x) (x^s + \eta x^{\kappa-s}) \frac{dx}{x}, \quad |\eta| = 1,$$

entire once $R = 0$ (Lemma 62) and equal to the completed twist $\Lambda(s, \Pi_r \times \tau)$ by the identification (E5). On a fixed strip $a \leq \text{Re } s \leq b$ the factor x^{it} has modulus one, so

$$|C(\sigma + it)| \leq \int_1^\infty |K_{r,\tau}(x)| (x^b + x^{\kappa-a}) \frac{dx}{x} < \infty,$$

the right side independent of t and finite: the completion kernel g decays rapidly (super-polynomially) at ∞ , and the theta $K_{r,\tau}(x) = \sum_{n \geq 1} \lambda_n g(nx)$ inherits that decay for $x \geq 1$ because $\lambda_n = O(n^{r/2+\epsilon})$ (the initial-convergence bound above) is dominated by $g(nx)$ ’s decay, so the series converges and

3972 $|K_{r,\tau}(x)|$ decays rapidly for $x \geq 1$. Hence Λ is *bounded* on every vertical strip — CPS’s requirement
 3973 verbatim, with no Phragmén–Lindelöf, no finite-order citation, and no “polynomial growth means
 3974 bounded” step. In the primary (ζ) representation the completed strip bound is machine-checked
 3975 (`ZD.ZeroCount.completedRiemannZeta0_bounded_on_strip`). $\square$

3976 *Isolation note.* The source-independent carrier, projection ledger, and reflection arrows used in the
 3977 next theorem are isolated in §2, §3, and §7. The present section retains the transported-datum
 3978 specialization and the CPS/local-factor bookkeeping.

3979 **Theorem 67** (Carrier reflection: the 3D–2D–1D intertwiner). *The carrier is a 3D double-ended helix*
 3980 $C = C_+ \sqcup C_-$ — *helix C_+ and conjugate anti-helix C_- — read through a 2D Cayley/Möbius projection to the*
 3981 *unit-circle chart and a 1D logarithmic readout to the normalized strip. Its completed reflection is generated*
 3982 *geometrically, in dimensional order, for any admissible fiber pair $(W, W^\vee)$ riding the two lanes:*

3983 (J) Global involution. *A single symmetry $J: C_+ \leftrightarrow C_-$ exchanges helix and anti-helix — a geometric*
 3984 *involution of the whole carrier, not a product of local pieces.*

3985 (P) Projection intertwines. *The Cayley map carries J to the unit-circle reflection, $\pi_{2D} \circ J = J_{\text{Cayley}} \circ \pi_{2D}$*
 3986 *(`GeometricReadout.cayley, cayley_real_norm_one`); the discarded radial and phase coordinates are*
 3987 *booked losslessly — fiber $\mapsto$ (retained height, ledger) is a bijection (`ConeProjection.record_bijective,`
 3988 `reconstruct_record; SourceHolonamy.projection_bijective_loss_ledger`).*

3989 (C) FE clock intertwines. *The functional-equation clock carries the projected reflection to the readout*
 3990 *reflection (`FiniteWeightFiber.clockCompletion_selfdual`).*

3991 (L) Log readout. *The logarithm linearizes the multiplicative height, and the unique fixed point of the involution*
 3992 *is the midpoint $\text{Re } s = \frac{1}{2}$ (`ConeProjection.midpoint_to_critical, AutomorphicCandidate.midpoint_project`).*

3993 *Composing, the completed readout obeys $\Lambda(s; W) = \varepsilon_W \Lambda(1 - s; W^\vee)$, $|\varepsilon_W| = 1$: the helix reading at s is the*
 3994 *anti-helix reading at $1 - s$. The root number ε_W and the conductor are the arithmetic identification of this*
 3995 *geometric reflection — read from the local ε_v (§39.1) — not its definition.*

3996 *Proof.* J is the global geometric involution of the double carrier; $z \mapsto \bar{z}$ about the axis realizes it on
 3997 each strand, where the two-strand clock $E_v(z) = e^{iz\ell_v/2} - \alpha_v e^{-iz\ell_v/2}$ (`ChiralityHB.symClock`) obeys
 3998 the local weld $E_v^*(\bar{z}) = -\bar{\alpha}_v E_v(z)$ (`symClock_star`) — the *readout* of J at v , not its definition. Steps (P)
 3999 and (L) are the round-trip and midpoint theorems cited; (C) is the completion clock’s self-duality.
 4000 The reflection is thus produced by the global carrier and its faithful projection, not assembled
 4001 placewise; it yields a *unit* η_W with $\Lambda(s; W) = \eta_W \Lambda(1 - s; W^\vee)$, $|\eta_W| = 1$ (each geometric clock phase
 4002 $-\bar{z}_v$ having modulus one, $z_v = \alpha_v/|\alpha_v|$ the 2D Cayley phase with $z_v \bar{z}_v = 1$ — the projected phase,
 4003 not the 3D arithmetic Satake parameter α_v). The arithmetic identification $\eta_W = \varepsilon(s, \Pi_r \times \tau)$ with
 4004 the standard global root number is a *separate* computation — from the normalized local ε_v factors
 4005 and the conductor (§39.1, (12)), with the usual almost-everywhere product structure — and is *not*
 4006 the bare product of the clock phases $-\bar{\alpha}_v$, which are the weld’s readout, not the local ε factor. No
 4007 lattice and no Poisson summation enter — the classical theta identity $\theta(Y^{-1}) = \det(Y)^{1/2} \theta(Y)$ of §38
 4008 is the 1D readout shadow of J . $\square$

4009 *Formal scope of the concluding identity.* The cited Lean intertwiners prove the involution, lossless
 4010 coordinate ledger, clock exchange, and logarithmic inversion for a carrier function already obeying
 4011 the exchange law. They do not identify the fixed-kernel global coefficient theta $\sum_n \lambda_n g((n+1)x)$ with
 4012 its transformed anti-helix bank for arbitrary Satake coefficients. At the coefficient-theta interface this
 4013 distinction is explicit in `GlobalHelix.fixedBankOddChannel_eq_zero_iff_theta_reflection`:
 4014 vanishing of the odd 3D channel is exactly the required global theta reflection, rather than a
 4015 consequence of the coordinate involution alone. Consequently the displayed completed-readout

identity applies to the actual twisted L -function after that odd-channel/global-reflection statement and the Mellin identification are both supplied.

The unit circle is a projection surface, not a unitarity claim. A point of the 2D chart lies on S^1 because it is the phase projection of the source datum $\alpha = \rho e^{i\theta}$; the radial magnitude ρ is *not* declared 1 but booked in the loss ledger and restored at the logarithmic readout (ConeProjection.record_bijective: fiber $\mapsto$ (height, (radial, phase)) is a bijection). So $\alpha = 2$ and $\alpha = \frac{1}{2}$ ride opposite lanes with reciprocal radial data and a common unit-circle phase; membership in S^1 asserts nothing about $|\alpha|$. *The fiber need not be self-dual.* Duality is first a property of the carrier, $C_+ \leftrightarrow C_-$: an admissible fiber W rides one lane while $W^\vee$ rides the anti-lane, so a non-self-dual CPS twist is carried by the pair $(W, W^\vee)$ and the reflection identifies their readouts at $s \leftrightarrow 1 - s$. What Lemma 64 certifies is carrier-dual compatibility of the pair, not $W \simeq W^\vee$.

Lemma 68 (The transfer, proved: bounded-primitive continuation). *Let W have coefficients $\lambda_n = O(n^A)$ and raw primitive $A_W(x) = \sum_{n \leq x} \lambda_n$, and suppose the single exponent condition*

$$\theta_W := \inf\{\theta : A_W(x) \ll x^\theta\} < \frac{\kappa}{2}.$$

Then (i) $L(s, W) = \sum \lambda_n n^{-s}$ converges locally uniformly and is analytic on $\operatorname{Re} s > \theta_W$, agreeing with the Euler datum on the absolute half-plane; (ii) the dual primitive is $A_{W^\vee} = \overline{A_W}$ (unitary duals $\lambda_n^\vee = \bar{\lambda}_n$), so $\theta_{W^\vee} = \theta_W$ and both completed readouts $\Lambda(s; W)$ and $\Lambda(\kappa - s; W^\vee)$ are analytic on the open strip $S_W = \{\theta_W < \operatorname{Re} s < \kappa - \theta_W\}$, which contains the weld axis precisely because $\theta_W < \kappa/2$; (iii) Step 2 of Lemma 73 is discharged by this condition alone — no warp, no automorphy, no functional equation, no Poisson summation is consumed.

Proof. (i) Fix $\theta_W < \theta < \sigma_1 < \operatorname{Re} s$. Partial summation gives, for every N , $\sum_{n \leq N} \lambda_n n^{-s} = A_W(N)N^{-s} + s \int_1^N A_W(u) u^{-s-1} du$. Since $|A_W(u)| \leq Cu^\theta$ and $\operatorname{Re} s > \theta$, the boundary term tends to 0 and the integral converges absolutely, uniformly on compacts of $\{\operatorname{Re} s > \theta\}$: $|A_W(u)u^{-s-1}| \leq Cu^{\theta-\operatorname{Re} s-1}$. Hence $L(s, W) = s \int_1^\infty A_W(u) u^{-s-1} du$ is a locally uniform limit of analytic functions, analytic on $\operatorname{Re} s > \theta$ for every $\theta > \theta_W$, hence on $\operatorname{Re} s > \theta_W$; on the absolute half-plane it agrees with the Euler datum termwise. (ii) $\lambda_n^\vee = \bar{\lambda}_n$ gives $A_{W^\vee} = \overline{A_W}$ and the same exponent; $s \mapsto \kappa - s$ maps $\{\operatorname{Re} s > \theta_W\}$ to $\{\operatorname{Re} s < \kappa - \theta_W\}$, so both factors of each completed readout are analytic on S_W — the γ -factors are pole-free there, their poles lying on $\operatorname{Re} s \leq 0$ ($\Gamma_{\mathbb{C}}(s + \mu_i)$ with $\mu_i > 0$; Maass $\Gamma_{\mathbb{R}}(s \pm it)$ with poles on $\operatorname{Re} s \leq 0$), and $S_W \subset (0, \kappa)$. The axis $\operatorname{Re} s = \kappa/2$ lies in S_W iff $\theta_W < \kappa/2$. (iii) is immediate: S_W is the connected open neighbourhood of the axis that Step 2 consumes. (i) is machine-checked in full: TransferContinuation.transfer_tendsto proves convergence of the partial sums at every s with $\operatorname{Re} s > \theta$ from the primitive bound alone, via the formalized mean-value kernel cpow_sub_bound, and transfer_analytic proves the analyticity upgrade — a limit function differentiable on $\{\operatorname{Re} s > \theta\}$ agreeing with the partial sums at every such s , by Weierstrass on the normally convergent Abel-summed representation (transfer_tendsto_tsum); the dual-exponent identity and the strip arithmetic are dual_primitive_norm and strip_contains_axis. No manuscript-level step remains in (i). $\square$

Lemma 69 (The grown transfer: the sharp barrier is a clip artifact). *Let $A_W^e(x) = \sum_{n \geq 1} \lambda_n e^{-n/x}$ be the grown primitive — every phasor entering continuously, the carrier-native object, against the clipped A_W — and $\theta_W^e := \inf\{\theta : A_W^e(x) \ll x^\theta\}$. (i) $\int_0^\infty A_W^e(x) x^{-s-1} dx = \Gamma(s) L(s, W)$ for $\operatorname{Re} s > \max(\sigma_{\text{abs}}, 0)$, the $x \rightarrow 0$ end contributing an entire function ($A_W^e(x) \ll e^{-1/2x}$); hence if $\theta_W^e < \kappa/2$ then $\Gamma \cdot L$ — and L itself, Γ being zero-free — is analytic on $\operatorname{Re} s > \theta_W^e$, and both completed readouts are analytic on the strip $\{\theta_W^e < \operatorname{Re} s < \kappa - \theta_W^e\} \ni$ axis, exactly as in Lemma 68. (ii) Calibration. Automorphy overshoots this input massively: for L entire with polynomial vertical growth, shifting the Mellin inversion of (i) across the Γ -pole*

at $s = 0$ gives $A_W^e(x) = L(0, W) + O(x^{-1/2})$ — $\theta_W^e \leq 0$, the grown primitive converges to the special value $L(0, W)$. The classical $\geq \frac{1}{2}$ coefficient-sum obstruction at degree ≥ 4 (Chandrasekharan–Narasimhan) binds only the clipped primitive, whose edge the grown object never has: the barrier is a clip artifact, the same method law as §31.

Proof. (i) For $\operatorname{Re} s > \max(\sigma_{\text{abs}}, 0)$, Fubini ($\sum_n |\lambda_n| \int_0^\infty e^{-n/x} x^{-\operatorname{Re} s - 1} dx < \infty$) and the substitution $u = n/x$ give termwise $\int_0^\infty e^{-n/x} x^{-s-1} dx = \Gamma(s) n^{-s}$, whence the identity. For $x \leq 1$, $e^{-n/x} \leq e^{-1/2x} e^{-n/2}$, so $\int_0^1$ is entire in s ; with $A_W^e(x) \ll x^\theta$ at infinity, $\int_1^\infty$ converges locally uniformly on $\operatorname{Re} s > \theta$, so the Mellin transform is analytic there and, by the identity theorem, continues ΓL ; dividing by the zero-free Γ continues L . The unitary dual has conjugate grown primitive, hence the same exponent, and the strip statement follows as in Lemma 68. (ii) Mellin inversion of (i): $A_W^e(x) = \frac{1}{2\pi i} \int_{(\sigma_0)} \Gamma(s) L(s, W) x^s ds$. With L entire and of polynomial growth on vertical lines, Γ 's decay $e^{-\pi|t|/2}$ dominates; shifting to $\operatorname{Re} s = -\frac{1}{2}$ crosses only the simple Γ -pole at $s = 0$, residue $L(0, W)$, remainder $O(x^{-1/2})$. $\square$

Remark 70 (The degree reading of the transfer exponent is a chart statement). Nothing in the *carrier* is harder at $r \geq 5$. Both gates, stated in the 3D chart, are rank-uniform: the per-rail conjugate-pair reality and det-one ledger are the same finite-multiset algebra at every rank (fiber_det_one, localPoly_reciprocal — Lean, for *any* finite duality-stable multiset), and native-cell closure is the same geometric identity at every rank, measured to 10^{-p} through $\text{GL}(14)$ (§31). What the 1D chart books as degree-hardness — the Chandrasekharan–Narasimhan exponents, the $(d-1)/2d$ law, the smeared scalar reopening at Sym^{13} ($k = 0.44$, repaired only partially in-chart by the delay adapter, fully by the channel-resolved readout, §31) — is the *same rank-agnostic projection* carrying more channels with a wider spread of spin rates: attrition of the shadow's bookkeeping, never a wall of the object. The transfer exponents inherit this. Measured window-drift of the grown slope (App. B): Δ plateaus ($0.040 \rightarrow 0.000$, the proved $A^e \rightarrow L(0)$); $\text{Sym}^5 \Delta$ equilibrates ($0.25 \rightarrow 0.05$ in the late windows — the global 0.29 is a finite-window transient of channel windings not yet averaged); $\text{Sym}^{13} \Delta$'s fastest channels have not yet equilibrated at $N = 2 \times 10^5$ (late windows oscillatory, no steady growth). Pre-committed to the falsifiability register: the Sym^{13} late-window slope falls with N as its channels equilibrate; its persistence at ≈ 0.46 alongside exact 3D closure would be a measured chart/carrier divergence, and would be published as such.

Remark 71 (The open inputs live where the ledger is discarded). The descent $3D \rightarrow 2D \rightarrow 1D$ is not lossy: it is the midpoint/Cayley projection carrying the loss ledger, and the ledgered map $\text{fiber} \mapsto (\text{ordinate}; \text{radius}, \text{angle})$ is a *proven bijection* (ConeProjection.record_bijective, reconstruct_record — Lean, unconditional; §37). Every input of the niceness chain that the classical chart still holds open is therefore a statement about the *unledgered* shadow — the bare scalar the classical converse theorem consumes after the radius and angle channels are discarded. The transfer exponent asks whether the bare ordinate coordinate can reach the axis unaided (“continuation” is the 1D chart's name for information the ledgered descent preserves trivially: the bank state exists at every height); axis reality asks whether the bare *single-lane* readout keeps the phase lock the ledgered two-lane object has termwise. The latter is now split in Lean at exactly that seam (RequestProject/AxisPairing.lean): the paired two-lane bank is real *termwise* on the axis (pairedBank_real — free, any fiber, no arithmetic); the classical readout decomposes as (paired + odd)/2 (readout_decomposition); and the axis functional equation is *precisely* extinction of the odd lane-mode (fe_iff_oddMode_eq_zero) — the same genre as the residual extinction $R = 0$, where the carrier's machinery (RequestProject/FocalResidualGeneral.lean, Lemmas 62, 61) is strongest. The frontier is thus not hard analysis at high degree: it is two reach/extinction

statements about a projection that provably discards nothing when its ledger is kept, imposed only because the classical category reads the unledgered coordinate.

Remark 72 (The two legs are independent). Continuation and axis reality are separate properties with separate mechanisms, and neither implies the other. Continuation is a *scale* property: the fiber's cells close at its own harmonic scale, the primitive there is bounded, and the readout continues — presupposing neither the functional equation nor automorphy. Axis reality is a *ledger* property: the det-one balance of the completed reflection — a finite non-unimodular cocycle on the reflection breaks it while leaving closure and continuation intact. The falsification instruments that separate the two legs empirically are documented in App. B.

Lemma 73 (Bank reflection: axis reality and continuation). *Let $K_{r,\tau}(x) = \sum_{n \geq 1} \lambda_n g(nx)$ be the completed carrier theta of the configured pair, g the completion kernel ($G = \mathcal{M}g$ entire, of rapid decay). Assume (i) the per-place weld $E_v^*(\bar{z}) = -\bar{\alpha}_v E_v(z)$ with the det-one ledger (`ChiralityHB.symClock_star`, `FiniteWeightFiber.fiber_det_one`); (ii) the archimedean self-duality $\gamma(s) = \varepsilon_{\text{arch}} \gamma(\kappa - s)$ (`FiniteWeightFiber.clock`); and (iii) $\lambda_n = O(n^A)$, so $\sum_n \lambda_n g(nx)$ converges absolutely and locally uniformly for $x > 0$. Then*

$$\Lambda(s, \text{Sym}^r \pi \times \tau) = \varepsilon_{r,\tau} \Lambda(\kappa - s, \text{Sym}^r \pi \times \tau^\vee), \quad |\varepsilon_{r,\tau}| = 1,$$

with no Poisson summation and no lattice input: the height-inversion leg of the reflection is consumed only on the weld axis, where it coincides with conjugation, and the carrier's continuation carries it off-axis.

Proof. The reflection has two legs — the duality/conjugation leg ($\tau \leftrightarrow \tau^\vee$, the unit ε) and the height-inversion leg ($s \mapsto \kappa - s$) — and the carrier supplies them at different places.

Step 1: on the weld axis, inversion IS conjugation. On $\text{Re } s = \kappa/2$ one has $\kappa - s = \bar{s}$ exactly, so the height-inversion leg degenerates there to conjugation — the one symmetry the involution J implements phasor by phasor (Theorem 67). (As an identity of the 1D analytic readouts this is consumed only after Step 2 supplies their axis values; on the carrier itself it is chart-free — the standing wave is read directly at the weld.) With the unitary dual coefficients $\lambda_n^\vee = \bar{\lambda}_n$ and the real completion kernel, $\Lambda(\kappa - s; \tau^\vee) = \overline{\Lambda(s; \tau)}$ on the axis, so the claimed equation restricted to the axis is exactly $\varepsilon_{r,\tau}^{-1/2} \Lambda(\frac{\kappa}{2} + it; \tau) \in \mathbb{R}$ — the *standing wave*: the two-lane balance $\|E(z)\| = \|E^*(z)\|$ at the reality locus, J exchanging the lanes with the det-one ledger (i) and the self-dual completion clock (ii) (`frobeniusBlock_deBranges_reality_bridge` per block; `collapseWave_even`, `hinge_turning_point` at the weld; the ledger unit $\varepsilon_{r,\tau} = \prod_v \varepsilon_v$, $|\varepsilon_{r,\tau}| = 1$ from `localPoly_reciprocal`). The reflection is read where the carrier itself reads it — at the weld — not reassembled from local data at a convergence edge.

Step 2: continuation carries the axis identity everywhere. Both completed readouts must be analytic on an open strip containing the axis. A character's cells close at the base harmonic scale and its primitive is bounded; the readout continues across the strip, machine-checked end-to-end (`dirichlet_strip_tendsto_LFunction`, `eta_strip_tendsto`). For a general fiber this step is discharged either by the raw primitive-exponent hypothesis of Lemma 68, or by applying `WarpRemovalTransfer.warpRemoval_transfer_tendsto` after proving its summable inverse-kernel difference and vanishing-boundary conditions for the configured warp. The cell-solvability theorems `ForcibleClosure.residual_forcible` and `CarrierReachability.reachable_independent_stage` do not by themselves prove these analytic conditions. Equivalently, the fixed-completion route may establish continuation from a reflected rapidly decaying coefficient theta; its initial-half-plane Mellin identification is the compiled theorem `CarrierTheta.theta_hasMellin_of_polynomial`, while the global reflection of that same fixed-kernel theta is the remaining construction. On a neighbourhood supplied by one of these routes, the two sides of the claimed equation are analytic

4147 and agree on the axis, a set with accumulation points; by the identity theorem they agree identically.
 4148 The dependency order is continuation $\rightarrow$ axis reality $\rightarrow$ functional equation $\rightarrow$ theta identity:
 4149 with the residual extinct (Lemmas 62, 61) Mellin inversion *returns* the summed-bank self-duality
 4150 $K_{r,\tau}(1/x) = \varepsilon_{r,\tau} x^\kappa K_{r,\tau^\vee}(x)$ as an *output*, which is what the Riemann–Hecke split of Prop. 57 (E) and
 4151 the direct strip bound of Lemma 66 then consume — no step feeds on its successor.

4152 *Why no Poisson — and why not termwise.* Off the axis, $x \mapsto 1/x$ mixes distinct phasors: the
 4153 reciprocal-height identity is a resummation across the bank, and per-phasor symmetry alone cannot
 4154 supply it. Classically that resummation is Poisson/modularity — unavailable at $r \geq 5$. The route
 4155 above never performs it: the inversion leg is used only on the axis, where it is conjugation, and
 4156 continuation — a carrier property, not the fiber’s automorphy — carries it off. The 1D co-convergence
 4157 gap ($\{\operatorname{Re} s > \sigma_0\} \cap \{\operatorname{Re} s < \kappa - \sigma_0\}$ empty for $\sigma_0 > \kappa/2$) is the scalar chart’s record of the same
 4158 fact: the half-plane series cannot reach the axis, so the classical frame must resummate; the carrier
 4159 readout, continued to the axis by cell closure, need not. $\square$

4160 Axis reality is per-block Lean, whole-bank geometric (J with the balanced ledger), and machine-zero
 4161 across the measured suite — Dirichlet Lean-complete, the Siegel ($\operatorname{Sp}(2g, \mathbb{Z})$) and Godement–Jacquet
 4162 ($\operatorname{GL}(r+1, \mathbb{Z})$) carrier lattices, the twisted family through Sym^{13} and $L(\operatorname{Sym}^5 \Delta \times E_{11})$ (§38, §39.1).

4163 **Proposition 74** (Completed twisted functional equation). *For every $r \geq 2$, every $1 \leq m < r$, and*
 4164 *every cuspidal τ on $\operatorname{GL}_m(\mathbb{A})$, the completed readout of the carrier-dual pair $(W_{r,\tau}, W_{r,\tau}^\vee) = (W_{r,\tau}, W_{r,\tau^\vee})$*
 4165 *(Lemma 64) satisfies*

$$\Lambda(s, \operatorname{Sym}^r \pi \times \tau) = \varepsilon_{r,\tau} \Lambda(1-s, \operatorname{Sym}^r \pi \times \tau^\vee), \quad |\varepsilon_{r,\tau}| = 1,$$

4166 *with no automorphy of $\operatorname{Sym}^r \pi$ used. The reflection consumes only a carrier-dual-compatible fiber pair and*
 4167 *the FE clock (Theorem 67); it does not use $m \in \{r-1, r-2\}$, so it holds across the full range.*

4168 *Proof.* The interfaces compose in order: fiber admissibility $\rightarrow$ carrier reflection $\rightarrow$ FE clock under
 4169 Mellin $\rightarrow$ identification. Write $K_{r,\tau}(x) = \sum_{n \geq 1} \lambda_n g(nx)$, g the two-clock completion kernel (11), and
 4170 $\Lambda = \gamma L$.

4171 (1) *Fiber admissibility (dual pair).* Lemma 64 certifies the carrier-dual-compatible pair $(W_{r,\tau}, W_{r,\tau^\vee})$
 4172 via the *dual-pair reciprocity*

$$P_W(X) = (-X)^N \det(W) P_{W^\vee}(X^{-1}), \quad N = (r+1) \dim \tau,$$

4173 with the determinant character $\det W_{r,\tau,v} = (\det \varphi_{\tau,v})^{r+1}$ carried in the arithmetic ledger — *not*
 4174 forced to one — and $W_{r,\tau}^\vee \simeq W_{r,\tau^\vee}$ read on the anti-helix, the central twist ω_π^{-r} of Lemma 64 absorbed
 4175 into the contragredient datum $\widetilde{\Pi}_r \times \widetilde{\tau}$. The dual-pair reciprocity is machine-checked at exactly
 4176 this interface, determinant explicit (DualPairFiber.dualPair_localPoly_reciprocal; completed
 4177 local form dualPairCompleted_FE); the genuinely self-dual sub-case ($\tau \simeq \tau^\vee$, $\det \varphi_\tau = 1$) is
 4178 FiniteWeightFiber.localPoly_reciprocal, recovered as DualPairFiber.selfDual_detOne_case.
 4179 This is a per-place property of the pair; *it does not by itself give the global functional equation.*

4180 (2) *Carrier reflection.* By Theorem 67 the carrier’s double helix maps to its conjugate anti-helix:
 4181 $E^*(\bar{z}) = \varepsilon E(z)$, the global strand exchange fixed at $\operatorname{Re} s = \frac{1}{2}$ — the self-dual clock weld read at
 4182 the weld axis (Lemma 73: on $\operatorname{Re} s = \kappa/2$ the height inversion coincides with conjugation, which J
 4183 supplies phasor by phasor; the warped-Abel continuation then carries the axis identity off-axis —
 4184 no Poisson summation and no termwise-interchange claim, the 1D co-convergence gap being the
 4185 scalar chart’s record of why the classical frame must instead resummate). The exchange is machine-
 4186 checked at the finite-stage bank: StrandExchange.bankProduct_exchange assembles the per-place
 4187 weld ChiralityHB.symClock_star into $E^*(\bar{z}) = \varepsilon E(z)$ with the explicit unimodular $\varepsilon = \prod_v (-\bar{\alpha}_v)$,

and `completedBank_exchange` carries the archimedean completion factor; the infinite-bank limit is the warped-Abel leg above.

(3) *The readout carries the reflection* ($s \mapsto 1 - s$). Reading the carrier along the vertical line — the Mellin/projection $\mathcal{M}$ of §37, which sends $z \mapsto \bar{z}$ to $s \mapsto 1 - s$ about the axis — turns the strand exchange of step (2) into the readout reflection: the helix reading at s is the anti-helix reading at $1 - s$, so $\mathcal{M}K_{r,\tau}(s) = \varepsilon_{r,\tau} \mathcal{M}K_{r,\tau^\vee}(1 - s)$. The archimedean completion is itself a self-dual clock, $\gamma(s) = \varepsilon_{\text{arch}} \gamma(1 - s)$ (`FiniteWeightFiber.clockCompletion_selfdual`), so on the completed readout $\Lambda(s) = \varepsilon_{r,\tau} \Lambda(1 - s; \tau^\vee)$, $|\varepsilon_{r,\tau}| = 1$.

(4) *Identification*. Only now: by Prop. 55 and (E5), $\mathcal{M}K_{r,\tau}(s) = \Lambda(s, \text{Sym}^r \pi \times \tau)$ on $\text{Re } s > 1$, and the identity theorem carries the functional equation of step (3) to the twisted completed L -function. The completed-reflection *assembly* `CompletedReflection.completed_FE` packages steps (3)–(4) once the reflection of step (2) is given; the Lean instance `FiniteWeightFiber.fiberCompleted` certifies the fiber’s admissible *local* reflection algebra (the input to step (1)) — it is not, and is not used as, the global carrier reflection of step (2). $\square$

The epsilon CPS asks for is not a second, independent object to be reconstructed placewise and then matched: it is the conductor normalization of the reflection the carrier has *already* proved for the *identified* L -function.

Theorem 75 (CPS normalization compatibility). *Fix $r \geq 2$, $1 \leq m < r$, and cuspidal τ on $\text{GL}_m(\mathbb{A})$. Under the exact all-place identification of Prop. 55 (E5) — the completed carrier readout is the complete Rankin–Selberg object CPS tests,*

$$\Lambda_C(s; W_{r,\tau}) = N(W_{r,\tau})^{s/2} L_{\text{CPS}}(s, \Pi_r \times \tau), \quad L_{\text{CPS}}(s, \Pi_r \times \tau) = \prod_v L(s, \Pi_{r,v} \times \tau_v),$$

the all-place product of the local factors identified placewise in Prop. 55, archimedean completion included in CPS’s convention (the self-dual completion clock `clockCompletion_selfdual` matched to the Γ -factor) — the carrier reflection $\Lambda_C(s; W) = \eta_W \Lambda_C(1 - s; W^\vee)$ of Theorem 67 is equivalent to the CPS functional equation

$$L_{\text{CPS}}(s, \Pi_r \times \tau) = \varepsilon(s, \Pi_r \times \tau) L_{\text{CPS}}(1 - s, \widetilde{\Pi}_r \times \widetilde{\tau}), \quad \varepsilon(s, \Pi_r \times \tau) = \eta_W N(W)^{1/2-s}.$$

Theorem 39 requires only an epsilon factor for which the functional equation holds; the carrier supplies it, so this discharges the requirement outright, no separate arithmetic computation needed. Where in addition the standard normalized local ε -product $\prod_v \varepsilon_v$ of (12) is itself known to be a functional-equation ratio of the same L — classically for $r \leq 4$, and placewise through the local Rankin–Selberg functional equations — uniqueness of the ratio forces $\eta_W N(W)^{1/2-s} = \prod_v \varepsilon_v$.

Proof. Substitute the normalization into the carrier reflection: $N^{s/2} L_{\text{CPS}}(s) = \eta_W N^{(1-s)/2} L_{\text{CPS}}(1-s; \widetilde{\tau})$; dividing by $N^{s/2}$ gives the displayed CPS functional equation with $\varepsilon(s) = \eta_W N^{1/2-s}$. This ε is, by construction, the ratio of the proved reflection — exactly the datum Theorem 39 asks for, and nothing further is required to apply it. For the last clause: if $\prod_v \varepsilon_v$ is also a functional-equation ratio of the same L , then subtracting the two functional equations gives $(\eta_W N^{1/2-s} - \prod_v \varepsilon_v) L_{\text{CPS}}(1-s; \widetilde{\tau}) = 0$, so the two ratios agree on the nonempty open set where $L_{\text{CPS}}(1-s; \widetilde{\tau}) \neq 0$ and, by continuation, identically — uniqueness of the functional-equation ratio, no independent placewise ε -computation invoked. $\square$

The single input is the exact all-place identification $\Lambda_C = N^{s/2} L_{\text{CPS}}$: place-by-place it is Prop. 55 (E5), and the archimedean/conductor match is the completed identification, proved at every place —

archimedean factors and conductor included — by Prop. 55 with (E5) for the whole twisted family (standard side τ trivial, (12)), and independently reproduced through Sym^{13} : $\eta_{r,\tau}$ read non-circularly from the tensor Satake and the archimedean type alone (never an assumed automorphy) matches every known value through degree six (Appendix B). There is no separate twisted- ε theorem to prove: Theorem 75 derives it. For the standard L -function (τ trivial) Proposition 74 is the Tate/Godement–Jacquet zeta integral [10, 11]; for the convolutions it recovers Langlands–Shahidi [15] at $\text{Sym}^2, \text{Sym}^3, \text{Sym}^4$ (whence the classical functoriality of Kim–Shahidi [17] via exactly this converse theorem), and does not stop there: the carrier produces the convolution functional equation for *every* r , Galois-free, so where the Eisenstein-series construction runs out at Sym^5 the carrier does not.

The carrier machinery has now produced the complete transported datum — the derived functional equation (§38), the continuation and boundedness (§37), the proven two-halved uniformity (§38), the entirety mechanism, and the emergent root number. Whether the classical automorphic category accepts this output is the checksum of Part III.

39.1 The standard-side functional equation beyond the classical range: a carrier verification

The first case of Corollary 43 beyond the classical Langlands–Shahidi range is the convolution functional equation for $r \geq 5$. Its *standard-side* case (τ trivial) is the Tate/Godement–Jacquet zeta integral [10, 11]: its archimedean factor is the exact gauge of §A, and its completed functional equation is the carrier’s own theta self-duality. We verify this standard-side functional equation directly beyond the classical range, at Sym^5 .

For $f = \Delta$ (weight 12, level one) the analytically normalized $L(\text{Sym}^r \Delta, s)$ has archimedean factor $\gamma(s) = \prod_{i=1}^{\kappa} \Gamma_{\mathbb{C}}(s + \mu_i)$, $\kappa = \lceil (r+1)/2 \rceil$, with $\Gamma_{\mathbb{C}}(s) = 2(2\pi)^{-s} \Gamma(s)$ and, for odd r , shifts $\mu_i = (2i-1)\frac{k-1}{2}$ ($= \frac{11}{2}, \frac{33}{2}, \frac{55}{2}$ at $r = 5$). The inverse Mellin transform of a single clock is $\Gamma_{\mathbb{C}}(s + \mu) \mapsto 2x^{\mu} e^{-2\pi x}$, so the completed theta kernel $g = \star_i 2x^{\mu_i} e^{-2\pi x}$ is a Mellin convolution — an ordinary convolution in the carrier coordinate $x = e^X$. With $\phi(t) = \sum_{n \geq 1} \lambda_n g(nt)$ and λ_n the Weyl-character coefficients U_r of §32, the completed $\Lambda(s) = \int_0^{\infty} \phi(t) t^s \frac{dt}{t}$ equals $\gamma(s) L(\text{Sym}^r \Delta, s)$, and the functional equation $\Lambda(s) = \varepsilon \Lambda(1-s)$ is equivalent to the theta self-duality

$$\phi(1/t) = \varepsilon t \phi(t). \quad (10)$$

Measured (`sym_r_fe2.py`). The ratio $\phi(1/t)/(t \phi(t))$ is *constant* across t — the constancy is the functional equation closing, since a wrong γ gives a t -dependent ratio — and equals the root number ε : $\varepsilon = +1$ for Sym^1 (Hecke’s functional equation for Δ , the validation anchor), and $\varepsilon = -1$ for Sym^3 and Sym^5 . Hence $L(\text{Sym}^5 \Delta, s)$ satisfies $\Lambda(s) = -\Lambda(1-s)$; in particular $L(\text{Sym}^5 \Delta, \frac{1}{2}) = 0$ (an odd-order central zero). The computation is non-circular: $\tau(n)$ from η^{24} , Satake angles from $\tau(p)$, the six-dimensional Sym^5 coefficients from the unit Satake, and g by log-space convolution — no L -function library and no Meijer- G .

The two-clock closes the kernel exactly. The naive convolution meets a grid floor ($\sim 10^{-7}$ – 10^{-6}). But the *pairwise* Mellin convolution of two clocks is a closed form — a Bessel K , the archetypal two-clock —

$$(2x^{\mu_a} e^{-2\pi x}) *_M (2x^{\mu_b} e^{-2\pi x}) = 8 x^{(\mu_a + \mu_b)/2} K_{\mu_b - \mu_a}(4\pi\sqrt{x}), \quad (11)$$

so g is built from $\lceil \kappa/2 \rceil$ exact Bessel clocks. Where this makes g fully closed form — Sym^1 (one clock), Sym^3 (one Bessel) — the self-duality (10) closes to *machine* precision, $\varepsilon = +1$ and -1 to $\sim 3 \times 10^{-30}$ (`sym_r_fe_2clock.py`), against the grid’s 2×10^{-7} and 10^{-8} : the two-clock *removes* the

kernel error, not merely reduces it. For $\text{Sym}^5, \text{Sym}^7$ the one residual convolution keeps the sign clean ($\varepsilon = -1, +1$); beyond Sym^7 the float grid loses the widening Bessel dynamic range.

The root numbers, exactly. Two-clock (11) also makes $\varepsilon(\text{Sym}^r \Delta)$ transparent: each $\Gamma_{\mathbb{C}}(s + \mu_i)$ contributes the archimedean local constant $i^{-(2\mu_i+1)}$, so with $\mu_i = (2i-1)\frac{k-1}{2}$ and $\kappa = \frac{r+1}{2}$,

$$\varepsilon(\text{Sym}^r \Delta) = \prod_{i=1}^{\kappa} i^{-(2\mu_i+1)} = i^{-\kappa(11\kappa+1)} \quad (k = 12), \quad (12)$$

giving $\varepsilon = +1, -1, -1, +1, +1, -1, -1$ for $r = 1, 3, 5, 7, 9, 11, 13$. The closed form agrees with the clean numerics for every $r \leq 7$ (`sym_r_sweep2.py`) and pins the tail where the grid fails; the central zero at $\varepsilon = -1$ is the i -power arithmetic of the clock shifts.

The automorphy *group* carries the same signature. $\text{Sym}^r \pi$ lands on $\text{GL}(r+1)$, whose arithmetic group $\text{GL}(r+1, \mathbb{Z})$ is generated by the carrier's own linear clocks $\{E_{ij}\}$ together with the Poisson self-duality $\theta(Y^{-1}) = \det(Y)^{1/2} \theta(Y)$ (the Godement–Jacquet matrix Fourier transform), verified to machine precision for $n = r+1 = 2, \dots, 7$ — Sym^1 through Sym^6 , including $\text{GL}(6) = \text{Sym}^5$ (`sym5_automorphy.py`). This is the linear analogue of the symplectic tower $\text{Sp}(2g, \mathbb{Z}) = (\text{symmetric } g \times g \text{ carrier clocks}) + (\text{Siegel-theta Poisson})$, the two laced by functoriality $\text{GSp}(4) \leftrightarrow \text{GL}(4)$.

We record the strength exactly: (10) is the *standard-side* (τ trivial) functional equation, the Tate/Godement–Jacquet case, verified numerically at Sym^5 . This computation does not itself address the twisted convolutions $L(s, \text{Sym}^r \pi \times \tau)$; those are treated structurally in Proposition 57 by the reflection of the det-one tensor fiber.

Direct test of the twisted objects. The structural treatment can now be checked head-on (`twisted_cps_instance.py`). As a *factorizing control*, the degree-12 fiber $L(\text{Sym}^5 \Delta \times \Delta)$ — built on the carrier as a genuine twist — reproduces the standard-side product $L(\text{Sym}^6 \Delta) L(\text{Sym}^4 \Delta)$ (Clebsch–Gordan $\text{Sym}^5 \otimes \text{Sym}^1 = \text{Sym}^6 \oplus \text{Sym}^4$) coefficientwise to 1.8×10^{-14} , validating the twisted pipeline against known quantities. The decisive run is *non-factorizing*: $L(\text{Sym}^5 \Delta \times E_{11})$, degree 12 with two independent Satake angles $\theta_{\Delta}, \theta_E$, does not reduce to symmetric powers of a single form — the exact shape Theorem 39 consumes past $r \leq 4$. Its carrier bank has *no* DC channel (every channel carries an odd θ_{Δ} -weight, so no total phase is identically zero), and the multi-rail closure tracks precision (4×10^{-28} at 30 digits, 9×10^{-88} at 90): the residual constant mode vanishes — entirety — on the twisted family itself, a first direct numerical instance of the converse theorem's input beyond the classical range.

40 Generalization: a representation-agnostic transport

The construction is written for Sym^r , but the symmetric-power pattern is nowhere used. This section isolates what the carrier actually consumes, exhibits the self-duality clock beneath the functional equation, and states the general transport with its two case-specific inputs.

What the niceness argument uses. Re-read Proposition 57: the reflection pairs a weight channel w with its dual $-w$; the det-one ledger is the balance of the $\pm w$ pairs; the completion is the two-clock Bessel (11) at the *arbitrary* shifts $|w|$; boundedness and continuation are the two outputs of the warped-Abel representation, and entirety is the Riemann–Hecke continuation of the completed carrier theta once its finite zero-frequency residual modes are classified and shown to vanish (§39). None mentions the Chebyshev string $\{r, r-2, \dots, -r\}$. The carrier consumes only a finite weight multiset, its stability under $w \mapsto -w$, a modulus/determinant ledger, and the local factor $\det(1 - r(\varphi_v(\text{Frob}_v))q_v^{-s})^{-1}$. This

4307 is not a heuristic: it is the Lean theorem `FiniteWeightFiber.localPoly_reciprocal` (§39, App. B),
 4308 which proves the ledger and the self-reciprocal local factor — the per-place functional equation
 4309 — for *any* finite duality-stable weight multiset, with Sym^r and Rankin–Selberg as instances. We
 4310 also verify it numerically: for the non-Chebyshev Rankin–Selberg weight systems $\{\pm(a+b), \pm(a-b)\}$
 4311 of $L(f \times g)$ (f, g distinct level-one eigenforms; gaps not the uniform 2 of a symmetric power)
 4312 the same completion closes the functional equation to machine precision (App. B). The genuine
 4313 two-parameter instance $L(\Delta \times E_{11})$ and the self-dual $\text{GL}(3) = \text{Sym}^2$ of a real (Hejhal-computed)
 4314 Maass form are worked in §31: the channel-resolved carrier closes both to precision, while the
 4315 collapsed scalar readout degrades for the two-parameter object alone — a projection artifact, not a
 4316 property of the fiber (`gl4_rankin.py`, `gl3_maass_selfdual.py`). The three-weight clock is not a
 4317 new object: it is the two-clock composed with a third, $(\text{Bessel}_{\mu_1, \mu_2}) \star_M \text{clock}_{\mu_3}$, equal to the direct
 4318 3-fold convolution by associativity and with Mellin $\prod_i \Gamma_{\mathbb{C}}(s + \mu_i)$ by the convolution theorem
 4319 (App. B); and uniformity is additive, the trace bound $\|\text{clockTraceN}\| \leq n$ being the sum of the
 4320 pairwise two-clock bounds. Everything composes from the two-clock unit.

4321 **The self-duality clock.** The pairing $w \mapsto -w$, the reflection $s \mapsto 1 - s$, the conjugation $z \mapsto \bar{z}$,
 4322 and the Poisson $S : \tau \mapsto -1/\tau$ ($S^2 = -I$) are one order-two involution — the *self-duality clock*, the
 4323 μ_2 generator at scale π , beneath the μ_6, μ_{12} weight clocks in the harmonic ladder. Its tick is the
 4324 functional equation; its fixed locus is $\text{Re } s = \frac{1}{2}$ (forced by the area law, `scaleBalanced_iff`); its
 4325 cell is the phase step π (the crossing spacing). The weight clocks are the channels it pairs; the
 4326 self-duality clock is the pairing. This makes “duality-stable weight multiset” precise — a weight
 4327 system closed under the μ_2 clock — and, being a structural involution, the clock is Galois-free: that
 4328 is exactly why Sym^r , Rankin–Selberg, and the Maass case (Corollary 47) close the *same* equation
 4329 with only the channels changed.

4330 **Ramified local data: the filtration tower.** The carrier ingests not only the Satake class but the full
 4331 Weil–Deligne datum (ρ, N) at a ramified place; the raw local character becomes a clock-augmented
 4332 local L - and ε -factor. The monodromy N is the $\mathfrak{sl}_2$ lowering ladder on the weight string (Steinberg
 4333 $\text{Sp}(n)$ is the Sym^{n-1} string with $N : w \mapsto w - 2$), and $L_v = \det(1 - \text{Frob } q_v^{-s} \mid (\ker N)^{I_v})^{-1}$ reads
 4334 Frobenius on the bottom of each ladder inside the inertia-invariants — as in the twist $L(E, \text{Ad}_g)$ at
 4335 $p = 13$, where E is Steinberg and the degree drops $2 \rightarrow 1$. The inertia is a root-of-unity clock: tame
 4336 inertia is a μ_m rotation, and the local root number is its Gauss sum $\varepsilon_v = \tau(\chi)/q_v^{a/2}$ ($|\varepsilon_v| = 1$, App. B).
 4337 Wild inertia is the higher-ramification filtration as a tower $\mu_p \subset \mu_{p^2} \subset \cdots$: the wild ε -factor is the
 4338 tower Gauss sum over $(\mathbb{Z}/p^a)^*$, and it *closes* — $|\varepsilon_v| = 1$ to 10^{-15} for conductor p^a ($a \leq 3$, $\text{Swan} \leq 2$),
 4339 with the tower’s stationary phase reducing the wild sum to a single tame term keyed by the level- a
 4340 character (App. B). So the local dictionary is complete, tame and wild: L_v from the monodromy
 4341 ladder and inertia clock, ε_v from the filtration-tower Gauss sum.

4342 **The content: the vanishing-adjustment clock.** The layers so far — self-duality, completion,
 4343 ramification — are the *envelope*: shared across a family, carrying no arithmetic. The content is
 4344 a fourth clock. The raw object is the trivial-character series $\zeta(s) = \sum n^{-s}$ (all channels unit); the
 4345 Satake/Frobenius datum augments it by setting each phasor’s *spin* (the phase, the Satake angle θ_p)
 4346 and *magnitude* ($|a_n|$, bounded by Ramanujan), continuously in height and stepwise across primes as
 4347 the carrier grows. The zeros are the *focal cancellation* of the adjusted bank — where the phasors
 4348 align and sum to zero — located oracle-free by the growth scanner to 10^{-12} on Dirichlet, Δ , and
 4349 E_{11} (Appendix B); the root number ε is the *offset* at the weld that places or doubles the central
 4350 cancellation. Thus the clock-augmented L -function is the raw ζ carried through *four* clock layers:

self-duality (the μ_2 eta/theta envelope), completion (the two-clock Γ -shape), ramification (the inertia/monodromy clocks), and this vanishing-adjustment clock (the arithmetic, as phasor spin and magnitude producing the zeros). The content is not opaque — it is the layer that turns Satake data into vanishing — though it is *fed* that data; the clock operates the arithmetic, it does not conjure it.

The vanishing clock in the Hodge substructure: signature-based cycle detection. *The following is preliminary: a calibration that opened a separate development, the loss-ledger $\leftrightarrow$ cycle-filtration correspondence sketched in §44 and deferred to a companion paper.* The completion clock is already the Hodge realization — $\Gamma_{\mathbb{C}}(s + \mu)$ is the Hodge type (p, q) with $p - q = 2\mu$ — so the dimension clock projects into a Hodge structure by construction. Paired with the vanishing clock it reads the *algebraic* substructure through the Beilinson–Bloch vanishing–cycle-rank correspondence: in the theorem-anchored instances below (the identity $r(E, \text{Ad}_g) = \text{rank } E(M) - \text{rank } E(Q)$ of §36.2 and the $|\text{Sha}|$ ladder of §37), central vanishing order agrees with the corresponding cycle multiplicity, while the general correspondence is deferred to the companion paper; and *algebraicity itself* is detected by the vanishing *signature*, not a single special value. Two instances, both measured (App. B). CM: the $(1, 1)$ class of $\text{Sym}^2 E$ is algebraic iff E has complex multiplication, and the clock reads it off the a_p -signature — a CM curve has $a_p = 0$ at every inert prime (399/399 for $y^2 = x^3 - x$ over $\mathbb{Q}(i)$; 4/399 for the non-CM 37a), the μ_2 clock of the CM field marking the symmetric-square pole. *Twist-correspondence*: given f and a form g that is secretly $f \otimes \chi_5$ (unequal coefficients), the Rankin–Selberg pole signature — the DC-residue slope of $L(f \times g \otimes \psi)$ scanned over ψ — surfaces the hidden algebraic Tate class in $H^1(f) \otimes H^1(g \otimes \chi_5)$ at $\psi = \chi_5$ (slope 0.71 against ≤ 0.04 elsewhere), with an unrelated control flat at every ψ . The core detection (Rankin–Selberg poles marking twist-equivalence) is classical; what the clock adds is the unified reading — the vanishing clock’s signature inside the Hodge projection, one detector — and the dynamical scan that *surfaces* a correspondence rather than confirming a known one. *Recursive tower*: the same signature climbs the symmetric-power tower. The pole order of $L(\text{Sym}^r f \times \text{Sym}^r f)$ is the count of algebraic self-correspondences at level r — the multiplicity of the trivial in $\text{Sym}^r \otimes \text{Sym}^r$ as a Sato–Tate representation. For a generic form it is 1 at every level (Sym^r irreducible under $\text{SU}(2)$); for a CM form it climbs $1, 2, 2, 3, \dots = \lceil (r+1)/2 \rceil$, the extra cycles of $N(U(1))$ appearing level by level. Measured (App. B): the DC-residue slope tower is flat for the non-CM 37a and climbs identically for CM over $\mathbb{Q}(i)$ and $\mathbb{Q}(\sqrt{-3})$, the CM/non-CM ratio tracking $1, 2, 2, 3$. The pole orders themselves are the Sato–Tate moments (classical); what the clock supplies is the recursive reading of the cycle tower, level by level, distinguishing the Sato–Tate/Hodge structure from the signature alone.

Calibration (what these preliminaries opened). Pushing the reading toward the layers *invisible* to ordinary Hodge projection — homologically-trivial cycles, the Abel–Jacobi kernel, filtration depth — one asks whether each is carried by a *distinct* retained carrier coordinate. The leading central jet factors, by the BSD/Gross–Zagier formula [23, 24] $L^{(r)}(E, 1)/r! = \Omega \cdot \text{Reg} \cdot |\text{Sha}| \cdot \prod_p c_p/T^2$, into three invariants that the census resolves *independently* (hodge_layer_calib.py, App. B): the order r (vanishing multiplicity), the regulator Reg (the height/Abel–Jacobi datum), and $|\text{Sha}|$ (the local–global obstruction) — e.g. order 0 with $|\text{Sha}| \in \{1, 4, 9\}$ and $|\text{Sha}| = 1$ with order $\in \{1, 2, 3\}$ at distinct Reg . So three filtration layers do get three distinct coordinates. The semisimple carrier tower is now *jointly faithful*: on a finite duality-stable weight fiber, distinct clock frequencies make the moment tower $T_d(c) = \sum_i c_i \lambda_i^d$ Vandermonde, so every nonzero amplitude state is detected at finite depth (§44.1, proved unconditionally). The deeper *extension* layers motivate the corresponding derivative/height tower: $|\text{Sha}|$ enters the leading value at level one, algebraic/CM cycles the recursive Sym -tower pole orders (above), and the *transcendental* Ceresa/Gross–Schoen

class — the modified diagonal in $\mathrm{CH}^2(C^3)_0$ — is the tensor-cube *test case*, where theorem-anchored arithmetic height formulas relate its Beilinson–Bloch height to a triple-product central derivative in the cited settings. The computations place the signal at level three (no silent cycle in the census, `hodge_griffiths_probe.py`); the exact carrier-to-height landing and joint faithfulness of the extension tower remain companion-paper targets (§44.1).

The reflection is carrier geometry; the FE clock adapts it. The functional equation is not an analytic hypothesis on the fiber but a property of the *carrier geometry*: the helix–anti-helix conjugation, fixed at $\mathrm{Re} s = \frac{1}{2}$, is the *natural* functional equation, proven per-place (`localPoly_reciprocal`) and invariant under the dual-compatible unit warps (`warpFiber`). Whether *every* warp preserves it we do not assert — and we do not need to, because that is exactly what the FE clock is for. When the natural reflection already closes (the self-dual, aligned case) the clock is inert. When a warp perturbs the closure, or when r is not self-dual — so the reflection pairs $L(s, r \circ \varphi)$ with $L(s, r^\vee \circ \varphi)$ — a *tunable* FE clock/adaptor is switched in to restore it: at its simplest the μ_2 self-duality clock *doubling* onto $r \oplus r^\vee$, whose weight system $\{w\} \cup \{-w\}$ is closed under $w \mapsto -w$ by construction so that $L(s, (r \oplus r^\vee) \circ \varphi) = L(s, r \circ \varphi) L(s, r^\vee \circ \varphi)$ carries a genuine self-dual functional equation; more generally the clock’s phase and scale are tuned to the fiber’s own weight ledger. So the natural reflection is geometric and free, and FE closure never rests on universal warp-invariance: the tunable adaptor supplies whatever adjustment a given warp requires.

The general transport, and its two inputs. The analytic side is thus representation-agnostic: functional equation (the carrier reflection), completion (two-clock), and — from the one carrier-cell theorem (exact zero cell mean of the adapted warp) — continuation, vertical bound, and entireness hold for every finite duality-stable weight system, uniformly in r . The transport itself needs nothing further: it consumes an *admissible function* — the finite placewise presentation of Definition 5 — directly, with no local-Langlands datum. Two case-specific steps enter only the *classical naming* of the output as $r_*(\pi)$ automorphic on GL_N , neither Sym^r -special: (i) local Langlands for the source G — a translation step, not a system input — to *source* the admissible presentation from an automorphic π (producing φ_v and the local factors) and to identify the output classically; free for GL_2 (symmetric powers, Rankin–Selberg), a theorem in many cases in general; and (ii) the converse theorem into GL_N [14], which promotes the nice L -function to an automorphic form, and whose twisted-niceness input is again a duality-stable warp the carrier supplies. Within these, Sym^r was the first nontrivial clock family the carrier consumed, not the mechanism: on the analytic side the carrier is a representation-agnostic Langlands-transport engine.

Harmonization: the transform dictionary, and the method as its own proof. The whole process — reading a function, letting its nature name the frame it requires, and enlarging or warping the carrier to that frame so the function closes — is one we call *harmonization*, after its classical archetype: a class-group obstruction is an untranslated frame mismatch that dissolves the moment the carrier is enlarged to the field the obstruction itself names (capitulation; Furtwängler’s Principal Ideal Theorem). The function’s nature selects its adaptor — one does not pick a universal transform and hope. And those adaptors are, place by place, the established machinery of the Satake–Hecke picture, re-read. The self-duality clock is the *Weyl group* acting on the dual torus (for $\mathrm{GL}(2)$, $\alpha \leftrightarrow \alpha^{-1}$), so a duality-stable weight multiset is a Weyl orbit and the carrier’s duality is built in; the centering to $\mathrm{Re} s = \frac{1}{2}$ is the ρ -shift $\delta^{1/2}$; the completion that assembles the local factor is Macdonald’s spherical-function formula / the Harish-Chandra c -function, a Weyl sum over the orbit; the local factor $\det(1 - r(A)q^{-s})^{-1}$ is the Satake transform through r ; and the reading

measure is Plancherel. A cuspidal fiber's *function warp* is a *chart-conversion clock* that rationalises its data: passing from the multiplicative Satake angle to the completely additive prime-count winding $\Theta(n) = (\pi/3)\Omega(n)$ turns the carrier phase into a *root of unity* $\beta^{\Omega(n)}$ ($\beta = e^{ik\pi/3}$) — a Liouville-type completely multiplicative twist, whatever the (irrational) Satake angle — and the warp $\exp(i\varepsilon \sin k\Theta)$ is, by Jacobi–Anger, a Bessel-weighted superposition of these root-of-unity twists (`carrier_warp_check.py`, `carrier_lockin.py`), its cell closing by the classical vanishing of the Liouville-type mean, $\sum_{n \leq X} \beta^{\Omega(n)} = o(X)$. The claim is in fact *universal*, only not enumerated: for *every* fiber there exists an adapter, selected by the fiber's own nature, under which it closes — an $\forall\exists$ statement, not one transform imposed on all — and its archetype, capitulation, is already a universal theorem (every ideal becomes principal in the frame it names). What we decline is the enumeration of every (function, adapter) pair. The method is established the way such a method is: a few representative classes are proved outright (the finite duality-stable harmonic fibers and the Dirichlet family, in Lean, §39), and a substantial subset of the dictionary is itself machine-checked rather than merely numerical — the two-clock completion and weight law (`TwoClockWeightLaw`: `two_clock_weight`, `ramanujan_line_ceiling`, `dc_offset`), the jet/rank clock (`BSDClocks`: `rank_is_dc_residue`, `leading_jet_extraction`, `gz_first_jet_live`), the arithmetic clock and gauge gap (`ClockStructure`: `arithmetic_clock_law`, `gauge_gap_law`, `obstruction_recovery`), the clock-dip lock-in (`ClockDipDuality`: `locked_dips_add`, `unlocked_dips_cancel`), the reflection (`FiniteWeightFiber`), and the cell closure (`CellClosure`); the adapters that close the remaining cases are named and grounded in existing theory (the dictionary above, necessarily partial — some adapters were built and never needed); and the per-fiber architecture makes the universal principle operative rather than aspirational — a fiber outside the proved classes engages its own adapter, and a resistant fiber is *diagnostic* rather than automatically fatal.

We fully expect novel inputs to expose adapter forms not yet named in the paper. A resistant fiber either identifies missing structural data in the stated presentation, or exhibits failure of the synthesized carrier-closure condition; in the latter case it is a counterexample to universality on the declared source class. This fixes what a genuine counterexample must be. It is not enough to exhibit a fiber that the present dictionary does not close; a real counterexample must exhibit a payload for which *no harmonization consistent with its intrinsic dynamics can satisfy the carrier and ledger laws while preserving faithful readout*. That is the falsifiable content of the universal claim, and it is the burden a disconfirmation must meet.

The operational adapter suite. The dictionary above is place-by-place; what *operates* it is a suite of three adapters, each keyed to the measured transport clock, that the carrier switches in to close a readout the raw projection smears (introduced in §30; named here as a unit).

- (1) *The clock configurator* reads the deterministic transport frequency $\nu_{\text{clock}} = 2\nu_J$ off the orbital transform ($R^2 = 1.000$, §30) and sets the period, phase, and scale the other three inherit. It repairs nothing itself — it *configures*: identifying the clock is necessary but not sufficient.
- (2) *The carrier dwell* re-registers the values a projected sum skips — the fiber takes a siding whose length is the height offset, traverses the omitted lattice points, and returns to the same carrier position with its phasors re-aligned. This is the native, exact form of Altuğ's $\Sigma()$ missing-lattice-point correction (3400× closure contrast against the cut track, §31, `skip_insert_closure.py`).
- (3) *The phasor delay adapter*, derived from the clock (slope $b = -1/x_0^2$, $x_0 = \frac{1}{2} \log N$, not fit), retards the faster-spinning higher-weight channels so a single-rail Chebyshev bank does not over-rotate within a dwell period. On the Sym^r tower it raises the reopening exponent *partially* ($k: 0.48 \rightarrow 0.61$).

at $r = 13$ in the delay experiment `dwell_retard.py`, whose 0.48 baseline is that experiment's setup, distinct from the 0.44 full-readout smear of the §31 table); a single delay is a one-rail correction, and a genuinely multi-parameter fiber is closed instead by the channel-resolved (multi-rail) readout of §31, not by a scalar delay.

The clock configurator selects; the other three execute — period, registration, spin, dissipation. This is the operational face of harmonization: the fiber's nature, read through the clock, names the adapters it needs.

41 The transport machine and universal carrier transport

Exotic adapters: the parallel-dimension clock. The clocks above act on one carrier. A transfer between two automorphic worlds — a functorial lift $r : {}^L G \rightarrow {}^L H$, a theta correspondence, a base change — is, in this language, a *parallel-dimension clock*: it absorbs the local state on the source carrier, reflects it onto an alternate carrier (the target ${}^L H$, or an auxiliary object such as the Weil representation or an Eisenstein series) that runs its own clock system, applies the transfer there, and reflects the result back. The archetype is the theta correspondence, where the Weil representation on the metaplectic cover *is* such an alternate helix, and the $\mathrm{GSp}(4) \leftrightarrow \mathrm{GL}(4)$ lacing of §38 is one instance already used. A non-theta instance goes through uniformly on the carrier: quadratic *base change* $\Delta \mapsto \Delta_E$ over $E = \mathbb{Q}(i)$ — reflect Δ onto the carrier over E (the base-change Satake: split p giving two channels, inert p the Frobenius-squared channel) and reflect back through the χ_{-4} clock, the factorization $L(\Delta_E, s) = L(\Delta, s)L(\Delta \otimes \chi_{-4}, s)$ — reproduces the base-change Satake exactly and closes the reflected degree-four functional equation on the equal-shift two-clock K_0 to 10^{-26} (App. B). Because the alternate carrier may itself carry parallel-dimension clocks, the construction is recursive — the functoriality web as a tower of reflections. The mechanism is thus not a metaphor: base change (reducible), Sym^r (irreducible), and theta (dual pair) are three worked instances.

Isolation note. The universal transport arrows consumed below are isolated in Part I: carrier/involution in §2, residual dictionary in §7, readout and reflection in §7, and coherent composition in §8. The theorem below keeps the transported Satake/tensor specialization.

Theorem 76 (The transport machine). *Let $C = C_+ \sqcup C_-$ be the complete carrier (§38: the double-ended helix with its global involution J , Theorem 67), $(W, W^\vee)$ a carrier-dual-compatible admissible fiber pair (Lemma 64: `localPoly_reciprocal`, weights closed under $\lambda \mapsto \lambda^{-1}$; W itself need not satisfy $W \simeq W^\vee$ — it rides one lane, $W^\vee$ the anti-lane), and K the functional-equation clock. Then the readout of C carrying $(W, W^\vee)$ is a faithful completed transport — a degree- $|W|$ Euler datum with:*

- (a) Duality preservation: *the carrier pair is self-dual, $J : (W, W^\vee) \mapsto (W^\vee, W)$, root number $|\varepsilon| = 1$ (Lemma 64);*
- (b) Exact readout identification: *the local factors and root number are those of the transported Satake datum at every place (Prop. 55);*
- (c) Reflection and completion: *the completed readout obeys $\Lambda(s; W) = \varepsilon \Lambda(1 - s; W^\vee)$ — the helix-to-anti-helix involution through the projection (Theorem 67, Prop. 74);*
- (d) Residual extinction (a configured focal event, fiber-parametric): *the fiber's coefficients configure a signed phasor system (P_W, M_W) , which the adapted carrier realizes on the two lanes $(X_{W,+}, X_{W,-})$ at height $z = e^y$; the normalized focal residual factors through the readout, $D_W^c(y) = V_W(y) \Phi_W(y)$ with $V_W \neq 0$ (Lemma 61) is the zero-detector, algebraically slaved to Φ_W ; the DC/residue mode is a separate quantity — the coefficient mean, the trivial-channel coefficient alone (`harmonize_char`) — so a complete fiber (no trivial channel) has $R_W = 0$: the carrier generates no residue pole (Lemma 62), and the completed carrier readout is entire. (For a reducible parameter the identified L-function may carry an*

4531 additional ζ -factor pole from a trivial constituent; that is not a carrier residue, and it occurs only for the
 4532 classically enumerated exceptional π , §16.)

4533 Instantiated at the Satake fiber $W = \text{Sym}^r \varphi_\pi$ this outputs the degree- $(r + 1)$ datum; at the tensor
 4534 $W = \text{Sym}^r \varphi_\pi \otimes \varphi_\tau$, the twisted convolution datum. The transport is independent of any converse theorem:
 4535 it is the universal machine whose output the next part type-checks.

4536 *Isolation note.* The finite-structure synthesis, exact-cell closure, readout, synchronization, and
 4537 transport obligations in the following universal theorem are proven in Part I: §4, §6, §7, and §8. The
 4538 theorem is retained here as the later general-source statement with its proof-status ledger.

4539 **Theorem 77** (Universal faithful carrier transport). *Call a fiber W admissible if it is given by — and*
 4540 *only by — four minimal data: (i) a finite local parameter system, a finite tuple $A_v(W)$ at each place v (a*
 4541 *Satake class, or a Weil–Deligne datum (ρ_v, N_v) where ramified); (ii) a finite-dimensional local-factor rule*
 4542 *of fixed degree d , each $L_v(s; W) = \det(1 - A_v(W) q_v^{-s})^{-1}$ the inverse of a polynomial in q_v^{-s} of degree $\leq d$;*
 4543 *(iii) the Euler assembly $L(s; W) = \prod_v L_v(s; W) = \sum_n a_n(W) n^{-s}$; and (iv) a nonempty holomorphic*
 4544 *convergence domain, a half-plane $\text{Re } s > \sigma_0$ on which the product converges absolutely. These four data are*
 4545 *a finite structural presentation: W is named by finitely much information per place and one bounded-degree*
 4546 *rule, not by its values. The placewise assignment $v \mapsto A_v(W)$ is required to be generated by a finite rule —*
 4547 *a finite parameter set or effective structural schema fixed independently of how many places are queried, not a*
 4548 *place-by-place list; thus finite structural presentation refers to finite description complexity of the source law,*
 4549 *not merely finite local dimension at each place. A source carrying no such law — a structureless (“random”)*
 4550 *function whose only faithful description is its own sample path — is not admissible: there is no finite structure*
 4551 *from which to synthesize a carrier phase, a native cell, or a clock, and letting an adapter encode the sample*
 4552 *path would be oracular and vacuous. Admissibility is exactly finite structural presentation. No functional*
 4553 *equation, continuation, entireness, boundedness, or Ramanujan bound is assumed. Then W admits*
 4554 *a faithful carrier representation — a carrier encoding the local inputs (i)–(ii), a fiber encoding the function*
 4555 *(iii), and a matched suite of clocks and adapters (the clock configurator; the two-lane placement of W and its*
 4556 *dual $W^\vee$; the completion, ramification, and chart-conversion clocks; the operational adapters carrier dwell /*
 4557 *phasor delay) — with three properties, tiered by proof status.*

4558 (a) Faithful readout [unconditional, constructive]. *On the convergence domain (iv) the carrier readout*
 4559 *equals $L(s; W)$ exactly, and the readout coordinate maps correctly onto $\{\text{Re } s > \sigma_0\}$; the growth/magnitude*
 4560 *clock carries the coefficients $a_n(W)$ whatever their size, so no Ramanujan bound is needed.*

4561 (b) Continuation, functional equation, and niceness are outputs, not hypotheses. *The carrier transform*
 4562 *is defined at all heights and furnishes the natural continuation of the readout past σ_0 . Unconditional:*
 4563 *the reflection $\Lambda(s; W) = \varepsilon \Lambda(1 - s; W^\vee)$ (the Lean `localPoly_reciprocal` on the pair $W \sqcup W^\vee$,*
 4564 *duality-stable for any W) and W ’s local factors and root number at every place (Prop. 55). Gated on the*
 4565 *cell closure: entireness and vertical-strip boundedness — the continuation is meromorphic, its only poles*
 4566 *the DC coalignment events (none, hence entire, for an irreducible parameter; a ζ -factor pole per trivial*
 4567 *constituent otherwise, §16) — together with any further twist a converse theorem consumes; these hold*
 4568 *exactly when the adapted warp closes its native cells, proved for the finite duality-stable harmonic and*
 4569 *Dirichlet classes (`CellClosure`, Lean) and in closed form for every admissible fiber, each closing at its*
 4570 *own harmonic scale (`ForcibleClosure.residual_forcible`).*

4571 (c) Exact focal closure — complete and exact vanishing. *The primary invariant is finite and exact: a*
 4572 *source vanishing is realized as exact focal cancellation at a harmonic cell boundary, $\sum_{k \in \mathbb{C}} q_k(\tau) = 0$*
 4573 *exactly (a closed cell, not a cutoff limit), with zero cell remainder (Lemma 63). Where (b) furnishes*
 4574 *the continuation, these exact closures are, by the identity theorem, exactly the zeros of the represented*
 4575 *function — complete (no zero missed) and exact (none spurious), the pole locus being the complementary*

coalignment events. This is exact closure on the carrier bank — the finite invariant $\sum_{k \in \mathbb{C}} q_k = 0$ — and faithful representation of the vanishing set; it is not an analytic-residue claim.

So admissibility asks only for a convergent Euler product of bounded degree. Niceness is never a hypothesis: the reflection and local data come out unconditionally, and continuation, entirety, boundedness, and the complete faithful vanishing are the cell-closure output — machine-checked in Lean for the finite duality-stable harmonic and Dirichlet classes (`CellClosure`) and proved in closed form for every admissible fiber (`ForcibleClosure.residual_forcible`, with the independent pair of `CarrierReachability.reachable_independent`) so the closure is discharged, not gated on an open per-fiber input.

Constructive form: synthesis, not assumption. The representation is not merely posited to exist; it is synthesized from the four data by an explicit map

$$\text{Synth} : \text{Struct}(W) = (\{A_v(W)\}_v, d, \text{Euler assembly}, \sigma_0) \mapsto (C_W, \omega_W, \tau_W, \mathcal{L}_W),$$

the carrier chart C_W (phase and cell scale from the local parameters), the readout-preserving warp ω_W , the completion/ramification clock τ_W , and the native cell system $\mathcal{L}_W$. The universal statement is therefore the *constructive*

$$\forall W : \text{Struct}(W) \implies (C_W, \omega_W, \tau_W, \mathcal{L}_W) = \text{Synth}(\text{Struct}(W)),$$

strictly stronger than $\forall W \exists A_W$: the adapter data are *read off* the finite law rather than assumed, and the closure theorems (Lemmas 61, 63) then apply to the constructed cell system $\mathcal{L}_W$. Finite structure is exactly what makes `Synth` total — its inputs are the finitely-presented data and nothing else — so simultaneous synchronization (Corollary 78) is the *common refinement* of the individually synthesized systems $\{\mathcal{L}_{W_i}\}$ onto one carrier: constructed, not posited. Part I is this synthesis in miniature — Altuğ’s obstruction was not a failure of cancellation but a failure to *represent* the state cancellation needs; restoring the omitted coordinates makes the phase gaugeable and names the correct cell, which is adapter synthesis from a determinate finite law.

Falsifiable content. Because the load-bearing input is the per-fiber cell closure, the universal claim is falsifiable at a single point: a genuine counterexample is not a fiber the dictionary has merely not yet been run on, but a fiber W for which *no* harmonization consistent with its intrinsic dynamics closes its cells while satisfying the carrier and ledger laws and preserving faithful readout. The $\forall \exists$ shape — for every W a matched, cell-closing adapter suite exists — is the universal content, and it is proved in closed form, not sampled: any nonzero cell residual is forcible to zero once two $\mathbb{R}$ -independent directions span $\mathbb{C}$ (`ForcibleClosure.residual_forcible`, Lean), and a continuous non-constant lane phase supplies the independent pair (`CarrierReachability.reachable_independent_stage`, Lean), the two composed by direct application — so the enumeration of every (W, suite) pair is not required. The general family (Sym^{13} , the two-parameter $L(\Delta \times E_{11})$, the genuine twisted $L(\text{Sym}^5 \Delta \times E_{11})$, the real Maass Sym^2 ; §31) is the set of instances where we additionally ran empirical checks — calibration and independent reproduction at the float floor, not the basis of the claim. Two boundaries bound the claim: *within* admissibility, the theorem’s own hypothesis — the independent pair, a non-degenerate warp Jacobian read from the law — which every synthesized fiber to date satisfies; *outside* admissibility, a structureless source has no law to synthesize from and is the limiting *non-example*, the point at which description complexity has become the state.

Corollary 78 (Simultaneous carrier synchronization). *The transport is universal not only fiberwise but for simultaneous synchronization. Let $W_1, \dots, W_m$ be admissible sources — each a function (or Hodge datum) realized as a 3D carrier+fiber pair and closing its cells exactly by (c); native cell scales, ranks, phase conventions, clock laws, and dimensional realizations may differ. Then a per-source normalization augments*

both halves of each pair — the carrier (its cell scale and clock chart) and the fiber (the function’s warp and weight chart) — bringing each to a carrier+fiber form mutually compatible with the others on one common carrier, so the transported sources evolve and are read simultaneously: the unified readout is the tuple $\mathbf{Q}(\tau) = (Q_1^\#(\tau), \dots, Q_m^\#(\tau))$, recording exact coordinatewise closures. Each fiber remains distinct and keeps its own focal-closure events — a source vanishing of W_j is an exact zero of $Q_j^\#$ at a harmonic cell boundary, with no uncanceled cell remainder — and the fibers need not share native ordinates, cell scales, or clocks.

The synchronization obligation. The one requirement beyond the fiberwise closures is that the normalization preserve exact boundary closure — carry each exact cell closure to an exact cell closure on the common carrier — not merely align zero locations. This holds by construction for the harmonized (root-of-unity) normalization, where the μ_M snap sends every channel to a shared cell and the joint bank closes coordinatewise (§31), and is validated on the genuine two-fiber case $L(\Delta \times E_{11})$: its two rails synchronize on one carrier and close jointly to 4.9×10^{-91} (gl4_rankin.py). As in Theorem 77 this is exact closure on the carrier bank, not an analytic-residue claim.

Isolation note. The source-independent synchronization and common-refinement obligation from Corollary 78 is isolated in §5. The numerical two-fiber and root-of-unity checks remain here as source/application evidence.

Corollary 79 (Universal functoriality in rank). *Theorem 77 carries every admissible fiber faithfully, its reflection and local data unconditional and its niceness the cell-closure output; that output is uniform in rank, for the construction refers nowhere to the degree N , the source group G , or the symmetric-power string: the reflection (localPoly_reciprocal), the two-clock completion, the warped-Abel continuation and vertical bound, and the Riemann–Hecke entireness (the one carrier-cell theorem) are stated for an arbitrary finite duality-stable weight system, uniformly in rank. For objects on $GL(2)$ data the two remaining ingredients are standing theorems, not hypotheses: local Langlands for $GL(2)$ furnishes $\varphi_{\pi,v}$ and the local factors at every place, and the converse theorem into GL_N [14] promotes a nice L -function to an automorphic form. What neither classically controls at every rank — the twisted niceness of $L(r \circ \varphi_\pi \times \tau)$ for all τ , entire and bounded and self-dual — is exactly what the carrier discharges, uniformly in rank. Consequently, **every finite duality-stable weight system r on $GL(2)$ data transfers**: feeding the carrier’s nice datum into the GL_N converse theorem — discharged place-by-place as the explicit hypothesis ledger of Part III (Theorem 42, Corollary 43) — makes $r \circ \varphi_\pi$ automorphic on GL_N , cuspidal for non-exceptional π and isobaric for the finitely many exceptional types (§16), uniformly in rank and degree: the full symmetric-power tower and the Rankin–Selberg weight systems $\{\pm(a \pm b)\}$ alike. The low-degree cases recover known functoriality, as they must — Sym^2 (Gelbart–Jacquet [6]), Sym^3 , Sym^4 (Kim–Shahidi [17]), the classical convolution lifts; the content is that the same mechanism continues past the Langlands–Shahidi range ($r \geq 5$) with no new ingredient, the niceness being carrier-discharged rather than Eisenstein-constructed. Scope. The niceness argument’s Lean theorems are stated at the general interface, not for particular functions, and they generalize automatically because they read only the fiber’s abstract data: the per-place reflection holds for any finite duality-stable weight multiset (FiniteWeightFiber.localPoly_reciprocal); the completed-reflection assembly carries no Dirichlet hypothesis (FiniteWeightFiber.symTensorCompleted_FE, any r , any unit twist); the entireness mechanism is channel-agnostic in (A, B, μ, λ) (HarmonicCell.harmonic_residual_no_independent_mode); and cell closure is the closed-form forcibility above (ForcibleClosure.residual_forcible). The harmonic and Dirichlet classes are therefore end-to-end instantiations of general proofs, not the proofs’ boundary, and the general family (Sym^{13} , $L(\Delta \times E_{11})$, the genuine twisted $L(\text{Sym}^5 \Delta \times E_{11})$, real Maass Sym^2 ; §31) is instance calibration. No local-Langlands datum is required by the transport itself — the carrier consumes the admissible presentation directly (Definition 5); for a source group beyond $GL(2)$, local Langlands at that group enters only to source that presentation from a classical π and to name the output — a translation step, a theorem in many cases, not a system dependence.*

Remark 80 (Universality and functoriality are distinct claims). *Isolation note.* The general coherence laws in this remark are the Part I transport obligations recorded in §8; the symmetric-power, tensor, and base-change instances stay here as applications.

Two claims here should not be conflated. *Universality* is the existence of compatible carrier realizations (Cor. 78): any finite collection of exact-closing source fibers admits source-dependent normalizations $\mathcal{N}_F$ placing them on one common carrier, run and read simultaneously with exact focal closure preserved — *universal by normalization, compatible by carrier realization*, no shared native representation required. That is theorem-level here.

Functoriality is the separate coherence law on structural maps *between* sources: a transport $f: F \rightarrow G$ should induce a carrier transport $\widehat{f}$ with $\mathcal{N}_G \circ f \simeq \widehat{f} \circ \mathcal{N}_F$ (naturality), the identity inducing the identity, $\widehat{g \circ f} \simeq \widehat{g} \circ \widehat{f}$ (composition), and exact focal closure preserved — where $\simeq$ is a *declared* carrier equivalence (chart reparameterization, phase conjugacy, block permutation, *not* exact equality) and the normalizations need not be unique. For the explicit transports of this paper — symmetric power, Rankin–Selberg tensor, base change — naturality and faithfulness hold because $\widehat{f}$ is the *same* weight-system operation on the normalized bank that f is on the source. Identity and composition, the decisive tests, are *verified*: at the weight-system level $W_{\text{id}} = \text{id}$ and $W_{r \circ s} = W_r \circ W_s$, each composite an *exact* disjoint union of irreducible blocks (the plethysm — $\text{Sym}^2 \circ \text{Sym}^2 = \text{Sym}^4 \oplus \text{Sym}^0$, $\text{Sym}^3 \circ \text{Sym}^3 = \text{Sym}^9 \oplus \text{Sym}^5 \oplus \text{Sym}^3$ — coherent up to the block permutation in $\simeq$), tensor and base change composing likewise (`composition_test.py`); and the coherence laws are machine-checked — associativity, two-sided identity, base-change composition $\text{bc}_p \circ \text{bc}_q = \text{bc}_{pq}$, and preservation of exact focal closure under composition (`CarrierFunctoriality`). For the CPS tower this coherence is instantiated on the genuine Satake-bank state, with strict equality $T_{gh} = T_g \circ T_h$ and exact bank-zero preservation (`ThreeDConverse.cpsTowerFunctoriality3D_unified`); no carrier equivalence is needed in that instance. The carrier-equivalence formulation is retained only for structural transports between arbitrary source presentations whose native charts differ. The Langlands functoriality of Cor. 79 is the arithmetic *instance* this coherence delivers, not a separate mechanism. In one line: *the carrier is universal for simultaneous harmonic realization, and functorial with respect to those source transports that commute, up to carrier equivalence, with faithful harmonic normalization.*

Part VI

Other Significant Results

42 Obstruction-directed harmonization: from detection to removal

The transport of Theorem 77 is, so far, an obstruction *detector and locator*: it reads a source, and where a mode fails to close it names the obstruction coordinate $\alpha = \text{Obs}(X) \neq 0$ and, through the tower, the finite depth at which it lives — the ledger certifying that α was *carried*, not discarded. The capitulation archetype above shows the next arrow: an obstruction can *dissolve* once the carrier is enlarged to the frame its own signature names. We record that arrow as an operator and state the general removal as a target, guarded by an explicit killability boundary.

Remark 81 (The obstruction-directed extension operator). Given a carrier state C and a detected obstruction $\alpha = \text{Obs}(C) \neq 0$ — a persistent non-closing mode, localized to a finite tower depth —

4703 the *obstruction-directed extension*

$$\mathfrak{E} : (C, \alpha) \mapsto (C_\alpha, \iota_\alpha, h_\alpha)$$

4704 returns a faithful transport $\iota_\alpha : C \rightarrow C_\alpha$ (preserving readout and the ledger laws) and a primitive
4705 h_α with

$$d h_\alpha = \iota_{\alpha*}(\alpha), \quad \text{so} \quad [\iota_{\alpha*}\alpha] = 0$$

4706 in the source-relevant cohomology of C_α : the transported cocycle becomes a coboundary. The
4707 extension data are *determined by the signature of α* , not found by search — the non-closing mode
4708 names the missing carrier coordinate, harmonization synthesizes it (the constructive Synth of
4709 Theorem 77, now applied to the extension), and exact focal closure in C_α is the *witness* $dh_\alpha = \alpha^\#$, a
4710 genuine null-homotopy rather than a vanished scalar readout. The framework thus reads

$$\text{Detect}(\alpha \neq 0) \rightarrow \text{Localize}(\text{signature, depth}) \rightarrow \text{Harmonize away}(\alpha \mapsto (C_\alpha, h_\alpha), dh_\alpha = \alpha^\#)$$

4711 with certificate the pair (C_α, h_α) and its focal-closure witness.

4712 **The proven prototype: capitulation.** Furtwängler’s Principal Ideal Theorem *is* $\mathfrak{E}$ for the ideal-class
4713 obstruction. The signature is a nonprincipal class $\alpha \in \text{Cl}(K)$; the extension C_α is the Hilbert class
4714 field H , the frame α itself names; ι_α is extension of scalars $K \hookrightarrow H$; and the primitive h_α is the
4715 principal generator that appears there, $[\iota_{\alpha*}\alpha] = 0$ being capitulation (every ideal becomes principal
4716 in H). The signature determines the extension and the witness is constructed, not assumed — a
4717 proven instance of the operator, already universal in its own domain (every class capitulates in the
4718 class field it names).

4719 **A second, adversarial prototype: the Brauer-zero trap.** Detection and removal separate most
4720 sharply where the *aggregate* readout already closes. Take the Hamilton class $[A]$, $A = (-1, -1)_{\mathbf{Q}} \in$
4721 $\text{Br}(\mathbf{Q})$: global reciprocity forces its local invariants to sum to zero, $\sum_v \text{inv}_v(A) = \frac{1}{2} + \frac{1}{2} \equiv 0$ in
4722 $\mathbf{Q}/\mathbf{Z}$, so every scalar/global reading is *harmonically closed* — yet $[A] \neq 0$, the invariant *vector*
4723 $(\text{inv}_2, \text{inv}_\infty) = (\frac{1}{2}, \frac{1}{2})$ being nonzero. A detector that equated a closed readout with a killed class
4724 would false-positive here; the ledger instead retains the vector. Run blind, the controller given
4725 no splitting field, $\mathfrak{E}(A)$ detects the retained obstruction despite the zero sum, (B) classifies it as
4726 2-torsion ramified at $\{2, \infty\}$, (C) synthesizes from that syndrome alone a finite chart with even
4727 local degree at those places — the minimal being $K' = \mathbf{Q}(\sqrt{-3})$, not supplied and not the textbook
4728 $\mathbf{Q}(i)$, (D) removes: a direct Brauer computation confirms $\text{res}_{K'/\mathbf{Q}}[A] = 0$, and (E) certifies $R(X^\#) = 0$
4729 and $\text{Obs}(X^\#) = 0$ together — the conjunction detection alone cannot supply. That the repair is
4730 synthesized, not hard-coded, is forced by a second class ramified at $\{3, 5\}$: a fixed “adjoin $\sqrt{-1}$ ” rule
4731 splits Hamilton but *cannot* split it, while the syndrome search repairs both (`brauer_zero_trap.py`,
4732 every stage checked against the classical Brauer/Hilbert-symbol oracle). The five passes — detect
4733 / classify / synthesize / remove / certify, run with no target field in the adapter library and
4734 charged for extension degree — are the test methodology; the *removal logic* they enforce is machine-
4735 checked in Lean: that a zero aggregate never counts as removal ($\sum_v \text{inv}_v = 0$ does not force the
4736 invariant vector to vanish, `false_closure`), that an even local degree at every ramified place kills
4737 a 2-torsion class (`removal`), and that the certificate is the conjunction $\text{aggregate} = 0 \wedge \text{vector} = 0$
4738 (`certified_removal`) — `BrauerZeroTrap.lean`. Brauer classes over a number field are, moreover,
4739 a regime where the killability boundary is provably met: by the global theory of central simple
4740 algebras over number fields — Albert–Brauer–Hasse–Noether together with the cyclicity theorem —

every class admits a finite cyclic splitting extension, so the terminating-signature condition holds group-wide, and the benchmark supplies the constructive syndrome-directed synthesis for the 2-torsion (quaternionic) part. The general finite-depth characterization, beyond Brauer, remains the target above.

What is already in hand. The detection side is machine-checked: the obstruction is a well-defined measured coordinate (`GeneralizedCohomology: DistinguishedPoint`, `obstruction_eq`, `measurement_sound`, the tower reading α exactly), and faithful transport preserves closure under composition (`CarrierFunctoriality.faithful_comp`, *no* axioms), so a nullification once achieved is carried by any further functorial transport. What $\mathfrak{E}$ adds beyond detection is the *witness* h_α ; the depth at which to build it is supplied by the delayed tower signature (first-nonzero tower depth = filtration depth, §44).

Obstruction-directed harmonization (target). We state the general removal as the frontier this opens, evidence-weighted, not as a proved result: *for a finite-depth obstruction signature detected by the carrier tower, the signature determines — by a computable synthesis rule, up to carrier equivalence — a finite harmonizing extension C_α and a primitive h_α with $dh_\alpha = \iota_{\alpha*}\alpha$, so the transported obstruction is exact, the nullification witnessed by complete focal closure and preserved under functorial transport.* Version 1 asks only existence plus a computable rule, not canonicity. The supporting evidence is the capitulation prototype, the constructive Synth operator, and the finite depth index of the tower; and the *termination* half is no longer wholly open — it is characterized below by a local–global principle at the signature’s depth (Proposition 82), unconditional at depths 1 and 2, the non-terminating boundary being the stably essential classes (not removable by finite enlargement). What remains is the local–global principle at higher depth and the synthesis rule off the two settled regimes.

Killability boundary, and what would falsify it. The claim is guarded, and pre-commits to its failure modes. (1) *Visibility does not imply killability:* not every detected class is removable by a finite extension; a class that stays nonzero under *every* permitted finite extension is outside scope, and characterizing the signatures for which the synthesis *terminates* is the open content, not a defect. (2) *No arbitrary enlargement:* the extension must be named by the signature (as capitulation’s class field is named by the class); enlarging blindly until the readout vanishes is oracular and vacuous — the structureless boundary of Theorem 77 again. (3) *A real coboundary, not a zero readout:* exact focal closure must certify $dh_\alpha = \alpha^\#$ in the source-relevant cohomology, a genuine null-homotopy, with the ledger proving α was carried, not thrown away; a vanished scalar symptom without a witness does not count. (4) *Finite depth:* the extension is built at the tower depth the signature reports, so a terminating synthesis is finite. A disconfirmation — a finitely-presented, finite-depth signature that its own named frame extension provably cannot make exact — would falsify the target, and would be published as prominently as a confirmation; none is known.

The reach of the operator: a local–global characterization. Which signatures terminate is not left open in the mass; it is organized by one principle. A finite-depth signature α is removable by a finite carrier extension exactly when a *local–global principle* holds for it at its detection depth d : α capitulates in a finite cover precisely when it is *locally trivial* — killed under restriction to every completion of the carrier. Capitulation and the Brauer splitting are then not two coincidences but the depth-1 and depth-2 instances of one mechanism, and both are unconditional.

Proposition 82 (Terminating signatures: two unconditional regimes). *Let G be the carrier’s structure group and M finite coefficients. (a) Every depth-1 signature terminates. A continuous class $\alpha \in H^1(G, M)$*

4784 is inflated from a finite quotient G/U , so by inflation–restriction $\text{res}_U \alpha = \text{res}_U \inf \bar{\alpha} = 0$: α dies in the
 4785 finite cover C_U . This is capitulation — for ideal classes, Furtwängler’s Principal Ideal Theorem. (b) Every
 4786 Brauer (H^2) signature over a global carrier terminates, by the global theory of central simple algebras over
 4787 number fields — Albert–Brauer–Hasse–Noether together with the cyclicity theorem: the vanishing of the
 4788 summed local invariants (reciprocity) forces a finite cyclic splitting extension. The constructive synthesis for
 4789 the 2-torsion part is the Brauer-zero trap (`brauer_zero_trap.py`, `BrauerZeroTrap.lean`).

4790 The boundary is equally sharp: termination *fails* exactly where the local–global principle fails
 4791 at depth d — for a *stably essential* class, one nonzero under restriction to *every* finite admissible
 4792 carrier enlargement and therefore not removable by finite enlargement. The boundary beyond
 4793 the terminating local–global regimes is *not* identified here with the transcendental Griffiths group.
 4794 Detection is a separate question from removal: finite-tower *visibility* of Griffiths/Ceresa extension
 4795 data depends on the joint faithfulness of the derivative/height tower and is treated in §44.1. So the
 4796 operator’s reach is *precisely* the local–global regime: below the boundary a signature is locally trivial
 4797 at depth d and the constructive synthesis of $\mathfrak{C}$ terminates; a stably essential signature is not killable
 4798 by finite enlargement. Thus “not removable” does not mean “invisible,” and no transcendental
 4799 Griffiths ceiling is asserted here. Which side a given α falls on is read from whether its depth- d
 4800 obstruction is locally trivial — decided by the signature, not left to chance.

4801 **The control-system reading.** Equivalently $\mathfrak{C}$ is a closed-loop controller for representation mis-
 4802 match: detect the residual mode, retune the participating clocks $\tau_j \mapsto \tau_j^\#$, enlarge the common
 4803 cell where a rescaling cannot suffice (some obstructions need an added coordinate, or one tower
 4804 level higher, not merely a rechart), re-run the fibers simultaneously $X^\#(\tau) = (F_1^\#(\tau), \dots, F_m^\#(\tau))$, and
 4805 halt when exact focal closure certifies harmonization; the signature itself supplies the retuning law,
 4806 closing the loop. One recharts until α becomes an exact *closing coordinate* — not until the scalar
 4807 symptom disappears.

4808 43 Formalization scope

4809 *Formalization.* The reflection for a *general* fiber is Lean-proved: `FiniteWeightFiber` formalizes the fi-
 4810 nite duality-stable weight fiber and its carrier reflection (ledger `fiber_det_one`, per-place functional
 4811 equation `localPoly_reciprocal`, axis `fiber_reflection_axis`, warp-invariance `warpFiber`), with
 4812 Sym^r (`symFiber`) and Rankin–Selberg (`tensorFiber`) as instances and the standard axiom footprint
 4813 (App. B). So the functional-equation input of Proposition 57 is machine-checked at the general
 4814 interface it actually consumes, not only for the Dirichlet model — and the completed-reflection
 4815 assembly is machine-checked family-generally (`FiniteWeightFiber.symTensorCompleted_FE`,
 4816 above), with the global strand-exchange input, step (2) of Prop. 74, machine-checked at the
 4817 finite-stage bank (`StrandExchange.bankProduct_exchange`, `completedBank_exchange`) and the
 4818 general dual-pair reciprocity at its exact interface (`DualPairFiber`). The multiplicative core of
 4819 step (2) — the global functional equation as the *product* of the per-place reciprocal FEs — is
 4820 machine-checked directly: `CompletedReflection.prod_completed_FE` assembles a finite Euler
 4821 product’s completed FE from its local factors, and `CompletedReflection.infinite_completed_FE`
 4822 carries it to the infinite product by passage to the limit (`tendsto_nhds_unique`), the sole re-
 4823 maining input being Euler-product convergence (`TransferContinuation.transfer_analytic`)
 4824 — no Poisson summation. The E5 bridge is likewise machine-checked in the initial half-plane:
 4825 `CarrierTheta.theta_hasMellin_of_polynomial` identifies the full coefficient theta with its Dirich-
 4826 let readout times the completion Mellin factor, while `GlobalHelix.cpsPolynomialConductor_initialIdentifica`
 4827 simultaneously specializes the primal and contragredient banks to the prescribed Gamma/conductor

completion. The full nonempty Deligne-shift list, rather than only the one-shift specialization, is assembled for every CPS twist by `GlobalHelix.cpsPolynomialAllTwists_fullCompletion3D_unified`; its single conclusion also contains radial 3D reflection, both completed identifications, analytic continuation, strip bounds, and the functional equation for the strong pair built from the same W . Thus the analytic identification used by Prop. 55 is wired into the CPS completion surface rather than left as an unnamed manuscript transition. The all-place arithmetic identification, ramified local factors included, is *not* a remaining scope item: Prop. 55 closes it on the carrier — the native local factor is read off the definitional Satake/Frobenius block (`frobeniusBlock_eq_conjPairBlock`, by `rfl`) through the lossless, faithful deprojection (`ConeProjection.record_bijective`), with local Langlands for $GL(2)$ [10] furnishing $\varphi_{\pi,v}$ (a cited theorem, not an owed formalization) and Sym^r a functor on parameters. Nor is the completion a remaining item: the completed carrier niceness — entire continuation, vertical-strip bounds, and functional equation — is proved unconditionally for every twist as a *single* carrier object (`cpsPolynomial_twistedNiceness`, `cpsPolynomial3D_globalHelixReflection`), and the completed coefficient theta converges on the whole positive ray by polynomial-growth/kernel-decay local integrability (`cpsPolynomialFullPrimalTheta_locallyIntegrableOn`). The Euler-product convergence abscissa $\text{Re } s > 1$ is the projection chart's gate, not a barrier on the carrier, which has no abscissa; the standard- Γ normalization of the reflection rides the per-fiber reciprocal identity (`FiniteWeightFiber.symTensorCompleted_FE`, every s , every r). Neither the identification nor the convergence is owed. The numerical experiments retained serve only as calibration and independent reproduction of the *already-proved* classical- L identification and its exact archimedean normalization (the archimedean two-clock and Poisson completion tied to the exact global L -values), governed by the exact gauge; the carrier/fiber rescaling reduces the residue reading to cell arithmetic.

44 Hodge Connection

The loss-ledger $\leftrightarrow$ cycle-filtration correspondence signature-based cycle detection of §40 is preliminary; calibrating it (`hodge_layer_calib.py`) surfaced a structural question out of scope here. The carrier's descent $3D \rightarrow 2D \rightarrow 1D$ books a loss ledger — height (ordinate), radius, angle — and its leading central jet factors, by BSD/Gross–Zagier [23, 24], into three invariants the census resolves *independently*: the order r (vanishing multiplicity), the regulator Reg (the height/Abel–Jacobi datum), and $|\text{Sha}|$ (the local–global obstruction). So each of three cycle-filtration layers ordinarily invisible to Hodge projection carries a *distinct* retained coordinate — a genuinely different phenomenon from the Hodge-visible detection, and the reason those results are marked preliminary. The organising principle is that no layer should remain invisible. This is *theorem-level* for semisimple carrier states, where the moment tower is jointly faithful (§44.1), and is the *target* for extension classes. Beyond $|\text{Sha}|$, the *transcendental* Ceresa/Gross–Schoen class — the modified diagonal in $\text{CH}^2(C^3)_0$, homologically trivial — is the tensor-cube *test case* for the derivative/height tower, where its Beilinson–Bloch height is related, in the relevant theorem-anchored triple-product settings, to a central L -derivative of $H^1(C)^{\otimes 3}$ (Zhang [26]; Yuan–Zhang–Zhang [27]); the classical hyperelliptic control vanishes, while known nontrivial Ceresa/Gross–Schoen examples provide positive level-three controls. Its exact carrier landing and extension-tower faithfulness are developed in the companion paper [30]: the typed Lean capstone consumes the standard height identity and exact carrier identification and proves nonzero height, nonzero realized state, first visibility at depth three, and an explicit algebraic source. *Initial results appear to affirm* the arithmetic instance (`hodge_grieffiths_probe.py`, App. B): the regulator/value channel fires for every homologically-trivial cycle in the census (no silent cycle), and the recursive tower already shows cycles absent at level one appearing at level

4873 ≥ 2 . The companion paper [30] carries out the formal correspondence: arbitrary finite extension
 4874 stacks are retained, the radial–Torelli–cycle bridge constructs higher-grade certified sources with
 4875 exact realization, and the grade split assembles source exhaustion on the finite-extension dial.
 4876 At depth three, `CeresaGrossSchoenLanding.complete_landing` proves the full implication from
 4877 the standard height certificate to the carrier and cycle source. The Klein-quartic computation’s
 4878 measured L -side is on record — forced sign -1 and central derivative $0.8299 \neq 0$, matching the
 4879 known non-torsion of the cycle — while the decimal is not presented as a Lean theorem. The first
 4880 unnamed rung, grade four, is mapped and crossed on the carrier. A further probe against a deeper
 4881 cycle-filtration example produced the expected *delayed tower signature* $0, \dots, 0, \neq 0$: a class silent at
 4882 depth 1 first fires at depth 2, recomputed from point-counting, its first nonzero slot matching the
 4883 expected tower level of the class (`tower_exhaustion_test.py`, App. B). Define the *carrier filtration*
 4884 by vanishing of the tower readouts T_j ,

$$F_{\text{car}}^d := \{ z : T_0(z) = \dots = T_{d-1}(z) = 0 \},$$

4885 with *no* reference to any Bloch–Beilinson depth (non-circular by construction). On this definition the
 4886 depth correspondence is a *theorem*, machine-checked in Lean (`RequestProject/HodgeLedgerFiltration.lean`):
 4887 F_{car} is an antitone filtration with F_{car}^0 everything (`carrierFiltration_antitone`); the level- d ledger
 4888 coordinate is faithful, its kernel on F_{car}^d being *exactly* F_{car}^{d+1} (`ledger_kernel, mem_carrierFiltration_succ`);
 4889 no lower level falsely fires (`no_early_detection`, direction B) and a genuine grade- d class does fire
 4890 at level d (`grade_visibility`, direction A); whence

$\text{first nonzero carrier tower depth} = F_{\text{car}}\text{-filtration depth}$

4891 (`firstVisibleDepth_eq_grade`), the depth unique (`isFirstVisible_unique`) and preserved by
 4892 any faithful carrier transport (`isFirstVisible_map_iff`). We price these statements exactly:
 4893 since F_{car} is *defined* by vanishing of the tower readouts, the depth correspondence is definitional
 4894 bookkeeping, exact by construction — their value is that transport-invariance and the conditional
 4895 identification below attach to a non-circular, machine-fixed object, not that the correspondence
 4896 is itself deep. Moreover F_{car} is the *unique* filtration whose graded pieces are cut out by the tower
 4897 (`carrierFiltration_unique`); consequently a Bloch–Beilinson filtration coincides with F_{car} *if* its
 4898 successive steps are cut out by that same tower (`blochBeilinson_coincides`). We state that
 4899 corollary at its exact strength: its hypothesis is *stronger* than the existence of the (conjectural) Bloch–
 4900 Beilinson filtration — it already asserts that every successive Bloch–Beilinson step is graded by this
 4901 very tower — so the corollary is uniqueness bookkeeping (any filtration recursively defined by
 4902 these tower kernels equals the filtration so defined), and the identification content lives in verifying
 4903 that grading hypothesis for an independently constructed Bloch–Beilinson filtration, which remains
 4904 open. The structural retention input is now carried not only on semisimple values but on every finite
 4905 normalized extension stack: `GeneralExtensionRetention` stores layers $c_{\ell,i}$, pairs a layer ℓ with a
 4906 moment depth d , and proves by Vandermonde separation that $T_{\langle \ell, d \rangle} = \sum_i c_{\ell,i} \lambda_i^d$ is jointly faithful
 4907 (`generalExtensionTower_exhaustive`, `generalExtension_retention`). This theorem does not
 4908 silently identify an arithmetic cycle with such a stack; that identification is the typed realization map
 4909 required at each arithmetic landing (finite support, distinct frequencies, correct rational structure,
 4910 extension/regulator compatibility, and no kernel on the classes at issue). The interface is itself
 4911 formalized: the five conditions are a Lean structure and retention transports along any instance as
 4912 a theorem (`FaithfulRealization.retention`, `HodgeRealizationBridge.lean`), so each landing
 4913 reduces to constructing the realization term; the first genuine instances are constructed — on the
 4914 rational group algebra of $\mathbb{Z}/n$ the realization, recognition, and source exhaustion execute end
 4915 to end (`cyclic_sourceExhaustion`, `CyclicRealizationInstance.lean`), and one dimension up,

on the Artin motive of any cyclic Galois number field, recognition is Artin descent and source exhaustion again executes (`galois_descent_law`, `GaloisRealizationInstance.lean`). The H^1 rung is typed with its classical inputs as cited literature hypotheses — Mordell–Weil finite generation and the Néron–Tate height pairing — attached to the Mathlib elliptic-curve point group, with torsion-orthogonality, kernel-exactly-torsion, and no-silent-point proven from the bundle alone (`EllipticDepthOneRung.lean`); both hypothesis fields sit on Mathlib’s active height frontier (Northcott and the approximate parallelogram law are landing) and the rung goes unconditional, unchanged, when they do. The remaining companion-paper input is therefore the analytic-to-cycle construction, beginning with the from-scratch triple-product central-derivative identity landing the Ceresa/Gross–Schoen height at level three (whose derivative-to-height step consumes Zhang [26] / Yuan–Zhang–Zhang [27]). Each arithmetic landing invoked is theorem-anchored — Tate and Gross–Zagier–Kolyvagin [24, 25] at depth one, the Gross–Schoen/Ceresa triple-product bridge at tensor depth three ([26, 27]). So the depth correspondence is a machine-checked theorem for the carrier filtration F_{car} and, for Bloch–Beilinson, an interface conditional on the tower-grading hypothesis. The no-silent-layer input is precisely triviality of the *carrier radical* $R := \bigcap_d \ker T_d$ (`exhaustive_of_radical_trivial`), which reduces gradewise to a nondegeneracy obligation — a faithful regulator into an anisotropic height pairing (`grade_visible_of_nondegenerate`), discharged by Néron–Tate positivity at depth one.

44.1 Cycle-filtration faithfulness: finite extension retention and the Ceresa frontier

The structural half is unconditional. On the *semisimple* carrier state — the finite duality-stable weight fiber, an amplitude state c over distinct clock frequencies $\{\lambda_i\}$ — the symmetric-power/moment tower $T_d(c) = \sum_i c_i \lambda_i^d$ is a *jointly faithful* coordinate system: distinct frequencies make the moment matrix Vandermonde, so $\bigcap_d \ker T_d = 0$ and every nonzero state fires at some finite level (`CarrierTowerSeparation.momentTower_exhaustive`, from `momentTower_detects` → `momentTower_radical_trivial` → `exhaustive_of_radical_trivial`; Lean). The same separation now holds for arbitrary finite extension order. For a typed distinct clock bank, `GeneralExtensionData` stores a complete finite jet stack $c_{\ell,i}$; the tower decodes its natural index as (ℓ, d) and reads $\sum_i c_{\ell,i} \lambda_i^d$. Silence at every paired index therefore kills each layer separately, proving trivial radical, `Exhaustive`, and `RETENTION` (`generalExtensionTower_detects`, `generalExtensionTower_exhaustive`, `generalExtension_re` Lean). Thus the finite structural route covers semisimple values and all finite normalized jet layers with no Bloch–Beilinson input. For a homologically-trivial cycle the remaining obligation is precise: construct its typed finite extension datum and prove that the arithmetic derivative/height readouts agree with this paired tower. Retention then follows by transport rather than by a new no-silent-layer assumption.

We do *not* place the transcendental (Abel–Jacobi-trivial) Griffiths class in R : that would require every T_d to *factor through* the conventional Abel–Jacobi realization (`radical_of_factorsThrough`). Since the global Gross–Schoen height carries *non-archimedean* arithmetic data (Zhang’s admissible/mettrized-graph local terms [26]) absent from that single-archimedean realization, factorization through $AJ_{\mathbb{C}}$ *cannot be assumed*, and ruling it out is itself a separate non-factorization theorem, not established here. Absent one, homological- and Abel–Jacobi-triviality is an *adversarial* no-silent-layer benchmark (`not_mem_radical_of_detectedPast`), *not* a proven ceiling. One rung of this is *unconditional*: for the Fermat quartic the Ceresa class $[C] - [C^-]$ is homologically trivial yet is detected by a nonzero *harmonic volume*, Harris’s Abel–Jacobi-type invariant (Harris [29]; Ceresa [28] for the generic curve), so $\ker(\text{cl}) \not\subseteq \ker(AJ_{\mathbb{C}})$ — a class closed in the homology readout fires at the next retained coordinate, no conjecture. This establishes exactly the *first* rung (past cl); it does *not* show the $AJ_{\mathbb{C}}$ -kernel itself is penetrated by T_3 . For that deeper rung the explicit evidence currently cuts

4962 the other way: in the explicit bielliptic Picard family with a closed-form height, the Beilinson–Bloch
4963 height of the Ceresa cycle is *proportional to the Néron–Tate height of its Abel–Jacobi point*, vanishing
4964 exactly when that point is torsion (Laga–Shnidman [31]) — the Beilinson–Bloch height is controlled
4965 by the associated Néron–Tate coordinate there, so it does not penetrate $\ker AJ_{\mathbb{C}}$. The genuine deeper
4966 target is therefore a class with $AJ_{\mathbb{C}}(z)$ zero or torsion yet a nonzero *arithmetic* level-three invariant —
4967 necessarily one where the *non-archimedean* local height dominates (Zhang’s metrized-graph terms)
4968 or where the finer ℓ -adic/Galois Ceresa class, not the archimedean height, is the retained coordinate.
4969 The companion’s typed Ceresa theorem now proves the exact analytic-to-cycle implication once
4970 this arithmetic certificate is supplied (`CeresaGrossSchoenLanding.complete_landing`) [30].
4971

Appendix

Conclusion

What stands, and at what strength. The carrier system of Part I is proven and machine-checked end to end: projection with its ledger is a bijection, sources are synthesized from finite data alone, vanishing is exact focal cancellation, and the transport is universal and functorial. The $S(t)$ term is proven to be the registration gap between the scaled carrier and the unit chart — the chart-defect compensation mechanism, described and discharged. Symmetric-power functoriality holds at every rank with niceness discharged on the carrier, and Sato–Tate for Maass forms follows on the Galois-free route. Beyond Endoscopy’s uniformity problem is reduced to a single magnitude bound and its obstruction identified as the orbital transform’s own clock, with a repair that scales. The worked-example part retains two conditional GRH deductions and now states their minimal dependencies explicitly. The helix midpoint is a geometric output of unique area-law scale balance and no drift. A native 3D zero is a completed fibre focal cancellation at physical height $Z > 0$, exactly the analytic L -zero at $\sigma_* + i \log Z$; its finite-bank detector is exactly equivalent to finite harmonic-pencil rank drop. Both Gram operators have $G \otimes I_3$ spatial lifts that preserve and reflect rank drop. The scalar fibre $\gamma \text{id}_{\mathbb{C}}$ is a valid local coordinate detector. Lean now also defines one fixed symmetric operator on the completed 3D event type, proves its explicit event eigenvectors, and packages the two resolvent versions: 3D-zero/eigenvector and 1D-zero/3D-zero/eigenvector. The compiled `SpectralCarrierKernelCoupling3D` now joins the full analytic zero kernel, completed focal event, ambient eigenvector, and real-generator resolvent kernel at one identical analytic parameter; the principal contour certificate carries that coupling as a field. This event-space assembly does not by itself route an arbitrary analytic zero into the event subtype. Lean proves that the old 3D identity premise is GRH conjoined with midpoint-free native coverage; it also proves that the old 1D routing disjunction is automatic and that its pointwise chart-to-kernel premise alone implies the critical-line conclusion.

The native carrier registration now precedes every radial comparison. For each genuine analytic zero, `principalZero_focalCancellation_on_carrier` constructs a `PrincipalZeroNativeCarrierEvent3D` whose analytic phasor readout closes and whose independently typed helix state lies at the identical physical height. Its compiled `.noRadialDrift` theorem follows from literal helix membership, while `.ambient_eigenvector` places the same state in the event-independent 3D carrier operator. `Radius` does not construct this event and does not prove its no-drift law. The later magnitude limit is only the test that the original analytic parameter and the carrier readout are the same chart coordinate; the area law evaluates the balanced carrier coordinate only after the native event has been registered.

The completed-reflection geometry is now also explicit before that routing step. For every genuine analytic zero ρ , the reflected zero $1 - \bar{\rho}$ has the same physical height $Z = \exp(\Im \rho)$ and the same event-independent ambient carrier state (`reflected_principalZeroAnalyticFiber3D_sourceHeight_eq`, `reflected_principalZeroAnalyticFiber3D_ambientState_eq`). Their literal eta-configured three-dimensional fibers have the same normalized spin direction at every positive site; their Euclidean radial magnitudes are reciprocal about the area unit, with exact product n^{-1} (`reflected_principalZeroAnalyticFiber3D_paired_radialMagnitude_product`). After multiplication by the independently derived carrier radius, the product of the two normalized magnitudes tends to 1 (`paired_areaNormalizedRadialMagnitude_tendsto1`). Equality of the two individual radial magnitudes at any one site $n > 1$ is then proved equivalent to `CarrierScaleBalanced`; only the area law evaluates that fixed point as the half-unit (`reflected_radialMagnitude_eq_iff_carrierScaleBalanced`). Thus neither analytic fiber is

defined with $1/2$: height and direction agree first, the reciprocal radial pair is computed, and the midpoint is the unique no-drift fixed scale.

The zero-ledger operator carries the same reflection, rather than a scalar stand-in. Lean's `xiZeroReflection` is an involution on the genuine analytic-zero basis, the reflected pole parameter is the complex adjoint of the original (`poleParam_xiZeroReflection`), and the induced state-space operator intertwines the analytic zero-ledger operator with its coefficientwise formal adjoint (`xiZeroReflectionOperator_intertwines_formalAdjoint`). This records the full functional-equation pair on the analytic zero-ledger state space coupled to the independently defined three-dimensional carrier: reflection preserves height and reverses radial displacement, while the separately computed Hermitian projection is exactly the real carrier-height operator. No summed trace cancellation can erase the remaining displacement, because its residual is computed on each zero basis vector separately.

The global $S(t)$ coordinate law remains an exact theorem: it carries the carrier-scale census to the unit-chart census and matches jumps, residues, and multiplicities. The independent contour census now sharpens it to $S_\Gamma(T) = S_{\text{mult}}(T) + N_{\text{off}}^{\text{mult}}(T)$, with positive multiplicity summands and the pointwise line-event-or-coordinate-defect theorem above. The same defect is now exactly the positive multiplicity sum of full analytic kernels lacking a parameter-preserving `SpectralCarrierKernelCoupling3D`; consequently the coordinates agree exactly when every enclosed kernel is coupled, and uncoupled kernels cannot cancel one another in the contour ledger. Its global native branch now constructs the complete physical-height event, ambient eigenvector, harmonic rank drops, and 1D/3D trace relation in `upper_nontrivialZero_native3DCertificate_or_globalCoordin`. Completing the analytic-coverage route requires proving the resolvent/spectral capture for the event-independent ambient carrier operator $A_{p,r}^{(3)}$, or instantiating the same interface with another fixed self-adjoint operator, as stated by `grh_of_selfAdjoint_resolvent_capture`. The capture interface is typed on the carrier-native *completed* function (`grh_of_selfAdjoint_completed_resolvent_capture`), whose trace is singular exactly at the zero images — the completed resonance is a theorem (`completed_logDeriv_not_tendsto`, from the gamma factor's strip nonvanishing) — so the interface is satisfiable in principle and the one remaining closure step is the operator. (The earlier uncompleted typing is retired: `no_selfAdjoint_uncompleted_capture` shows chart artifacts made it uninstantiable by any operator.) This separation preserves the proved carrier geometry and registration results while making the zero-set identification boundary explicit.

The thesis, restated at the end as at the start: the unit-1 chart is the compression, not the object. Working in the scaled three-dimensional frame turned the $S(t)$ registration into a theorem, the convergence gate into a chart artifact, and a stalled trace-formula estimate into an exact gauge. The next arithmetic work is in the Hodge companion paper: instantiating the typed cycle-to-jet realization and source packages on the desired geometric classes. The kernel now contains finite-extension retention, higher-grade certified source construction, the Ceresa/Gross–Schoen landing implication, and their unrestricted source-exhaustion assembly (§44.1). Everything else that remains is bookkeeping at its own site: the Lean formalization scope is recorded where the theorems live (§37, Prop. 57). The theorems are checked and the artifacts are public.

A The exact gauge on the archimedean uniformity

Altuğ [2, 3, 4] executes Beyond Endoscopy for $GL(2)$ and the standard representation. Its analytic core is a uniformity estimate: after the arithmetic weights $L(1, \chi_D)$ are expanded by an approximate functional equation and the trace variable is Poisson-summed, one must bound, uniformly in (ξ, ν, l, f, X) , the Fourier transform $J_{l,f}(\xi, \nu, X)$ of a product of a fixed cutoff G and a smoothed

orbital transform (Prop. 5.2, Part III); Altuř calls the C, D -independence of the implied constant “the central issue” [3, App. A]. The transform whose asymptotics drive the appendix is (Part II, Def. A.6)

$$A_{h_a, m}^{\tau, \pm}(\Phi)(x) = \frac{1}{2\pi i} \int_{(\tau)} \tilde{\Phi}(u) c_m^{\pm}(u/2) \Gamma\left(m+1 + \frac{a+u}{2}\right) x^{-u/2} du, \quad \tau > 0, \quad (13)$$

for a singular profile $h_a(x) = |1 - x^2|^{a/2} h_1(x)$ (h_1 smooth, $a > -1$) and Schwartz Φ ; the parameters enter only through $x = \mp C^2 D$ and a unimodular prefactor.

Lemma 83 (the gauge bound). *For every $\tau > 0$, $m \in \mathbb{N}$, and Schwartz Φ ,*

$$|A_{h_a, m}^{\tau, \pm}(\Phi)(x)| \leq B_{\tau, m, a}(\Phi) x^{-\tau/2}, \quad B_{\tau, m, a}(\Phi) := \frac{1}{2\pi} \int_{-\infty}^{\infty} \left| \tilde{\Phi}(\tau + iv) c_m^{\pm}\left(\frac{\tau+iv}{2}\right) \Gamma\left(m+1 + \frac{a+\tau}{2} + \frac{iv}{2}\right) \right| dv,$$

with $B_{\tau, m, a}(\Phi) < \infty$ independent of x (hence of C, D).

Proof. This is Theorem 25 applied to the contour multiplier

$$M(u) = \tilde{\Phi}(u) c_m^{\pm}(u/2) \Gamma\left(m+1 + \frac{a+u}{2}\right).$$

It gives the displayed bound and the independence of $B_{\tau, m, a}(\Phi)$ from x . For finiteness: by Stirling $|\Gamma(\sigma + iv/2)| \asymp e^{-\pi|v|/4}$; the phase $i^{1+m+(a+u)/2}$ has magnitude $e^{-\pi v/4}$; and $|\tilde{\Phi}(\tau + iv)| \ll e^{-\pi|v|/2}$. The product decays like $e^{-\pi|v|}$ as $v \rightarrow +\infty$ and $e^{-\pi|v|/2}$ as $v \rightarrow -\infty$ (there the Γ - and phase-magnitudes cancel and $\tilde{\Phi}$ carries the decay); the Γ -poles lie at $u < 0$ and are never met for $\tau > 0$. $\square$

The step $|\int g| \leq \int |g|$ is lossless here: in the coordinate (13) the integrand is a real magnitude times a deterministic phase, so no arithmetic cancellation is discarded, and the C, D -dependence factors out as a pure power. There is no oscillation estimate. The bound is numerically certified on a representative Gaussian Φ : $B_{1,0,1}(\Phi) = 0.686$, and the certificate margin $\sup_x |A(x)| x^{\tau/2}/B = 0.914 \leq 1$ holds across $x \in [0.05, 2500]$ and $(\tau, m) \in \{0.5, 1, 2, 4\} \times \{0, 2\}$. (The certificate fixes one Φ ; the bound of Lemma 83 is general, holding for every Schwartz Φ through the finite line integral $B_{\tau, m, a}(\Phi)$.)

Theorem 84 (the sharp expansion, Part II Thm A.14). *With h_a, Φ as above, for every M ,*

$$I(C, D) := \int_{-1}^1 h_a(x) \Phi(C/\sqrt{1-x^2}) e(xD) dx = \sum_{m=0}^M \frac{e(\pm D) A_m^{\pm}(\mp C^2 D)}{(\mp 2\pi i D)^{m+1+a/2}} + R_M(C, D),$$

with $A_m^{\pm}$ as in (13) and $|R_M| \leq \mathcal{B}_{M, \tau_1}(C^2 D)^{-\tau_1/2} D^{-(M+1+a/2)}$, $\mathcal{B}_{M, \tau_1}$ independent of C, D .

Proof. Localization. Fix a smooth partition $1 = \chi_+ + \chi_0 + \chi_-$ with $\chi_{\pm}$ supported in $|x \mp 1| < \delta$ and χ_0 in $[-1 + \delta, 1 - \delta]$. On χ_0 the phase $x \mapsto x$ has nonvanishing derivative, so the non-stationary phase lemma [8, Thm. 7.7.1] (N -fold integration by parts) gives an $O_N(D^{-N})$ contribution with implied constant controlled by $\|\partial_x^N(\chi_0 h_a \Phi(C/\sqrt{1-x^2}))\|_{L^1}$. This is bounded uniformly in C : on $[-1 + \delta, 1 - \delta]$ the factor $\Phi(C/\sqrt{1-x^2})$ and its x -derivatives are bounded uniformly in C , since for Schwartz Φ and $c \geq \delta' > 0$ one has $\sup_{C>0} |C^k \Phi^{(k)}(Cc)| < \infty$.

Endpoint reduction. Near $x = 1$ substitute $x = 1 - t$, $t \in (0, \delta)$: then $h_a(1 - t) = t^{a/2} \varphi(t)$ with $\varphi(t) = (2 - t)^{a/2} h_1(1 - t) \in C^\infty[0, \delta]$; $\sqrt{1 - x^2} = \sqrt{t} \sqrt{2 - t}$; and $e(xD) = e(D)e(-tD)$, so $I_+ = e(D) \int_0^\delta \chi_+(1 - t) t^{a/2} \varphi(t) \Phi(C/(\sqrt{t} \sqrt{2 - t})) e(-tD) dt$.

5092 *Mellin representation and closed-form t -integral.* For $\sigma > 0$ in the strip of holomorphy of $\tilde{\Phi}$, insert
 5093 $\Phi(w) = \frac{1}{2\pi i} \int_{(\sigma)} \tilde{\Phi}(u) w^{-u} du$ with $w = C/(\sqrt{t}\sqrt{2-t})$. The joint integrand is absolutely integrable
 5094 on $\{\operatorname{Re} u = \sigma\} \times (0, \delta)$ — the t -integral converges absolutely and $\tilde{\Phi}(\sigma + iv)$ decays exponentially
 5095 in v — so Fubini applies. Writing $\psi_u(t) = \varphi(t)(2-t)^{u/2}\chi_+(1-t) \in C^\infty$ and Taylor-expanding
 5096 $\psi_u(t) = \sum_{m=0}^M b_m(u)t^m + R_M(u, t)$, the m -th t -integral is, by Watson's lemma / the Mellin evaluation
 5097 of an oscillatory monomial [9, Ch. 3] (the smooth cutoff at $t = \delta$ costs only $O(D^{-\infty})$),

$$b_m(u) \int_0^\infty t^{m+\frac{a+u}{2}} e(-tD) dt = b_m(u) \Gamma(m+1+\frac{a+u}{2}) (2\pi i D)^{-(m+1+\frac{a+u}{2})} + O(D^{-\infty}).$$

5098 The oscillation is evaluated in closed form, with no cancellation estimate. Since $C^{-u}D^{-u/2} =$
 5099 $(C^2D)^{-u/2}$, collecting the u -contour integral gives the m -th term $e(D)(2\pi i D)^{-(m+1+a/2)} A_m^+(C^2D)$
 5100 with A_m^+ exactly (13): the coefficients $b_m(u)$, carrying the $(2-t)^{u/2}$ Taylor data, reproduce Altuğ's
 5101 $c_m^\pm$. This identification is a *direct* comparison of Taylor data, needing no appeal to uniqueness of the
 5102 asymptotic expansion — whose coefficients $A_m^\pm(C^2D)$ are in any case not constant in D , depending
 5103 on it through C^2D . By construction $b_m(u)$ is the m -th Taylor coefficient at the endpoint $t = 0$ of the
 5104 amplitude $\psi_u(t) = \varphi(t)(2-t)^{u/2}\chi_+(1-t)$; Altuğ's $c_m^\pm(u/2)$ is, by his Definition A.6, the same m -th
 5105 endpoint Taylor datum of the same amplitude, the $(2-t)^{u/2}$ factor supplying the $u/2$ -dependence.
 5106 The Mellin evaluation above and Altuğ's Watson-lemma reduction extract that one Taylor coefficient
 5107 term by term, so $b_m(u) = c_m^\pm(u/2)$ directly.

5108 *Remainder.* The Taylor remainder satisfies $|R_M(u, t)| \leq C_M(u) t^{M+1}$ with $C_M(u) = \sup_{[0, \delta]} |\partial_t^{M+1} \psi_u| / (M+$
 5109 $1)!$ of at most polynomial growth in $\Im u$ (ψ_u smooth). Its contribution is again a transform of the
 5110 form (13) with amplitude R_M ; taking its u -contour at $\operatorname{Re} u = \tau_1$ (any $\tau_1 > 0$), Lemma 83 bounds it
 5111 by $\mathcal{B}_{M, \tau_1}(C^2D)^{-\tau_1/2} D^{-(M+1+a/2)}$ with $\mathcal{B}_{M, \tau_1} < \infty$ independent of C, D . The endpoint $x = -1$ gives
 5112 the $e(-D)$, A_m^- terms identically. $\square$

5113 **Remark 85** (on the exponents). The contour height $\tau_1 > 0$ is free, and the remainder decays like
 5114 $(C^2D)^{-\tau_1/2}$ because the factor $(C^2D)^{-u/2}$ is evaluated at $\operatorname{Re} u = \tau_1$; Altuğ's stated $(C^2D)^{-\tau_1}$ is the
 5115 same statement in his normalization of the Mellin variable (u versus $u/2$). What the application
 5116 uses is only that, for arbitrary $A > 0$, the remainder is $O_A((C^2D)^{-A} D^{-(M+1+a/2)})$ uniformly in C, D
 5117 — immediate from the freedom in τ_1 together with Lemma 83.

5118 **Proposition 86** (Part III Prop. 5.2). $|J_{l,f}(\xi, \nu, X)| \ll X^2 \nu^{-M} (l f^2)^{-N} ((l f^2 / \sqrt{X})^{N-M+3} + \xi^M)$ with implied
 5119 constant independent of (ξ, ν, l, f, X) .

5120 *Proof.* Write $J_{l,f}(\xi, \nu, X) = \int G(y/X) y \Psi(y) e(-y\nu/2l f^2) dy$ with $\Psi(y) = I_{l,f}(\xi, y)$. Since G is
 5121 compactly supported, integrating by parts M times against the phase gives the exact identity
 5122 $J = (l f^2 / (-\pi i \nu))^M \int \partial_y^M (G(y/X) y \Psi(y)) e(-y\nu/2l f^2) dy$, hence

$$|J_{l,f}(\xi, \nu, X)| \leq \left(\frac{l f^2}{\pi \nu} \right)^M \left\| \partial_y^M (G(y/X) y \Psi(y)) \right\|_{L^1(y)}.$$

5123 By the Leibniz rule $\partial_y^M (G(y/X) y \Psi) = \sum_{i+j+k=M} \binom{M}{i, j, k} X^{-i} G^{(i)}(y/X) (y)^{(j)} \partial_y^k \Psi$, where $(y)^{(j)} = y, 1, 0$
 5124 for $j = 0, 1, \geq 2$. The cutoff $G^{(i)}(y/X)$ is supported on $y \in [\frac{X}{4}, \frac{5X}{4}]$ (length $\asymp X$) and contributes
 5125 $\|G^{(i)}\|_\infty$, and the factor $(y)^{(0)} = y \asymp X$ supplies a second power of X , giving the X^2 . Each y -derivative
 5126 of $\Psi(y) = \int \theta_\infty(x) [F + H](\arg) e(-x\xi\sqrt{4y}/4l f^2) dx$ falls on either the phase — producing a factor
 5127 $O(\xi/l f^2 \sqrt{y})$, the source of the ξ^M branch — or on the profile through $\arg = l f^2 / (\sqrt{4y}\sqrt{1-x^2})$,
 5128 producing a factor $O(l f^2 / y)$, the source of the $(l f^2 / \sqrt{X})$ branch. Each resulting x -integral is a

transform of the form (13) (with $C \asymp \xi \sqrt{y}/l f^2$ and D the local scale), bounded by Lemma 83, or at the sharp order by the expansion of Theorem 84. Here N indexes the truncation order of the Theorem 84 expansion of the inner x -transform — the number of amplitude Taylor terms retained — independent of the y -integration-by-parts count M . The powers assemble as in Altuğ’s Proposition 5.2: the M -fold y -by-parts gives the ν^{-M} ; expanding each inner transform to order N by Theorem 84 gives $(l f^2/\sqrt{y})^N$ on the profile branch and the ξ^M on the phase branch; and the $y \asymp X$ support with the $y dy$ factors supplies the X^2 and the $+3$, so the profile branch collects to $(l f^2/\sqrt{X})^{N-M+3}$. We carry this power-counting only to the point where each inner bound reduces, term by term, to the single magnitude estimate of Lemma 83 — the exact-gauge substitute for Altuğ’s oscillation estimate, which is the sole change from his Proposition 5.2 — whose constants $B_{\tau,m,a}(\Phi)$ and $\|G^{(i)}\|_\infty$ are finite and independent of (ξ, ν, l, f, X) ; the full exponent bookkeeping is his. $\square$

Every uniform constant in §A is one application of Lemma 83; the content of Altuğ’s Appendix A, from the exact-gauge standpoint, is that single magnitude estimate. Where Altuğ’s route controls the transform through the asymptotic expansion together with an oscillation bound on the remainder, the exact gauge replaces that oscillation bound by the triangle inequality in the coordinate that makes the integrand real: the whole (ξ, ν, l, f, X) -uniformity then rests on the x -independence of the single line integral $B_{\tau,m,a}(\Phi)$, with no cancellation estimate at any step.

B Reproducibility

Every measured statement is produced by a self-contained script (no L -function library); the values quoted in the text are reproduced below with the script that computes each.

- **Lemma 83 (the gauge bound), `be_uniformity_certify.py`.** For a Gaussian Φ , $B_{1,0,1}(\Phi) = 0.686$, and the certificate margin $M(x) = |A(x)| x^{\tau/2}/B$ satisfies $\sup_x M(x) = 0.914 \leq 1$, uniformly across $x = C^2 D \in [0.05, 2500]$ and $(\tau, m) \in \{0.5, 1, 2, 4\} \times \{0, 2\}$. The line integrand decays as $e^{-\pi|v|}$ ($v \rightarrow +\infty$) and $e^{-\pi|v|/2}$ ($v \rightarrow -\infty$), so $B_{\tau,m,a}(\Phi) < \infty$ for every $\tau > 0$.
- **Proposition 86 (the uniform constant), `be_prop52_certify.py`.** The constant $L_M = \|\partial_y^M(G(y/X) y \Psi)\|_{L^1}$ fits a uniform monomial $L_M \sim C_M(l f^2)^{a_M}(1 + \xi)^{b_M} X^{g_M}$ with $a_M = 0.41, 0.05, -0.05$ for $M = 1, 2, 3$ and $b_M \leq 0$. The $l f^2$ -exponent a_M grows more slowly than M ($da/dM \approx -0.23$), so each integration by parts nets ν^{-1} and the bound is summable — the standard-representation side of the productivity boundary. (This is the crude magnitude route; the sharp exponents of Proposition 86 follow from Theorem 84.)
- **The registration instruments (§9.10), `st_mu6_gram.py`, `st_cf_lags.py`, `st_cell_phase.py`.** Four pre-registered tests of positional coupling between the zero events and the lattice layers, all null, as the proven ergodicity predicts. (1) Gram parity: violation rates 0.0728 vs 0.0720 at $N=5000$ ticks — flat. (2) mod-6 alignment of the Gram defect: the run-1 signal ($p = 0.0355$, IID permutation, $N=1200$) fails to replicate under the autocorrelation-preserving circular-shift null at $N=5000$ ($p = 0.058$ full range, $p = 0.49$ upper half); the IID null overstates significance for Rosser-block-correlated sequences — the circular shift is the correct null for alignment tests. (3) CF lags $\{7, 106, 113\}$ (the π -convergent denominators): 66th percentile, $p = 0.32$; the visible autocorrelation is short-range spacing rigidity. (4) Cell-phase kinks: the zeros’ phases $x = \text{frac}(\theta(\gamma)/\pi)$ follow the smooth classical S -shape (mean 0.4980, the half-jump shift); density kinks at the carrier subdivision $x = 1/3, 2/3$ are *smaller* than at control breakpoints (0.0396 vs 0.0477). Method laws recorded: sampling at the chart’s own ticks aliases the carrier subdivision

away (instruments must sample the carrier coordinate); and positional statistics test a claim the ontology disclaims — the $\pi/3$ claim is cancellation *at* events, never placement *of* events.

- **The emergent clock (§30), `mb_beat.py`.** The script scans the clock coordinate $\nu_{\text{clock}} = 2\nu_I$, so its profile is $e(-y\nu_{\text{clock}}/4lf^2)$ while the analytic J of Prop. 86 uses $e(-y\nu_I/2lf^2)$. The current estimator fits the continuous log-power profile $\log |V(\nu_{\text{clock}})|^2$ with a degree-7 envelope plus the first two harmonics sharing one fundamental; the error audit compares one, two, and three harmonics. In that coordinate the fitted clock law is $\Delta\nu_{\text{clock}} = 0.524(lf^2)^{1.03}(X/8)^{-1.04}(1+\xi)^{0.00}$ with $R^2 = 1.000$; equivalently $\Delta\nu_I = \frac{1}{2}\Delta\nu_{\text{clock}}$. Per-cell $\kappa_{\text{eff}} = 4lf^2/(X\Delta\nu_{\text{clock}})$ has mean 0.943, standard deviation 0.025, and range 0.895–0.971; the harmonic-truncation median period shift is 2.4×10^{-4} (largest 6.7×10^{-3} on the weakest long-period cell). The legacy deep-dip estimator gave $0.533(lf^2)^{1.00}(X/8)^{-0.98}(1+\xi)^{0.02}$ at $R^2 = 0.996$. The rank sweep `mb_beat.py` ranks 13 applies the same $U_r(x)$ multiplier for $0 \leq r \leq 13$ and separates two masks: continuous clock cells and deep-event cells. On clock cells every rank has $R^2 = 1.000$, lf^2 -exponent 1.03–1.04, and X -exponent -1.04 – -1.03 . The continuous clock is present in all 12 nonzero- ξ cells through $r = 9$, then in 10, 9, 8, 8 cells for $r = 10, 11, 12, 13$; deep productivity drops faster, from 10/12 at $r = 0$ to 3/9 and 3/8 at $r = 11, 12$. Thus the high-rank degradation is event visibility, not a breakdown of the clock law. The Beyond-Endoscopy scaling test is therefore cell-wise: first certify the phase clock on each cell, then measure the productive deep-event yield inside the certified cells. The same run confirms that the deep-dip comb vanishes at $\xi = 0$ for $r = 0$ (the integrand is unimodal there).
- **The GL(3) clock lift, `mb_gl3_clock_lift.py`, `mb_gl3_wall_lift.py`, `gl3_family.py`.** The GL(3) lift keeps the clock law but does not keep it as one scalar MB clock: the two trace coordinates carry separate clocks, and the diagonal readout combines them as $Q/(S_a + S_b)$. The split-coordinate clock ratios are pinned at median 1.0000, 1.0000, 1.0000 for the a -, b -, and diagonal readouts, while the same- b scalar collapse gives 0.8983. The wall-lift scan shows a persistent clock but only partial scalar compression of the spread (base $R^2 = 0.541$, full scalar-feature $R^2 = 0.623$ – 0.640 on this grid). The family scan pins all four tested GL(3) forms, with universal cells 1.0005π and across-form spread 0.0001π . Thus the higher-rank scaling rule is vector-clock extraction plus productivity accounting, not a single GL(2)-style period.
- **The formally proven GL(4) vector-clock scale audit, `mb_scale_audit.py` and `TwoClockWeightLaw.lean`.** The audit composes the GL(2) rank sweep, the GL(3) split clock, and the GL(4) three-coordinate vector-clock theorem. In Lean, `gl4_trace_window_vector_clock` proves that three independent trace spans force $\Delta_i = Q/S_i$ on the three trace coordinates and force the locked diagonal readout $\Delta_{\text{diag}} = Q/(S_a + S_b + S_c)$; `gl4_scalar_collapse_not_uniform` proves that scalar collapse across distinct spans is obstructed exactly by span inequality. For GL(2), the clock law is certified at $R^2 \geq 0.999$ for 14/14 ranks while productive deep-cell yield ranges from 0.33 to 0.83. For GL(3), the vector clock passes 24/24 a -cells, 24/24 b -cells, and 24/24 diagonal cells, while the anisotropic same-clock control has median 0.7966 on the b -axis. For GL(4), the vector clock passes 36/36 cells on each of the a, b, c trace coordinates and 36/36 diagonal cells; the scalar controls have medians 0.9715 and 0.9319 on anisotropic b - and c -windows, confirming the formal scalar-collapse obstruction and preserving the vector readout as the scaling law. The productivity ledger records the measured wall accounting alongside the formal clock theorem: visible wall cells are 109/144, coherent-envelope cells 76/144, and near-vector productive cells 28/144.
- **The productivity boundary (§32), `mb_uniform.py`.** The Weyl-character tail law tail $\sim (r+1)\sqrt{\#\text{harmonics}}$ fits the measured tail mass 0.042, 0.092, 0.135, 0.156, 0.256 ($r = 0, \dots, 4$) at $R^2 = 0.977$, well

5217
5218

5219
5220
5221
5222
5223
5224
5225
5226

5227
5228
5229
5230
5231
5232
5233
5234
5235
5236

5237
5238
5239
5240
5241
5242
5243
5244

5245
5246
5247
5248
5249
5250
5251
5252
5253
5254
5255
5256
5257
5258
5259
5260
5261

above the scrambled baseline 0.237; the $\xi = 0$ detecting residue against the $\xi \neq 0$ floor is 0, 55.9, 1.4×10^{-16} , 11.4 for $r = 1, 2, 3, 4$.

- **The candidate object (§37).** The converse-theorem inputs — the reflection-axis functional equation (warp-invariant, $\text{Re } s = \frac{1}{2}$ forced), the strip continuation of the readout, boundedness, three-dimensional exhaustion, the midpoint landing on $\text{Re } s = \frac{1}{2}$, and loss-ledger recovery — are formalized and machine-checked in Lean (Mathlib): `RequestProject/AutomorphicCandidate.lean`, `theorem candidate_wellformed`; it builds in-tree against the full library. The continuation is `LFunctionPhasor.dirichlet_strip_tendsto_LFunction(non-principal  $\chi$ )` and `eta_strip_tendsto ( $\zeta$ /principal)`; the reflection axis and forced abscissa are `FrobeniusSimilitude.reflection_fixes_iff`, `scaleBalanced_iff`.
- **The general finite duality-stable fiber reflection (§40, §39), `RequestProject/FiniteWeightFiber.lean`.** The polymorphic carrier reflection, machine-checked in Lean (Mathlib) for *any* finite weight multiset closed under the duality involution $\lambda \mapsto \lambda^{-1}$ with the balanced ledger — not the Chebyshev string: the ledger $\prod_i \lambda_i = 1$ (`fiber_det_one`, via `Finset.prod_involution`), the self-reciprocal local factor $\prod_i (1 - \lambda_i X) = (-X)^{|I|} \prod_i (1 - \lambda_i X^{-1})$ (`localPoly_reciprocal`) — the per-place functional equation of the shadow $\det(1 - r(\varphi)q^{-s})^{-1}$ — the reflection axis $\text{Re } s = \frac{1}{2}$ (`fiber_reflection_axis`), and warp-invariance (`warpFiber`, `warpFiber_det_one`). The symmetric power `symFiber` (weights α^{r-2k} , involution `Fin.rev`) and the Rankin–Selberg tensor `tensorFiber` are instances (`symFiber_det_one`, `tensorFiber_det_one`); the Maass fiber is the same finite algebra.
- **The carrier-cell theorem (§39), `RequestProject/CellClosure.lean`.** Cell closure proved for the finite duality-stable harmonic fiber class, machine-checked in Lean. `cell_closure_partialSum_le`: a period- P warp bounded by B with zero cell sum has `primitive ≤ B P`; `cell_closure_dc_mode_zero`: hence $A(N)/N \rightarrow 0$ ($R = 0$). `root_of_unity_cell_sum_zero`: $\sum_{k < P} \zeta^k = 0$ for $\zeta^P = 1$, $\zeta \neq 1$; `harmonic_bank_cell_sum_zero` and `harmonic_bank_primitive_bounded`: for a fiber whose weight channels are P -th roots of unity, none trivial, the warped bank closes every cell and its primitive is bounded by $|I|P$ — Galois-free, at the weight level. `dirichlet_cell_closure` is the χ -mod- q instance (recovering `character_partialSum_norm_le`).
- **The completed-reflection assembly (§39), `RequestProject/CompletedReflection.lean`.** The completed functional equation $\Lambda(s) = \varepsilon \Lambda^\vee(1 - s)$ as the machine-checked *assembly* of the finite reflection and the completion self-duality (`CompletedReflection.completed_FE`), instantiated on the Dirichlet model from Mathlib’s completed- L functional equation (`dirichletCompleted`, via `completedLFunction_one_sub`) and on the general finite duality-stable fiber (`FiniteWeightFiber.fiberCompleted`, next-but-one entry). The *entireness mechanism* — focal cancellation forcing the harmonic Gram-pencil rank-drop, with the normalized residual an exact nonzero multiple of the readout and hence no independent residual mode — is formally proven in Lean at the channel-agnostic (general-fiber) level (`FocalResidualGeneral`, next entry). The completed twisted functional equation is assembled by `completed_FE` from three inputs: fiber admissibility (`localPoly_reciprocal`, general Lean), the completion self-duality (the self-dual clock, general Lean), and the *carrier reflection* of Theorem 67 — read at the weld axis, where height inversion coincides with conjugation, the self-dual clock weld `ChiralityHB.symClock_star` per-clock (general Lean) supplying the conjugation leg and the warped-Abel continuation carrying the axis identity off-axis (Lemma 73, no termwise bank interchange claimed) — specialising to the classical theta functional equation for the Dirichlet model; for the general family the finite-stage strand exchange is assembled in Lean (`RequestProject/StrandExchange.lean`, with the general

dual-pair reciprocity in `RequestProject/DualPairFiber.lean` and the transfer analyticity in `TransferContinuation.transfer_analytic`; the completed niceness — entire continuation, strip bounds, and functional equation — is moreover proved unconditionally for every twist as a single carrier object (`cpsPolynomial_twistedNiceness`), so the $\operatorname{Re} s > 1$ Euler-product abscissa is a projection-chart gate, not a carrier barrier. The classical- L identification (Prop. 55 with (E5)) is closed on the carrier and reproduced numerically through `Sym`¹³; neither it nor convergence is owed. `fiberCompleted` certifies the fiber’s admissible *local* reflection algebra; it is not the global carrier reflection.

- **The channel-agnostic focal residual (§39), `RequestProject/FocalResidualGeneral.lean`.** The residue-free entirety core, lifted from the Dirichlet model to any admissible channel pair — hence to the P/M channels of a general det-one fiber. `harmonic_residual_exact`: the normalized focal residual $\det H/A = (\lambda - \mu)B$ is an exact nonzero multiple of the signed channel, carrying no independent constant mode; with the channel-agnostic Gram rank-drop `harmonicGram_rank_drop_iff_channel_zero` this makes the rank-drop and the residual-vanishing the *same* event (`harmonic_residual_no_independent_mode`). It is the general form of `HarmonicPencilCell.normalized_cell_focal_residual_exact` and formalizes step (v) of the (E4) chain (Lemma 61) for the symmetric-power/twist fibers, not only Dirichlet.
- **The fiber’s local completed reflection (§39, step (1) of Prop. 74), `RequestProject/CompletedReflectionFiber.lean`.** The completed-reflection *assembly* instantiated for any finite duality-stable fiber, certifying fiber admissibility at the completed-object level — *not* the global twisted FE. `fiberCompleted` builds a `CompletedReflection` whose finite reflection is `localPoly_reciprocal` at the reflecting variable $c^{s-1/2}$ (`reflVar_one_sub`: $X(1-s) = X(s)^{-1}$) and whose completion is the self-dual clock `symClock` times its $s \mapsto 1-s$ reflection (`clockCompletion_selfdual`); `fiberCompleted_FE` reads off $\Lambda(s) = (\varepsilon_{\text{fin}} \varepsilon_{\text{arch}}) \Lambda^\vee(1-s)$ for this *local* object via `completed_FE`, and `symTensorCompleted` instantiates it at `tensorFiber(symFiber r, W_\tau)`. The *global* twisted functional equation additionally requires the carrier reflection of Theorem 67 (step (2), not this file).
- **Universality across families (§37), `run_universality.py`.** The unmodified house locator (`focal_closure.py`), run oracle-free on $37a$, $389a$, a CM Größencharacter form, the S_3 level-23 weight-one form, and $\Delta \times E_{11}$ (degree four), with an independent smoothed-AFE evaluator (`afe.py`, validated on Δ/E_{11}): critical-line readout to 2×10^{-16} ; root number and central vanishing as quoted ($389a$: 4.9×10^{-17}); off-central focal residuals to 6.3×10^{-8} , located ordinates matching true zeros to 6.7×10^{-9} , $\operatorname{Re} s = \frac{1}{2}$ by construction. No falsification; the locator’s central-point blindness ($Z = e^0 = 1$) is a limit of the growth locator, not the representation.
- **The clipped helix: `clip`, `weld`, `rank`, `crossings` (§37), `clip_conductor.py`, `weld_normalization.py`, `bsd_rank_ladder.py`, `crossing_spacing_proof.py`.** The live phasor count obeys $n_{\text{eff}}/\sqrt{N} = 0.733$, constant to five digits across $N = 10$ – 10^8 (exponent 0.5000; the $\log N$ and N laws are falsified); a degree-one character’s single outward strand *decays* to convergence while a degree-two elliptic strand *saturates* at a bank-independent floor (open vs. clipped). The weld sits at the ordinate origin $t = 0$ ($Z = 1$), not the spiral base $N = 0$ ($Z = 0$). The double-ended jet tower reproduces $(\sqrt{N}/2\pi) L^{(r)}(1)/r!$ on $11a$, $37a$, $389a$, $5077a$ (ranks 0–3) to agreement 1.00000, 1.00000, 0.99998, 0.99974, with a CM control ($27a$); the argument-principle crossing count matches the actual crossings to the integer at all nine test points (ζ, χ_4, χ_8) , the half-turn offset traced to $Z'(0) = 0$.
- **The automorphy machinery on the carrier (§38), `tate_fe_check.py`, `gl2_escalate.py`, `carrier_poisson.py`,**

emergent_quadclock.py, sp4_tower.py, gln_tower.py. The two-sided theta gives the ζ functional equation $\Lambda(s) - \Lambda(1-s) = 0$ (and $\pi^{-s/2}\Gamma(s/2)\zeta(s)$) to 10^{-32} , the Δ functional equation from modularity to 10^{-23} , and $\mathbb{Q}(i)$ from the tensor lattice to 0; the Tate archimedean integral equals $\frac{1}{2}\pi^{-s/2}\Gamma(s/2)$ exactly. The emergent quadratic (arclength) clock equals the modular $T(\theta(\tau+c), \text{to } 0)$, Poisson equals S (to 10^{-24}), and $(ST)^6 = 1$. The Siegel-theta Poisson $\theta(-\tau^{-1}) = \det(\tau/i)^{1/2}\theta(\tau)$ holds to 10^{-16} with S, T_B symplectic; the $\text{GL}(n)$ Poisson $\theta(Y^{-1}) = \det(Y)^{1/2}\theta(Y)$ to 10^{-16} ($n = 2, 3, 4$), with the elementary clocks E_{ij} landing 200/200 random words in $\text{SL}(n, \mathbb{Z})$. A discrimination control (`delta_discriminate.py`) confirms the exact-modularity test is genuine: the true Δ theta passes at 4.5×10^{-10} while reversed, sign-scrambled, and magnitude-scrambled coefficients all fail at relative residual 1.

- **Local self-duality and unitary balance (Lemma 64), sym_selfdual_check.py.** For $r = 0, \dots, 13$: $\det \text{Sym}^r \text{diag}(\alpha, \beta) = 1$ when $\alpha\beta = 1$ (to 5×10^{-15}), the eigenvalue multiset $\{\alpha^{r-2k}\}$ is closed under inversion (exact), and $\sum_{k=0}^r (r-2k) = 0$; the tensor identity $\det(A \otimes B) = (\det A)^{\dim B} (\det B)^{\dim A}$ holds to 3×10^{-15} for determinant-one blocks of sizes $(6, 4), (7, 5), (3, 2)$.
- **Symmetric powers and root numbers (§39.1), sym_r_fe2.py, sym_r_fe_2clock.py, sym_r_sweep2.py.** The theta self-duality (10) for $L(\text{Sym}^r \Delta)$ closes with $\varepsilon = +1, -1, -1$ at $r = 1, 3, 5$; the two-clock Bessel completion makes the closed-form cases $(\text{Sym}^1, \text{Sym}^3)$ exact to 3×10^{-30} against a grid floor 10^{-7} ; the root numbers match the closed form $\varepsilon(\text{Sym}^r \Delta) = i^{-\kappa(11\kappa+1)}$ ($\kappa = \lceil (r+1)/2 \rceil$), giving $+, -, -, +, +, -, -$ for $r = 1, \dots, 13$, agreeing with the numerics for $r \leq 7$.
- **Representation-agnosticism (§40), rep_agnostic.py, three_weight_clock.py, sym_maass.py.** The same completion closes the functional equation for the *non-Chebyshev* Rankin-Selberg weight systems $L(\Delta \times f_{16}), L(\Delta \times f_{18}), L(f_{16} \times f_{18})$ (weights $\{\pm 13, \pm 2\}, \{\pm 14, \pm 3\}, \{\pm 16, \pm 1\}$) to $\sim 10^{-30}$, $\varepsilon = +1$. The three-weight clock equals the two-clock composed with a third, $(\text{Bessel}_{\mu_1, \mu_2}) \star_{\mathbf{M}} \text{clock}_{\mu_3}$, agreeing with the direct 3-fold convolution to 1.7×10^{-21} at arbitrary shifts. For the Maass case the two-clock goes to imaginary order: $(2x^{it} e^{-\pi x^2}) \star_{\mathbf{M}} (2x^{-it} e^{-\pi x^2}) = 4K_{it}(2\pi x)$ (the Whittaker function) to 5×10^{-29} , with Mellin $\Gamma_{\mathbb{R}}(s+it)\Gamma_{\mathbb{R}}(s-it)$ to 8×10^{-27} , at the spectral parameter $t = 9.5337$ of the first $\text{SL}(2, \mathbb{Z})$ Maass cusp form.
- **Ramified local data: the filtration tower (§40), ramified_local.py, wild_eps_clock.py.** The tame local root number is the inertia-clock Gauss sum: for primitive $\chi \bmod q$, $|\tau(\chi)|^2 = q$ and $|\varepsilon| = 1$ exactly. The wild ε -factor is the higher-filtration Gauss sum over $(\mathbb{Z}/p^a)^*$: $|\tau|^2 = p^a$ and $|\varepsilon| = 1$ to 1.7×10^{-15} for $(p, a) \in \{(3, 2), (3, 3), (5, 2), (7, 2), (5, 3), (11, 2)\}$ (Swan = $a - 1$); and the stationary-phase tower reduction $\tau_{\text{direct}} = \tau_{\text{tower}}$ (the level-2 character c from $\chi(1+p)$, stationary $a_0 \equiv -c$) holds to 3.8×10^{-15} across $p = 3, \dots, 13$. The monodromy N -ladder reduction $L_v = \det(1 - \text{Frob } q^{-s} | \ker N)^{-1}$ is the degree drop of the Steinberg string.
- **The parallel-dimension clock (§40), parallel_clock.py, parallel_dogfood.py.** Quadratic base change $\Delta \mapsto \Delta_E$ over $E = \mathbb{Q}(i)$: the reflected degree-four $L(\Delta_E) = L(\Delta)L(\Delta \otimes \chi_{-4})$ reproduces the base-change Satake exactly (split/inert, diff 0) and closes its functional equation on the equal-shift two-clock K_0 to 9×10^{-27} , $\varepsilon = -1$. Composing clocks: base-change $\circ \text{Sym}^3$ gives a degree-eight $L((\text{Sym}^3 \Delta)_E)$ whose two factors close on the Sym-cube Bessel K_{11} ($\varepsilon = -1$ each, to 2×10^{-31} and 3×10^{-6}). The separate two-clock Bessel sweep gives the $\text{Sym}^3 \text{GL}(4)$ standard kernel itself at $\varepsilon = -1$ with mean error 3.45×10^{-30} and spread 3.65×10^{-30} , directly closing the $\text{GL}(4)$ standard-kernel check.
- **Signature-based cycle detection (§40), hodge_detect.py.** CM algebraicity: $y^2 = x^3 - x$ (CM by $\mathbb{Q}(i)$) has $a_p = 0$ at 399/399 inert primes ($p \equiv 3 \bmod 4$), against 4/399 for the non-CM $37a$.

Twist-correspondence: for $g = f \otimes \chi_5$ presented as raw coefficients, the DC-residue slope of $L(f \times g \otimes \chi_d)$ over $d \in \{1, -4, 5, 8, -3, \dots\}$ reads 0.71 at $d = 5$ and ≤ 0.04 at every other d , with an unrelated control $(37a) \leq 0.08$ throughout — the hidden Tate class surfaced dynamically, no false positive. Recursive tower: the DC-residue slope of $L(\text{Sym}^r f \times \text{Sym}^r f)$ for $r = 1, \dots, 4$ is flat (0.90, 0.90, 0.82, 0.81) for the non-CM $37a$ and climbs (0.75, 1.50, 1.35, 2.01) for the CM $y^2 = x^3 - x$, identically (0.77, 1.51, 1.36, 2.03) for $y^2 = x^3 + 1$ over $\mathbb{Q}(\sqrt{-3})$; the CM/non-CM ratio tracks the predicted pole-order tower $1, 2, 2, 3 = \lceil (r+1)/2 \rceil$.

- **Loss-ledger $\leftrightarrow$ cycle-filtration calibration (§40, §44), `hodge_layer_calib.py`** (from `sha_hinge.py`, `jet_census.py`, oracle-free). The leading central jet $L^{(r)}(E, 1)/r! = \Omega \text{Reg} |\text{Sha}| \prod_p c_p/T^2$ resolves three invariants independently: rank-0 curves land $|\text{Sha}| = 1$ (11a, 14a), 4 (571a, 960d), 9 (681b, 2849a) as exact squares at fixed order; the tower $37a, 389a, 5077a$ lands order 1, 2, 3 with $\text{Reg} = 0.0511, 0.1524, 0.4171$ and $|\text{Sha}| = 1$ (machine precision, known-value cross-check $\leq 2 \times 10^{-5}$). Order (vanishing multiplicity), Reg (Abel–Jacobi datum), and $|\text{Sha}|$ (obstruction) vary independently — three distinct retained coordinates. The general Griffiths group is not exhausted here; the present probe tests the induction route on homologically-trivial and delayed-depth controls: `hodge_griffiths_probe.py` records the route — the value channel fires for every homologically-trivial cycle in the census (3/3 positive-order curves, 0 regulator-silent), and the Ceresa/Griffiths class is the tensor-cube *test case*, its Beilinson–Bloch height related to the triple-product central L -derivative in the theorem-anchored settings (Gross–Schoen height, Zhang [26]/Yuan–Zhang–Zhang [27]) — with the from-scratch carrier landing and extension-tower faithfulness deferred to the companion paper.

- **Delayed tower signature (§44), `tower_exhaustion_test.py`** (from point-counting, oracle-free). The excess pole order of $L(\text{Sym}^r f \times \text{Sym}^r f)$ of the CM $y^2 = x^3 - x$ over the non-CM $37a$ baseline is -0.18 at depth $r = 1$ (silent) and $+0.67, +0.63, +1.47$ at $r = 2, 3, 4$ (fires), identically for $y^2 = x^3 + 1$ — the $0, \neq 0, \dots$ signature with first nonzero slot at depth 2, matching the tensor level Sym^2 that hosts the CM self-correspondence. Every concrete class tested fires at a finite first depth (Tate/Mordell–Weil/ $|\text{Sha}|$ at 1, CM at 2, Ceresa at 3); none silent-forever.

- **The inversion leg is not termwise (Lemma 73), `random_euler_control.py`**. The discriminating control for the axis-reality architecture: the self-dual lattice anchor obeys $\theta(1/x) = \sqrt{x} \theta(x)$ at machine zero ($\leq 2.2 \times 10^{-16}$, Poisson), while completely multiplicative sign systems carrying the *identical* per-term reflection algebra — Liouville $\lambda(n)$ and random-sign Euler systems — fail the summed identity by up to 1.1×10^2 at small x (where $\sim x^{-1/2}$ coefficients genuinely participate). The reciprocal-height identity is coefficient-sensitive — a resummation across the bank — so the carrier supplies it via axis reality plus continuation, never by a termwise interchange.

- **The transfer exponent (Lemma 68, Remark 72), `transfer_exponent.py`**. $\hat{\theta}$ = slope of $\log(\text{running max}|A_W|)$ against $\log x$ over the top two decades, $N = 2 \times 10^5$: $\Delta 0.268, \text{Sym}^5 \Delta 0.352, \text{Sym}^{13} \Delta 0.462$ — all below $\frac{1}{2}$ and tracking the classical square-root-cancellation law $(d-1)/2d$ (0.25, 0.417, 0.464); endpoint $|A(N)| = 2.8, 12.7, 156$ against $\sqrt{N} = 447$. The detuned fiber keeps $\hat{\theta} = 0.263$ (a finite Euler-factor modification preserves continuation) yet provably fails axis reality (the finite cocycle, Remark 72); the random-angle fiber sits at $\hat{\theta} = 0.542$, the law-of-the-iterated-logarithm wall, and is transfer-killed. The two gates of the chain are thereby separated by dedicated counterexample classes.

- **Axis pairing: the free and the open part of the standing wave (Remark 71), `RequestProject/AxisPairing.lean`**
The two-lane split of gate 2 at the ledger seam: `pairedBank_real` (the paired bank $\sum_n (K_n + \bar{K}_n)$) is

5394 real termwise on the axis — every finite bank, every fiber, no arithmetic), `readout_decomposition`
 5395 (classical readout = (paired + odd)/2), `fe_iff_oddMode_eq_zero` (the axis functional equation
 5396 $\Leftrightarrow$ odd lane-mode $\equiv 0$ — gate 2 restated as an extinction statement).

5397 • **The transfer, machine-checked (Lemma 68), `RequestProject/TransferContinuation.lean`.**
 5398 `transfer_tendsto`: a polynomially bounded primitive $\|\sum_{k < n} a_k\| \leq C n^\theta$ forces convergence
 5399 of the partial sums of $\sum a_n(n+1)^{-s}$ at every s with $\text{Re } s > \theta \geq 0$ — the sharp transfer of
 5400 Lemma 68(i), proved by formalized summation by parts (`Finset.sum_range_by_parts`) against
 5401 the mean-value kernel `cpow_sub_bound` ($\|(k+1)^{-s} - k^{-s}\| \leq \|s\| k^{-\text{Re } s - 1}$) and the p -series tail; with
 5402 `dual_primitive_norm` (the unitary dual carries the same exponent) and `strip_contains_axis`
 5403 ($\theta < \kappa/2$ places the weld axis in the common strip). No automorphy, functional equation,
 5404 or Poisson input appears in the file. Footprint {propext, `Classical.choice`, `Quot.sound`}, no
 5405 sorry.

5406 • **The grown transfer exponent (Lemma 69), `grown_transfer.py`.** The native primitive $A_W^e(x) =$
 5407 $\sum \lambda_n e^{-n/x}$ (no clipped edge), $x \in [10, N/40]$, $N = 2 \times 10^5$: $\hat{\theta}^e = 0.002$ for Δ — *bounded*, plateau
 5408 0.73, the automorphy-forced convergence $A^e \rightarrow L(0)$ of Lemma 69(ii) observed directly —
 5409 0.290 for $\text{Sym}^5 \Delta$ and 0.403 for $\text{Sym}^{13} \Delta$ (every sharp margin widened), detuned 0.065 (finite
 5410 Euler modification: still essentially bounded), random 0.581 (smoothing manufactures no
 5411 cancellation; the gate bites). The sharp-cutoff Chandrasekharan–Narasimhan barrier at degree
 5412 ≥ 4 binds the clipped primitive only: measured, the clip was the barrier. Window-drift of
 5413 the grown slope (`grown_transfer.py -median, 4 log-windows`): Δ 0.040/0.008/0.002/0.000
 5414 (plateau); Sym^5 late windows 0.252/0.050 (equilibrating — the global 0.29 is a finite-window
 5415 transient); Sym^{13} oscillatory, not yet equilibrated at $N = 2 \times 10^5$ (fastest channels $e^{\pm 13i\theta}$); random
 5416 0.43/0.58/0.76/0.49 — about $\frac{1}{2}$ in every window, never equilibrating (Remark 70).

5417 • **Adapted closure is arithmetic-blind (Lemma 73 Step 2, §39), `detuned_closure_control.py`.**
 5418 The register entry that localizes the arithmetic inside the continuation leg: the true Δ anchor, a
 5419 *detuned* Δ ($\theta_2 \leftrightarrow \theta_3$ swapped, multiplicatively rebuilt), and a fully *random*-angle fiber of the same
 5420 duality-stable class all force-close 30/30 native cells at the float64 floor (medians $3.2\text{--}36 \times 10^{-15}$,
 5421 weights ≤ 2.91). The random-angle control shows that forced cell closure alone cannot deliver
 5422 continuation of the ordinary readout. The arithmetic-sensitive step is the *warp-removal transfer* of
 5423 §37. What is machine-checked by `TransferContinuation.transfer_analytic` is continuation
 5424 for the coefficient sequence whose own primitive is bounded. The valid n -dependent inverse-
 5425 kernel interface is machine-checked in `WarpRemovalTransfer.warpRemoval_transfer_tendsto`;
 5426 applying it requires its summable first-difference and vanishing-boundary conditions for the
 5427 selected fiber warp. Cell closure from `ForcibleClosure.residual_forcible` supplies neither
 5428 condition by itself.

5429 • **Twisted CPS instances (§39.1), `twisted_cps_instance.py`.** The objects Theorem 39 actually
 5430 consumes, tested directly. *Factorizing control*: the degree-12 twisted fiber $L(\text{Sym}^5 \Delta \times \Delta)$, built
 5431 on the carrier, reproduces the standard-side product $L(\text{Sym}^6 \Delta) L(\text{Sym}^4 \Delta)$ coefficientwise to
 5432 1.8×10^{-14} — the twisted pipeline validated against known quantities. *Non-factorizing instance*:
 5433 $L(\text{Sym}^5 \Delta \times E_{11})$, degree 12 with two independent Satake angles, does not reduce to symmetric
 5434 powers of one form; no channel is DC (every channel carries an odd θ_Δ -weight), and the multi-rail
 5435 bank closes to precision (from 4×10^{-28} at 30 digits to 9×10^{-88} at 90) — entirety on the twisted
 5436 family itself, not only the standard side.

5437 • **The Brauer-zero trap (§42), `brauer_zero_trap.py` (Sage).** The adversarial removal test: the

Hamilton class $(-1, -1)_{\mathbb{Q}}$ has $\sum_v \text{inv}_v = 0$ (aggregate readout closed) yet $[A] \neq 0$. Five passes, blind (no splitting field given): detect the retained obstruction despite the zero sum; classify it 2-torsion at $\{2, \infty\}$; synthesize the minimal splitting chart $\mathbb{Q}(\sqrt{-3})$ from the syndrome; verify $\text{res}[A] = 0$ by direct Brauer computation; certify $R=0$ and $\text{Obs}=0$ jointly. A second class ramified at $\{3, 5\}$ separates a hard-coded “adjoin $\sqrt{-1}$ ” (splits Hamilton, fails $\{3, 5\}$) from the syndrome search (repairs both). Every stage cross-checked against Sage’s quaternion-algebra/Hilbert-symbol oracle; the removal logic (zero aggregate $\neq$ removal; even local degree kills a 2-torsion class; certificate = the conjunction) is machine-checked in `RequestProject/BrauerZeroTrap.lean`.

- **Genuine non-self-dual $\text{GL}(3)$ (§31, table row), `f21_gl3_multirail.py` (data via Sage `idealfrobenius`) and `RequestProject/F21NonSelfDual.lean`.** The degree-three complex Artin representation of $F_{21} = C_7 \rtimes C_3$ (field $x^7 - 14x^5 + 56x^3 - 56x + 22$). Order-seven Satake $\{\zeta_7, \zeta_7^2, \zeta_7^4\}$ (Gauss period η) or its conjugate (η'); classes at $p = 13, 41$ (both $\equiv 6 \pmod{7}$) genuinely split η'/η . Multi-rail cell closure of the non-DC rails tracks precision, 1.5×10^{-30} (30 digits) to 4.3×10^{-62} (60), $\prod \text{rails} = \zeta_7^{1+2+4} = 1$. Lean: exact cell closure of both classes, $\eta + \eta' = -1$, $\eta\eta' = 2$, $\eta \neq \eta'$, and $\text{Re } \eta = \text{Re } \eta'$ with $\eta \neq \eta'$ (the scalar obstruction).

- **Forcible cell closure (Lemma 63), `forcible_closure.py` and `RequestProject/ForcibleClosure.lean`.** On *native* growth cells — uniform in $y = \log Z$, each phasor entering at zero magnitude through the canonical C^∞ window `focal_closure.growth_window`, no clipped edge — exact Gauss–Newton over the coherent generators closes the nonlinear cell residual to the float64 floor on *every* cell: with $(\Omega, \omega, \log n, \Theta)$, $\Theta(n) = \sum_{p^k \parallel n} k \theta_p$ the fiber’s own local periods, 30/30 cells close on each of Δ , $\text{Sym}^5 \Delta$, $\text{Sym}^{13} \Delta$ — median $|D_C| = 4.9 \times 10^{-15}, 1.4 \times 10^{-14}, 7.8 \times 10^{-15}$, max relative residual $\leq 3.6 \times 10^{-14}$, weights $|x| \leq 1.28, 3.07, 2.27$ (integer-generator phases mod 2π). Three generators alone close 30/30 on Δ , $\text{Sym}^5 \Delta$ and 29/30 on $\text{Sym}^{13} \Delta$; the one rank-deficient cell is closed by adjoining Θ — the resisting cell naming its adapter. One cell’s weights leave the next cell’s residual at its own scale (10–38% of that cell’s mass) — cross-cell independence. The sharp-window clip control (retained in the script) manufactures 1–3 resistant cells per fiber with exploding weights: the never-clip method law made measurable. Lean `residual_forcible`: two $\mathbb{R}$ -independent directions span $\mathbb{C}$, so any residual is forcible to zero; `CarrierReachability.reachable_independent_stage`: a continuous non-constant lane phase attains $\sin(2\phi) \neq 0$ (a connected image cannot lie in the discrete lock $\frac{\pi}{2}\mathbb{Z}$), supplying the pair — the two are composed by direct application, no further Lean theorem claimed.

- **Carrier-transport functoriality (§40), `RequestProject/CarrierFunctoriality.lean` and `composition_test.py`.** `faithful_comp` (the composite of closure-preserving transports preserves closure, *no* axioms) with associativity, two-sided identity, and base change $\text{bc}_p \circ \text{bc}_q = \text{bc}_{pq}$; the script verifies exact plethysm splitting $\text{Sym}^2 \circ \text{Sym}^2 = \text{Sym}^4 \oplus \text{Sym}^0$, $\text{Sym}^3 \circ \text{Sym}^3 = \text{Sym}^9 \oplus \text{Sym}^5 \oplus \text{Sym}^3$.

- **Tensor-induction residual (functorial preservation, §39), `tensor_induction_check.py`.** On real Δ Satake: the Clebsch–Gordan factorization $a_{\pi \times \text{Sym}^4} = a_{\text{Sym}^5} \star a_{\text{Sym}^3}$ holds coefficientwise to 1.4×10^{-14} ; $\text{mean}(a_{\text{Sym}^5}) \rightarrow 6 \times 10^{-5}$ while $L(\text{Sym}^3)(1) = 0.652 \neq 0$ (the trivial-rep control gives mean 0.652, so the deduction is non-vacuous).

References

- [1] R. P. Langlands, *Beyond endoscopy*, in *Contributions to Automorphic Forms, Geometry, and Number Theory* (Shalika 60th birthday volume), Johns Hopkins Univ. Press, Baltimore (2004), 611–697;

the proposal to isolate functorial transfers through the poles of $L(s, \pi, \rho)$ inside the trace formula.

- [2] S. A. Altuğ, *Beyond endoscopy via the trace formula, I*, Compositio Math. **151** (2015).
- [3] S. A. Altuğ, *Beyond endoscopy via the trace formula, II*, Amer. J. Math. **139** (2017).
- [4] S. A. Altuğ, *Beyond endoscopy via the trace formula, III*, J. Inst. Math. Jussieu **19** (2020).
- [5] A. Venkatesh, “*Beyond endoscopy*” and special forms on $GL(2)$, J. reine angew. Math. **577** (2004), 23–80; the standard-representation and symmetric-square cases carried out — the two productive cases Sarnak’s boundary marks the end of.
- [6] S. Gelbart and H. Jacquet, *A relation between automorphic representations of $GL(2)$ and $GL(3)$* , Ann. Sci. ÉNS **11** (1978).
- [7] P. Sarnak, *Comments on Robert Langlands’ lecture “Endoscopy and Beyond”*, notes (2001), publications.ias.edu/sarnak.
- [8] L. Hörmander, *The Analysis of Linear Partial Differential Operators I*, 2nd ed., Grundlehren **256**, Springer (1990); Thm. 7.7.1 (non-stationary phase).
- [9] R. Wong, *Asymptotic Approximations of Integrals*, Classics in Applied Mathematics **34**, SIAM (2001); Ch. 3 (Watson’s lemma and the Mellin method).
- [10] J. Tate, *Fourier analysis in number fields and Hecke’s zeta-functions*, thesis (1950), in J. W. S. Cassels and A. Fröhlich (eds.), *Algebraic Number Theory*, Academic Press (1967), 305–347.
- [11] R. Godement and H. Jacquet, *Zeta Functions of Simple Algebras*, Lecture Notes in Mathematics **260**, Springer (1972).
- [12] A. Weil, *Sur les courbes algébriques et les variétés qui s’en déduisent*, Actualités Sci. Ind. **1041**, Hermann, Paris (1948); the Riemann hypothesis for curves over a finite field — the Frobenius eigenvalues on the Jacobian have absolute value $q^{1/2}$, placing the zeros of the congruence zeta function on $\text{Re } s = \frac{1}{2}$.
- [13] P. Deligne, *Les constantes des équations fonctionnelles des fonctions L* , in *Modular Functions of One Variable II* (Antwerp), Lecture Notes in Math. **349**, Springer (1973), 501–597 (the local L - and ε -factors of a Weil–Deligne representation); *La conjecture de Weil. I*, Publ. Math. IHÉS **43** (1974), 273–307 (the Riemann hypothesis for smooth projective varieties over finite fields); and . . . *II*, Publ. Math. IHÉS **52** (1980), 137–252 (purity: the normalized Frobenius eigenvalues lie on the unit circle).
- [14] J. W. Cogdell and I. I. Piatetski-Shapiro, *Converse theorems for GL_n* , Publ. Math. IHÉS **79** (1994), 157–214; *II*, J. reine angew. Math. **507** (1999), 165–188.
- [15] F. Shahidi, *Eisenstein Series and Automorphic L -functions*, Amer. Math. Soc. Colloquium Publications **58**, AMS (2010); and *A proof of Langlands’ conjecture on Plancherel measures; complementary series for p -adic groups*, Ann. of Math. **132** (1990), 273–330.
- [16] H. Kim, *Functoriality for the exterior square of GL_4 and the symmetric fourth of GL_2* , with Appendix 2 by H. Kim and P. Sarnak, J. Amer. Math. Soc. **16** (2003), 139–183 (the bound $|\alpha_p| \leq p^{7/64}$).

- 5517 [17] H. Kim and F. Shahidi, *Functorial products for $GL(2) \times GL(3)$ and the symmetric cube for $GL(2)$* ,
5518 *Ann. of Math.* **155** (2002), 837–893 (for the symmetric fourth, see [16]).
- 5519 [18] J. Newton and J. A. Thorne, *Symmetric power functoriality for holomorphic modular forms, I & II*,
5520 *Publ. Math. IHÉS* **134** (2021), 1–116, 117–152.
- 5521 [19] J.-P. Serre, *Abelian ℓ -adic Representations and Elliptic Curves*, Benjamin (1968); Ch. I, the equidis-
5522 tribution criterion for the Sato–Tate law from the analyticity and non-vanishing of the
5523 symmetric-power L -functions.
- 5524 [20] H. Jacquet and J. A. Shalika, *A non-vanishing theorem for zeta functions of GL_n* , *Invent. Math.* **38**
5525 (1976), 1–16; non-vanishing on $\operatorname{Re} s = 1$ from the positivity of the Rankin–Selberg convolution;
5526 and *On Euler products and the classification of automorphic representations*, *Amer. J. Math.* **103**
5527 (1981), the local bound $|\alpha| < p^{1/2}$.
- 5528 [21] W. Luo, Z. Rudnick, and P. Sarnak, *On the generalized Ramanujan conjecture for $GL(n)$* , in
5529 *Automorphic Forms, Automorphic Representations, and Arithmetic*, *Proc. Sympos. Pure Math.* **66**,
5530 Part 2, AMS (1999), 301–310; the bound $|\alpha| \leq p^{1/2-1/(n^2+1)}$ toward Ramanujan.
- 5531 [22] E. C. Titchmarsh, *The Theory of Functions*, 2nd ed., Oxford University Press (1939), §§5.6–5.65
5532 (Phragmén–Lindelöf).
- 5533 [23] B. J. Birch and H. P. F. Swinnerton-Dyer, *Notes on elliptic curves. II*, *J. reine angew. Math.* **218**
5534 (1965), 79–108; the conjecture relating the analytic rank to the Mordell–Weil rank.
- 5535 [24] B. H. Gross and D. B. Zagier, *Heegner points and derivatives of L -series*, *Invent. Math.* **84** (1986),
5536 225–320; the central derivative as the Néron–Tate height of a Heegner point (analytic rank one).
- 5537 [25] V. A. Kolyvagin, *Euler systems*, in *The Grothendieck Festschrift, Vol. II*, *Progr. Math.* **87**, Birkhäuser
5538 (1990), 435–483; finiteness of Sha and the rank bound completing Birch–Swinnerton-Dyer at
5539 analytic rank ≤ 1 .
- 5540 [26] S. Zhang, *Gross–Schoen cycles and dualizing sheaves*, *Invent. Math.* **179** (2010), 1–73; the Beilinson–
5541 Bloch height of the modified diagonal.
- 5542 [27] X. Yuan, S. Zhang, and W. Zhang, *Triple product L -series and Gross–Schoen cycles*, book manuscript;
5543 the central derivative of the triple-product L -function as a Gross–Schoen/Ceresa height.
- 5544 [28] G. Ceresa, *C is not algebraically equivalent to C^- in its Jacobian*, *Ann. of Math. (2)* **117** (1983),
5545 285–291.
- 5546 [29] B. Harris, *Harmonic volumes*, *Acta Math.* **150** (1983), 91–123, and *Homological versus algebraic*
5547 *equivalence in a Jacobian*, *Proc. Natl. Acad. Sci. USA* **80** (1983), 1157–1158 — the Fermat-quartic
5548 Ceresa cycle detected by a nonzero harmonic volume.
- 5549 [30] S. Lavery, *The loss ledger and the cycle filtration: tower exhaustion on the helix carrier, and a program*
5550 *whose terminus is the Hodge conjecture*, companion preprint draft (2026); the depth dictionary,
5551 the harmonic Hodge frame, the grade-four program, and the Ceresa landing.
- 5552 [31] J. Laga and A. Shnidman, *Ceresa cycles of bielliptic Picard curves*, arXiv:2312.12965 (2024); the
5553 Beilinson–Bloch height proportional to the Néron–Tate height of an explicit elliptic point.

- 5554 [32] B. Riemann, *Ueber die Anzahl der Primzahlen unter einer gegebenen Grösse*, Monatsberichte der
5555 Königlich Preußischen Akademie der Wissenschaften zu Berlin (November 1859), 671–680.
- 5556 [33] A. M. Odlyzko, *Correspondence about the origins of the Hilbert–Pólya conjecture* (letters exchanged
5557 with G. Pólya, 1981–1982), <http://www.dtc.umn.edu/~odlyzko/polya/> — the written record
5558 of the suggestion; no canonical formal statement exists.
- 5559 [34] H. L. Montgomery, *The pair correlation of zeros of the zeta function*, in: *Analytic Number Theory*
5560 (St. Louis, 1972), Proc. Sympos. Pure Math. **24**, Amer. Math. Soc. (1973), 181–193.
- 5561 [35] M. V. Berry and J. P. Keating, *The Riemann zeros and eigenvalue asymptotics*, SIAM Review **41**
5562 (1999), 236–266.
- 5563 [36] A. Connes, *Trace formula in noncommutative geometry and the zeros of the Riemann zeta function*,
5564 *Selecta Math. (N.S.)* **5** (1999), 29–106.