

Topology

Samuel Lindskog

September 5, 2024

Open and closed sets

Definition 1.1 (Metric). A *metric* on a set X is a real-valued function d on $X \times X$ that has the following properties:

- (a) For all $x, y \in X$, $d(x, y) \geq 0$.
- (b) $d(x, y) = 0$ iff $x = y$.
- (c) For all $x, y \in X$, $d(x, y) = d(y, x)$.
- (d) For all $x, y, z \in X$, $d(x, z) \leq d(x, y) + d(y, z)$.

Definition 1.2 (Metric space). A metric space (X, d) is a set X equipped with a metric d on X .

Definition 1.3 (Subspace). If (X, d) is a metric space and Y is a subset of X , then the restriction d' of d to $Y \times Y$ is a metric on Y , and (Y, d') is called a subspace of (X, d) .

Remark. Any set X can be made into a discrete metric space by associating with X the metric d defined by

$$d(x, y) = \begin{cases} 1, & x \neq y \\ 0, & x = y \end{cases}$$

Definition 1.4 (Open ball). The open ball $B(x, r)$ with center $x \in X$ and radius $r > 0$ is defined by

$$B(x, r) = \{y \in X \mid d(x, y) < r\}.$$

Definition 1.5 (Interior point). Let Y be a subset of X . A point $x \in X$ is an interior point of Y if there exists $r > 0$ such that $B(x, r) \subseteq Y$. The set of interior points of Y is the interior of Y , and it is denoted by $\text{int}(Y)$.¹

$$^1 \text{int}(Y) \subseteq Y.$$

Definition 1.6 (Open subset). A subset Y of X is open if $\text{int}(Y) = Y$.

Theorem 1.1. Any open ball $B(x, r)$ in a metric space X is an open subset of X .

Proof: Suppose $y \in B(x, r)$. Then $d(x, y) < r$, and $0 < r - d(x, y)$. Suppose $z \in B(y, r - d(x, y))$. It follows from the definition of a metric that $d(x, z) \leq d(x, y) + d(y, z)$, so $d(x, z) \leq d(x, y) + (r - d(x, y)) = r$, so $z \in B(x, r)$. \square

Theorem 1.2. The union of a family of open subsets of a metric space X is an open subset of X .

Proof: Suppose $\{U_\alpha\}_{\alpha \in A}$ a family of open subsets of X . If $x \in \bigcup_{\alpha \in A} U_\alpha$, then $\exists \alpha (x \in U_\alpha)$, so there exists an open ball $B(x, r)$ such that $B(x, r) \subseteq U_\alpha$. Because $x \in U_\alpha \Rightarrow x \in \bigcup_{\alpha \in A} U_\alpha$, then $B(x, r) \subseteq \bigcup_{\alpha \in A} U_\alpha$. \square

Theorem 1.3. A subset U of a metric space X is open iff U is a union of open balls in X .

Proof: Theorem 1.1 and 1.3 prove the left implication. If U is an open subset of X , then for all $x \in U$, there exists $r(x) > 0$ such that $B(x, r(x)) \subseteq U$, so $\bigcup_{x \in U} B(x, r(x)) = U$. \square

Theorem 1.4. The intersection of any finite number of open subsets of a metric space is open.

Proof: Suppose $x \in \bigcap_{n=1}^m U_n$, a finite union of open subsets of a metric space. Then for all n , there exists $r(n) > 0$ such that $B(x, r(n)) \subseteq U_n$. Let $r = \min(r(1) \dots r(m))$. Then for all n we see $B(x, r) \subseteq B(x, r(n))$ and thus $B(x, r) \subseteq \bigcap_{n=1}^m U_n$. \square

Theorem 1.5. Let Y be a subspace of a metric space X . Then a subset U of Y is open in Y iff $U = V \cap Y$ for some open subset V of X .

Proof: Suppose $x \in V \cap Y$. Then there exists an open ball in X with radius $r(x)$ such that $B(x, r(x)) \subseteq V$, and $x \in Y$. Because $Y \subseteq X$ we see that $Y \cap B(x, r(x)) = \{y \in X \cap Y \mid d(x, y) < r(x)\} = \{y \in Y \mid d(x, y) < r(x)\}$, by definition an open ball in Y . Trivially $V \cap Y \subseteq \bigcap_{x \in V \cap Y} Y \cap B(x, r(x))$ and by definition the reverse is true.

To prove the converse, suppose $x \in U$. Then there exists an open ball in Y with radius $r(x)$ such that $B(x, r(x)) \subseteq U$. It follows from conclusions reached above that if $B'(x, r(x))$ is open in X , then $B'(x, r(x)) \cap Y = B(x, r(x))$. Let $V = \bigcup_{x \in U} B'(x, r(x))$. Then $V \cap Y \subseteq U$, and $x \in U \Rightarrow x \in V$. \square

Definition 1.7 (Adherent point). Let Y be a subset of a metric space X . A point $x \in X$ is adherent to Y if for all $r > 0$

$$B(x, r) \cap Y \neq \emptyset$$

Definition 1.8 (Closure). The closure of Y denoted by \bar{Y} , consists of all points in X that are adherent to Y .²

$$^2 Y \subseteq \bar{Y}.$$

Definition 1.9 (Closed subset). The subset Y is closed if $Y = \bar{Y}$.³

³ The empty set \emptyset and X are closed subsets of X . Interestingly, X is also open in X .

Theorem 1.6. If Y is a subset of a metric space X , then the closure of Y is closed, i.e.

$$\overline{\bar{Y}} = \bar{Y}$$

Proof: \bar{Y} contains all $x \in X$ such that for all $r > 0$ in $B(x, r) \cap Y \neq \emptyset$. Let $y \in X$ with $B(y, r') \cap \bar{Y} \neq \emptyset$ for $r' > 0$. Suppose to the

contrary that there does not exist $x \in X$ such that $x = y$. Then there exists $a = \min(d(x, y)) > 0$ such that $\forall x (x \notin B(y, a))$, therefore $B(y, a) \cap \bar{Y} = \emptyset$, a contradiction. \square

Theorem 1.7. A subset Y of a metric space X is closed iff the complement of Y is open.

Proof: If Y is closed, then Y contains all $x \in X$ such that for all $r > 0$, $B(x, r) \cap Y \neq \emptyset$. Therefore iff $y \in Y^c$ the negation is true, i.e. there exists $r' > 0$ such that $B(y, r') \cap Y = \emptyset$, and because $Y^c \cup Y = X$ we have $B(y, r') \subset Y^c$ and Y^c is open. \square

Theorem 1.8. The intersection of any family of closed sets is closed. The union of any finite family of closed sets is closed.

Proof: Let $\{Y_\alpha\}$ be a family of closed sets in X , and $\alpha \in A$, the number of elements in $\{Y_\alpha\}$. Following the fact that a union of open subsets is open, and the intersection of finite open subsets is open, as well as the previous theorem, we see

$$\begin{aligned} X \setminus \bigcup_{\alpha \in A} Y_\alpha &= \bigcap_{\alpha \in A} X \setminus Y_\alpha \\ X \setminus \bigcap_{\alpha \in A} Y_\alpha &= \bigcup_{\alpha \in A} X \setminus Y_\alpha \end{aligned}$$

\square

Definition 1.10 (Convergent sequence). A sequence $\{x_n\}_{n=1}^\infty$ in a metric space X converges to $x \in X$ if

$$\lim_{n \rightarrow \infty} d(x_n, x) = 0$$

In this case, x is the limit of $\{x_n\}$ and we write $x_n \rightarrow x$, or

$$\lim_{n \rightarrow \infty} x_n = x.$$

Lemma 1.1. The limit of a convergent sequence in a metric space is unique

Proof: Let $\lim_{n \rightarrow \infty} x_n = x, y$ and suppose to the contrary that $x \neq y$. Then $d(x, y) > 0$ and for all $\epsilon > 0$ there exists δ such that $d(x_n, x)$ and $d(x_n, y)$ are both less than $\frac{\epsilon}{2}$. But then if $\epsilon < d(x, y)$ then $d(x_n, x) + d(x_n, y) < d(x, y)$, a contradiction. \square

Theorem 1.9. Let Y be a subset of the metric space X , then $x \in X$ is adherent to Y iff there is a sequence in Y that converges to x .

Proof: If x is adherent to Y , then $\forall r > 0$, $B(x, r) \cap Y \neq \emptyset$, i.e. for all r there exists $y \in Y$ such that $d(x, y_n) < r$. Using this fact we can construct a sequence that converges to x . Let $y_n \in Y$, and $\{y_n\}$ be a sequence such that for all $\epsilon > 0$ there exists $N \in \mathbb{N}$ such that $n > N$ implies $d(x, y) < \epsilon$.

Let $\{y_n\}$ be a sequence with $y_n \in Y$, and let $x \in X$. Let $\{y_n\}$ be such that for all $\epsilon > 0$, $n \in \mathbb{N}$ with $n > N$ implies $d(x, y_n) < \epsilon$. Then for all $r > 0$ there exists $r = \epsilon$ such that $y_n \in B(x, r)$, and thus $B(x, r) \cap Y \neq \emptyset$ for all $r > 0$. \square

Completeness

Definition 2.1 (Cauchy sequence). A sequence $\{x_n\}_{n=1}^{\infty}$ in a metric space X is a Cauchy sequence if

$$\lim_{m, n \rightarrow \infty} d(x_n, x_m) = 0.$$

Lemma 2.1. A convergent sequence is a Cauchy sequence.

Proof: Suppose $\{x_n\}$ a sequence that converges to x . Then

$$\forall \epsilon > 0, \exists n, m > N (d(x_n, x), d(x_m, x) < \epsilon).$$

If we choose N such that $d(x_n, x), d(x_m, x) < \frac{\epsilon}{2}$ then

$$d(x_n, x) + d(x_m, x) < \epsilon \Rightarrow d(x_n, x_m) < \epsilon.$$

\square

Lemma 2.2. If $\{x_n\}$ is a Cauchy sequence and if there is a subsequence $\{x_{n_k}\}_{k=1}^{\infty}$ of $\{x_n\}$ that converges to x , then $\{x_n\}$ converges to x .

Proof: Suppose $\{x_n\}$ a convergent sequence and $\{x_{n_k}\}_{k=1}^{\infty}$ a subsequence which converges to x then

$$\forall \delta > 0, \exists N (n_k > N \Rightarrow d(x_{n_k}, x) < \delta)$$

$$\forall \epsilon > 0, \exists M (n > M \wedge n_k > M, N \Rightarrow d(x_n, x_{n_k}) < \epsilon).$$

Because $d(x_n, x_{n_k}) + d(x_{n_k}, x) < \epsilon + \delta$ then $d(x_n, x) < \epsilon + \delta$. \square

Definition 2.2 (Complete metric space). A metric space X is complete if every Cauchy sequence in X converges.

Theorem 2.1. A complete subspace Y of a metric space X is closed in X

Proof: If $x \in \bar{Y}$, then $\forall r > 0, \exists B(x, r)$ such that $B(x, r) \cap Y \neq \emptyset$, so $\exists y \in Y$ such that $d(x, y) < r$. It follows there exists a Cauchy sequence $\{y_n\}$ in Y with limit x such that $\forall r, \exists N (n > N \Rightarrow d(x, y_n) < r)$. And because every Cauchy sequence in Y converges, $x \in Y$ and $\bar{Y} = Y$. \square

Definition 2.3 (Uniform convergence). Let $\{f_n\}_{n=1}^{\infty}$ be a sequence of functions from a set S to a metric space X and let f be a function from S to X . The sequence $\{f_n\}$ converges uniformly to f on S if for each $\epsilon > 0$ there exists an integer N such that $d(f_n(s), f(s)) < \epsilon$ for all integers $n \geq N$ and for all $s \in S$.

Definition 2.4. A sequence $\{f_n\}$ of functions from S to X is a Cauchy sequence of functions if for each $\epsilon > 0$ there exists an integer N such that

$$d(f_n(s), f_m(s)) < \epsilon, \quad \text{all } s \in S, n, m \geq N.$$

Theorem 2.2. Let S be a set, and let X be a complete metric space.

If $\{f_n\}_{n=1}^{\infty}$ is a Cauchy sequence of functions from S to X , then there exists a function f from S to X such that $\{f_n\}$ converges uniformly to f .

Definition 2.5 (Dense subsets). A subset T of a metric space X is dense in X if $\overline{T} = X$.

Theorem 2.3 (Baire Category Theorem). Let $\{U_n\}_{n=1}^{\infty}$ be a sequence of dense open subsets of a complete metric space X . Then $\bigcap_{n=1}^{\infty} U_n$ is also dense in X .