

Advanced Calculus

Samuel Lindskog

September 20, 2024

Test 1

Definition 1.1 (Relation). A relation between A and B is any subset R of $A \times B$. If $(a, b) \in R$, then we say aRb .

Definition 1.2 (Equivalence Relation). A relation R on a set S is an equivalence relation if it has the following properties for all x, y, z in S :

1. xRx (reflexive property)
2. $xRy \Rightarrow yRx$ (Symmetric property)
3. $xRy \wedge yRz \Rightarrow xRz$ (Transitive property)

A partition of a set S is a collection \mathcal{P} of nonempty subsets such that

1. $x \in S \Rightarrow x \in \bigcup_{A \in \mathcal{P}} A$
2. $\forall A, B \in \mathcal{P}, A \neq B \Rightarrow A \cap B = \emptyset$

Definition 1.3 (Function). Let A and B be sets. A function from A to B is a nonempty relation $f \subseteq A \times B$ that satisfies the following two conditions:

1. $\forall a \in A, \exists b \in B, (a, b) \in f$
2. $(a, b) \in f \wedge (a, c) \in f \Rightarrow b = c$

Definition 1.4 (Upper bound). Let $S \subseteq \mathbb{R}$. If there exists a real number m such that $m \geq s$ for all $s \in S$, then m is an upper bound of S .

Definition 1.5 (Maximum). If an upper bound m of S is a member of S , then m is called the maximum of S .

Definition 1.6 (Supremum). Let S be a nonempty subset of \mathbb{R} . If S is bounded above, then the least upper bound of S is called the supremum. Thus $m = \sup S$ iff

1. $\forall s \in S, m \geq s$
2. $m' < m \Rightarrow \exists s' \in S, s' > m'$

Axiom 1.1 (Completeness of \mathbb{R}). Every nonempty subset S of \mathbb{R} that is bounded above has a least upper bound, i.e. $\sup S$ exists.