

(19)



(11)

EP 3 247 711 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
09.03.2022 Bulletin 2022/10

(51) International Patent Classification (IPC):
C07D 405/14 ^(2006.01) **A61K 31/4439** ^(2006.01)
A61P 31/10 ^(2006.01)

(21) Application number: **16703009.7**

(52) Cooperative Patent Classification (CPC):
A61P 31/10; C07D 405/14

(22) Date of filing: **14.01.2016**

(86) International application number:
PCT/US2016/013356

(87) International publication number:
WO 2016/118396 (28.07.2016 Gazette 2016/30)

(54) **NOVEL SALTS AND POLYMORPHS OF SCY-078**

NEUE SALZE UND POLYMORPHE VON SCY-078

NOUVEAUX SELS ET POLYMORPHES DE SCY-078

(84) Designated Contracting States:
**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR**

(30) Priority: **19.01.2015 PCT/CN2015/070967**

(43) Date of publication of application:
29.11.2017 Bulletin 2017/48

(73) Proprietor: **Scynexis, Inc.**
Research Triangle Park, NC 27709-2878 (US)

(72) Inventor: **ZHANG, Yi**
Suzhou
Jiangsu 215123 (CN)

(74) Representative: **Finnegan Europe LLP**
1 London Bridge
London SE1 9BG (GB)


(56) References cited:
WO-A1-2010/019203

EP 3 247 711 B1

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description

[0001] SCY-078 (or "compound 1") is a glucan synthase inhibitor useful as an antifungal compound. SCY-078 is useful for treating, among other things, *Invasive Candidiasis* and *Invasive Aspergillosis*. Other antimicrobial utilities of SCY-078 are disclosed, for example, in U.S. Patent No. 8,188,085. SCY-078 has the following chemical structure:



[0002] WO2010/019203 discloses novel derivatives of enfumafungin, along with their pharmaceutically acceptable salts, hydrates and prodrugs. Also disclosed are compositions comprising such compounds, methods of preparing such compounds and methods of using such compounds as antifungal agents and/or inhibitors of (1,3)- β -D-glucan synthase. The disclosed compounds, their pharmaceutically acceptable salts, hydrates and prodrugs, as well as compositions comprising such compounds, salts, hydrates and prodrugs, are useful for treating, and/or preventing fungal infections and associated diseases and conditions.

[0003] The present disclosure is directed to citrate salts of SCY-078 and polymorphs of those salts. In another embodiment, the salts and polymorphs thereof exhibit certain kinetic solubilities. Higher kinetic solubilities can be importance in formulations, such as intravenous formulations. In yet another embodiment, the salts and polymorphs thereof exhibit certain hygroscopicities. Hygroscopy has been found to play an important role in the preparation of solid dosage forms and the selection of excipients. The present disclosure is further directed to, among other things, methods of preparing citrate salts of SCY-078 and polymorphs thereof.

[0004] The present disclosure is additionally directed to, among other things, pharmaceutical compositions comprising citrate salts of SCY-078 and polymorphs thereof. In another embodiment, the disclosure relates to methods of preparing pharmaceutical compositions comprising citrate salts of SCY-078 and polymorphs thereof suitable for injection or intravenous administration. In yet another embodiment, the present disclosure relates to citrate salts of the present invention for use in treating fungal infections by administering citrate salts of SCY-078 and polymorphs thereof.

BRIEF DESCRIPTION OF DRAWINGS

[0005]

Fig. 1 is an X-ray Powder Diffraction "XRPD" pattern of SCY-078 phosphate from Example 1 (not according to the invention).

Fig. 2 is a DSC curve and a TGA curve of SCY-078 phosphate from Example 1 (not according to the invention).

Fig. 3 is an XRPD of SCY-078 crystalline freebase (MeOH desolvate), batch 1 from Example 2 (not according to the invention).

Fig. 4 is a DSC curve and a TGA curve of SCY-078 crystalline freebase (MeOH desolvate), batch 1 from Example 2 (not according to the invention).

Fig. 5 is an XRPD of SCY-078 crystalline freebase (MeOH desolvate), batch 2 from Example 2 (not according to the invention).

Fig. 6 is a DSC curve and a TGA curve of SCY-078 amorphous freebase from Example 3 (not according to the invention).


Fig. 7 is an XRPD of SCY-078 HCl Type A from Example 7 (not according to the invention).

Fig. 8 is a DSC curve and a TGA curve of SCY-078 HCl Type A from Example 7 (not according to the invention).

Fig. 9 is an XRPD of SCY-078 citrate Type A from Example 8.

Fig. 10 is a DSC curve and a TGA curve of SCY-078 citrate Type A from Example 8.

Fig. 11 is an XRPD of SCY-078 hippurate Type A from Example 9 (not according to the invention).



[0009] The term "HCl" refers to hydrochloric acid. The term "Ca" refers to calcium.

[0010] The terms "pharmaceutically acceptable salt" and the like should be understood to include, but not limited to, citrate salts, hippurate salts, fumarate salts, glycolate salts, mesylate salts, and calcium salts.

[0011] As used here, phrases such as "SCY-078 salt," "SCY-078 salts," "salt of SCY-078," "salts of SCY-078," "pharmaceutically acceptable salt of SCY-078," and "pharmaceutically acceptable salts thereof" should be understood to be salts in various forms, for example, the polymorphs disclosed herein. In addition, as used here, phrases such as "SCY-078 phosphate," "SCY-078 citrate," "SCY-078 hippurate," "SCY-078 glycolate," "SCY-078 mesylate," "SCY-078 fumarate," and "SCY-078 calcium" should be understood to be salts in various forms, for example, the polymorphs disclosed herein.

[0012] The term "solvent" and the like refer to any appropriate aqueous or organic solvent. Solvents include, but are not limited to, methanol, acetic acid, tetrahydrofuran, 2-methyl-tetrahydrofuran, 1,4-dioxane, n-methyl-2-pyrrolidone, dimethyl sulfoxide, dimethylacetamide, isopropyl alcohol, acetonitrile, acetone, ethyl acetate, water and mixtures thereof.

[0013] The term "pharmaceutically acceptable carrier" and the like refer to an ingredient that is compatible with SCY-078 and is not harmful to a patient's health. Pharmaceutically acceptable carriers include, but are not limited to, one or more of the following: aqueous vehicles and solvents, such as water, saline solutions, and alcohols; buffers; surface active agents; dispersing agents; inert diluents; preservatives; suspending agents; emulsifying agents; demulcents; thickening agents; emulsifying agents; antioxidants; and stabilizing agents. Other additional ingredients that may be included in the pharmaceutical compositions of the disclosure are generally known in the art and may be described, for example, in Remington's Pharmaceutical Sciences, Mack Publishing Co., Easton, Pa.

[0014] The term "injection" and the like refer to the insertion of a composition into the body by syringe, hollow needle, or the like. The term "injection" and the like include, but are not limited to, intravenous injections, including those entailing administering using an IV bag containing a diluent.

[0015] The term "effective amount" refers to an amount of the active ingredient that, when administered to a subject, alleviates at least some of the symptoms or stops the progression of the identified disease or condition. The terms "disease" or "condition" include, but are not limited to, infections such as fungal infections. Exemplary dosage amounts can be found, for example, in U.S. Patent No. 8,188,085.

[0016] The term "Å" refers to angstroms. Terms such as "2θ" or "2 Th." refer to degrees 2 theta.

[0017] The XRPD peaks recited herein should be understood to reflect a precision of ± 0.2 for the 2 theta peaks, and the equivalent precision for d-spacings as per Bragg's law. The present disclosure also fully incorporates section 941 of the United States Pharmacopeia. The National Formulary from 2014 (USP 37/NF 32, volume 1) relating to characterization of crystalline and partially crystalline solids by X-ray Powder Diffraction.

[0018] The present disclosure relates to, among other things, citrate salts of SCY-078. In a further embodiment, the salt is SCY-078 citrate Type A.

[0019] The present disclosure further relates to citrate salts of SCY-078 that have a chemical purity of at least 90%. In another embodiment, citrate salts of SCY-078 have a chemical purity of at least 95%. In a further embodiment, citrate salts of SCY-078 have a chemical purity of at least 98%. In yet another embodiment, citrate salts of SCY-078 have a chemical purity of at least 99%.

[0020] The present disclosure additionally relates to citrate salts of SCY-078 that have a kinetic solubility of at least 2 mg/mL at 4 hours in dextrose buffer at pH 5.5. In one embodiment, the citrate salts of SCY-078 have a kinetic solubility of at least 4 mg/mL at 4 hours in dextrose buffer at pH 5.5. In another embodiment, the citrate salts of SCY-078 have a kinetic solubility of from 2 mg/mL to 5 mg/mL at 4 hours in dextrose buffer at pH 5.5. In yet another embodiment, the citrate salts of SCY-078 have a kinetic solubility of from 4 mg/mL to 5 mg/mL at 4 hours in dextrose buffer at pH 5.5. In still another embodiment, the citrate salts of SCY-078 have a kinetic solubility greater than SCY-078 (as a freebase) at 4 hours in dextrose buffer at pH 5.5.

[0106] For precipitates, the solids were isolated and analyzed by XRPD. Clear solutions were evaporated slowly to dryness at room temperature.

[0107] The salt study (Table 8) showed that seven crystalline salts (eight crystal forms) of SCY-078 were found: HCl Type A, citrate Type A, hippurate Type A, fumarate Type A, fumarate Type B, glycolate Type A, mesylate Type A, and Ca salt Type A. Four crystal forms of SCY-078 freebase were discovered during the salt study and were identified as freebase ("FB") Type A, B, C, and D.

Table 8

Acid \ Solvent	EtOH (A)	IPA (B)	ACN (C)	Acetone (D)	EtOAc (E)	THF/H ₂ O (19:1, v/v) (F)
HCl	oil	oil	HCl salt Type A	HCl salt Type A	amorphous	oil
H ₃ PO ₄	oil	oil	oil	oil	oil	oil
Maleic Acid	amorphous	amorphous	FB Type A	amorphous	amorphous	amorphous
Citric Acid	amorphous	amorphous	Citrate Type A	amorphous	FB Type A	amorphous
Hippuric Acid	FB Type B	FB Type B	Hippurate Type A	Hippurate Type A	Hippurate Type A	Low crystallinity
Adipic Acid	amorphous	FB Type A	FB Type A	FB Type C	amorphous	amorphous
Fumaric Acid	amorphous	amorphous	Fumarate Type A	FB Type D	Fumarate Type B	amorphous
Glutaric Acid	amorphous	FB Type A	FB Type A	FB Type C	FB Type A	amorphous
Glycolic Acid	FB Type B	FB Type B	FB Type B	FB Type C	FB Type B	Glycolate Type A
D-Glutamic Acid	MeOH desolvate+acid	FB Type A	FB Type A	FB Type C	FB Type A	amorphous
Acetic Acid	oil	oil	amorphous	amorphous	amorphous	oil
Mucic Acid	amorphous	amorphous	FB Type D	FB Type D	FB Type D	amorphous
L-Malic Acid	amorphous	amorphous	FB Type A	FB Type C	FB Type A	amorphous
Benzoic acid	MeOH Desolvate	FB Type A	FB Type A	FB Type C	FB Type A	amorphous
Methanesulfonic Acid	amorphous	amorphous	FB Type B	Mesylate Type A	FB Type B	amorphous
Malonic Acid	amorphous	amorphous	FB Type A	amorphous	FB Type A	amorphous
Ethanesulfonic Acid	amorphous	oil	FB Type B	FB Type B	FB Type B	amorphous
Ca(OH) ₂	Ca(OH) ₂	Ca(OH) ₂	FB Type A	FB Type C	Ca(OH) ₂	Ca salt Type A

All the above salt formation experiments were performed at RT using 1.0 mole equivalency of acid.
FB: freebase

EXAMPLE 7 (not according to the invention)


[0108] SCY-078 HCl Type A: SCY-078 HCl Type A prepared from the salt study in Example 6 was characterized by XRPD, DSC, and TGA (Figs. 7-8). The resulting XRPD pattern evidenced that SCY-078 HCl Type A is weakly crystalline and has a unique form as compared to the freebase MeOH desolvate. The 2 theta and d-spacing values are summarized in Table 9. The DSC curve displayed an endotherm at 48.5 °C (onset temperature). The TGA curve showed a weight loss of 14.4% before 130 °C.

TABLE 9

Pos. [°2Th.]	Height [cts]	FWHM Left [°2Th.]	d-spacing [Å]	Rel. Int. [%]
5.635436	167.080400	0.153504	15.68264	21.28
9.023307	62.106750	0.307008	9.80065	7.91
11.313240	785.111500	0.179088	7.82150	100.00

Claims

1. A citrate salt of compound 1:



2. The citrate salt of claim 1, wherein the citrate salt of compound 1 comprises

Type A citrate crystal form, wherein the Type A citrate crystal form has an XRPD pattern comprising one or more peaks at d-spacings of 11.86, 7.70, 7.09, 6.71, 5.90, and 5.29 Å and/or an XRPD pattern comprising one or more peaks at degrees 2 theta of 7.45, 11.49, 12.49, 13.19, 15.02, and 16.75; or

Type B citrate crystal form, wherein the Type B citrate crystal form has an XRPD pattern comprising one or more peaks at d-spacings of 15.89, 12.77, 7.54, 5.82, and 5.27 Å and/or an XRPD pattern comprising one or more peaks at degrees 2 theta of 5.56, 6.92, 11.73, 15.23, and 16.81; or

Type E citrate crystal form, wherein the Type E citrate crystal form has an XRPD pattern comprising one or more peaks at d-spacings of 12.18, 7.74, 6.27, 5.62, and 5.43 Å and/or an XRPD pattern comprising one or more peaks at degrees 2 theta of 7.26, 11.44, 14.14, 15.76, and 16.33; or

Type F citrate crystal form, wherein the Type F citrate crystal form has an XRPD pattern comprising one or more peaks at d-spacings of 24.32 and 5.00 Å and/or an XRPD pattern comprising one or more peaks at degrees 2 theta of 3.63 and 17.74; or

Type M citrate crystal form, wherein the Type M citrate crystal form has an XRPD pattern comprising one or more peaks at d-spacings of 12.04, 7.69, 6.25, 5.61, and 5.41 Å and/or an XRPD pattern comprising one or more peaks at degrees 2 theta of 7.34, 11.51, 14.17, 15.80, and 16.37; or

Type N citrate crystal form, wherein the Type N citrate crystal form has an XRPD pattern comprising one or more peaks at d-spacings of 12.51, 7.77, 6.85, 6.27, 5.84, 5.45, and 4.79 Å and/or an XRPD pattern comprising one or more peaks at degrees 2 theta of 7.07, 11.38, 12.92, 14.13, 15.16, 16.26, and 18.51; or

Type O citrate crystal form, wherein the Type O citrate crystal form has an XRPD pattern comprising one or more peaks at d-spacings of 12.48, 7.43, and 5.29 Å and/or an XRPD pattern comprising one or more peaks at degrees 2 theta of 7.08, 11.91, and 16.76; or

Type Q citrate crystal form, wherein the Type Q citrate crystal form has an XRPD pattern comprising one or more peaks at d-spacings of 14.03, 12.83, 7.81, 7.54, and 5.21 Å and/or an XRPD pattern comprising one or more peaks at degrees 2 theta of 6.30, 6.89, 11.33, 11.73, and 17.01; or


Type R citrate crystal form, wherein the Type R citrate form has an XRPD pattern comprising one or more peaks at d-spacings of 14.39, 6.05, 5.40, and 5.00 Å and/or an XRPD pattern comprising one or more peaks at degrees 2 theta of 6.14, 14.64, 16.41, and 17.74; or

Type S citrate crystal form, wherein the Type S citrate crystal form has an XRPD pattern comprising one or more peaks at d-spacings of 16.03, 12.12, 7.37, and 5.27 Å and/or an XRPD pattern comprising one or more peaks at degrees 2 theta of 5.51, 7.30, 12.00, and 16.81.

3. The citrate salt of claim 1, wherein the citrate salt of compound 1 comprises

Type A citrate crystal form, wherein the Type A citrate crystal form has an XRPD pattern comprising one or more peaks at d-spacings of 11.86, 7.70, 7.09, 6.71, 5.90, and 5.29 Å and/or an XRPD pattern comprising one or more peaks at degrees 2 theta of 7.45, 11.49, 12.49, 13.19, 15.02, and 16.75.

4. The citrate salt of claim 1, wherein the citrate salt of compound 1



2. Citratsalz nach Anspruch 1, wobei das Citratsalz von Verbindung 1 Folgendes umfasst


Citrat-Kristallform Typ A, wobei die Citrat-Kristallform Typ A ein XRPD-Muster aufweist, umfassend ein oder mehrere Peaks in d-Abständen von 11,86, 7,70, 7,09, 6,71, 5,90 und 5,29 Å und/oder ein XRPD-Muster, umfassend ein oder mehrere Peaks bei Grad 2-Theta von 7,45, 11,49, 12,49, 13,19, 15,02 und 16,75; oder Citrat-Kristallform Typ B, wobei die Citrat-Kristallform Typ B ein XRPD-Muster aufweist, umfassend ein oder mehrere Peaks in d-Abständen von 15,89, 12,77, 7,54, 5,82 und 5,27 Å und/oder ein XRPD-Muster, umfassend ein oder mehrere Peaks bei Grad 2-Theta von 5,56, 6,92, 11,73, 15,23 und 16,81; oder Citrat-Kristallform Typ E, wobei die Citrat-Kristallform Typ E ein XRPD-Muster aufweist, umfassend ein oder mehrere Peaks in d-Abständen von 12,18, 7,74, 6,27, 5,62 und 5,43 Å und/oder ein XRPD-Muster, umfassend ein oder mehrere Peaks bei Grad 2-Theta von 7,26, 11,44, 14,14, 15,76 und 16,33; oder Citrat-Kristallform Typ F, wobei die Citrat-Kristallform Typ F ein XRPD-Muster aufweist, umfassend ein oder mehrere Peaks in d-Abständen von 24,32 und 5,00 Å und/oder ein XRPD-Muster, umfassend ein oder mehrere Peaks bei Grad 2-Theta von 3,63 und 17,74; oder Citrat-Kristallform Typ M, wobei die Citrat-Kristallform Typ M ein XRPD-Muster aufweist, umfassend ein oder mehrere Peaks in d-Abständen von 12,04, 7,69, 6,25, 5,61 und 5,41 Å und/oder ein XRPD-Muster, umfassend ein oder mehrere Peaks bei Grad 2-Theta von 7,34, 11,51, 14,17, 15,80 und 16,37; oder Citrat-Kristallform Typ N, wobei die Citrat-Kristallform Typ N ein XRPD-Muster aufweist, umfassend ein oder mehrere Peaks in d-Abständen von 12,51, 7,77, 6,85, 6,27, 5,84, 5,45 und 4,79 Å und/oder ein XRPD-Muster, umfassend ein oder mehrere Peaks bei Grad 2-Theta von 7,07, 11,38, 12,92, 14,13, 15,16, 16,26 und 18,51; oder Citrat-Kristallform Typ O, wobei die Citrat-Kristallform Typ O ein XRPD-Muster aufweist, umfassend ein oder mehrere Peaks in d-Abständen von 12,48, 7,43 und 5,29 Å und/oder ein XRPD-Muster, umfassend ein oder mehrere Peaks bei Grad 2-Theta von 7,08, 11,91 und 16,76; oder Citrat-Kristallform Typ Q, wobei die Citrat-Kristallform Typ Q ein XRPD-Muster aufweist, umfassend ein oder mehrere Peaks in d-Abständen von 14,03, 12,83, 7,81, 7,54 und 5,21 Å und/oder ein XRPD-Muster, umfassend ein oder mehrere Peaks bei Grad 2-Theta von 6,30, 6,89, 11,33, 11,73 und 17,01; oder Citrat-Kristallform Typ R, wobei die Citrat-Kristallform Typ R ein XRPD-Muster aufweist, umfassend ein oder mehrere Peaks in d-Abständen von 14,39, 6,05, 5,40 und 5,00 Å und/oder ein XRPD-Muster, umfassend ein oder mehrere Peaks bei Grad 2-Theta von 6,14, 14,64, 16,41 und 17,74; oder Citrat-Kristallform Typ S, wobei die Citrat-Kristallform Typ S ein XRPD-Muster aufweist, umfassend ein oder mehrere Peaks in d-Abständen von 16,03, 12,12, 7,37 und 5,27 Å und/oder ein XRPD-Muster, umfassend ein oder mehrere Peaks bei Grad 2-Theta von 5,51, 7,30, 12,00 und 16,81.

3. Citratsalz nach Anspruch 1, wobei das Citratsalz von Verbindung 1 Folgendes umfasst

Citrat-Kristallform Typ A, wobei die Citrat-Kristallform Typ A ein XRPD-Muster aufweist, umfassend ein oder mehrere Peaks in d-Abständen von 11,86, 7,70, 7,09, 6,71, 5,90 und 5,29 Å und/oder ein XRPD-Muster, umfassend ein oder mehrere Peaks bei Grad 2-Theta von 7,45, 11,49, 12,49, 13,19, 15,02 und 16,75.

4. Citratsalz nach Anspruch 1, wobei das Citratsalz der Verbindung 1

im Wesentlichen aus Citrat-Kristallform Typ A besteht; oder
mindestens 98 % Citrat-Kristallform Typ A umfasst; oder
mindestens 99 % Citrat-Kristallform Typ A umfasst;



2. Sel de citrate selon la revendication 1, ledit sel de citrate du composé 1 comprenant


une forme cristalline de citrate de type A, ladite forme cristalline de citrate de type A présentant un motif XRPD comprenant un ou plusieurs pics au niveau des espacements d de 11,86, 7,70, 7,09, 6,71, 5,90 et 5,29 Å et/ou un motif XRPD comprenant un ou plusieurs pics à 7,45, 11,49, 12,49, 13,19, 15,02 et 16,75 degrés 2 θ ; ou une forme cristalline de citrate de type B, ladite forme cristalline de citrate de type B présentant un motif XRPD comprenant un ou plusieurs pics au niveau des espacements d de 15,89, 12,77, 7,54, 5,82 et 5,27 Å et/ou un motif XRPD comprenant un ou plusieurs pics à 5,56, 6,92, 11,73, 15,23 et 16,81 degrés 2 θ ; ou une forme cristalline de citrate de type E, ladite forme cristalline de citrate de type E présentant un motif XRPD comprenant un ou plusieurs pics au niveau des espacements d de 12,18, 7,74, 6,27, 5,62 et 5,43 Å et/ou un motif XRPD comprenant un ou plusieurs pics à 7,26, 11,44, 14,14, 15,76 et 16,33 degrés 2 θ ; ou une forme cristalline de citrate de type F, ladite forme cristalline de citrate de type F présentant un motif XRPD comprenant un ou plusieurs pics au niveau des espacements d de 24,32 et 5,00 Å et/ou un motif XRPD comprenant un ou plusieurs pics à 3,63 et 17,74 degrés 2 θ ; ou une forme cristalline de citrate de type M, ladite forme cristalline de citrate de type M présentant un motif XRPD comprenant un ou plusieurs pics au niveau des espacements d de 12,04, 7,69, 6,25, 5,61 et 5,41 Å et/ou un motif XRPD comprenant un ou plusieurs pics à 7,34, 11,51, 14,17, 15,80 et 16,37 degrés 2 θ ; ou une forme cristalline de citrate de type N, ladite forme cristalline de citrate de type N présentant un motif XRPD comprenant un ou plusieurs pics au niveau des espacements d de 12,51, 7,77, 6,85, 6,27, 5,84, 5,45 et 4,79 Å et/ou un motif XRPD comprenant un ou plusieurs pics à 7,07, 11,38, 12,92, 14,13, 15,16, 16,26 et 18,51 degrés 2 θ ; ou une forme cristalline de citrate de type O, ladite forme cristalline de citrate de type O présentant un motif XRPD comprenant un ou plusieurs pics au niveau des espacements d de 12,48, 7,43 et 5,29 Å et/ou un motif XRPD comprenant un ou plusieurs pics à 7,08, 11,91 et 16,76 degrés 2 θ ; ou une forme cristalline de citrate de type Q, ladite forme cristalline de citrate de type Q présentant un motif XRPD comprenant un ou plusieurs pics au niveau des espacements d de 14,03, 12,83, 7,81, 7,54 et 5,21 Å et/ou un motif XRPD comprenant un ou plusieurs pics à 6,30, 6,89, 11,33, 11,73 et 17,01 degrés 2 θ ; ou une forme cristalline de citrate de type R, ladite forme cristalline de citrate de type R présentant un motif XRPD comprenant un ou plusieurs pics au niveau des espacements d de 14,39, 6,05, 5,40 et 5,00 Å et/ou un motif XRPD comprenant un ou plusieurs pics à 6,14, 14,64, 16,41 et 17,74 degrés 2 θ ; ou une forme cristalline de citrate de type S, ladite forme cristalline de citrate de type S présentant un motif XRPD comprenant un ou plusieurs pics au niveau des espacements d de 16,03, 12,12, 7,37 et 5,27 Å et/ou un motif XRPD comprenant un ou plusieurs pics à 5,51, 7,30, 12,00 et 16,81 degrés 2 θ .

3. Sel de citrate selon la revendication 1, ledit sel de citrate du composé 1 comprenant

une forme cristalline de citrate de type A, ladite forme cristalline de citrate de type A présentant un motif XRPD comprenant un ou plusieurs pics au niveau des espacements d de 11,86, 7,70, 7,09, 6,71, 5,90 et 5,29 Å et/ou un motif XRPD comprenant un ou plusieurs pics à 7,45, 11,49, 12,49, 13,19, 15,02 et 16,75 degrés 2 θ .

4. Sel de citrate selon la revendication 1, ledit sel de citrate du composé 1

étant constitué essentiellement de la forme cristalline de citrate de type A ; ou comprenant au moins 98 % de la forme cristalline de citrate de type A ; ou



XRPD of SCY-078 phosphate

Figure 1