


[0010] US 2015/164865 (D1) describes oral formulations of diphenylsulfonamide endothelin and angiotensin II receptor agonists to treat elevated blood pressure and diabetic nephropathy. Komers et al. (Am J Physiol Regul Integr Comp Physiol 310(10):R877-84, 2016) (D2) describes dual inhibition of renin-angiotensin-aldosterone system and endothelin-1 in the treatment of chronic kidney disease (CKD). Savage et al. (J. Am. Soc. Nephrol. 24(3): 364-175, 2013) (D8) describes expert guidelines for the management of Alport syndrome, and thin basement membrane nephropathy.

BRIEF SUMMARY

[0011] The present invention is defined by the claims, and is directed at a pharmaceutical composition comprising a compound having structure (I),



(I)

or a pharmaceutically acceptable salt thereof, for use in a method of treating Alport syndrome.


[0012] To the extent that other compositions and/or other uses are disclosed herein, they are mentioned merely for reference purposes. For example, pharmaceutical compositions comprising a compound having structure (I), or a pharmaceutically acceptable salt thereof, may be used in methods of treating a kidney disease or disorder in a subject in need thereof, the methods comprising administering to said subject said pharmaceutical composition (i) in an amount sufficient to achieve a urine protein to creatinine ("UP/C") ratio of less than or equal to 1.5 g/g; (ii) in an amount sufficient to achieve or maintain a UP/C ratio of less than or equal to 1.5 g/g; or (iii) at a dosing regimen sufficient to achieve or maintain a UP/C ratio of less than or equal to 1.5 g/g.

[0013] A first disclosure is directed to pharmaceutical compositions comprising a compound having structure (I), or a pharmaceutically acceptable salt thereof, for use in therapeutic methods of (i) maintaining a UP/C ratio at less than or equal to 1.5 g/g in a subject in need thereof, the method comprising administering to said subject said pharmaceutical composition in an amount sufficient to maintain a UP/C ratio of less than or equal to 1.5 g/g; or (ii) reducing a UP/C ratio to less than or equal to 1.5 g/g in a subject in need thereof, the method comprising administering to said subject said pharmaceutical composition in an amount sufficient to reduce said patient's UP/C ratio to less than or equal to 1.5 g/g.

[0014] A second disclosure is directed to methods of treating a kidney disease or disorder in a subject in need thereof, the methods comprising administering to the subject a pharmaceutical composition comprising a compound having structure (I),

Scheme I

Part 1



Part 2

