




Formula (I).

[0006] Milvexian has the chemical name (5R,9S)-9-(4-(5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl)-6-oxopyrimidin-1(6H)-yl)-21-(difluoromethyl)-5-methyl-21H-3-aza-1(4,2)-pyridina-2(5,4)-pyrazolacyclonaphan-4-one. Milvexian and a method of preparing milvexian are described in U.S. patent 9,453,018, which is hereby incorporated by reference in its entirety.

[0007] An amorphous solid dispersion (ASD) composition of milvexian in one or more polymers prepared by solvent-based spray drying methods, capsules thereof, and roller compaction tablets thereof prepared by dry granulation methods have been described in WO2020210629, which is hereby incorporated by reference in its entirety.

[0008] Tablets are composed of one or more active compounds and of tableting excipients, such as diluents, binders, lubricants and disintegrating agents. The active compound and the excipients are generally provided in the form of powders which are subjected to tableting, with or without preliminary treatment. There are currently three general methods of tablet preparation: (1) wet-granulation method, (2) dry-granulation method (slugging or roller compaction), and (3) direct compression (Williams et al., "Strategies to Address Low Drug Solubility in Discovery and Development", 2013, Pharmacol. Rev., V65, pp.215-499). All three methods have limitations.

[0009] Wet-granulation method is the most widely used method. Its popularity is due to the greater probability that the granules will meet all the physical requirements for the manufacture of good tablets. Its main limitations are the number of separate steps involved and the time and labor necessary to carry out the procedure. The steps involved in the wet-granulation method are: (1) weighing, (2) mixing, (3) granulation, (4) screening the damp mass after granulation, (5) drying (6) dry screening (7) lubrication, and (8) compression.