

A new spatial model for points above a threshold

December 11, 2014

1 Introduction

In most climatological applications, researchers are interested in learning about the average behavior of different climate variables (e.g. ozone, temperature, rainfall). However, averages do not help regulators prepare for the unusual events that only happen once every 100 years. For example, it is important to have an idea of how much rain will come in a 100-year floor in order to construct strong enough river levees to protect lands from flooding.

Unlike multivariate normal distributions, it is challenging to model multivariate extreme value distributions (e.g. generalized extreme value and generalized Pareto distribution) because few closed-form expressions exist for the density in more than two-dimensions (Coles and Tawn, 1991). Given this limitation, pairwise composite likelihoods have been used when modeling dependent extremes (Padoan et al., 2010; Blanchet and Davison, 2011; Huser, 2013).

One way around the multi-dimensional limitation of multivariate extreme value distributions is to use skew elliptical distributions to model dependent extreme values (Genton, 2004; Zhang and El-Shaarawi, 2010; Padoan, 2011). Due to their flexibility, the skew-normal and skew- t distribution offer a flexible way to handle non-symmetric data within a framework of multivariate normal and multivariate t -distributions. As with the spatial Gaussian process, the skew-normal distribution is also asymptotically independent; however, the skew- t does demonstrate asymptotic dependence (Padoan, 2011). Although asymptotic dependence is desirable between sites that are near one another, one drawback to the skew- t is that sites remain asymptotically dependent even at far distances.

In this paper, we present a model that has marginal distributions with flexible tails, demonstrates asymptotic dependence for observations at sites that are near to one another, and has computation on the order of Gaussian models for large space-time datasets. Specifically, our contribution is to incorporate thresholding and random spatial partitions using a multivariate skew- t distribution. The advantage of using a thresholded model as opposed to a non-thresholded model is that it allows for the tails of the distribution to inform the predictions in the tails (DuMouchel, 1983). The random spatial partition alleviates the long-range spatial dependence seen by the skew- t .

The paper is organized as follows. Section 2.1 is a brief review of the spatial skew- t process. In section 3.3, we build upon the traditional skew- t by incorporating censoring to focus on tails, partitioning to remove long-range asymptotic dependence, and extending the model to space-time data. The computing is described in section 4. In section 5, we present a simulation study that examines the predictive capabilities of this model compared with a naïve Gaussian method. We then compare our method to Gaussian and max-stable methods with a data analysis of ozone measurements from the eastern US in section 6. The final section provides brief discussion and direction for future research.

2 Spatial skew processes

Many types of data demonstrate some level of skewness and therefore should be modeled with distributions that allow for asymmetry. The skew-elliptical family of distributions provides models that are mathematically tractable while introducing a slant parameter to account for asymmetric data (Genton, 2004). A brief review of the additive process by which a skew- t process is created is given here.

2.1 Skew- t process

Let $Y(\mathbf{s})$ be the observation at spatial location $\mathbf{s} = (s_1, s_2)$. The spatial skew- t process can be written

$$Y(\mathbf{s}) = \mathbf{X}(\mathbf{s})^T \boldsymbol{\beta} + \lambda \sigma |z| + \sigma v(\mathbf{s}) \quad (1)$$

where $\mathbf{X}(\mathbf{s})$ is a set of spatial covariates at site \mathbf{s} , $\boldsymbol{\beta}$ is a set of regression parameters, $\lambda \in \mathcal{R}$ is a parameter controlling skew, $z \sim N(0, 1)$, $\sigma^2 \sim \text{IG}(a, b)$ is an inverse gamma random variable, and $v(\mathbf{s})$ is a spatial Gaussian process with mean zero and variance one. A common spatial correlation function is the Matérn with

$$\text{cor}(v(\mathbf{s}), v(\mathbf{t})) = \gamma I(\mathbf{s} = \mathbf{t}) + (1 - \gamma) \frac{1}{\Gamma(\nu) 2^{\nu-1}} \left(\sqrt{2\nu} \frac{h}{\rho} \right)^\nu K_\nu \left(\sqrt{2\nu} \frac{h}{\rho} \right) \quad (2)$$

where ρ is the spatial range, ν is the smoothness, γ is the proportion of variance accounted for by the spatial variation, K_ν is a modified Bessel function of the second kind, and $h = \|\mathbf{s} - \mathbf{t}\|$.

Let $\mathbf{Y} = [Y(\mathbf{s}_1), \dots, Y(\mathbf{s}_n)]^T$ be a set of observations at a finite collection of locations $\mathbf{s}_1, \dots, \mathbf{s}_n$. After marginalizing over both z and σ ,

$$\mathbf{Y} \sim \text{ST}_n(\mathbf{X}\boldsymbol{\beta}, \boldsymbol{\Omega}, \boldsymbol{\alpha}, 2a), \quad (3)$$

that is, \mathbf{Y} follows an n -dimensional skew- t process with location $\mathbf{X}\boldsymbol{\beta}$, correlation matrix $\boldsymbol{\Omega}$, slant parameters $\boldsymbol{\alpha}$ and degrees of freedom $2a$, where $\mathbf{X} = [\mathbf{X}(\mathbf{s}_1)^T, \dots, \mathbf{X}(\mathbf{s}_n)^T]$, $\boldsymbol{\Omega} = \boldsymbol{\omega} \bar{\boldsymbol{\Omega}} \boldsymbol{\omega}$, $\boldsymbol{\omega} = \text{diag} \left(\frac{1}{\sqrt{ab}}, \dots, \frac{1}{\sqrt{ab}} \right)$, $\bar{\boldsymbol{\Omega}} = (\boldsymbol{\Sigma} + \lambda^2 \mathbf{1}\mathbf{1}^T)$, $\boldsymbol{\Sigma}$ is a positive definite correlation matrix, $\boldsymbol{\alpha} = \lambda(1 + \lambda^2 \mathbf{1}^T \boldsymbol{\Sigma}^{-1} \mathbf{1})^{-1/2} \mathbf{1}^T \boldsymbol{\Sigma}^{-1}$ is a vector of slant parameters. This process is desirable because of its flexible tail that is controlled by the skewness parameter λ and degrees of freedom $2a$. Furthermore, the marginal distributions at each location also follow a univariate skew- t distribution (Azzalini and Capitanio, 2013).

2.2 Extremal dependence

One measure of extremal dependence is the χ statistic (Padoan, 2011). The χ statistic for the upper tail is given by $\lim_{c \rightarrow \infty} \Pr(Y(\mathbf{s}_1) > c | Y(\mathbf{s}_2) > c)$. For a stationary spatial process, we can write the χ coefficient as

$$\chi(h) = \lim_{c \rightarrow \infty} \Pr(Y(\mathbf{s}) > c | Y(\mathbf{t}) > c). \quad (4)$$

If $\chi(h) = 0$, then observations are asymptotically independent at distance h . For Gaussian processes, $\chi(h) = 0$ regardless of the distance, so they are not suitable for modeling spatially-dependent extremes. Unlike the Gaussian process, the skew- t process is asymptotically dependent. However, one problem with the spatial skew- t process is that $\lim_{h \rightarrow \infty} \chi(h) > 0$. This occurs because all observations, both near and far, share the same z and σ terms. Therefore, the skew- t process is not ideal for spatial analysis of large geographic regions where we expect only local spatial dependence. The explicit expression for $\chi(h)$ (Padoan, 2011) and a proof of this are given in Appendix A.4.

3 Spatiotemporal skew- t model for extremes

In this section, we propose extensions to the skew- t process to model spatial extremes over a large geographic region by introducing censoring to focus on tail behavior and a random partition similar to Kim et al. (2005) to remove long-range asymptotic dependence.

3.1 Censoring to focus on the tails

To avoid bias in estimating tail parameters, we model censored data. Let

$$\tilde{Y}(\mathbf{s}) = \begin{cases} Y(\mathbf{s}) & \delta(\mathbf{s}) = 1 \\ T & \delta(\mathbf{s}) = 0 \end{cases} \quad (5)$$

be the censored observation at site \mathbf{s} where $Y(\mathbf{s})$ is the uncensored observation, $\delta(\mathbf{s}) = I[Y(\mathbf{s}) > T]$, and T is a pre-specified threshold value. Then, assuming the uncensored data $Y(\mathbf{s})$ are observations from a skew- t process, we update values censored below the threshold using standard Bayesian missing data methods as described in Section 4.

3.2 Partitioning to remove long-range asymptotic dependence

We handle the problem of long-range asymptotic dependence with a random partition model. As discussed in Section 2, the source of long-range dependence is the shared z and σ . Therefore, to alleviate this dependence, we allow z and σ to vary by site. Then, the model becomes

$$Y(\mathbf{s}) = \mathbf{X}(\mathbf{s})^T \boldsymbol{\beta} + \lambda \sigma(\mathbf{s}) |z(\mathbf{s})| + \sigma(\mathbf{s}) v(\mathbf{s}). \quad (6)$$

Let $\mathbf{w} = (w_1, w_2)$ be the location of a spatial knot. To model spatial variation, consider a set of spatial knots $\mathbf{w}_1, \dots, \mathbf{w}_K$ from a homogeneous Poisson process with intensity μ over spatial domain $\mathcal{D} \in \mathcal{R}^2$. The knots define a random daily partition of \mathcal{D} by subregions P_1, \dots, P_K defined as

$$P_k = \{\mathbf{s} : k = \arg \min_{\ell} \|\mathbf{s} - \mathbf{w}_{\ell}\|\}. \quad (7)$$

All $z(\mathbf{s})$ and $\sigma(\mathbf{s})$ for sites in subregion k are assigned common values

$$z(\mathbf{s}) = z_k \quad (8)$$

$$\sigma(\mathbf{s}) = \sigma_k, \quad (9)$$

and the z_k and σ_k^2 are distributed as $z_k \stackrel{iid}{\sim} N(0, 1)$ and $\sigma^2 \stackrel{iid}{\sim} \text{IG}(a, b)$. So, within each partition, $Y(\mathbf{s})$ follows the spatial skew- t process defined in Section 2. Across partitions, the $Y(\mathbf{s})$ remain correlated via the correlation function for $v(\mathbf{s})$ because it spans the partitions.

When incorporating the random daily partition, conditional on knots $\mathbf{w}_1, \dots, \mathbf{w}_K$, the χ statistic for two sites \mathbf{s} and \mathbf{t} in partitions k_s and k_t respectively is

$$\begin{aligned} \chi(h) &= I(k_s = k_t) \chi_{\text{skew-}t}(h) + I(k_s \neq k_t) \chi_{\text{Gaus}}(h) \\ &= I(k_s = k_t) \chi_{\text{skew-}t}(h) \end{aligned} \quad (10)$$

where $\chi_{\text{skew-}t}(h)$ is the χ statistic for a skew- t process, $\chi_{\text{Gaus}}(h)$ is the χ statistic for a Gaussian process, and $h = \|\mathbf{s} - \mathbf{t}\|$. Therefore, sites in different subregions are asymptotically independent. Marginally, over the knots $\mathbf{w}_1, \dots, \mathbf{w}_K$, $\chi(h) = \pi(h) \chi_{\text{skew-}t}(h)$, where $\pi(h) = \Pr(k_s = k_t)$ is the probability that two sites separated by distance h are in the same partition. So, to show that $\lim_{h \rightarrow \infty} \chi(h) = 0$, we need only know that $\lim_{h \rightarrow \infty} \pi(h) = 0$. A proof of this is given in Appendix A.3.

Figure 1 shows $\chi(h)$ for varying data settings when $\mathbf{s} \in [0, 1] \times [0, 1]$. The plot clearly shows that without partitioning, in a skew- t process, all sites remain asymptotically dependent. However, upon incorporating partitioning into the model, the asymptotic dependence is only shared by sites that are close to one another.

Figure 1: $\chi(h)$ for different settings. I need to replace the plot here.

3.3 Extension to space-time data

When using daily measurements, the assumption of temporal independence is inappropriate. There are several places where temporal dependence could be incorporated in the model, including the residual $v_t(\mathbf{s})$. However, we choose to allow for temporal dependence in the \mathbf{w} , z , and σ terms because these terms dictate the tail behavior which is our primary focus. In this section, we extend (6) to the spatiotemporal setting. Let

$$Y_t(\mathbf{s}) = \mathbf{X}_t(\mathbf{s})^T \boldsymbol{\beta} + \lambda \sigma_t(\mathbf{s}) |z_t(\mathbf{s})| + \sigma_t(\mathbf{s}) v_t(\mathbf{s}), \quad (11)$$

where $t \in \{1, \dots, T\}$ denotes the day of each observation. Let $\mathbf{w}_{tk} = (w_{tk1}, w_{tk2})$ be a spatial knot on day t , and let w_{t1}, \dots, w_{tK} be a collection of spatial knots on day t . As in section 3.2, these knots define a daily partition P_{t1}, \dots, P_{tK} , and for $\mathbf{s} \in P_{tk}$,

$$z_t(\mathbf{s}) = z_{tk} \quad (12)$$

$$\sigma_t(\mathbf{s}) = \sigma_{tk}. \quad (13)$$

We use an AR(1) time series model for w_{tk} , z_{tk} , and σ_{tk} . The time series model must be specified after a transformation to preserve the skew- t process at each time point. We first transform the spatial knots from \mathcal{D} to \mathcal{R}^2 as follows. Let

$$w_{tki}^* = \Phi^{-1} \left[\frac{(w_{tki} - \min(\mathbf{s}_i))}{\text{range}(\mathbf{s}_i)} \right], \quad i = 1, 2 \quad (14)$$

where Φ is a univariate standard normal density function, and $\mathbf{s}_i = [s_{1i}, \dots, s_{ni}]$. Then $\mathbf{w}_{tk}^* \in \mathcal{R}^2$. We use a copula on the $\sigma_t^2(\mathbf{s})$ to ensure that the marginal distributions of $\sigma_t^2(\mathbf{s})$ are inverse gamma. Let

$$\sigma_t^{2*}(\mathbf{s}) = \Phi^{-1} \{ \text{IG}[\sigma_t^2(\mathbf{s})] \} \quad (15)$$

where IG is the distribution function for an $\text{IG}(a, b)$ random variable. The AR(1) process for each tail parameter is $\mathbf{w}_{1k}^* \sim N_w(0, 1)$, $z_{1k} \sim N(0, \sigma_{1k}^2)$, $\sigma_{1k}^{2*} \sim N(0, 1)$, and for $t > 1$ the time series is modeled as

$$\mathbf{w}_{tk}^* | \mathbf{w}_{t-1,k}^* \sim N_2 [\phi_w \mathbf{w}_{t-1,k}^*, (1 - \phi_w^2)] \quad (16)$$

$$z_{tk} | z_{t-1,k} \sim N [\phi_z z_{t-1,k}, \sigma_{tk}^2 (1 - \phi_z^2)] \quad (17)$$

$$\sigma_{tk}^{2*} | \sigma_{t-1,k}^{2*} \sim N [\phi_\sigma \sigma_{t-1,k}^{2*}, (1 - \phi_\sigma^2)] \quad (18)$$

where $|\phi_w| < 1$, $0 < \phi_z < 1$, and $|\phi_\sigma| < 1$. We restrict ϕ_z to be positive because in the model, we assume $Y(\mathbf{s})$ is a function of $|z_t(\mathbf{s})|$. These are stationary time series models with marginal distributions $\mathbf{w}_k^* \sim N_2(0, 1)$, $z_k \sim N(0, \sigma_k^2)$, $\sigma_k^{2*} \sim N(0, 1)$. After transformation back to the original space, $\mathbf{w}_{tk} \sim \text{Unif}(\mathcal{D})$, $\sigma_{tk}^2 \sim \text{IG}(a, b)$. For each day, the model is identical to the spatial-only model in (6) by construction.

3.4 Hierarchical model

Conditioned on $z_{tk}(\mathbf{s})$, $\sigma_{tk}^2(\mathbf{s})$, and P_{tk} , the marginal distributions are Gaussian and the joint distribution multivariate Gaussian. However, we do not fix the partitions, they are treated as unknown and updated in the MCMC. We model this with a Bayesian hierarchical model as follows. Let $\mathbf{w}_{t1}, \dots, \mathbf{w}_{tK}$ be a set of daily spatial knots in a spatial domain of interest, \mathcal{D} , and P_{tk} as defined in (7).

123

Then

$$Y_t(\mathbf{s}) \mid z_t(\mathbf{s}), \sigma^2(\mathbf{s}), P_{tk}, \alpha, \beta, \Theta = \mathbf{X}_t(\mathbf{s})^T \beta + \lambda |z_t(\mathbf{s})| + \sigma_t(\mathbf{s}) v_t(\mathbf{s}) \quad (19)$$

$$z_t(\mathbf{s}) = z_{tk} \text{ if } \mathbf{s} \in P_{tk} \quad (20)$$

$$\sigma_t^2(\mathbf{s}) = \sigma_{tk}^2 \text{ if } \mathbf{s} \in P_{tk} \quad (21)$$

$$\lambda \sim N(0, 10) \quad (22)$$

$$v_t(\mathbf{s}) \mid \Theta \sim \text{Matérn}(0, \Sigma) \quad (23)$$

$$z_{tk} \mid z_{t-1,k}, \sigma_{tk}^2 \sim N(\phi_z z_{t-1,k}, \sigma_{tk}^2 (1 - \phi_z^2)) \quad (24)$$

$$\sigma_{tk}^{2*} \mid \sigma_{t-1,k}^{2*} \sim N(\phi_\sigma \sigma_{t-1,k}^{2*}, (1 - \phi_\sigma^2)) \quad (25)$$

$$\mathbf{w}_{tk}^* \mid \mathbf{w}_{t-1,k}^* \sim N_2(\phi_w \mathbf{w}_{t-1,k}^*, (1 - \phi_w^2)) \quad (26)$$

124 where $\Theta = \{\rho, \nu, \gamma\}$, and Σ is a Matérn covariance matrix as described in Section 2.1.

125 4 Computation

126 First, we impute values below the threshold. Then, we update Θ using Metropolis Hastings or Gibbs sam-
 127 pling when appropriate. Finally, we make spatial predictions using conditional multivariate normal results
 128 and the fact that the distribution of $Y_t(\mathbf{s}) \mid \Theta, z(\mathbf{s})$ is the usual multivariate normal distribution with a
 129 Matérn spatial covariance structure.

130 We can use Gibbs sampling to update $Y_t(\mathbf{s})$ for censored observations that are below the threshold T .
 131 After conditioning on $\lambda, z_t(\mathbf{s})$ and non-censored observations, $Y_t(\mathbf{s})$ has truncated normal full conditionals.
 132 So we sample $Y_t(\mathbf{s}) \sim N_{(-\infty, T)}(\mathbf{X}\beta, \Sigma)$. After imputing the censored observations, we update the model
 133 parameters. To update the model parameters, we use standard Gibbs updates for parameters when possible.
 134 In the case Gibbs sampling is not possible, parameters are updated using a random-walk Metropolis Hastings
 135 algorithm. See Appendices A.1 and A.2 for details regarding the MCMC. The final step of the computation
 136 is to use Bayesian Kriging to generate a predictive distribution for $Y_t(\mathbf{s}^*)$ at prediction location \mathbf{s}^* . This
 137 step is similar to the imputation for censored observations except that the full conditionals are no longer
 138 truncated at T .

139 5 Simulation study

140 In this section, we conduct a simulation study to investigate how the number of partitions and the level of
 141 thresholding impact the accuracy of predictions made by the model.

142 5.1 Design

143 For all simulation designs, we generate data from the model in Section 3.2 using $n_s = 144$ sites and
 144 $n_t = 50$ independent days. The sites are generated Uniform($[0, 10] \times [0, 10]$). We generate data from 6
 145 different simulation designs:

- 146 1. Gaussian marginal, $K = 1$ knot
- 147 2. T marginal, $K = 1$ knot
- 148 3. T marginal, $K = 5$ knots
- 149 4. Skew- t marginal, $K = 1$ knots
- 150 5. Skew- t marginal, $K = 5$ knots

6. Max-stable.

In the first five designs, the $v_t(\mathbf{s})$ terms are generated using a Matérn covariance with smoothness parameter $\nu = 0.5$ and spatial range $\rho = 0.1$. For the covariance matrices in designs 1 – 5, the proportion of the variance accounted for by the spatial variation is $\gamma = 0.9$ while the proportion of the variance accounted for by the nugget effect is 0.1. In the first design, $\sigma^2 = 2$ is used for all days. For designs 2 – 4, $\sigma_{tk}^2 \stackrel{iid}{\sim} \text{IG}(3, 8)$. For designs 1 – 3, we set $\lambda = 0$. For designs four and five, $\lambda = 3$ was used, and the z_t are generated as described in (8). In the sixth design, we generate from a spatial max-stable distribution (Reich and Shaby, 2012) with parameters $\mu = 1, \sigma = 1, \xi = 0.2$ and 144 spatial knots on a regular lattice in the square $[1, 9] \times [1, 9]$. **I know I need more here about the max-stable method** In all six designs, the mean $\mathbf{X}\beta = 10$ is assumed to be constant across space.

$M = 50$ data sets are generated for each design. For each data set we fit the data using

1. Gaussian marginal, $K = 1$ knots
2. Skew- t marginal, $K = 1$ knots, $T = -\infty$
3. T marginal, $K = 1$ knots, $T = q(0.80)$
4. Skew- t marginal, $K = 5$ knots, $T = -\infty$
5. T marginal, $K = 5$ knots, $T = q(0.80)$

where $q(0.80)$ is the 80th sample quantile of the data. The design matrix \mathbf{X} includes an the intercept with a prior of $\beta \sim \text{N}(0, 10)$. The spatial covariance parameters have priors $\log(\nu) \sim \text{N}(-1.2, 1)$, $\gamma \sim \text{U}(0, 1)$, $\log(\rho) \sim \text{N}(-2, 1)$. The skewness parameter has prior $\lambda \sim \text{N}(0, 2)$. The residual variance terms have priors $\sigma_t^2(\mathbf{s}) \sim \text{IG}(0.1, 0.1)$. The knots have priors $\mathbf{w} \sim \text{Unif}(\mathcal{D})$. We do not fit the data using the max-stable methods from Reich and Shaby (2012) because of the time it takes.

5.2 Cross validation

Models were compared using cross validation with 100 sites used as training sites and 30 sites withheld for testing. The model was fit using the training set, and predictions were generated at the testing site locations. Because one of the primary goals of this model is to predict extreme events, we use Brier scores to select the model that best fits the data (Gneiting and Raftery, 2007). The Brier score for predicting exceedance of a threshold c is given by $[e(c) - P(c)]^2$ where $e(c) = I[y > c]$ is an indicator function indicating that a test set value, y , has exceeded the threshold, c , and $P(c)$ is the predicted probability of exceeding c . We average the Brier scores over all test sites and days. For the Brier score, a lower score indicates a better fit.

5.3 Results

We compared the Brier scores for exceeding 11 different thresholds for each dataset. The thresholds used for the Brier scores are extreme quantiles from the simulated data. Figure 2 gives the Brier score relative to the Brier score for the Gaussian method calculated as

$$\text{BS}_{\text{rel}} = \frac{\text{BS}_{\text{method}}}{\text{BS}_{\text{Gaussian}}}.$$

For data that are relatively symmetric, the standard Gaussian method outperforms the other methods. However, when the data come from a skewed distribution (settings 3 – 6), the partitioned and thresholded methods outperform the Gaussian method. Additionally, for both of the skew- t settings ($K = 1, K = 5$), the skew- t method with the appropriate number of partitions performs the best. Finally, the simulation results




Figure 2: Brier scores relative to the Gaussian method for simulation study results. A ratio lower than 1 indicates that the method outperforms relative to the Gaussian method.

suggest that for data from a skew- t distribution, thresholded methods provide some improvement over non-thresholded models for the extreme thresholds. With the exception of the Brier score for $c = q(0.995)$ in setting 6, the improvement over the Gaussian method is statistically significant using a Wilcoxon signed rank test ($\alpha = 0.05$).

6 Data analysis

To illustrate this method, we consider the daily maximum 8-hour ozone measurements for July 2005 at 735 Air Quality System (AQS) monitoring sites in the eastern United States as the response (see Figure 3). For each site, we also have covariate information containing the estimated ozone from the Community Multi-scale Air Quality (CMAQ) modeling system. Initially, we fit a linear regression assuming a mean function of

$$\mathbf{X}\boldsymbol{\beta} = \beta_0 + \beta_1 \cdot \text{CMAQ}_t(\mathbf{s}). \quad (27)$$




Figure 3: Ozone values on 10 July 2005 (left) Q-Q plot of the residuals (right)

The data from July 10 are shown in Figure 3 along with a Q-Q plot of the residuals compared to a skew- t distribution with 8 d.f. and $\alpha = 1$.

The $\hat{\chi}(h)$ and $\hat{\chi}(t)$ plots estimate the spatial and temporal dependence using sample data. For $\hat{\chi}(h)$, we take the sample proportion of all pairs of sites within distance h of one another that are above a threshold T . I know I need another sentence or two here about $\hat{\chi}(t)$ indicate that the residuals still demonstrate spatial and temporal dependence (see Figure 4)

6.1 Model comparisons

We fit the model using Gaussian and skew- t marginal distributions, $K = 1, 5, 10, 15$ partitions, with $Y(s)$ censored at $T = 0, 50, 75, 90$ ppb as described in Section ???. We also include a max-stable analysis using the method by ????. All methods assume the mean function given in (27). For each model, Brier scores and quantile scores were averaged over all sites and days to obtain a single quantile score for each dataset. At a particular threshold or quantile level, the model that fits the best is the one with the lowest score. Looking at the heatplot in Figure 5, we can see that the methods using no thresholding tend to outperform those methods that do use thresholding.




Figure 4: $\hat{\chi}(h)$ plot for the residuals. The vertical line indicates the distance after which observations no longer demonstrate dependence (left) $\hat{\chi}(t)$ plot for the residuals (right)




Figure 5: Heatplot of quantile scores for the 99th quantile (left) Quantile scores of Gaussian, skew- t ($K=10$, $T=q(0)$), and Max-stable (right)

6.2 Results

7 Conclusions

Acknowledgments

A Appendices

A.1 MCMC details

The MCMC sampling for the model 3.4 is done using R (<http://www.r-project.org>). Whenever possible, we select conjugate priors (see Appendix A.2); however, for some of the parameters, no conjugate prior distributions exist. When no conjugate prior distribution exists, we use a random walk Metropolis Hastings update step. In each Metropolis Hastings update, we tune the algorithm to give acceptance rates near 0.40.

Spatial knot locations

For each day, we update the spatial knot locations, $\mathbf{w}_1, \dots, \mathbf{w}_K$, using a Metropolis Hastings block update. Because the spatial domain is bounded, we generate candidate knots using the transformed knots $\mathbf{w}_1^*, \dots, \mathbf{w}_K^*$ (see section 3.3) and a random walk bivariate Gaussian candidate distribution

$$\mathbf{w}_k^{*(c)} \sim N(\mathbf{w}_k^{*(r-1)}, s^2 I_2)$$

where $\mathbf{w}_k^{*(r-1)}$ is the location for the transformed knot at MCMC iteration $r - 1$, s is a tuning parameter, and I_2 is an identity matrix. After candidates have been generated for all K knots, the acceptance ratio is

$$R = \left\{ \frac{l[Y_t(\mathbf{s}|\mathbf{w}_1^{(c)}, \dots, \mathbf{w}_K^{(c)}, \dots)]}{l[Y_t(\mathbf{s}|\mathbf{w}_1^{(r-1)}, \dots, \mathbf{w}_K^{(r-1)}, \dots)]} \right\} \times \left\{ \frac{\prod_{k=1}^K \phi(\mathbf{w}_k^{(c)})}{\prod_{k=1}^K \phi(\mathbf{w}_k^{(r-1)})} \right\} \times \left\{ \frac{\prod_{k=1}^K p(\mathbf{w}_k^{*(c)})}{\prod_{k=1}^K p(\mathbf{w}_k^{*(r-1)})} \right\}$$

where l is the likelihood given in (19), and $p(\cdot)$ is the prior either taken from the time series given in (3.3) or assumed to be uniform over \mathcal{D} . The candidate knots are accepted with probability $\min\{R, 1\}$.

Spatial random effects

If there is temporal dependence amongst the observations, then we update z_{tk} using a Metropolis Hastings update. First, we generate a candidate using a random walk Gaussian candidate distribution

$$z_{tk}^{(c)} \sim N(z_{tk}^{(r-1)}, s^2)$$

where $z_{tk}^{(r-1)}$ is the value at MCMC iteration $r - 1$, and s is a tuning parameter. The acceptance ratio is

$$R = \left\{ \frac{l[Y_t(\mathbf{s})|z_{tk}^{(c)}, \dots]}{l[Y_t(\mathbf{s})|z_{tk}^{(r-1)}, \dots]} \right\} \times \left\{ \frac{p[z_{tk}^{(c)}]}{p[z_{tk}^{(r-1)}]} \right\}.$$

The candidate is accepted with probability $\min\{R, 1\}$.

Variance terms

When there is more than one site in a partition, then we update σ_{tk}^2 using a Metropolis Hastings update. First, we generate a candidate for σ_{tk}^2 using an $\text{IG}(a^*/s, b^*/s)$ candidate distribution in an independence Metropolis Hastings update where $a^* = (n_{tk} + 1)/2 + a$, $b^* = [Y_{tk}^T \Sigma_{tk}^{-1} Y_{tk} + z_{tk}^2]/2 + b$, n_{tk} is the number of sites in partition k on day t , and Y_{tk} and Σ_{tk}^{-1} are the observations and precision matrix for partition k on day t . The acceptance ratio is

$$R = \left\{ \frac{l[Y_t(\mathbf{s}) | \sigma_{tk}^{2(c)}, \dots]}{l[Y_t(\mathbf{s}) | \sigma_{tk}^{2(r-1)}]} \right\} \times \left\{ \frac{l[z_{tk} | \sigma_{tk}^{2(c)}, \dots]}{l[z_{tk} | \sigma_{tk}^{2(r-1)}, \dots]} \right\} \times \left\{ \frac{p[\sigma_{tk}^{2(c)}]}{p[\sigma_{tk}^{2(r-1)}]} \right\} \times \left\{ \frac{c[\sigma_{tk}^{2(r-1)}]}{c[\sigma_{tk}^{2(c)}]} \right\}$$

where $p[\cdot]$ is the prior either taken from the time series given in (3.3) or assumed to be $\text{IG}(a, b)$, and $c[\cdot]$ is the candidate distribution. The candidate is accepted with probability $\min\{R, 1\}$.

Spatial covariance parameters

We update the three spatial covariance parameters, $\log(\rho)$, $\log(\nu)$, γ , using a Metropolis Hastings block update step. First, we generate a candidate using a random walk Gaussian candidate distribution

$$\log(\rho)^{(c)} \sim \text{N}(\log(\rho)^{(r-1)}, s^2)$$

where $\log(\rho)^{(r-1)}$ is the value at MCMC iteration $r - 1$, and s is a tuning parameter. Candidates are generated for $\log(\nu)$ and γ in a similar fashion. The acceptance ratio is

$$R = \left\{ \frac{\prod_{t=1}^T l[Y_t(\mathbf{s}) | \rho^{(c)}, \nu^{(c)}, \gamma^{(c)}, \dots]}{\prod_{t=1}^T l[Y_t(\mathbf{s}) | \rho^{(r-1)}, \nu^{(r-1)}, \gamma^{(r-1)}, \dots]} \right\} \times \left\{ \frac{p[\rho^{(c)}]}{p[\rho^{(r-1)}]} \right\} \times \left\{ \frac{p[\nu^{(c)}]}{p[\nu^{(r-1)}]} \right\} \times \left\{ \frac{p[\gamma^{(c)}]}{p[\gamma^{(r-1)}]} \right\}.$$

All three candidates are accepted with probability $\min\{R, 1\}$.

A.2 Posterior distributions

Conditional posterior of $z_{tk} \mid \dots$

If knots are independent over days, then the conditional posterior distribution of $|z_{tk}|$ is conjugate. For simplicity, drop the subscript t , let $z_{tk}^* = |z_{tk}|$, and define

$$R(\mathbf{s}) = \begin{cases} Y(\mathbf{s}) - X(\mathbf{s})\beta & s \in P_l \\ Y(\mathbf{s}) - X(\mathbf{s})\beta - \alpha z(\mathbf{s}) & s \notin P_l \end{cases}$$

Let

$$R_1 = \text{the vector of } R(\mathbf{s}) \text{ for } s \in P_l$$

$$R_2 = \text{the vector of } R(\mathbf{s}) \text{ for } s \notin P_l$$

$$\Omega = \Sigma^{-1}.$$

Then

$$\begin{aligned} \pi(z_l | \dots) &\propto \exp \left\{ -\frac{1}{2} \left[\begin{pmatrix} R_1 - \alpha z_l^* \mathbf{1} \\ R_2 \end{pmatrix}^T \begin{pmatrix} \Omega_{11} & \Omega_{12} \\ \Omega_{21} & \Omega_{22} \end{pmatrix} \begin{pmatrix} R_1 - \alpha z_l^* \mathbf{1} \\ R_2 \end{pmatrix} + \frac{z_l^{*2}}{\sigma_l^2} \right] \right\} I(z_l > 0) \\ &\propto \exp \left\{ -\frac{1}{2} [\Lambda_l z_l^{*2} - 2\mu_l z_l^*] \right\} \end{aligned}$$

256 where

$$\begin{aligned}\mu_l &= \alpha(R_1^T \Omega_{11} + R_2^T \Omega_{21})\mathbf{1} \\ \Lambda_l &= \alpha^2 \mathbf{1}^T \Omega_{11} \mathbf{1} + \frac{1}{\sigma_l^2}.\end{aligned}$$

257 Then $Z_l^* | \dots \sim N_{(0,\infty)}(\Lambda_l^{-1} \mu_l, \Lambda_l^{-1})$

258 **Conditional posterior of $\beta | \dots$**

259 Let $\beta \sim N_p(0, \Lambda_0)$ where Λ_0 is a precision matrix. Then

$$\begin{aligned}\pi(\beta | \dots) &\propto \exp \left\{ -\frac{1}{2} \beta^T \Lambda_0 \beta - \frac{1}{2} \sum_{t=1}^T [\mathbf{Y}_t - X_t \beta - \alpha |z_t|]^T \Omega [\mathbf{Y}_t - X_t \beta - \alpha |z_t|] \right\} \\ &\propto \exp \left\{ -\frac{1}{2} \left[\beta^T \Lambda_\beta \beta - 2 \sum_{t=1}^T [\beta^T X_t^T \Omega (\mathbf{Y}_t - \alpha |z_t|)] \right] \right\} \\ &\propto N(\Lambda_\beta^{-1} \mu_\beta, \Lambda_\beta^{-1})\end{aligned}$$

260 where

$$\begin{aligned}\mu_\beta &= \sum_{t=1}^T [X_t^T \Omega (\mathbf{Y}_t - \alpha |z_t|)] \\ \Lambda_\beta &= \Lambda_0 + \sum_{t=1}^T X_t^T \Omega X_t.\end{aligned}$$

261 **Conditional posterior of $\sigma^2 | \dots$**

262 In the case where $L = 1$ and temporal dependence is negligible, then σ^2 has a conjugate posterior distribu-
263 tion. Let $\sigma_t^2 \stackrel{iid}{\sim} \text{IG}(\alpha_0, \beta_0)$. For simplicity, drop the subscript t . Then

$$\begin{aligned}\pi(\sigma^2 | \dots) &\propto (\sigma^2)^{-\alpha_0 - 1/2 - n/2 - 1} \exp \left\{ -\frac{\beta_0}{\sigma^2} - \frac{|z|^2}{2\sigma^2} - \frac{(\mathbf{Y} - \boldsymbol{\mu})^T \Sigma^{-1} (\mathbf{Y} - \boldsymbol{\mu})}{2\sigma^2} \right\} \\ &\propto (\sigma^2)^{-\alpha_0 - 1/2 - n/2 - 1} \exp \left\{ -\frac{1}{\sigma^2} \left[\beta_0 + \frac{|z|^2}{2} + \frac{1}{2} (\mathbf{Y} - \boldsymbol{\mu})^T \Sigma^{-1} (\mathbf{Y} - \boldsymbol{\mu}) \right] \right\} \\ &\propto \text{IG}(\alpha^*, \beta^*)\end{aligned}$$

264 where

$$\begin{aligned}\alpha^* &= \alpha_0 + \frac{1}{2} + \frac{n}{2} \\ \beta^* &= \beta_0 + \frac{|z|^2}{2} + \frac{1}{2} (\mathbf{Y} - \boldsymbol{\mu})^T \Sigma^{-1} (\mathbf{Y} - \boldsymbol{\mu}).\end{aligned}$$

265 In the case that $L > 1$, a random walk Metropolis Hastings step will be used to update σ_{lt}^2 .

266 **Conditional posterior of $\lambda \mid \dots$**

267 Let $\lambda \sim N(0, \tau_\lambda)$ where τ_λ is a precision term. Then

$$\begin{aligned}\pi(\lambda \mid \dots) &\propto \exp \left\{ -\frac{1}{2} \tau_\lambda \lambda^2 + \sum_{t=1}^T \frac{1}{2} [\mathbf{Y}_t - X_t \beta - \lambda |z_t|]^T \Omega [\mathbf{Y}_t - X_t \beta - \lambda |z_t|] \right\} \\ &\propto \exp \left\{ -\frac{1}{2} [\lambda^2 (\tau_\lambda + \sum_{t=1}^T |z_t|^T \Omega |z_t|^T) - 2\lambda \sum_{t=1}^T |z_t|^T \Omega (\mathbf{Y}_t - X_t \beta)] \right\} \\ &\propto \exp \left\{ -\frac{1}{2} [\tau_\lambda^* \lambda^2 - 2\mu_\lambda] \right\}\end{aligned}$$

268 where

$$\begin{aligned}\mu_\lambda &= \sum_{t=1}^T |z_t|^T \Omega (\mathbf{Y}_t - X_t \beta) \\ \tau_\lambda^* &= t_\alpha + \sum_{t=1}^T |z_t|^T \Omega |z_t|.\end{aligned}$$

269 Then $\lambda \mid \dots \sim N(\tau_\lambda^{*-1} \mu_\lambda, \tau_\lambda^{*-1})$

270 **A.3 Proof that $\lim_{h \rightarrow \infty} \pi(h) = 0$**

271 Let $N(A)$ be the number of knots in A , the area between sites \mathbf{s}_1 and \mathbf{s}_2 . Consider a spatial Poisson process
272 with intensity $\mu(A)$. So,

$$P[N(A) = k] = \frac{\mu(A)^k \exp\{-\mu(A)\}}{k!}.$$

273 Then for any finite k , $\lim_{h \rightarrow \infty} P[N(A) = k] = 0$ because $\lim_{h \rightarrow \infty} \mu(A) = \infty$. With each additional knot
274 in A , the chance that \mathbf{s}_1 and \mathbf{s}_2 will be in the same partition will decrease, because partition membership
275 is defined by the closest knot to a site. Therefore, $\lim_{h \rightarrow \infty} \pi(h) = 0$.

276 **A.4 Skew- t distribution**

277 **Univariate extended skew- t distribution**

278 We say that Y follow a univariate extended skew- t distribution with location $\xi \in \mathcal{R}$, scale $\omega > 0$, skew
279 parameter $\alpha \in \mathcal{R}$, extended parameter $\tau \in \mathcal{R}$, and degrees of freedom ν if has distribution function

$$f_{\text{EST}}(y) = \omega^{-1} \frac{f_T(z; \nu)}{F_T(\tau / \sqrt{1 + \alpha^2}; \nu)} F_T \left[(\alpha z + \tau) \sqrt{\frac{\nu + 1}{\nu + z^2}}; 0, 1, \nu + 1 \right] \quad (28)$$

280 where $f_T(t; \nu)$ is a univariate Student's t with ν degrees of freedom, $F_T(t; \nu) = P(T < t)$, and $z = (y - \xi)/\omega$.
281 In the case that $\tau = 0$, then Y follows a univariate skew- t distribution.

Multivariate skew- t distribution

If $\mathbf{Z} \sim \text{ST}_d(0, \bar{\boldsymbol{\Omega}}, \boldsymbol{\alpha}, \eta)$ is a d -dimensional skew- t distribution, and $\mathbf{Y} = \boldsymbol{\xi} + \boldsymbol{\omega}\mathbf{Z}$, where $\boldsymbol{\omega} = \text{diag}(\omega_1, \dots, \omega_d)$, then the density of Y at y is

$$f_y(\mathbf{y}) = \det(\boldsymbol{\omega})^{-1} f_z(\mathbf{z}) \quad (29)$$

where

$$f_z(\mathbf{z}) = 2t_d(\mathbf{z}; \bar{\boldsymbol{\Omega}}, \eta) T \left[\boldsymbol{\alpha}^T \mathbf{z} \sqrt{\frac{\eta + d}{\nu + Q(\mathbf{z})}}; \eta + d \right] \quad (30)$$

$$\mathbf{z} = \boldsymbol{\omega}^{-1}(\mathbf{y} - \boldsymbol{\xi}) \quad (31)$$

where $t_d(\mathbf{z}; \bar{\boldsymbol{\Omega}}, \eta)$ is a d -dimensional Student's t -distribution with scale matrix $\bar{\boldsymbol{\Omega}}$ and degrees of freedom η , $Q(\mathbf{z}) = \mathbf{z}^T \bar{\boldsymbol{\Omega}}^{-1} \mathbf{z}$ and $T(\cdot; \eta)$ denotes the univariate Student's t distribution function with η degrees of freedom (Azzalini and Capitanio, 2013).

Extremal dependence

For a bivariate skew- t random variable $\mathbf{Y} = [Y(\mathbf{s}), Y(\mathbf{t})]^T$, the $\chi(h)$ statistic (Padoan, 2011) is given by

$$\chi(h) = \bar{F}_{\text{EST}} \left\{ \frac{[x_1^{1/\eta} - \varrho(h)]\sqrt{\eta+1}}{\sqrt{1-\varrho(h)^2}}; 0, 1, \alpha_1, \tau_1, \eta+1 \right\} + \bar{F}_{\text{EST}} \left\{ \frac{[x_2^{1/\eta} - \varrho(h)]\sqrt{\eta+1}}{\sqrt{1-\varrho(h)^2}}; 0, 1, \alpha_2, \tau_2, \eta+1 \right\}, \quad (32)$$

where \bar{F}_{EST} is the univariate survival extended skew- t function with zero location and unit scale, $\varrho(h) = \text{cor}(y_1, y_2)$, $\alpha_j = \alpha_i \sqrt{1-\varrho^2}$, $\tau_j = \sqrt{\eta+1}(\alpha_j + \alpha_i \varrho)$, and $x_j = F_T(\bar{\alpha}_i \sqrt{\eta+1}; 0, 1, \eta) / F_T(\bar{\alpha}_j \sqrt{\eta+1}; 0, 1, \eta)$ with $j = 1, 2$ and $i = 2, 1$ and where $\bar{\alpha}_j = (\alpha_j + \alpha_i \varrho) / \sqrt{1 + \alpha_i^2 [1 - \varrho(h)^2]}$.

Proof that $\lim_{h \rightarrow \infty} \chi(h) > 0$

Consider the bivariate distribution of $\mathbf{Y} = [Y(\mathbf{s}), Y(\mathbf{t})]^T$, with $\varrho(h)$ given by (2). So, $\lim_{h \rightarrow \infty} \varrho(h) = 0$. Then

$$\lim_{h \rightarrow \infty} \chi(h) = \bar{F}_{\text{EST}} \left\{ \sqrt{\eta+1}; 0, 1, \alpha_1, \tau_1, \eta+1 \right\} + \bar{F}_{\text{EST}} \left\{ \sqrt{\eta+1}; 0, 1, \alpha_2, \tau_2, \eta+1 \right\}. \quad (33)$$

Because the extended skew- t distribution is not bounded above, for all $\bar{F}_{\text{EST}}(x) = 1 - F_{\text{EST}} > 0$ for all $x < \infty$. Therefore, for a skew- t distribution, $\lim_{h \rightarrow \infty} \chi(h) > 0$.

References

- Azzalini, A. and Capitanio, A. (2013) *The Skew-Normal and Related Families*. Institute of Mathematical Statistics Monographs. Cambridge University Press.
- Blanchet, J. and Davison, A. C. (2011) Spatial modeling of extreme snow depth. *The Annals of Applied Statistics*, **5**, 1699–1725.

- 304 Coles, S. G. and Tawn, J. A. (1991) Modelling Extreme Multivariate Events. *Journal of the Royal Statistical*
305 *Society: Series B (Methodological)*, **53**, 377–392.
- 306 DuMouchel, W. H. (1983) Estimating the stable index α in order to measure tail thickness: a critique. *The*
307 *Annals of Statistics*, **11**, 1019–1031.
- 308 Genton, M. G. (2004) *Skew-Elliptical Distributions and Their Applications: A Journey Beyond Normality*.
309 Statistics (Chapman & Hall/CRC). Taylor & Francis.
- 310 Gneiting, T. and Raftery, A. E. (2007) Strictly Proper Scoring Rules, Prediction, and Estimation. *Journal of*
311 *the American Statistical Association*, **102**, 359–378.
- 312 Huser, R. (2013) *Statistical Modeling and Inference for Spatio-Temporal Extremes*. Ph.D. thesis.
- 313 Kim, H.-M., Mallick, B. K. and Holmes, C. C. (2005) Analyzing Nonstationary Spatial Data Using Piece-
314 wise Gaussian Processes. *Journal of the American Statistical Association*, **100**, 653–668.
- 315 Padoan, S. A. (2011) Multivariate extreme models based on underlying skew- and skew-normal distribu-
316 tions. *Journal of Multivariate Analysis*, **102**, 977–991.
- 317 Padoan, S. A., Ribatet, M. and Sisson, S. A. (2010) Likelihood-Based Inference for Max-Stable Processes.
318 *Journal of the American Statistical Association*, **105**, 263–277.
- 319 Reich, B. J. and Shaby, B. A. (2012) A hierarchical max-stable spatial model for extreme precipitation. *The*
320 *Annals of Applied Statistics*, **6**, 1430–1451.
- 321 Zhang, H. and El-Shaarawi, A. (2010) On spatial skewGaussian processes and applications. *Environmetrics*,
322 **21**, 33–47.