

## 1 (Azzalini, 1996)

Consider  $\mathbf{Y} \in \mathcal{R}^k$  that is distributed as MVN and is independent from  $Y_0 \sim N(0, 1)$ . Then for

$$\begin{pmatrix} Y_0 \\ \mathbf{Y} \end{pmatrix} \sim N_{k+1} \left\{ 0, \begin{pmatrix} 1 & 0 \\ 0 & \mathbf{\Psi} \end{pmatrix} \right\} \quad (1)$$

Then,  $Z_j = \delta_j |Y_0| + \sqrt{1 - \delta_j^2} Y_j, j = 1, \dots, k$  is skewed normal and its density is

$$f_k(z) = 2\phi_k(z; \mathbf{\Omega})\Phi(\mathbf{\alpha}^t \mathbf{z}) \quad \mathbf{z} \in \mathcal{R}^k \quad (2)$$

## 2 (Azzalini and Capitanio, 1999)

### 3 (Branco and Dey, 2001)

4 Modeling distributions that can account for skewness and heavy tails.

#### 5 Multivariate elliptical

6 Notation:  $\mathbf{X} \sim El_k(\boldsymbol{\mu}, \mathbf{\Sigma}; \phi)$  means that  $\mathbf{X} \in \mathcal{R}^k$  follows an elliptical distribution with location vector  
7  $\boldsymbol{\mu} \in \mathcal{R}^k$ , a dispersion matrix  $\mathbf{\Sigma} \in \mathcal{R}^{k \times k}$  and characteristic function  $\phi$ . If the density exists, then it is given  
8 by

$$f(\mathbf{x} | \boldsymbol{\mu}, \mathbf{\Sigma}) = |\mathbf{\Sigma}|^{-1/2} g^{(k)}[(\mathbf{x} - \boldsymbol{\mu})^T \mathbf{\Sigma}^{-1}(\mathbf{x} - \boldsymbol{\mu})] \quad (3)$$

#### 9 Multivariate skew elliptical

10 Notation:  $\mathbf{Y} \sim SE_k(\boldsymbol{\mu}, \mathbf{\Omega}, \boldsymbol{\delta}; \phi)$  means that  $\mathbf{Y} \in \mathcal{R}^k$  follows a skew-elliptical distribution with location  
11 vector  $\boldsymbol{\mu} \in \mathcal{R}^k$ , a dispersion matrix  $\mathbf{\Sigma} \in \mathcal{R}^{k \times k}$ , characteristic function  $\phi$  and skewness parameter  $\boldsymbol{\delta}$ . If the  
12 density exists, then it is given by

$$f_{\mathbf{Y}}(\mathbf{y}) = 2f_{g^{(k)}}(\mathbf{y})F_{g_q(\mathbf{y})}(\boldsymbol{\lambda}^T(\mathbf{y} - \boldsymbol{\mu})) \quad (4)$$

#### 13 Multivariate skew normal distribution

14 For a multivariate skew normal distribution, the density function is

$$f_{\mathbf{Y}}(\mathbf{y}) = 2\phi_k(\mathbf{y}; \boldsymbol{\mu}, \mathbf{\Omega})\Phi(\boldsymbol{\lambda}^T(\mathbf{y} - \boldsymbol{\mu})) \quad (5)$$

15 where

$$\boldsymbol{\lambda}^T = \frac{\boldsymbol{\delta}^T \mathbf{\Omega}^{-1}}{(1 - \boldsymbol{\delta}^T \mathbf{\Omega}^{-1} \boldsymbol{\delta})^{1/2}} \quad (6)$$

#### 16 Multivariate skew $t$ distribution

17 For a multivariate skew  $t$  distribution, the density function is

$$f_{\mathbf{Y}}(\mathbf{y}) = 2f_{\nu, \tau}(\mathbf{y}; \boldsymbol{\mu}; \mathbf{\Omega})F_{\nu^*, \tau^*}(\boldsymbol{\lambda}^T(\mathbf{y} - \boldsymbol{\mu})) \quad (7)$$

## Other densities mentioned

- Skew logistic
- Skew stable distribution
- Skew exponential power distribution
- Skew Pearson Type II distribution

(Sahu et al., 2003)

(Gupta et al., 2004)

p. 189 The general multivariate skew normal distribution has density function

$$2\phi_k(z; \Omega)\Phi(\alpha^T z) \quad (z \in \mathcal{R}^k) \quad (8)$$

where  $\phi_k(z; \Omega)$  is a  $k$ -dimensional process with mean zero, and correlation matrix  $\Omega$ ,  $\Phi(\cdot)$  is the  $N(0, 1)$  distribution, and  $\alpha \in \mathcal{R}^k$  is a shape term.

(Allard and Naveau, 2007)

(Zhang and ElShaarawi, 2010)

(Minozzo and Ferracuti, 2012)

p. 164 Let  $U_t \sim N(0, 1)$  and  $V_t(\mathbf{s}) \sim MVN$  with mean 0, and variance 1, then

$$\mathbf{Y}_t(\mathbf{s}) = \sigma\delta|U_t| + \sigma\sqrt{1 - \delta^2}V_t(\mathbf{s}) \quad (9)$$

follows a

## References

- Allard, D. and Naveau, P. (2007) A New Spatial Skew-Normal Random Field Model. *Communications in Statistics - Theory and Methods*, **36**, 1821–1834.  
URL<http://www.tandfonline.com/doi/abs/10.1080/03610920601126290>.
- Azzalini, A. (1996) The multivariate skew-normal distribution. *Biometrika*, **83**, 715–726.  
URL<http://biomet.oxfordjournals.org/content/83/4/715.short>  
<http://biomet.oupjournals.org/cgi/doi/10.1093/biomet/83.4.715>.
- Azzalini, a. and Capitanio, a. (1999) Statistical applications of the multivariate skew normal distribution. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, **61**, 579–602.  
URL<http://doi.wiley.com/10.1111/1467-9868.00194>.

- 43 Branco, M. D. and Dey, D. K. (2001) A General Class of Multivariate  
 44 Skew-Elliptical Distributions. *Journal of Multivariate Analysis*, **79**, 99–113.  
 45 URL<http://linkinghub.elsevier.com/retrieve/pii/S0047259X00919602>.
- 46 Gupta, A. K., González-Faras, G. and Domnguez-Molina, J. (2004) A multivari-  
 47 ate skew normal distribution. *Journal of Multivariate Analysis*, **89**, 181–190.  
 48 URL<http://linkinghub.elsevier.com/retrieve/pii/S0047259X03001313>.
- 49 Minozzo, M. and Ferracuti, L. (2012) On the existence of some skew-normal  
 50 stationary processes. *Chilean Journal of Statistics (ChJS)*, **3**, 157–170.  
 51 URL<http://chjs.deuv.cl/Vol13N2/ChJS-03-02-04.pdf>.
- 52 Sahu, S. K., Dey, D. K. and Branco, M. D. (2003) A new class of multivariate skew distributions  
 53 with applications to bayesian regression models. *Canadian Journal of Statistics*, **31**, 129–150.  
 54 URL<http://doi.wiley.com/10.2307/3316064>.
- 55 Zhang, H. and ElShaarawi, A. (2010) On spatial skewGaussian processes and applications. *Environmetrics*,  
 56 33–47. URL<http://onlinelibrary.wiley.com/doi/10.1002/env.982/abstract>.