

Genome-Wide Significant Linkage of Schizophrenia-Related Neuroanatomical Trait to 12q24

Emma Sprooten,^{1,2*} Cota Navin Gupta,³ Emma E.M. Knowles,^{1,2} D. Reese McKay,^{1,2} Samuel R. Mathias,^{1,2} Joanne E. Curran,^{4,5} Jack W. Kent Jr.,^{4,5} Melanie A. Carless,^{4,5} Marcio A. Almeida,^{4,5} Thomas D. Dyer,^{4,5} Harald H.H. Göring,^{4,5} Rene L. Olvera,⁶ Peter Kochunov,⁷ Peter T. Fox,⁸ Ravi Duggirala,^{4,5} Laura Almasy,^{4,5} Vince D. Calhoun,^{1,3,9,10} John Blangero,^{4,5} Jessica A. Turner,^{3,11} and David C. Glahn^{1,2}

¹Department of Psychiatry, Yale University School of Medicine, New Haven, Connecticut

²Olin Neuropsychiatry Research Center, Institute of Living, Hartford Hospital, Connecticut

³The Mind Research Network, Albuquerque, New Mexico

⁴Department of Genetics, Texas Biomedical Research Institute, San Antonio, Texas

⁵South Texas Diabetes and Obesity Institute, University of Texas Rio Grande Valley School of Medicine, Brownsville, Texas

⁶Department of Psychiatry, University of Texas Health Science Center San Antonio, San Antonio, Texas

⁷Department of Psychiatry, University of Maryland School of Medicine, Maryland Psychiatric Research Center, Baltimore, Maryland

⁸Research Imaging Institute, University of Texas Health Science Center San Antonio, San Antonio, Texas

⁹Department of Psychiatry, University of New Mexico, Albuquerque, New Mexico

¹⁰Department of Electrical and Computer Engineering, University of New Mexico, Albuquerque, New Mexico

¹¹Department of Psychology and Neuroscience Institute, Georgia State University, Atlanta, Georgia

Manuscript Received: 21 December 2014; Manuscript Accepted: 31 July 2015

The insula and medial prefrontal cortex (mPFC) share functional, histological, transcriptional, and developmental characteristics, and they serve higher cognitive functions of theoretical relevance to schizophrenia and related disorders. Meta-analyses and multivariate analysis of structural magnetic resonance imaging (MRI) scans indicate that gray matter density and volume reductions in schizophrenia are the most consistent and pronounced in a network primarily composed of the insula and mPFC. We used source-based morphometry, a multivariate technique optimized for structural MRI, in a large sample of randomly ascertained pedigrees (N = 887) to derive an insula–mPFC component and to investigate its genetic determinants. Firstly, we replicated the insula–mPFC gray matter component as an independent source of gray matter variation in the general population, and verified its relevance to schizophrenia in an

How to Cite this Article:

Sprooten E, Gupta CN, Knowles EE, McKay DR, Mathias SR, Curran JE, Kent Jr JW, Carless MA, Almeida MA, Dyer TD, Göring HH, Olvera RL, Kochunov P, Fox PT, Duggirala R, Almasy L, Calhoun VD, Blangero J, Turner JA, Glahn DC. 2015. Genome-Wide Significant Linkage of Schizophrenia-Related Neuroanatomical Trait to 12q24.

Am J Med Genet Part B 168B:678–686.

Conflict of interests: None

Grant sponsor: National Institute of Mental Health; Grant numbers: MH0708143 (Principal Investigator [PI]: DCG), MH078111 (PI: JB), MH083824 (PI: DCG & JB), 1R01MH094524 (NIMH, PIs: VC & JAT); Grant sponsor: COBRE; Grant number: P20GM103472 (PI:VC); Grant sponsor: SOLAR; Grant numbers: MH59490 (PI: JB), EB015611 (PI: PK); Grant sponsor: National Center for Research Resources; Grant numbers: C06 RR13556, C06 RR017515; Grant sponsor: AT&T; Grant sponsor: National Center for Research Resources; Grant number: S10 RR029392.

*Correspondence to:

Emma Sprooten, Ph.D., Olin Neuropsychiatry Research Center, Whitehall Research Building, Institute of Living, 200 Retreat Ave, Hartford, CT 06106. E-mail: emma.sprooten@yale.edu

Article first published online in Wiley Online Library (wileyonlinelibrary.com): 5 October 2015

DOI 10.1002/ajmg.b.32360

independent case-control sample. Secondly, we showed that the neuroanatomical variation defined by this component is largely determined by additive genetic variation ($h^2 = 0.59$), and genome-wide linkage analysis resulted in a significant linkage peak at 12q24 (LOD = 3.76). This region has been of significant interest to psychiatric genetics as it contains the Darier's disease locus and other proposed susceptibility genes (e.g., *DAO*, *NOS1*), and it has been linked to affective disorders and schizophrenia in multiple populations. Thus, in conjunction with previous clinical studies, our data imply that one or more psychiatric risk variants at 12q24 are co-inherited with reductions in mPFC and insula gray matter concentration.

© 2015 Wiley Periodicals, Inc.

Key words: extended pedigrees; magnetic resonance imaging; insula; medial prefrontal cortex; quantitative trait locus

INTRODUCTION

Schizophrenia is a heritable disorder [Sullivan et al., 2003] but the genetic variation accounting for its inheritance is complex and difficult to characterize. Many genetic markers, both common and rare, are thought to contribute to genetic risk for schizophrenia and related disorders [Gratten et al., 2014]. Efforts to localize susceptibility variants and investigate their downstream effects on protein synthesis and interactions are hindered by the heterogeneity and complexity of the clinical phenotype, and by the small effect sizes of the common variants typically identified by large-scale association studies. Family-based designs offer increased power to identify genetic variants, especially rare variants with potentially larger effect sizes [Williams and Blangero, 1999]. In addition, intermediate phenotypes that are heritable and genetically associated with a clinical diagnosis can facilitate variant localization, both because of their quantitative nature and because of their assumed proximity to the genetic effects [Glahn et al., 2007; Gottesman and Gould, 2003]. Simultaneously, these endophenotypes [Gottesman and Gould, 2003] provide insights into the variants' influences on biological processes, yielding clues to pathological mechanisms that contribute to the expression of the clinical phenotype.

Neuroanatomical traits derived from magnetic resonance imaging (MRI) are logical endophenotypes as their selection can be informed by a large body of the literature in clinical samples, and they are likely to be biologically intermediate between genes' functions and their more remote effects on behavioral phenotypes [Glahn et al., 2007]. Despite marked clinical and methodological heterogeneity, multiple meta-analyses of voxel-based morphometry studies in schizophrenia have identified the insula and the medial prefrontal cortex (mPFC) as the neuroanatomical regions most consistently associated with schizophrenia across all published case-control studies [Glahn et al., 2008; Fornito et al., 2009; Bora et al., 2011; Palaniyappan et al., 2012; Shepherd et al., 2012].

Source-based morphometry (SBM) [Xu et al., 2009] is a multivariate method that decomposes gray matter concentration images, derived from T_1 -weighted MRI scans, into spatially inde-

pendent sources. The outcome is a matrix of weights for each individual on each source map, which can be used as dependent variables instead of voxel-wise values. As such, SBM dramatically reduces the number of comparisons typically performed in voxel-based analyses. SBM also addresses other common problems in MRI analysis, including the choice of smoothness kernel and the non-stationarity of image smoothness [Hayasaka and Nichols, 2003]. Remarkably, the application of SBM has repeatedly identified a single component comprising the mPFC and the insula—the same regions as the case-control meta-analyses (15–19)—as the most affected anatomical network in schizophrenia patients [Gupta et al., 2015; Xu et al., 2009; Kasperek et al., 2010; Turner et al., 2012]. Thus, following an endophenotype strategy, quantification and localization of the genetic influences on this schizophrenia-associated neuroanatomical trait could provide testable candidate genes and generate novel hypotheses about mechanisms underlying susceptibility for schizophrenia.

Here, we used SBM in a large sample of randomly ascertained pedigrees. Our aims were threefold: (i) to replicate the schizophrenia-associated insula–mPFC source as a spatially independent component in a new sample representative of the general population; (ii) to estimate its heritability; and (iii) to localize this genetic influence to specific genomic regions using linkage analysis.

MATERIALS AND METHODS

GOBS Extended Pedigree Sample

Participants were individuals of Mexican American ancestry who took part in the genetics of brain structure and function study (GOBS) [Olvera et al., 2011; McKay et al., 2014], which is an extension of the San Antonio family study [Mitchell et al., 1996]. Individuals were randomly selected from the community with the only constraints that they were part of a large family of Mexican–American ancestry and lived within the San Antonio region. For the present analysis, subjects were excluded for MRI contraindications, documented medical history of neurological illness, or any neurological event visible on the T_1 -weighted scans (see Source-Based Morphometry Section). Of the participants in the final analysis, 22 self-reported history of a neurological event or illness (18 stroke, 1 Parkinson's disease, 3 multiple sclerosis, 1 brain surgery), but excluding these individuals did not change the pattern of results presented. After quality control procedures (see below), T_1 -weighted scans were available for 887 individuals (532 female), from 69 pedigrees ranging from 2 to 90 family members, and 46 singletons. Participants were between 18 and 85 years old (mean = 44; standard deviation (SD) = 15).

History of axis-1 disorders was assessed using the Mini-International Neuropsychiatric Interview [Sheehan et al., 1998]. Of the individuals included in the main analysis, four had a diagnosis of schizophrenia, 15 of bipolar disorder, and four of schizoaffective disorder. For our main results, we performed additional analyses excluding these individuals (see Results Section). Additionally, 297 participants had a history of major depression, and 118 had a history of an anxiety disorder, and we performed additional analyses co-varying for these diagnoses.

MR Imaging and Processing in GOBS

An MRI protocol optimized for cortical gray matter measurements [Kochunov and Davis, 2009], with a retrospective motion correction technique [Kochunov et al., 2006], was used. For each participant, seven T_1 -weighted scans were obtained in a Siemens 3 Tesla Trio scanner located at the Research Imaging Institute, University of Texas Health Science Center, using a magnetization prepared sequence with an adiabatic inversion contrast-forming pulse (scan parameters: TE/TR/TI = 3.04/2100/785 ms, flip angle = 11°). As in Kochunov et al. [2006], for each subject, the seven volumes were co-registered and averaged.

Upon visual inspection, five individuals were excluded for neurological abnormalities, and one for a scanner artifact. The resulting images were further processed in SPM5 (<http://www.fil.ion.ucl.ac.uk/spm/software/spm5/>), using the same parameters as in Turner et al. [Gupta et al., 2015; Turner et al., 2012] and Xu et al. [2009]. Gray matter maps were non-linearly normalized, resliced to a 2 mm³ MNI template, and segmented into gray matter, white matter, and cerebrospinal fluid [Ashburner and Friston, 2000]. The accuracy of the segmentations and normalizations was ensured by visual inspection and by calculating correlations with the average normalized gray matter map for the entire sample. Five individuals were excluded because of segmentation or normalization problems.

Source-Based Morphometry

SBM [Xu et al., 2009; Kasperek et al., 2010] (<http://mialab.mrn.org/software/gift/>) is a multivariate method that decomposes structural images into spatially distinct sources using independent component analysis [Bell and Sejnowski, 1995]. The decomposition of the subject-by-voxel matrix (X) results in a subject-by-component mixing matrix (W), which contains the weights of the subjects on each component; and a component-by-voxel source matrix (C), which contains the loadings of each voxel for each component. This decomposition¹ can be noted as follows [Calhoun et al., 2001]:

$$X = W * C$$

And, therefore,

$$W = X * C^{-1}$$

For a subject i , the weight of a component j reflects the overall gray matter concentration for that component map of n voxels. More quantitatively, the weight can be conceived of as the sum of each voxel's observed gray matter value multiplied by each voxel's loading on the component:

$$W_{i,j} = \sum_{k=1}^n X_{i,k} C_{k,j}^{-1}$$

¹In practice, this decomposition of interest is preceded by an initial principal component analysis step on the subject-by-voxel matrix X to obtain a square matrix RX , as explained in Calhoun et al. (32). For clarity, we have omitted this step from the current explanation and also note the inverse of C as C^{-1} even though it would technically be non-square if we omitted the PCA step.

These weights can be used as dependent variables in subsequent analyses.

The optimal number of components for our data was estimated according to an information criteria algorithm [Li et al., 2007]. To determine the stability of the decomposition, ICASSO [Himberg et al., 2004] was used, with random value initiation and bootstrapping options, for 20 repetitions.² The stability indices for all components were higher than 0.97.

Relevance to Schizophrenia: Application to an Independent Case-Control Cohort

While we had strong a priori evidence for the involvement of insula and mPFC gray matter in schizophrenia [Glahn et al., 2008; Fornito et al., 2009; Bora et al., 2011; Palaniyappan et al., 2012; Shepherd et al., 2012; Turner et al., 2012], we also directly verified the relevance of the currently derived component to brain morphology of schizophrenia in a separate dataset of 936 healthy control participants (HC) and 784 patients with schizophrenia (SCZ), aggregated from eight independent studies. More detailed information about the case-control sample are presented in the Supplementary Materials and in Gupta et al. [2015].

Using spatio-temporal regression, a method available in the SBM toolbox, we obtained weighting scores for each individual in the case-control dataset for the components identified using the GOBS data. By entering these as the dependent variable in a regression with study site, diagnosis, and their interactions as factors, we tested directly whether gray matter concentration defined by the component extracted from the GOBS dataset was reduced in patients with schizophrenia.

In addition, we quantified the similarity between the components derived from the GOBS and the case-control datasets by calculating (i) pairwise correlations of the voxel loading values across maps from both datasets; and (ii) Dice coefficients of the thresholded maps at $z > 3$, where the Dice coefficient is defined as twice the number of voxels with the same value in both maps divided by total number of voxels (within the masks).

Heritability Analysis

All quantitative genetics analyses were performed in SOLAR [Almasy and Blangero, 1998], which decomposes the variance of a trait into genetic and environmental components by modeling the covariance between individuals as a function of their genetic proximity. Typically, the trait variance is decomposed into an additive genetic effects (heritability), covariate effects (sex, age, age², age \times sex, age² \times sex), and residual environmental effects. We also tested for cubic effects of age, but these were negligible ($P > 0.9$) and dropped from the model. The significance of each variance component is assessed by a likelihood-ratio test

²Note that in this case, bootstrapping is applied to estimate the degree of variability across iterations, and not—as commonly is the case—to estimate a null-distribution, which would require many more iterations.

comparing the final model to the model without the variable of interest. Using the subject's weights on the insula-mPFC component as a trait, this method yielded an index of the overall heritability (h^2) of gray matter concentration within the anatomical regions of the component. Prior to running SOLAR, eight outliers ($\text{mean} \pm 3 \times \text{SD}$) were removed from the data and the remaining weights were transformed using an inverse normalization transformation.

Linkage and Association Analyses

Linkage analysis was performed in SOLAR, by adding location-specific identity-by-descent (IBD) information to the above heritability model. IBD means that two individuals within the same pedigree not only share the same genotype but also inherited it from the same founder. Here, for the GOBS pedigrees, the IBD matrices were estimated as in Curran et al. [2013], using the Loki package [Heath, 1997]. In brief, for 15,000 SNPs³ across the genome, which were selected to be in linkage equilibrium ($r < 0.2$), Loki applies Markov Chain Monte Carlo sampling methods to empirically estimate the pairwise IBD probabilities for each SNP between each individual [Heath, 1997].

The significance of the contribution of each locus is quantified by a LOD score, defined as the logarithm (base 10) of the ratio of the likelihood of the model with the locus-specific IBD matrix to the model without this component (i.e., the same model used to test heritability). The LOD threshold for genome-wide significance was determined a priori for the complex pedigree structure of GOBS. This calculation is based on Gaussian models of the probability of crossover rates under the null distribution [Feingold et al., 1993], given our pedigree structure, the number of SNPs in our IBD matrix, and known Haldane maps. Given our pedigree structure and distribution of markers, a LOD of 2.9 is required for genome-wide significant linkage (genome-wide $\alpha < 0.05$).

To further determine whether any specific SNPs were driving the observed linkage, association analysis was performed for all common SNPs under the linkage peak (defined as all consecutive loci with LODs greater than half the maximum LOD). DNA was extracted from lymphocytes, genotyped using Illumina beadchips (Human1M-Duo Beadchip; or HumanHap550 BeadChip in tandem with HumanHap450S Beadchip), and checked for accordance with Mendelian consistency as described previously [Sprouten et al., 2014]. To account for pedigree structure, association analysis was performed in SOLAR, using methods identical to [Sprouten et al., 2014]. In brief, the minor allele dosage of each tagging SNP is added as a covariate to the model used for heritability. A corrected P -value was calculated according to family-wise error rate at 5%, by calculating in SOLAR the effective number of independent SNPs under the peak, adjusted for linkage disequilibrium (LD) [Moskvina and Schmidt, 2008].

³Note that here the SNPs are not used to test for association testing but merely to represent an independent locus for IBD estimation to perform linkage.

RESULTS

Identification and Validation of Insula-mPFC Component

The method of Li [Li et al., 2007] estimated that the GOBS gray matter images were optimally explained by 21 independent sources.

As hypothesized, one component (Fig. 1) closely resembled the insula-mPFC clusters resulting from the aforementioned voxel-based morphometry meta-analyses [Bora et al., 2011; Palaniyappan et al., 2012; Shepherd et al., 2012], as well as previous SBM studies [Gupta et al., 2015; Turner et al., 2012; Xu et al., 2009]. The bilateral insular parts of this component contained voxels in the insula and temporal pole, extending to the inferior frontal, orbitofrontal, opercular, and superior temporal gyri. The frontal cluster of the component contained voxels in the anterior cingulate and paracingulate gyrus, frontal pole, medial frontal cortex, and superior frontal gyrus. A small number of voxels were negatively correlated with the mPFC and insula, mostly containing white matter in the superior parietal lobes and splenium.

In line with qualitative comparisons to the literature, spatio-temporal regression in the case-control dataset revealed that the weights on this insula-mPFC component were highly significantly different between schizophrenia patients and healthy participants ($F = 292.69$, $P = 1.19 \times 10^{-60}$), in the absence of site-by-diagnosis interactions ($F = 1.24$, $P = 0.22$). The correlation coefficient of the loadings across voxels between this map and the map derived from the case-control data was 0.58 and the Dice coefficient was 0.97, both indicating a high degree of overlap.

The present paper focuses on the insula-mPFC cluster. We provide information about all 21 component maps including heritability estimates and case-control statistics in the Supplementary Materials.

Quantitative Trait Localization

The polygenic model in SOLAR using the weights of the insula-mPFC component as quantitative trait revealed that gray matter in this region is significantly heritable ($h^2 = 0.59$; $P = 1.78 \times 10^{-15}$). Linkage analysis resulted in a genome-wide significant peak on chromosome 12 at 12q24 (12q24.11–12q24.23; maximum LOD = 3.76; Fig. 2).

The linkage peak contained 392 tagging SNPs. Taking into account LD, the peak-wide corrected P -value for SNP associations was 1.64×10^{-4} . None of the SNPs were peak-wide significant (Fig. S2). The strongest association ($P = 7.71 \times 10^{-4}$) was found for rs7133582, an intronic SNP in a transcription factor binding site in *KSR2*. Several other nearby SNPs had modest to strong associations with the insula-mPFC trait, altogether spanning 12q24.21–12q24.23.

Pedigree-specific LODs were all < 1 , indicating that linkage was not driven by any specific pedigree. The heritability and the linkage results remained similar when excluding individuals with self-reported history of neurological events or illness, schizophrenia, schizoaffective disorder, and bipolar disorder ($h^2 = 0.52$; LOD = 3.19) and when co-varying for history of anxiety disorders ($h^2 = 0.60$; LOD = 3.06) and major depressive disorder ($h^2 = 0.60$; LOD = 2.97). There were no significant effects of history of major




FIG. 1. The insula–mPFC component derived from SBM analysis in GOBS. Insula–mPFC component map showing the voxels that primarily contribute to the component of interest, and that covary highly with one another across individuals in the GOBS sample. Voxels loading positively ($z > 2.5$) on the component are colored red-to-yellow, and voxels loading negatively ($z < -2.5$) are colored blue-to-light blue. Images are in radiological convention. [Color figure can be seen in the online version of this article, available at <http://wileyonlinelibrary.com/journal/ajmgb>]

depression ($P > 0.8$) or anxiety disorders ($P > 0.3$) on the component's weights.

To further assess the value of the SBM approach, we also extracted gray matter concentration averages within a binarized mask of the component map ($z > 3$) for each individual. This univariate phenotype was less heritable ($h^2 = 0.47$, $P = 3.08 \times 10^{-11}$), and the maximum LOD score on chromosome 12 was found at the same marker at 12q24 as the multivariate phenotype, but was much lower ($\text{LOD} = 1.82$).

DISCUSSION

We used multivariate analyses of MRI images to extract a gray matter component comprising the insula and the mPFC, which have previously been shown to be the most pronounced [Gupta et al., 2015; Turner et al., 2012] and most consistently implicated gray

matter regions in schizophrenia [Bora et al., 2011; Fornito et al., 2009; Glahn et al., 2007; Palaniyappan et al., 2012; Shepherd et al., 2012]. We directly confirmed that gray matter defined by our empirically derived component was reduced in patients with schizophrenia from an independent case-control sample. Next, we found that the overall gray matter concentration in this component is heritable, and following genome-wide linkage analysis, we identified a quantitative trait locus for this component at 12q24.

Replication and Interpretation of Insula-mPFC Component

So far, SBM has been predominantly applied to case-control studies in schizophrenia [Gupta et al., 2015; Turner et al., 2012; Xu et al., 2009]. Together these studies investigated three independent samples, in four separate analyses, all of which




FIG. 2. Linkage peak on chromosome 12 at 12q24 for weights on the insula–medial prefrontal cortex component. LOD scores plotted against location on chromosome 12. The maximum LOD is 3.76 at 12q24. The threshold for genome-wide significance is at 2.9, as indicated by the dashed line. [Color figure can be seen in the online version of this article, available at <http://wileyonlinelibrary.com/journal/ajmg>] b]

highlight the insula–mPFC component as the most important gray matter component in association with schizophrenia. The replication and heritability of this component in a sample representative of the general population supports the reliability of this technique, and indicates that gray-matter variation in this network is influenced by genetic factors.

The notion of the mPFC–insula as a coherent network is consistent with histological, anatomical, functional, and developmental similarities between these regions. The insula and the mPFC display similarly high rates of cortical thickening during neonatal development [Lyall et al., 2014] and throughout childhood [Sowell et al., 2004]. Histologically, they are also similar most notably because of the distinct and exclusive presence of von Economo neurons in these regions [Butti et al., 2013]. As key regions of the limbic system, and more specifically the “salience network” [Seeley et al., 2007], the mPFC and insula interact intensely to serve higher order cognitive processes such as social and self-awareness, intuition, error monitoring, and interoception. As such, their dysfunction has been postulated to lie at the core of the experience of psychotic symptoms [Kapur, 2003; Palaniyappan et al., 2012; Pu et al., 2012]. In a complimentary theory, interactions between superior temporal, inferior frontal regions, and the mPFC have long been hypothesized as key to the experience of auditory verbal hallucinations [Fletcher et al., 1999; Stephan et al., 2009].

Locus 12q24

We identified a 10 Mb region at 12q24 linked to the insula–mPFC gray matter phenotype. This region has been of great interest to psychiatric genetics since Craddock et al. [1994] reported a cosegregation of bipolar disorder and Darier’s disease, a skin disease that is caused by mutations in *ATP2A2* [Bashir et al., 1993;

Craddock et al., 1993] (Fig. S2). Subsequently, dozens of studies reported linkage of 12q24 to affective disorders in the UK, Ireland, Germany, Denmark, Canada, Iceland, and Finland in locations spanning from 12q22 to 12q24 [Jones et al., 2002; Ekholm et al., 2003; McInnis et al., 2003; Morissette et al., 1999]. Fewer but better powered studies have linked schizophrenia to 12q24 with loci mostly concentrated in 12q24.11–12q24.31 (113–128 Mb; Assembly GRCh37/hg19) [Moises et al., 1995; DeLisi et al., 2002; Williams et al., 2003; Faraone et al., 2006; Bulayeva et al., 2007; Holmans et al., 2009]. This narrowed region matches our mPFC–insula locus, as well as another linkage result in relation to neurocognitive performance in schizophrenia patients [Lien et al., 2010].

In our association study, we were unable to localize our linkage signal to any specific variants. The strongest association was found for rs7133582, a SNP in a transcription factor binding site of *KSR2*, a functionally poorly characterized gene that is involved in the MAPK and ERK signaling pathways [Dougherty et al., 2009]. Similarly, despite the considerable interest in the region, previous studies have not been able to unequivocally identify specific genes that may drive linkage results at 12q24. Although in the most recent genome-wide association analysis of schizophrenia, the Psychiatric Genomics Consortium identified a top SNP in the Darier’s gene *ATP2A2* with $P < 10^{-9}$ (Schizophrenia Working Group of the Psychiatric Genomics 2014), in the original Darier’s disease pedigrees risk for psychiatric disorders did not map to *ATP2A2* itself [Jacobsen et al., 2001], and the co-segregation of psychiatric symptoms with Darier’s disease was thought to be due to nearby variation in LD [Jones et al., 2002]. Other efforts to identify specific neuropsychiatric risk genes at 12q24 overall yielded inconclusive results [Dawson et al., 1995; Jacobsen et al., 1996; Shink et al., 2005]. Several large-scale family studies highlight the genetic heterogeneity at 12q24 [Bulayeva et al., 2007; McInnis et al., 2003; Shink et al., 2005], and it has been suggested that the wider 12q23–12q24 region contains multiple genes that may influence neuropsychiatric phenotypes [Shink et al., 2005; Barden et al., 2006]. The original fine-mapping studies suggested *CUX2*, *FAM109A* (or *FLJ32356*) [Glaser et al., 2005], *P2RX7*, *CAMKK2*, [Barden et al., 2006], and *LINC00944* (“Slynar gene”) [Kalsi et al., 2006; Buttenschon et al., 2010]. However 12q24 also contains the schizophrenia candidate genes *DAO* [Verrall et al., 2010] and *NOS1* [Cui et al., 2010; Silberberg et al., 2010; Wockner et al., 2014]. Interestingly, during fetal development, *NOS1* is transiently highly expressed in the mPFC and insula only [Funk and Kwan, 2014], and variants in *NOS1* have been associated with prefrontal morphology and function [Rose et al., 2012]. Finally, rs7294919, the genome-wide association with hippocampal volume identified by the ENIGMA and CHARGE consortia [Bis et al., 2012; Stein et al., 2012], also lies under our linkage peak.

Strengths and Limitations

An advantage of our multivariate voxel-based analysis is that it gave rise to a reliably identifiable and data-driven single trait that was known to be relevant to schizophrenia based on a large body of the pre-existing literature. However, a limitation of all voxel-based techniques is that the concentration values bear an indirect relationship to the physiological, morphological, and cellular

properties within each voxel. Both regional thickness and surface area contribute independently to voxel-wise gray matter concentrations [Winkler et al., 2010; Rimol et al., 2012], and likely local cortical curvature and white matter morphology also play a role. As such, an understanding of the biological mechanisms of the genetic effects we identified requires further specification of the morphological properties that contribute to the schizophrenia-related phenotype we investigated here.

Secondly, there is a multitude of a priori evidence and a strong rationale for the relevance of our quantitative trait to schizophrenia, which we also directly confirmed in an independent case-control dataset. However, a limitation of our study, as of most intermediate phenotype studies, is that neither endophenotype-disease associations nor quantitative genetics analyses directly test the assumption that the endophenotype lies on the causal pathway from genetic risk variants to development of disease. This is a general limitation of the field that can only truly be addressed by longitudinal designs, which are difficult to obtain to the same quantity as the data we present here, and to this date lack the statistical power and/or suitable pedigree structures for gene localization and identification.

Thirdly, while our linkage peak was not driven by any specific families within our sample, the detection of a genetic signal that is potentially obscured by locus heterogeneity (whether caused by rare or common variants) is likely facilitated by the use of a quantitative intermediate phenotype and the recruitment of large pedigrees. However, as discussed, we were unable to localize the genetic effects to any specific genes or variants. Large-scale family-based analysis of deep sequence data may be necessary to obtain more definitive answers regarding which and to what extent specific genes at 12q24 influence brain morphology and neuropsychiatric phenotypes.

Conclusions

There is compelling evidence that gray matter concentration in the insula and mPFC is reduced in patients with schizophrenia and in their unaffected family members. The genomic region 12q24 has been linked to psychiatric disorders in multiple populations worldwide and contains many genes of interest for neuropsychiatric phenotypes. Our findings indicate that genetic variation in this region also contributes to gray matter concentration in the insula and mPFC in the general population. Thus, mPFC and insula morphology are likely neuroanatomical correlates of schizophrenia that are co-inherited with schizophrenia susceptibility variants at 12q24.

ACKNOWLEDGMENTS

Financial support for this study was provided by the National Institute of Mental Health Grants: MH0708143 (Principal Investigator [PI]: DCG), MH078111 (PI: JB), MH083824 (PI: DCG & JB), and 1R01MH094524 (NIMH, PIs: VC & JAT). Each of these grants contributed to the design and conduct of the study; collection, management, analysis, and interpretation of the data; and preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication. Financial support from COBRE P20GM103472 (PI:VC) contributed to analysis and inter-

pretation of the data; and preparation, review, and approval of the manuscript. SOLAR is supported by MH59490 (PI: JB) and EB015611 (PI: PK), and contributed to analysis and interpretation of the data; and preparation, review, or approval of the manuscript. This investigation was conducted, in part, in facilities constructed with support from Research Facilities Improvement Program Grant Numbers C06 RR13556 and C06 RR017515 from the National Center for Research Resources. We thank participants of the San Antonio Family Study and our research staffs. The supercomputing facilities used for this work at the AT&T Genomics Computing Center were supported, in part, by a gift from the AT&T Foundation and by the National Center for Research Resources Grant Number S10 RR029392.

REFERENCES

- Almasy L, Blangero J. 1998. Multipoint quantitative-trait linkage analysis in general pedigrees. *Am J Hum Genet* 62:1198–1211.
- Ashburner J. 2000. Voxel-based morphometry-the methods. *NeuroImage* 805–821.
- Barden N, Harvey M, Gagne B, Shink E, Tremblay M, Raymond C, Labbe M, et al. 2006. Analysis of single nucleotide polymorphisms in genes in the chromosome 12 Q24. 31 region points to P2RX7 as a susceptibility gene to bipolar affective disorder. *Am J Med Genet B Neuropsychiatr Genet* 141B(4):374–382.
- Bashir R, Munro CS, Mason S, Stephenson A, Rees JL, Strachan T. 1993. Localisation of a gene for Darier's disease. *Hum Mol Genet* 2(11):1937–1939.
- >Bell AJ, Sejnowski TJ. 1995. An information-maximization approach to blind separation and blind deconvolution. *Neural Comput* 7(6):1129–1159.
- Bis JC, DeCarli C, Smith AV, van der Lijn F, Crivello F, Fornage M, Debette S, et al. 2012. Common variants at 12q14 and 12q24 are associated with hippocampal volume. *Nat Genet* 44(5):545–551.
- Bora E, Fornito A, Radua J, Walterfang M, Seal M, Wood SJ, Yucel M, et al. 2011. Neuroanatomical abnormalities in schizophrenia: A multimodal voxelwise meta-analysis and meta-regression analysis. *Schizophr Res* 127(1–3):46–57.
- Bulayeva KB, Glatt SJ, Bulayev OA, Pavlova TA, Tsuang MT. 2007. Genome-wide linkage scan of schizophrenia: A cross-isolate study. *Genomics* 89(2):167–177.
- Buttenschon HN, Foldager L, Flint TJ, Olsen IM, Deleuran T, Nyegaard M, Hansen MM, et al. 2010. Support for a bipolar affective disorder susceptibility locus on chromosome 12q24.3. *Psychiatr Genet* 20(3):93–101.
- Butti C, Santos M, Uppal N, Hof PR. 2013. Von Economo neurons: Clinical and evolutionary perspectives. *Cortex* 49(1):312–326.
- Calhoun VD, Adali T, Pearlson GD, Pekar JJ. 2001. Spatial and temporal independent component analysis of functional MRI data containing a pair of task-related waveforms. *Hum Brain Mapp* 13(1):43–53.
- Craddock N, Dawson E, Burge S, Parfitt L, Mant B, Roberts Q, Daniels J, et al. 1993. The gene for Darier's disease maps to chromosome 12q23-q24.1. *Hum Mol Genet* 2(11):1941–1943.
- Craddock N, Owen M, Burge S, Kurian B, Thomas P, McGuffin P. 1994. Familial cosegregation of major affective disorder and Darier's disease (keratosis follicularis). *Br J Psychiatry* 164(3):355–358.
- Cui H, Nishiguchi N, Yanagi M, Fukutake M, Mouri K, Kitamura N, Hashimoto T, et al. 2010. A putative cis-acting polymorphism in the

- NOS1 gene is associated with schizophrenia and NOS1 immunoreactivity in the postmortem brain. *Schizophr Res* 121(1–3):172–178.
- Curran JE, McKay DR, Winkler AM, Olvera RL, Carless MA, Dyer TD, Kent JW Jr. et al. 2013. Identification of pleiotropic genetic effects on obesity and brain anatomy. *Hum Hered* 75(2–4):136–143.
- Dawson E, Gill M, Curtis D, Castle D, Hunt N, Murray R, Powell J. 1995. Genetic association between alleles of pancreatic phospholipase A2 gene and bipolar affective disorder. *Psychiatr Genet* 5(4):177–180.
- DeLisi LE, Shaw SH, Crow TJ, Shields G, Smith AB, Larach VW, Wellman N, et al. 2002. A genome-wide scan for linkage to chromosomal regions in 382 sibling pairs with schizophrenia or schizoaffective disorder. *Am J Psychiatry* 159(5):803–812.
- Dougherty MK, Ritt DA, Zhou M, Specht SI, Monson DM, Veenstra TD, Morrison DK. 2009. KSR2 is a calcineurin substrate that promotes ERK cascade activation in response to calcium signals. *Mol Cell* 34(6):652–662.
- Ekelund J, Kieseppa T, Hiekkalinna T, Partonen T, Paunio T, Perola M, et al. 2003. Evidence of susceptibility loci on 4q32 and 16p12 for bipolar disorder. *Hum Mol Genet* 12(15):1907–1915.
- Ewald H, Kruse TA, Mors O. 2003. Genome wide scan using homozygosity mapping and linkage analyses of a single pedigree with affective disorder suggests oligogenic inheritance. *Am J Med Genet B, Neuropsychiatr Genet* 120B(1):63–71.
- Faraone SV, Hwu HG, Liu CM, Chen WJ, Tsuang MM, Liu SK, Shieh MH, et al. 2006. Genome scan of Han Chinese schizophrenia families from Taiwan: Confirmation of linkage to 10q22.3. *Am J Psychiatry* 163(10):1760–1766.
- Feingold E, Brown PO, Siegmund D. 1993. Gaussian models for genetic linkage analysis using complete high-resolution maps of identity by descent. *Am J Hum Genet* 53(1):234–251.
- Fletcher P, McKenna PJ, Friston KJ, Frith CD, Dolan RJ. 1999. Abnormal cingulate modulation of fronto-temporal connectivity in schizophrenia. *NeuroImage* 9(3):337–342.
- Fornito A, Yucel M, Patti J, Wood SJ, Pantelis C. 2009. Mapping grey matter reductions in schizophrenia: An anatomical likelihood estimation analysis of voxel-based morphometry studies. *Schizophr Res* 108(1–3):104–113.
- Funk OH, Kwan KY. 2014. Nitric oxide signaling in the development and evolution of language and cognitive circuits. *Neurosci Res* 86:77–87.
- Glahn DC, Laird AR, Ellison-Wright I, Thelen SM, Robinson JL, Lancaster JL, Bullmore E, et al. 2008. Meta-analysis of gray matter anomalies in schizophrenia: Application of anatomic likelihood estimation and network analysis. *Biol Psychiatry* 64(9):774–781.
- Glahn DC, Thompson PM, Blangero J. 2007. Neuroimaging endophenotypes: Strategies for finding genes influencing brain structure and function. *Hum Brain Mapp* 28(6):488–501.
- Glaser B, Kirov G, Green E, Craddock N, Owen MJ. 2005. Linkage disequilibrium mapping of bipolar affective disorder at 12q23-q24 provides evidence for association at CUX2 and F L32356. *Am J Med Genet B Neuropsychiatr Genet* 132B(1):38–45.
- Gottesman II, Gould TD. 2003. The endophenotype concept in psychiatry: Etymology and strategic intentions. *Am J Psychiatry* 160(4):636–645.
- Gratten J, Wray NR, Keller MC, Visscher PM. 2014. Large-scale genomics unveils the genetic architecture of psychiatric disorders. *Nat Neurosci* 17(6):782–790.
- Green E, Elvidge G, Jacobsen N, Glaser B, Jones I, O'Donovan MC, Kirov G, et al. 2005. Localization of bipolar susceptibility locus by molecular genetic analysis of the chromosome 12q23-q24 region in two pedigrees with bipolar disorder and Darier's disease. *Am J Psychiatry* 162(1):35–42.
- Green EK, Grozeva D, Raybould R, Elvidge G, Macgregor S, Craig I, Farmer A, et al. 2009. P2RX7: A bipolar and unipolar disorder candidate susceptibility gene? *Am J Med Genet B Neuropsychiatr Genet* 150B(8):1063–1069.
- Gupta CN, Calhoun VD, Rachkova S, Chen J, Liu J, Segall J, Franke B, et al. Patterns of gray matter abnormalities in schizophrenia based on an international mega-analysis. *Schizophr Bull* In press.
- Heath SC. 1997. Markov chain segregation and linkage analysis for oligogenic models. *Am J Hum Genet* 61:748–760.
- Hayasaka S, Nichols TE. 2003. Validating cluster size inference: Random field and permutation methods. *NeuroImage* 20(4):2343–2356.
- Himberg J, Hyvarinen A, Esposito F. 2004. Validating the independent components of neuroimaging time series via clustering and visualization. *NeuroImage* 22(3):1214–1222.
- Holmans PA, Riley B, Pulver AE, Owen MJ, Wildenauer DB, Gejman PV, Mowry BJ, et al. 2009. Genomewide linkage scan of schizophrenia in a large multicenter pedigree sample using single nucleotide polymorphisms. *Mol Psychiatry* 14(8):786–795.
- Jacobsen N, Daniels J, Moorhead S, Harrison D, Feldman E, McGuffin P, Owen MJ, et al. 1996. Association study of bipolar disorder at the phospholipase A2 gene (PLA2A) in the Darier's disease (DAR) region of chromosome 12q23-q24.1. *Psychiatr Genet* 6(4):195–199.
- Jacobsen NJ, Franks EK, Elvidge G, Jones I, McCandless F, O'Donovan MC, Owen MJ, et al. 2001. Exclusion of the Darier's disease gene, ATP2A2, as a common susceptibility gene for bipolar disorder. *Mol Psychiatry* 6(1):92–97.
- Jones I, Jacobsen N, Green EK, Elvidge GP, Owen MJ, Craddock N. 2002. Evidence for familial cosegregation of major affective disorder and genetic markers flanking the gene for Darier's disease. *Mol Psychiatry* 7(4):424–427.
- Kalsi G, McQuillin A, Degn B, Lundorf MD, Bass NJ, Lawrence J, Choudhury K, et al. 2006. Identification of the Slynar gene (AY070435) and related brain expressed sequences as a candidate gene for susceptibility to affective disorders through allelic and haplotypic association with bipolar disorder on chromosome 12q24. *Am J Psychiatry* 163(10):1767–1776.
- Kapur S. 2003. Psychosis as a state of aberrant salience: A framework linking biology, phenomenology, and pharmacology in schizophrenia. *Am J Psychiatry* 160(1):13–23.
- Kasperek T, Marecek R, Schwarz D, Prikryl R, Vanicek J, Mikl M, Ceskova E. 2010. Source-based morphometry of gray matter volume in men with first-episode schizophrenia. *Hum Brain Mapp* 31(2):300–310.
- Kochunov P, Davis M. 2009. Development of structural MR brain imaging protocols to study genetics and maturation. *Methods* 50(3):136–146.
- Kochunov P, Lancaster JL, Glahn DC, Purdy D, Laird AR, Gao F, Fox P. 2006. Retrospective motion correction protocol for high-resolution anatomical MRI. *Hum Brain Mapp* 27(12):957–962.
- Li YO, Adali T, Calhoun VD. 2007. Estimating the number of independent components for functional magnetic resonance imaging data. *Hum Brain Mapp* 28(11):1251–1266.
- Lien YJ, Liu CM, Faraone SV, Tsuang MT, Hwu HG, Hsiao PC, Chen WJ. 2010. A genome-wide quantitative trait loci scan of neurocognitive performances in families with schizophrenia. *Genes Brain Behav* 9(7):695–702.
- Lyall AE, Shi F, Geng X, Woolson S, Li G, Wang L, Hamer RM, et al. 2014. Dynamic development of regional cortical thickness and surface area in early childhood. *Cereb Cortex*.

- McInnis MG, Lan TH, Willour VL, McMahon FJ, Simpson SG, Addington AM, MacKinnon DF, et al. 2003. Genome-wide scan of bipolar disorder in 65 pedigrees: Supportive evidence for linkage at 8q24, 18q22, 4q32, 2p12, and 13q12. *Mol Psychiatry* 8(3):288–298.
- McKay DR, Knowles EE, Winkler AA, Sprooten E, Kochunov P, Olvera RL, Curran JE, et al. 2014. Influence of age, sex and genetic factors on the human brain. *Brain Imaging Behav* 8(2):143–152.
- Mitchell BD, Kammerer CM, Blangero J, Mahaney MC, Rainwater DL, Dyke B, Hixson JE, et al. 1996. Genetic and environmental contributions to cardiovascular risk factors in Mexican Americans. The San Antonio family heart study. *Circulation* 94(9):2159–2170.
- Moises HW, Yang L, Kristbjarnarson H, Wiese C, Byerley W, Macciardi F, Arolt V, et al. 1995. An international two-stage genome-wide search for schizophrenia susceptibility genes. *Nat Genet* 11(3):321–324.
- Morissette J, Villeneuve A, Bordeleau L, Rochette D, Laberge C, Gagne B, Laprise C, et al. 1999. Genome-wide search for linkage of bipolar affective disorders in a very large pedigree derived from a homogeneous population in quebec points to a locus of major effect on chromosome 12q23-q24. *Am J Med Genet* 88(5):567–587.
- Moskvina V, Schmidt KM. 2008. On multiple-testing correction in genome-wide association studies. *Genet Epidemiol* 32(6):567–573.
- Olvera RL, Bearden CE, Velligan DI, Almasy L, Carless MA, Curran JE, Williamson DE, et al. 2011. Common genetic influences on depression, alcohol, and substance use disorders in Mexican-American families. *Am J Med Genet B Neuropsychiatr Genet* 156B(5):561–568.
- Palaniyappan L, Balain V, Liddle PF. 2012. The neuroanatomy of psychotic diathesis: A meta-analytic review. *J Psychiatr Res* 46(10):1249–1256.
- Pu W, Li L, Zhang H, Ouyang X, Liu H, Zhao J, Li L, et al. 2012. Morphological and functional abnormalities of salience network in the early-stage of paranoid schizophrenia. *Schizophr Res* 141(1):15–21.
- Rimol LM, Nesvag R, Hagler DJ Jr., Bergmann O, Fennema-Notestine C, Hartberg CB, Haukvik UK, et al. 2012. Cortical volume, surface area, and thickness in schizophrenia and bipolar disorder. *Biol Psychiatry* 71(6):552–560.
- Rose EJ, Greene C, Kelly S, Morris DW, Robertson IH, Fahey C, Jacobson S, O'Doherty J, Newell FN, McGrath J, Bokke A, Garavan H, Frodl T, Gill M, Corvin AP, Donohoe G. 2012. The NOS1 variant rs6490121 is associated with variation in prefrontal function and grey matter density in healthy individuals. *NeuroImage* 60(1):614–622.
- Schizophrenia Working Group of the Psychiatric Genomics C. 2014. Biological insights from 108 schizophrenia-associated genetic loci. *Nature* 511(7510):421–427.
- Seeley WW, Menon V, Schatzberg AF, Keller J, Glover GH, Kenna H, Reiss AL, et al. 2007. Dissociable intrinsic connectivity networks for salience processing and executive control. *J Neurosci* 27(9):2349–2356.
- Sheehan DV, Lecrubier Y, Sheehan KH, Amorim P, Janavs J, Weiller E, Hergueta T, et al. 1998. The Mini-International Neuropsychiatric Interview (M.I.N.I.): The development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *J Clin Psychiatry* 59:22–33. quiz 34–57.
- Shepherd AM, Laurens KR, Matheson SL, Carr VJ, Green MJ. 2012. Systematic meta-review and quality assessment of the structural brain alterations in schizophrenia. *Neurosci Biobehav Rev* 36(4):1342–1356.
- Shink E, Harvey M, Tremblay M, Gagne B, Belleau P, Raymond C, Labbe M, et al. 2005. Analysis of microsatellite markers and single nucleotide polymorphisms in candidate genes for susceptibility to bipolar affective disorder in the chromosome 12 Q24. 31 region. *Am J Med Genet B Neuropsychiatr Genet* 135B(1):50–58.
- Silberg G, Ben-Shachar D, Navon R. 2010. Genetic analysis of nitric oxide synthase 1 variants in schizophrenia and bipolar disorder. *Am J Med Genet B Neuropsychiatr Genet* 153B(7):1318–1328.
- Sowell ER, Thompson PM, Leonard CM, Welcome SE, Kan E, Toga AW. 2004. Longitudinal mapping of cortical thickness and brain growth in normal children. *J Neurosci* 24(38):8223–8231.
- Sprooten E, Knowles EE, McKay DR, Goring HH, Curran JE, Kent JW, Jr., Carless MA, et al. 2014. Common genetic variants and gene expression associated with white matter microstructure in the human brain. *NeuroImage* 97:252–261.
- Stein JL, Medland SE, Vasquez AA, Hibar DP, Senstad RE, Winkler AM, Toro R, et al. 2012. Identification of common variants associated with human hippocampal and intracranial volumes. *Nat Genet* 44(5):552–561.
- Stephan KE, Friston KJ, Frith CD. 2009. Dysconnection in schizophrenia: From abnormal synaptic plasticity to failures of self-monitoring. *Schizophr Bull* 35(3):509–527.
- Sullivan PF, Kendler KS, Neale MC. 2003. Schizophrenia as a complex trait: Evidence from a meta-analysis of twin studies. *Arch Gen Psychiatry* 60(12):1187–1192.
- Turner JA, Calhoun VD, Michael A, van Erp TG, Ehrlich S, Segall JM, Gollub RL, et al. 2012. Heritability of multivariate gray matter measures in schizophrenia. *Twin Res Hum Gen* 15(3):324–335.
- Verrall L, Burnet PW, Betts JF, Harrison PJ. 2010. The neurobiology of D-amino acid oxidase and its involvement in schizophrenia. *Mol Psychiatry* 15(2):122–137.
- Williams JT, Blangero J. 1999. Comparison of variance components and sibpair-based approaches to quantitative trait linkage analysis in unselected samples. *Genet Epidemiol* 16(2):113–134.
- Williams NM, Norton N, Williams H, Ekholm B, Hamshere ML, Lindblom Y, Chowdari KV, et al. 2003. A systematic genomewide linkage study in 353 sib pairs with schizophrenia. *Am J Hum Genet* 73(6):1355–1367.
- Winkler AM, Kochunov P, Blangero J, Almasy L, Zilles K, Fox PT, Duggirala R, et al. 2010. Cortical thickness or grey matter volume? The importance of selecting the phenotype for imaging genetics studies. *NeuroImage* 53(3):1135–1146.
- Wockner LF, Noble EP, Lawford BR, Young RM, Morris CP, Whitehall VL, Voisey J. 2014. Genome-wide DNA methylation analysis of human brain tissue from schizophrenia patients. *Trans Psychiatry* 4:e339.
- Xu L, Groth KM, Pearlson G, Schretlen DJ, Calhoun VD. 2009. Source-based morphometry: The use of independent component analysis to identify gray matter differences with application to schizophrenia. *Hum Brain Mapp* 30(3):711–724.

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of this article at the publisher's web-site.