

Anomalies are in the Physical World

Samuel Epstein*

July 13, 2022

Abstract

The Independence Postulate (IP) is a finitary Church Turing Thesis, postulating that certain finite and infinite sequences cannot be found in nature, i.e. have high “physical addresses”. In this paper we show that IP implies that anomalies are found in the physical world.

Anomalies

$\mathbf{K}(x|y)$ is the conditional prefix Kolmogorov complexity. For probability p over \mathbb{N} , the deficiency of randomness is $\mathbf{d}(a|p) = \lfloor -\log p(a) \rfloor - \mathbf{K}(a)$. $\mathbf{I}(a; \mathcal{H}) = \mathbf{K}(a) - \mathbf{K}(a|\mathcal{H})$, where \mathcal{H} is the halting sequence. An elementary probability measure over \mathbb{N} has finite support and a range in \mathbb{Q} . Stochasticity is

$$\begin{aligned} \Lambda(a|b) &= \min\{\mathbf{K}(Q|b) + 3 \log \max\{\mathbf{d}(a|Q, b), 1\} \\ &\quad : Q \text{ is an elementary probability measure}\}. \\ \Lambda(a|b) &< \Lambda(a) + O(\log \mathbf{K}(b)). \end{aligned}$$

The following definition is from [Lev74] .

Definition 1 (Information) $\mathbf{I}(\alpha : \beta) = \log \sum_{x,y} 2^{\mathbf{K}(x) + \mathbf{K}(y) - \mathbf{K}(x,y) - \mathbf{K}(x|\alpha) - \mathbf{K}(y|\beta)}$.

The Independence Postulate [Lev84, Lev13] statement is:

IP: Let α be a sequence defined with an n -bit mathematical statement, and a sequence β can be located in the physical world with a k -bit instruction set. Then $\mathbf{I}(\alpha : \beta) < k + n + c$ for some small absolute constant c .

It is well known in the literature that stochastic numbers have high mutual information with the halting sequence. One detailed proof is in [Eps21].

Lemma 1 $\Lambda(x) <^{\log} \mathbf{I}(x; \mathcal{H})$.

Lemma 2 For computable probability p over \mathbb{N} and for $D \subset \mathbb{N}$, $|D| = 2^s$, $s < \max_{a \in D} \mathbf{d}(a|p) + \Lambda(D) + \mathbf{K}(s) + \mathbf{K}(p) + O(\log \mathbf{K}(s)\mathbf{K}(p))$.

Proof. We relativize the universal Turing machine to p and s . Let Q be an elementary probability measure that realizes $\Lambda(D)$. Let $d = \mathbf{d}(D|Q)$. Let $F \subseteq \mathbb{N}$ be a random set where each element of F is selected independently with probability $cd2^{-s}$, where $c \in \mathbb{N}$ is chosen later. $\mathbf{E}[p(F)] \leq cd2^{-s}$. Furthermore

$$\begin{aligned} &\mathbf{E}[Q(\{G : |G| = 2^s, G \cap F = \emptyset\})] \\ &\leq \sum_G Q(G)(1 - cd2^{-s})^{2^s} < e^{-cd}. \end{aligned}$$

Thus finite $W \subset \mathbb{N}$ can be chosen such that $p(W) \leq 2cd2^{-s}$ and $Q(\{G : |G| = 2^s, G \cap W = \emptyset\}) \leq e^{1-cd}$. $D \cap W \neq \emptyset$, otherwise, using the Q -test, $t(G) = e^{cd-1}$ if $(|G| = 2^s, G \cap W = \emptyset)$ and $t(G) = 0$ otherwise, we have

$$\begin{aligned} \mathbf{K}(D|Q, d, c) &<^+ -\log Q(D) - (\log e)cd \\ (\log e)cd &<^+ -\log Q(D) - \mathbf{K}(D|Q) + \mathbf{K}(d, c) \\ (\log e)cd &<^+ d + \mathbf{K}(d, c), \end{aligned}$$

which is a contradiction for large c . Thus there is an $a \in D \cap W$, where

$$\begin{aligned} \mathbf{K}(a) &<^+ -\log p(a) + \log d - s + \mathbf{K}(d) + \mathbf{K}(Q) \\ s &<^+ \mathbf{d}(a|p) + \Lambda(D). \end{aligned}$$

Removing the relativization of p and s

$$\begin{aligned} s &< -\log p(a) - \mathbf{K}(a|p, s) + \Lambda(D|p, s) \\ s &< \max_{a \in D} \mathbf{d}(a|p) + \Lambda(D) + \mathbf{K}(s) + \mathbf{K}(p) \\ &\quad + O(\log \mathbf{K}(s)\mathbf{K}(p)). \end{aligned}$$

*JP Theory Group. samepst@jpththeorygroup.org

□

For $\tau \in \mathbb{N}^{\mathbb{N}}$, $\langle \tau \rangle = \langle \tau[1] \rangle \langle \tau[2] \rangle \langle \tau[3] \rangle \dots$. Let $\tau(n)$ be the first 2^n unique numbers found in τ . The sequence τ is assumed to have an infinite amount of unique numbers, and represents a series of observations.

Theorem 1 *For probability p over \mathbb{N} and $\tau \in \mathbb{N}^{\mathbb{N}}$, let $s_{\tau,p}$ be the smallest number such that $\max_{a \in \tau(n)} \mathbf{d}(a|p) > n - 4\mathbf{K}(n) - s_{\tau,p}$. Then $s_{\tau,p} <^{\log} \mathbf{I}(\langle \tau \rangle : \mathcal{H}) + \mathbf{K}(p)$.*

Proof. By Lemmas 1 and 2, and the fact that $\mathbf{I}(x; \mathcal{H}) <^+ \mathbf{I}(\alpha : \mathcal{H}) + \mathbf{K}(x|\alpha)$,

$$\begin{aligned} n &< \max_{a \in \tau(n)} \mathbf{d}(a|p) + \mathbf{I}(\tau(n); \mathcal{H}) + \mathbf{K}(p) + 2\mathbf{K}(n) \\ &\quad + O(\log(\mathbf{I}(\tau(n); \mathcal{H}) + \mathbf{K}(p))), \\ n &< \max_{a \in \tau(n)} \mathbf{d}(a|p) + 4\mathbf{K}(n) + \mathbf{I}(\langle \tau \rangle : \mathcal{H}) + \mathbf{K}(p) \\ &\quad + O(\log(\mathbf{I}(\langle \tau \rangle : \mathcal{H}) + \mathbf{K}(p))), \\ n - 4\mathbf{K}(n) - (\mathbf{I}(\langle \tau \rangle : \mathcal{H}) + \mathbf{K}(p)) \\ &\quad + O(\log(\mathbf{I}(\langle \tau \rangle : \mathcal{H}) + \mathbf{K}(p))) < \max_{a \in \tau(n)} \mathbf{d}(a|p). \end{aligned}$$

□

Let k be the physical address of an infinite sequence of numbers $\tau \in \mathbb{N}^{\mathbb{N}}$. The halting sequence can be described by a small mathematical statement. By Theorem 1 and **IP**,

$$s_{\tau,p} <^{\log} \mathbf{I}(\langle \tau \rangle : \mathcal{H}) + \mathbf{K}(p) <^{\log} k + c + \mathbf{K}(p).$$

Thus sequences τ with large s_{τ} , as defined in Theorem 1, will have large physical addresses. So it's hard to find observations which do not have large anomalies, and impossible to find observations with no anomalies.

References

- [Eps21] Samuel Epstein. All sampling methods produce outliers. *IEEE Transactions on Information Theory*, 67(11):7568–7578, 2021.
- [Lev74] L. A. Levin. Laws of Information Conservation (Non-growth) and Aspects of the Foundations of Probability Theory. *Problemy Peredachi Informatsii*, 10(3):206–210, 1974.
- [Lev84] L. A. Levin. Randomness conservation inequalities; information and independence in mathematical theories. *Information and Control*, 61(1):15–37, 1984.
- [Lev13] L. A. Levin. Forbidden information. *J. ACM*, 60(2), 2013.