

Quantum Decoherence Mostly Results in White Noise

Samuel Epstein
samepst@jpththeorygroup.org

April 4, 2023

Abstract

An overwhelming majority of quantum (pure and mixed) states, when undertaking decoherence, will result in a classical probability with no algorithmic information. Thus most quantum states decohere into white noise. This can be seen as a consequence of the vastness of Hilbert spaces.

Information non-growth laws say information about a target source cannot be increased with randomized processing. In classical information theory, we have [CT91]

$$I(g(X):Y) \leq I(X:Y).$$

where g is a randomized function, X and Y are random variables, and I is the mutual information function. Thus processing a channel at its output will not increase its capacity. Information conservation carries over into the algorithmic domain, with the inequalities [Lev84, Eps22]

$$\mathbf{I}(f(x):y) <^+ \mathbf{I}(x:y); \quad \mathbf{I}(f(a);\mathcal{H}) <^+ \mathbf{I}(a;\mathcal{H}).$$

The information function is $\mathbf{I}(x:y) = \mathbf{K}(x) + \mathbf{K}(y) - \mathbf{K}(x,y)$, where \mathbf{K} is Kolmogorov complexity. The other term is $\mathbf{I}(a;\mathcal{H}) = \mathbf{K}(a) - \mathbf{K}(a|\mathcal{H})$, where $\mathcal{H} \in \{0,1\}^\infty$ is the halting sequence. These inequalities ensure target information cannot be obtained by processing. If for example the second inequality was not true, then one can potentially obtain information about \mathcal{H} with simple functions. Obtaining information about \mathcal{H} violates the Independence Postulate, (see [Lev13]). Information non growth laws can be extended to signals [Eps23b] which can be modeled as probabilities over \mathbb{N} or Euclidean space¹. The “signal strength” of a probability p over \mathbb{N} is measured by its self information.

$$\mathbf{I}_{\text{Prob}}(p:p) = \log \sum_{i,j} 2^{\mathbf{I}(i:j)} p(i)p(j).$$

A signal, when undergoing randomized processing f , will lose its cohesion². Thus any signal going through a classical channel will become less coherent [Eps23b].

$$\mathbf{I}_{\text{Prob}}(f(p):f(p)) <^+ \mathbf{I}_{\text{Prob}}(p:p).$$

In Euclidean space, probabilities that undergo convolutions with probability kernels will lose self information. For example a signal spike at a random position will spread out when convoluted with the Gaussian function, and lose self information. The above inequalities deal with classical

¹In [Eps23b] probabilities over $\{0,1\}^\infty$ and T_0 second countable topologies were also studied.

²A probability p , when processed by a channel $f : \{0,1\}^* \times \{0,1\}^* \rightarrow \mathbb{R}_{\geq 0}$ is a new probability $fp(x) = \sum_z f(x|z)p(z)$.

transformations. One can ask, is whether, quantum information processing can add new surprises to how information signals occur and evolve.

One can start with the prepare-and-measure channel, also known as a Holevo-form channel. Alice starts with a random variable X that can take values $\{1, \dots, n\}$ with corresponding probabilities $\{p_1, \dots, p_n\}$. Alice prepares a quantum state, corresponding to density matrix ρ_X , chosen from $\{\rho_1, \dots, \rho_n\}$ according to X . Bob performs a measurement on the state ρ_X , getting a classical outcome, denoted by Y . Though it uses quantum mechanics, this is a classical channel $X \rightarrow Y$. So using the above inequality, cohesion will deteriorate regardless of X 's probability, with

$$\mathbf{I}_{\text{Prob}}(Y : Y) <^+ \mathbf{I}_{\text{Prob}}(X : X).$$

There remains a second option, constructing a signal directly from a mixed state. This involves constructing a mixed state, i.e. density matrix σ , and then performing a POVM measurement³ E on the state, inducing the probability $E\sigma(\cdot)$. However from [Eps23b], for elementary (even enumerable) probabilities $E\sigma$,

$$\mathbf{I}_{\text{Prob}}(E\sigma : E\sigma) <^+ \mathbf{K}(\sigma, E).$$

Thus for simply defined density matrices and measurements, no signal can appear. So experiments that are simple will result in simple measurements, or white noise. However it could be that a larger number of uncomputable pure or mixed states produce coherent signals. However, theorems in [Eps23a] say otherwise, in that given a POVM measurement E , a vast majority of pure and mixed states will have negligible self-information. Thus for uniform distributions Λ and μ over pure and mixed states⁴⁵,

$$\int 2^{\mathbf{I}_{\text{Prob}}(E|\psi\rangle : E|\psi\rangle)} d\Lambda = O(1); \quad \int 2^{\mathbf{I}_{\text{Prob}}(E\sigma : E\sigma)} d\mu(\sigma) = O(1).$$

This can be seen as a consequence of the vastness of Hilbert spaces as opposed to the limited discriminatory power of quantum measurements. In addition, there could be non-uniform distributions of pure or mixed states that could be of research interest. In quantum decoherence, a quantum state becomes entangled with the environment, losing decoherence. The off diagonal elements of the mixed state become dampened, as the state becomes more like a classical mixture of states. Let p_σ be the idealized classical probability that σ decoheres to, with $p_\sigma(i) = \sigma_{ii}$. The following theorem from [Eps23a] states that for an overwhelming majority of pure or mixed states σ , p_σ is noise, that is, has negligible self-information.

$$\int 2^{\mathbf{I}_{\text{Prob}}(p_{|\psi\rangle} : p_{|\psi\rangle})} d\Lambda = O(1); \quad \int 2^{\mathbf{I}_{\text{Prob}}(p_\sigma : p_\sigma)} d\mu(\sigma) = O(1).$$

This is to be expected, with one supporting fact being for an n qubit space, $i \in \{1, \dots, 2^n\}$, $\mathbf{E}_\Lambda[p_{|\psi\rangle}(i)] = 2^{-n}$. With Algorithmic Information Theory, we've taken this fact one step further, showing that $p_{|\psi\rangle}$ has no (in the exponential) self-algorithmic information and cannot be processed by deterministic or randomized means to produce a more coherent signal. In addition, it appears a more direct proof than the one in [Eps23a] of the first decoherence inequality could be possible.

These results mark a continuing investigation of algorithmic information and physics. The results of this survey show that the boundary between the quantum world and the classical world is permeated by white noise, so how is information transmitted from the former to the latter?

³A POVM measurement E is a collection of positive-semi definite Hermitian matrices $\{E_k\}$ such that $\sum_k E_k = 1$. Given a state σ , E induces a probability over the measurements of the form $E\sigma(k) = \text{Tr} E_k \sigma$.

⁴The mixed state integral is $\int f(\sigma) d\mu(\sigma) = \int_{\Delta_M} \int_{\Lambda_1} \dots \int_{\Lambda_M} f\left(\sum_{i=1}^M p_i |\psi_i\rangle \langle \psi_i|\right) d\Lambda_1 \dots d\Lambda_M d\eta(p_1, \dots, p_M)$, where η is any distribution over the M -simplex Δ_M .

⁵The proof to these inequalities is in the running for the strangest in AIT, relying on a lower computable combination of *upper* computable tests.

References

- [CT91] T. Cover and J. Thomas. *Elements of Information Theory*. Wiley-Interscience, New York, NY, USA, 1991.
- [Eps22] S. Epstein. The outlier theorem revisited. *CoRR*, abs/2203.08733, 2022.
- [Eps23a] S. Epstein. An Introduction to Algorithmic Information Theory and Quantum Mechanics. 2023. <http://www.jptheorygroup.org/doc/AITQM.pdf>.
- [Eps23b] S. Epstein. On the algorithmic information between probabilities. *CoRR*, abs/2303.07296, 2023.
- [Lev84] L. A. Levin. Randomness conservation inequalities; information and independence in mathematical theories. *Information and Control*, 61(1):15–37, 1984.
- [Lev13] L. A. Levin. Forbidden information. *J. ACM*, 60(2), 2013.