

# On Kolmogorov Structure Functions

Sam Epstein\*

May 31, 2024

## Abstract

All strings with low mutual information with the halting sequence will have flat Kolmogorov Structure Functions, in the context of Algorithmic Statistics. Assuming the Independence Postulate, strings with non-negligible information with the halting sequence are purely mathematical constructions, and cannot be found in the physical world.

## 1 Introduction

In statistics, one tries to determine a model (such as a parameter for a distribution) from data which is assumed to have noise. In the Minimum Description Principle [Gru07], the model that describes information with the shortest code is assumed to be the best model. The data is described as a two part code, where the first part is the model and the second part is the noise. In one of his last works, Kolmogorov suggested a two part code for individual strings  $x \in \{0, 1\}^*$  based off Kolmogorov Complexity. The first part (the model) is a set  $D$  containing  $x$ , the second part (the noise) is the code of  $x$  given  $D$ , of size  $\lceil \log |D| \rceil$ . Other works examined probabilities and also total computable functions as models [Vit02]. Kolmogorov suggested the following *structure function* at the Tallinn conference in Estonia, 1973.

$$\mathbf{H}_k(x) = \{\log |S| : x \in S, \mathbf{K}(S) \leq k\}.$$

The function  $\mathbf{K}$  is the prefix Kolmogorov complexity. This definition is used for the following function, which is a central definition of *Algorithmic Statistics* [VS15, VS17, VV04],

$$k \mapsto k + \mathbf{H}_k(x) - \mathbf{K}(x).$$

This function's equivalence to several other definitions is the main theorem of Algorithmic Statistics [SSV24]. Furthermore, Theorem 1 of [VS17] showed that any shape of the structure function is possible.

The structure function is flat for all strings with low mutual information with the halting sequence. Assuming the *Independence Postulate*, [Lev84, Lev13], strings with non-negligible mutual information with the halting sequence are exotic, in that they cannot be found in nature. Such strings are purely mathematical constructions.

---

\*samepst@jpththeorygroup.org

## 2 Bounds

We review the results of [GTV01], in particular Theorem of III.24, which I don't think is widely known in the literature.  $\mathbf{m}(x)$  is the algorithmic probability. The amount of information that the halting sequence  $\mathcal{H} \in \{0,1\}^\infty$  has about  $x \in \{0,1\}^*$  is  $\mathbf{I}(x; \mathcal{H}) = \mathbf{K}(x) - \mathbf{K}(x|\mathcal{H})$ . We use  $x <^+ y$ ,  $x >^+ y$  and  $x =^+ y$  to denote  $x < y + O(1)$ ,  $x + O(1) > y$  and  $x = y \pm O(1)$ , respectively. In addition,  $x <^{\log} y$  and  $x >^{\log} y$  denote  $x < y + O(\log y)$  and  $x + O(\log x) > y$ , respectively. For  $x, y \in \{0,1\}^*$ ,  $x \sqsubseteq y$  if  $y = xz$  for some  $z \in \{0,1\}^*$ .  $[A] = 1$  if mathematical statement  $A$  is true, and  $[A] = 0$  otherwise.

Let  $S_k = \{x : \mathbf{K}(x) \leq k\}$ . Let  $N_k = |S_k|$  where  $\log N_k =^+ k - \mathbf{K}(k)$ , due to [GTV01]. Let  $I_k^x$  be the index of  $x$  in an enumeration of  $S_k$ . For  $\mathbf{K}(x) = k$ , let  $m_x$  be the longest joint prefix of  $I_k^x$  and  $N_k$ . So  $m_x 0 \sqsubseteq I_k^x$  and  $m_x 1 \sqsubseteq N_k$ . Let  $S_x = \{y : m_x 0 \sqsubseteq I_k^y\}$ . So

$$\begin{aligned} \log |S_x| &=^+ k - \mathbf{K}(k) - \|m_x\| \\ \mathbf{K}(S_x) &<^+ \mathbf{K}(k) + \mathbf{K}(m_x) <^+ \mathbf{K}(k) + \|m_x\| + \mathbf{K}(\|m_x\|). \end{aligned}$$

**Theorem 1** ([GTV01]).

$$\|m_x\| < \mathbf{K}(\mathbf{K}(x)) + \mathbf{I}(x; \mathcal{H}) + O(\log \mathbf{I}(x; \mathcal{H})).$$

*Proof.* Let  $\nu(y) = c[\mathbf{K}(y) \leq k] \mathbf{m}(y) 2^{\|m_y\|} / (\|m_y\|^2)$ . For proper choice of  $c$ ,  $\nu$  is a semimeasure and computable relative to  $\mathcal{H}$  and  $k$ . So  $\mathbf{K}(x|\mathcal{H}, k) <^+ -\log \nu(x) =^+ \mathbf{K}(x) - \|m_x\| + 2 \log \|m_x\|$ .  $\square$

Note that with some additional effort, the  $\mathbf{K}(\mathbf{K}(x))$  term can be eliminated.

**Corollary 1.** For  $x \in \{0,1\}^*$ ,  $n = \mathbf{K}(x)$ , for all  $m \leq n$ ,  $m \in \mathbb{W}$ , there is a set  $S \ni x$  such that  $|S| = 2^m$  and  $\mathbf{K}(S) + m <^{\log} n + \mathbf{I}(x; \mathcal{H})$ .

**Claim 1.** Thus there exists a set  $S \ni x$  such that  $\mathbf{K}(S) < 2\mathbf{K}(\mathbf{K}(x)) + \mathbf{I}(x; \mathcal{H}) + O(\log \mathbf{I}(x; \mathcal{H}))$  and  $\mathbf{K}(S) + \log |S| <^+ \mathbf{K}(x) + \mathbf{K}(\mathbf{K}(x)) + O(\log \mathbf{I}(x; \mathcal{H}))$ . This fact combined with the following proposition characterizes the Kolmogorov Structure Function.

**Proposition 1.** Let  $S \ni x$ . For all  $s < \log |S|$  there exists a set  $S' \ni x$  such that  $|S'| \leq |S| 2^{-s}$  and  $\mathbf{K}(S') <^+ \mathbf{K}(S) + s + \mathbf{K}(s)$ .

## References

- [Gru07] P. Grunwald. *The Minimum Description Length Principle*. The MIT Press, 2007.
- [GTV01] P. Gács, J. Tromp, and P. Vitányi. Algorithmic statistics. *Information Theory, IEEE Transactions on*, 47:2443 – 2463, 2001.

- [Lev84] L. A. Levin. Randomness conservation inequalities; information and independence in mathematical theories. *Information and Control*, 61(1):15–37, 1984.
- [Lev13] L. A. Levin. Forbidden information. *J. ACM*, 60(2), 2013.
- [SSV24] A. Semenov, A. Shen, and N. Vereshchagin. Kolmogorov’s Last Discovery? (Kolmogorov and Algorithmic Statistics). *Theory of Probability & Its Applications*, 68(4):582–606, 2024.
- [Vit02] P. Vitányi. Meaningful information. In *Algorithms and Computation*, pages 588–599, Berlin, Heidelberg, 2002. Springer Berlin Heidelberg.
- [VS15] N. Vereshchagin and A. Shen. *Algorithmic Statistics Revisited*, pages 235–252. Springer International Publishing, 2015.
- [VS17] N. Vereshchagin and A. Shen. *Algorithmic Statistics: Forty Years Later*, pages 669–737. Springer International Publishing, 2017.
- [VV04] N. Vereshchagin and P. Vitányi. Kolmogorov’s structure functions and model selection. *IEEE Transactions on Information Theory*, 50(12):3265–3290, 2004.