

Towards a General Theory of Algorithmic Physics

Samuel Epstein
samepst@jptheorygroup.org

September 2, 2023

Abstract

This article presents a survey of published and unpublished material of the intersection of algorithmic information theory, quantum mechanics, and thermodynamics. It is, to the author's knowledge, the first of its type. Three different notions of the algorithmic content of quantum states are reviewed. Notions of algorithmic quantum typicality and mutual information are introduced. The relationship between algorithmic information and quantum measurements is explored. One of the surprising results is that an overwhelming majority of quantum (pure and mixed) states, when undertaking decoherence, will result in a classical probability with no algorithmic information. Thus most quantum states decohere into white noise. A quantum analogue of Martin L f random sequence is reviewed. Algorithmic Information Theory presents new complications for the Many Worlds Theory, as it conflicts with the Independence Postulate. When algorithmically complicated processes are ruled out, measurements are required to produce distributions over quantum states that have cloneable information.

As for thermodynamics, new definitions of algorithmic coarse and fine grained entropy are introduced. Algorithmic fine grained entropy obeys the chain rule. The function oscillates during the course of dynamics. Small fluctuations are common and larger fluctuations are more rare. Coarse grained entropy is shown to be an excellent approximation to fine grained entropy. It is shown to oscillate in the presence of dynamics as well. Marginal algorithmic thermodynamic entropies cannot be synchronized. Synchronized oscillations occur during the course of discrete dynamics with algorithmic fine grained entropy with respect to two volume measures. Properties of discrete ergodic dynamics are proven, in that, in the limit, a state will be in a partition cell with frequency equal to its measure.

Contents

I	Introduction	5
1	Introduction	6
1.1	Some Surprises About Algorithmic Physics	6
1.1.1	Quantum States Have No Self-Information.	6
1.1.2	Algorithmic Thermodynamic Entropy and Fluctuations.	7
1.1.3	Typicality and Newtonian Physics	7
1.2	Conventions	8
1.2.1	Algorithmic Information Theory	8
1.2.2	Algorithmic Information Between Probabilities	10
II	Quantum Mechanics	12
2	Introduction	13
3	Background	15
3.1	Algorithmic Information Theory	15
3.1.1	Algorithmic Information Between Probabilities	16
3.2	Quantum Mechanics	17
3.2.1	Quantum Operations	18
4	Quantum Complexity	19
4.1	Three Measures of the Algorithmic Content of Individual Quantum States	19
4.1.1	BVL Complexity	19
4.1.2	Vitányi Complexity	19
4.1.3	Gács Complexity	20
4.2	Properties of Universal Matrix and Gács Complexity	20
4.3	No Cloning Theorem	21
4.4	Addition Inequality	22
4.5	Subadditivity, Strong Subadditivity, Strong Superadditivity	23
4.6	Relation Between Complexities	24
4.6.1	Vitányi Complexity and Gács Complexity	24
4.6.2	BVL Complexity and Gács Complexity	25
4.7	Quantum EL Theorem	26
4.7.1	Computable Projections	27
4.7.2	Quantum Data Compression	28

5	Quantum Typicality	29
5.1	Definition of Quantum Randomness Deficiency	29
5.1.1	Uncomputable Mixed States	30
5.2	Conservation Over Quantum Operations	31
5.3	A Quantum Outlier Theorem	31
5.3.1	Computable Projections	33
6	Quantum Information	35
6.1	Definition of Quantum Algorithmic Information	35
6.2	Paucity of Self-Information	36
6.2.1	Pure States	36
6.2.2	Mixed States	37
6.3	Information Nongrowth	38
6.3.1	Algorithmic No-Cloning Theorem	39
6.3.2	Purification	40
6.3.3	Decoherence	40
7	Quantum Measurements	42
7.1	Typicality and Measurements	42
7.2	Information and Measurements	42
7.3	Algorithmic Contents of Measurements	43
7.3.1	Algorithmic Content of Decoherence	45
7.4	PVMs	45
8	Infinite Quantum Spin Chains	46
8.1	Infinite Quantum Bit Sequences	46
8.1.1	NS Randomness	47
8.2	Closure Properties	47
8.3	Gács Complexity and NS Random Sequences	48
8.4	Encodings of States	49
8.5	Quantum Operation Complexity	49
8.6	Initial Segment Incompressibility	49
8.7	Quantum Ergodic Sources	51
8.8	Measurement Systems	53
8.9	NS Solovay States	54
9	The Many Worlds Theory	56
9.1	Many Worlds Theory	56
9.1.1	Branching Worlds	57
9.1.2	Deriving the Born Rule	57
9.2	Violating the Independence Postulate	58
9.3	Conclusion	59
10	Conclusion	61
10.1	Signals from Classical and Quantum Sources	61
10.2	Apriori Distributions	62
10.3	Measurements Before Information Cloning	64

III	Thermodynamics	66
11	Computable Measure Spaces	69
11.1	Enumerative Lattices	69
11.2	Computing with Measures	70
11.3	Computable Measure Space	73
11.4	Randomness	75
12	Algorithmic Fine Grained Entropy	78
12.1	Thermodynamic Information	79
12.2	Entropy Balance	80
12.3	Maxwell's Demon	80
12.4	Distribution of Algorithmic Fine Grained Entropy	81
12.5	Addition Equality	83
13	Oscillation of Algorithmic Fine Grained Entropy	86
13.1	On Exotic Sets of Natural Numbers	86
13.2	On Exotic Sets of Reals	87
13.3	Oscillations Occur	88
13.4	Oscillations are Rare	91
14	Discrete Dynamics	93
14.1	Synchronized Oscillations	93
14.2	Ergodic Dynamics	94
14.2.1	Single Points	94
14.2.2	Indicator Functions	95
14.2.3	Lower Computable Functions	97
15	Dynamics on Product Spaces	99
15.1	Continuous Dynamics	99
15.1.1	Exotic Intervals	99
15.1.2	No Synchronization Under Continuous Dynamics	100
15.2	Ergodic Dynamics	100
15.2.1	Effective Convergence Time of Ergodic Dynamics	100
15.2.2	Busy Beaver Functions	102
15.2.3	Marginal Entropies of Ergodic Dynamics	102
16	Algorithmic Coarse Grained Entropy	105
16.1	Fundamentals of Coarse Grained Entropy	105
16.2	Entropy Variance in Cells	109
17	Computability of Algorithmic Entropy	111
17.1	Uncomputability of Decreasing Entropy	111
17.2	Non-Approximability of Algorithmic Entropy	111
17.2.1	Kolmogorov Complexity is Exotic	111
17.2.2	Algorithmic Fine Grain Entropy Cannot be Approximated	113
17.3	Computability of Measures	113
17.4	Exotic Dynamics	115

IV	Newtonian Gravity	117
18	Typicality in Newtonian Gravity	118
18.1	Introduction	118
18.2	Preliminaries	118
18.3	Properties of Universal Tests	120
18.4	Conservation of Randomness	121
18.5	Comparable Definitions of Typicality	122
18.6	Compact Subspaces	123
18.7	Orbits	124
V	Appendix	132
A	Information Between Basis States	133
B	An Extended Coding Theorem	136
B.1	Left-Total Machines	136
B.2	Extended Coding Theorem	138

Part I

Introduction

Chapter 1

Introduction

1.1 Some Surprises About Algorithmic Physics

This manuscript deals with the application of algorithmic information theory to physics, namely quantum information theory and thermodynamics. The reader is assumed to be familiar with these three areas, and the reader is referred to the books [LO97, Wil13, She15]. The main references to this manuscript are [G01, Gac94, HR09, SBC01, Vit01, NS19, Vai98, Eve57, Eps20, Vit00, BvL01] and unpublished material from the author. In particular, the references of special import are [G01, Gac94]. The reference [G01] introduces the central quantum matrix μ , which is used to define quantum complexity, quantum mutual information, and to prove properties of quantum measurements. The reference [Gac94] introduces algorithmic (coarse and fine grained) thermodynamic entropy with the key insight that it is the negative logarithm of a universal lower computable test over the phase space. We now present some interesting facts about algorithmic physics that are detailed in this manuscript.

1.1.1 Quantum States Have No Self-Information.

All strings of high Kolmogorov complexity have high self information, with $\mathbf{I}(x : x) =^+ \mathbf{K}(x)$. However the situation is much different in the quantum world, with respect to the definition of mutual information of quantum mixed states σ and ρ introduced in Chapter 6: $\mathbf{I}(\sigma : \rho)$. Almost all pure states $|\psi\rangle$ have low $\mathbf{I}(|\psi\rangle : |\psi\rangle)$. Indeed, let Λ be the uniform distribution over n qubit pure states:

$$\int 2^{\mathbf{I}(|\psi\rangle : |\psi\rangle)} d\Lambda = O(1).$$

This upper bound has several consequences, one being that given a (POVM) measurement, its application to overwhelming majority of quantum states produces white noise, as shown in Chapter 7. In addition, an overwhelming majority of quantum states decohere into random noise. These results are a consequence of the vastness of Hilbert spaces opposed to the limited discretionary power of measurements. Conservation inequalities prevent any type of post-processing of the measured information. As discussed in Chapter 10, the only means to infuse quantum self information is with a projection operation caused by a quantum measurement.

1.1.2 Algorithmic Thermodynamic Entropy and Fluctuations.

Thermodynamic entropy is subject to fluctuations. It will spend most of its time at its maximum value, will exhibit frequent small fluctuations, and rarer large fluctuations. In this manuscript, we show that algorithmic fine grained entropy exhibits such oscillations, and even go one step further in proving the existence of synchronized oscillations for discrete dynamics.

The phase space Ω describes all possible states of the dynamic system, such as all the particles momentums and positions. The phase space is paired with (not necessarily probability) computable measure μ that represents the volume of the space. Like classical thermodynamic entropy, algorithmic fine grained entropy is defined with respect to a particular measure μ and phase space, denoted $\mathbf{H}_\mu(x)$ over $x \in \Omega$.

Due to the Liouville theorem, the dynamics of the system are volume invariant. In this manuscript, it is proved that during the course of such dynamics, algorithmic fine grain thermodynamic have oscillations. Small dips in \mathbf{H}_ν are frequent, and larger dips are more rare. We get the following inequalities, where \mathbf{K} is the prefix-free Kolmogorov complexity. This parallels the discrete ergodic transformation case, detailed in Chapter 14.

Let L be the Lebesgue measure over \mathbb{R} , and (\mathcal{X}, μ) be a computable measure space and $\alpha \in \mathcal{X}$ with finite mutual information with the halting sequence. For transformation group G^t acting on \mathcal{X} , there are constants c_1 and c_2 with

$$2^{-n-\mathbf{K}(n)-c_1} < L\{t \in [0, 1] : \mathbf{H}_\mu(G^t \alpha) < \log \mu(\mathcal{X}) - n\} < 2^{-n+c_2}.$$

The above result proves the existence of oscillations in thermodynamic entropy. However the situation becomes more complicated if one were to examine product spaces. If a product state is typical of the space then its marginal entropies cannot oscillate in synch. For every number there is a time where the marginal entropies differ by more than that amount. Thus, marginal algorithmic thermodynamic entropies cannot be synchronized. This result can be found in Chapter 15, where there exists an analogous theorem for discrete ergodic transforms.

Let $(\mathcal{X} \times \mathcal{Y}, \mu \times \nu)$ be a computable product measure space. Let G^t be a transformation group. Let $(\alpha, \beta) \in \mathcal{X} \times \mathcal{Y}$ with $(\alpha^t, \beta^t) = G^t(\alpha, \beta)$. If $\mathbf{H}_\mu(\alpha, \beta) < -\infty$ and (α, β) has finite mutual information with the halting sequence then $\sup_{t \in [0, 1]} |\mathbf{H}_\mu(\alpha^t) - \mathbf{H}_\nu(\beta^t)| = \infty$.

1.1.3 Typicality and Newtonian Physics

This manuscript has the first, to the author's knowledge, application of algorithmic information theory to Newtonian Physics. Given a system representing a finite number of mass points, an infinite measure κ is defined equal to the magnitude of the gravitational vector field. In addition, a universal lower computable κ -test \mathbf{T}_κ is defined that represents an atypicality score of points in this space. Points at the center of mass points have infinite atypicality, and somewhat surprisingly, \mathbf{T}_κ functions like a typical universal test over compact spaces with suitable computability properties. This machinery can be applied to orbits, which are one dimensional rings around mass points. As a object orbits a body of large mass, it will oscillate in algorithmic typicality, much like with oscillations of algorithmic fine grained entropy over dynamics.

Let L be the one-dimensional Lebesgue measure. Let κ be a computable system with mass point $z \in \mathbb{R}^3$ and O be an orbit centered at z whose encoding has finite mutual information with the halting sequence. There is a $c \in \mathbb{N}$ where for all $n \in \mathbb{N}$,

$$2^{-n-\mathbf{K}(n)-c} < L\{x : x \in O, \mathbf{T}_\kappa(x) > 2^n\}.$$

The above result shows that non-exotic orbits will oscillate in typicality. The following result extends this result to two orbits. Given two orbits with two starting points that are not atypical or exotic, the typicality of the points will be out of sync as they orbit the mass points.

Let κ be a computable system and $O_1 \ni x_1$ and $O_2 \ni x_2$ be two orbits. If the points and encoding of the orbits $(x_1, x_2, \text{Enc}(O_1), \text{Enc}(O_2))$ is typical of the space $\mathbb{R}^3 \times \mathbb{R}^3 \times \{0, 1\}^\infty \times \{0, 1\}^\infty$ and has finite information with the halting sequence then given two transformation groups G_1^t and G_2^t around O_1 and O_2 , we have $\sup_{t \in [0, 1]} |\mathbf{T}_\kappa(G_1^t x_1) - \mathbf{T}_\kappa(G_2^t x_2)| = \infty$.

1.2 Conventions

The following section details the conventions in algorithmic information theory, which will be used in both the quantum mechanics section and the thermodynamics section. We use \mathbb{N} , \mathbb{Z} , \mathbb{Q} , \mathbb{R} , $\{0, 1\}$, $\{0, 1\}^*$, and $\{0, 1\}^\infty$ to denote natural numbers, integers, rational numbers, reals, bits, finite strings, and infinite sequences. $\{0, 1\}^{*\infty} \stackrel{\text{def}}{=} \{0, 1\}^* \cup \{0, 1\}^\infty$. $\|x\|$ denotes the length of the string. Let $X_{\geq 0}$ and $X_{> 0}$ be the sets of non-negative and of positive elements of X . $[A] \stackrel{\text{def}}{=} 1$ if statement A holds, else $[A] \stackrel{\text{def}}{=} 0$. The set of finite bit-strings is denoted by $\{0, 1\}^*$. For set of strings $A \subseteq \{0, 1\}^*$, $\llbracket A \rrbracket = \{x\alpha : x \in A, \alpha \in \{0, 1\}^\infty\}$. When it is clear from the context, we will use natural numbers and other finite objects interchangeably with their binary representations.

The i th bit of $\alpha \in \{0, 1\}^{*\infty}$ is denoted α_i , and its n bit prefix is denoted $\alpha_{\leq n}$. $\langle x \rangle \in \{0, 1\}^*$ for $x \in \{0, 1\}^*$ is the self-delimiting code that doubles every bit of x and changes the last bit of the result. For positive real functions f , by $<^+ f$, $>^+ f$, $=^+ f$, and $<^{\log} f$, $>^{\log} f$, $\sim f$ we denote $\leq f + O(1)$, $\geq f - O(1)$, $= f \pm O(1)$ and $\leq f + O(\log(f+1))$, $\geq f - O(\log(f+1))$, $= f \pm O(\log(f+1))$. Furthermore, $<^* f$, $>^* f$ denotes $< O(1)f$ and $> f/O(1)$. The term $\overset{*}{=} f$ is used to denote $>^* f$ and $<^* f$.

A probability measure Q over $\{0, 1\}^*$ is elementary if it has finite support and range that is a subset of rationals. Elementary probability measures can be encoded into finite strings $\langle Q \rangle$ in the standard way.

1.2.1 Algorithmic Information Theory

$T_y(x)$ is the output of algorithm T (or \perp if it does not halt) on input $x \in \{0, 1\}^*$ and auxiliary input $y \in \{0, 1\}^{*\infty}$. T is prefix-free if for all $x, s \in \{0, 1\}^*$ with $s \neq \emptyset$, either $T_y(x) = \perp$ or $T_y(xs) = \perp$. The complexity of $x \in \{0, 1\}^*$ with respect to T_y is $\mathbf{K}_T(x|y) \stackrel{\text{def}}{=} \inf\{\|p\| : T_y(p) = x\}$.

There exist optimal for \mathbf{K} prefix-free algorithms U , meaning that for all prefix-free algorithms T , there exists $c_T \in \mathbb{N}$, where $\mathbf{K}_U(x|y) \leq \mathbf{K}_T(x|y) + c_T$ for all $x \in \{0, 1\}^*$ and $y \in \{0, 1\}^{*\infty}$. For example, one can take a universal prefix-free algorithm U , where for each prefix-free algorithm T , there exists $t \in \{0, 1\}^*$, with $U_y(tx) = T_y(x)$ for all $x \in \{0, 1\}^*$ and $y \in \{0, 1\}^{*\infty}$. $\mathbf{K}(x|y) \stackrel{\text{def}}{=} \mathbf{K}_U(x|y)$ is the Kolmogorov complexity of $x \in \{0, 1\}^*$ relative to $y \in \{0, 1\}^{*\infty}$.

The chain rule is $\mathbf{K}(x, y) =^+ \mathbf{K}(x) + \mathbf{K}(y|x, \mathbf{K}(x))$. The algorithmic probability is $\mathbf{m}(x|y) = \sum\{2^{-\|p\|} : U_y(p) = x\}$. By the coding theorem $\mathbf{K}(x|y) =^+ -\log \mathbf{m}(x|y)$. The amount of mutual

information between two strings x and y is $\mathbf{I}(x : y) = \mathbf{K}(x) + \mathbf{K}(y) - \mathbf{K}(x, y)$. By the chain rule $\mathbf{K}(x, y) =^+ \mathbf{K}(x) + \mathbf{K}(y|x, \mathbf{K}(x))$. The halting sequence $\mathcal{H} \in \{0, 1\}^\infty$ is the infinite string where $\mathcal{H}_i \stackrel{\text{def}}{=} [U(i) \text{ halts}]$ for all $i \in \mathbb{N}$. The amount of information that \mathcal{H} has about $x \in \{0, 1\}^*$ is $\mathbf{I}(x; \mathcal{H}) = \mathbf{K}(x) - \mathbf{K}(x|\mathcal{H})$. The randomness deficiency of $x \in \{0, 1\}^*$ with respect to elementary probability P over $\{0, 1\}^*$ is $\mathbf{d}(x|P) = \lfloor -\log P(x) - \mathbf{K}(x|\langle P \rangle) \rfloor$. We say $t : \{0, 1\}^* \rightarrow \mathbb{R}_{\geq 0}$ is a P -test, for some probability P , if $\sum_x t(x)P(x) \leq 1$. Let \mathbf{t}_P be a universal lower computable P -test, where for any other lower computable P -test t , $\mathbf{t}_P(x) \geq^* \mathbf{m}(t)t(x)$. Then by the universality of the deficiency of randomness, [G01], $\mathbf{d}(x|P) =^+ \log \mathbf{t}_P(x)$. The transform of a probability Q by $f : \{0, 1\}^* \rightarrow \{0, 1\}^*$, is the probability fQ , where $fQ(x) = \sum_{f(y)=x} Q(y)$. Both randomness deficiency and information enjoy conservation inequalities.

Theorem 1 (See [G01]) $\mathbf{d}(f(x)|fQ) <^+ \mathbf{d}(x|Q)$.

Theorem 2 ([Lev84]) $\mathbf{I}(f(x) : y) <^+ \mathbf{I}(x : y)$.

Proof. Due to the chain rule, $\mathbf{K}(x, y, z) <^+ \mathbf{K}(x, \mathbf{K}(x)) + \mathbf{K}(y|x, \mathbf{K}(x)) + \mathbf{K}(z|x, \mathbf{K}(x)) =^+ \mathbf{K}(x, y) + \mathbf{K}(x, z) - \mathbf{K}(x)$, since $\mathbf{K}(y, z|t) <^+ \mathbf{K}(y|t) + \mathbf{K}(z|t)$. So $\mathbf{I}((z, x) : y) >^+ \mathbf{I}(x : y)$. The statement follows from $\mathbf{I}(z : y) =^+ \mathbf{I}((z, x) : y)$ for $x = A(z)$ since z and $(z, A(z))$ are computable from each other.

Lemma 1 ([Eps22]) $\mathbf{I}(f(a); \mathcal{H}) <^+ \mathbf{I}(a; \mathcal{H}) + \mathbf{K}(f)$.

Proof.

$$\mathbf{I}(a; \mathcal{H}) = \mathbf{K}(a) - \mathbf{K}(a|\mathcal{H}) >^+ \mathbf{K}(a, f(a)) - \mathbf{K}(a, f(a)|\mathcal{H}) - \mathbf{K}(f).$$

The chain rule applied twice results in

$$\begin{aligned} \mathbf{I}(a; \mathcal{H}) + \mathbf{K}(f) &>^+ \mathbf{K}(f(a)) + \mathbf{K}(a|f(a), \mathbf{K}(f(a))) - (\mathbf{K}(f(a)|\mathcal{H}) + \mathbf{K}(a|f(a), \mathbf{K}(f(a)|\mathcal{H}), \mathcal{H})) \\ &=^+ \mathbf{I}(f(a); \mathcal{H}) + \mathbf{K}(a|f(a), \mathbf{K}(f(a))) - \mathbf{K}(a|f(a), \mathbf{K}(f(a)|\mathcal{H}), \mathcal{H}) \\ &=^+ \mathbf{I}(f(a); \mathcal{H}) + \mathbf{K}(a|f(a), \mathbf{K}(f(a))) - \mathbf{K}(a|f(a), \mathbf{K}(f(a)), \mathbf{K}(f(a)|\mathcal{H}), \mathcal{H}) \\ &>^+ \mathbf{I}(f(a); \mathcal{H}). \end{aligned}$$

□

The stochasticity of a string $x \in \{0, 1\}^*$ is $\mathbf{Ks}(x) = \min_{\text{Elementary } Q} \mathbf{K}(Q) + 3 \log \max\{\mathbf{d}(x|Q), 1\}$. Strings with high stochasticity measures are exotic, in that they have high mutual information with the halting sequence. A proof to the following result can be found in Lemma 28 of Appendix B.

Lemma 2 ([Lev16, Eps21b]) $\mathbf{Ks}(x) <^{\log} \mathbf{I}(x; \mathcal{H})$.

The following definition is from [Lev74].

Definition 1 (Information) For infinite sequences $\alpha, \beta \in \{0, 1\}^\infty$, their mutual information is defined to be $\mathbf{I}(\alpha : \beta) = \log \sum_{x, y \in \{0, 1\}^*} 2^{\mathbf{I}(x:y) - \mathbf{K}(x|\alpha) - \mathbf{K}(y|\beta)}$.

It is easy to see that $\mathbf{I}(f(\alpha) : \beta) <^+ \mathbf{I}(\alpha : \beta) + \mathbf{K}(f)$.

Definition 2 (Randomness Deficiency of Infinite Sequences) The deficiency of randomness of an infinite sequence $\alpha \in \{0, 1\}^\infty$, relative to $x \in \{0, 1\}^{*\infty}$ and (not necessarily probability) measure μ over $\{0, 1\}^\infty$ is

$$\mathbf{D}(\alpha|\mu, x) = \sup_n -\log \mu(\alpha[0 \dots n]) - \mathbf{K}(\alpha[0 \dots n]|\mu, x).$$

We say $\mathbf{D}(\alpha|\mu) = \mathbf{D}(\alpha|\mu, \emptyset)$.

1.2.2 Algorithmic Information Between Probabilities

We can generalize from information from strings to information between arbitrary probability measures over strings.

Definition 3 (Information, Probabilities)

For semi-measures p and q over $\{0, 1\}^*$, $\mathbf{I}_{\text{Prob}}(p : q) = \log \sum_{x, y \in \{0, 1\}^*} 2^{\mathbf{I}(x:y)} p(x) q(y)$.

Definition 4 (Channel) A channel $f : \{0, 1\}^* \times \{0, 1\}^* \rightarrow \mathbb{R}_{\geq 0}$ has $f(\cdot|x)$ being a probability measure over $\{0, 1\}^*$ for each $x \in \{0, 1\}^*$. For probability p , channel f , $fp(x) = \sum_z f(x|z)p(z)$.

Lemma 3 Let ψ_a be an enumerable semi-measure, semi-computable relative to a .

$$\sum_c 2^{\mathbf{I}(\langle a, c \rangle : b)} \psi_a(c) \stackrel{*}{<} 2^{\mathbf{I}(a:b)} / \mathbf{m}(\psi).$$

Proof. This requires a slight modification of the proof of Proposition 2 in [Lev84], by requiring ψ to have a as auxilliary information. For completeness, we reproduce the proof. We need to show $\mathbf{m}(a, b) / (\mathbf{m}(a) \mathbf{m}(b)) \stackrel{*}{>} \sum_c (\mathbf{m}(a, b, c) / (\mathbf{m}(b) \mathbf{m}(a, c))) \mathbf{m}(\psi) \psi_a(c)$, or $\sum_c (\mathbf{m}(a, b, c) / \mathbf{m}(a, c)) \mathbf{m}(c|a) \stackrel{*}{<} \mathbf{m}(a, b) / \mathbf{m}(a)$, since $\mathbf{m}(c|a) \stackrel{*}{>} \mathbf{m}(\psi) \psi_a(c)$. Rewrite it $\sum_c \mathbf{m}(c|a) \mathbf{m}(a, b, c) / \mathbf{m}(a, c) \stackrel{*}{<} \mathbf{m}(a, b) / \mathbf{m}(a)$ or $\sum_c \mathbf{m}(c|a) \mathbf{m}(a) \mathbf{m}(a, b, c) / \mathbf{m}(a, c) \stackrel{*}{<} \mathbf{m}(a, b)$. The latter is obvious since $\mathbf{m}(c|a) \mathbf{m}(a) \stackrel{*}{<} \mathbf{m}(a, c)$ and $\sum_c \mathbf{m}(a, b, c) \stackrel{*}{<} \mathbf{m}(a, b)$. \square

Theorem 3 For probabilities p and q over $\{0, 1\}^*$, computable channel f , $\mathbf{I}_{\text{Prob}}(fp : q) <^+ \mathbf{I}_{\text{Prob}}(p : q)$.

Proof. Using Lemma 4,

$$\mathbf{I}_{\text{Prob}}(fp : q) = \log \sum_{x, y} 2^{\mathbf{I}(x:y)} \sum_z f(x|z) p(z) q(y) <^+ \log \sum_{y, z} q(y) p(z) \sum_x 2^{\mathbf{I}((x, z):y)} f(x|z).$$

Using Lemma 5,

$$\mathbf{I}_{\text{Prob}}(fp : q) <^+ \log \sum_{z, y} q(y) p(z) 2^{\mathbf{I}(z:y)} =^+ \mathbf{I}_{\text{Prob}}(p : q).$$

\square

Thus processing cannot increase information between two probabilities. If the the probability measure is concentrated at a single point, then it contains self-information equal to the complexity of that point. If the probability measure is spread out, then it is white noise, and contains no self-information. Some examples are as follows.

Example 1

- In general, a probability p , will have low $\mathbf{I}_{\text{Prob}}(p : p)$ if it has large measure on simple strings, or low measure on a large number of complex strings, or some combination of the two.
- If probability p is concentrated on a single string x , then $\mathbf{I}_{\text{Prob}}(p : p) = \mathbf{K}(x)$.
- The uniform distribution over strings of length n has self information equal to (up to an additive constant) $\mathbf{K}(n)$.
- There are semi-measures that have infinite self information. Let α_n be the n bit prefix of a Martin L f random sequence α and $n \in [2, \infty)$. Semi-measure $p(x) = [x = \alpha_n] n^{-2}$ has $\mathbf{I}_{\text{Prob}}(p : p) = \infty$.

- The universal semi-measure \mathbf{m} has no self information.

Example 2 (Uniform Spread) *An example channel f has $f(\cdot|x)$ be the uniform distribution over strings of length $\|x\|$. This is a canonical spread function. Thus if p is a probability measure concentrated on a single string, then $\mathbf{I}_{\text{Prob}}(p : p) = \mathbf{K}(x)$, and $\mathbf{I}(fp : fp) =^+ \mathbf{K}(\|x\|)$. Thus f results in a decrease of self-information of p . This decrease of information occurs over all probabilities and computable channels.*

Part II

Quantum Mechanics

Chapter 2

Introduction

Classical information theory studies the communication of bits across a noisy channel. Quantum Information Theory (QIT) studies the kind of information (“quantum information”) which is transmitted by microparticles from a preparation device (sender) to a measuring apparatus (receiver) in a quantum mechanical experiment—in other words, the distinction between carriers of classical and quantum information becomes essential. The notion of a qubit can be defined at an abstract level, without giving preference to any particular physical system such as a spin-1/2 particle or a photon. Qubits behave very differently than bits. To start, qubits can be in a linear superposition between 0 and 1. Qubits can have entanglement, where two objects at a distance become a single entity. The study of entanglement and in particular the question how it can be quantified is therefore a central topic within quantum information theory. However, due to the no-cloning theorem [WZ82], instant communication is not possible. Some other aspects of QIT are as follows.

1. **Quantum Computing:** includes hardware (quantum computers), software, algorithm such as Shor’s factoring algorithm or Grover’s algorithm, and applications.
2. **Quantum Communication:** quantum networking, quantum internet, quantum cryptography.
3. **Applications in Physics:** applications to convex optimizations, black holes, and exotic quantum phases of matter.
4. **Quantum Shannon Theory:** quantum channels, quantum protocols, quantum information and entropy.

One aspect of Quantum Shannon Theory (QST) that has had relatively little study is its relationship to Algorithmic Information Theory (AIT). AIT, in part, is the study of the information content of individual strings. A string is random if it cannot be compressed with respect to a universal Turing machine. This paper surveys the current state of research of QST and AIT and provides unpublished results from the author. Hopefully it will convince the reader that there is a fruitful area of research of QST and AIT. Some areas of this intersection include algorithmic content of quantum states, how typical a quantum state is with respect to a quantum source, and how to quantify the algorithmic content of a measurement. One can also gain further insight into quantum transformations, such as purification, decoherence, and approximations to quantum cloning.

As this survey will show, there are some aspects of AIT that directly transfer over to quantum mechanics. This includes comparable definitions of complexity, and conservation inequalities. In addition, there exist quantum versions of the EL Theorem, [Lev16, Eps19c] and the Outlier Theorem, [Eps21b]. However there are some aspects of AIT that are different in the context of quantum

mechanics. This includes the fact the self information of most quantum pure states is zero, with $\mathbf{I}(|\psi\rangle : \psi) \approx 0$. This has implications on the algorithmic content of measurements and decoherence. The main quantum mechanical areas covered in this manuscript are

- **Chapter 3:** This chapter covers the background material needed for the article. A new definition, the algorithmic information between probabilities, is introduced and shown to have information non-growth properties with respect to randomized processing.
- **Chapter 4:** Three different algorithmic measures of quantum states are covered. Their properties are described, including an addition inequality, a Quantum EL Theorem, and a generalized no-cloning theorem. Multiple relationships between the complexities are proven.
- **Chapter 5:** The notion of the algorithmic typicality of one quantum state with respect to another quantum state is introduced. Typicality is conserved with respect to quantum operations. A quantum outlier theorem is proven. This states that non-exotic projections must have atypical pure states in their images.
- **Chapter 6:** The definition of quantum algorithmic information is introduced. Quantum information differs from classical algorithmic information in that an overwhelming majority of pure states have negligible self-information. Information is conserved over quantum operations, with implications to quantum cloning, quantum decoherence, and purification.
- **Chapter 7:** Quantum algorithmic information upper bounds the amount of classical information produced by quantum measurements. Given a quantum measurement, for an overwhelming majority of pure states, the measurement will be random noise. An overwhelming majority of quantum pure states, when undertaking decoherence, will result in a classical probability with no algorithmic information.
- **Chapter 8:** A quantum equivalent to Martin L f random sequence is introduced. Such quantum random states have incompressible initial segments with respect to a new measure quantum complexity called Quantum Operation Complexity. This complexity term measures the cost of approximating a state with a classical and quantum component.
- **Chapter 9:** This chapter shows the Many Worlds Theory and AIT are in conflict, as shown through the existence of a finite experiment that measures the spin of a large number of electrons. After the experiment there are branches of positive probability which contain forbidden sequences that break the Independence Postulate, a postulate in AIT.
- **Chapter 10:** This chapter concludes the quantum mechanical section of the manuscript with a discussion of the boundary between quantum information and classical information. We show that measurements are necessary to produce distributions over quantum states that have cloneable information.
- **Appendix A:** Properties of the quantum information of basis states are proven.
- **Appendix B:** An extended coding theorem is proved with applications to proving inequalities of quantum complexities and the relation between dynamics and coarse grained entropy.

Chapter 3

Background

The reader is assumed to be familiar with both algorithmic information theory and quantum information theory, but we review the core terms. We use \mathbb{N} , \mathbb{Z} , \mathbb{Q} , \mathbb{R} , $\{0, 1\}$, $\{0, 1\}^*$, and $\{0, 1\}^\infty$ to denote natural numbers, integers, rational numbers, reals, bits, finite strings, and infinite sequences. $\{0, 1\}^{*\infty} \stackrel{\text{def}}{=} \{0, 1\}^* \cup \{0, 1\}^\infty$. $\|x\|$ denotes the length of the string. Let $X_{\geq 0}$ and $X_{> 0}$ be the sets of non-negative and of positive elements of X . $[A] \stackrel{\text{def}}{=} 1$ if statement A holds, else $[A] \stackrel{\text{def}}{=} 0$. The set of finite bit-strings is denoted by $\{0, 1\}^*$. For set of strings $A \subseteq \{0, 1\}^*$, $\llbracket A \rrbracket = \{x\alpha : x \in A, \alpha \in \{0, 1\}^\infty\}$. When it is clear from the context, we will use natural numbers and other finite objects interchangeably with their binary representations.

The i th bit of $\alpha \in \{0, 1\}^{*\infty}$ is denoted α_i , and its n bit prefix is denoted $\alpha_{\leq n}$. $\langle x \rangle \in \{0, 1\}^*$ for $x \in \{0, 1\}^*$ is the self-delimiting code that doubles every bit of x and changes the last bit of the result. For positive real functions f , by $<^+ f$, $>^+ f$, $=^+ f$, and $<^{\log} f$, $>^{\log} f$, $\sim f$ we denote $\leq f + O(1)$, $\geq f - O(1)$, $= f \pm O(1)$ and $\leq f + O(\log(f+1))$, $\geq f - O(\log(f+1))$, $= f \pm O(\log(f+1))$. Furthermore, $<^* f$, $>^* f$ denotes $< O(1)f$ and $> f/O(1)$. The term $\stackrel{*}{=} f$ is used to denote $>^* f$ and $<^* f$.

A probability measure Q over $\{0, 1\}^*$ is elementary if it has finite support and range that is a subset of rationals. Elementary probability measures can be encoded into finite strings $\langle Q \rangle$ in the standard way.

3.1 Algorithmic Information Theory

$T_y(x)$ is the output of algorithm T (or \perp if it does not halt) on input $x \in \{0, 1\}^*$ and auxiliary input $y \in \{0, 1\}^{*\infty}$. T is prefix-free if for all $x, s \in \{0, 1\}^*$ with $s \neq \emptyset$, either $T_y(x) = \perp$ or $T_y(xs) = \perp$. The complexity of $x \in \{0, 1\}^*$ with respect to T_y is $\mathbf{K}_T(x|y) \stackrel{\text{def}}{=} \inf\{\|p\| : T_y(p) = x\}$.

There exist optimal for \mathbf{K} prefix-free algorithms U , meaning that for all prefix-free algorithms T , there exists $c_T \in \mathbb{N}$, where $\mathbf{K}_U(x|y) \leq \mathbf{K}_T(x|y) + c_T$ for all $x \in \{0, 1\}^*$ and $y \in \{0, 1\}^{*\infty}$. For example, one can take a universal prefix-free algorithm U , where for each prefix-free algorithm T , there exists $t \in \{0, 1\}^*$, with $U_y(tx) = T_y(x)$ for all $x \in \{0, 1\}^*$ and $y \in \{0, 1\}^{*\infty}$. $\mathbf{K}(x|y) \stackrel{\text{def}}{=} \mathbf{K}_U(x|y)$ is the Kolmogorov complexity of $x \in \{0, 1\}^*$ relative to $y \in \{0, 1\}^{*\infty}$.

The algorithmic probability is $\mathbf{m}(x|y) = \sum\{2^{-\|p\|} : U_y(p) = x\}$. By the coding theorem $\mathbf{K}(x|y) =^+ -\log \mathbf{m}(x|y)$. The amount of mutual information between two strings x and y is $\mathbf{I}(x : y) = \mathbf{K}(x) + \mathbf{K}(y) - \mathbf{K}(x, y)$. By the chain rule $\mathbf{K}(x, y) =^+ \mathbf{K}(x) + \mathbf{K}(y|x, \mathbf{K}(x))$. The halting sequence $\mathcal{H} \in \{0, 1\}^\infty$ is the infinite string where $\mathcal{H}_i \stackrel{\text{def}}{=} [U(i) \text{ halts}]$ for all $i \in \mathbb{N}$. The amount of information that \mathcal{H} has about $x \in \{0, 1\}^*$ is $\mathbf{I}(x; \mathcal{H}) = \mathbf{K}(x) - \mathbf{K}(x|\mathcal{H})$. The randomness deficiency of $x \in \{0, 1\}^*$ with respect to elementary probability P over $\{0, 1\}^*$ is $\mathbf{d}(x|P) =$

$\lfloor -\log P(x) - \mathbf{K}(x|\langle P \rangle) \rfloor$. we say $t : \{0,1\}^* \rightarrow \mathbb{R}_{\geq 0}$ is a P -test, for some probability P , if $\sum_x t(x)P(x) \leq 1$. Let \mathbf{t}_P be a universal lower computable P -test, where for any other lower computable P -test t , $\mathbf{t}_P(x) \stackrel{*}{>} \mathbf{m}(t)t(x)$. Then by the universality of the deficiency of randomness, [G01], $\mathbf{d}(x|P) =^+ \log \mathbf{t}_P(x)$. The transform of a probability Q by $f : \{0,1\}^* \rightarrow \{0,1\}^*$, is the probability fQ , where $fQ(x) = \sum_{f(y)=x} Q(y)$. Both randomness deficiency and information enjoy conservation inequalities.

Theorem 4 (See [G01]) $\mathbf{d}(f(x)|fQ) <^+ \mathbf{d}(x|Q)$.

Theorem 5 ([Lev84]) $\mathbf{I}(f(x) : y) <^+ \mathbf{I}(x : y)$.

Lemma 4 ([Eps22]) $\mathbf{I}(f(a); \mathcal{H}) <^+ \mathbf{I}(a; \mathcal{H}) + \mathbf{K}(f)$.

Lemma 5 Let ψ_a be an enumerable semi-measure, semi-computable relative to a .
 $\sum_c 2^{\mathbf{I}(\langle a,c \rangle : b)} \psi_a(c) <^* 2^{\mathbf{I}(a:b)} / \mathbf{m}(\psi)$.

Proof. This requires a slight modification of the proof of Proposition 2 in [Lev84], by requiring ψ to have a as auxilliary information. For completeness, we reproduce the proof. We need to show $\mathbf{m}(a,b)/(\mathbf{m}(a)\mathbf{m}(b)) \stackrel{*}{>} \sum_c (\mathbf{m}(a,b,c)/(\mathbf{m}(b)\mathbf{m}(a,c))) \mathbf{m}(\psi)\psi_a(c)$, or $\sum_c (\mathbf{m}(a,b,c)/\mathbf{m}(a,c)) \mathbf{m}(c|a) <^* \mathbf{m}(a,b)/\mathbf{m}(a)$, since $\mathbf{m}(c|a) \stackrel{*}{>} \mathbf{m}(\psi)\psi_a(c)$. Rewrite it $\sum_c \mathbf{m}(c|a)\mathbf{m}(a,b,c)/\mathbf{m}(a,c) <^* \mathbf{m}(a,b)/\mathbf{m}(a)$ or $\sum_c \mathbf{m}(c|a)\mathbf{m}(a)\mathbf{m}(a,b,c)/\mathbf{m}(a,c) <^* \mathbf{m}(a,b)$. The latter is obvious since $\mathbf{m}(c|a)\mathbf{m}(a) <^* \mathbf{m}(a,c)$ and $\sum_c \mathbf{m}(a,b,c) <^* \mathbf{m}(a,b)$. \square

The stochasticity of a string $x \in \{0,1\}^*$ is $\mathbf{Ks}(x) = \min_{\text{Elementary } Q} \mathbf{K}(Q) + 3 \log \max\{\mathbf{d}(x|Q), 1\}$. Strings with high stochasticity measures are exotic, in that they have high mutual information with the halting sequence.

Lemma 6 ([Lev16, Eps21b]) $\mathbf{Ks}(x) <^{\log} \mathbf{I}(x; \mathcal{H})$.

3.1.1 Algorithmic Information Between Probabilities

We can generalize from information from strings to information between arbitrary probability measures over strings.

Definition 5 (Information, Probabilities)

For semi-measures p and q over $\{0,1\}^*$, $\mathbf{I}_{\text{Prob}}(p : q) = \log \sum_{x,y \in \{0,1\}^*} 2^{\mathbf{I}(x:y)} p(x)q(y)$.

Definition 6 (Channel) A channel $f : \{0,1\}^* \times \{0,1\}^* \rightarrow \mathbb{R}_{\geq 0}$ has $f(\cdot|x)$ being a probability measure over $\{0,1\}^*$ for each $x \in \{0,1\}^*$. For probability p , channel f , $fp(x) = \sum_z f(x|z)p(z)$.

Theorem 6 For probabilities p and q over $\{0,1\}^*$, computable channel f , $\mathbf{I}_{\text{Prob}}(fp : q) <^+ \mathbf{I}_{\text{Prob}}(p : q)$.

Proof. Using Lemma 4,

$$\mathbf{I}_{\text{Prob}}(fp : q) = \log \sum_{x,y} 2^{\mathbf{I}(x:y)} \sum_z f(x|z)p(z)q(y) <^+ \log \sum_{y,z} q(y)p(z) \sum_x 2^{\mathbf{I}((x,z):y)} f(x|z).$$

Using Lemma 5,

$$\mathbf{I}_{\text{Prob}}(fp : q) <^+ \log \sum_{z,y} q(y)p(z)2^{\mathbf{I}(z:y)} =^+ \mathbf{I}_{\text{Prob}}(p : q).$$

□

Thus processing cannot increase information between two probabilities. If the the probability measure is concentrated at a single point, then it contains self-information equal to the complexity of that point. If the probability measure is spread out, then it is white noise, and contains no self-information. Some examples are as follows.

Example 3

- In general, a probability p , will have low $\mathbf{I}_{\text{Prob}}(p : p)$ if it has large measure on simple strings, or low measure on a large number of complex strings, or some combination of the two.
- If probability p is concentrated on a single string x , then $\mathbf{I}_{\text{Prob}}(p : p) = \mathbf{K}(x)$.
- The uniform distribution over strings of length n has self information equal to (up to an additive constant) $\mathbf{K}(n)$.
- There are semi-measures that have infinite self information. Let α_n be the n bit prefix of a Martin L f random sequence α and $n \in [2, \infty)$. Semi-measure $p(x) = [x = \alpha_n]n^{-2}$ has $\mathbf{I}_{\text{Prob}}(p : p) = \infty$.
- The universal semi-measure \mathbf{m} has no self information.

Example 4 (Uniform Spread) An example channel f has $f(\cdot|x)$ be the uniform distribution over strings of length $\|x\|$. This is a canonical spread function. Thus if p is a probability measure concentrated on a single string, then $\mathbf{I}_{\text{Prob}}(p : p) = \mathbf{K}(x)$, and $\mathbf{I}(fp : fp) =^+ \mathbf{K}(\|x\|)$. Thus f results in a decrease of self-information of p . This decrease of information occurs over all probabilities and computable channels.

3.2 Quantum Mechanics

We use the standard model of qubits used throughout quantum information theory. We deal with finite N dimensional Hilbert spaces \mathcal{H}_N , with bases $|\alpha_1\rangle, |\alpha_2\rangle, \dots, |\alpha_n\rangle$. We assume $\mathcal{H}_{n+1} \supseteq \mathcal{H}_n$ and the bases for \mathcal{H}_n are the beginning of that of \mathcal{H}_{n+1} . An n qubit space is denoted by $\mathcal{Q}_n = \bigotimes_{i=1}^n \mathcal{Q}_1$, where qubit space \mathcal{Q}_1 has bases $|0\rangle$ and $|1\rangle$. For $x \in \Sigma^n$ we use $|x\rangle \in \mathcal{Q}_n$ to denote $\bigotimes_{i=1}^n |x[i]\rangle$. The space \mathcal{Q}_n has 2^n dimensions and we identify it with \mathcal{H}_{2^n} .

A pure quantum state $|\phi\rangle$ of length n is represented as a unit vector in \mathcal{Q}_n . Its corresponding element in the dual space is denoted by $\langle\phi|$. The tensor product of two vectors is denoted by $|\phi\rangle \otimes |\psi\rangle = |\phi\rangle |\psi\rangle = |\phi\psi\rangle$. The inner product of $|\psi\rangle$ and $\langle\phi|$ is denoted by $\langle\psi|\phi\rangle$.

The symbol Tr denotes the trace operation. The conjugate transpose of a matrix M is denoted by M^* . For Hermitian matrix with eigenvalue decomposition $A = \sum a_i |\psi_i\rangle \langle\psi_i|$, $|A| = \sum |a_i| |\psi_i\rangle \langle\psi_i|$. The tensor product of two matrices is denoted by $A \otimes B$. Projection matrices are Hermitian matrices with eigenvalues in $\{0, 1\}$. For tensor product space $\mathcal{G}_X \otimes \mathcal{G}_Y$, the partial trace is denoted by Tr_Y . For $B^X = \text{Tr}_Y B$, $\text{Tr}(A \cdot B^X) = \text{Tr}((A \otimes I) \cdot B)$, which is used frequently throughout the paper. For positive semidefinite matrices, σ and ρ we say $\sigma \leq \rho$ if $\rho - \sigma$ is positive semidefinite. For positive semidefinite matrices A, B, C , if $A \leq B$ then $\text{Tr}AC \leq \text{Tr}BC$. Mixed states are represented by density matrices, which are, self adjoint, positive semidefinite, operators

of trace 1. A semi-density matrix has non-negative trace less than or equal to 1. The von Neumann entropy of a density matrix σ with orthogonal decomposition $\sum p_i |\psi_i\rangle \langle \psi_i|$ is $S(\sigma) = -\sum p_i \log p_i$.

A pure quantum state $|\phi\rangle$ and (semi)density matrix σ are called *elementary* if their real and imaginary components have rational coefficients. Elementary objects can be encoded into strings or integers and be the output of halting programs. Therefore one can use the terminology $\mathbf{K}(|\phi\rangle)$ and $\mathbf{K}(\sigma)$, and also $\mathbf{m}(|\phi\rangle)$ and $\mathbf{m}(\sigma)$.

We say program $q \in \{0, 1\}^*$ lower computes positive semidefinite matrix σ if, given as input to universal Turing machine U , the machine U reads $\leq \|q\|$ bits and outputs, with or without halting, a sequence of elementary semi-density matrices $\{\sigma_i\}$ such that $\sigma_i \leq \sigma_{i+1}$ and $\lim_{i \rightarrow \infty} \sigma_i = \sigma$. A matrix is lower computable if there is a program that lower computes it.

3.2.1 Quantum Operations

A quantum operation is the most general type of operation than can be applied to a quantum state. In Chapters 5 and 6, conservation inequalities will be proven with respect to quantum operations. A map transforming a quantum state σ to $\varepsilon(\sigma)$ is a quantum operation if it satisfies the following three requirements

1. The map of ε is positive and trace preserving, with $\text{Tr}(\sigma) = \text{Tr}(\varepsilon(\sigma))$.
2. The map is linear with $\varepsilon(\sum_i p_i \sigma_i) = \sum_i p_i \varepsilon(\sigma_i)$.
3. The map is completely positive, were any map of the form $\varepsilon \otimes \mathbf{M}$ acting on the extended Hilbert space is also positive.

Another means to describe quantum operations is through a series of operators. A quantum operation ε on mixed state σ_A can be seen as the appending of an ancilla state σ_b , applying a unitary transform U , then tracing out the ancilla system with

$$\varepsilon(\sigma_A) = \text{Tr}_B (U(\sigma_A \otimes \sigma_B)U^*). \quad (3.1)$$

A third way to characterize a quantum operation is through Kraus operators, which can be derived using an algebraic reworking of Equation 3.1. Map ε is a quantum operation iff it can be represented (not necessarily uniquely) using a set of matrices $\{M_i\}$ where $\varepsilon(\sigma) = \sum_i M_i \sigma M_i^*$ and $\sum_i M_i^* M_i \leq I$, where I is the identity matrix.

A quantum operation ε is elementary iff it admits a represented of the form in Equation 3.1 where B , U , and σ_B are each elementary, in that they each can be encoded with a finite string. The encoding of an elementary quantum operation is denoted by $\langle \varepsilon \rangle = \langle B \rangle \langle U \rangle \langle \sigma_B \rangle$. Each elementary quantum operation admits an elementary Kraus operator representation $\{M_i\}$, in that each M_i is an elementary matrix. This elementary Kraus operator is computable from $\langle \varepsilon \rangle$.

Chapter 4

Quantum Complexity

4.1 Three Measures of the Algorithmic Content of Individual Quantum States

The formal study of Algorithmic Information Theory and Quantum Mechanics began with the introduction of three independent measures of the algorithmic content of a mixed or pure quantum state, detailed in the papers [BvL01, G01, Vit01]. BVL complexity [BvL01] measures the complexity of a pure quantum state $|\psi\rangle$ by the length of the smallest input to a universal quantum Turing machine that outputs a good approximation of $|\psi\rangle$. Vitányi complexity [Vit01] measures the entropy of a pure state $|\psi\rangle$ as the amount of classical information needed to reproduce a good approximation of $|\psi\rangle$. Gács complexity measures the entropy of a pure or mixed quantum state by using a quantum analogue of the universal semi-measure \mathbf{m} .

4.1.1 BVL Complexity

BVL complexity, introduced in [BvL01], uses a universal quantum Turing machine to define the complexity of a pure quantum state. The input and output tape of this machine consists of symbols of the type 0, 1, and #. The input is an ensemble $\{p_i\}$ of pure states $|\psi_i\rangle$ of the same length n , where $p_i \geq 0$ and $\sum_i p_i = 1$. This ensemble can be represented as a mixed state of n qubits. If, during the operation of the quantum Turing machine, all computational branches halt at a time t , then the contents on the output tape are considered the output of the quantum Turing machine. The BVL Complexity of a pure state, $\mathbf{Hbvl}[\epsilon](|\psi\rangle)$ is the size of the smallest (possibly mixed state) input to a universal quantum Turing machine such that fidelity between the output and $|\psi\rangle$ is at least ϵ . The fidelity between a mixed state output σ and $|\psi\rangle$ is $\langle\psi|\sigma|\psi\rangle$. We require that the input quantum state be elementary. We also require that universal quantum Turing machine be conditioned on the number of qubits n , on a classical auxiliary tape.

4.1.2 Vitányi Complexity

Vitányi complexity [Vit01] is a measure of the algorithmic information content of a pure state $|\psi\rangle$. It is equal to the minimum of the Kolmogorov complexity of an approximating elementary pure state $|\phi\rangle$ summed with a score of their closeness. We use a slightly different definition than the original [Vit01], in that we use a classical machine and not a quantum machine for the approximation. Let N be the dimension of the Hilbert space.

$$\mathbf{Hv}(|\psi\rangle) = \min_{\text{Elementary } |\theta\rangle \in \mathcal{H}_N} \mathbf{K}(|\theta\rangle | N) - \log |\langle\psi|\theta\rangle|^2.$$

4.1.3 Gács Complexity

Gács complexity [G01] is defined using the following universal lower computable semi-density matrix, parametered by $x \in \{0, 1\}^*$, with

$$\mu_x = \sum_{\text{Elementary } |\phi\rangle \in \mathcal{H}_N} \mathbf{m}(|\phi\rangle |x, N) |\phi\rangle \langle \phi|.$$

The parameter N represents the dimension of the Hilbert space. We use μ_X to denote the matrix μ over the Hilbert space denoted by symbol X . The Gács entropy of a mixed state σ , conditioned on $x \in \{0, 1\}^*$ is defined by

$$\mathbf{Hv}(\sigma|x) = \lceil -\log \text{Tr} \mu_x \sigma \rceil.$$

We use the following notation for pure states, with $\mathbf{Hg}(|\phi\rangle |x) = \mathbf{Hg}(|\phi\rangle \langle \phi| |x)$. For empty x we use the notation $\mathbf{Hg}(\sigma)$. This definition generalizes \underline{H} in [G01] to mixed states. Note that in [G01], there is another measure of quantum algorithmic entropy, \overline{H} , which we will not cover in this paper. An infinite version of algorithmic entropy can be found at [BOD14].

4.2 Properties of Universal Matrix and Gács Complexity

The matrix μ is important in algorithmic information theory and quantum mechanics, as it is the foundation for the information term defined in Chapter 6. The following theorem shows that the lower computable semi-density matrix μ is universal. It is greater than any other lower computable matrix, weighted by their complexity. This parallels the classical case, where universal measure \mathbf{m} majorizes lower computable semi measure p , with $\mathbf{m}(x) \stackrel{*}{>} \mathbf{m}(p)p(x)$. This theorem is used throughout the paper, and will not be explicitly cited.

Theorem 7 ([G01]) *Let $q \in \{0, 1\}^*$, and the dimension of the Hilbert space, N , compute lower compute semi-density matrix A . Then $\mu \stackrel{*}{>} \mathbf{m}(q|N)A$.*

Proof. A can be composed into a sum $\sum_i p(i) |\psi_i\rangle \langle \psi_i|$, where each $|\psi_i\rangle$ is elementary, p is a semi-measure, with $\sum_i p(i) \leq 1$, and p is computable from q . Thus since p is computable from q and N ,

$$A = \sum_i p(i) |\psi_i\rangle \langle \psi_i| \stackrel{*}{<} \mathbf{m}(p|N)^{-1} \sum_i \mathbf{m}(i|N) |\psi_i\rangle \langle \psi_i| \stackrel{*}{<} \mathbf{m}(q|N)^{-1} \sum_i \mathbf{m}(i|N) |\psi_i\rangle \langle \psi_i| \stackrel{*}{<} \mu / \mathbf{m}(q|N).$$

□

Theorem 8 ([G01]) $\mu_{ii} \stackrel{*}{=} \mathbf{m}(i|N)$.

Proof. The matrix $\rho = \sum_i \mathbf{m}(i|N) |i\rangle \langle i|$ is lower computable, so $\rho \stackrel{*}{<} \mu$ so $\mu_{ii} \stackrel{*}{>} \mathbf{m}(i|N)$. Furthermore, $f(i) = |i\rangle \mu \langle i|$ is a lower computable semi-measure, so $\mathbf{m}(i|N) \stackrel{*}{>} \mu_{ii}$. □

Theorem 9 ([G01]) $\text{Tr}_Y \mu_{XY} \stackrel{*}{=} \mu_X$.

Proof. Let $\rho = \text{Tr}_Y \mu_{XY}$, which is a lower computable semi-density matrix because one can enumerate elementary pure states $|\psi\rangle\langle\psi|$ in the space XY , take their partial trace, $\text{Tr}_T |\psi\rangle\langle\psi|$, and add the resulting pure or mixed state to the sum ρ . Thus $\rho <^* \mu_X$. Let $\sigma = \mu_X \otimes |\psi\rangle\langle\psi|$, where $|\psi\rangle$ is a reference elementary state. Thus $\sigma <^* \mu_{XY}$ so

$$\mu_X = \text{Tr}_Y \sigma <^* \text{Tr}_Y \mu_{XY}.$$

□

Theorem 10 ([G01]) $\text{Hg}(\sigma) <^+ \text{Hg}(\sigma \otimes \rho)$.

Proof. Note that this theorem is not less general than that of Theorem 12, because both σ and ρ can be non-elementary. Using Theorem 9 and the properties of partial trace,

$$2^{-\text{Hg}(\sigma)} >^* \text{Tr} \sigma \mu_X >^* \text{Tr} \sigma \text{Tr}_Y \mu_{XY} >^* \text{Tr}(\sigma \otimes I) \mu_{XY} >^* \text{Tr}(\sigma \otimes \rho) \mu_{XY} \stackrel{*}{=} 2^{-\text{Hg}(\sigma \otimes \rho)}.$$

4.3 No Cloning Theorem

In classical algorithmic information theory, one can easily reproduce a string x , with

$$\mathbf{K}(x) =^+ \mathbf{K}(x, x).$$

However the situation is much different in the quantum case. Due to the no-cloning theorem, [WZ82] one cannot clone a quantum state. The following theorem generalizes this no-go result, by showing there exist tensor products $|\psi\rangle^m$ that has significantly more **Hg** measure than $|\psi\rangle$. The following theorem presents a new proof to this result.

Theorem 11 ([G01]) $\log \binom{m+N-1}{m} <^+ \max_{|\psi\rangle} \text{Hg}(|\psi\rangle^{\otimes m}) <^+ \mathbf{K}(m) + \log \binom{m+N-1}{m}$

Proof. Let \mathcal{H}_N be an N dimensional Hilbert space and let \mathcal{H}_N^m be an m -fold tensor space of \mathcal{H}_N . Let $\text{Sym}(\mathcal{H}_N^m)$ be the subspace of \mathcal{H}_N^m consisting of all pure states of the form $|\psi\rangle^{\otimes m}$. The subspace $\text{Sym}(\mathcal{H}_N^m)$ is spanned by M basis vectors, where M is the number of multisets of size m from the set $\{1, \dots, N\}$. This is because for each such multiset $S = \{i_1, \dots, i_m\}$, one can construct a basis vector $|\psi_S\rangle$ that is the normalized superposition of all basis vectors of $\text{Sym}(\mathcal{H}_N^m)$ that are permutations of S . If $S' \neq S$, then $|\psi_S\rangle$ is orthogonal to $|\psi_{S'}\rangle$. Thus the dimension of $\text{Sym}(\mathcal{H}_N^m)$ M , is $\binom{m+N-1}{m}$ because choosing a multiset is the same as splitting an interval of size m into N intervals. For the upper bounds, let P_S be the projector onto $\text{Sym}(\mathcal{H}_N^m)$. If $|\psi\rangle \in \text{Sym}(\mathcal{H}_N^m)$, then $\langle\psi| P_S |\psi\rangle = 1$ so

$$\text{Hg}(|\psi\rangle) <^+ \mathbf{K}(P_S/M | N^m) - \log \langle\psi| \frac{1}{M} P_S / M |\psi\rangle <^+ \mathbf{K}(m) + \log \binom{m+N-1}{m}.$$

For the lower bound, let $c = \max_{|\psi\rangle \in \mathcal{H}_N} \text{Hg}(|\psi\rangle^{\otimes m})$. We have for all $|\psi\rangle \in \mathcal{H}_N$

$$\text{Tr} \mu |\psi\rangle^m \langle\psi|^m >^* 2^{-c}. \quad (4.1)$$

Let Λ be the uniform distribution on the unit sphere of \mathcal{H}_N . And let

$$\rho = \int |\psi\rangle^m \langle\psi|^m d\Lambda.$$

$\text{Tr}\rho = \int \text{Tr} |\psi\rangle^m \langle\psi|^m d\Lambda = \int d\Lambda = 1$. Furthermore for $|\phi\rangle^m, |\nu\rangle^m \in \text{Sym}(\mathcal{H}_N^m)$, with unitary transform U such that $U^m |\psi\rangle^m = |\rho\rangle^m$, we have

$$\langle\nu|^n \rho |\nu\rangle^n = \int \langle\phi|^m (U^{*m} |\psi\rangle^m \langle\psi|^m U^m) |\phi\rangle^m d\Lambda = \int \langle\phi^m | \psi^m \rangle \langle\psi^m | \phi^m \rangle^m d\Lambda = \langle\phi|^n \rho |\phi\rangle^n.$$

For any pure state $|\psi\rangle \in \mathcal{H}_N^m$, such that $\langle\psi| P_S |\psi\rangle = 0$, then $\langle\psi| \rho |\psi\rangle = 0$. Thus $\rho = P_S/M$. Integrating Equation 4.1, by $d\Lambda$ results in

$$2^{-c} <^* \text{Tr} \mu \rho \stackrel{*}{=} \text{Tr} \mu P_S/M \stackrel{*}{=} \binom{m+N-1}{m}^{-1} \\ c >^+ \log \binom{m+N-1}{m}.$$

□

4.4 Addition Inequality

The addition theorem for classical entropy asserts that the joint entropy for a pair of random variables is equal to the entropy of one plus the conditional entropy of the other, with $\mathcal{H}(\mathcal{X}) + \mathcal{H}(\mathcal{Y}|\mathcal{X}) = \mathcal{H}(\mathcal{X}, \mathcal{Y})$. For algorithmic entropy, the chain rule is slightly more nuanced, with $\mathbf{K}(x) + \mathbf{K}(y|x, \mathbf{K}(x)) =^+ \mathbf{K}(x, y)$. An analogous relationship cannot be true for Gács entropy, \mathbf{Hg} , since as shown in Theorem 11, there exists elementary $|\phi\rangle$ where $\mathbf{Hg}(|\phi\rangle |\phi\rangle) - \mathbf{Hg}(|\phi\rangle)$ can be arbitrarily large, and $\mathbf{Hg}(|\phi\rangle / |\phi\rangle) =^+ 0$. However, the following theorem shows that a chain rule inequality does hold for \mathbf{Hg} .

For $n^2 \times n^2$ matrix A , let $A[i, j]$ be the $n \times n$ submatrix of A starting at position $(n(i-1) + 1, n(j-1) + 1)$. For example for $n = 2$ the matrix

$$A = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ 9 & 10 & 11 & 12 \\ 13 & 14 & 15 & 16 \end{bmatrix}$$

has $A[1, 1] = \begin{bmatrix} 1 & 2 \\ 5 & 6 \end{bmatrix}$, $A[1, 2] = \begin{bmatrix} 3 & 4 \\ 7 & 8 \end{bmatrix}$, $A[2, 1] = \begin{bmatrix} 9 & 10 \\ 13 & 14 \end{bmatrix}$, $A[2, 2] = \begin{bmatrix} 11 & 12 \\ 15 & 16 \end{bmatrix}$.

For $n^2 \times n^2$ matrix A and $n \times n$ matrix B , let M_{AB} be the $n \times n$ matrix whose (i, j) entry is equal to $\text{Tr} A[i, j] B$. For any $n \times n$ matrix C , in can be seen that $\text{Tr} A(C \otimes B) = \text{Tr} M_{AB} C$. Furthermore if A is lower computable and B is elementary, then M_{AB} is lower computable.

For elementary semi density matrices ρ , we use $\langle\rho, \mathbf{Hg}(\rho)\rangle$ to denote the encoding of the pair of an encoded ρ and an encoded natural number $\mathbf{Hg}(\rho)$.

Theorem 12 ([Eps19a]) *For semi-density matrices σ, ρ , elementary ρ , $\mathbf{Hg}(\rho) + \mathbf{Hg}(\sigma | \langle\rho, \mathbf{Hg}(\rho)\rangle) <^+ \mathbf{Hg}(\sigma \otimes \rho)$.*

Proof. Let μ_{2n} be the universal lower computable semi density matrix over the space of $2n$ qubits, $\mathcal{Q}_{2n} = \mathcal{Q}_n \otimes \mathcal{Q}_n = \mathcal{Q}_A \otimes \mathcal{Q}_B$. Let μ_n be the universal matrix of the space over n qubits. We define the following bilinear function over complex matrixes of size $n \times n$, with $T(\nu, \delta) = \text{Tr} \mu_{2n}(\nu \otimes \delta)$. For fixed ρ , $T(\nu, \rho)$ is of the form $T(\nu, \rho) = \text{Tr} M_{\mu_{2n}\rho} \nu$. The matrix $M_{\mu_{2n}\rho}$ has trace equal to

$$\begin{aligned} \text{Tr} M_{\mu_{2n}\rho} &= T(\rho, I) \\ &= \text{Tr} \mu_{2n}(\rho \otimes I) \\ &= \text{Tr} ((\text{Tr}_{\mathcal{Q}_B} \mu_{2n}) \rho) \\ &\stackrel{*}{=} \text{Tr} \mu_n \rho \\ &\stackrel{*}{=} 2^{-\mathbf{Hg}(\rho)}, \end{aligned}$$

using Theorem 9, which states $\text{Tr}_Y \mu_{XY} \stackrel{*}{=} \mu_X$. By the definition of M , since μ_{2n} and ρ are positive semi-definite, it must be that $M_{\mu_{2n}\rho}$ is positive semi-definite. Since the trace of $M_{\mu_{2n}\rho}$ is $\stackrel{*}{=} 2^{-\mathbf{Hg}(\rho)}$, it must be that up to a multiplicative constant, $2^{\mathbf{Hg}(\rho)} M_{\mu_{2n}\rho}$ is a semi-density matrix.

Since μ is lower computable and ρ is elementary, by the definition of M , $2^{\mathbf{Hg}(\rho)} M_{\mu_{2n}\rho}$ is lower computable relative to the string $\langle \rho, \mathbf{Hg}(\rho) \rangle$. Therefore we have that $2^{\mathbf{Hg}(\rho)} M_{\mu_{2n}\rho} \stackrel{*}{<} \mu_{\langle \rho, \mathbf{Hg}(\rho) \rangle}$. So we have that $-\log \text{Tr} 2^{\mathbf{Hg}(\rho)} M_{\mu_{2n}\rho} \sigma = -\mathbf{Hg}(\rho) - \log T(\sigma, \rho) \stackrel{+}{=} \mathbf{Hg}(\sigma \otimes \rho) - \mathbf{Hg}(\rho) \stackrel{+}{>} -\log \mu_{\langle \rho, \mathbf{Hg}(\rho) \rangle} \sigma \stackrel{+}{=} \mathbf{Hg}(\sigma | \langle \rho, \mathbf{Hg}(\rho) \rangle)$. \square

4.5 Subadditivity, Strong Subadditivity, Strong Superadditivity

Theorem 13 ([G01]) $\mathbf{Hg}(\sigma)$ is subadditive, with $\mathbf{Hg}(\sigma \otimes \rho) <^+ \mathbf{Hg}(\sigma) + \mathbf{Hg}(\rho)$.

Proof.

$$\begin{aligned} &2^{-\mathbf{Hg}(\sigma) - \mathbf{Hg}(\rho)} \\ &= (\text{Tr} \mu_X \sigma) (\text{Tr} \mu_Y \rho) \\ &= \text{Tr}(\sigma \otimes \rho) (\mu_X \otimes \mu_Y) \\ &\stackrel{*}{>} \text{Tr}(\sigma \otimes \rho) (\mu_{XY}) \\ &\stackrel{*}{=} 2^{-\mathbf{Hg}(\sigma \otimes \rho)}. \end{aligned}$$

\square

A function \mathbf{L} from quantum mixed states to whole numbers is strongly subadditive if there exists a constant $c \in \mathbb{N}$ such that for all mixed states ρ_{123} , $\mathbf{L}(\rho_{123}) + \mathbf{L}(\rho_2) < \mathbf{L}(\rho_{12}) + \mathbf{L}(\rho_{23}) + c$. Similarly \mathbf{L} is strongly superadditive if there exists a constant $c \in \mathbb{N}$ such that for all mixed states ρ_{123} , $\mathbf{L}(\rho_{12}) + \mathbf{L}(\rho_{23}) < \mathbf{L}(\rho_{123}) + \mathbf{L}(\rho_2) + c$.

Theorem 14 \mathbf{Hg} is not strongly subadditive.

Proof. We fix the number of qubits n , and for $i \in [1..2^n]$, $|i\rangle$ is the i th basis state of the n qubit space. Let $|\psi\rangle = \sum_{i=1}^{2^n} 2^{-n/2} |i\rangle |i\rangle$. The pure state $|\psi\rangle$ is elementary, with $\mathbf{K}(|\psi\rangle |2^{2n}) =^+ 0$. We define the the $3n$ qubit mixed state $\rho_{123} = .5 |\psi\rangle \langle \psi| \otimes |1\rangle \langle 1| + .5 |1\rangle \langle 1| \otimes |\psi\rangle \langle \psi|$. $\rho_{12} = .5 |\psi\rangle \langle \psi| + .5 |1\rangle \langle 1| \otimes 2^{-n} I$. $\rho_{23} = .5 * 2^{-n} I \otimes |1\rangle \langle 1| + .5 |\psi\rangle \langle \psi|$. $\rho_2 = 2^{-n} I$. $\mathbf{Hg}(\rho_{12}) =^+ -\log \text{Tr} \mu^{2n} \rho_{12} <^+ -\log \text{Tr} \mu^{2n} |\psi\rangle \langle \psi| <^+ -\log \mathbf{m}(|\psi\rangle |2^{2n}) | \langle \psi | \psi \rangle |^2 <^+ 0$. Similarly, $\mathbf{Hg}(\rho_{23}) =^+ 0$. $\mathbf{Hg}(\rho_2) =^+ n$. So $\mathbf{Hg}(\rho_{123}) + \mathbf{Hg}(\rho_2) >^+ n$ and $\mathbf{Hg}(\rho_{12}) + \mathbf{Hg}(\rho_{23}) =^+ 0$, proving that \mathbf{Hg} is not strongly subadditive. \square

Theorem 15 \mathbf{Hg} is not strongly superadditive.

Proof. We fix the number of qubits n , and for $i \in [1..2^n]$, $|i\rangle$ is the i th basis state of the n qubit space. Let $|\phi\rangle = \sum_{i=1}^{2^n} 2^{-n/2} |i\rangle |i\rangle |i\rangle$, with $\mathbf{K}(|\phi\rangle |2^{3n}) = 0$. Let $\sigma_{123} = |\phi\rangle \langle \phi|$. $\sigma_{12} = \sigma_{23} = \sum_{i=1}^{2^n} 2^{-n} |i\rangle \langle i| \otimes |i\rangle \langle i|$. $\mathbf{Hg}(\sigma_{123}) =^+ -\log \text{Tr} \sigma_{123} \mu^{3n} <^+ -\log \text{Tr} \mathbf{m}(|\phi\rangle |2^{3n}) | \langle \phi | \phi \rangle |^2 <^+ 0$. Let D be a unitary transform where $D |i\rangle |i\rangle = |i\rangle |1\rangle$ and $\mathbf{K}(D |2^{2n}) =^+ 0$. So $\mathbf{Hg}(\sigma_{12}) =^+ \mathbf{Hg}(D \sigma_{12} D^*) =^+ \mathbf{Hg}(2^{-n} I \otimes |1\rangle \langle 1|) =^+ n - \log \text{Tr}(I \otimes |1\rangle \langle 1|) \mu^{2n}$. By Theorem 8 and properties of partial trace, $\mathbf{Hg}(2^{-n} I \otimes |1\rangle \langle 1|) =^+ n - \log \text{Tr} |1\rangle \langle 1| \mu^n =^+ n$. So $\mathbf{Hg}(\sigma_{12}) = \mathbf{Hg}(\sigma_{23}) =^+ n$. So $\mathbf{Hg}(\sigma_{123}) + \mathbf{Hg}(\sigma_2) <^+ n$, and $\mathbf{Hg}(\sigma_{12}) + \mathbf{Hg}(\sigma_{23}) >^+ 2n$, proving that \mathbf{Hg} is not strongly superadditive. \square

4.6 Relation Between Complexities

4.6.1 Vitányi Complexity and Gács Complexity

By definition $\mathbf{Hg}(|\psi\rangle) <^+ \mathbf{Hv}(|\psi\rangle)$. In fact, as shown in the following theorem, Vitányi complexity is bounded with respect to Gács complexity.

Theorem 16 ([GÓ1]) $\mathbf{Hg}(|\psi\rangle) <^+ \mathbf{Hv}(|\psi\rangle) <^{\log} 4\mathbf{Hg}(|\psi\rangle)$.

Proof. For semi-density matrix A with eigenvectors $\{|a_i\rangle\}$ and decreasing eigenvectors $\{a_i\}$ with $\langle \psi | A | \psi \rangle \geq 2^{-k}$ and $|\psi\rangle = \sum c_i |a_i\rangle$, let A_m be a projector onto the m largest eigenvectors. Let m be the first i where $a_i \leq 2^{-k-1}$. Since $\sum a_i \leq 1$, we have $m \leq 2^{k+1}$. Since

$$\sum_{i \geq m} a_i |c_i|^2 < 2^{-k-1} \sum_i |c_i|^2 = 2^{-k-1},$$

we have

$$\langle \psi | A_m | \psi \rangle \geq \sum_{i < m} |c_i|^2 \geq \sum_{i < m} a_i |c_i|^2 \geq 2^{-k} - \sum_{i \geq m} a_i |c_i|^2 > 2^{-k-1}.$$

Thus there is some $i \leq m$ such that $|\langle \psi | a_i \rangle|^2 \geq 2^{-2k-2}$. Let $\nu = \text{Tr} \mu$ and $\nu_k \in \mathbb{Q}$ be a rational created from the first k digits of ν . Let $\hat{\mu}$ be a lower approximation of μ , with trace greater than ν_k . So $\mathbf{K}(\hat{\mu}) <^{\log} k$. Thus if $\langle \psi | \mu | \psi \rangle \geq 2^{-k}$, then $\langle \psi | \hat{\mu} | \psi \rangle \geq 2^{-k-1}$. Thus there is an eigenvector $|u\rangle$ of $\hat{\mu}$ of complexity $\mathbf{K}(|u\rangle | N) <^{\log} 2k$ and $|\langle \psi | u \rangle|^2 \stackrel{*}{>} 2^{-2k}$, so

$$\mathbf{Hv}(|\psi\rangle) \leq \mathbf{K}(|u\rangle | N) - \log |\langle \psi | u \rangle|^2 <^{\log} 4k <^{\log} 4\mathbf{Hg}(|\psi\rangle).$$

\square

We now describe an infinite encoding scheme for an arbitrary (not necessarily elementary) quantum pure state $|\psi\rangle$. This scheme is defined as an injection between the set of pure states and $\{0, 1\}^\infty$. We define $\langle\langle |\psi\rangle \rangle\rangle$ to be an ordered list of the encoded tuples $\langle\langle |\theta\rangle \rangle, q, [|\langle \psi | \theta \rangle|^2 \geq q]]$, over all elementary states $|\theta\rangle$ and rational distances $q \in \mathbb{Q}_{>0}$. The following theorem states that only exotic pure states will have a Vitányi complexity much greater than Gács complexity. States are exotic if they have high mutual information, \mathbf{I} , with the halting sequence $\mathcal{H} \in \{0, 1\}^\infty$.

Lemma 7 For pure quantum state $|\psi\rangle$,
 $\min_{|\phi\rangle} \mathbf{K}(|\phi\rangle) - \log |\langle \psi | \phi \rangle|^2 <^{\log} -\log \sum_{|\phi\rangle} \mathbf{m}(|\phi\rangle) |\langle \psi | \phi \rangle|^2 + \mathbf{I}(\langle\langle |\psi\rangle \rangle\rangle : \mathcal{H})$.

Proof. Let \mathcal{D} be a finite set of elementary pure states, computable from $\langle|\psi\rangle\rangle$ and the value $g = \lceil -\log \sum_{|\phi\rangle} \mathbf{m}(|\phi\rangle) |\langle\psi|\phi\rangle|^2 \rceil$ such that $-\log \sum_{|\theta\rangle \in \mathcal{D}} \mathbf{m}(|\theta\rangle) |\langle\psi|\theta\rangle|^2 \leq g + 1$. It is computable because there exists an algorithm that can find \mathcal{D} by the following method. The algorithm enumerates all elementary states $|\theta\rangle$. This algorithm approximates the algorithmic probabilities $\mathbf{m}(|\theta\rangle)$ (from below) with $\widehat{\mathbf{m}}(|\theta\rangle)$. This algorithm uses $\langle|\psi\rangle\rangle$ to approximate $|\langle\theta|\psi\rangle|^2$ from below with $|\widehat{\langle\theta|\psi\rangle}|^2$. This algorithm stops when it finds a finite set \mathcal{D} such that $-\log \sum_{|\theta\rangle \in \mathcal{D}} \widehat{\mathbf{m}}(|\theta\rangle) |\widehat{\langle\theta|\psi\rangle}|^2 \leq g + 1$. Thus we have that $\mathbf{K}(\mathcal{D}|g, \langle|\psi\rangle\rangle) = O(1)$. Let $f: \mathcal{D} \rightarrow \mathbb{W}$ be a elementary function such that $|\log |\langle\psi|\theta\rangle|^2 - f(|\theta\rangle)| \leq 1$. One such f is computable relative to $\langle|\psi\rangle\rangle$, and g . Firstly this is because D is computable from $\langle|\psi\rangle\rangle$ and g . The individual values of f are computable from $\langle|\psi\rangle\rangle$, since $|\langle\psi|\theta\rangle|^2$ can be computed to any degree of accuracy. So $\mathbf{K}(f|g, \langle|\psi\rangle\rangle) = O(1)$ and $-\log \sum_{|\theta\rangle \in \mathcal{D}} \mathbf{m}(|\theta\rangle) 2^{-f(|\theta\rangle)} \leq g + 2$. One then has that

$$\begin{aligned} \min_{|\phi\rangle} \mathbf{K}(|\phi\rangle) - \log |\langle\psi|\phi\rangle|^2 &<^+ \min_{\theta \in \mathcal{D}} \mathbf{K}(|\theta\rangle) + f(|\theta\rangle) \\ &<^{\log} -\log \sum_{|\theta\rangle \in \mathcal{D}} \mathbf{m}(|\theta\rangle) 2^{-f(|\theta\rangle)} + \mathbf{I}(\langle f \rangle; \mathcal{H}). \end{aligned} \quad (4.2)$$

$$<^{\log} g + \mathbf{I}(\langle f \rangle; \mathcal{H}) \quad (4.3)$$

$$<^{\log} g + \mathbf{I}(\langle|\psi\rangle\rangle; \mathcal{H}) + \mathbf{K}(\langle f \rangle | \langle|\psi\rangle\rangle) \quad (4.4)$$

$$<^{\log} g + \mathbf{I}(\langle|\psi\rangle\rangle; \mathcal{H}) + \mathbf{K}(g)$$

$$<^{\log} -\log \sum_{|\phi\rangle} \mathbf{m}(|\phi\rangle) |\langle\psi|\phi\rangle|^2 + \mathbf{I}(\langle|\psi\rangle\rangle; \mathcal{H}).$$

Inequality 4.2 is due to Theorem 81. Inequality 4.3 is due to the definition of f and \mathcal{D} . Inequality 4.4 is due to the definition of \mathbf{I} , where $\mathbf{I}(x; \mathcal{H}) <^+ \mathbf{I}(\alpha; \mathcal{H}) + \mathbf{K}(x|\alpha)$.

Theorem 17 $\mathbf{Hg}(|\psi\rangle) <^+ \mathbf{Hv}(|\psi\rangle) <^{\log} \mathbf{Hg}(|\psi\rangle) + \mathbf{I}(|\psi\rangle; \mathcal{H}|n)$.

Proof. This follows directly from Lemma 7, relativized to n , and the fact that $\mathbf{Hg}(|\psi\rangle) =^+ -\log \text{Tr} \mu |\psi\rangle \langle\psi| =^+ -\log \sum_{|\phi\rangle} \mathbf{m}(|\phi\rangle |n) |\langle\phi|\psi\rangle|^2$.

4.6.2 BVL Complexity and Gács Complexity

Theorem 18 ([Eps20]) For pure state $|\psi\rangle \in \mathcal{Q}_n$, $\mathbf{Hg}(|\psi\rangle) <^+ \mathbf{Hbvl}[\epsilon](|\psi\rangle) + \mathbf{K}(\mathbf{Hbvl}^\epsilon(|\psi\rangle) - \log \epsilon)$.

Proof. For each k and t in \mathbb{N} , let $\mathcal{H}_{k,t}$ be the smallest linear subspace spanning elementary k -qubit inputs to the universal quantum Turing machine M of size k that halt in t steps, outputting a n qubit mixed state. As shown in [Mul08], if $t \neq t'$, then $\mathcal{H}_{k,t}$ is perpendicular to $\mathcal{H}_{k,t'}$. Let $P_{k,t}$ be the projection onto $\mathcal{H}_{k,t}$. For each k and t , the universal quantum Turing machine defines a completely positive map $\Psi_{k,t}$ over $\mathcal{H}_{k,t}$, where $\Psi_{k,t}(\nu) = \rho$ implies that the quantum Turing machine, with semi-density matrix ν of length k on the input tape will output the n qubit mixed state ρ and halt in time t . Let ρ be a k qubit mixed state that minimizes $k = \mathbf{Hbvl}[\epsilon](|\psi\rangle)$ in time

t .

$$\begin{aligned}
\rho &\leq P_{k,t} \\
2^{-k}\rho &\leq 2^{-k}P_{k,t} \\
\Psi_{k,t}2^{-k}\rho &\leq \Psi_{k,t}2^{-k}P_{k,t} \\
\Psi_{k,t}2^{-k}\rho &\leq \sum_t \Psi_{k,t}2^{-k}P_{k,t}
\end{aligned}$$

The semi density matrix $\sum_t \Psi_{k,t}2^{-k}P_{k,t}$ is lower computable relative to k , so

$$\begin{aligned}
\mathbf{m}(k|N)2^{-k}\Psi_{k,t}\rho &\leq \mathbf{m}(k|N)\sum_t \Psi_{k,t}2^{-k}P_{k,t} <^* \mu \\
\mathbf{m}(k|N)2^{-k}\langle\psi|\Psi_{k,t}(\rho)|\psi\rangle &<^* \langle\psi|\mu|\psi\rangle \\
k + \mathbf{K}(k|N) - \log \epsilon &>^+ \mathbf{Hg}(|\psi\rangle).
\end{aligned}$$

□

Theorem 19 ([Eps20]) $\mathbf{Hbvl}[2^{-\mathbf{Hg}(|\psi\rangle)-O(\log \mathbf{Hg}(|\psi\rangle))}] (|\psi\rangle) <^{\log} \mathbf{Hg}(|\psi\rangle)$.

Proof. We use reasoning from Theorem 7 in [G01]. From Theorem 7 in [Eps20] there exists a ρ such that $\mathbf{K}(\rho|N) - \log \langle\psi|\rho|\psi\rangle <^{\log} \mathbf{Hg}(|\psi\rangle)$. Let $\lceil -\log \langle\psi|\rho|\psi\rangle \rceil = m$. Let $|u_1\rangle, |u_2\rangle, |u_3\rangle, \dots$ be the eigenvectors of ρ with eigenvalues $u_1 \geq u_2 \geq u_3 \dots$. For $y \in \mathbb{N}$, let $\rho_y = \sum_{i=1}^y u_i |u_i\rangle \langle u_i|$. We expand $|\psi\rangle$ in the basis of $\{|u_i\rangle\}$ with $|\psi\rangle = \sum_i c_i |u_i\rangle$. So we have that $\sum_i u_i |c_i|^2 \geq 2^{-m}$. Let $s \in \mathbb{N}$ be the first index i with $u_i < 2^{-m-1}$. Since $\sum_i u_i \leq 1$, it must be that $s \leq 2^{m+2}$. So

$$\begin{aligned}
\sum_{i \geq s} u_i |c_i|^2 &< 2^{-m-1} \sum_i |c_i|^2 \leq 2^{-m-1}, \\
\langle\psi|\rho_{2^{m+2}}|\psi\rangle &\geq \text{Tr} \langle\psi|\rho_s|\psi\rangle > \sum_{i < s} u_i |c_i|^2 \geq 2^{-m} - \sum_{i \geq s} u_i |c_i|^2 > 2^{-m-1}.
\end{aligned}$$

We now describe a program to the universal quantum Turing machine that will construct $\rho_{2^{m+2}}$. The input is an ensemble $\{u_i\}_{i=1}^{2^{m+2}}$ of vectors $\{|cB(i)\rangle\}$, where $B(i)$ is the binary encoding of index $i \in \mathbb{N}$ which is of length $m+2$. Helper code c of size $=^+ \mathbf{K}(p|N)$ transforms each $|cB(i)\rangle$ into $|u_i\rangle$. Thus the size of the input is $<^+ \mathbf{K}(p|N) + m <^{\log} \mathbf{Hg}(|\psi\rangle)$. The fidelity of the approximation is $\langle\psi|\rho_{2^{m+2}}|\psi\rangle > 2^{-m-1} \geq 2^{-\mathbf{Hg}(|\psi\rangle)-O(\log \mathbf{Hg}(|\psi\rangle))}$. □

4.7 Quantum EL Theorem

In this paper we prove a Quantum EL Theorem. In AIT, the EL Theorem [Lev16, Eps19d] states that sets of strings that contain no simple member will have high mutual information with the halting sequence.

Theorem 20 ([Lev16, Eps19c])

For finite set $D \subset \{0, 1\}^*$, $\min_{x \in D} \mathbf{K}(x) <^{\log} -\log \sum_{x \in D} \mathbf{m}(x) + \mathbf{I}(D; \mathcal{H})$.

It has many applications, including that all sampling methods produce outliers [Eps21b]. The Quantum EL Theorem states that non exotic projections of large rank must have simple quantum

pure states in their images. By non exotic, we mean the coding of the projection has low information with the halting sequence. The Quantum EL Theorem has the following consequence.

Claim. *As the von Neumann entropy associated with the quantum source increases, the lossless quantum coding projectors have larger rank and thus must have simpler (in the algorithmic quantum complexity sense) pure states in their images.*

Theorem 21 (Quantum EL Theorem [Eps23b]) *Fix an n qubit Hilbert space. Let P be a elementary projection of rank $> 2^m$. Then, relativized to (n, m) , $\min_{|\phi\rangle \in \text{Image}(P)} \mathbf{Hv}(|\phi\rangle) <^{\log} 3(n - m) + \mathbf{I}(\langle P \rangle; \mathcal{H})$.*

Proof. We assume P has rank 2^m . Let Q be the elementary probability measure that realized the stochasticity, $\mathbf{Ks}(P)$, of an encoding of P . We can assume that every string in the support of Q encodes a projection of rank 2^m . We sample N independent pure states according to the uniform distribution Λ on the n qubit space. For each pure state $|\psi_i\rangle$ and projection R in the support of Q , the expected value of $\langle \psi_i | R | \psi_i \rangle$ is

$$\int \langle \psi_i | R | \psi_i \rangle d\Lambda = \text{Tr} R \int |\psi_i\rangle \langle \psi_i| d\Lambda = 2^{-n} \text{Tr} R I = 2^{m-n}.$$

Let random variable $X_R = \frac{1}{N} \sum_{i=1}^N \langle \psi_i | R | \psi_i \rangle$ be the average projection size of the random pure states onto the projection R . Since $\langle \psi_i | R | \psi_i \rangle \in [0, 1]$ with expectation 2^{m-n} , by Hoeffding's inequality,

$$\Pr(X_R \leq 2^{m-n-1}) < \exp \left[-N 2^{-2(m-n)-1} \right]$$

Let $d = \mathbf{d}(P|Q)$. Thus if we set $N = d 2^{2(m-n)+1}$, we can find N elementary n qubit states such that $Q(\{R : X_R \leq 2^{m-n-1}\}) \leq \exp(-d)$, where X_R is now a fixed value and not a random variable. Thus $X_P > 2^{m-n-1}$ otherwise one can create a Q -expectation test, t , such that $t(R) = \exp d$. This is a contradiction because

$$1.44d <^+ \log(P) <^+ \mathbf{d}(P|Q) <^+ d,$$

for large enough d which we can assume without loss of generality. Thus there exists i such that $\langle \psi_i | P | \psi_i \rangle \geq 2^{m-n-1}$. Thus $|\phi\rangle = P|\psi_i\rangle / \sqrt{\langle \psi_i | P | \psi_i \rangle}$ is in the image of P and $|\langle \psi_i | \phi \rangle|^2 = \langle \psi_i | P | \psi_i \rangle \geq 2^{m-n-1}$. The elementary state $|\psi_i\rangle$ has classical Kolmogorov complexity $\mathbf{K}(|\psi_i\rangle) <^{\log} \log N + \mathbf{K}(Q, d) <^{\log} 2(m - n) + \mathbf{Ks}(P)$. Thus by Lemma 6,

$$\begin{aligned} & \min\{\mathbf{Hv}(|\psi\rangle) : |\psi\rangle \in \text{Image}(P)\} \\ & \leq \mathbf{Hv}(|\phi\rangle) \\ & <^{\log} \mathbf{K}(|\psi_i\rangle) + |\langle \psi_i | \phi \rangle|^2 \\ & <^{\log} 3(n - m) + \mathbf{Ks}(P) \\ & <^{\log} 3(n - m) + \mathbf{I}(P; \mathcal{H}). \end{aligned}$$

□

4.7.1 Computable Projections

Theorem ?? is in terms of elementary described projections and can be generalized to arbitrarily computable projections. For a matrix M , let $\|M\| = \max_{i,j} |M_{i,j}|$ be the max norm. A program

$p \in \{0,1\}^*$ computes a projection P of rank ℓ if it outputs a series of rank ℓ projections $\{P_i\}_{i=1}^\infty$ such that $\|P - P_i\| \leq 2^{-i}$. For computable projection operator P , $\mathbf{I}(P; \mathcal{H}) = \min\{\mathbf{K}(p) - \mathbf{K}(p|\mathcal{H}) : p \text{ is a program that computes } P\}$.

Corollary 1 ([Eps23b]) *Fix an n qubit Hilbert space. Let P be a computable projection of rank $> 2^m$. Then, relativized to (n, m) , $\min_{|\phi\rangle \in \text{Image}(P)} \mathbf{Hv}(|\phi\rangle) <^{\log} 3(n - m) + \mathbf{I}(P; \mathcal{H})$.*

Proof. Let p be a program that computes P . There is a simply defined algorithm A , that when given p , outputs P_n such that $\min_{|\psi\rangle \in \text{Image}(P)} \mathbf{Hv}(|\psi\rangle) =^+ \min_{|\psi\rangle \in \text{Image}(P_n)} \mathbf{Hv}(|\psi\rangle)$. Thus by Lemma 4, one gets that $\mathbf{I}(P_n; \mathcal{H}) <^+ \mathbf{I}(P; \mathcal{H})$. The corollary follows from Theorem 21. \square

4.7.2 Quantum Data Compression

The Quantum EL Theorem can be used to address open issues in Quantum Information Theory. In [G01] the following remark was made.

Remark 1 ([G01]) *Maybe the study of the problem for quantum description complexity helps with the understanding of the problem for von Neumann entropy, and its relation to coding tasks of quantum information theory.*

A quantum source consists of a set of pure quantum states $\{|\psi_i\rangle\}$ and their corresponding probabilities $\{p_i\}$, where $\sum_i p_i = 1$. The pure states are not necessarily orthogonal. The sender, Alice wants to send the pure states to the receiver, Bob. Let $\rho = \sum_i p_i |\psi_i\rangle \langle \psi_i|$ be the density matrix associated with the quantum source. Let $S(\rho)$ be the von Neumann entropy of ρ . By Schumacher compression, [Sch95], in the limit of $n \rightarrow \infty$, Alice can compress n qubits into $S(\rho)n$ qubits and send these qubits to Bob with fidelity approaching 1. For example, if the message consists of n photon polarization states, we can compress the initial qubits to $nS(\rho)$ photons. Alice cannot compress the initial qubits to $n(S(\rho) - \delta)$ qubits, as the fidelity will approach 0. The qubits are compressed by projecting the message onto a typical subspace of rank $nS(\rho)$ using a projector P . The projection occurs by using a quantum measurement consisting of P and a second projector $(I - P)$, which projects onto a garbage state.

The results of this paper says that as $S(\rho)$ increases, there must be simple states in the range of P . There is no way to communicate a quantum source with large enough $S(\rho)$ without using simple quantum states.

Chapter 5

Quantum Typicality

5.1 Definition of Quantum Randomness Deficiency

In [G01], the quantum notion of randomness deficiency was introduced. This quantum randomness deficiency measures the algorithmic atypicality of a pure or mixed quantum state ρ with respect to a second quantum mixed state σ . Mixed states σ are used to model random mixtures $\{p_i\}$ of pure states $\{|\psi_i\rangle\}$, so quantum randomness deficiency is a score of how atypical a quantum state is with respect to a mixture. We first describe typicality with respect to computable σ , and then generalize to uncomputable σ .

Given a density matrix σ , a σ -test is a lower computable matrix T such that $\text{Tr} T \sigma = 1$. Let \mathcal{T}_σ be the set of all σ -tests. If σ is computable, there exists a universal σ test \mathbf{t}_σ , that is lower computable relative to the number of qubits n , $\text{Tr} \sigma \mathbf{t}_\sigma \leq 1$, and for every lower computable σ test T , $O(1) \mathbf{t}_\sigma > \mathbf{m}(T|\sigma)T$.

This universal test can be computed the following manner, analogously to the classical case (see [G21]). A program enumerates all strings p and lower computes $\mathbf{m}(p|\sigma)$. The program then runs p and continues with the outputs as long as p outputs a series of positive semi-definite matrices T_i such that $\text{Tr} T_i \sigma \leq 1$ and $T_i \leq T_{i+1}$. If p outputs something other than this sequence or does not halt, the sequence is frozen. $\mathbf{t}_\sigma = \sum_p \mathbf{m}(p|\sigma) \lim_i T_i$ is the weighted sum of all such outputs of programs p .

Definition 7 (Quantum Randomness Deficiency) For mixed states σ and ρ , computable σ , $\mathbf{d}(\rho|\sigma) = \log \text{Tr} \mathbf{t}_\sigma \rho$.

The quantum randomness deficiency, among other interpretations, is score of how typical a pure state is with respect to an algorithmically generated quantum source. Indeed, suppose there is a computable probability P over encodings of elementary orthogonal pure states $\{|\psi_i\rangle\}$ of orthogonal pure states $\{|\psi_i\rangle\}$, with corresponding density matrix $\sigma = \sum_i P(\langle|\psi_i\rangle) |\psi_i\rangle \langle\psi_i|$. Then there is a lower-computable σ -test $T = \sum_i 2^{\mathbf{d}(\langle|\psi_i\rangle)P} |\psi_i\rangle \langle\psi_i|$ with $O(1) \mathbf{t}_\sigma > T$. Thus $\mathbf{d}(|\psi_i\rangle|\sigma) >^+ \mathbf{d}(\langle|\psi_i\rangle)P$, giving high scores to pure states $|\psi_i\rangle$ which are atypical of the source. In general the $\mathbf{d}(|\phi\rangle|\sigma)$ score for arbitrary $|\phi\rangle$ will be greater than a combination of $\mathbf{d}(\cdot|P)$ scores, with $\mathbf{d}(|\phi\rangle|\sigma) >^+ \log \sum 2^{\mathbf{d}(\langle|\psi_i\rangle)P} |\langle\phi|\psi_i\rangle|^2$. In fact \mathbf{d} is equivalent to the classical definition of randomness deficiency when σ is purely classical, i.e. only diagonal.

Theorem 22 For diagonal $\sigma = \sum_i p(i) |i\rangle \langle i|$, $\mathbf{d}(|i\rangle|\sigma) =^+ \mathbf{d}(i|p)$.

Proof. The positive semi-definite matrix $T = \sum_i 2^{\mathbf{d}(i|p)} |i\rangle \langle i|$ is a σ -test, so $T \leq^* \mathbf{t}_\sigma$ and thus $\mathbf{d}(|i\rangle|\sigma) \geq^+ \log \langle i|T|i\rangle =^+ \mathbf{d}(i|p)$. The function $t(i) = \langle i|\mathbf{t}_\sigma|i\rangle$ is a lower computable p -test, so $\mathbf{d}(i|P) \geq^+ \mathbf{d}(|i\rangle|\sigma)$. \square

The following theorem shows that randomness deficiency $\mathbf{d}(\rho|\sigma)$ parallels the classical definition of randomness deficiency, $\mathbf{d}(x|P) = \log \mathbf{m}(x)/P(x)$.

Theorem 23 ([G01]) *Relativized to elementary σ , $\log \mathbf{d}(\rho|\sigma) =^+ \log \text{Tr} \rho \sigma^{-1/2} \boldsymbol{\mu} \sigma^{-1/2} \sigma$*

Proof. The matrix $\sigma^{1/2} \mathbf{t}_\sigma \sigma^{1/2}$ is a lower-computable semi density matrix, so $\mathbf{t}_\sigma \leq^* \sigma^{-1/2} \boldsymbol{\mu} \sigma^{-1/2}$. This implies $\text{Tr} \mathbf{t}_\sigma \rho \leq^* \text{Tr} \rho \sigma^{-1/2} \boldsymbol{\mu} \sigma^{-1/2}$. \square

5.1.1 Uncomputable Mixed States

We now extend \mathbf{d} to uncomputable σ . For uncomputable σ , \mathcal{T}_σ is not necessarily enumerable, and thus a universal lower computable randomness test does not necessarily exist, and cannot be used to define the σ deficiency of randomness. So in this case, the deficiency of randomness is instead defined using an aggregation of σ -tests, weighted by their lower algorithmic probabilities. The lower algorithmic probability of a lower computable matrix σ is $\underline{\mathbf{m}}(\sigma|x) = \sum \{\mathbf{m}(q|x) : q \text{ lower computes } \sigma\}$. Let $\mathfrak{T}_\sigma = \sum_{\nu \in \mathcal{T}_\sigma} \underline{\mathbf{m}}(\nu|n) \nu$.

Definition 8 (Quantum Randomness Deficiency (Uncomputable States)) *The randomness deficiency of ρ with respect to σ is $\mathbf{d}(\rho|\sigma) = \log \text{Tr} \mathfrak{T}_\sigma \rho$.*

If σ is computable, then Definition 8 equals Definition 7. By definition, \mathfrak{T}_σ is universal, since for every lower computable σ -test ν , $\underline{\mathbf{m}}(\nu) \nu < \mathfrak{T}_\sigma$.

Theorem 24 *For n qbit density matrices σ , ρ , ν , and ξ , $\mathbf{d}(\sigma|\rho) + \mathbf{d}(\nu|\xi) <^+ \mathbf{d}(\sigma \otimes \nu|\rho \otimes \xi)$.*

Proof.

$$\begin{aligned}
\mathbf{d}(\sigma|\rho) + \mathbf{d}(\nu|\xi) &= \log \text{Tr} \sum_{\rho' \in \mathcal{T}_\rho} \underline{\mathbf{m}}(\rho') \rho' \sigma + \log \text{Tr} \sum_{\xi' \in \mathcal{T}_\xi} \underline{\mathbf{m}}(\xi') \xi' \nu \\
&= \log \text{Tr} \left(\left(\sum_{\rho' \in \mathcal{T}_\rho} \underline{\mathbf{m}}(\rho') \rho' \right) \otimes \left(\sum_{\xi' \in \mathcal{T}_\xi} \underline{\mathbf{m}}(\xi') \xi' \right) \right) (\sigma \otimes \nu) \\
&= \log \text{Tr} \left(\sum_{\rho' \in \mathcal{T}_\rho, \xi' \in \mathcal{T}_\xi} \underline{\mathbf{m}}(\rho') \underline{\mathbf{m}}(\xi') \rho' \otimes \xi' \right) (\sigma \otimes \nu) \\
&<^+ \log \text{Tr} \left(\sum_{\rho' \in \mathcal{T}_\rho, \xi' \in \mathcal{T}_\xi} \underline{\mathbf{m}}(\rho' \otimes \xi') \rho' \otimes \xi' \right) (\sigma \otimes \nu) \\
&<^+ \log \text{Tr} \left(\sum_{\kappa \in \mathcal{T}_{\rho \otimes \xi}} \underline{\mathbf{m}}(\kappa) \kappa \right) (\sigma \otimes \nu) \\
&=^+ \mathbf{d}(\sigma \otimes \nu|\rho \otimes \xi).
\end{aligned}$$

\square

5.2 Conservation Over Quantum Operations

Proposition 1 *For semi-density matrix ν , relativized to a finite set of elementary matrices $\{M_i\}$, $\underline{\mathbf{m}}(\sum_i M_i^* \nu M_i | N) \stackrel{*}{>} \underline{\mathbf{m}}(\nu | N)$.*

Proof. For every string q that lower computes ν , there is a string q_M of the form rq , that lower computes $\sum_i M_i^* \nu M_i$. This string q_M uses the helper code r to take the intermediary outputs ξ_i of q and outputs the intermediary output $\sum_i M_i^* \xi_i M_i$. Since q_M has access to $\{M_i\}$ on the auxiliary tape, $\mathbf{m}(q_M | N) \stackrel{*}{>} \mathbf{m}(q | N)$.

$$\begin{aligned} \underline{\mathbf{m}}(\nu | N) &= \sum \{ \mathbf{m}(q | N) : q \text{ lower computes } \nu \} \\ &\stackrel{*}{<} \sum \{ \mathbf{m}(q_M | N) : q \text{ lower computes } \nu \} \\ &\stackrel{*}{<} \sum \{ \mathbf{m}(q' | N) : q' \text{ lower computes } \sum_i M_i^* \nu M_i \} \\ &\stackrel{*}{<} \underline{\mathbf{m}} \left(\sum_i M_i^* \nu M_i / n \right). \end{aligned}$$

□

The following theorem shows conservation of randomness with respect to elementary quantum operations. It generalizes Theorems 2 and 3 from [Eps19c].

Theorem 25 (Randomness Conservation) *Relativized to elementary quantum operation ε , for semi-density matrices ρ, σ , $\mathbf{d}(\varepsilon(\rho) | \varepsilon(\sigma)) \stackrel{+}{<} \mathbf{d}(\rho | \sigma)$.*

Proof. Since the universal Turing machine is relativized to ε , there is an elementary Kraus operator $\{M_i\}$ that can be computed from ε where $\varepsilon(\xi) = \sum_i M_i \xi M_i^*$. If ν is a $\sum_i M_i \rho M_i^*$ -test, with $\nu \in \mathcal{T}_{\sum_i M_i \rho M_i^*}$, then $\sum_i M_i^* \nu M_i$ is a ρ -test, with $\sum_i M_i^* \nu M_i \in \mathcal{T}_\rho$. This is because by assumption $\text{Tr} \nu \sum_i M_i \rho M_i^* \leq 1$. So by the cyclic property of trace $\text{Tr} \sum_i M_i^* \nu M_i \rho \leq 1$. Therefore since $\sum_i M_i^* \nu M_i$ is lower computable, $\sum_i M_i^* \nu M_i \in \mathcal{T}_\rho$. From Proposition 1, $\underline{\mathbf{m}}(\sum_i M_i^* \nu M_i | n) \stackrel{*}{>} \underline{\mathbf{m}}(\nu | n)$. So we have the following inequality

$$\begin{aligned} \mathbf{d} \left(\sum_i M_i \sigma M_i^* \middle| \sum_i M_i \rho M_i^* \right) &= \log \sum_{\nu \in \mathcal{T}_{\sum_i M_i \rho M_i^*}} \underline{\mathbf{m}}(\nu | N) \text{Tr} \nu \sum_i M_i \sigma M_i^* \\ &\stackrel{+}{<} \log \sum_{\nu \in \mathcal{T}_{\sum_i M_i \rho M_i^*}} \underline{\mathbf{m}} \left(\sum_i M_i^* \nu M_i | N \right) \text{Tr} \sum_i M_i^* \nu M_i \sigma \\ &\stackrel{+}{<} \mathbf{d}(\sigma | \rho). \end{aligned}$$

□

5.3 A Quantum Outlier Theorem

One recent result in the classical randomness deficiency case is that sampling methods produce outliers [Eps21b]. There are several proofs to this result, with one of them derived from the fact

that large sets of natural numbers with low randomness deficiencies are exotic, in that they have high mutual information with the halting sequence.

In this paper, we prove a quantum version of this result. Projections of large rank must contain pure quantum states in their images that are outlying states. Otherwise, the projections are exotic, in that they have high mutual information with the halting sequence. Thus quantum coding schemes that use projections, such as Schumacher compression, must communicate using outlier quantum states. The classical and quantum theorems are analogous, but their proofs are very different!

Theorem 26 ([Eps23c]) *Relativized to an n qubit mixed state σ , for elementary 2^m rank projector P , $3m - 2n <^{\log} \max_{|\phi\rangle \in \text{Image}(P)} \mathbf{d}(|\phi\rangle|\sigma) + \mathbf{I}(\langle P \rangle; \mathcal{H})$.*

Proof. We relativize the universal Turing machine to $\langle \sigma \rangle$ and $(3m - 2n)$. Thus it is effectively relativized to m , n , and σ . Let elementary probability measure Q and $d \in \mathbb{N}$ realize $\mathbf{Ks}(P)$, where $d = \max\{\mathbf{d}(P|Q), 1\}$. Without loss of generality we can assume that the support of Q is elementary projections of rank 2^m . There are $d2^{n-m+2}$ rounds. For each round we select an σ -test T , that is of dimension 1, $\text{Tr}\sigma T \leq 1$, and for a certain Q -probability of projections B , $\text{Tr}TB$ is large. We now describe the selection process.

Select a random test T to be $2^{m-2}|\psi\rangle\langle\psi|$, where $|\psi\rangle$ is an n qubit state chosen uniformly from the unit sphere, with distribution Λ .

$$\mathbf{E}[\text{Tr}T\sigma] = 2^{m-2} \int \text{Tr}\langle\psi|\sigma|\psi\rangle d\Lambda = 2^{m-2}\text{Tr}\sigma \int |\psi\rangle\langle\psi| d\Lambda = 2^{m-n-2}\text{Tr}\sigma = 2^{m-n-2}.$$

Thus the probability that T is a σ -test is $\geq 1 - 2^{m-n-2}$. Let I_m be an n -qubit identity matrix with only the first 2^m diagonal elements being non-zero. Let $K_m = I - I_m$. Let $p = 2^{m-n}$ and $\hat{T} = T/2^{m-2}$. For any projection B of rank 2^m ,

$$\begin{aligned} & \Pr(\text{Tr}B\hat{T} \leq .5p) \\ &= \Pr(\text{Tr}I_m\hat{T} \leq .5p) \\ &= \Pr(\text{Tr}K_m\hat{T} \geq 1 - .5p) \\ & \mathbf{E}[\text{Tr}K_m\hat{T}] = 1 - p \\ & \Pr(\text{Tr}K_m\hat{T} \geq 1 - .5p) \leq (1 - p)/(1 - .5p) \\ & \Pr(\text{Tr}B\hat{T} \geq .5p) = 1 - \Pr(\text{Tr}K_m\hat{T} \geq 1 - .5p) \\ & \geq 1 - (1 - p)/(1 - .5p) \\ & = .5p/(1 - .5p) \geq .5p \\ & \Pr(\text{Tr}BT \geq 2^{2m-n-3}) \geq .5p. \end{aligned}$$

Let Ω be the space of all matrices of the form $2^{m-2}|\phi\rangle\langle\phi|$. Let R be the uniform distribution over Ω . Let $[A, B]$ be 1 if $\text{Tr}AB > 2^{2m-n-3}$, and 0 otherwise. By the above equations, for all $A \in \text{Support}(Q)$, $\int_{\Omega}[A, B]dR(B) \geq .5p$. So $\sum_A \int_{\Omega}[A, B]Q(A)dR(B) \geq .5p$. For Hermitian matrix A , $\{A\}$ is 1 if $\text{Tr}A\sigma \leq 1$, and 0 otherwise. So $\int_{\Omega}\{A\}dR(A) \geq (1 - p2^{-2})$. Let $f = \max_T\{T\} \sum Q(A)[T, A]$.

So

$$\begin{aligned}
.5p &\leq \sum_A \int_{\Omega} [A, B] Q(A) dR(B) \\
&= \sum_A \int_{\Omega} \{B\} Q[A, B](A) dR(B) + \sum_A \int_{\Omega} (1 - \{B\}) [A, B] Q(A) dR(B) \\
&\leq \sum_A \int_{\Omega} \{B\} [A, B] Q(A) dR(B) + \int_{\Omega} (1 - \{B\}) dR(B) \\
&\leq \sum_A \int_{\Omega} \{B\} [A, B] Q(A) dR(B) + p2^{-2} \\
p/4 &\leq \sum_A \int_{\Omega} \{B\} [A, B] Q(A) dR(B) = \int_{\Omega} \left(\{B\} \sum_A [A, B] Q(A) \right) dR(B) \leq \int_{\Omega} f dR(B) \\
p/4 &\leq f.
\end{aligned}$$

Thus for each round i , the lower bounds on f proves there exists a one dimensional matrix $T_i = 2^{m-2} |\psi\rangle \langle \psi|$ such that $\text{Tr} T_i \sigma \leq 1$ and $\sum_R \{Q(R) : \text{Tr} T_i R \geq 2^{2m-n-3}\} \geq p/4 = 2^{m-n-2}$. Such a T_i is selected, and the the Q probability is conditioned on those projections B for which $[T_i, B] = 0$, and the next round starts. Assuming that there are $d2^{n-m+2}$ rounds, the Q measure of projections B such there does not exist a T_i with $[T_i, B] = 1$ is

$$\leq (1 - p/4)^{d2^{n-m+2}} \leq e^{-d}.$$

Thus there exists a T_i such that $[T_i, P] = 1$, otherwise one can create a Q test t that assigns e^d to all projections B where there does not exist T_i with $[T_i, B] = 1$, and 0 otherwise. Then $t(P) = e^d$ so

$$1.44d < \log t(P) <^+ \mathbf{d}(P|Q) <^+ d.$$

This is a contradiction, because without loss of generality, one can assume d is large. Let $T_i = 2^{m-2} |\psi\rangle \langle \psi|$ with $[T_i, P] = 1$. Let $|\phi\rangle = P|\psi\rangle / \sqrt{\langle \psi| P |\psi\rangle}$. So $\langle \phi| T_i |\phi\rangle \geq 2^{2m-n-3}$ and $|\phi\rangle$ is in the image of P . Thus by Lemma 6,

$$\begin{aligned}
2m - n &<^+ \log \langle \phi| T_i |\phi\rangle \\
2m - n &<^+ \log \max_{|\phi\rangle \in \text{Image}(P)} \langle \phi| T_i |\phi\rangle \\
2m - n &<^+ \max_{|\phi\rangle \in \text{Image}(P)} \mathbf{d}(P|\sigma) + \mathbf{K}(T_i) \\
2m - n &<^+ \max_{|\phi\rangle \in \text{Image}(P)} \mathbf{d}(P|\sigma) + (n - m) + \log d + \mathbf{K}(d) + \mathbf{K}(Q) \\
2m - n &<^+ \max_{|\phi\rangle \in \text{Image}(P)} \mathbf{d}(P|\sigma) + (n - m) + \mathbf{Ks}(P) \\
3m - 2n &<^{\log} \max_{|\phi\rangle \in \text{Image}(P)} \mathbf{d}(P|\sigma) + \mathbf{I}(P; \mathcal{H}).
\end{aligned}$$

Note that due to the fact that the left hand side of the equation is $(3m - 2n)$ and it has log precision, this enables one to condition the universal Turing machine to $(3m - 2n)$. \square

5.3.1 Computable Projections

Theorem ?? is in terms of elementary described projectcctions and can be generalized to arbitrarily computable projections. For a matrix M , let $\|M\| = \max_{i,j} |M_{i,j}|$ be the max norm. A program

$p \in \{0,1\}^*$ computes a projection P of rank ℓ if it outputs a series of rank ℓ projections $\{P_i\}_{i=1}^\infty$ such that $\|P - P_i\| \leq 2^{-i}$. For computable projection operator P , $\mathbf{I}(P; \mathcal{H}) = \min\{\mathbf{K}(p) - \mathbf{K}(p|\mathcal{H}) : p \text{ is a program that computes } P\}$.

Corollary 2 ([Eps23c]) *Relativized to an n qubit mixed state σ , for computable 2^m rank projector P , $3m - 2n <^{\log} \max_{|\phi\rangle \in \text{Image}(P)} \mathbf{d}(|\phi\rangle | \sigma) + \mathbf{I}(\langle P \rangle; \mathcal{H})$.*

Proof. Let p be a program that computes P . There is a simply defined algorithm A , that when given p and σ , outputs P_n such that $\max_{|\psi\rangle \in \text{Image}(P)} \mathbf{d}(|\psi\rangle | \sigma) =^+ \max_{|\psi\rangle \in \text{Image}(P_n)} \mathbf{d}(|\psi\rangle | \sigma)$. Thus by Lemma 4, one gets that $\mathbf{I}(P_n; \mathcal{H}) <^+ \mathbf{I}(P; \mathcal{H})$. The corollary follows from Theorem ?? . \square

Chapter 6

Quantum Information

6.1 Definition of Quantum Algorithmic Information

For a pair of random variables, \mathcal{X} , \mathcal{Y} , their mutual information is defined to be $\mathbf{I}(\mathcal{X} : \mathcal{Y}) = \mathcal{H}(\mathcal{X}) + \mathcal{H}(\mathcal{Y}) - \mathcal{H}(\mathcal{X}, \mathcal{Y}) = \mathcal{H}(\mathcal{X}) - \mathcal{H}(\mathcal{X}|\mathcal{Y}) = \sum_{x,y} p(x,y) \log p(x,y)/p(x)p(y)$. This represents the amount of correlation between \mathcal{X} and \mathcal{Y} . Another interpretation is that the mutual information between \mathcal{X} and \mathcal{Y} is the reduction in uncertainty of \mathcal{X} after being given access to \mathcal{Y} .

Quantum mutual information between two subsystems described by states ρ_A and ρ_B of a composite system described by a joint state ρ_{AB} is $I(A : B) = S(\rho_A) + S(\rho_B) - S(\rho_{AB})$, where S is the Von Neumann entropy. Quantum mutual information measures the correlation between two quantum states.

As stated in Chapter 3, The algorithmic information between two strings is defined to be $\mathbf{I}(x : y) = \mathbf{K}(x) + \mathbf{K}(y) - \mathbf{K}(x, y)$. By definition, it measures the amount of compression two strings achieve when grouped together.

The three definitions above are based off the difference between a joint aggregate and the separate parts. Another approach is to define information between two semi-density matrices as the deficiency of randomness over $\boldsymbol{\mu} \otimes \boldsymbol{\mu}$, with the mutual information of σ and ρ being $\mathbf{d}(\sigma \otimes \rho | \boldsymbol{\mu} \otimes \boldsymbol{\mu})$. This is a counter argument for the hypothesis that the states are independently chosen according to the universal semi-density matrix $\boldsymbol{\mu}$. This parallels the classical algorithmic case, where $\mathbf{I}(x : y) =^+ \mathbf{d}((x, y) | \mathbf{m} \otimes \mathbf{m}) =^+ \mathbf{K}(x) + \mathbf{K}(y) - \mathbf{K}(x, y)$. However to achieve the conservation inequalities, a further refinement is needed, with the restriction of the form of the $\boldsymbol{\mu} \otimes \boldsymbol{\mu}$ tests. Let $\mathcal{C}_{C \otimes D}$ be the set of all lower computable matrices $A \otimes B$, such that $\text{Tr}(A \otimes B)(C \otimes D) \leq 1$. Let $\mathfrak{C}_{C \otimes D} = \sum_{A \otimes B \in \mathcal{C}_{C \otimes D}} \underline{\mathbf{m}}(A \otimes B | N) A \otimes B$.

Definition 9 (Information) *The mutual information between two semi-density matrices σ , ρ is defined to be $\mathbf{I}(\sigma : \rho) = \log \text{Tr} \mathfrak{C}_{\boldsymbol{\mu} \otimes \boldsymbol{\mu}}(\sigma \otimes \rho)$.*

Up to an additive constant, information is symmetric.

Theorem 27 $\mathbf{I}(\sigma : \rho) =^+ \mathbf{I}(\rho : \sigma)$.

Proof. This follows from the fact that for every $A \otimes B \in \mathcal{C}_{\boldsymbol{\mu} \otimes \boldsymbol{\mu}}$, the matrix $B \otimes A \in \mathcal{C}_{\boldsymbol{\mu} \otimes \boldsymbol{\mu}}$. Furthermore, since $\underline{\mathbf{m}}(A \otimes B | N) \stackrel{*}{=} \underline{\mathbf{m}}(B \otimes A | N)$, this guarantees that $\text{Tr} \mathfrak{C}_{\boldsymbol{\mu} \otimes \boldsymbol{\mu}}(\sigma \otimes \rho) \stackrel{*}{=} \text{Tr} \mathfrak{C}_{\boldsymbol{\mu} \otimes \boldsymbol{\mu}}(\rho \otimes \sigma)$, thus proving the theorem. \square

6.2 Paucity of Self-Information

6.2.1 Pure States

For classical algorithmic information, for all $x \in \{0, 1\}^*$,

$$\mathbf{I}(x : x) =^+ \mathbf{K}(x).$$

As shown in this section, this property differs from the quantum case, where there exists quantum states with high descriptonal complexity and negligible self information. In fact this is the case for most quantum states, where for most n qubit pure states $|\psi\rangle$,

$$\mathbf{Hg}(|\psi\rangle) \approx n, \quad \mathbf{I}(|\psi\rangle : |\psi\rangle) \approx 0.$$

The following theorem states that the information between two elementary states is not more than the combined length of their descriptions.

Theorem 28 *For elementary ρ and σ , $\mathbf{I}(\rho : \sigma) <^+ \mathbf{K}(\rho|N) + \mathbf{K}(\sigma|N)$.*

Proof. Assume not. Then for any positive constant c , there exists semi-density matrices ρ and σ , such that

$$c\mathbf{m}(\rho|N)\mathbf{m}(\sigma|N)2^{\mathbf{I}(\rho:\sigma)} = c\mathbf{Trm}(\rho|N)\mathbf{m}(\sigma|N)\mathfrak{C}_{\mu\otimes\mu}(\rho\otimes\sigma) > 1.$$

By the definition of μ , $\mathbf{m}(\rho|N)\rho <^* \mu$ and $\mathbf{m}(\sigma|N)\sigma <^* \mu$. Therefore by the definition of the Kronecker product, there is some positive constant d such that for all ρ and σ , $d\mathbf{m}(\rho|N)\mathbf{m}(\sigma|N)(\rho\otimes\sigma) < (\mu\otimes\mu)$, and similarly

$$d\mathbf{Trm}(\rho|N)\mathbf{m}(\sigma|N)\mathfrak{C}_{\mu\otimes\mu}(\rho\otimes\sigma) < \mathbf{Tr}\mathfrak{C}_{\mu\otimes\mu}(\mu\otimes\mu).$$

By the definition of \mathfrak{C} , it must be that $\mathbf{Tr}\mathfrak{C}_{\mu\otimes\mu}\mu\otimes\mu \leq 1$. However for $c = d$, there exists a ρ and a σ , such that

$$\mathbf{Tr}\mathfrak{C}_{\mu\otimes\mu}\mu\otimes\mu > d\mathbf{Trm}(\rho|N)\mathbf{m}(\sigma|N)\mathfrak{C}_{\mu\otimes\mu}(\rho\otimes\sigma) > 1,$$

causing a contradiction. □

Theorem 29 ([Eps19b]) *Let Λ be the uniform distribution on the unit sphere of \mathcal{H}_N .*

1. $\mathbf{Hg}(I/N) =^+ \log N$,
2. $\mathbf{I}(I/N : I/N) <^+ 0$,
3. $\int 2^{-\mathbf{Hg}(|\psi\rangle)} d\Lambda \stackrel{*}{=} N^{-1}$,
4. $\int 2^{\mathbf{I}(|\psi\rangle : |\psi\rangle)} d\Lambda <^+ 0$.

Proof. (1) follows from $\mathbf{Hg}(I/N) = + - \log \text{Tr} \boldsymbol{\mu} I / N = + \log N - \log \text{Tr} \boldsymbol{\mu} = + \log N$. (2) is due to Theorem 28, with $\mathbf{I}(I/N : I/N) <^+ 2\mathbf{K}(I/N|N) <^+ 0$. (3) uses the fact that $\rho = \int |\psi\rangle \langle \psi| d\Lambda = I/N$, because $\text{Tr} \rho = 1$, and $\langle \psi | \rho | \psi \rangle = \langle \phi | \rho | \phi \rangle$. Thus $\int 2^{-\mathbf{Hg}(|\psi\rangle)} d\Lambda \stackrel{*}{=} \int \text{Tr} \boldsymbol{\mu} |\psi\rangle \langle \psi| d\Lambda \stackrel{*}{=} \text{Tr} \boldsymbol{\mu} \int |\psi\rangle \langle \psi| d\Lambda \stackrel{*}{=} N^{-1}$. (4) uses the proof of Theorem 11, which states $\int |\psi\rangle \langle \psi| \otimes |\psi\rangle \langle \psi| d\Lambda = \int |\psi\psi\rangle \langle \psi\psi| d\Lambda = \binom{N+1}{2}^{-1} P$, where P is the projection onto the space of pure states $|\psi\psi\rangle$. So

$$\begin{aligned}
\int 2^{\mathbf{I}(|\psi\rangle : |\psi\rangle)} d\Lambda &= \int \text{Tr} \mathfrak{C}_{\boldsymbol{\mu} \otimes \boldsymbol{\mu}} |\psi\rangle \langle \psi| \otimes |\psi\rangle \langle \psi| d\Lambda \\
&= \text{Tr} \mathfrak{C}_{\boldsymbol{\mu} \otimes \boldsymbol{\mu}} \int |\psi\rangle \langle \psi| \otimes |\psi\rangle \langle \psi| d\Lambda \\
&= \text{Tr} \mathfrak{C}_{\boldsymbol{\mu} \otimes \boldsymbol{\mu}} \binom{N+1}{2}^{-1} P \\
&\stackrel{*}{<} \text{Tr} \mathfrak{C}_{\boldsymbol{\mu} \otimes \boldsymbol{\mu}} N^{-2} I \\
&\stackrel{*}{=} 2^{\mathbf{I}(I/N : I/N)} \\
&<^+ 0.
\end{aligned}$$

□

6.2.2 Mixed States

The results of the previous section can be extended to mixed states. Given a uniform measure over mixed states, an overwhelming majority of such states contain no algorithmic self information. Let Λ be the uniform distribution of the unit sphere of \mathcal{H}_N . Fix any number $M \in \mathbb{N}$. Let the M -simplex be

$$\Delta_M = \{(p_i)_{1 \leq i \leq M} | p_i \geq 0, p_1 + \dots + p_M = 1\}.$$

Let η be any distribution over Δ_M . Let

$$\mu \left(\sum_{i=1}^M p_i |\psi_i\rangle \langle \psi_i| \right) = \eta(p_1, \dots, p_M) \prod_{i=1}^M \Lambda(|\psi_i\rangle),$$

Theorem 30 $\int 2^{\mathbf{I}(\sigma : \sigma)} d\mu(\sigma) <^+ 0$.

Proof.

$$\begin{aligned}
&\int 2^{\mathbf{I}(\sigma : \sigma)} d\mu(\sigma) \\
&= \text{Tr} \mathfrak{C}_{\boldsymbol{\mu} \otimes \boldsymbol{\mu}} \int_{\Delta_M} \int_{\Lambda_1} \dots \int_{\Lambda_M} \left(\sum_{i=1}^M p_i |\psi_i\rangle \langle \psi_i| \right) \otimes \left(\sum_{i=1}^M p_i |\psi_i\rangle \langle \psi_i| \right) d\Lambda_1 \dots d\Lambda_M d\eta(p_1, \dots, p_M) \\
&= \text{Tr} \mathfrak{C}_{\boldsymbol{\mu} \otimes \boldsymbol{\mu}} \int_{\Delta_M} \int_{\Lambda_1} \dots \int_{\Lambda_M} \left(\sum_{i,j=1}^M p_i p_j |\psi_i\rangle \langle \psi_i| \otimes |\psi_j\rangle \langle \psi_j| \right) d\Lambda_1 \dots d\Lambda_M d\eta(p_1, \dots, p_M) \\
&= \text{Tr} \mathfrak{C}_{\boldsymbol{\mu} \otimes \boldsymbol{\mu}} \int_{\Delta_M} \int_{\Lambda} \sum_{i=1}^M p_i^2 |\psi\psi\rangle \langle \psi\psi| d\Lambda d\eta(p_1, \dots, p_M) \\
&\quad + \text{Tr} \mathfrak{C}_{\boldsymbol{\mu} \otimes \boldsymbol{\mu}} \int_{\Delta_M} \int_{\Lambda_1} \int_{\Lambda_2} \sum_{i,j \in \{1, \dots, M\}, i \neq j} 2p_i p_j |\psi_1\rangle \langle \psi_1| \otimes |\psi_2\rangle \langle \psi_2| d\Lambda_1 d\Lambda_2 d\eta(p_1, \dots, p_M).
\end{aligned}$$

The first term is not greater than

$$\begin{aligned} & \text{Tr} \mathfrak{C}_{\mu \otimes \mu} \int_{\Delta_M} \int_{\Lambda} \sum_{i=1}^M |\psi \psi\rangle \langle \psi \psi| d\Lambda d\eta(p_1, \dots, p_M) \\ &= \text{Tr} \mathfrak{C}_{\mu \otimes \mu} \int_{\Lambda} \sum_{i=1}^M |\psi \psi\rangle \langle \psi \psi| d\Lambda. \end{aligned}$$

At this point, reasoning from the proof of Theorem 29 can be used to show that this term is $O(1)$. The second term is not greater than

$$\begin{aligned} & \text{Tr} \mathfrak{C}_{\mu \otimes \mu} \int_{\Delta_M} \int_{\Lambda_1} \int_{\Lambda_2} \left(\sum_i p_i \right) \left(\sum_i p_i \right) |\psi_1\rangle \langle \psi_1| \otimes |\psi_2\rangle \langle \psi_2| d\eta(p_1, \dots, p_M) \\ &= \text{Tr} \mathfrak{C}_{\mu \otimes \mu} \int_{\Delta_M} \int_{\Lambda_1} \int_{\Lambda_2} |\psi_1\rangle \langle \psi_1| \otimes |\psi_2\rangle \langle \psi_2| d\eta(p_1, \dots, p_M) \\ &= \text{Tr} \mathfrak{C}_{\mu \otimes \mu} \int_{\Lambda_1} \int_{\Lambda_2} |\psi_1\rangle \langle \psi_1| \otimes |\psi_2\rangle \langle \psi_2| \\ &= \text{Tr} \mathfrak{C}_{\mu \otimes \mu} (I/N \otimes I/N). \end{aligned}$$

Again, at this point, reasoning from the proof of Theorem 29 can be used to show that this term is $O(1)$.

6.3 Information Nongrowth

Classical algorithmic information non-growth laws asserts that the information between two strings cannot be increased by more than a constant depending on the computable transform f , with $\mathbf{I}(f(x) : y) < \mathbf{I}(x : y) + O_f(1)$ (Theorem 5). Conservation inequalities have been extended to probabilistic transforms and infinite sequences. The following theorem shows information non-growth in the quantum case; information cannot increase under quantum operations, the most general type of transformation that a mixed or pure quantum state can undergo. The following theorem shows information nongrowth with respect to elementary quantum operations. It generalizes Theorems 5 and 10 from [Eps19c].

Theorem 31 (Information Conservation) *Relativized to elementary quantum operation ε , for semi-density matrices ρ, σ , $\mathbf{I}(\varepsilon(\rho) : \sigma) <^+ \mathbf{I}(\rho : \sigma)$.*

Proof. Since the universal Turing machine is relativized to ε , there is an elementary Kraus operator $\{M_i\}$ that can be computed from ε where $\varepsilon(\xi) = \sum_i M_i \xi M_i^*$. Given density matrices A, B, C and D , we define $\mathbf{d}'(A \otimes B | C \otimes D) = \log \mathfrak{C}_{C \otimes D} A \otimes B$. Thus $\mathbf{I}(\sigma : \rho) = \mathbf{d}'(\sigma \otimes \rho | \mu \otimes \mu)$. The semi-density matrix $\sum_i M_i \mu M_i^*$ is lower semicomputable, so therefore $\sum_i M_i \mu M_i^* \stackrel{*}{<} \mu$ and also $(\sum_i M_i \mu M_i^* \otimes \mu) \stackrel{*}{<} \mu \otimes \mu$. So if $E \otimes F \in \mathcal{C}_{\mu \otimes \mu}$ then $\text{Tr}(E \otimes F)(\mu \otimes \mu) \leq 1$, implying that $\text{Tr}(E \otimes F)(\sum_i M_i \mu M_i^* \otimes \mu) < O(1)$. Thus there is a positive constant c , where $c(E \otimes F) \in$

$\mathcal{C}_{(\sum_i M_i \mu M_i^*) \otimes \mu}$. So we have

$$\begin{aligned} \mathbf{d}' \left(\sum_i M_i \sigma M_i^* \otimes \rho \middle| \mu \otimes \mu \right) &= \log \sum_{E \otimes F \in \mathcal{C}_{\mu \otimes \mu}} \underline{\mathbf{m}}(E \otimes F | N) \text{Tr}(E \otimes F) \left(\sum_i M_i \sigma M_i^* \otimes \rho \right) \\ &<^+ \log \sum_{E \otimes F \in \mathcal{C}_{\mu \otimes \mu}} \underline{\mathbf{m}}(c(E \otimes F) | N) \text{Tr}(E \otimes F) \left(\sum_i M_i \sigma M_i^* \otimes \rho \right) \\ &<^+ \mathbf{d}' \left(\sum_i M_i \sigma M_i^* \otimes \rho \middle| \sum_i M_i \mu M_i^* \otimes \mu \right). \end{aligned}$$

Using the reasoning of the proof of Theorem 25 on the elementary Kraus operator $\{M_i \otimes I\}$ and \mathbf{d}' , where \mathcal{C} replaces \mathcal{T} , we have that

$$\mathbf{d}' \left(\sum_i M_i \sigma M_i^* \otimes \rho \middle| \sum_i M_i \mu M_i^* \otimes \mu \right) <^+ \mathbf{d}'(\sigma \otimes \rho \middle| \mu \otimes \mu).$$

Therefore we have that

$$\begin{aligned} \mathbf{I} \left(\sum_i M_i \sigma M_i^* : \rho \right) &= \mathbf{d}' \left(\sum_i M_i \sigma M_i^* \otimes \rho \middle| \mu \otimes \mu \right) \\ &<^+ \mathbf{d}' \left(\sum_i M_i \sigma M_i^* \otimes \rho \middle| \sum_i M_i \mu M_i^* \otimes \mu \right) \\ &<^+ \mathbf{d}'(\sigma \otimes \rho \middle| \mu \otimes \mu) =^+ \mathbf{I}(\sigma : \rho). \end{aligned}$$

□

6.3.1 Algorithmic No-Cloning Theorem

The no-cloning theorem states that every unitary transform cannot clone an arbitrary quantum state. However some unitary transforms can clone a subset of pure quantum states. For example, given basis states $|1\rangle, |2\rangle, |3\rangle, \dots$ there is a unitary transform that transforms each $|i\rangle |0\rangle$ to $|i\rangle |i\rangle$. In addition, there exists several generalizations to the no-cloning theorem, showing that imperfect clones can be made. In [BH96], a universal cloning machine was introduced that can clone an arbitrary state with the fidelity of 5/6. Theorem 11 shows a generalization of the no-cloning theorem using Gács complexity.

Given the information function introduced in this chapter, a natural question to pose is whether a considerable portion of pure states can use a unitary transform to produce two states that share a large amount of shared information. The following theorem answers this question in the negative. It states that the amount of information created between states with a unitary transform is bounded by the self information of the original state.

Theorem 32 ([Eps19b]) *Let $C|\psi\rangle|0^n\rangle = |\phi\rangle|\varphi\rangle$, where C is an elementary unitary transform. Relativized to C , $\mathbf{I}(|\phi\rangle : |\varphi\rangle) <^+ \mathbf{I}(|\psi\rangle : |\psi\rangle)$.*

Proof. We have the inequalities

$$\mathbf{I}(|\phi\rangle : |\varphi\rangle) <^+ \mathbf{I}(|\phi\rangle |\varphi\rangle) : |\phi\rangle |\varphi\rangle <^+ \mathbf{I}(|\psi\rangle |0^n\rangle : |\psi\rangle |0^n\rangle) <^+ \mathbf{I}(|\psi\rangle : |\psi\rangle),$$

where the first inequality is derived using partial trace, the second inequality is derived using the unitary transform C , and the third inequality is derived by appending of an environment, all constituting quantum operations, whose conservation of information is proven in Theorem 31. \square

Theorem 32, combined with the paucity of self-information in pure states (Theorem 29) shows that only a very sparse set of pure states can, given any unitary transform, duplicate algorithmic information.

6.3.2 Purification

Every mixed state can be considered a reduced state of a pure state. The purification process is considered physical, so the extended Hilbert space in which the purified state resides in can be considered the existing environment. It should therefore be possible to regard our system with its mixed state as part of a larger system in a pure state. In this section we proof that the purifications of two mixed states will contain more information than the reduced states.

Purification occurs in the following manner, starting with a density matrix $\rho = \sum_{i=1}^n p_i |i\rangle \langle i|$. A copy of the space is defined with orthonormal basis $\{|i'\rangle\}$. In this instance the purification of ρ is $|\psi\rangle = \sum_{i=1}^n \sqrt{p_i} |i\rangle \otimes |i'\rangle$. For a density matrix ρ of size n , let \mathcal{P}_ρ^m be the set of purifications of ρ of dimension $m \geq 2n$.

Corollary 3 For all $|\psi_\sigma\rangle \in \mathcal{P}_\sigma^n$, $|\psi_\rho\rangle \in \mathcal{P}_\rho^n$, $\mathbf{d}(\sigma|\rho) <^+ \mathbf{d}(|\psi_\sigma\rangle : |\psi_\rho\rangle)$.

Corollary 4 For all $|\psi_\sigma\rangle \in \mathcal{P}_\sigma^n$, $|\psi_\rho\rangle \in \mathcal{P}_\rho^n$, $\mathbf{I}(\sigma : \rho) <^+ \mathbf{I}(|\psi_\sigma\rangle : |\psi_\rho\rangle)$.

This all follows from conservation of randomness (Theorem 25) and information (Theorem 31) over quantum operations, which includes the partial trace function.

6.3.3 Decoherence

In quantum decoherence, a quantum state becomes entangled with the environment, losing decoherence. The off diagonal elements of the mixed state become dampened, as the state becomes more like a classical mixture of states.

The single qubit example is as follows. The system is in state $|\psi_Q\rangle = \alpha|0\rangle + \beta|1\rangle$ and the environment is in state $|\psi_E\rangle$. The initial state is $|\psi_{QE}\rangle = |\psi_Q\rangle \otimes |\psi_E\rangle = \alpha|0, \psi_E\rangle + \beta|1, \psi_E\rangle$. The combined system undergoes a unitary evolution U , becoming entangled, with the result $U|\psi_{QE}\rangle = \alpha|0, E_1\rangle + \beta|1, E_2\rangle$. The density matrix is $\rho_{QE} = |\alpha|^2|0, E_1\rangle\langle 0, E_1| + |\beta|^2|1, E_2\rangle\langle 1, E_2| + \alpha^*\beta|1, E_2\rangle\langle 0, E_1| + \alpha\beta^*|0, E_1\rangle\langle 1, E_2|$. The partial trace over the environment yields

$$\rho_Q = |\alpha|^2|0\rangle\langle 0| \langle E_1|E_1\rangle + |\beta|^2|1\rangle\langle 1| \langle E_2|E_2\rangle + \alpha^*\beta|1\rangle\langle 0| \langle E_2|E_1\rangle + \alpha\beta^*|0\rangle\langle 1| \langle E_1|E_2\rangle.$$

We have $\langle E_1|E_1\rangle = \langle E_2|E_2\rangle = 1$. Two environment-related terms are time dependent and can be described by an exponential decay function

$$\langle E_1|E_2\rangle = e^{-\gamma(t)}.$$

The larger the decay, the more off diagonal terms are suppressed. So

$$\rho_Q \approx \begin{pmatrix} |\alpha|^2 & \alpha^* \beta e^{-\gamma(t)} \\ \alpha \beta^* e^{-\gamma(t)} & |\beta|^2 \end{pmatrix}.$$

The above example can be generalized to n qubit density matrices. Let $\text{Decohere}(\sigma, t)$ be a decoherence operation that dampens the off-diagonal elements of σ with decay t . By definition, Decohere is a quantum operation. Randomness is conserved over decoherence. Thus if two states decohere, the first state does not increase in algorithmic atypicality with respect to the second state.

Corollary 5 $\mathbf{d}(\text{Decohere}(\sigma, t) | \text{Decohere}(\rho, t)) <^+ \mathbf{d}(\sigma | \rho)$.

This is a corollary to Theorem 25. When a state loses coherence into the environment will not gain information with any other state.

Corollary 6 *For semi-density matrices σ and ρ , $\mathbf{I}(\text{Decohere}(\sigma, t) : \text{Decohere}(\rho, t)) <^+ \mathbf{I}(\sigma : \rho)$.*

Chapter 7

Quantum Measurements

In quantum mechanics, measurements are modeled by POVMs. A POVM E is a finite or infinite set of positive definite matrices $\{E_k\}$ such that $\sum_k E_k = I$. For a given semi-density matrix σ , a POVM E induces a semi measure over integers, where $E\sigma(k) = \text{Tr} E_k \sigma$. This can be seen as the probability of seeing measurement k given quantum state σ and measurement E . An elementary POVM E has a program q such that $U(q)$ outputs an enumeration of $\{E_k\}$, where each E_k is elementary. A quantum instrument with respect to POVM E , is a quantum operation Φ_E that takes a state σ to a set of outcomes and their probabilities, $\Phi_E(\sigma) = \sum_k E(\sigma(k)) |k\rangle \langle k|$.

7.1 Typicality and Measurements

Theorem 33 shows that measurements can increase only up to a constant factor, the deficiency of randomness of a quantum state with respect to another quantum state. The classical deficiency of randomness of a probability with respect to a another probability is denoted as follows.

Definition 10 (Deficiency, probabilities (Folklore)) For probabilities p and q over $\{0,1\}^\infty$, $\mathbf{d}(q|p) = \log \sum_x q(x) \mathbf{m}(x)/p(x)$.

Note that in the following theorem, $\mathbf{d}(E\sigma|E\rho)$ term represents the classical deficiency of randomness of a semimeasure $E\sigma$ with respect to a computable probability measure $E\rho$.

Theorem 33 ([Eps19b]) For density matrices σ, ρ , relativized to elementary ρ and POVM E , $\mathbf{d}(E\sigma|E\rho) <^+ \mathbf{d}(\sigma|\rho)$.

Proof. $2^{\mathbf{d}(E\sigma|E\rho)} = \sum_k (\text{Tr} E_k \sigma) \mathbf{m}(k|N) / (\text{Tr} E_k \rho) = \text{Tr} (\sum_k (\mathbf{m}(k|N) / \text{Tr} E_k \rho) E_k) \sigma = \text{Tr} \nu \sigma$, where the matrix $\nu = (\sum_k (\mathbf{m}(k|N) / \text{Tr} E_k \rho) E_k)$ has $\nu \in \mathcal{T}_\rho$, since ν is lower computable and $\text{Tr} \nu \leq 1$. So $2^{\mathbf{d}(\sigma|\rho)} \geq \underline{\mathbf{m}}(\nu|N) \text{Tr} \nu \sigma = \underline{\mathbf{m}}(\nu|N) 2^{\mathbf{d}(E\sigma|E\rho)}$. Since $\underline{\mathbf{m}}(\nu|N) >^* 1$, $\mathbf{d}(E\sigma|E\rho) <^+ \mathbf{d}(\sigma|\rho)$.

7.2 Information and Measurements

Given two mixed states σ and ρ and POVM E , the mutual information between the probabilities of $E\sigma$ and $E\rho$, from Definition 5, is $\mathbf{I}_{\text{Prob}}(E\sigma : E\rho)$. The following theorem states that given two states, the classical (algorithmic) information between the probabilities generated by two quantum measurements is less, up to a logarithmic factor, than the information of the two states. Thus \mathbf{I} represents an upper bound on the amount of classical algorithmic information that can be extracted between two states.

Theorem 34 *Relative to POVMS E and F , $\mathbf{I}_{\text{Prob}}(E\sigma : F\rho) <^{\log} \mathbf{I}(\sigma : \rho)$.*

Note that since the universal Turing machine is relativized to E and F , all \mathbf{K} and \mathbf{m} are conditioned to the number of qubits N . Quantum instruments with respect to POVMS E and F produces two mixed states $\Psi_E(\sigma) = \sum_{i=1}^m E_i(\sigma) |i\rangle \langle i|$ and $\Psi_F(\rho) = \sum_{j=1}^m F_j(\rho) |j\rangle \langle j|$, where, without loss of generality, m can be considered a power of 2. By Theorem 8, the (i, i) th entry of $\boldsymbol{\mu}$ is $\stackrel{*}{=} \mathbf{m}(i)$, so $\mathcal{T}_{ij} = 2^{\mathbf{K}(i)+\mathbf{K}(j)-O(1)} |i\rangle \langle i| |j\rangle \langle j|$ is a $\boldsymbol{\mu} \otimes \boldsymbol{\mu}$ test, with $\text{Tr} \mathcal{T}_{i,j}(\boldsymbol{\mu} \otimes \boldsymbol{\mu}) < 1$. So, using the fact that $x/\log x$ is convex,

$$\begin{aligned}
\mathbf{I}(\sigma : \rho) &>^+ \mathbf{I}(\Psi_E(\sigma) : \Psi_F(\rho)) \\
&>^+ \log \sum_{i,j} \mathbf{m}(\mathcal{T}_{i,j}) \mathcal{T}_{i,j} \Psi_E(\sigma) \otimes \Psi_F(\rho) \\
&>^+ \log \sum_{i,j} 2^{\mathbf{K}(i)+\mathbf{K}(j)} \mathbf{m}(i, j, \mathbf{K}(i) + \mathbf{K}(j)) E_i(\sigma) F_j(\rho) \\
&>^+ \log \sum_{i,j} 2^{\mathbf{I}(i:j) - \mathbf{K}(\mathbf{I}(i:j))} E_i(\sigma) F_j(\rho) \\
&>^+ \log \sum_{i,j} 2^{\mathbf{I}(i:j)} \mathbf{I}(i : j)^{-O(1)} E_i(\sigma) F_j(\rho) \\
&>^{\log} \log \sum_{i,j} 2^{\mathbf{I}(i:j)} E_i(\sigma) F_j(\rho) \\
&>^{\log} \mathbf{I}_{\text{Prob}}(E\sigma : F\rho).
\end{aligned}$$

Corollary 7 *For density matrices ρ and σ , and $i, j \in \mathbb{N}$, relativized to POVMS E and F , $\mathbf{I}(i : j) + \log E_i(\rho) F_j(\sigma) <^{\log} \mathbf{I}(\rho : \sigma)$.*

7.3 Algorithmic Contents of Measurements

This section shows the limitations of the algorithmic content of measurements of pure quantum states. Theorem 35 says that given a measurement apparatus E , the overwhelming majority of pure states, when measured, will produce classical probabilities with no self-information, i.e. random noise. Theorem 6 shows that there is no randomized way to process the probabilities to produce more self-information, i.e. process the random noise. This is independent of the number of measurement outcomes of E .

To prove this result, we need to define an upper-information term \mathcal{I} that is defined using *upper computable* tests. We say a semi-density matrix ρ is upper computable if there a program $q \in \{0, 1\}^*$ such that when given to the universal Turing machine U , outputs, with or without halting, a finite or infinite sequence of elementary matrices ρ_i such that $\rho_{i+1} \preceq \rho_i$ and $\lim_{i \rightarrow \infty} \rho_i = \rho$. If U reads $\leq \|q\|$ bits on the input tape, then we say p upper computes ρ . The upper probability of an upper computable mixed state σ is defined by $\overline{\mathbf{m}}(\sigma/x) = \sum \{\mathbf{m}(q/x) : q \text{ upper computes } \sigma\}$.

Let $\mathcal{G}_{C \otimes D}$ be the set of all upper computable matrices (tests) of the form $A \otimes B$, where $\text{Tr}(A \otimes B)(C \otimes D) \leq 1$. Let $\mathfrak{G}_{C \otimes D} = \sum_{A \otimes B \in \mathcal{G}_{C \otimes D}} \overline{\mathbf{m}}(A \otimes B/n)(A \otimes B)$ be an aggregation of upper computable $C \otimes D$ tests of the form $A \otimes B$, weighted by their upper probability.

Definition 11 *The upper information between semi-density matrices A and B is $\mathcal{I}(A : B) = \log \text{Tr} \mathfrak{G}_{\boldsymbol{\mu} \otimes \boldsymbol{\mu}}(A \otimes B)$.*

Proposition 2 $\mathcal{I}(I/N : I/N) = O(1)$.

Proof. $1 \geq \text{Tr} \mathfrak{G}_{\mu \otimes \mu}(\mu \otimes \mu) \stackrel{*}{>} \text{Tr} \mathfrak{G}_{\mu \otimes \mu}(I/N \otimes I/N) \stackrel{*}{>} 2^{\mathbf{I}(I/N:I/N)}.$ \square

Lemma 8

- Let Λ be the uniform distribution on the unit sphere of an n qubit space.
 $\int 2^{\mathcal{I}(|\psi\rangle:|\psi\rangle)} d\Lambda = O(1),$
- $\int 2^{\mathcal{I}(\sigma:\sigma)} d\mu(\sigma) = O(1).$

Proof. The proof follows identically to that of Theorems 29 and 30, with reference to Proposition 2. \square

Lemma 9 ([Eps21a]) *Relativized to POVM E , $\mathbf{I}_{\text{Prob}}(E\sigma:E\sigma) <^+ \mathcal{I}(\sigma:\sigma).$*

Proof. Note that all complexity terms are relativized to N , due to the relativization of E . Since $z(k) = \text{Tr} \mu E_k$ is lower semi-computable and $\sum_k z(k) < 1$, $\mathbf{m}(k) \stackrel{*}{>} \text{Tr} \mu E_k$, and so $1 > 2^{\mathbf{K}(k)-O(1)} \text{Tr} \mu E_k$. So $\nu_{i,j} = 2^{\mathbf{K}(i)+\mathbf{K}(j)-O(1)}(E_i \otimes E_j) \in \mathcal{G}_{\mu \otimes \mu}$, with $\bar{\mathbf{m}}(\nu_{i,j}) \stackrel{*}{>} \mathbf{m}(i, j).$

$$\begin{aligned} \mathcal{I}(\sigma:\sigma) &= \log \sum_{A \otimes B \in \mathcal{G}_{\mu \otimes \mu}} \bar{\mathbf{m}}(A \otimes B)(A \otimes B)(\sigma \otimes \sigma) \\ &>^+ \log \text{Tr} \sum_{i,j} \nu_{i,j} \bar{\mathbf{m}}(\nu_{i,j})(\sigma \otimes \sigma) \\ &>^+ \log \sum 2^{\mathbf{K}(i)+\mathbf{K}(j)} \mathbf{m}(i, j) E\sigma(i) E\sigma(j) \\ &>^+ \mathbf{I}_{\text{Prob}}(E\sigma:E\sigma). \end{aligned}$$

\square

Theorem 35 ([Eps21a]) *Let Λ be the uniform distribution on the unit sphere of an n qubit space. Relativized to POVM E , $\int 2^{\mathbf{I}_{\text{Prob}}(E|\psi\rangle:E|\psi\rangle)} d\Lambda = O(1).$*

Proof. By Lemma 9, $2^{\mathcal{I}(|\psi\rangle:|\psi\rangle)} \stackrel{*}{>} 2^{\mathbf{I}_{\text{Prob}}(E|\psi\rangle:E|\psi\rangle)}$. From Lemma 8, $\int 2^{\mathcal{I}(|\psi\rangle:|\psi\rangle)} d\Lambda = O(1)$. The integral $\int 2^{\mathbf{I}_{\text{Prob}}(E|\psi\rangle:E|\psi\rangle)} d\Lambda$ is well defined because $2^{\mathbf{I}_{\text{Prob}}(E|\psi\rangle:E|\psi\rangle)} = \text{Tr} \sum_{i,j} \mathbf{m}(i, j) \nu_{i,j} (|\psi\rangle \langle \psi| \otimes |\psi\rangle \langle \psi|)$, for some matrices $\nu_{i,j}$ which can be integrated over Λ . \square

Theorem 36 *Relativized to POVM E , $\int 2^{\mathbf{I}_{\text{Prob}}(E\sigma:E\sigma)} d\mu(\sigma) = O(1).$*

Proof. By Lemma 9, $2^{\mathcal{I}(\sigma:\sigma)} \stackrel{*}{>} 2^{\mathbf{I}_{\text{Prob}}(E\sigma:E\sigma)}$. From Lemma 8, $\int 2^{\mathcal{I}(\sigma:\sigma)} d\mu(\sigma) = O(1).$ \square

An implication of Theorems 35 and 36 is that for an overwhelming majority of quantum states, the probabilities induced by a measurement will have negligible self information.

7.3.1 Algorithmic Content of Decoherence

Decoherence was explained in Section 6.3.3. In the idealized case, decoherence transforms an arbitrary density matrix σ into a classical probability, with the off-diagonal terms turned to 0. Let p_σ be the classical probability that σ decoheres to, with $p_\sigma(i) = \sigma_{ii}$. The following corollary to Theorem 35, for an overwhelming majority of pure or mixed states σ , p_σ is noise, that is, has negligible self-information. The corollary follows from the fact that there is a POVM E , where $E_i = |i\rangle\langle i|$ with $E_i|\psi\rangle = p_{|\psi\rangle}(i)$.

Corollary 8 *Let Λ be the uniform distribution on the unit sphere of an n qubit space.*

- $\int 2^{\mathbf{I}_{\text{Prob}}(p_{|\psi\rangle} : \mathcal{P}_{|\psi\rangle})} d\Lambda = O(1),$
- $\int 2^{\mathbf{I}_{\text{Prob}}(p_\sigma : p_\sigma)} d\mu(\sigma) = O(1).$

7.4 PVMs

Quantum measurements is also of the form of PVMs, or projection value measures. A PVM $P = \{\Pi_i\}$ is a collection of projectors Π_i with $\sum_i \Pi_i = I$, and $\text{Tr} \Pi_i \Pi_j = 0$ when $i \neq j$. When a measurement occurs, with probability $\langle \psi | \Pi_i | \psi \rangle$, the value i is measured, and the state collapses to

$$|\psi'\rangle = \Pi_i |\psi\rangle / \sqrt{\langle \psi | \Pi_i | \psi \rangle}.$$

Further measurements of $|\psi'\rangle$ by P will always result in the i measurement, so $P|\psi'\rangle(i) = 1$. To look at the effects of a measurement operation on the algorithmic information theoretic properties of a state, take a PVM, $F = \{\Pi_i\}_{i=1}^{2^{n-c}}$, where n is the number of qubits of the Hilbert space. Let Λ_F be the distribution of pure states when F is measured over the uniform distribution Λ over n qubit spaces. Thus Λ_F represents the F -collapsed states from Λ .

Theorem 37 $n - 2c <^+ \log \int 2^{\mathbf{I}_{\text{Prob}}(F : |\psi\rangle : F|\psi\rangle)} d\Lambda_F.$

Proof. Note that $\int \langle \psi | \Pi_i | \psi \rangle d\Lambda = \text{Dim}(\Pi_i) 2^{-n}$. Furthermore, let $\kappa \subset \{1, \dots, 2^{n-c}\}$ be the set of numbers $a \in \kappa$ such that $\mathbf{K}(a) >^+ n - c$. So $|\kappa| >^* 2^{n-c}$. We have that if $\langle \psi | \Pi_i | \psi \rangle = 1$ then $\mathbf{I}_{\text{Prob}}(F|\psi) : F|\psi\rangle) = \mathbf{I}_{\text{Prob}}(j \mapsto [i = j] : j \mapsto [i = j]) = \mathbf{I}(i : i) =^+ \mathbf{K}(i).$

$$\begin{aligned} & \int 2^{\mathbf{I}(F : |\psi\rangle : F|\psi\rangle)} d\Lambda_F \\ &= \sum_{i=1}^{2^{n-c}} \text{Dim}(\Pi_i) 2^{-n} 2^{\mathbf{K}(i)} \\ &>^* \sum_{i \in \kappa} \text{Dim}(\Pi_i) 2^{-n} 2^{n-c} \\ &>^* |\kappa| 2^{-n} 2^{n-c} \\ &>^* 2^{n-2c}. \end{aligned}$$

Chapter 8

Infinite Quantum Spin Chains

A qubit abstracts the properties of a single spin $1/2$ particle. A complex system can be described by the collection of qubits, which model properties of superposition and entanglement. It can be convenient to consider a system's *thermodynamic limit*, which is the limit of infinite system size. This model is an infinite quantum spin chain. In the study of infinite quantum spin chains one can make a distinction between local and global effects. In addition, one does not need to consider boundary conditions.

A Martin L f random sequence is the accepted definition in AIT for a random infinite sequence. Can one define a quantum Martin L f infinite quantum state? This chapter shows that this can be answered in the affirmative, and even landmark theorems in AIT like the Levin-Schnorr theorem can transfer over to the quantum domain.

We first review Martin L f random sequences. A Martin L f test is an effective null set of the form $\bigcap_n G_n$, where the measure of open set G_n of the Cantor space goes toward zero. An infinite sequence passes a Martin L f test if it is not contained in its null set. A Martin L f random infinite sequence passes all Martin L f tests. Let MLR be the set of Martin L f random sequences.

In [NS19], the set of random infinite quantum states was introduced, which we call a NS random state. Just like the classical setting, a NS random state passes allso-called NS tests. An NS test is a quantum analog to Martin L f tests, and it is defined by projections instead of open sets.

8.1 Infinite Quantum Bit Sequences

Before we introduce NS random sequences, we revisit the notion of C^* algebras and functional states. A C^* algebra, \mathcal{M} , is a Banach algebra and a function $*$: $\mathcal{M} \rightarrow \mathcal{M}$ such that

- For every $x \in \mathcal{M}$, $x^{**} = x$.
- For every $x, y \in \mathcal{M}$, $(x + y)^* = x^* + y^*$ and $(xy)^* = y^*x^*$.
- For every $\lambda \in \mathbb{C}$ and $x \in \mathcal{M}$, $(\lambda x)^* = \bar{\lambda}x^*$.
- For all $x \in \mathcal{M}$, $\|x^*x\| = \|x\|\|x^*\|$.

A C^* algebra \mathcal{M} is unital if it admits a multiplicative identity $\mathbf{1}$. A state over unital \mathcal{M} is a positive linear functional $Z : \mathcal{M} \rightarrow \mathbb{C}$ such that $Z(\mathbf{1}) = 1$. States are used to define NS random sequences. The set of states of \mathcal{M} is denoted by $S(\mathcal{M})$. A state is tracial if $Z(x^*x) = Z(xx^*)$, for all $x \in \mathcal{M}$.

The C^* algebra over matrices of size 2^k over \mathbb{C} is denoted by \mathcal{M}_k . Each state $\rho \in S(\mathcal{M}_k)$, can be identified with a density matrix S such that $\rho(X) = \text{Tr}SX$, for all $X \in \mathcal{M}$. States that cannot

be represented as the convex combination of other states are called pure states. Otherwise they are called mixed states. States are used interchangeably with density matrices, depending on the context. The tracial state $\tau_n \in S(\mathcal{M}_n)$ corresponds to the matrix $2^{-n}I_{2^n}$. The algebra \mathcal{M}_∞ is the direct limit of the ascending sequence of \mathcal{M}_n . A state $Z \in S(\mathcal{M}_\infty)$ over \mathcal{M}_∞ can be seen as a sequence of density matrices $\{\rho_n\}$ that are coherent under partial traces, with $\text{Tr}_{\mathcal{M}_{n+1}}\rho_{n+1} = \rho_n$. We use $Z|_n$ to denote the restriction of state Z to the algebra \mathcal{M}_n . There is a unique tracial state $\tau \in S(\mathcal{M}_\infty)$, where $\tau|_n = \tau_n$. A projection $p \in \mathcal{M}_\infty$ is a self adjoint positive element such that $p = p^2$. A special projection $p \in \mathcal{M}_n$ is a projection represented by an elementary matrix.

8.1.1 NS Randomness

An NS Σ_1^0 set is a computable sequence of special projections $\{p_i\}$ in \mathcal{M}_∞ with $p_i \leq p_{i+1}$ over all i . For state ρ and NS Σ_1^0 set G , $\rho(G) = \sup_i \rho(p_i)$.

We define NS tests. But initially, we will provide the definition for the classical Martin L f random sequence, to provide a point of reference. A classical Martin L f test, is a sequence $\{U_n\}$ of uniformly Σ_1^0 sets of infinite sequences $U_n \subseteq \{0,1\}^\infty$ such that $\mu(U_n) \leq 2^{-n}$. An infinite sequence $\alpha \in \{0,1\}^\infty$ is Martin-L f random if there is no Martin L f test $\{U_n\}$ such that $\alpha \in \bigcap_n U_n$. There is a universal Martin L f test $\{V_n\}$ such that if $\alpha \notin \bigcap_n V_n$, then α is random.

Mirroring the classical case, a NS test is an effective sequence of NS Σ_1^0 sets $\langle G^r \rangle$ such that $\tau(G^r) \leq 2^{-r}$. Unlike a classical test, which can either pass or fail a sequence, a NS test can pass a quantum state up to a particular order. For $\delta \in (0,1)$, state $Z \in S(\mathcal{M}_\infty)$ fails test $\langle G^r \rangle$ at order δ if $Z(G^r) > \delta$ for all r . Otherwise Z passes the test at order δ . We say Z passes a NS test if it passes it at all orders $\delta \in (0,1)$.

A state is NS random if it passes every NS test at every order.

Theorem 38 ([NS19]) *There exists a universal NS test $\langle R^n \rangle$, where for each NS test $\langle G^k \rangle$ and each state Z and for each n there exists a k such that $Z(R^n) \geq Z(G^k)$.*

Proof. Let $\langle G_n^k \rangle_{n=1}^\infty$ be an enumeration of NS tests, performed analogously to the classical case (see [G 1]). Furthermore let $G_m^e = \langle p_{m,r}^e \rangle_{r \in \mathbb{N}}$. For each $k, n \in \mathbb{N}$, let $q_k^n = \bigvee_{e+n+1 \leq k} p_{e+n+1,k}^e$. Thus $q_k^n \leq q_{k+1}^n$ and $\tau q_k^n \leq \sum_e \tau(p_{e+n+1,k}^e) \leq 2^{-n}$. The universal NS test is $R^n = \langle q_k^n \rangle_{k \in \mathbb{N}}$. Since $\tau(R^r) \leq 2^{-r}$, $\langle R^n \rangle$ is a NS test. For a set e ,

$$\rho(R^n) = \sup_k \rho(q_k^n) \geq \sup_k \rho(p_{n+e+1,k}^e) = \rho(G_{n+e+1}^e).$$

□

A state Z is NS random if it passes the test $\langle R^n \rangle$. More information about $\langle R^r \rangle$ can be found in [NS19].

8.2 Closure Properties

The set of NS random sequences has closure properties over (possibly noncomputable) convex combinations, as shown in the following theorem.

Theorem 39 *Every convex combination $Z = \sum_i \alpha_i Z_i$ of NS random states Z_i , with $\sum_i \alpha_i = 1$ and $\alpha_i \geq 0$, is NS random.*

Proof. Given an NS test $\langle G^r \rangle = \langle p_t^r \rangle$, there exists a NS test $\langle H^r \rangle$ such that for all states Z , $\inf_r Z(H^r) \geq \inf_r Z(G^r)$ and $H^r \supseteq H^{r+1}$. This is by setting H^r equal to $\bigvee_{i \geq r} G^i$. More formally, $\langle H^r \rangle = \langle q_t^r \rangle$, where $q_t^r = \bigvee_{i=1}^t p_t^{r+i}$. Thus there exists a universal NS test $\langle L^r \rangle$ such that $L^r \supseteq L^{r+1}$. Assume that Z is not NS random. Then

$$\begin{aligned} \lim_{r \rightarrow \infty} Z(L^r) &> 0 \\ \lim_{r \rightarrow \infty} \sum_i \alpha_i Z_i(L^r) &> 0 \\ \sum_i \alpha_i \lim_{r_i \rightarrow \infty} Z_i(L^{r_i}) &> 0. \end{aligned}$$

So there exists an i such that $\lim_{r \rightarrow \infty} Z_i(L^r) > 0$, and thus Z_i is not NS random.

8.3 Gács Complexity and NS Random Sequences

In this section, we characterize NS random states in terms of Gács complexity, \mathbf{Hg} .

Theorem 40 *Given state $Z \in \mathcal{M}_\infty$, and program p that enumerates infinite set $A \subseteq \mathbb{N}$, then $\sup_{n \in \mathbb{N}} n - \mathbf{Hg}(Z \upharpoonright n) <^+ \sup_{n \in A} n - \mathbf{Hg}(Z \upharpoonright n) + \mathbf{K}(p)$.*

Proof. There exists a program p' of size $\|p\| + O(1)$ that outputs a list $\{a_n\} \subseteq A$ such that $n < a_n$. For a given a_n , $\sigma = 2^{n-a_n} \mu_n \otimes I_{a_n-n}$ is a lower computable $2^{a_n} \times 2^{a_n}$ semi-density matrix. There is a program $q = q' \langle a_n, n \rangle$ that lower computes σ where q' is helper code that uses the encodings of a_n and n . By the universal properties of μ , we have the inequality $\mathbf{m}(q|a_n)\sigma <^* \mu_{a_n}$. So, using properties of partial trace,

$$\begin{aligned} a_n + \log \mathbf{m}(q|a_n) \text{Tr} \sigma Z \upharpoonright a_n &<^+ a_n + \log \text{Tr} \mu(Z \upharpoonright a_n) \\ a_n + \log \text{Tr} 2^{n-a_n} (\mu_n \otimes I_{a_n-n}) Z \upharpoonright a_n - \mathbf{K}(q|a_n) &<^+ a_n + \log \text{Tr} \mu(Z \upharpoonright a_n) \\ n + \log \text{Tr} (\mu_n \otimes I_{a_n-n}) Z \upharpoonright a_n - \mathbf{K}(\langle n, a_n \rangle | a_n) &<^+ a_n + \log \text{Tr} \mu(Z \upharpoonright a_n) \\ n + \log \text{Tr} (\mu_n \text{Tr}_n Z \upharpoonright a_n) - \mathbf{K}(p'|a_n) &<^+ a_n + \log \text{Tr} \mu(Z \upharpoonright a_n) \\ n - \mathbf{Hg}(Z \upharpoonright n) &<^+ a_n - \mathbf{Hg}(Z \upharpoonright a_n) + \mathbf{K}(p). \end{aligned}$$

So $\sup_{n \in \mathbb{N}} n - \mathbf{Hg}(Z \upharpoonright n) <^+ \sup_{a_n \in \{a_n\}} a_n - \mathbf{Hg}(Z \upharpoonright a_n) + \mathbf{K}(p) <^+ \sup_{n \in A} n - \mathbf{Hg}(Z \upharpoonright n) + \mathbf{K}(p)$.

Theorem 41 *Suppose for state Z , and for infinite enumerable set $A \subseteq \mathbb{N}$, $\sup_{n \in A} n - \mathbf{Hg}(Z \upharpoonright n) < \infty$. Then Z is NS random.*

Proof. Suppose Z is not NS random. Let $L^r = \langle p_t^r \rangle$ be the universal NS test. So $\text{Rank}(p_n^r) \leq 2^{n-r}$. Thus $\inf_r Z(L^r) = \delta > 0$. For each r , there exists an n such that $\text{Tr}(p_n^r z_n) \geq \delta$, where $z_n = Z \upharpoonright n$. Since $2^{r-n} p_n^r$ is a computable semi-density matrix given n and r , $\mathbf{m}(r|n) 2^{r-n} p_n^r <^* \mu$. So $\mathbf{m}(r|n) 2^{r-n} \delta <^* \text{Tr} \mu z_n$, which implies that $\mathbf{Hg}(Z \upharpoonright n) <^+ n - r + \mathbf{K}(r|n)$. Since this property holds for all $r \in \mathbb{N}$, $\sup_n n - \mathbf{Hg}(Z \upharpoonright n) = \infty$. From Theorem 40, $\sup_{n \in A} n - \mathbf{Hg}(Z \upharpoonright n) = \infty$.

8.4 Encodings of States

Let $[Z] \in \{0,1\}^\infty$ be an encoding of the state Z described as follows. For each n , let $e(n,m)$ be the m th enumeration of a pair (p,k) consisting of a special projection p of \mathcal{M}_n and a rational $0 \leq k \leq 1$. For $[Z]$, the i th bit, where $i = 2^m m$ for maximum n , corresponds to 1 if and only if $\text{Tr} p Z \upharpoonright n > k$, where (p,k) is the pair enumerated by $e(n,m)$. We say that state $Z \in \mathcal{QH}$ if and only if the halting sequence can be computed from $[Z]$.

8.5 Quantum Operation Complexity

In a canonical algorithmic information theory example, Alice wants to send a single text message x to Bob. Alice sends a program q to Bob such that $x = U(q)$, where U is a fixed universal Turing machine. The cost of the transmission is the length of q . Alice can minimize cost by sending $\mathbf{K}(x)$ bits to Bob, where \mathbf{K} is the Kolmogorov complexity function.

We now look at the quantum case. Suppose that Alice wants to send a (possibly mixed) n qubit quantum state σ to Bob, represented as an density matrix over \mathbb{C}^{2^n} , or an element of $S(\mathcal{M}_n)$. Alice has access to two channels, a quantum channel and a classical channel. Alice can choose to send $m \leq n$ qubits ρ on the quantum channel and classical bits $q \in \{0,1\}^*$ on the classical channel, describing an elementary quantum operation η , where $U(q) = [\eta]$. Bob then applies η to ρ to produce $\sigma' = \eta(\rho)$. Bob is not required to produce σ exactly. Instead the fidelity of the attempt is measured by the trace distance between σ and σ' . The trace distance D between two matrices A and B is $D(A,B) = \frac{1}{2} \|A - B\|_{\text{Tr}}$, with $\|A\|_{\text{Tr}} = \text{Tr}|A|$. We use $\mathcal{O}_{m,n}$ to denote the set of elementary quantum operations that take m qubit quantum states to $n \geq m$ qubit quantum states.

Definition 12 For n qubit density matrix σ , the quantum operation complexity at accuracy ϵ is $\mathbf{Hoc}^\epsilon(\sigma) = \min\{\mathbf{K}([\eta]) + m : \eta \in \mathcal{O}_{m,n}, \xi \in S(\mathcal{M}_m), D(\sigma, \eta(\xi)) < \epsilon\}$.

8.6 Initial Segment Incompressibility

Due to Levin and Schnorr, [Lev74, Sch71] α is random iff there is an r such that for all n , $\mathbf{K}(\alpha_{\leq n}) \geq n - r$, where $\alpha_{\leq n}$ is a prefix of α of size n , and \mathbf{K} is prefix free Kolmogorov complexity. In this section, we prove a quantum analog to this result. We show that NS states that are NS random have incompressible prefixes with respect to quantum operation complexity. Theorem 42 builds upon the proof of the Theorem 4.4 in [NS19] using quantum operation complexity \mathbf{Hoc} .

Theorem 42 Let Z be a state on \mathcal{M}_∞ .

1. Let $1 > \epsilon > 0$, and suppose Z passes each NS test at order $1 - \epsilon$. Then there is an r where for all n , $\mathbf{Hoc}^\epsilon(Z \upharpoonright_n) > n - r$.
2. Let $1 > \epsilon > 0$ be lower computable and Z fails some NS test at order $1 - \epsilon$. Then either $Z \in \mathcal{QH}$ or for all r , there is an n where $\mathbf{Hoc}^{\sqrt{\epsilon}}(Z \upharpoonright_n) < n - r$.

Proof. (1). Let $\mathbf{K}_t(x)$ be the smallest program to produce x in time t . Let $s(n,r,t)$ be the set of pure n qubit states $\rho \in S(\mathcal{M}_n)$ such that there exists a quantum operation $\eta \in \mathcal{O}_{z,n}$ and pure state $\sigma \in S(\mathcal{M}_z)$ such that $\rho = \eta(\sigma)$ and $\mathbf{K}_t([\eta]) + z \leq n - r$. Let $p(n,r,t)$ be the orthogonal projection in \mathcal{M}_n with minimum $\tau(p(n,r,t))$ such that $\rho(p(n,r,t)) = 1$ for all $\rho \in s(n,r,t)$. Let $p(r,t) = \sup_{n \leq t} p(n,r,t)$. So $p(r,t)$ is in \mathcal{M}_t and $p(r,t)$ is computable from r and t and $p(r,t) \leq p(r,t+1)$.

Let $b(y, n, z)$ be the number of programs of length y which outputs an encoding of an elementary quantum operation $\eta \in \mathcal{O}_{z,n}$. Let $b(y, n)$ be the number of programs of length y which outputs an encoding of an elementary quantum operation $\eta \in \mathcal{O}_{z,n}$, for any $z \leq n$. So

$$\begin{aligned}
\text{Range}(p(n, r, t)) &\leq \sum_{y+z \leq n-r} b(y, n, z) 2^z \\
\tau(p(n, r, t)) &\leq \sum_{y+z \leq n-r} b(y, n, z) 2^{z-n} \\
&\leq \sum_{y+z \leq n-r} b(y, n, z) 2^{-y-r} \\
&\leq \sum_{y=1}^{n-r} b(y, n) 2^{-y-r} \\
\tau(p(r, t)) &\leq \sum_{n=1}^{\infty} \tau(p(n, r, t)) \\
&\leq \sum_{n=1}^{\infty} \sum_{y=1}^{n-r} b(y, n) 2^{-y-r} \\
&= 2^{-r} \sum_{n=1}^{\infty} \sum_{y=1}^{n-r} b(y, n) 2^{-y} \leq 2^{-r}.
\end{aligned}$$

So for NS Σ_1^0 set G^r enumerated by the sequence $\{p(r, t)\}_t$, $\langle G^r \rangle$ is a NS test. For each r suppose there is an n such that $\mathbf{Hoc}^\epsilon(Z \upharpoonright n) \leq n - r$. So there is an elementary quantum operation $\eta \in \mathcal{O}_{z,n}$ and input $\rho \in S(\mathcal{M}_z)$ such that $\mathbf{K}([\eta]) + z \leq n - r$ and $D(Z \upharpoonright n, \eta(\rho)) < \epsilon$. So $\eta(\rho)$ is in the range $p(n, r, t)$ for some t and so $\text{Tr} \eta(\rho) p(n, r, t) = 1$. This implies $1 - \epsilon < Z(p(n, r, t)) \leq Z(G^r)$. Since this is for all r , Z fails the test at order $1 - \epsilon$.

(2). Let $\mathbf{bb}(n)$ be the longest running time of a halting program of length $\leq n$. Let $\langle L^r \rangle$ be the universal NS test, where each L^r is enumerated by $\{p_t^r\}$, with $p_t^r \in \mathcal{M}_{n(r,t)}$. Assume there is an infinite number of r where $\text{Tr} Z \upharpoonright n(r, \mathbf{bb}(r/2)) p_{\mathbf{bb}(r/2)}^r > 1 - \epsilon$. Fix one such r and let $n = n(r, \mathbf{bb}(r/2))$, and $p = p_{\mathbf{bb}(r/2)}^r$. Projection p has eigenvectors $\{u_i\}$ and kernel spanned by $\{v_i\}$. Thus $2^{-r} \geq \tau(p)$. Let $p' \geq p$ with $p' \in \mathcal{M}_n$ such that each u_i is in the range of p' and $\{v_i\}_{i=1}^k$ is in the range of p' such that k is minimized such that $\tau(p') = 2^{-r}$. Thus $\text{Tr} Z \upharpoonright n(p') > 1 - \epsilon$. The eigenvectors of p' are $\{w_i\}_{i=1}^{2^{n-r}}$ and its kernel is spanned by the vectors $\{y_i\}_{i=1}^{2^n - 2^{n-r}}$. Let $z' = \text{Proj}(Z \upharpoonright n; p')$ be a density matrix with eigenvalues $v_i \in \mathbb{R}$ corresponding to eigenvectors w_i . For $i \in [1, 2^n]$, let $B(i) \in \{0, 1\}^*$ be an encoding of n bits of the number i , with $B(1) = 0^{(n)}$, $B(2) = 10^{(n-1)}$, and $B(2^n) = 1^{(n)}$. Let U be a $2^n \times 2^n$ unitary matrix, of the form $U = \sum_{i=1}^{2^{n-r}} |B(i)\rangle \langle w_i| + \sum_{i=1}^{2^n - 2^{n-r}} |B(i + 2^{n-r})\rangle \langle y_i|$.

Proposition 3 ([NS19]) *Let $\text{Proj}(s; h) = \frac{1}{\text{Tr}[sh]} hsh$. Let p be a projection in M_n and σ be a density matrix in M_n . If $\alpha = \text{Tr} p\sigma$ and $\sigma' = \text{Proj}(\sigma; p)$ then $D(\sigma, \sigma') \leq \sqrt{1 - \alpha}$.*

Proof. Let $|\psi_\sigma\rangle$ be a purification of σ . Then $\alpha^{-\frac{1}{2}} p |\psi_\sigma\rangle$ is a purification of σ' . Uhlmann's theorem states $F(\sigma, \sigma') \geq \alpha^{-\frac{1}{2}} \langle \psi_\sigma | p |\psi_\sigma \rangle = \alpha^{\frac{1}{2}}$, where F is fidelity, with $F(\sigma, \sigma') = \text{Tr} \sqrt{\sqrt{\sigma'} \sigma \sqrt{\sigma'}}$. Thus the proposition follows from $D(\sigma, \sigma') \leq \sqrt{1 - F(\sigma, \sigma')}$. \square

Thus for the diagonal $2^{n-r} \times 2^{n-r}$ matrix σ with entries $\{v_i\}_{i=1}^{2^{n-r}}$, $z' = U(\sigma \otimes |0^r\rangle\langle 0^r|)U^*$. By Proposition 3, since $1 - \epsilon < \text{Tr}(p'Z \upharpoonright n)$ and $z' = \text{Proj}(z_n; p')$, it must be that $D(z', Z \upharpoonright n) < \sqrt{\epsilon}$. Thus using quantum operation $\eta = (U, |0^r\rangle\langle 0^r|, \emptyset)$ and input σ ,

$$\begin{aligned} \mathbf{Hoc}^\epsilon(Z \upharpoonright n) &\leq \text{Dim}(\sigma) + \mathbf{K}([\eta]) \\ &\leq n - r + \mathbf{K}([(U, |0^r\rangle\langle 0^r|, \emptyset)]) \\ &<^+ n - r + \mathbf{K}(n, r) \\ &<^+ n - r + \mathbf{K}(\mathbf{bb}(r/2), r) \\ &<^+ n - r + r/2 + \mathbf{K}(r) \\ &<^+ n - r/3. \end{aligned}$$

Thus for every r there exists an n where $\mathbf{Hoc}^\epsilon(Z \upharpoonright n) < n - r$. This is because the additive constant of the above equation is not dependent on r .

Otherwise there is some R where for all $r \geq R$, and $q < \mathbf{bb}(r/2)$, $\text{Tr}Z_{n(r,q)}p_{n(r,q)}^r \leq 1 - \epsilon$. Thus given R , $\langle L^r \rangle$, $[Z]$, and a lower enumeration of ϵ , one can iterate through each $r \geq R$ and return an s such that $\text{Tr}Z_{n(r,s)}p_{n(r,s)}^r > 1 - \epsilon$. This is because the set of rational numbers Q such that $q > 1 - \epsilon$ for all $q \in Q$ can be enumerated and the set $V = \{v : \text{Tr}Z_{n(r,v)}p_{n(r,v)}^r > q, q \in Q\}$ can be enumerated using the infinite encoding $[Z] \in \{0, 1\}^\infty$. The returned s is the first enumerated element of V . This number s has the property that $s \geq \mathbf{bb}(r/2)$, and can be used to compute the prefix of the halting sequence over all programs of length $\leq r/2$ as every such program that will halt will do so in less than s steps. Thus the halting sequence is computable relative to $[Z]$ and thus $Z \in \mathcal{QH}$.

Corollary 9 *Let state $Z \notin \mathcal{QH}$. Then Z is NS random iff for all $0 < \epsilon < 1$, there is an r , where for all n , $\mathbf{Hoc}^\epsilon(Z \upharpoonright n) > n - r$.*

Proof. Assume Z is NS random. Then for all $0 < \epsilon < 1$, Z passes each NS test at order $1 - \epsilon$. Then by Theorem 42 (1), for all $0 < \epsilon < 1$ there is an r where for all n , $\mathbf{Hoc}^\epsilon(Z \upharpoonright n) > n - r$. Assume Z is not NS random. Then there is some rational $0 < \delta < 1$ such that Z fails some NS test at order $1 - \delta$. Then by Theorem 42 (2), for $\epsilon = \sqrt{\delta}$, for all r , there is an n where $\mathbf{Hoc}^\epsilon(Z \upharpoonright n) < n - r$.

8.7 Quantum Ergodic Sources

In [Bru78], Brudno proved that for ergodic measures η over bi-infinite sequences, for η -almost all sequences, the rate of the Kolmogorov complexity of their finite prefixes approaches the entropy rate of η . Therefore the average compression rate of sequences produced by η is not more than its entropy rate. In [BKM⁺06], a quantum version of Brudno's theorem was introduced relating, in a similar fashion, Von Neumann entropy and BVL complexity (using the fidelity measure). The results provide two bounds with respect to two variants of **Hbvl**: approximate-scheme complexity and finite accuracy complexity.

In this subsection we provide a quantum variant of Brudno's theorem with respect to quantum communication complexity \mathbb{R}^ϵ . Differently from the **Hbvl** results, the bounds provided below are for almost all n , invariant to the accuracy term ϵ .

We define the quasilocal C^* algebra \mathcal{A}_∞ , which differs only from \mathcal{M}_∞ in that it is a doubly infinite product space over \mathbb{Z} . In particular, \mathcal{A} is the C^* algebra of qbits, i.e. 2×2 matrices acting on \mathbb{C}^2 . For finite $\Lambda \subset \mathbb{Z}$, $\mathcal{A}_\Lambda = \bigotimes_{z \in \Lambda} \mathcal{A}_z$.

The quasilocal C^* algebra \mathcal{A}_∞ is defined to be the norm closure of $\bigcup_{\Lambda \subset \mathbb{Z}} \mathcal{A}_\Lambda$. For states Ψ over \mathcal{A}_∞ , we use $\Psi|_n$ to denote Ψ restricted to the finite subalgebra $\mathcal{A}_{\{1, \dots, n\}}$ of \mathcal{A}_∞ . The right shift T is a $*$ -automorphism on \mathcal{A}_∞ uniquely defined by its actions on local observables $T : a \in \mathcal{A}_{\{m, \dots, n\}} \mapsto \mathcal{A}_{\{m+1, \dots, n+1\}}$. A quantum state Ψ is stationary if for all $a \in \mathcal{A}_\infty$, $\Psi(a) = \Psi(T(a))$. The set of shift-invariant states on \mathcal{A}_∞ is convex and compact in the weak* topology. The extremal points of this set are called ergodic states.

Lemma 10 *Let R_j be the smallest subspace spanned by pure states produced by elementary quantum operations $\eta \in \mathcal{O}_{z,n}$ with $\mathbf{K}(\eta) + z < j$. Then $\text{Dim}(R_j) < 2^j$.*

Proof. Let $b(y, z)$ be the number of programs of length y that outputs an elementary quantum operation $\eta \in \mathcal{O}_{z,z}$ over the Hilbert space \mathcal{H}_{2^n} . Let $b(y)$ be the number of programs of length y that outputs an elementary quantum operation $\mathcal{O}_{z,n}$ over the Hilbert space \mathcal{H}_{2^n} .

$$\begin{aligned} \text{Dim}(R_j) &\leq \sum_{y+z < j} b(y, z) 2^z \\ &= 2^j \sum_{y+z < j} b(y, z) 2^{z-j} \\ &< 2^j \sum_{y,z} b(y, z) 2^{-y} \\ &= 2^j \sum_y b(y) 2^{-y} \\ &\leq 2^j. \end{aligned}$$

Theorem 43 *Let Ψ be an ergodic state with mean entropy h . For all $\delta > 0$, for almost all n , there is an orthogonal projector $P_n \in \mathcal{A}_n$ such that for all $\epsilon > 0$,*

1. $\Psi|_n(P_n) > 1 - \delta$.
2. For all one dimensional projectors $p \leq P_n$, $\mathbf{Hoc}^\epsilon(p)/n \in (h - \delta, h + \delta)$.

Proof. Let $\delta' < \delta'' < \delta$. From [BDK⁺05], there is a sequence of projectors $P'_n \in \mathcal{A}_n$ where for almost all n , $\Psi|_n(P'_n) > 1 - \delta'$, for all one dimensional projectors $p' \leq P'_n$, $2^{-n(h+\delta')} < \Psi|_n(p') < 2^{-n(h-\delta')}$, and $2^{n(h-\delta'')} < \text{Tr} P'_n < 2^{n(h+\delta')}$. Let S'_n be the subspace that P'_n projects onto. Let R_n be the smallest subspace spanned by all pure states produced by an elementary quantum operation $\eta \in \mathcal{O}_{g,n}$, where $\mathbf{K}(\eta) + g < n(h - \delta'')$. Let Q_n be the projector onto R_n . By Lemma 10, $\text{Dim}(R_n) < 2^{n(h-\delta'')}$. Let S_n be the largest subspace of S'_n that is orthogonal to R_n . Let P_n be the orthogonal projector onto S_n . So for sufficiently large n , $\Psi|_n(P_n) \geq \Psi|_n(P'_n) - \text{Dim}(R_n) 2^{-n(h-\delta')} > 1 - \delta' - 2^{n(h-\delta'')} 2^{-n(h-\delta')} = 1 - \delta' - 2^{n(\delta'-\delta'')} > 1 - \delta$, for large enough n .

By definition, since P_n is orthogonal to R_n , for all ϵ , for all one dimensional projectors $p \leq P_n$, $\mathbf{Hoc}^\epsilon(p) \geq n(h - \delta'') > n(h - \delta)$. Furthermore, all such p can be produced from an elementary quantum operation η that maps $\lceil n(h + \delta') \rceil$ length pure states into S_n . Therefore for large enough n , $\mathbf{Hoc}^\epsilon(p) \leq \mathbf{K}(\eta) + \lceil n(h + \delta') \rceil <^+ \mathbf{K}(n, h) + \lceil n(h + \delta') \rceil < n(h + \delta)$.

8.8 Measurement Systems

We note that pre-measures are of the form $\gamma : \{0, 1\}^* \rightarrow \mathbb{R}_{\geq 0}$, where $\gamma(x) = \gamma(x0) + \gamma(x1)$. By the Carathéodory's Extension Theorem, each such pre-measure can be uniquely extended to a measure Γ over $\{0, 1\}^\infty$. In Chapter 7, measurements of finite collections of qubits are studied. This section deals with measurement measurement systems, which can be applied to infinite quantum states.

Definition 13 (Measurement System ([Bho21])) *An α -computable measurement system $B = \{(|b_0^n\rangle, |b_1^n\rangle)\}$ is a sequence of orthonormal bases for \mathcal{Q}_1 such that each $|b_i^n\rangle$ is elementary and the sequence $\langle |b_1^n\rangle, |b_0^n\rangle \rangle_{n=1}^\infty$ is α -computable.*

Note that the above definition can be generalized to a sequence of PVMs. We now define the application of a measurement system B to an infinite quantum state Z which produces a pre-measure p . Let ρ_n be the density matrix associated with $Z \upharpoonright n$. For the first bit, we use the standard definition of measurement, where

$$p(i) = \text{Tr} |b_i^1\rangle \langle b_i^1| \rho_1.$$

Given ρ_2 , if i is measured on the first bit, then the resulting state would be

$$\rho_2^i = \frac{(|b_i^1\rangle \langle b_i^1| \otimes I) \rho_2 (|b_i^1\rangle \langle b_i^1| \otimes I)}{\text{Tr}(|b_i^1\rangle \langle b_i^1| \otimes I) \rho_2}$$

So

$$\begin{aligned} p(ij) &= p(i)p(j|i) \\ &= (\text{Tr} |b_i^1\rangle \langle b_i^1| \rho_1) \text{Tr} (I \otimes |b_j^2\rangle \langle b_j^2|) \left(\frac{(|b_i^1\rangle \langle b_i^1| \otimes I) \rho_2 (|b_i^1\rangle \langle b_i^1| \otimes I)}{(|b_i^1\rangle \langle b_i^1| \otimes I) \rho_2} \right) \end{aligned}$$

Since $\text{Tr}_2 \rho_2 = \rho_1$, $\text{Tr} |b_i^1\rangle \langle b_i^1| \rho_1 = \text{Tr} (|b_i^1\rangle \langle b_i^1| \otimes I) \rho_2$. Therefore

$$p(ij) = \text{Tr} \rho_2 (|b_i^1 b_j^2\rangle \langle b_i^1 b_j^2|).$$

More generally for $x \in \{0, 1\}^n$, we define the pre-measure p to be

$$p(x) = \text{Tr} \rho_n |\otimes_{i=1}^n b_{x_i}^i\rangle \langle \otimes_{i=1}^n b_{x_i}^i|;$$

It is straightforward to see that p is a pre-measure, with $p(x) = p(x0) + p(x1)$. Let μ_Z^B be the measure over $\{0, 1\}^\infty$ derived from the described pre-measure, using measurement system B and state Z . We recall that MLR is the set of Martin L f random sequences.

Definition 14 (Bhojraj Random) *A state Z is Bhojraj Random if for any computable measurement system B , $\mu_Z^B(\text{MLR}) = 1$.*

Theorem 44 ([Bho21]) *All NS Random states are Bhojraj Random states,*

Proof. Let state Z be NS random. Let $\{\rho_n\}$ be the density matrices associated with Z . Suppose not. Then there is $\delta \in (0, 1)$ and computable measurement system $B = \{|b_0^n\rangle, |b_1^n\rangle\}_{n=1}^\infty$ where $\mu_Z^B(\{0, 1\}^\infty \setminus \text{MLR}) > \delta$. Let $\{S^m\}$ be a universal ML test. Without loss of generality, this test is of the form

$$S^m = \bigcup_{m \leq i} \llbracket A_i^m \rrbracket,$$

where $\llbracket A_i^m \rrbracket \subseteq \llbracket A_{i+1}^m \rrbracket$, and $A_i^m \{\tau_1^{m,i}, \dots, \tau_{k^{m,i}}^{m,i}\} \subset \{0, 1\}^i$ for some $0 \leq k^{m,i} \leq 2^{i-m}$. Thus $\mu(S^m) \leq 2^{-m}$, where μ is the uniform distribution over $\{0, 1\}^\infty$. We define an NS test as follows. For all m and i , with $m \leq i$, let $\tau_a = \tau_a^{m,i}$ and define the special projection

$$p_i^m = \sum_{a \leq k^{m,i}} |\otimes_{q=1}^i b_{\tau_a[q]}^q\rangle \langle \otimes_{q=1}^i b_{\tau_a[q]}^q|.$$

We define $P^m = \{p_i^m\}_{m \leq i}$ we have that $\langle P^m \rangle$ is an NS Test. The special tests p_i^m is uniformly computable in i and m since B and A_i^m are uniformly computable in i and m . Since $\llbracket A_i^m \rrbracket \subseteq \llbracket A_{i+1}^m \rrbracket$, $\text{Range}(p_i^m) \subseteq \text{Range}(p_{i+1}^m)$. So P^m is an NS Σ_1^0 set for all m . Since $k^{m,i} \leq 2^{i-m}$ for all m and i , this implies $\tau(P^m) \leq 2^{-m}$ for all m .

For all m , $\{0, 1\}^\infty \setminus \text{MLR} \subseteq S^m$. Since by assumption $\mu_Z^B(\{0, 1\}^\infty \setminus \text{MLR}) > \delta$, for all m there exists $i(m) > m$ such that

$$\mu_Z^B(\llbracket A_{i(m)}^m \rrbracket) > \delta.$$

Fix an m and $i = i(m)$ and let $A_i^m = \{\tau_1, \dots, \tau_{k^{m,i}}\}$, where $k^{m,i} \leq 2^{i-m}$. Let p be the pre-measure associated with μ_Z^B . So we have

$$\delta < \sum_{a \leq k^{m,i}} p(\tau_a) = \sum_{a \leq k^{m,i}} \text{Tr} \rho_i |\otimes_{q=1}^i b_{\tau_a[q]}^q\rangle \langle \otimes_{q=1}^i b_{\tau_a[q]}^q| = \text{Tr} \rho_i \sum_{a \leq k^{m,i}} |\otimes_{q=1}^i b_{\tau_a[q]}^q\rangle \langle \otimes_{q=1}^i b_{\tau_a[q]}^q|$$

So we see that for all m there is an i such that

$$\delta < \text{Tr} \rho_i p_i^m \leq Z(P^m).$$

So $\inf_m Z(P^m) > \delta$, contradicting that Z is NS random. \square

Theorem 45 ([[Bho21](#)]) *There are states that are Bhojraj random and not NS Random.*

8.9 NS Solovay States

A NS Solovay test is a sequence of NS Σ_0^1 sets $\langle G_n \rangle$ such that $\sum_n \tau(G_n) < \infty$. A state Z fails a quantum NS test $\langle G^r \rangle$ at order $\delta \in (0, 1)$ if there is an infinite number of $r \in R$ such that $\inf_{r \in R} Z(G^r) > \delta$. Otherwise state Z passes the quantum NS test at order δ . A quantum state Z is NS Solovay random if it passes all NS Solovay tests at all orders. The following theorem shows the equivalence of NS randomness and NS Solovay randomness with respect to every order δ . Given a special projection p , NS Σ_0^1 set $Q = \{q_n\}$, and state Z , we define $Z(p \setminus Q) = \inf_n Z(p \setminus q_n)$. In [[Bho21](#)], it was proven that NS randomness is equivalent to NS Solovay randomness.

Proposition 4 *Given a special projection p , NS Σ_0^1 set Q , and state Z , $Z(p) - Z(Q) \leq Z(p \setminus Q) \leq Z(p)$.*

The proof is straightforward.

Theorem 46 *If a state Z fails an NS test at order δ then it fails an NS Solovay test at order δ .*

Proof. Assume that state Z fails a NS test $\langle G^r \rangle$ at order δ . Since $\sum_r \tau(G^r) \leq 1$, and each G^r is an NS Σ_1^0 set, $\langle G^r \rangle$ is a NS Solovay test. Furthermore since $\inf_r Z(G^r) \geq \delta$, there exists an infinite number of r such that $Z(G^r) > \delta$. Thus Z fails a NS Solovay test at order δ . \square

Theorem 47 *For all $\delta' < \delta$, if a state Z fails an NS Solovay test at order δ then it fails an NS test at order δ' .*

Proof. Assume state Z fails NS Solovay test $\langle G^r \rangle$ at order δ . Given $\langle G^r \rangle$, where $G^r = \langle p_n^r \rangle_{n \in \mathbb{N}}$, we construct an NS test $\langle H^r \rangle$ as follows. There exists an m such that $\sum_{n > m} \tau(G^n) \leq 1$. Fix r . Enumerate all unordered sets of $r + 1$ natural numbers $\{D_n^r\}_{n \in \mathbb{N}}$, $D_n^r \subset \mathbb{N}$, with infinite repetition.

$$H^r = \{q_n^r\}, q_n^r = \bigvee_{m < n} q_m^r \bigvee \left(\bigwedge_{t \in D_n^r} p_n^t \right).$$

Each H^r can be seen to be an NS Σ_1^0 set. Furthermore $\tau(H^r) \leq \sum_{t > r} Z(G^t) \leq \sum_{t > r} 2^{-t} = 2^{-r}$. So $\langle H^r \rangle$ is an NS test. For each r , $Z(H^r) > \delta'$. Assume not. Then there exists a k such that $Z(H^k) \leq \delta'$. Since Z fails $\langle G^r \rangle$ at order δ , there exists an infinite number of $r \in R$ and $n_r \in \mathbb{N}$ such that $Z(p_{n_r}^r) \geq \delta''$, for some $\delta' < \delta'' < \delta$. We reorder the NS Solovay test $\langle G^r \rangle$ such that r ranges over solely R . Let $z_r = p_{n_r}^r$. Let $D_{n,k}$ be the set of all unordered subsets of $\{1, \dots, n\}$ of size k . For $k > n$ let $F_{n,k} = \emptyset$. Let

$$F_{n,k} = \left(\bigvee_{A \in D_{n,k}} \bigwedge_{r \in A} z^r \right) \setminus \bigvee_{s > k} F_{n,s}.$$

So for all $n \in \mathbb{N}$, using Proposition 4,

$$\begin{aligned} & n(\delta'' - \delta') \\ & \leq \sum_{r=1}^n (Z(z^r) - Z(H^k)) \\ & \leq \sum_{r=1}^n Z(z^r \setminus H^k) \\ & \leq \sum_{r=1}^n Z \left(\bigvee_{s=1}^k F_{n,s} \wedge z^r \right) \end{aligned} \tag{8.1}$$

Equation 8.1 is due to the fact that for $s > k$ there is a t where we have $\text{Range}(F_{n,s}) \leq \text{Range}(q_s^k)$. Let $F_{n,s,r} = F_{n,s} \wedge z^r$, with for a fixed $s \leq k$, $\sum_{i=1}^n Z(F_{n,s,i}) \leq s$.

$$\begin{aligned} & n(\delta'' - \delta') \\ & \leq \sum_{r=1}^n Z \left(\bigvee_{s=1}^k F_{n,s,r} \right) \\ & = \sum_{s=1}^k \sum_{r=1}^n Z(F_{n,s,r}) \\ & \leq \sum_{s=1}^k s = O(k^2). \end{aligned}$$

This is a contradiction for large enough n . \square

Corollary 10 *A quantum state is NS random if and only if it is NS Solovay random.*

Chapter 9

The Many Worlds Theory

The Many Worlds Theory (**MWT**) was formulated by Hugh Everett [Eve57] as a solution to the measurement problem of Quantum Mechanics. Branching (a.k.a splitting of worlds) occurs during any process that magnifies microscopic superpositions to the macro-scale. This occurs in events including human measurements such as the double slit experiments, or natural processes such as radiation resulting in cell mutations.

One question is if **MWT** causes issues with the foundations of computer science. The physical Church Turing Thesis (**PCTT**) states that any functions computed by a physical system can be simulated by a Turing machine. A straw man argument for showing **MWT** and **PCTT** are in conflict is an experiment that measures the spin of an unending number of electrons, with each measurement bifurcating the current branch into two sub-branches. This results in a single branch in which the halting sequence is outputted. However this branch has Born probability converging to 0, and can be seen as a deviant, atypical branch.

In fact, conflicts do emerge between **MWT** and Algorithmic Information Theory. In particular, the Independence Postulate (**IP**) is a finitary Church-Turing thesis, postulating that certain infinite and *finite* sequences cannot be found in nature, a.k.a. have high “addresses”. If a forbidden sequence is found in nature, an information leak will occur. However **MWT** represents a theory in which such information leaks can occur. This blog entry covers the main arguments of this conflict.

9.1 Many Worlds Theory

Some researchers believe there is an inherent problem in quantum mechanics. On one hand, the dynamics of quantum states is prescribed by unitary evolution. This evolution is deterministic and linear. On the other hand, measurements result in the collapse of the wavefunction. This evolution is non-linear and nondeterministic. This conflict is called the measurement problem of quantum mechanics.

The time of the collapse is undefined and the criteria for the kind of collapse are strange. The Born rule assigns probabilities to macroscopic outcomes. The projection postulate assigns new microscopic states to the system measured, depending on the the macroscopic outcome. One could argue that the apparatus itself should be modeled in quantum mechanics. However it’s dynamics is deterministic. Probabilities only enter the conventional theory with the measurement postulates.

MWT was proposed by Everett as a way to remove the measurement postulate from quantum mechanics. The theory consists of unitary evolutions of quantum states without measurement collapses. For **MWT**, the collapse of the wave function is the change in dynamical influence of one

part of the wavefunction over another, the decoherence of one part from the other. The result is a branching structure of the wavefunction and a collapse only in the phenomenological sense.

9.1.1 Branching Worlds

An example of a branching of universes can be seen in an idealized experiment with a single electron with spin $|\phi_\uparrow\rangle$ and $|\phi_\downarrow\rangle$. This description can be found in [SBKW10]. There is a measuring apparatus \mathcal{A} , which is in an initial state of $|\psi_{\text{ready}}^{\mathcal{A}}\rangle$. After \mathcal{A} reads spin-up or spin-down then it is in state $|\psi_{\text{reads spin } \uparrow}^{\mathcal{A}}\rangle$ or $|\psi_{\text{reads spin } \downarrow}^{\mathcal{A}}\rangle$, respectively. The evolution for when the electron is solely spin-up or spin-down is

$$\begin{aligned} |\phi_\uparrow\rangle \otimes |\psi_{\text{ready}}^{\mathcal{A}}\rangle &\xrightarrow{\text{unitary}} |\phi_\uparrow\rangle \otimes |\psi_{\text{reads spin } \uparrow}^{\mathcal{A}}\rangle \\ |\phi_\downarrow\rangle \otimes |\psi_{\text{ready}}^{\mathcal{A}}\rangle &\xrightarrow{\text{unitary}} |\phi_\downarrow\rangle \otimes |\psi_{\text{reads spin } \downarrow}^{\mathcal{A}}\rangle. \end{aligned}$$

Furthermore, one can model the entire quantum state of an observer \mathcal{O} of the apparatus, with

$$\begin{aligned} &|\phi_\uparrow\rangle \otimes |\psi_{\text{ready}}^{\mathcal{A}}\rangle \otimes |\xi_{\text{ready}}^{\mathcal{O}}\rangle \\ &\xrightarrow{\text{unitary}} |\phi_\uparrow\rangle \otimes |\psi_{\text{reads spin } \uparrow}^{\mathcal{A}}\rangle \otimes |\xi_{\text{ready}}^{\mathcal{O}}\rangle \\ &\xrightarrow{\text{unitary}} |\phi_\uparrow\rangle \otimes |\psi_{\text{reads spin } \uparrow}^{\mathcal{A}}\rangle \otimes |\xi_{\text{reads spin } \uparrow}^{\mathcal{O}}\rangle \\ &|\phi_\downarrow\rangle \otimes |\psi_{\text{ready}}^{\mathcal{A}}\rangle \otimes |\xi_{\text{ready}}^{\mathcal{O}}\rangle \\ &\xrightarrow{\text{unitary}} |\phi_\downarrow\rangle \otimes |\psi_{\text{reads spin } \downarrow}^{\mathcal{A}}\rangle \otimes |\xi_{\text{ready}}^{\mathcal{O}}\rangle \\ &\xrightarrow{\text{unitary}} |\phi_\downarrow\rangle \otimes |\psi_{\text{reads spin } \downarrow}^{\mathcal{A}}\rangle \otimes |\xi_{\text{reads spin } \downarrow}^{\mathcal{O}}\rangle. \end{aligned}$$

For the general case, the electron is in a state $|\phi\rangle = a|\phi_\uparrow\rangle + b|\phi_\downarrow\rangle$, where $|a|^2 + |b|^2 = 1$. In this case, the final superposition would be of the form:

$$\begin{aligned} &a|\phi_\uparrow\rangle \otimes |\psi_{\text{reads spin } \uparrow}^{\mathcal{A}}\rangle \otimes |\xi_{\text{reads spin } \uparrow}^{\mathcal{O}}\rangle \\ &+ b|\phi_\downarrow\rangle \otimes |\psi_{\text{reads spin } \downarrow}^{\mathcal{A}}\rangle \otimes |\xi_{\text{reads spin } \downarrow}^{\mathcal{O}}\rangle. \end{aligned}$$

This is a superposition of two branches, each of which describes a perfectly reasonable physical story. This bifurcation is one method on how the quantum state of universe bifurcates into two branches.

9.1.2 Deriving the Born Rule

In my opinion, one of the main problems of **MWT** is its reconciliation of the Born rule, for which no proposed solution has universal consensus. In standard quantum mechanics, measurements are probabilistic operations. Measurements on a state vector $|\psi\rangle$, which is a unit vector over Hilbert space \mathcal{H} , are self-adjoint operators \mathcal{O} on \mathcal{H} . Observables are real numbers that are the spectrum $\text{Sp}(\mathcal{O})$ of \mathcal{O} . A measurement outcome is a subset $E \subseteq \text{Sp}(\mathcal{O})$ with associated projector P_E on \mathcal{H} . Outcome E is observed on measurement of \mathcal{O} on $|\psi\rangle$ with probability $P(E) = \langle\psi|P_E|\psi\rangle$. This is known as the Born rule. After this measurement, the new state becomes $P_E|\psi\rangle / \sqrt{\langle\psi|P_E|\psi\rangle}$. This is known as the projection postulate.

However, the Born rule and the projection postulate are not assumed by **MWT**. The dynamics are totally deterministic. Each branch is equally real to the observers in it. To address these issues, Everett first derived a typicality-measure that weights each branch of a state's superposition. Assuming a set of desirable constraints, Everett derived the typicality-measure to be equal to the norm-squared of the coefficients of each branch, i.e. the Born probability of each branch. Everett then drew a distinction between typical branches that have high typicality-measure and exotic atypical branches of decreasing typicality-measure. For the repeated measurements of the spin of an electron $|\phi\rangle = a|\phi_{\uparrow}\rangle + b|\phi_{\downarrow}\rangle$, the relative frequencies of up and down spin measurements in a typical branch converge to $|a|^2$ and $|b|^2$, respectively. The notion of typicality can be extended to measurements with many observables.

In a more recent resolution to the relation between **MWT** and probability, Deutsch introduced a decision theoretic interpretation [Deu99] that obtains the Born rule from the non-probabilistic axioms of quantum theory and non-probabilistic axioms of decision theory. Deutsch proved that rational actors are compelled to adopt the Born rule as the probability measure associated with their available actions. This approach is highly controversial, as some critics say the idea has circular logic.

Another attempt uses subjective probability [Vai98]. The experimenter puts on a blindfold before he finishes performing the experiment. After he finishes the experiment, he has uncertainty about what world he is in. This uncertainty is the foundation of a probability measure over the measurements. However, the actual form of the probability measure needs to be postulated:

Probability Postulate. *An observer should set his subjective probability of the outcome of a quantum experiment in proportion to the total measure of existence of all worlds with that outcome.*

Whichever explanation of the Born rule one adopts, the following section shows there is an issue with **MWT** and **IP**. There exist branches of substantial Born probability where information leaks occurs.

9.2 Violating the Independence Postulate

In [Lev84, Lev13], the Independence Postulate, **IP**, was introduced:

Let $\alpha \in \{0,1\}^{\infty}$ be a sequence defined with an n -bit mathematical statement (e.g., in Peano Arithmetic or Set Theory), and a sequence $\beta \in \{0,1\}^{*\infty}$ can be located in the physical world with a k -bit instruction set (e.g., ip-address). Then $\mathbf{I}(\alpha : \beta) < k + n + c_{\text{IP}}$, for some small absolute constant c_{IP} .*

The **I** term is an information measure in Algorithmic Information Theory. For this blog, the information term we use is $\mathbf{I}(x : y) = \mathbf{K}(x) + \mathbf{K}(y) - \mathbf{K}(x, y)$, where \mathbf{K} is the prefix-free Kolmogorov complexity. We can use this definition of **I** because we only deal with finite sequences.

Let Ω_m be the first m bits of Chaitin's Omega (the probability that a universal Turing machine will halt). We have $m <^+ \mathbf{K}(\Omega_m)$. Furthermore Ω_m can be described by a mathematical formula of size $O(\log m)$. Thus by **IP**, where $\Omega_m = \alpha = \beta$, Ω_m can only be found with addresses of size at least $m - O(\log m)$.

IP can be violated in the following idealized experiment measuring the spin $|\phi_\uparrow\rangle$ and $|\phi_\downarrow\rangle$ of N isolated electrons. We denote $|\phi_0\rangle$ for $|\phi_\uparrow\rangle$ and $|\phi_1\rangle$ for $|\phi_\downarrow\rangle$. The “address” (in the sense of **IP**) of this experiment is $< O(\log N)$. The measuring apparatus will measure the spin of N electrons in the state $|\phi\rangle = \frac{1}{2}|\phi_\uparrow\rangle + \frac{1}{2}|\phi_\downarrow\rangle$. There is a measuring apparatus \mathcal{A} with initial state of $|\psi^{\mathcal{A}}\rangle$, and after reading N spins of N electrons, it is in the state $|\psi^{\mathcal{A}}[x]\rangle$, where $x \in \{0, 1\}^N$, whose i th bit is 1 iff the i th measurement returns $|\phi_1\rangle$.

The experiment evolves according to the following unitary transformation:

$$\bigotimes_{i=1}^N |\phi\rangle \otimes |\psi^{\mathcal{A}}\rangle \xrightarrow{\text{unitary}} \sum_{a_1, \dots, a_N \in \{0, 1\}^N} 2^{-N/2} \bigotimes_{i=1}^N |\phi_{a_i}\rangle \otimes |\psi^{\mathcal{A}}[a_1 a_2 \dots a_N]\rangle.$$

If the bits returned are Ω_N then a memory leak of size $N - O(\log N)$ has occurred, because Ω_N has been located by the address of the experiment, which is $O(\log N)$. Thus

$$\text{Born-Probability(a memory leak of size } N - O(\log N) \text{ occurred)} \geq 2^{-N}.$$

9.3 Conclusion

There are multiple variations of **MWT** when it comes to consistency across universes. In one formulation, all universes conform to the same physical laws. In another model, each universe has its own laws, for example different values of gravity, etc. However, the experiment in the previous section shows that mathematics itself is different between universes, regardless of which model is used. In some universes, **IP** holds and there is no way to create information leaks. In other universes information leaks occur, and there are tasks where randomized algorithms fail but non-algorithmic physical methods succeeds. One such task is finding new axioms of mathematics. This was envisioned as a possibility by Gödel [Gö61], but there is a universal consensus of the impossibility of this task. Not any more! In addition, because information leaks are finite events, the Born probability of worlds containing them is not insignificant. In such worlds, **IP** cannot be formulated, and the foundations of Algorithmic Information Theory itself become detached from reality.

Formulated another way, let us suppose the Born probability is derived from the probability postulate. We have a “blindfolded mathematician” who performs the experiment described above. Before the mathematician takes off her blindfold, she states the Independence Postulate. By the probability postulate, with measure 2^{-N} over all worlds, there is a memory leak of size $N - O(\log N)$ and **IP** statement by the mathematician is in error.

As a rebuttal, one can, with non-zero probability, just flip a coin N times and get N bits of Chaitin's Omega. Or more generally, how does one account for a probability P over finite or infinite sequences learning information about a forbidden sequence β with good probability? Due

to probabilistic conservation laws [Lev74, Lev84], we have

$$\Pr_{\alpha \sim P} [\mathbf{I}(\alpha : \beta) > \mathbf{I}(\langle P \rangle : \beta) + m] \stackrel{*}{<} 2^{-m}.$$

Thus the probability of a single event creating a leak is very small. However if many events occur, then the chances of a memory leak grows. However as there is many events, to locate one such leak, one will probably need a long address to find the leak, balancing out the **IP** equation.

This still leaves open the possibility of a memory leak occurring at an event with a small address. Since there are a small number of events that have a small address, the probability of a significant memory leak is extremely small. In physics one can postulate away events with extremely small probabilities. For example, the second law of thermodynamics states that entropy is non-decreasing, postulating away the extremely unlikely event that a large system suddenly decreases in thermodynamic entropy.

There is no way to postulate such memory leaks in **MWT**. Assuming the *probability postulate*, probability is a measure over the space of possible worlds. Thus when Bob now threatens to measure the spin of N particles, Alice now knows 2^{-N} of the resultant worlds will contain N bits of Chaitin's Omega, violating **IP**.

Chapter 10

Conclusion

10.1 Signals from Classical and Quantum Sources

Information non-growth laws say information about a target source cannot be increased with randomized processing. In classical information theory, we have

$$I(g(X):Y) \leq I(X:Y).$$

where g is a randomized function, X and Y are random variables, and I is the mutual information function. Thus processing a channel at its output will not increase its capacity. Information conservation carries over into the algorithmic domain, with the inequalities

$$\mathbf{I}(f(x):y) <^+ \mathbf{I}(x:y); \quad \mathbf{I}(f(a);\mathcal{H}) <^+ \mathbf{I}(a;\mathcal{H}).$$

These inequalities ensure target information cannot be obtained by processing. If for example the second inequality was not true, then one can potentially obtain information about the halting sequence \mathcal{H} with simple functions. Obtaining information about \mathcal{H} violates the Independence Postulate, discussed in Chapter 9. Information non growth laws can be extended to signals [Eps23a] which can be modeled as probabilities over \mathbb{N} or Euclidean space¹. The “signal strength” of a probability p over \mathbb{N} is measured by its self information.

$$\mathbf{I}_{\text{Prob}}(p:p) = \log \sum_{i,j} 2^{\mathbf{I}(i:j)} p(i)p(j).$$

A signal, when undergoing randomized processing f (see Section 3.1.1), will lose its cohesion. Thus any signal going through a classical channel will become less coherent.

$$\mathbf{I}_{\text{Prob}}(f(p):f(p)) <^+ \mathbf{I}_{\text{Prob}}(p:p).$$

In Euclidean space, probabilities that undergo convolutions with probability kernels will lose self information. For example a signal spike at a random position will spread out when convoluted with the Gaussian function, and lose self information. The above inequalities deal with classical transformations. One can ask, is whether, quantum information processing can add new surprises to how information signals occur and evolve.

One can start with the prepare-and-measure channel, also known as a Holevo-form channel. Alice starts with a random variable X that can take values $\{1, \dots, n\}$ with corresponding probabilities $\{p_1, \dots, p_n\}$. Alice prepares a quantum state, corresponding to density matrix ρ_X , chosen

¹In [Eps23a] probabilities over $\{0,1\}^\infty$ and T_0 second countable topologies were also studied.

from $\{\rho_1, \dots, \rho_n\}$ according to X . Bob performs a measurement on the state ρ_X , getting a classical outcome, denoted by Y . Though it uses quantum mechanics, this is a classical channel $X \rightarrow Y$. So using the above inequality, cohesion will deteriorate regardless of X 's probability, with

$$\mathbf{I}_{\text{Prob}}(Y : Y) <^+ \mathbf{I}_{\text{Prob}}(X : X).$$

There remains a second option, constructing a signal directly from a mixed state. This involves constructing a mixed state, i.e. density matrix σ , and then performing a measurement E on the state, inducing the probability $E\sigma(k) = \text{Tr}\sigma E_k$. However from [Eps23a], for elementary (even enumerable) probabilities $E\sigma$,

$$\mathbf{I}_{\text{Prob}}(E\sigma : E\sigma) <^+ \mathbf{K}(\sigma, E).$$

Thus for simply defined density matrices and measurements, no signal will appear. So experiments that are simple will result in simple measurements, or white noise. However it could be that a larger number of uncomputable pure or mixed states produce coherent signals. Theorems 35 and 36 say otherwise, in that the POVM measurement E of a vast majority of pure and mixed states will have negligible self-information. Thus for uniform distributions Λ and μ over pure and mixed states (see Section 6.2.2),

$$\int 2^{\mathbf{I}_{\text{Prob}}(E|\psi\rangle : E|\psi\rangle)} d\Lambda = O(1); \quad \int 2^{\mathbf{I}_{\text{Prob}}(E\sigma : E\sigma)} d\mu(\sigma) = O(1).$$

This can be seen as a consequence of the vastness of Hilbert spaces as opposed to the limited discriminatory power of quantum measurements. In addition, there could be non-uniform distributions of pure or mixed states that could be of research interest. In quantum decoherence, a quantum state becomes entangled with the environment, losing decoherence. The off diagonal elements of the mixed state become dampened, as the state becomes more like a classical mixture of states. Let p_σ be the idealized classical probability that σ decoheres to, with $p_\sigma(i) = \sigma_{ii}$. Corollary 8 states that for an overwhelming majority of pure or mixed states σ , p_σ is noise, that is, has negligible self-information.

$$\int 2^{\mathbf{I}_{\text{Prob}}(p_{|\psi\rangle} : p_{|\psi\rangle})} d\Lambda = O(1); \quad \int 2^{\mathbf{I}_{\text{Prob}}(p_\sigma : p_\sigma)} d\mu(\sigma) = O(1).$$

This is to be expected, with one supporting fact being for an n qubit space, $i \in \{1, \dots, 2^n\}$, $\mathbf{E}_\Lambda[p_{|\psi\rangle}(i)] = 2^{-n}$. With Algorithmic Information Theory, we've taken this fact one step further, showing that $p_{|\psi\rangle}$ has no (in the exponential) self-algorithmic information and cannot be processed by deterministic or randomized means to produce a more coherent signal. In addition, it appears a more direct proof of the first decoherence inequality could be possible.

However the measurement process has a surprising consequence, in that the wave function collapse causes an massive uptake in algorithmic signal strength. Let F be a PVM, of size 2^{n-c} , of an n qubit space and let Λ_F be the distribution of pure states when F is measured over the uniform distribution Λ . Thus Λ_F represents the F -collapsed states from Λ . Theorem 37 states

$$n - 2c <^{\log \log} \int 2^{\mathbf{I}_{\text{Prob}}(F|\psi\rangle : F|\psi\rangle)} d\Lambda_F.$$

10.2 Apriori Distributions

To avoid the pitfall of a signalless distribution that only produces white noise, we can conjecture a new apriori distribution for quantum states that is not signalless. Note that we are dealing with



Figure 10.1: Each box on the top row represents an n qubit Hilbert space, with the shaded rectangles being the subspaces of the PVM projectors. Thus there are three PVMs. The self-information majorizes these subspaces, inversely weighted by the PVM's complexity.

measures over the density operator space and not directly with density operators because we are measuring properties, such as self-information, over all possible (pure or mixed) states. Properties of this apriori distribution can be discerned by working backwards. Indeed, suppose there are a set of (possibly infinite) systems $\{|\psi_i\rangle\}$, where for each system $|\psi_i\rangle$, a measurement occurs, producing a discernable signal. By Theorem 34, this implies the states $|\psi_i\rangle$ have high $\mathbf{I}(|\psi_i\rangle : |\psi_i\rangle)$, where \mathbf{I} is the information function between mixed states introduced in Definition 9. Thus any universal quantum apriori distribution over these systems must be weighted toward states with high self information. One candidate is an probability measure ξ over pure states where

$$\xi(|\psi\rangle) \propto 2^{\mathbf{I}(|\psi\rangle : |\psi\rangle)}.$$

However this area of research is still ongoing. Another clue to this universal quantum apriori distribution is the measurement operation, which as shown above, causes an uptake in signal strength. Take a PVM measurement F , which procures a value i from a state $|\psi\rangle$, projecting to a new state $|\psi'\rangle$. $P|\psi'\rangle(i) = 1$. By Corollary 7, this new state $|\psi'\rangle$ has self information

$$\mathbf{K}(i) <^{\log} \mathbf{I}(|\psi'\rangle : |\psi'\rangle).$$

The error term is on the order of $\mathbf{K}(P)$. Most of the measurement values i of P will be random, i.e. have large $\mathbf{K}(i)$ (just look at the Kolmogorov complexity of the first 2^n numbers!). Thus simple quantum measurements increase the self information of most measured quantum states (see Figure 10.1). So this fact, and Theorem 29, leads us to the following conclusion.

Take a distribution over density operators, such as Λ , where an overwhelming majority of states have negligible self-information. When each such state in its support is mea-

sured with a simple apparatus, the result is new a distribution where most of the states have substantial self-information.

However, the situation is reversed for quantum channels. A quantum state that is transformed by a quantum operation will not increase in self-information. So by Theorem 31, we get the following claim, where equality occurs if the quantum operation is a unitary transform.

Given any distribution over density operators, if all the density matrices its support are transformed by a simple quantum operation, then the resultant distribution will give more measure to mixed states with less self-information.

Thus simple measurements with many operators can only increase self-information, simple quantum operations can only decrease self-information, and simple unitary transforms leave the self-information unaltered. If the operation is complex, then nothing so far has been proven.

10.3 Measurements Before Information Cloning

The no-cloning theorem states that every unitary transform cannot clone an arbitrary quantum state. However there is the possibility of copying information from a subset of states. By “copying information”, we mean that two measurements of two states will produce two values that are similar. More formally, the information cloned from a state $|\psi\rangle$ relative to unitary transform U , and POVMs E and F is,

$$\mathbf{I}_{\text{Clone}}(|\psi\rangle) = \mathbf{I}_{\text{Prob}}(E|\phi_1\rangle : F|\phi_2\rangle), \text{ where } U|\psi\rangle|0\rangle = |\phi_1\rangle|\phi_2\rangle.$$

This represents the shared signal strength between $|1\rangle$ and $|2\rangle$ when the states² were created from a unitary transform U of $|\psi\rangle$ tensored with an ancilla state $|0\rangle$. Note that by Theorems 32 and 34, cloneable information is less than self information, with

$$\mathbf{I}_{\text{Clone}}(|\psi\rangle) <^{\log} \mathbf{I}(|\psi\rangle : |\psi\rangle).$$

The question is, given an initial distribution over density operators with low expected $\mathbf{I}_{\text{Clone}}$, what sort of transform is required to increase this expectation. In this section, we discuss necessary conditions of this transform. We require the following two assumptions.

Assumption (1): The initial distribution has low expected self information. Theorem 29 shows there is a large set of natural distributions that have this property. Any distribution Ω that is less than $2^c\Lambda$ will have $\log \int 2^{\mathbf{I}(|\psi\rangle : |\psi\rangle)} d\Omega <^+ c$. Another way to interpret this assumption is through parameterized distributions. Let P be a probability over parameters θ over pure state distribution, $\Gamma(|\psi\rangle|\theta)$. The distribution is balanced, where $\int \Gamma(|\psi\rangle|\theta) dP(\theta) = \Lambda(|\psi\rangle)$. Then because of Theorem 29,

$$P(\{\theta : \mathbf{E}_{|\psi\rangle \sim \Gamma(\cdot|\theta)}[\mathbf{I}(|\psi\rangle : |\psi\rangle)] \geq m\}) \leq 2^{-m+1}.$$

Assumption (2): The universal Turing machine is relativized to all the transforms and operators. This assumption states that for a system, the operations are known quantities. This is congruent with quantum information theory, in which actors are seen to compute unitary transforms or quantum operations. It is asumed that these actors have knowledge of the transforms.

²Note this definition can be generalized to arbitrary states, with $\mathbf{I}_{\text{Prob}}(E\text{Tr}_2\sigma : F\text{Tr}_1\sigma)$, where $\sigma = \varepsilon(|\psi\rangle)$, for quantum operation ε .



Figure 10.2: The initial distribution has low self information and cloneable information. A measurement increases the self information and potentially increases the cloneable information.

How do you create a distribution with high expected $\mathbf{I}_{\text{Clone}}$, where most states can have cloneable information? Any transform that increases cloneable information must increase self-information. However Theorem 31, along with assumption (2) bars quantum operations as a means to create self-information, as the complexity of the quantum operations is $O(1)$. Thus the only way to potentially increase self-information is to perform a measurement, which as Theorems 34 and 37 show, often times cause an uptake in self-information (see Figure 10.2). This is also discussed in the quotes of Section 10.2. Thus we get the following claim.

Measurements are required to produce distributions over quantum states that have cloneable information.

For example, take the starting distribution to be the uniform measure over pure states, Λ . Let $E = F = \{|i\rangle\langle i|\}$ be POVM measurements over projectors to the basis states and let U be any unitary transform such that $U|i\rangle|0\rangle = |i\rangle|i\rangle$ for $i \in \{1, \dots, 2^n\}$. By Theorems 29 and 32, we have that

$$\int 2^{\mathbf{I}_{\text{Clone}}(|\psi\rangle)} d\Lambda = O(1).$$

Now suppose we apply the measurement $G = E$ to Λ , producing a new distribution Λ_G concentrated evenly among the basis states, where $\Lambda_G(|i\rangle) = 2^{-n}$. Thus we have that $\mathbf{I}_{\text{Clone}}(|i\rangle) = \mathbf{I}_{\text{Prob}}(E|i) : F|i\rangle) = \mathbf{K}(i)$. Since there are $2^{n-O(1)}$ basis states $|i\rangle$ where $n <^+ \mathbf{K}(i)$, we have the following uptake in cloneable information.

$$n <^+ \log \int 2^{\mathbf{I}_{\text{Clone}}(|\psi\rangle)} d\Lambda_G.$$

Other such applications can be seen as generalizations from this extreme example. Future work involves determining how tightly self information covers cloneable information.

Part III

Thermodynamics

Classical thermodynamics is the study of substances and changes to their properties such as volume, temperature, and pressure. Substances, such as a gas or a liquid, is modeled as a point in a phase space. The phase space, \mathcal{X} , is modeled by a computable metric space, [HR09], and a volume measure μ , is modeled by a computable (not necessarily probabilistic) positive measure over \mathcal{X} . The dynamics are modeled by a one dimensional transformation group G^t , indexed by $t \in \mathbb{R}$. Due to Louville’s theorem, the dynamics are measure-preserving, where $\mu(G^t A) = \mu(A)$, for all Borel sets $A \subseteq \mathcal{X}$.

Whether quantum or classical, the known laws of physics are reversible. Thus the dynamics G of our system are also reversible, in that if $\beta = G^t \alpha$, then there is some t' such that the original state can found with $\alpha = G^{t'} \beta$. Thus if given a set of particles with position and velocity, by reversing the velocities, a previous state can be found. This is contradiction to the second law of thermodynamics, which states,

The total entropy of a system either increases or remains constant in any spontaneous process; it never decreases.

This conforms to our experiences of broken vases never reforming. To reconcile this difference, Boltzmann introduced *macro-states*, Π_i , indexed by $i \in \mathbb{N}$, which groups states together by macroscopic parameters, with corresponding Boltzmann entropy $S(\Pi_i) = k_B \ln \mu(\Pi_i)$. By definition, a vast majority of typical states will experience an increase in Boltzmann entropy.

In [Gac94], coarse grained entropy was introduced as an algorithmic update to Boltzmann entropy. This formulation was made to be independent of the choice of parameters of the macro state. In this paper, we introduce a modified version of coarse grained entropy. We also model the thermodynamic entropy of a micro-state with algorithmic methods. The micro-state of a system contains the information of the entire physical state. For example, the microstate of a system of N molecules is a point

$$(q_1, \dots, q_{3N}, p_1, \dots, p_{3N}) \in \mathbb{R}^{6N}$$

where q_i are the position coordinates and p_i are the momentum coordinates. The set of states, \mathbb{R}^{3N} is a computable metric space. To model the entropy of the state, we use slight variant to algorithmic fine-grained entropy H_μ in [Gac94], using symbol \mathbf{H}_μ . This entropy measure captures the level of disorder of the state. Continuing the example above, if all the particles are at rest, then the thermodynamic entropy of the state of

$$(q_1, \dots, q_{3N}, 0, \dots, 0)$$

is expected to be very low.

The evolution of the system will be thermodynamic like if it spends most of the time close to its maximum value, from which it exhibits frequent small fluctuations and rarer large fluctuations.

In this paper, using the algorithmic definition of thermodynamic entropy, \mathbf{H}_μ , we prove that such fluctuations *have* to occur, and the greater the fluctuation, the lesser its measure.

Throughout the part on thermodynamics, conditions will be proven for states, sets of states, dynamics, and measures to have high or infinite mutual information in the halting sequence. Assuming the Independence Postulate (see Chapter 9), such constructs are non-physical and can be considered non-realizable in nature. This also goes for the part of Quantum Mechanics, in particular the Quantum EL Theorem 21 and the Quantum Outlier Theorem 26.

The thermodynamics section of this manuscript is arranged as follows.

- **Chapter 11:** Computable metric spaces and their relation to randomness is detailed. This material is the foundation for which algorithmic coarse and fine grained entropy is based upon. This chapter is a modification to the work in [HR09] to arbitrary positive measures and dual measure spaces.
- **Chapter 12:** Algorithmic fine grained entropy is introduced. This is a modification to the definition in [Gac94], using computable measure theory. An entropy balance lemma is proven with applications to Maxwell's demon. We detail a result from [G21] that algorithmic fine grain entropy is conformant to the addition inequality analagous to that of string algorithmic information theory.
- **Chapter 13:** In this chapter, algorithmic fine grained entropy is proved to oscillate in the presence of dynamics, regardless of the choice of phase space and volume measure.
- **Chapter 14:** Discrete dynamics are studied in this chapter. It is proved that given a phase space and two different volume measures of it, the algorithmic thermodynamic entropy will oscillate in a synchronized fashion with respect to both volume measures.
- **Chapter 15:** It is shown in the course of dynamics on product spaces, for typical states, the marginal entropies cannot be in sync. This is true for continuous dynamics and discrete ergodic dynamics.
- **Chapter 16:** Algorithmic coarse grained entropy is defined and shown to be an excellent approximation to algorithmic fine grained entropy. Algorithmic coarse grained entropy is proved to oscillate in the presence of dynamics.
- **Chapter 17:** In this chapter, computability properties of algorithmic thermodynamics is studied. It is proved that there is no computable means to select times where dynamics produce states of continually decreasing entropy. It is also proved that algorithmic fine grained entropy cannot be approximated.

Chapter 11

Computable Measure Spaces

The results in the thermodynamics uses computable metric spaces and computable measure spaces. We use the constructs from [HR09], generalizing from probability measures to arbitrary nonnegative measures and from binary representations to dual binary representations.

Definition 15 *A computable metric space consists of a triple (X, \mathcal{S}, d) , where*

- *X is a separable complete metric space.*
- *\mathcal{S} is an enumerable list of dense ideal points \mathcal{S} in X .*
- *d is a distance metric that is uniformly computable over points in \mathcal{S} .*

For $x \in X$, $r \in \mathbb{Q}_{>0}$ a ball is $B(x, r) = \{y : d(x, y) < r\}$. The ideal points induce a sequence of enumerable ideal balls $B_i = \{B(s_i, r_j) : s_i \in \mathcal{S}, r_j \in \mathbb{Q}_{>0}\}$. We have $\overline{B}(s_i, r_j) = \{y : d(x, y) \leq r\}$, which might not equal the closure of $B(s_i, r_j)$ if there are isolated points. A sequence of ideal points $\{x_n\} \subseteq Y$ is said to be a fast Cauchy sequence if $d(x_n, x_{n+1}) < 2^{-n}$ for all $n \in \mathbb{N}$. A point x is computable there is a computable fast Cauchy sequence converging to x . Each computable function f between computable metric spaces X and \mathcal{Y} has an algorithm \mathfrak{A} such that if $f(x) = y$ then for all fast Cauchy sequences \vec{x} for x , $\mathfrak{A}(\vec{x})$ outputs an encoding of a fast Cauchy sequence for y .

Proposition 5 *For $x \in X$, the following statements are equivalent.*

1. *x is a computable point.*
2. *Each $d(x, s_i)$ are upper semi-computable uniformly in i .*
3. *$d_x = d(x, \cdot) \rightarrow \mathbb{R}^+$ is a computable function.*

11.1 Enumerative Lattices

An enumerative lattice is the tuple (X, \leq, \mathcal{P}) where (X, \leq) is a complete lattice and \mathcal{P} is a numbered set such that if $x \in X$ then $x = \sup P$, for some $P \subseteq \mathcal{P}$. An element $x \in X$ is called lower-computable if there is some enumeration of $\{p : p \leq x\}$.

Proposition 6 *Let (X, \leq, \mathcal{P}) be an enumerative lattice. There is an enumeration of all its lower-computable elements .*

Proof. Using the universal Turing machine, enumerate all enumerable sets. That is, for each enumerable E , there is an i such that $E = \{U(i, n) : n \in \mathbb{N}\}$. Thus for each lower-computable element x has an enumerated set E such that $x = \sup\{p_i : i \in E\}$. \square

Definition 16 Given two enumerative lattices, Y and Z , a function $f : Y \rightarrow Z$ is Scott continuous if it is monotonic and $\sup f(\vec{p}) = f(\sup \vec{p})$ for every increasing sequence \vec{p} . We say f is bi-lower-computable, if there exists a computable method that given an enumerable sequence $\vec{p} = (p_{n_1}, p_{n_2}, \dots)$ with $y = \sup \vec{p}$, outputs a sequence $\vec{q} = \{q_{n_1}, q_{n_2}, \dots\}$ such that $f(y) = \sup \vec{q}$.

Proposition 7 If a function $f : Y \rightarrow Z$ is Scott-continuous and all $f(\sup(p_{n_1}, \dots, p_{n_k}))$ are lower-computable uniformly in (n_1, \dots, n_k) , then f is bi-lower-computable.

Proof. Let $\vec{p} = (p_{n_1}, p_{n_2}, \dots)$ be a sequence such that $y = \sup \vec{p} \in Y$. An algorithm works with access to \vec{p} works as follows. For all k , it lower computes $f(\sup(p_{n_1}, p_{n_2}, \dots))$, which is possible due to the assumption of the Proposition. The supremum of this sequence is $\sup_k f(\sup\{p_{n_1}, \dots, p_{n_k}\})$, which is lower computable due to f being Scott continuous. Thus the enumerated sequence is a lower description of $f(y)$. \square

Definition 17 Given a computable metric space (X, d, \mathcal{L}) and an enumerative lattice (Y, \leq, \mathcal{P}) , we denote \mathcal{F} to be the step functions from X to Y , where

$$f_{(i,j)}(x) = \begin{cases} p_j & \text{if } x \in B_i \\ \perp & \text{otherwise.} \end{cases}$$

We define $\mathcal{C}(X, Y)$ as the closure of \mathcal{F} under pointwise suprema, with pointwise ordering \sqsubseteq . It immediately follows that $(\mathcal{C}(X, Y), \sqsubseteq, \mathcal{F})$ is an enumerative lattice. A function $f : X \rightarrow Y$ is lower-computable if it is a lower-computable element of the enumerative lattice. $(\mathcal{C}(X, Y), \sqsubseteq, \mathcal{F})$.

Example 5 The set $\mathbb{R}^+ = [0, \infty) \cup \{\infty\}$ has an enumerative lattice $(\mathbb{R}^+, \leq, \mathbb{Q}^+)$ which induces a lattice $\mathcal{S}(X, \mathbb{R}^+)$ of positive lower semi-continuous functions from X to \mathbb{R}^+ . Its lower-computable elements are the lower semi-computable functions.

Definition 18 A subset A of X is semi-decidable if it is an r.e. open set.

Proposition 8 Let (X, d_X, S_X) and (Y, d_Y, S_Y) be computable metric spaces. A partial function $f : D \subseteq X \rightarrow Y$ is computable if and only if the preimages of ideal balls are uniformly r.e. open in D sets. So for all i , $f^{-1}(B_i) = U_i \cap D$ where U_i is an r.e. open set uniformly in i . \square

11.2 Computing with Measures

The computable metric space of all Borel probability measures over X is $\mathcal{M}(X)$. If X is separable and compact then so is $\mathcal{M}(X)$. The ideal points of $\mathcal{M}(X)$ are \mathcal{D} , the set of probability measures that are concentrated on finitely many points with rational values. The distance metric on $\mathcal{M}(X)$ is the *Prokhorov metric*, defined as follows.

Definition 19 (Prokhorov metric)

$$\pi(\mu, \nu) = \inf \{ \epsilon \in \mathbb{R}^+ : \mu(A) \leq \nu(A^\epsilon) \text{ for Borel set } A \},$$

where $A^\epsilon = \{x : d(x, A) < \epsilon\}$.

In thermodynamics, the measure function representing the volume is not necessarily a probability measure. Thus the results of [HR09] needs to be extended to nonnegative measures of arbitrary size to prove a result about thermodynamics. Let $(\mathbb{R}^+, d_{\mathbb{R}}, \mathbb{Q}^+)$ be the computable metric space where $\mathbb{R}^+ = [0, \infty)$ is the complete separable metric space and nonnegative rationals \mathbb{Q}^+ consists of the ideal points. The distance function is $d_{\mathbb{R}}(x, y) = |x - y|$, which is obviously computable over $\mathbb{Q}_{\geq 0}$. The space of nonnegative Borel measures over a computable metric space is the space $\mathfrak{M}(X) = \mathcal{M}(X) \times \mathbb{R}_{\geq 0}$, the product space of the space of probability measures of X , $\mathcal{M}(X)$, with the space of nonnegative reals. We identify a measure with a pair $(\mu, m) \in \mathfrak{M}(X)$ where $\mu \in \mathcal{M}(X)$ is a Borel probability measure over X , and $m \in \mathbb{R}^+$ is the size of the measure.

The distance function of \mathfrak{M} is

$$d_{\mathfrak{M}}((\mu, m), (\nu, n)) = \max\{\pi(\mu, \nu), d_{\mathbb{R}}(m, n)\},$$

where π is the Prokhorov metric (see Definition 19). The ideal points of $\mathfrak{M}(X)$ is the set $\mathcal{D}_{\mathfrak{M}}$ of all finite points with nonnegative rational values. This definition is different from the ideal points in $\mathcal{M}(X)$ in that they don't have to sum to 1. The computable measures of $\mathfrak{M}(X)$ are its constructive points, with respect to a fast Cauchy description. From this definition, the results of Theorem ?? apply directly to arbitrary measures $\mu \in \mathfrak{M}(X)$.

Proposition 9 *The tuple $(\mathfrak{M}(X), d_{\mathfrak{M}}, \mathcal{D}_{\mathfrak{M}})$ is a computable metric space.*

Proof. Let (μ_i, v_i) and (μ_j, v_j) be two ideal points of $(\mathfrak{M}(X), d_{\mathfrak{M}}, \mathcal{D}_{\mathfrak{M}})$, where μ_i and μ_j are two probability measures over X , assigning rational measure to a finite number of ideal points. In addition $v_i, v_j \in \mathbb{Q}^+$. If U is a r.e. open subset of X , $\mu_i(U)$ is lower-computable uniformly in i and U . This is because of $(s_{n_1}, q_{m_1}), \dots, (s_{n_k}, q_{m_k})$ are the mass points of μ_i with their weights then $\mu_i(U) = \sum_{s_{n_j} \in U} q_{m_j}$. As all $s_{n_j} \in U$ can be enumerated from a description of U this sum is lower computable. So $\mu_i(B_{i_1} \cup \dots B_{i_k})$ is lower-computable and $\mu_i(\overline{B_{i_1}} \cup \dots \overline{B_{i_k}})$ is upper semi-computable, uniformly in i and (i_1, \dots, i_k) .

We show that $\pi(\mu_i, \mu_j)$ is computable uniformly in (i, j) . Since μ_i is an ideal measure concentrated over S_i , we have $\pi(\mu_i, \mu_j) = \inf\{\epsilon \in \mathbb{Q} : \forall A \subset S_i, \mu_i(A) < \mu_j(A^\epsilon) + \epsilon\}$. Since μ_j is an ideal measure and A^ϵ is a finite union of open ideal balls, $\mu_j(A^\epsilon)$ is lower computable, uniformly in ϵ and j , so $\pi(\mu_i, \mu_j)$ is upper computable, uniformly in (i, j) . The term $\pi(\mu_i, \mu_j)$ is lower computable, uniformly in (i, j) because $\pi(\mu_i, \mu_j) = \sup\{\epsilon \in \mathbb{Q} : \exists A \subset S_i, \mu_i(A) > \mu_j(A^{\bar{\epsilon}})\}$, with $A^{\bar{\epsilon}} = \{x : d(x, A) \leq \epsilon\}$, and using the upper semi-computability of $\mu_j(A^{\bar{\epsilon}})$.

In addition, it easy to see that $d_{\mathbb{R}}(v_i, v_j)$ is computable. Thus the following term is computable.

$$d_{\mathfrak{M}}((\mu_i, v_i), (\mu_j, v_j)) = \max\{\pi(\mu_i, \mu_j), d_{\mathbb{R}}(v_i, v_j)\}.$$

□

Claim 1 *If $(\mu, m) \in \mathfrak{M}(X)$ is a computable measure, then $(\mu, 1) \in \mathfrak{M}(X)$ is computable as well. This follows from taking the fast Cauchy sequence for (μ, m) and normalizing each ideal point in the series.*

For a metric space X , let τ be the set of all open sets of X . The valuation operator $v : \mathfrak{M}(X) \times \tau \rightarrow \mathbb{R}^+$ maps $((\mu, m), U)$ to $m\mu(U)$. More formally, for the first argument, v takes a $\mathfrak{M}(X)$ fast Cauchy sequence to a measure (μ, m) , $m \in \mathbb{R}^+$, and a sequence of ideal balls B_i such that $U = \bigcup_i B_i$ and outputs $\{x : x < m\mu(U)\}$.

Proposition 10 *The valuation operator v is bi-lower computable, in the second argument.*

Proof. Since $v((\mu, m), \cdot)$ is Scott-continuous, due to Proposition 7, the proof is satisfied if we show that $v((\mu, m), \cdot)$ is uniformly lower-computable on finite union of ideal balls. For ideal probability measure $\mu_i \in \mathcal{M}(X)$, due to the proof of Proposition 9, $\mu_i(B_{i_1} \cup \dots \cup B_{i_k})$ is lower computable, uniformly in (i, i_1, \dots, i_k) .

Let $((\mu_{k_n}, m_n))_{n \in \mathbb{N}}$ be a description of a (not necessarily probability) measure $(\mu, m) \in \mathfrak{M}(X)$. Thus $\pi(\mu_{k_n}, \mu) \leq \epsilon_n$ and $|m_n - m| \leq \epsilon_n$, where $\epsilon_n = 2^{-n+1}$. For $n \in \mathbb{N}$ and $U = B(s_{i_1}, q_{j_1}) \cup \dots \cup B(s_{i_k}, q_{j_k})$ we have

$$U_n = \bigcup_{m \leq k} B(s_{i_m}, q_{j_m} - \epsilon_n).$$

We have $U_{n-1}^{\epsilon_n} \subseteq U_n$ and $U_n^{\epsilon_n} \subseteq U$, where $A^\epsilon = \{x : d(x, A) < \epsilon\}$. We will show that $\mu(U) = \sup_n (\mu_{j_n}(U_n) - \epsilon_n)$. Since $\pi(\mu_{j_n}, \mu) \leq \epsilon_n$, and we have that $\mu_{j_n}(U_n) \leq \mu(U) + \epsilon_n$ for all n , so $\mu(U) \geq \sup_n (\mu_{j_n}(U_n) - \epsilon_n)$. Similarly, we have $\mu(U_{n-1}) \leq \mu_{j_n}(U_n) + \epsilon_n$, for all n . So as $n \rightarrow \infty$, $\mu(U) \leq \sup_n (\mu(U_{n-1}) - 2\epsilon_n) \leq \sup_n (\mu_{j_n}(U_n) - \epsilon_n)$. Thus $\mu(U) = \sup_n \mu_{j_n}(U_n) - \epsilon_n$ is lower computable. In addition $m = \sup_n m_n - \epsilon_n$ is lower computable $v((\mu, m), U) = m\mu(U)$ is lower computable, uniformly in (i, i_1, \dots, i_k) . \square

Proposition 11 For measure (μ, m) , if m is computable and measure $\mu(B_{i_1} \cup \dots \cup B_{i_k})$ is uniformly lower computable in (i_1, \dots, i_k) then (μ, m) is computable.

Proof. We show that $\pi(\mu_n, \mu)$ is upper computable uniformly in n and then apply Proposition 5. Since $\pi(\mu, \mu) < \epsilon$ iff $\mu_n(A) < \mu(A^\epsilon) + \epsilon$ for all $A \subset S_n$ where S_n is the finite support of μ_n , and $\mu(A^\epsilon)$ is lower computable (as A^ϵ is a finite union of open ideal balls) $\pi(\mu_n, \mu)$ is semi-decidable, uniformly in n and ϵ . Furthermore, for any ideal point $(\mu_n, m_n) \in \mathfrak{M}(\mathcal{X})$, since m is computable $d_{\mathfrak{M}}((\mu_n, m_n), (\mu, m)) = \max\{\pi(\mu_n, \mu), d_{\mathbb{R}}(m_n, m)\}$ is upper computable so Proposition 5 can be applied. Thus one can construct a fast sequence of ideal measures converging to (μ, m) . \square

For the Cantor space $\{0, 1\}^\infty$ with the standard metric space structure, the ideal balls are the cylinders $x\{0, 1\}^\infty$, for $x \in \{0, 1\}^*$.

Corollary 11 If a measure $(\mu, m) \in \mathfrak{M}(\{0, 1\}^\infty)$ is computable and m is computable, then the cylinders are uniformly computable.

Proposition 12 The integral operator $\int : \mathfrak{M}(X) \times \mathcal{C}(X, \mathbb{R}^+) \rightarrow \mathbb{R}^+$ is bi-lower computable, in the second argument.

Proof. Let $(\mu, m) \in \mathfrak{M}(X)$. The integral of a finite supremum of steps functions can be expressed by induction on the number functions, starting with $m \int f_{(i,j)} d\mu = mq_j \mu(B_i)$ and

$$m \int \sup \{f_{(i_1, j_1)}, \dots, f_{(i_k, j_k)}\} d\mu = mq_{j_z} \mu(B_{i_1} \cup \dots \cup B_{i_k}) + m \int \sup \{f_{(i_1, j'_1)}, \dots, f_{(i_k, j'_k)}\} d\mu$$

where $q_{j_z} = \min\{q_{j_1}, \dots, q_{j_k}\}$ and $q_{j'_i} = q_{j_i} - q_{j_z}$. Since $f_{(i_z, j'_z)}$ is zero, it can be removed. It is easy to see that m can be computed, and by Proposition 10, the measure of finite balls can be lower computed, uniformly in $(B_{i_1}, \dots, B_{i_m})$. For any measure (μ, m) , the operator $m \int d_\mu : \mathcal{C}(X, \mathbb{R}^+) \rightarrow \mathbb{R}^+$ is Scott continuous, so by Proposition 7, the operator is bi-lower computable. \square

Corollary 12 Let $(f_i)_i$ be a sequence of uniformly computable functions, such that the function $(i, x) \mapsto f_i(x)$ is computable. If f_i is bounded by M_i computable uniformly in i , then the function $((\mu, m), i) \mapsto m \int f_i \mu$ is computable.

Proof. $f_i + M$ and $M_i - f_i$ are uniformly lower computable, so $m \int f_i d\mu = m \int (f + M_i) d\mu - m M_i = m M_i - m \int (M_i - f_i) d\mu$ is lower and upper computable by Proposition 12.

11.3 Computable Measure Space

Definition 20

1. A dual measure space $(\mathcal{X}, (\mu, m), (\nu, n))$ is a computable metric space \mathcal{X} and two computable Borel measures, (μ, m) , and (ν, n) over \mathcal{X} . A measure space $(\mathcal{X}, (\mu, m))$ is dual measure space $(\mathcal{X}, (\mu, m), (\mu, m))$.
2. A constructive G_δ -set is a set of the form $\bigcap_n U_n$ where $(U_n)_n$ is a sequence of uniformly r.e. open sets.
3. For computable measure space $(\mathcal{X}, (\mu, m))$ and computable metric space \mathcal{Y} , a function $f : \mathcal{X} \rightarrow \mathcal{Y}$ is almost computable if it is computable on a constructive G_δ set of measure m .
4. A morphism of computable probability spaces $Q : (\mathcal{X}, (\mu, m)) \rightarrow (\mathcal{Y}, (\nu, n))$ is an almost computable measure-preserving function $Q : D_Q \subset \mathcal{X} \rightarrow \mathcal{Y}$, where $\mu(Q^{-1}(A)) = \nu(A)$ for all Borel sets A . An isomorphism (Q, R) is a pair of morphisms such that $Q \circ R = \text{id}$ on $R^{-1}(D_Q)$ and $R \circ Q = \text{id}$ on $Q^{-1}(D_R)$.
5. A dual binary representation of dual computable measure space $(\mathcal{X}, (\mu, m), (\nu, n))$ is a tuple $(\delta, \mu_\delta, \nu_\delta)$ where (μ_δ, m) and (ν_δ, n) are computable (not necessarily probability) measures on $\{0, 1\}^\infty$ and $\delta : (\{0, 1\}^\infty, (\mu_\delta, m)) \rightarrow (\mathcal{X}, (\mu, m))$ and $\delta : (\{0, 1\}^\infty, (\nu_\delta, n)) \rightarrow (\mathcal{X}, (\nu, n))$ are surjective morphisms. Denoting $\delta^{-1}(x)$ to be the set of expansion of $x \in X$:
 - There is a dense full-measure constructive G_δ -set D of points have a unique expansion.
 - $\delta^{-1} : D \rightarrow \delta^{-1}(D)$ is computable.
 - (δ, δ^{-1}) is an isomorphism.
6. A binary representation of computable measure space $(\mathcal{X}, (\mu, m))$ is a dual representation of the dual computable measure space $(\mathcal{X}, (\mu, m), (\mu, m))$.
7. A set A is almost decidable with respect to measures (μ, m) and (ν, n) if there are two. r.e. open sets U and V such that $U \subset A$, $V \subseteq A^c$, $U \cup V$ is dense and has full μ and ν measure. We say the elements of a sequence $\{A_i\}$ are uniformly almost decidable with respect to (μ, m) and (ν, n) if there are two sequences $\{U_i\}$ and $\{V_i\}$ of uniformly r.e. sets satisfying the above conditions.

The follow proof of existence of an almost decidable set is from [GHR11].

Lemma 11 Let X be \mathbb{R} or \mathbb{R}^+ or $[0, 1]$. Let (μ, m) and (ν, n) be a computable measures on X . Then there is a sequence of uniformly computable reals $(x_n)_n$ which is dense in X and such that $\mu(\{x_n\}) = \nu(\{x_n\}) = 0$ for all n .

Proof. Let I be a closed rational interval. Let $M = \max\{m, n\}$. We construct $x \in I$ such that $\mu(\{x\}) = \nu(\{x\}) = 0$. To do this, we construct inductively a nested sequence of closed intervals J_k of μ and ν measure $< M2^{-k+1}$, with $J_0 = I$. Suppose $J_k = [a, b]$ has been constructed, with $\mu(J_k) < M2^{-k+1}$ and $\nu(J_k) < 2^{-k+1}$. Let $m = (b - a)/16$ and $\ell = (b - a)/64$: by the Markov inequality one of the intervals $[a + jm + \ell, a + (j + 1)m - \ell]$ $j \in \{0, \dots, 15\}$ must have μ and ν measure $< M2^{-k}$ and since these measures are upper computable, it can be found effectively, and we denote it J_{k+1} . By enumerating all dyadic intervals $(I_n)_n$, one can construct $x_n \in I_n$ uniformly. \square

Corollary 13 *Let $(\mathcal{X}, (\mu, m), (\nu, m))$ be a dual measure space and $(f_i)_i$ be a sequence of uniformly computable real valued functions on X . There is a sequence of uniformly computable reals $(x_n)_n$ which is dense in \mathbb{R} such that $\mu(\{f_i^{-1}(x_n)\}) = \nu(\{f_i^{-1}(x_n)\}) = 0$ over all i, n .*

Proof. We define the uniformly computable measure (μ_i, m) where $\mu_i = \mu \circ f_i^{-1}$ and (ν_i, n) where $\nu_i = \nu \circ f_i^{-1}$. Define measure (λ, m) , $\lambda = \sum 2^{-i} \mu_i$ and (γ, n) , $\gamma = \sum 2^{-i} \nu_i$. By Proposition 11, (λ, m) and (γ, m) are computable measures so by Lemma 11 there is a sequence of uniformly computable reals $(x_n)_n$ which is dense in \mathbb{R} such that $\lambda(\{x_n\}) = \gamma(\{x_n\}) = 0$ for all i, n . \square

Corollary 14 *There is a sequence of uniformly computable reals $(r_n)_{n \in \mathbb{N}}$ such that $(B(s_i, r_i))$ is a basis of almost decidable balls.*

Proof. Apply Corollary 13 to $(f_i)_i$ defined by $f_i(x) = d(s_i, x)$. \square

Every ideal ball can be expressed as a r.e. union of almost decidable balls, and vice-versa. So the two bases are constructively equivalent.

Definition 21 *A set D is an ad-set if it is a finite union of almost decidable balls, with $D = B_{i_1} \cup \dots \cup B_{i_k}$. We have $\overline{\overline{D}} = \overline{\overline{B_{i_1}}} \cup \dots \cup \overline{\overline{B_{i_k}}}$, which may differ than the closure of D if there are isolated points.*

Proposition 13 *For computable measure space (\mathcal{X}, μ) and ad-set D , $\mu(D)$ is computable.*

Proof. This follows from Proposition 11, which implies $\mu(D)$ and $\mu(X \setminus \overline{\overline{D}})$ being lower computable, noting the fact that all almost decidable balls have borders of null measure.

We fix computable measures (μ, m) and (ν, n) , and their computable representations. We denote $B(s_i, r_n)$ by B_k where $k = \langle i, n \rangle$ and r_n is the sequence defined in 14. Let $C_k = X \setminus \overline{\overline{B}}(s_i, r_n)$. For $w \in \{0, 1\}^*$, the cell $\Gamma(w)$ is defined by $\Gamma(\epsilon) = X$, $\Gamma(w0) = \Gamma(w) \cap C_i$ and $\Gamma(w1) = \Gamma(w) \cap B_i$, where ϵ is the empty word and $i = \|w\|$. This is an almost decidable set, uniformly in w .

Theorem 48 *Every dual measure space $(\mathcal{X}, (\mu, m), (\nu, n))$ has a dual binary representation.*

Proof. We construct an encoding function $b : D \rightarrow \{0, 1\}^\infty$, a decoding function $\delta : D_\delta \rightarrow X$, and show that δ is a multi binary representation, with $b = \delta^{-1}$.

Let $D = \cap_i B_i \cup C_i$. The set D is a full-measure constructive G_δ -set. Define the computable function $b : D \rightarrow \{0, 1\}^\infty$ with

$$b(x)_i = \begin{cases} 1 & \text{if } x \in B_i \\ 0 & \text{if } x \in C_i. \end{cases}$$

Let $x \in D$: $\omega = b(x)$ is also characterized by $\{x\} = \cap_i \Gamma(\omega_{0\dots i-1})$. b can be computed from $\Gamma(\cdot)$. Let (μ_δ, m) and (ν_δ, n) computable measures over $\{0, 1\}^\infty$, where $\mu_\delta = \mu \circ b^{-1}$, and $\nu_\delta = \nu \circ b^{-1}$. Let D_δ be the set of binary sequences ω such that $\cap_i \Gamma(\omega_{0\dots i-1})$ is a singleton. The decoding function $\delta : D_\delta \rightarrow X$ is defined by

$$\delta(\omega) = x \text{ if } \cap_i \overline{\Gamma(\omega_{0\dots i-1})} = \{x\}.$$

The next steps are to prove that δ is a surjective morphism. The center and radius of the ball B_i will be s_i and r_i , respectively. We say n is an i -witness for ω if $r_i < 2^{-(n+1)}$, $\omega[i] = 1$, and $\Gamma(\omega[0..i]) \neq \emptyset$. We first prove that

$$D_\delta = \cap_n \{\omega \in \{0, 1\}^\infty : \omega \text{ has a } n\text{-witness}\}.$$

Let $\omega \in D_\delta$ and $x = \delta(\omega)$. For every n , $x \in D(s_i, r_i)$ for some i with $r_i \leq 2^{-(n+1)}$. Since $x \in \overline{\Gamma(\omega[0\dots i])}$, we have $\Gamma(\omega[0\dots i]) \neq \emptyset$ and $\omega[i] = 1$. So i is an n -witness for ω . Conversely if ω has a n -witness i_n for all n , since $\overline{\Gamma[0\dots i_n]} \subseteq \overline{B_{i_n}}$ with radius going to zero, the sequence $\overline{\Gamma(\omega[0\dots n])}$ of closed balls has a non-empty intersection, due to the completeness of the space, and it is a singleton.

$\delta : D_\delta \rightarrow X$ is computable. For each n , find an n -witness i_n of ω : the sequence $(s_{i_n})_n$ is a fast Cauchy sequence converge to $\delta(\omega)$. In addition, δ is surjective: each $x \in X$ has at least one expansion. We construct by induction a sequence $\omega = \omega[0]\omega[1]\dots$ such that for all i , $x \in \overline{\Gamma(\omega[0\dots i])}$. Let $i \geq 0$ and suppose that $\omega[0\dots i-1]$ has been constructed. Since $B_i \cup C_i$ is open and dense and $\Gamma(\omega_{0\dots i-1})$ is open, $\overline{\Gamma(\omega_{0\dots i-1})} = \overline{\Gamma(\omega_{0\dots i-1}) \cap (B_i \cup C_i)} = \overline{\Gamma\omega_{0\dots i-1}0} \cup \overline{\Gamma\omega_{0\dots i-1}1}$, so for some $\omega[i] \in \{0, 1\}$, has $x \in \overline{\Gamma(\omega_{0\dots i})}$. So $x \in \cap_i \overline{\Gamma(\omega_{0\dots i-1})}$. Since $(B_i)_i$ is a basis and $\omega_i = 1$ whenever $x \in B_i$, ω is an expansion of x .

11.4 Randomness

Definition 22 For a measure $(\mu, m) \in \mathfrak{M}(X)$, a (μ, m) ML randomness test is a sequence of uniformly r.e. open sets $(U_n)_n$, satisfying $m\mu(U_n) \leq 2^{-n}$. The set $\cap_n U_n$ is a null measure set and is called a μ -effective null set. An alternative definition of null sets uses integrals (see [G21]), with a slight modification as measures are being used. Given a measure $(\mu, m) \in \mathfrak{M}(X)$ a μ -randomness test is a (μ, m) computable element of $\mathcal{C}(X, \overline{\mathbb{R}}^+)$ such that $m \int t d\mu \leq 1$. Any subset of $\{x \in X : t(x) = \infty\}$ is called a μ -effective null set. The two definitions of null sets are equivalent. A point $x \in X$ is (μ, m) -ML random if it is in no effective null set. A uniform randomness test is a computable function T from $\mathfrak{M}(X)$ to $\mathcal{C}(X, \overline{\mathbb{R}}^+)$ such that $m \int T^{(\mu, m)} d\mu \leq 1$.

Using proposition 6, let $(H_i)_{i \in \mathbb{N}}$ be an enumeration of all lower computable elements of the enumerative lattice $\mathcal{C}(\mathfrak{M}(X), \mathcal{C}(X, \overline{\mathbb{R}}^+))$, such as $H_i \sup_k f_\phi$ where $\phi : \mathbb{N}^2 \rightarrow \mathbb{N}$ is some recursive function and f_n are step functions.

Lemma 12 There is a computable function $T : \mathbb{N} \times \mathfrak{M}(X) \rightarrow \mathcal{C}(X, \overline{\mathbb{R}}^+)$ with

- For all i , $T_i = T(i, \cdot)$ is a uniform randomness test.
- If $\int m H_i((\mu, m)) d\mu < 1$ for some (μ, m) , then $T_i(\mu) = H_i(\mu)$.

Proof. To enumerate only tests, we'd like to be able to semi-decide $m \int \sup_{k < n} f_{\phi(i, k)}((\mu, m)) d\mu < 1$. But $m \sup_{k < n} f_{\phi(i, k)}((\mu, m))$ is only lower computable (relative to (μ, m)). Let \mathcal{Y} be a computable metric space. For an ideal point $s \in Y$ and positive rationals q, r, ϵ , define the hat function:

$$h_{q, s, r, \epsilon}(y) = q[1 - [d(y, s) - r]^+ / \epsilon]^+,$$

where $[a]^+ = \max\{0, a\}$. This is a continuous function whose value is q in $B(s, r)$ and 0 outside $B(s, r + \epsilon)$. It is easy to see there is a number $(h_n)_{n \in \mathbb{N}}$ of all the hat functions. They are equivalent to step function in the enumerative lattice $\mathcal{C}(Y, \mathbb{R}^+)$. The step functions can be constructed as the supremum of such function $f_{(i,j)} = \sup\{h_{q_j, s, r-\epsilon, \epsilon: 0 < \epsilon < r}\}$ with $B_i = B(s, r)$ and conversely.

We let $Y = \mathfrak{M}(X) \times X$ endowed with the canonical computable metric structure. By “curryfication” it provides functions $h_n \in \mathcal{C}(\in \mathfrak{M}(X), \mathcal{C}(X, \overline{\mathbb{R}}^+))$ with which the H_i can be expressed: there is a recursive function $\phi : \mathbb{N}^2 \rightarrow \mathbb{N}$ such that for all i , $H_i = \sup_k h_{\phi(i,k)}$.

In addition, $h_n((\mu, m))$ is bound by a constant computable from n and independent of (μ, m) . Hence, by Corollary 12, the integration operator $\int : \mathfrak{M}(X) \times \mathbb{N} \rightarrow [0, 1]$ which maps $((\mu, m) \langle i_1, \dots, i_k \rangle)$ to $m \int \sup\{h_{i_1}((\mu, m)), \dots, h_{i_k}((\mu, m))\} d\mu$ is computable. Thus $T(i, (\mu, m)) = \sup\{H_i^k((\mu, m)) : m \int H_i^k((\mu, m)) \leq 1\}$ where $H_i^k = \sup_{n < k} h_{\phi(i,n)}$. Since $m \int H_i^k((\mu, m)) d\mu$ can be computed from i, k , and a description of (μ, m) , T is a computable function from $\mathbb{N} \times \mathfrak{M}(X)$ to $\mathcal{C}(X, \overline{\mathbb{R}}^+)$. \square

Theorem 49 *There is a universal uniform randomness test, that is a uniform test \mathbf{t} such that for every uniform test T , there is a constant $c > 0$ with $\mathbf{t} > \mathbf{m}(T)T$.*

Proof. Using Lemma 12, the universal test is defined by $\mathbf{t} = \sum_i \mathbf{m}(i)T_i$: since every T_i is a uniform randomness test, \mathbf{t} is also a uniform randomness test. In addition, for every uniform test T , there is an i such that $T = T_i = H_i$. \square

The following corollary is due to [G21] with the proofs adapted to uniform tests. Let $F : \mathcal{Y} \times \mathfrak{M}(X) \rightarrow \mathbb{Z} \cup \{-\infty, \infty\}$ be lower computable, where \mathcal{Y} is a computable metric space. An F uniform randomness test R is a computable function from $\mathcal{Y} \times \mathfrak{M}(X)$ to $\mathcal{C}(X, \overline{\mathbb{R}}^+)$ such that $m \int R^{(y, (\mu, m))} d\mu \leq 2^{-F(y, (\mu, m))}$.

Corollary 15 *There exists a universal F uniform test \mathbf{r} such that $m \int \mathbf{r}^{(y, (\mu, m))} d\mu \leq 2^{-F(y, (\mu, m))}$ and for every F uniform test R , $\mathbf{m}(R | \langle \overrightarrow{y} \rangle) R_y \leq^* \mathbf{r}_y$.*

Proof. The proof follows analogously to that of Lemma 12, except $T(i, y, (\mu, m)) = \sup\{H_i^k(y, (\mu, m)) : m \int H_i^k(y, (\mu, m)) \leq 2^{-F(y, (\mu, m))}\}$ where $H_i^k = \sup_{n < k} h_{\phi(i, n, \langle \overrightarrow{y} \rangle)}$. The term $\phi(i, n, \langle \overrightarrow{y} \rangle)$ is the partial recursive function being given the numbers i and n , and an encoding of a fast Cauchy sequence for $y \in \mathcal{Y}$.

Corollary 16 *Let $\overline{\mathbb{Z}} = \mathbb{Z} \cup \{-\infty, \infty\}$. For computable metric spaces \mathcal{X} and \mathcal{Y} , consider computable functions $\mathcal{Y} \rightarrow \mathcal{C}(X, \overline{\mathbb{R}}^+)$. There is a modification to a computable $w' : \mathfrak{M}(X) \times \mathcal{Y} \times \overline{\mathbb{Z}} \rightarrow \mathcal{C}(X, \overline{\mathbb{R}}^+)$ such that for all $y \in \mathcal{Y}$, $(\nu, m) \in \mathfrak{M}(X)$, $m \in \overline{\mathbb{Z}}$, $n \int w'((\nu, \nu), y, m, x) d\nu \leq 2^{-m+1}$ and if $n \int w((\nu, \nu), y, x) d\nu \leq 2^{-m+1}$.*

Proof. For every w , there is an i such that $w((\nu, n), x, y) = \sup_k H_k((\nu, n), x, y)$, where $H_k = \sup_{j < k} h_{\phi(i, j, \langle \overrightarrow{y} \rangle)}$, where h is defined in the proof of Lemma 12. The transformed function is

$$w'((\nu, n), y, m, x) = \sup_k \{H_k((\nu, n), x, y) : n \int H_k((\nu, n), x, y) d\nu \leq 2^{-m}\}.$$

\square

Definition 23 (Randomness Deficiency) We recall that the deficiency of randomness of an infinite sequence $\alpha \in \{0,1\}^\infty$ with respect to a computable measure (P, p) over $\{0,1\}^\infty$ is defined to be

$$\mathbf{D}(\alpha|(P, p), x) = \log \sup_n \mathbf{m}(\alpha[0..n]|x)/p \cdot P(\alpha[0..n]).$$

We have $\mathbf{D}(\alpha|(P, p)) = \mathbf{D}(\alpha|(P, p), \emptyset)$. By [G21], $2^{\mathbf{D}}$ is a lower-computable (P, p) -test, in that

$$p \int_{\{0,1\}^\infty} 2^{\mathbf{D}(\alpha|(P, p))} dP(\alpha) = O(1).$$

Proposition 14 Let $(\{0,1\}^\infty, (\mu, m))$ be a computable measure space. Then $\mathbf{t}_{(\mu, m)}(\alpha) \stackrel{*}{=} 2^{\mathbf{D}(\alpha|(\mu, m))}$.

Proof. There is a constructive element H of $\mathcal{C}(\in \mathfrak{M}(X), \mathcal{C}(X, \overline{R}^+))$, such that $H((\nu, n), \alpha) = 2^{\mathbf{D}(\alpha|(\mu, m))}$. Thus there is some i , such that $H_i = H$. Furthermore T_i is a uniform test satisfying $T_i((\mu, m)) = 2^{\mathbf{D}(\alpha|(\mu, m))}$ because $m \int T_i(\mu, m) d\mu = m \int H((\mu, m)) d\mu < 1$.

As described in [G21], $2^{\mathbf{D}(\alpha|(\mu, m))}$ is a universal lower computable test. Thus $\mathbf{t}_{(\mu, m)}(\alpha) \stackrel{*}{<} 2^{\mathbf{D}(\alpha|(\mu, m))}$.

Proposition 15 For computable measure space $(X, (\mu, m))$, every random point lies in every r.e. open set of full measure.

Proof. Let $U = \bigcup_{(i,j) \in E} B(s_i, q_j)$ be a r.e. open set of measure m , with $E \subseteq \mathbb{N}$ being r.e. Let F be the r.e. set $\{(i, k) : \exists j, (i, j) \in E, q_k < q_j\}$. Let

$$U_n = \bigcup_{(i,k) \cap [0,n]} B(s_i, q_k) \text{ and } V_n^\mathcal{C} = \bigcup_{(i,k) \cap [0,n]} \overline{\overline{B}}(s_i, q_k).$$

Then U_n and V_n are r.e. uniformly in n , $U_n \nearrow U$ and $U^\mathcal{C} = \bigcap_n V_n$. As $\mu(U_n)$ is lower semi-computable uniformly in n , a sequence $(n_i)_{i \in \mathbb{N}}$ can be computed such that $m\mu(U_{n_i}) > 1 - 2^{-i}$. Thus $m\mu(V_{n_i}) < 2^{-i}$ and $U^\mathcal{C} = \bigcap_i V_{n_i}$ is a μ -ML test. Thus every (μ, m) -random point is in U . \square

Lemma 13 Let $Q : D \subset X \rightarrow \mathcal{Y}$ be a morphism of equal computable measure spaces $(X, (\mu, m))$ and $(\mathcal{Y}, (\nu, m))$, with universal tests $\mathbf{t}_{(\mu, m)}$ and $\mathbf{t}_{(\nu, m)}$. Then there is some c with the following properties. If $x \in X$ and $\mathbf{t}_{(\mu, m)}(x) < \infty$, then $Q(x)$ is defined and $\mathbf{t}_{(\nu, m)}(Q(x)) \stackrel{*}{<} c\mathbf{t}_{(\mu, m)}(x)$.

Proof. Assuming $\mathbf{t}_{(\mu, m)}(x) < \infty$, then x is a random point then $x \in D$, because due to Proposition 15, every random point lies in every r.e. open set of full measure, and D is an intersection of full-measure r.e. open sets. Thus $Q(x)$ is defined.

We have that $\mathbf{t}_{(\nu, m)} \circ Q \in \mathcal{C}(X, \overline{R}^+)$ because there is an algorithm that enumerates all finite prefixes of fast Cauchy sequences to Q and enumerates all resultant outputted ideal balls. Then the algorithm sees which outputted ideal balls B are in the values ideal balls (B', v) enumerated by $\mathbf{t}_{(\nu, m)}$. If $B \subseteq B'$, then the algorithm outputs (B, v) .

Since $\mu(D) = 1$, $\int \mathbf{t}_{(\nu, m)} \circ Q d\mu$ is well defined. As Q is measure-preserving, $m \int \mathbf{t}_{(\nu, m)} \circ Q du = m \int \mathbf{t}_{(\nu, m)} d\nu \leq 1$. Hence $\mathbf{t}_{(\nu, m)} \circ Q$ is a μ -test, so there exists $c \in \mathbb{N}$ with $\mathbf{t}_{(\nu, m)} \circ Q \stackrel{*}{<} c\mathbf{t}_{(\mu, m)}$. \square

Corollary 17 Let $(Q, R) : (\mathcal{X}, \mu) \rightleftharpoons (\mathcal{Y}, \nu)$ be an isomorphism of computable measure spaces, with universal tests \mathbf{t}_μ and \mathbf{t}_ν . Then there is a $c \in \mathbb{N}$ where $\mathbf{t}_\nu(Q(x)) = \mathbf{t}_\mu(x) \pm c$ and $\mathbf{t}_\mu(R(y)) = \mathbf{t}_\nu(y) \pm c$.

Chapter 12

Algorithmic Fine Grained Entropy

In this chapter we introduce the central term of algorithmic thermodynamics: algorithmic fine grain entropy \mathbf{H}_μ . We also show some canonical properties of \mathbf{H}_μ , originating from [G21], with modifications to the proofs as needed to be compatible with Chapter 11. In this chapter and in subsequent ones, we represent (not necessarily probabilistic) measures as μ , dropping the (μ, m) notation.

Definition 24 *Given a measure space (\mathcal{X}, μ) , its corresponding algorithmic fine grained entropy is $\mathbf{H}_\mu(x) = -\log \mathbf{t}_\mu(x)$.*

The term \mathbf{H}_μ is bounded from above by $\log \mu(X)$ and can take arbitrary negative values, including infinitely negative values. If x is in a μ -nullset then $\mathbf{H}_\mu(x) = -\infty$.

Proposition 16 *If μ is a probability measure, then $\mathbf{H}_\nu(y) >^+ \mathbf{H}_{\mu,\nu}(x, y)$.*

Proof. $2^{-\mathbf{H}_\nu(y)}$ is a test for $\mu \times \nu$, since $\mu^x \nu^y 2^{-\mathbf{H}_\nu(y)} \leq \mu^x 1 = 1$. □

Definition 25 (Computable Transformation Group) *A one dimensional transformation group G^t , parameterized by $t \in \mathbb{R}$ over a measure space (\mathcal{X}, μ) where each G^t is a homeomorphism of \mathcal{X} onto itself, where $G^t(G^s(x)) = G^{t+s}(x)$. And $G^t x$ is continuously simultaneously in x and t . G is measure preserving, where $\mu(G^t(A)) = \mu(A)$, for all Borel sets A . Furthermore there is a program that when given an encoding of a fast Cauchy sequence of $t \in \mathbb{R}$ and $x \in \mathcal{X}$, outputs an encoding of a fast Cauchy sequence of $G^t x$.*

Example 6 (Particles in a Box) *We detail a canonical example: particles in a box, as depicted in Figure 12.1. Let there be a box containing N particles. Each particle has a position $p \in [0, M] \times [0, M] \times [0, M]$ and momentum $m \in [-M/2, M/2] \times [-M/2, M/2] \times [-M/2, M/2]$ and no other degrees of freedom. The state space X is the position and volume of all particles, and using the Lebesgue measure μ , the total volume is $\mu(X) = M^{6N}$. A state of this space can be represented as $\omega \in X$, with*

$$(\omega.q_1, \dots, \omega.q_{3N}, \omega.p_1, \dots, \omega.p_{3N}).$$

When dynamics are applied to such a point, the particles will move from their positions according to their momentum. If ω contains a computable coordinate, either an x, y, z value of momentum or position, then $\mathbf{H}_\mu(\omega) = -\infty$. Indeed suppose it is a position coordinate k , where $\omega.p_k = \epsilon$ for some computable value $\epsilon \in [0, M]$ (it works similarly for a momentum coordinate). We define the test $t(\alpha) = \sum_n [|\alpha.p_n - \epsilon| < 2^{-n-1}] \mathbf{m}(n) 2^n / M^{6N}$. Thus since ϵ is computable, so is t , and since $t(\omega) = \infty$, $\mathbf{H}_\mu(\omega) = -\infty$.

Proposition 17 *For rational $t \in \mathbb{Q}$, $\mathbf{H}_\mu(G^t x) - \mathbf{H}_\mu(x) <^+ \mathbf{K}(t)$.*



Figure 12.1: A canonical example for algorithmic fine grained entropy. The space consists of the momentum and position of N particles.

Proof. This is because, since G is measure preserving, $\mathbf{t}_\mu(G^{-t}\omega)$ is a μ -test of complexity $\mathbf{K}(t)$. Thus $\mathbf{m}(t)\mathbf{t}_\mu(G^{-t}\omega) \stackrel{*}{<} \mathbf{t}_\mu(\omega)$.

12.1 Thermodynamic Information

Information between a point of the metric space and a binary sequence is introduced as well as the information between two points in metric spaces. The term $\mathbf{H}_\mu(\alpha|t)$ is the fine grained algorithmic entropy of α when the universal Turing machine is relativized to the sequence t .

Definition 26 (Information) Let (\mathcal{X}, μ) and (\mathcal{Y}, ν) be computable measure spaces. For $\alpha \in \mathcal{X}$, $\beta \in \mathcal{Y}$ and $t \in \{0, 1\}^* \cup \{0, 1\}^\infty$,

- $\mathbf{I}(\alpha; t) = \mathbf{H}_\mu(\alpha) - \mathbf{H}_\mu(\alpha|t)$.
- $\mathbf{I}(\alpha : \beta) = \mathbf{H}_\mu(\alpha) + \mathbf{H}_\nu(\beta) - \mathbf{H}_{\mu \times \nu}(\alpha, \beta)$.

If dynamics are used to increase or decrease algorithmic thermodynamic entropy by a non trivial amount, then the encoded dynamics shares algorithmic information with the ending or starting state, respectively. Proposition 18 is due to [Gac94], with usage of algorithmic fine grained entropy. Put another way,

if you want to increase the entropy of a state, you need information about its ending state and if you want to decrease the entropy of a state, you need information about its starting state.

Proposition 18 $-\mathbf{I}(\alpha; t) <^+ \mathbf{H}_\mu(G^t\alpha) - \mathbf{H}_\mu(\alpha) <^+ \mathbf{I}(G^t\alpha; t)$.

Proof. By definition

$$\begin{aligned} \mathbf{I}(\alpha; t) &= \mathbf{H}_\mu(\alpha) - \mathbf{H}_\mu(\alpha|t) \\ \mathbf{I}(G^t\alpha; t) &= \mathbf{H}_\mu(G^t\alpha) - \mathbf{H}_\mu(G^t\alpha|t). \end{aligned}$$

Since the function $2^{-\mathbf{H}_\mu(G^t\alpha)}$ is a μ test,

$$\mathbf{H}_\mu(G^t\alpha) >^+ \mathbf{H}_\mu(\alpha|t) = \mathbf{H}_\mu(\alpha) - \mathbf{I}(\alpha; t)..$$

Which gives us the first inequality. In addition $2^{-\mathbf{H}_\mu(G^t\alpha)}$ is also a μ test, so

$$\mathbf{H}_\mu(G\alpha) >^+ \mathbf{H}_\mu(G^t\alpha|t) = \mathbf{H}_\mu(G^t\alpha) - \mathbf{I}(G^t\alpha; t),$$

which gives the second inequality. \square

Proposition 19 (Conservation of Information) $\mathbf{I}(G^t\alpha : \beta) <^+ \mathbf{I}(\alpha : \beta)$.

Proof. We have that G^t and $G^t \times \text{Id}$ are μ and $\mu \times \nu$ preserving so $2^{-\mathbf{H}_\mu(G^t\alpha)}$ is a μ test and $2^{-\mathbf{H}_\mu((G^t \times \text{Id})(\alpha, \beta))}$ is a $\mu \times \nu$ test. So $\mathbf{H}_\mu(\alpha) >^+ \mathbf{H}(G^t\alpha)$ and $\mathbf{H}_{\mu, \nu}(\alpha, \beta) <^+ \mathbf{H}(\mu, \nu)(G^t\alpha, \beta)$. So

$$\begin{aligned} \mathbf{I}(G^t\alpha : \beta) &= \mathbf{H}_\mu(G^t\alpha : \beta) + \mathbf{H}_n(\beta) - \mathbf{H}_{\mu, \nu}(G^t\alpha, \beta) \\ &<^+ \mathbf{H}_\mu(\alpha : \beta) + \mathbf{H}_n(\beta) - \mathbf{H}_{\mu, \nu}(\alpha, \beta). \end{aligned}$$

\square

12.2 Entropy Balance

The following section is due to [Gac94]. Lets say there exists two independent systems (\mathcal{X}, μ) and (\mathcal{Y}, ν) represented as computable measure spaces that are put under joint dynamics G . We show that under mild assumptions, an increase of entropy in one subsystem implies a decrease in entropy in another system. Let $(\alpha_t, \beta_t) = G^t(\alpha, \beta)$, and $\Delta\mathbf{H}_\mu(\alpha) = \mathbf{H}_\mu(\alpha_t) - \mathbf{H}_\mu(\alpha)$, and similarly for $\Delta\mathbf{H}_\nu(\beta)$.

Lemma 14 $\Delta\mathbf{H}_\mu(\alpha) + \Delta\mathbf{H}_\nu(\beta) >^+ \mathbf{I}(\alpha_t : \beta_t) - \mathbf{I}(\alpha : \beta) - \mathbf{I}((\alpha, \beta); t)$.

Proof. Due to Proposition 18 applied to (α, β) , $\Delta\mathbf{H}_{\mu \times \nu}(\alpha, \beta) >^+ -\mathbf{I}((\alpha, \beta); t)$. So

$$\begin{aligned} \mathbf{H}_\mu(\alpha_t) + \mathbf{H}_\nu(\beta_t) &= \mathbf{H}_{\mu \times \nu}(\alpha_t, \beta_t) + \mathbf{I}((\alpha_t, \beta_t); t) \\ &>^+ \mathbf{H}_{\mu \times \nu}(\alpha, \beta) - \mathbf{I}((\alpha, \beta); t) + \mathbf{I}(\alpha_t : \beta_t) \\ &=^+ \mathbf{H}_\mu(\alpha) + \mathbf{H}_\nu(\beta) + \mathbf{I}(\alpha_t : \beta_t) - \mathbf{I}(\alpha : \beta) - \mathbf{I}((\alpha : \beta); t). \end{aligned}$$

\square

The last term is almost always negligible. If one wants to lower the thermodynamic entropy of a state, the information of the state must be encoded into the dynamics or an independent environment can be coupled with the system which will absorb the entropy.

12.3 Maxwell's Demon

We revisit Maxwell's demon, providing yet another interpretation. This is done by reworking Lemma 14 to the specific case of binary sequences. For the recording space, we use the set $\{0, 1\}^\infty$ of infinite sequences with any computable probability measure λ over $\{0, 1\}^\infty$. Thus by Proposition 14, $\mathbf{H}_\lambda(\alpha) =^+ -\mathbf{D}(\alpha|\lambda)$, where \mathbf{D} is the deficiency of randomness. We couple the computable measure space $(\{0, 1\}^\infty, \lambda)$ with a typical system (\mathcal{X}, μ) , such as where the phase space is the

momentum and position of N particles, for large N . We couple a starting state $\alpha \in \{0,1\}^\infty$, with recording state $\beta \in \{0,1\}^\infty$ that has room to record information, for example, where λ is the uniform measure and $\beta = 0^{1000}\kappa$, for some ML random string κ . The states are independent, with $\mathbf{I}(\alpha : \beta) \approx 0$. The marginal states are assumed independent, with $\mathbf{I}(\alpha : \beta) \approx 0$. Joint dynamics are applied to get $(\alpha^t, \beta^t) = G(\alpha, \beta)$. By Lemma 14,

$$\mathbf{H}_\mu(\alpha^t) - \mathbf{H}_\mu(\alpha) >^+ \mathbf{D}(\beta|\lambda) - \mathbf{D}(\beta^t|\lambda) - \mathbf{I}((\alpha, \beta); t).$$

Again, for most times, $\mathbf{I}((\alpha, \beta); t)$ will be negligible. Thus after α decreases in algorithmic fine grain thermodynamic entropy, the contents of the register fills up, with a decrease in its deficiency of randomness \mathbf{D} . This shows that one benefit of an algorithmic formulation of thermodynamics is that pure algorithmic information and thermodynamic entropy can be exchanged in the course of joint dynamics. A graphical depiction of this phenomenon can be seen in Figure 12.2.

12.4 Distribution of Algorithmic Fine Grained Entropy

We say that measure ν is absolutely continuous with respect to μ , $\nu \ll \mu$ if $\mu(A) = 0$ implies $\nu(A) = 0$ for all $A \subseteq X$. The Radon–Nikodym theorem states that if $\nu \ll \mu$ there exists a measurable (over the Borel sets of \mathcal{X}) function f , uniquely defined up to a μ -nullset, such that for any measurable set $A \subseteq X$,

$$\nu(A) = \int_A f d\mu.$$

The function f can be written as $\frac{d\mu}{d\nu}$ or $\frac{\mu(dx)}{\nu(dx)}$. If $\mu \ll \nu$, then $\frac{\nu(dx)}{\mu(dx)} = \left(\frac{\mu(dx)}{\nu(dx)}\right)^{-1}$. If $\nu \ll \mu \ll$ then $\frac{d\nu}{d\lambda} = \frac{d\nu}{d\mu} \frac{d\mu}{d\lambda}$. We use the short hand $\mu^x f(x) = \int f d\mu$. We define

$$\mathcal{H}_\nu(\mu) = - \int \log \left(\frac{d\mu}{d\nu} \right) d\mu = -\mu^x \log \frac{\mu(dx)}{\nu(dx)} = -\nu^x f(x) \log f(x).$$

If both ν and μ are probability measures, then $-\mathcal{H}_\nu(\mu) = \mathcal{D}(\mu\|\nu)$, where \mathcal{D} is the Kullback–Leibler divergence. The following proposition shows that $\mathcal{H}_\nu(\mu)$ is non-positive when ν and μ are probability measures.

Proposition 20 *Over a space X ,*

$$\mathcal{H}_\nu(\mu) \leq -\mu(X) \log \frac{\mu(X)}{\nu(X)}.$$

Proof. We use the inequality $-a \ln a \leq -a \ln b + b - a$. Letting $a = f(x)$ and $b = \mu(X)/\nu(X)$ and integrating by ν gives us:

$$\begin{aligned} (\ln 2) \mathcal{H}_\nu(\mu) &= -\nu^x f(x) \ln f(x) \leq -\mu(X) \ln \frac{\nu(X)}{\mu(X)} + \frac{\mu(X)}{\nu(X)} \nu(X) - \mu(X) \\ &= -\mu(X) \ln \frac{\mu(X)}{\nu(X)}. \end{aligned}$$

□

Theorem 50 *For computable metric space \mathcal{X} , let μ be measure that that $\mu(X) \geq 1$. Then*

$$\mathcal{H}_\nu(\mu) \leq \mu^x \mathbf{H}_\nu(x).$$



Figure 12.2: A graphical depiction of an algorithmic interpretation of Maxwell's demon. Particles in a box is paired with a register of information, represented as an infinite sequence with a high deficiency of randomness. After dynamics occur, the subsystem of the particle box moves to a state of low entropy, in this case with particles in a smaller region. Due to entropy balance Lemma 14, the register will fill up, moving to a sequence with low deficiency of randomness.

Proof. Let δ be the measure with density $\mathbf{t}_\nu(x)$ with respect ν , with $\mathbf{t}_\nu(x) = \frac{\delta(dx)}{\nu(dx)}$. Since $\nu^x \mathbf{t}_\nu(x) \leq 1$, $\delta(X) \leq 1$. Since \mathbf{t} is a universal uniform test, $\mathbf{t}_n u(x) > 0$. Thus $\delta \ll \nu$, so by properties of the Radon-Nikodym derivative, $\frac{\nu(dx)}{\delta(dx)} = \left(\frac{\delta(dx)}{\nu(dx)}\right)^{-1}$. Using properties of the Radon-Nikodym derivative and Proposition 20,

$$\begin{aligned}\nu(\mu) &= -\mu^x \log \frac{\mu(dx)}{\nu(dx)} \\ -\mu^x \mathbf{H}_\nu(x) &= \mu^x \log \frac{\delta(dx)}{\nu(dx)} = -\mu^x \log \frac{\nu(dx)}{\delta(dx)} \\ \mathcal{H}_\nu(\mu) - \mu^x \mathbf{H}_\nu(x) &= -\mu^x \log \frac{\mu(dx)}{\delta(dx)} \leq -\mu(X) \log \frac{\mu(X)}{\delta(X)} \leq 0.\end{aligned}$$

□

12.5 Addition Equality

For a computable measures μ and ν over a computable metrix space \mathcal{X} , the term $\mathbf{H}_\mu(x|\nu)$ is equal to $\mathbf{H}_\mu(x)$ when the universal Turing machine is given access to a fast Cauchy sequence to ν in the measure space $\mathfrak{M}(X)$. By Proposition 10, this means algorithms can lower compute the ν measure of effectively open sets.

Proposition 21 $\mathbf{H}_\mu(x|\nu) <^+ -\log \nu^y 2^{-\mathbf{H}_{\mu,\nu}(x,y)}$.

Proof. Let $f(x, \mu, \nu) = -\log \nu^y 2^{-\mathbf{H}_{\mu,\nu}(x,y)}$. The function f is upper computable and has $\mu^x 2^{-f(x,\mu,\nu)} \leq 1$. Due to the universal properties of \mathbf{t}_μ and thus minimum property of \mathbf{H}_μ , the inequality is proven. □

Proposition 22 For a computable function $f : N^2 \rightarrow \mathbb{N}$,

$$\mathbf{H}_\mu(x|y) <^+ \mathbf{K}(z) + \mathbf{H}_\mu(x|f(y, z)).$$

Proof. The function

$$g_\mu(x, y) = \sum_z 2^{-\mathbf{H}_\mu(x|f(y,z)) - \mathbf{K}(z)},$$

is lower computable and $\mu^x g_\mu(x, y) \leq \sum_z 2^{-\mathbf{K}(z)} \leq 1$. So $g_\mu(x, y) \stackrel{*}{<} 2^{-\mathbf{H}_\mu(x|y)}$. The left hand side is a summation, so the inequality holds for each element of the sum, proving the proposition. □

Algorithmic fine grained entropy is bounded from above. It can take arbitrarily low negative values, and even $-\infty$.

Proposition 23

1. $\mathbf{H}_\mu(x|\lceil \log \mu(X) \rceil) <^+ \log \mu(X)$.
2. $\mathbf{H}_\mu(x) <^+ \log \mu(X) + \mathbf{K}(\lceil \log \mu(X) \rceil)$.

Proof. For (1), We use the μ -test $t_\mu(x) = 1/\lceil \mu(X) \rceil$, where $\int t_\mu d\mu \leq 1$. Thus $-\log \mu(X) <^+ \log t_\mu(x) <^+ \log \mathbf{t}_\mu(x \lceil \mu(X) \rceil)$. For (2), we use Proposition 22. \square

Proposition 24 *If $i < j$, then*

$$i + \mathbf{H}_\mu(x|i) <^+ j + \mathbf{H}_\mu(x|j).$$

Proof. Using Proposition 22, with $f(i, n) = i + n$, we have

$$\mathbf{H}_\mu(x|i) - \mathbf{H}_\mu(x|j) <^+ \mathbf{K}(j - i) <^+ j - i.$$

The following proposition has a different proof to that of [G21], where Corollary 16 has been introduced, leveraging the results in Chapter 11

Proposition 25 *Let $F : \mathcal{Y} \times \mathcal{M} \rightarrow \mathbb{Z} \cup \{-\infty, \infty\}$. By Corollary 15, among F uniform tests $g_\nu(x, y)$ with $\nu^x g_\nu(x, y) \leq 2^{-F_\nu(y)}$ there is a maximal F uniform test f within a multiplicative constant. For all x with $F_\nu(y) > -\infty$,*

$$f_\nu(x, y) \stackrel{*}{=} 2^{-F_\nu(y)} \mathbf{t}_\nu(x|y, F_\nu(y)).$$

Proof. To prove the inequality $\stackrel{*}{>}$, let $g_\nu(x, y, m) = \max_{i \geq m} 2^{-i} \mathbf{t}_\nu(x|y, i)$. This function is lower computable, and decreasing in m . Let $g_\nu(x, y) = g_\nu(x, y, F_\nu(y))$ is lower semicomputable since F is upper semi-computable. The multiplicative form of Proposition 24 implies

$$\begin{aligned} g_\nu(x, y, m) &\stackrel{*}{=} 2^{-m} \mathbf{t}_\nu(x|y, m) \\ g_\nu(x, y) &\stackrel{*}{=} 2^{-F_\nu(y)} \mathbf{t}_\nu(x|y, F_\nu(y)). \end{aligned}$$

Since \mathbf{t}_ν is a test:

$$\begin{aligned} \nu^x 2^{-m} \mathbf{t}_\nu(x|y, m) &\leq 2^{-m} \\ \nu^x g_\nu(x, y) &\stackrel{*}{<} 2^{-F_\nu(y)}, \end{aligned}$$

which implies $g_\nu(x, y) \stackrel{*}{<} f_\nu(x, y)$ by the optimality of $f_\nu(x, y)$. For the upper bound, consider all lower semicomputable functions $\phi(x, y, m, v)$. By Corollary 16, $f'_\nu(x, y, m)$ be the modification of f , which is a lower computable function such that $\nu^x f'_\nu(x, y, m) \leq 2^{-m+1}$ and if $\nu^x f_\nu(x, y) \leq 2^{-m}$ then $f'_\nu(x, y, m) = f_\nu(x, y)$. The function $2^{m-1} f'_\nu(x, y, m)$ is a uniform test conditioned on y, m so it has $\stackrel{*}{<} \mathbf{t}_\nu(x|y, m)$. Substituting $F_\nu(y)$ for m , we have that $\nu^x f_\nu(x, y) \leq 2^{-m}$ and so

$$f_\nu(x, y) = f'_\nu(x, y, F_\nu(y)) \stackrel{*}{<} 2^{-F_\nu(y)+1} \mathbf{t}_\nu(x|y, F_\nu(y)).$$

\square

Theorem 51

$$\mathbf{H}_{\mu \times \nu} =^+ \mathbf{H}_\mu(x|\nu) + \mathbf{H}_\nu(y|x, \lceil \mathbf{H}_\mu(x|\nu) \rceil, \mu).$$

Proof. We first prove the $<^+$ inequality. Let $G_{\mu,\nu}(x, y, m) = \min_{i \geq m} i + \mathbf{H}_\nu(y|x, i, \mu)$, which is upper computable and increasing in m . So the function

$$G_{\mu,\nu}(x, y) = G_{\mu,\nu}(x, y, \lceil \mathbf{H}_\mu(x|\nu) \rceil).$$

which is also upper computable because m is replaced with an upper computable function $\lceil \mathbf{H}_\mu(x|\nu) \rceil$. Proposition 22 implies

$$\begin{aligned} G_{\mu,\nu}(x, y, m) &=^+ m + \mathbf{H}_\nu(y|x, m, \mu), \\ G_{\mu,\nu}(x, y, m) &=^+ \mathbf{H}_\mu(x|\nu) + \mathbf{H}_\nu(y|x, \mathbf{H}_\mu(x|\nu), \mu). \end{aligned}$$

So

$$\begin{aligned} \nu^y 2^{-m - \mathbf{H}_\nu(y|x, m, \mu)} &\leq 2^{-m} \\ \nu^y 2^{-G_{\mu,\nu}(x, y)} &\stackrel{*}{<} 2^{-\mathbf{H}_\mu(x|\nu)}. \end{aligned}$$

Integrating over x gives $\mu^x \nu^y 2^{-G_{\mu,\nu}(x, y)} \stackrel{*}{<} 1$, implying $\mathbf{H}_{\mu,\nu}(x, y) <^+ G_{\mu,\nu}(x, y)$.

To prove the inequality $>^+$, let $f_\nu(x, y, \mu) = 2^{-\mathbf{H}_{\mu,\nu}(x, y)}$. Proposition 21 implies there exists $c \in \mathbb{N}$ with $\nu^y f_\nu(x, y, \mu) \leq 2^{-\mathbf{H}_\mu(x|\nu) + c}$. Let $F_\nu(x, \mu) = \lceil \mathbf{H}_\mu(x|\nu) \rceil$. Note that if h is a lower computable function such that $\nu^y h(x, y, \mu) \stackrel{*}{<} 2^{-\mathbf{H}_\mu(x|\nu)}$, then $\mu^x \nu^y h(x, y, \mu) \stackrel{*}{<} \mu^x \mathbf{t}_\mu(x|\nu) \stackrel{*}{<} 1$, so $h \stackrel{*}{<} f$. Proposition 25 (substituting y for x and (x, μ) for y) gives

$$\mathbf{H}_{\mu,\nu}(x, y) = -\log f_\nu(x, y, \mu) >^+ F_\nu(x, \mu) + \mathbf{H}_\nu(y|x, F_\nu(x, \mu), \mu).$$

□

Chapter 13

Oscillation of Algorithmic Fine Grained Entropy

In this chapter, it is proven that the algorithmic fine grained entropy of states will oscillate in the presence of continuous dynamics. The theorem matches oscillations proven to occur in the case of discrete ergodic dynamics, seen in Chapter 14. To prove this result, properties about sets of finite and infinite sequences need to be proven, which is done in Sections 13.1 and 13.2.

13.1 On Exotic Sets of Natural Numbers

Lemma 15 *For computable probabilities p, q over \mathbb{N} , $D \subset \mathbb{N}$, $|D| = 2^s$, $s < \max_{a \in D} \min\{\mathbf{d}(a|p), \mathbf{d}(a|q)\} + \mathbf{I}(D; \mathcal{H}) + O(\mathbf{K}(\mathbf{I}(D; \mathcal{H}), p, q, s))$.*

Proof. We relativize the universal Turing machine to $\langle s, p, q \rangle$. Let Q be a probability measure that realizes $\mathbf{K}_s(D)$, with $d = \max\{\mathbf{d}(D|Q), 1\}$. Let $F \subseteq \mathbb{N}$ be a random set where each element $a \in \mathbb{N}$ is selected independently with probability $cd2^{-s}$, where $c \in \mathbb{N}$ is chosen later. $\mathbf{E}[p(F)] = \mathbf{E}[q(F)] \leq cd2^{-s}$. Furthermore

$$\mathbf{E}[Q(\{G : |G| = 2^s, G \cap F = \emptyset\})] \leq \sum_G Q(G)(1 - cd2^{-s})^{2^s} < e^{-cd}.$$

Thus finite $W \subset \mathbb{N}$ can be chosen such that $p(W) \leq 4cd2^{-s}$, $q(W) \leq 4cd2^{-s}$, and $Q(\{G : |G| = 2^s, G \cap W = \emptyset\}) \leq e^{2-cd}$. $D \cap W \neq \emptyset$, otherwise, using the Q -test, $t(G) = e^{cd-1}$ if $(|G| = 2^s, G \cap W = \emptyset)$ and $t(G) = 0$ otherwise, we have

$$\begin{aligned} \mathbf{K}(D|Q, d, c) &<^+ -\log Q(D) - (\log e)cd \\ (\log e)cd &<^+ -\log Q(D) - \mathbf{K}(D|Q) + \mathbf{K}(d, c) \\ (\log e)cd &<^+ d + \mathbf{K}(d, c), \end{aligned}$$

which is a contradiction for large enough c . Thus there is an $a \in D \cap W$, where

$$\begin{aligned} \mathbf{K}(a) &<^+ \min\{-\log q(a), -\log p(a)\} + \log d - s + \mathbf{K}(d) + \mathbf{K}(Q) \\ s &<^+ \min\{\mathbf{d}(a|p), \mathbf{d}(a|q)\} + \mathbf{K}_s(D). \end{aligned}$$

Making the relativization of $\langle s, p, q \rangle$ explicit, and using Lemma 6 results in

$$\begin{aligned} s &<^+ \min\{\mathbf{d}(a|p), \mathbf{d}(a|q)\} + \mathbf{K}s(D) + O(\mathbf{K}(s, p, q)) \\ s &< \max_{a \in D} \min\{\mathbf{d}(a|p), \mathbf{d}(a|q)\} + \mathbf{K}s(D) + O(\mathbf{K}(s, p, q)) \\ s &< \max_{a \in D} \min\{\mathbf{d}(a|p), \mathbf{d}(a|q)\} + \mathbf{I}(D; \mathcal{H}) + O(\mathbf{K}(\mathbf{I}(D; \mathcal{H}), s, p, q)). \square \end{aligned}$$

13.2 On Exotic Sets of Reals

Let $\Omega = \sum\{2^{-\|p\|} : U(p) \text{ halts}\}$ be Chaitin's Omega, $\Omega_n \in \mathbb{Q}_{\geq 0}$ be the rational formed from the first n bits of Ω , and $\Omega^t = \sum\{2^{-\|p\|} : U(p) \text{ halts in time } t\}$. For $n \in \mathbb{N}$, let $\mathbf{bb}(n) = \min\{t : \Omega_n < \Omega^t\}$. $\mathbf{bb}^{-1}(m) = \arg \min_n \{\mathbf{bb}(n-1) < m \leq \mathbf{bb}(n)\}$. Let $\Omega[n] \in \{0, 1\}^*$ be the first n bits of Ω .

Lemma 16 For $n = \mathbf{bb}^{-1}(m)$, $\mathbf{K}(\Omega[n]|m, n) = O(1)$.

Proof. For a string x , let $BB(x) = \inf\{t : \Omega^t > 0.x\}$. Enumerate strings of length n , starting with 0^n , and return the first string x such that $BB(x) \geq m$. This string x is equal to $\Omega[n]$, otherwise let y be the largest common prefix of x and $\Omega[n]$. Thus $BB(y) = \mathbf{bb}(\|y\|) \geq BB(x) \geq m$, which means $\mathbf{bb}^{-1}(m) \leq \|y\| < n$, causing a contradiction. \square

The following lemma, while lengthy, is a series of straightforward application of inequalities.

Lemma 17 For computable probabilities P, Q , over $\{0, 1\}^\infty$, $Z \subset \{0, 1\}^\infty$, $|Z| = 2^s$, $s < \max_{\alpha \in Z} \min\{\mathbf{D}(\alpha|P), \mathbf{D}(\alpha|Q)\} + \mathbf{I}(\langle Z \rangle : \mathcal{H}) + O(\mathbf{K}(s, P, Q) + \log \mathbf{I}(\langle Z \rangle; \mathcal{H}))$.

Proof. We relativize the universal Turing machine to s , which can be done due to the precision of the theorem. Let $Z_n = \{\alpha[0..n] : \alpha \in Z\}$ and $m = \arg \min_m |Z_m| = |Z|$. Let $n = \mathbf{bb}^{-1}(m)$ and $k = \mathbf{bb}(n)$. Let p and q be probabilities over $\{0, 1\}^*$, where $p(x) = [\|x\| = k]P(x)$ and $\langle p \rangle = \langle k, P \rangle$ and let $q(x) = [\|x\| = k]Q(x)$ and $\langle q \rangle = \langle k, P \rangle$. Using $D = Z_k$, Lemma 15, relativized to k , produces $x \in Z_k$, where

$$\begin{aligned} s &< \min\{\mathbf{d}(x|p), \mathbf{d}(x|q)\} + \mathbf{I}(Z_k; \mathcal{H}|k) + O(\mathbf{K}(\mathbf{I}(Z_k; \mathcal{H}|k), q, p|k)) \\ &< \max_{\alpha \in Z} \min\{\mathbf{D}(\alpha|P), \mathbf{D}(\alpha|Q)\} + \mathbf{K}(Z_k|k) + \mathbf{K}(k) - \mathbf{K}(Z_k|k, \mathcal{H}) + O(\mathbf{K}(\mathbf{I}(Z_k; \mathcal{H}|k), q, p|k)). \\ &< \max_{\alpha \in Z} \min\{\mathbf{D}(\alpha|P), \mathbf{D}(\alpha|Q)\} + \mathbf{K}(Z_k|k) + \mathbf{K}(k) - \mathbf{K}(Z_k|k, \mathcal{H}) + O(\mathbf{K}(P, Q) + \log \mathbf{I}(Z_k; \mathcal{H}|k)). \end{aligned}$$

Since $\mathbf{K}(k) <^+ n + \mathbf{K}(n)$, by the chain rule,

$$\begin{aligned} &\mathbf{K}(Z_k|k) + \mathbf{K}(k) \\ &<^+ \mathbf{K}(Z_k|k, \mathbf{K}(k)) + \mathbf{K}(\mathbf{K}(k)|k) + \mathbf{K}(k) \\ &< \mathbf{K}(Z_k, k) + O(\log n) \\ &< \mathbf{K}(Z_k) + O(\log n). \end{aligned}$$

So

$$s < \max_{\alpha \in Z} \min\{\mathbf{D}(\alpha|P), \mathbf{D}(\alpha|Q)\} + \mathbf{K}(Z_k) - \mathbf{K}(Z_k|k, \mathcal{H}) + O(\log n + \mathbf{K}(P, Q) + \log \mathbf{I}(Z_k; \mathcal{H}|k)).$$

Since $\mathbf{K}(k|n, \mathcal{H}) = O(1)$, $\mathbf{K}(Z_k|\mathcal{H}) <^+ \mathbf{K}(Z_k|k, \mathcal{H}) + \mathbf{K}(n)$,

$$s < \max_{\alpha \in Z} \min\{\mathbf{D}(\alpha|P), \mathbf{D}(\alpha|Q)\} + \mathbf{I}(Z_k; \mathcal{H}) + O(\log n + \mathbf{K}(P, Q) + \log \mathbf{I}(Z_k; \mathcal{H}|k)).$$

Furthermore since $\mathbf{I}(Z_k; \mathcal{H}|k) + \mathbf{K}(k) < \mathbf{I}(Z_k; \mathcal{H}) + O(\log n)$,

$$s < \max_{\alpha \in Z} \min\{\mathbf{D}(\alpha|P), \mathbf{D}(\alpha|Q)\} + \mathbf{I}(Z_k; \mathcal{H}) + O(\log n + \mathbf{K}(P, Q)) + O(\log \mathbf{I}(Z_k; \mathcal{H})).$$

By Lemma 16, $\mathbf{K}(\Omega[n]|Z_k) <^+ \mathbf{K}(n)$ so by Lemma 4,

$$n <^{\log} \mathbf{I}(\Omega[n]; \mathcal{H}) <^{\log} \mathbf{I}(Z_k; \mathcal{H}) + \mathbf{K}(n) <^{\log} \mathbf{I}(Z_k; \mathcal{H}).$$

The above equation used the common fact that the first n bits of Ω has $n - O(\log n)$ bits of mutual information with \mathcal{H} . So

$$s < \max_{\alpha \in Z} \min\{\mathbf{D}(\alpha|P), \mathbf{D}(\alpha|Q)\} + \mathbf{I}(Z_k; \mathcal{H}) + O(\mathbf{K}(P, Q) + \log \mathbf{I}(Z_k; \mathcal{H})).$$

By the definition of mutual information \mathbf{I} between infinite sequences

$$\mathbf{I}(Z_k; \mathcal{H}) <^+ \mathbf{I}(Z : \mathcal{H}) + \mathbf{K}(Z_k|Z) <^{\log} \mathbf{I}(Z : \mathcal{H}) + \mathbf{K}(k|Z).$$

Now m is simple relative to Z and by Lemma 16, $\Omega[n]$ is simple relative to m and n . Furthermore k is simple relative to $\Omega[n]$. Therefore $\mathbf{K}(Z_k|Z) <^+ \mathbf{K}(n)$. So

$$s < \max_{\alpha \in Z} \min\{\mathbf{D}(\alpha|P), \mathbf{D}(\alpha|Q)\} + \mathbf{I}(Z : \mathcal{H}) + O(\log n) + O(\mathbf{K}(P, Q) + \log \mathbf{I}(Z; \mathcal{H}))$$

$$s < \max_{\alpha \in Z} \min\{\mathbf{D}(\alpha|P), \mathbf{D}(\alpha|Q)\} + \mathbf{I}(Z : \mathcal{H}) + O(\mathbf{K}(s, P, Q) + \log \mathbf{I}(Z; \mathcal{H})).$$

□

13.3 Oscillations Occur

Definition 27 (Mutual Information with the Halting Sequence) *An encoding of a fast Cauchy sequence \vec{x} is $\langle \vec{x} \rangle \in \{0, 1\}^\infty$, with $\langle \vec{x} \rangle = \langle x_1 \rangle \langle x_2 \rangle \dots$. Each $x_i \in \mathcal{S}$ is an ideal point, and $\langle x_i \rangle$ is its order in the enumeration of \mathcal{S} . Each point $x \in \mathcal{X}$ has a certain mutual information with the halting sequence $\mathbf{I}(x : \mathcal{H}) = \inf\{\mathbf{I}(\langle \vec{x} \rangle : \mathcal{H}) : \langle \vec{x} \rangle \text{ is a fast Cauchy sequence for } x\}$.*

Theorem 52 ([Ver21, Lev74]) *Let P_ρ be a family of probability distributions over $\{0, 1\}^\infty$, indexed by $\rho \in \{0, 1\}^\infty$. Assume that there is a Turing machine T such that for all $\rho \in \{0, 1\}^\infty$ computes P_ρ having oracle access to ρ . By “compute” we mean all the measures of the cylinder sets $P_\rho(x\{0, 1\}^\infty)$, can be computed, uniformly in $x \in \{0, 1\}^*$. Then there is a constant $c_T > 0$ solely dependent on T such that*

$$P_\rho\{\gamma : \mathbf{I}(\langle \gamma, \rho \rangle : \mathcal{H}) > m\} < 2^{\mathbf{I}(\rho; \mathcal{H}) - m + c_T}.$$

The following theorem shows oscillations of entropy occur during dynamics. Theorem 70 in Chapter 17 is more general and the proof is more straightforward, but the bounds are looser. Note that the measure preserving requirement for the transformation group is not need for Theorem 53.

Theorem 53 (Oscillation of Thermodynamic Entropy) *Let L be the Lebesgue measure over \mathbb{R} , (\mathcal{X}, μ) be a computable measure space, $\alpha \in \mathcal{X}$, with finite $\mathbf{I}(\alpha : \mathcal{H})$. For transformation group G^t acting on \mathcal{X} , there is a constant c with $L\{t \in [0, 1] : \mathbf{H}_\mu(G^t \alpha) < \log \mu(X) - n\} > 2^{-n - \mathbf{K}(n) - c}$.*

Proof. We first assume not. There exists (G^t, \mathcal{X}) and computable measure space (\mathcal{X}, μ) and there exists $\alpha \in X$ such that for all $c \in \mathbb{N}$, there exists n , where

$$\begin{aligned} L(\{t \in [0, 1] : \mathbf{H}_\mu(G^t \alpha) < \log \mu(\mathcal{X}) - n\}) &< 2^{-n-\mathbf{K}(n)-c} \\ L(\{t \in [0, 1] : n - \log \mu(\mathcal{X}) < \log \mathbf{t}_\mu(G^t \alpha)\}) &< 2^{-n-\mathbf{K}(n)-c}. \end{aligned}$$

We sample $2^{n+\mathbf{K}(n)+c-1}$ elements F by choosing a time t uniformly between $[0, 1]$. The probability that all samples $\beta \in F$ have $\mathbf{t}_\mu(G^\beta \alpha) \leq n - \log \mu(\mathcal{X})$ is

$$\begin{aligned} &\prod_{i=1}^{|F|} L\{t \in [0, 1] : \log \mathbf{t}_\mu(G^t \alpha) \leq n - \log \mu(\mathcal{X})\} \\ &\geq (1 - |F| 2^{-n-\mathbf{K}(n)-c}) \\ &\geq (1 - 2^{n+\mathbf{K}(n)+c-1} 2^{-n-\mathbf{K}(n)-c}) \\ &\geq 1/2. \end{aligned}$$

Let $(\{0, 1\}^\infty, \Gamma)$ be the Cantor space with the uniform measure. The binary representation (see Theorem 48) creates an isomorphism (ϕ, ϕ^{-1}) of computable probability spaces between the spaces $(\{0, 1\}^\infty, \Gamma)$ and $([0, 1], L)$. It is the canonical function $\phi(\gamma) = 0.\gamma$. Thus for all Borel sets $A \subseteq [0, 1]$, $\Gamma(\phi^{-1}(A)) = L(A)$. Since $\{t \in [0, 1] : \log \mathbf{t}_\mu(G^t \alpha) \leq n - \log \mu(\mathcal{X})\}$ is closed,

$$L\{t \in [0, 1] : \log \mathbf{t}_\mu(G^t \alpha) \leq n - \log \mu(\mathcal{X})\} = \Gamma\{\gamma \in \{0, 1\}^\infty : \log \mathbf{t}_\mu(G^{\phi(\gamma)} \alpha) \leq n - \log \mu(\mathcal{X})\}.$$

So

$$1/2 \leq \prod_{i=1}^{|F|} \Gamma\{\gamma \in \{0, 1\}^\infty : \log \mathbf{t}_\mu(G^{\phi(\gamma)} \alpha) \leq n - \log \mu(\mathcal{X})\}.$$

Let (δ, μ_δ) be a binary representation (see Definition 20), for the computable measure space (\mathcal{X}, μ) . Thus μ_δ is a computable (not necessarily probability) measure over $\{0, 1\}^\infty$. By Lemma 13, there is a $c' > 0$, where

$$\prod_{i=1}^{|F|} \Gamma\{\gamma \in \{0, 1\}^\infty : \log \mathbf{t}_{\mu_\delta}(\delta^{-1}(G^{\phi(\gamma)} \alpha)) \leq n - \log \mu(\mathcal{X}) + c'\} \geq 1/2.$$

Let $f : \{0, 1\}^\infty \times \{0, 1\}^\infty \rightarrow \{0, 1\}^\infty$, where $f(\gamma, \langle \vec{\zeta} \rangle) = \delta^{-1}(G^{\phi(\gamma)} \zeta)$. Note, $f(\gamma, \langle \vec{\zeta} \rangle)$ can be undefined when $\mathbf{t}_\mu(G^{\phi(\gamma)} \zeta) = \infty$, because the morphism δ^{-1} is only proven to be defined on a constructive G_δ set of full measure which includes random points. Let $\xi = \langle \vec{\alpha} \rangle$ be an encoding of a fast Cauchy sequence $\vec{\alpha}$ such that $\mathbf{I}(\xi : \mathcal{H}) < \infty$. The sequence ξ is guaranteed to exist because the assumption of the theorem statement. So

$$\prod_{i=1}^{|F|} \Gamma\{\gamma \in \{0, 1\}^\infty : \log \mathbf{t}_{\mu_\delta}(f(\gamma, \xi)) \leq n - \log \mu(\mathcal{X}) + c'\} \geq 1/2.$$

By Proposition 14, (and also updating c')

$$\prod_{i=1}^{|F|} \Gamma\{\gamma \in \{0, 1\}^\infty : \mathbf{D}(f(\gamma, \xi) | \mu_\delta) \leq n - \log \mu(\mathcal{X}) + c' + \mathbf{K}(\mu_\delta)\} \geq 1/2.$$

Let $\bar{\mu}_\delta(\alpha) = \mu_\delta(\alpha)/\mu_\delta(\{0,1\}^\infty)$, which is a computable probability measure over $\{0,1\}^\infty$.

$$\prod_{i=1}^{|F|} \Gamma\{\gamma \in \{0,1\}^\infty : \mathbf{D}(f(\gamma, \xi)|\bar{\mu}_\delta) \leq n + c' + \mathbf{K}(\mu_\delta)\} \geq 1/2.$$

Let Γ^{n+c} be a computable distribution over the product of $1 + 2^{n+\mathbf{K}(n)+c-1}$ independent probability measures over $\{0,1\}^\infty$, encoding into a $\{0,1\}^\infty$ in the standard way. The first probability distribution gives measure 1 to ξ and the last $2^{n+\mathbf{K}(n)+c}$ probability measures are the uniform distribution Γ over $\{0,1\}^\infty$. So

$$\Gamma^{n+c}(\text{Encoding of } 1 + 2^{n+\mathbf{K}(n)+c-1} \text{ elements with the first encoded sequence being } \xi \\ \text{and the rest of encoded sequences } \beta \text{ has } \mathbf{D}(f(\beta, \xi)|\bar{\mu}_\delta) \leq n + c' + \mathbf{K}(\mu_\delta)) \geq 1/2.$$

Let $n^* = \langle n, \mathbf{K}(n) \rangle$. There is an infinite sequence $\eta = \langle n, \mathbf{K}(n), c \rangle \xi$ and a Turing machine T , such that T computes Γ^{n+c} when given oracle access to η . By Theorem 52, with the universal Turing machine relativized to n^* , and folding the constants together,

$$\begin{aligned} & \Gamma^{n+c}(\{\gamma : \mathbf{I}(\gamma : \mathcal{H}|n^*) > m\}) \\ & < \Gamma^{n+c}(\{\gamma : \mathbf{I}(\langle \gamma, \eta \rangle : \mathcal{H}|n^*) >^+ m\}) \\ & <^* 2^{-m+\mathbf{I}(\eta:\mathcal{H}|n^*)+c_T} \\ & <^* 2^{-m+\mathbf{K}(n, \mathbf{K}(n), c|n^*)+\mathbf{I}(\xi:\mathcal{H}|n^*)+c_T} \\ & <^* 2^{-m+\mathbf{K}(c)}. \end{aligned}$$

Therefore,

$$\Gamma^{n,c}(\{\gamma : \mathbf{I}(\gamma : \mathcal{H}|n^*) >^+ \mathbf{K}(c)\}) \leq 1/4.$$

Thus, by probabilistic arguments, there exists $\kappa \in \{0,1\}^\infty$, such that $\kappa = \langle D, \xi \rangle$, where $D \subset \{0,1\}^\infty$ and $|D| = 2^{n+\mathbf{K}(n)+c-1}$ and each $\beta \in D$ has $\mathbf{D}(f(\beta, \xi)|\bar{\mu}_\delta) \leq n+c'+\mathbf{K}(\mu_\delta)$ and $\mathbf{I}(\kappa : \mathcal{H}|n^*) <^+ \mathbf{K}(c)$. Thus since $\mathbf{K}(f(D, \xi)|\kappa, n^*) = O(1)$ we have $\mathbf{I}(f(D, \xi) : \mathcal{H}|n^*) <^+ \mathbf{I}(\kappa : \mathcal{H}|n^*) <^+ \mathbf{K}(c)$. By Lemma 17, relativized to n^* , on the set $D' = f(D, \xi)$ and probability $\bar{\mu}_\delta$, there exists constants $d, f \in \mathbb{N}$ where

$$\begin{aligned} m = \log |D| & < \max_{\beta \in D'} \mathbf{D}(\beta|\bar{\mu}_\delta, n^*) + 2\mathbf{I}(D' : \mathcal{H}|n^*) + d\mathbf{K}(m|v) + f\mathbf{K}(\bar{\mu}_\delta|n^*) \\ m & < \max_{\beta \in D'} \mathbf{D}(\beta|\bar{\mu}_\delta) + \mathbf{K}(n) + 2\mathbf{I}(D' : \mathcal{H}|n^*) + d\mathbf{K}(m|n^*) + f\mathbf{K}(\mu_\delta|n^*) \\ & <^+ \max_{\beta \in D'} \mathbf{D}(\beta|\bar{\mu}_\delta) + \mathbf{K}(n) + 2\mathbf{K}(c) + d\mathbf{K}(m|v) + f\mathbf{K}(\mu_\delta|n^*) \\ & <^+ n + \mathbf{K}(n) + d\mathbf{K}(m|v) + 2\mathbf{K}(c) + (f+1)\mathbf{K}(\mu_\delta). \end{aligned} \tag{13.1}$$

Therefore:

$$\begin{aligned} m &= n + \mathbf{K}(n) + c - 1 \\ \mathbf{K}(m|n^*) &<^+ \mathbf{K}(c). \end{aligned} \tag{13.2}$$

Plugging Equation 13.2 back into Equation 16.1 results in

$$\begin{aligned} n + \mathbf{K}(n) + c &<^+ n + \mathbf{K}(n) + 2\mathbf{K}(c) + d(\mathbf{K}(c) + O(1)) + (f+1)\mathbf{K}(\mu_\delta) \\ c &<^+ (2+d)\mathbf{K}(c) + dO(1) + (f+1)\mathbf{K}(\mu_\delta). \end{aligned}$$

This result is a contradiction for sufficiently large c solely dependent \mathcal{X} , G , μ , and the universal Turing machine. \square

Corollary 18 (Oscillation of Marginal Thermodynamic Entropy) *Let L be the Lebesgue measure over \mathbb{R} , $(\mathcal{X} \times \mathcal{Y}, \mu \times \nu)$ be a computable product measure space, $(\alpha, \beta) \in \mathcal{X} \times \mathcal{Y}$, with finite $\mathbf{I}((\alpha, \beta) : \mathcal{H})$. For transformation group G^t acting on $\mathcal{X} \times \mathcal{Y}$, there is a constant c where if $(\alpha^t, \beta^t) = G^t(\alpha, \beta)$, then $L\{t \in [0, 1] : \mathbf{H}_\mu(\alpha^t) < \log \mu(X) - n\} > 2^{-n - \mathbf{K}(n) - c}$.*

13.4 Oscillations are Rare

The following lemma shows that Theorem 53 is tight. It is the same statement as Theorem 6 in [Gac94], with an updated proof.

Lemma 18 *Let L be the Lebesgue measure over \mathbb{R} , (\mathcal{X}, μ) be a computable measure space, and $\alpha \in \mathcal{X}$. For transformation group G^t acting on \mathcal{X} , there is a constant c where $L\{t \in [0, 1] : \mathbf{H}_\mu(G^t \alpha) < \mathbf{H}_\mu(\alpha) - m\} < 2^{-m+c}$.*

Proof. Since

$$\int_{\mathcal{X}} \int_{[0,1]} 2^{-\mathbf{H}_{\mu \times L}(\alpha, t)} dL(t) d\mu(\alpha) = \int_{\mathcal{X}} \int_{[0,1]} \mathbf{t}_{\mu \times L}(\alpha, t) dL(t) d\mu(\alpha) \leq 1,$$

the function $f(\alpha) = \int_{[0,1]} 2^{-\mathbf{H}_{\mu \times L}(\alpha, t)} dL(t)$ is a μ -test. So

$$\int_{[0,1]} 2^{-\mathbf{H}_{\mu \times L}(\alpha, t)} dt = f(\alpha) \stackrel{*}{<} \mathbf{t}_\mu(\alpha) \stackrel{*}{=} 2^{-\mathbf{H}_\mu(\alpha)}.$$

So

$$\begin{aligned} \{t \in [0, 1] : 2^{-\mathbf{H}_{\mu \times L}(\alpha, t)} > 2^{m - \mathbf{H}_\mu(\alpha)}\} &\stackrel{*}{<} 2^{-m} \\ \{t \in [0, 1] : \mathbf{H}_{\mu \times L}(\alpha, t) < \mathbf{H}_\mu(\alpha) - m\} &\stackrel{*}{<} 2^{-m} \end{aligned}$$

$\mathbf{H}_{\mu \times L}(\alpha, t) <^+ \mathbf{H}_\mu(G^t \alpha)$ because

$$\begin{aligned} &\int_{[0,1]} \int_{\mathcal{X}} \mathbf{t}_\mu(G^t \alpha) d\mu(\alpha) dL(t) \\ &= \int_{[0,1]} \int_{\mathcal{X}} \mathbf{t}_\mu(\alpha) d\mu(G^{-t} \alpha) dL(t) \\ &= \int_{[0,1]} \int_{\mathcal{X}} \mathbf{t}_\mu(\alpha) d\mu(\alpha) dL(t), \\ &= \int_{[0,1]} 1 dL(t) \\ &\leq 1, \end{aligned}$$

which means $\mathbf{t}_\mu(G^t \alpha) \stackrel{*}{<} \mathbf{t}_{\mu \times L}(\alpha, t)$ and thus $2^{-\mathbf{H}_\mu(G^t \alpha)} \stackrel{*}{<} 2^{-\mathbf{H}_{\mu \times L}(\alpha, t)}$. Thus

$$\{t \in [0, 1] : \mathbf{H}_\mu(G^t \alpha) < \mathbf{H}_\mu(\alpha) - m\} \stackrel{*}{<} 2^{-m}.$$

□

Combining Theorem 53 and Lemma 18 together, one gets a full characterization of the dynamics of states in the phase space. This corollary was shown in the introduction.

Corollary 19 *Let L be the Lebesgue measure over \mathbb{R} , and (\mathcal{X}, μ) be a computable measure space, and $\alpha \in \mathcal{X}$ with finite $\mathbf{I}(\alpha : \mathcal{H})$. For transformation group G^t acting on \mathcal{X} , there are constants c_1 and c_2 with*

$$2^{-n-\mathbf{K}(n)-c_1} < L\{t \in [0, 1] : \mathbf{H}_\mu(G^t \alpha) < \log \mu(\mathcal{X}) - n\} < 2^{-n+c_2}.$$

Chapter 14

Discrete Dynamics

In this chapter, results about discrete dynamics are detailed. Theorem 22 shows that oscillations occur in ergodic dynamics. This closely mirrors Corollary 19, which characterizes continuous dynamics.

Definition 28 (Discrete Dynamics) *Discrete dynamics is modeled by a transform group G^t from Definition 25, but with $t \in \mathbb{Z}$, being an integer. We assume there no $\alpha \in \mathcal{X}$ with a finite orbit.*

Proposition 26 *The sets $\{\alpha : \mathbf{H}_\mu(\alpha) \neq -\infty\}$ and $\{\alpha : \mathbf{H}_\mu(\alpha) = -\infty\}$ are conserved under a discrete transformation group G .*

Proof. If $\alpha \in B$, then α is in an effective null set $\bigcap_n U_n$. Thus $G^t \alpha$ is in the null set $\bigcap_n G^t U_n$ and thus $\mathbf{H}_\mu(G^t \alpha) = -\infty$. The same reasoning for $-t$ proves the case when $\mathbf{H}_\mu(\alpha) \neq -\infty$.

14.1 Synchronized Oscillations

Discrete dynamics will visit states with ever increasing \mathbf{t}_μ and \mathbf{t}_ν score. Given a finite set $D \subset \mathcal{X}$, with $D = \{\alpha_i\}_{i=1}^n$, its mutual information with the halting sequence is defined by $\mathbf{I}(D : \mathcal{H}) = \inf_{\vec{\alpha}_1, \dots, \vec{\alpha}_n} \mathbf{I}(\langle \vec{\alpha}_1, \dots, \vec{\alpha}_n \rangle : \mathcal{H})$, which is the infimum over all encoded fast Cauchy sequences to members of D .

Lemma 19 *Given dual computable measure space (\mathcal{X}, μ, ν) , there is a constant $c_{\mathcal{X}, \mu, \nu}$, with universal uniform tests \mathbf{t}_μ and \mathbf{t}_ν , such that $U = \mu(\mathcal{X}) = \nu(\mathcal{X})$, computable U , for a finite set $Z \subset \mathcal{X}$ with $n = \lceil \log |Z| \rceil$,*

$$n < \log \max_{\alpha \in Z} \min\{\mathbf{t}_\mu(\alpha), \mathbf{t}_\nu(\alpha)\} + U + \mathbf{I}(\langle Z \rangle : \mathcal{H}) + O(\log \mathbf{I}(\langle Z \rangle : \mathcal{H}) + \mathbf{K}(n) + c_{\mathcal{X}, \mu, \nu}).$$

Proof. Let $(\{0, 1\}^\infty, \mu_\delta, \nu_\delta)$ be a dual binary representation that are isomorphic to computable measure spaces (\mathcal{X}, μ) and (\mathcal{X}, ν) , with $\delta : (\{0, 1\}^\infty, \mu_\delta) \rightarrow (\mathcal{X}, \mu)$ and $\delta : (\{0, 1\}^\infty, \nu_\delta) \rightarrow (\mathcal{X}, \nu)$. If $\max_{\alpha \in Z} \min\{\mathbf{t}_\mu(\alpha), \mathbf{t}_\nu(\alpha)\} = \infty$, then the lemma is proven. Thus for all $\alpha \in Z$, either $\mathbf{t}_\mu(\alpha) < \infty$ or $\mathbf{t}_\nu(\alpha) < \infty$, so by Lemma 13, $\delta^{-1}(\alpha)$ is defined. Let $\bar{\mu}_\delta = \mu_\delta/U$ and $\bar{\nu}_\delta = \nu_\delta/U$ be computable probability measures over $\{0, 1\}^\infty$. Let $W = \delta^{-1}(Z) \subset \{0, 1\}^\infty$. By Theorem 17 applied to W , $\bar{\mu}_\delta$, and $\bar{\nu}_\delta$ with $s = n - O(1)$, gives

$$s < \max_{\alpha \in W} \min\{\mathbf{D}(\alpha | \bar{\mu}_\delta), \mathbf{D}(\alpha | \bar{\nu}_\delta)\} + \mathbf{I}(W : \mathcal{H}) + O(\log \mathbf{I}(W : \mathcal{H}) + \mathbf{K}(s) + c_{\mathcal{X}, \mu, \nu}).$$

Due to Proposition 14,

$$\begin{aligned} n &< \max_{\alpha \in W} \min\{\log \mathbf{t}_{\overline{\mu\delta}}(\alpha), \log \mathbf{t}_{\overline{\nu\delta}}(\alpha)\} + \mathbf{I}(W : \mathcal{H}) + O(\log \mathbf{I}(W : \mathcal{H}) + \mathbf{K}(n) + c_{\mathcal{X},\mu}), \\ n &< \max_{\alpha \in W} \min\{\log \mathbf{t}_{\mu\delta}(\alpha), \log \mathbf{t}_{\nu\delta}(\alpha)\} + U + \mathbf{I}(W : \mathcal{H}) + O(\log \mathbf{I}(W : \mathcal{H}) + \mathbf{K}(n) + c_{\mathcal{X},\mu}). \end{aligned}$$

Since $(\{0, 1\}^\infty, \mu_\delta)$ is isomorphic to (\mathcal{X}, μ) and $(\{0, 1\}^\infty, \nu_\delta)$ is isomorphic to (\mathcal{X}, ν) , due to Lemma 13,

$$n < \max_{\alpha \in Z} \min\{\log \mathbf{t}_\mu(\alpha), \log \mathbf{t}_\nu(\alpha)\} + U + \mathbf{I}(W : \mathcal{H}) + O(\log \mathbf{I}(W : \mathcal{H}) + \mathbf{K}(n) + c_{\mathcal{X},\mu}).$$

Given any encoding of the fast Cauchy sequences of the members of Z , one can compute W with δ^{-1} , thus $\mathbf{K}(W|Z) = O(1)$, so

$$n < \max_{\alpha \in Z} \min\{\log \mathbf{t}_\mu(\alpha), \log \mathbf{t}_\nu(\alpha)\} + U + \mathbf{I}(Z : \mathcal{H}) + O(\log \mathbf{I}(Z : \mathcal{H}) + \mathbf{K}(n) + c_{\mathcal{X},\mu}).$$

□

Theorem 54 *Let (\mathcal{X}, μ, ν) be a dual computable measure space, with $U = \mu(\mathcal{X}) = \nu(\mathcal{X})$ and $\alpha \in \mathcal{X}$, with finite $\mathbf{I}(\alpha : \mathcal{H})$. For discrete time dynamics G^t , there is a c such that*

$$\max_{\gamma \in G^{\{1, \dots, 2^n\}} \alpha} \max\{\mathbf{H}_\mu(\gamma), \mathbf{H}_\nu(\gamma)\} < U - n + O(\mathbf{K}(n)) + c.$$

Proof. Let $Z_n = G^{\{1, \dots, 2^n\}} \alpha$. Lemma 19, applied to (\mathcal{X}, μ, ν) and Z_n , results in $\gamma \in Z_n$ such that

$$n < \min\{\log \mathbf{t}_\mu(\gamma), \log \mathbf{t}_\nu(\gamma)\} + U + \mathbf{I}(Z_n : \mathcal{H}) + O(\log \mathbf{I}(Z_n : \mathcal{H}) + \mathbf{K}(n) + c_{\mathcal{X},\mu,\nu,\alpha}).$$

Since $\mathbf{I}(Z_n : \mathcal{H}) <^+ \mathbf{I}(\{\alpha\} : \mathcal{H}) + \mathbf{K}(n)$,

$$n < \min\{\log \mathbf{t}_\mu(\gamma), \log \mathbf{t}_\nu(\gamma)\} + U + \mathbf{I}(\alpha : \mathcal{H}) + O(\log \mathbf{I}(\alpha : \mathcal{H}) + \mathbf{K}(n) + c_{\mathcal{X},\mu,\nu,\alpha,G}).$$

The theorem is proven by noting $\mathbf{I}(\alpha : \mathcal{H}) < \infty$. □

14.2 Ergodic Dynamics

For a measure space (\mathcal{X}, μ) a function $T : X \rightarrow X$ is ergodic if it is measure preserving and all the invariant sets have measure 0 or $\mu(X)$. The following theorem adapts Theorem y of [BDH⁺12] to computable measure spaces using the recommendations of the proof sketch of Theorem 12. Comparable results can be found in [FMN12]. The main difference between this proof and that of Theorem 7 is that overlapping open balls are used instead of cylinders.

14.2.1 Single Points

Theorem 55 *Let (\mathcal{X}, μ) be a computable measure space. Let $T : X \rightarrow X$ be an computable ergodic function. Let A be an effectively open subset of X , where $\mu(A) < \mu(X)$. Let A^* be the set of points $x \in X$ such that $T^i(x) \in A$ for all $i \geq 0$. Then $-\mathbf{H}_\mu(x) = -\infty$ for all $x \in A^*$.*

It is sufficient to prove A^* is an effectively null set, introduced in Definition 22. We recall that from Corollary 14, there is an enumeration $\{B_i\}$ of the basis of “almost decidable” of open balls such that their borders have null μ -measure. Let $\nu(x) = \mu(x)/\mu(X)$ be a computable probability measure over \mathcal{X} , due to Claim 1. Let r be a real number such that $\nu(A) < r < 1$. Given an enumerated ball B_j , we want to find an n such that $\nu(B_j \cap \bigcap_{i \leq n} T^{-i}(A)) \leq r\nu(B)$. Note that it could be that $B_j \cap B_k \neq \emptyset$ for $j \neq k$. This gives an effective open cover of $A^* \cap I$ having measure at most $r\nu(B_j)$. For each j you iterate the process until you get the an effectively open cover of $B_j \cap A^*$ with measure $< r2^{-j}\nu(B_j)$. Thus the union of all effectively open covers of A^* has measure less than r . This process is repeated without end to get an ν effectively null set.

To estimate $\nu(B \cap \bigcap_{i \leq n} T^{-i}(A))$, we note that it does not exceed $\min_i \nu(B \cap T^{-i}(A))$ which does not exceed $\frac{1}{n+1} \sum_{i \leq n} \nu(B \cap T^{-i}(A))$. This average,

$$\frac{1}{n+1} [\nu(B \cap A) + \nu(B \cap T^{-1}(A)) + \dots + \nu(B \cap T^{-n}(A))] \quad (14.1)$$

is equal to

$$\frac{1}{n+1} [\nu(T^{-n}(B) \cap T^{-n}(A)) + \nu(T^{-(n-1)}(B) \cap T^{-n}(A)) + \dots + \nu(B \cap T^{-n}(A))],$$

because G is measure preserving. The latter expression is the scalar product of the indicator function of $T^{-n}(A)$ and the average $a_n = (\mathbf{1}_0 + \dots + \mathbf{1}_n)/(n+1)$, where $\mathbf{1}_i$ is the indicator of $T^{-i}(B)$.

As $n \rightarrow \infty$, the average a_n converges in L_2 to the constant function $\nu(B)$ due to von Neumann’s mean ergodic theorem. By Cauchy-Schwarz inequality this means the the scalar product converges to $\nu(A)\nu(B)$, so it does not exceed $r\nu(B)$ for n large enough.

It remains to find an effective value for n for which the L_2 -distance between a_n and the constant function $\nu(B)$ is small. Note that for all i the set $T^{-i}(B)$ is an effectively open set of measure $\nu(B)$, and, since T is measure preserving, $\nu(B)$ is computable. There for any i and $\epsilon > 0$, one can uniformly approximate $T^{-i}(B)$ by its ad-set subset U (see Definition 21) where $\nu(T^{-i}(B) \setminus U) < \epsilon$ can be computed, due to Proposition 13. This means that the L_2 -distance a_n and the constant function $\nu(B)$ can be computed effectively, so one can continue computing this value until it finds and n such that the average (Equation 14.1) is less than $r\nu(B)$. We then have $\nu(B \cap \bigcap_{i \leq n} T^{-i}(A)) < r\nu(B)$. \square

The above theorem has implications for algorithmic coarse grain entropy and in particular Theorem 63 which says that if a state travels through enough partitions (effective open sets) then oscillations will occur. Theorem 55 says that a state $x \in X$, with $\mathbf{H}_\mu(x) \neq -\infty$, under ergodic dynamics will travel through all the partitions if there are finitely many of them or an ever increasing number of partitions if there are infinite many of them.

14.2.2 Indicator Functions

The following theorem adapts Theorem 8 from [BDH⁺12] to computable measure space using the proof sketch in Theorem 12.

Theorem 56 *Let (\mathcal{X}, μ) be a computable measure space. Let $T : X \rightarrow X$ be a computable ergodic function. Let U be an effectively open set. If $\mathbf{H}_\mu(\omega) \neq -\infty$ then*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} \mathbf{1}_U(T^k(\omega)) = \mu(U)/\mu(X).$$

This also applies to effectively closed sets.

Proof. Let $\nu(x) = \mu(x)/\mu(X)$ be a computable probability measure over \mathcal{X} due to Claim 1. Let $g_n(\omega) = \frac{1}{n} \sum_{k=0}^{n-1} \mathbf{1}_U(T^k(\omega))$ be the frequency of U elements among the first n iterations of ω . We first prove $\limsup g_n(\omega) \leq \nu(U)$. We then prove $\liminf g_n(\omega) \geq \nu(U)$.

Let $r > \nu(U)$ be some rational number and $G_N = \{\omega : (\exists n \geq N) g_n(\omega) > r\}$ be the set of points where some far enough frequency exceeds r . The set G_N is an effectively open set; the functions g_n are lower computable uniformly in n ; the condition $g_n(\omega) > r$ is enumerable. The set G_N is decreasing in N . By the classical Birkhoff's pointwise ergodic theorem that $\nu(\bigcap_N G_N) = 0$ as the sequence of functions g_n converges to $\nu(U) < r$ ν -almost everywhere. So there exists N , where $\nu(G_N) < 1$. We can then apply Theorem 55 to effectively open set G_N we get that for $\omega \in U$ with $\mathbf{H}_\mu(\omega) \neq -\infty$ and k such that $T^k(\omega) \notin G_N$. So $\limsup_n g_n(T^k(\omega)) \leq r$. Since finite number of iterations does not change \limsup , we have $g_n(\omega) \leq r$. Since r was an arbitray rational number greater than $\nu(U)$ so $\limsup g_n(\omega) \geq \nu(U)$.

(2) We now prove that $\liminf g_n(\omega) \geq \nu(U)$. Since U is open it is a countable union of almost decidable balls. Taking ad-set $D \subset U$. We can apply the previous statement to $X \setminus \overline{D}$. It says the orbit of a point ω with $\mathbf{H}_\mu(\omega) \neq -\infty$ will be in \overline{D} with frequency at least $\nu(\overline{D}) = \nu(D)$. Since $\nu(D)$ can be arbitrarily close to $\nu(U)$, we have that $\liminf g_n(\omega) \geq \nu(U)$. \square

Proposition 27 *For computable non-atomic measure space (\mathcal{X}, μ) ,*

1. *Every set E with $\mu(E) > 0$ contains sets of arbitrarily small positive measure.*
2. *For any $\delta \in [0, \mu(X)]$, there exists an open set A where $\frac{1}{2}\delta \leq \mu(A) \leq \delta$.*

Proof. (1) Let $B_1 \subset E$ be a set such that $0 < \mu(B_1) < \mu(E)$. Then either $\mu(B_1) \leq \mu(E)/2$ and we set $A_1 = B_1$ or $\mu(X \setminus B_1) \leq \mu(E)/2$, and we set $A_1 = X \setminus B_1$. Now repeat the process with A_1 instead of E , obtaining a measurable subset A_2 of A_1 , with $0 < \mu(A_2) < \mu(E)/4$. Continuing in this way we see that X contains subsets with arbitrary small measure.

(2) We prove the existence of a measurable set with the desired property and use the fact that μ is regular to imply this set can be open. Let \mathcal{C} be the collection of measurable subsets A of X for which $\mu(A) < \frac{1}{2}\delta$. If \mathcal{C} is not closed under unions then the lemma is proved. For example, if $A, B \in \mathcal{C}$ but $A \cup B \notin \mathcal{C}$ then $\frac{1}{2}\delta \leq \mu(A \cup B) \leq \delta$. Therefore \mathcal{C} is closed under binary unions. Taking limits, this implies \mathcal{C} is closed under countable unions.

Let $\beta = \sup_{C \in \mathcal{C}} \mu(C)$. There exists a sequence of sets $\{B_i\}$ for which $\mu(B_n) \nearrow \beta$. Let $B = \bigcup B_n$, this implies $\mu(B) = \beta$ and since $B \in \mathcal{C}$, we have $\beta < \frac{1}{2}\delta$. But then by (2) we can find a subset $E \subseteq X \setminus B$ whose measure is less than $\frac{1}{2}\delta - \beta$, which would imply $B \cup E \in \mathcal{C}$ contradicting the fact that B attains $\sup_{C \in \mathcal{C}} \mu(C)$ \square

Proposition 28 *Given non-atomic computable measurable space (\mathcal{X}, μ) , there is a $c \in \mathbb{N}$, where for all n , $\mu(X)2^{-n-\mathbf{K}(n)-c} < \mu(\{x : \mathbf{H}_\mu(x) < \log \mu(X) - n\}) < \mu(X)2^{-n}$.*

Proof. By Proposition 27, for every $\delta \in [0, \mu(X)]$, there exists an open set A , with $\frac{1}{2}\delta \leq \mu(A) \leq \delta$. Thus one can uniformly, in $n \in \mathbb{W}$, enumerate an effectively open sets $\{D_n\}$ such that $\mathbf{m}(n)\mu(X)2^{-n-1} < \mu(D_n) < \mathbf{m}(n)\mu(X)2^{-n}$ such that $D_n \cap D_m = \emptyset$ if $n \neq m$, where $\mu(X)$ is computable because μ is computable. The reasoning is as follows.

Let $\{\hat{D}_n\}$ be current ad-sets all originally \emptyset such that $\hat{\mathbf{m}}(n)2^{-n-1}\mu(X) < \mu(\hat{D}_n) < \hat{\mathbf{m}}(n)\mu(X)2^{-n}$, where $\hat{\mathbf{m}}$ is a lower approximation of \mathbf{m} . One can lower compute the interval

$$[\mathbf{m}(n)\mu(X)2^{-n-1}, \mathbf{m}(n)\mu(X)2^{-n}]$$

for all n , and if the interval shifts by some rational amount, by Proposition 27, one can add an ad-set $D \subseteq X \setminus \bigcup_{i=1}^{\infty} \widehat{D}_i$, such that $\widehat{\mathbf{m}}(n)\mu(X)2^{-n-1} < D \cup \widehat{D}_n < \widehat{\mathbf{m}}(n)\mu(X)2^{-n-1}$, and then set $\widehat{D}_n = D \cup \widehat{D}_n$, and continue with the enumeration.

Let the μ -test $t(\alpha) = \sup_{n:\alpha \in D_n} 2^{n-\log \mu(X)}$. Thus since t is lower computable and $\int_X t d\mu \leq \sum_n \mu(D_n)2^n/\mu(X) = \sum_n \mathbf{m}(n) < 1$, we have that $t <^* \mathbf{t}_\mu$. Since $\mu\{x : \mathbf{t}_\mu(x) > 2^n/\mu(X)\} < \mu(X)2^{-n}$, we get that there exists $c \in \mathbb{N}$, with $\mu(X)2^{-n-\mathbf{K}(n)-c} < \mu(\{x : \mathbf{H}_\mu(x) < \log \mu(X) - n\}) < \mu(X)2^{-n}$. \square

Corollary 20 *Given non-atomic computable measurable space (\mathcal{X}, μ) , there is a $c \in \mathbb{N}$, where for all n , $\mu(X)2^{-n-\mathbf{K}(n)-c} < \mu(\{x : \mathbf{t}_\mu(x) > 2^n/\mu(X)\}) < \mu(X)2^{-n}$.*

The following shows that during the course of ergodic dynamics, the state will be guaranteed to oscillate in its algorithmic fine grained thermodynamic entropy. Small oscillations are frequent, and larger fluctuations are more rare. This theorem parallels Theorem 53 in its inequalities.

Theorem 57 (Discrete Oscillations) *Let (\mathcal{X}, μ) be a non-atomic computable measure space. There is a $c \in \mathbb{N}$ with the following properties. Let $T : X \rightarrow X$ be a computable ergodic function, and $U_n = \{x : \mathbf{H}_\mu(x) < \log \mu(X) - n\}$. If $\omega \in X$ has $\mathbf{H}_\mu(\omega) \neq -\infty$,*

$$2^{-n-\mathbf{K}(n)-c} < \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{t=0}^{n-1} \mathbf{1}_{U_n}(T^t(\omega)) < 2^{-n}.$$

Proof. By Proposition 28, there is a c where $\mu(X)2^{-n-\mathbf{K}(n)-c} < \mu(U_n) < \mu(X)2^{-n}$. By Theorem 56, $\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} \mathbf{1}_{U_n}(T^k(\omega)) = \mu(U_n)/\mu(X)$. So

$$2^{-n-\mathbf{K}(n)-c} < \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} \mathbf{1}_{U_n}(T^k(\omega)) < 2^{-n}.$$

14.2.3 Lower Computable Functions

Theorem 58 *Let (\mathcal{X}, μ) be a computable measure space. Let $T : X \rightarrow X$ be a computable ergodic function. Let $f : X \rightarrow \overline{\mathbb{R}}^+ \cup \infty$ be lower computable. If $\mathbf{H}_\mu(\omega) \neq -\infty$,*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} f(T^k(\omega)) = \frac{1}{\mu(X)} \int f d\mu.$$

Proof. Let $\nu = \mu/\mu(X)$ be a computable measure over \mathcal{X} , due to Claim 1. Let f be a lower computable function with a finite integral. Let $f_n = \frac{1}{n}(f + \dots + f \cdot T^{n-1})$. Let $r > \int f d\nu$ be a rational number and $G_N = \{\omega : (\exists n \geq N) f_n(\omega) > r\}$. The set G_N is effectively open and $\nu(\bigcup_N G_N) = 0$ as $f_n(\omega) = \int f d\nu < r$ for ν -almost every ω by the classical version of Birkoff's ergodic theorem. As a result, there exists N where $\nu(G_N) < 1$. By Theorem 55, if $\mathbf{H}_\nu(\omega) \neq -\infty$, then there exists k such that $T^k(\omega) \notin G_N$. So $\limsup f_n(T^k(\omega)) \leq r$ and $\limsup f_n(\omega) = \limsup f_n(T^k(\omega)) \leq r$. Since $r > \int f d\nu$ can be arbitrarily close to the integral, we have that $\limsup f_n(\omega) <^+ \int f d\nu = \frac{1}{\mu(X)} \int f d\mu$.

It remains to prove that $\liminf f_n(\omega) \geq \int d\nu$. This is true for any lower semicontinuous f . Consider some lower bound for f that is of the form $\hat{f}(\omega) = \sum_{i=1}^n c_n c_n \mathbf{1}_{B_n}(\omega)$, where each B_n is an almost decidable ball. For these basic functions the statement of the theorem is true using the reasoning of Theorem 56, and their integrals can be arbitrarily close to $\int f\nu = \frac{1}{\mu(X)} \int f d\mu$. \square

Corollary 21 *Let (\mathcal{X}, μ) be a computable measure space and $T : X \rightarrow X$ be a computable ergodic function. For $\omega \in X$ with $\mathbf{H}_\mu(\omega) \neq -\infty$, then*

$$\lim_{n \rightarrow \infty} \mu(X) \sum_{t=0}^{n-1} 2^{-\mathbf{H}_\mu(T^t \omega)} < 1.$$

Proof. This follows from Theorem 58 and the fact that $2^{-\mathbf{H}_\mu(\omega)} = \mathbf{t}_\mu(\omega)$ is lower computable and $\int \mathbf{t} d\mu < 1$. \square

Chapter 15

Dynamics on Product Spaces

Theorem 53 proves the existence of oscillations in thermodynamic entropy. However the situation becomes more complicated if one to exam product spaces. If a product state is typical of the space then its marginal entropies cannot oscillate in synch. For every number there is a time where the marginal entropies differ by more than that amount. In this chapter, we prove this fact for two cases. The first case is using continuous dynamics and the second case is using discrete ergodic dynamics. In both such cases, marginal entropies cannot be synchronized.

15.1 Continuous Dynamics

15.1.1 Exotic Intervals

Definition 29 *An r -interval $v \subseteq (0, 1)$ is a finite collection of open intervals with rational endpoints.*

Lemma 20 *For r -interval v , Lebesgue measure L , there exists a rational number $r \in v$ such that $\mathbf{K}(r) <^{\log} -\log L(v) + \mathbf{I}(\langle r \rangle; \mathcal{H})$.*

Proof. Let $n = \lceil -\log L(v) \rceil$. We condition the universal Turing machine on n , which can be done given the precision of the theorem. Let Q be an elementary probability measure that realizes the stochasticity of v (see Chapter 1), and $d = \mathbf{d}(r|Q)$ and $\mathbf{Ks}(x) = \mathbf{K}(Q) + 3 \log \max\{\mathbf{d}(x|Q), 1\}$. Without loss of generality, the support of Q can be assumed to consist entirely of r -intervals s with $n = \lceil -\log L(s) \rceil$. We sample $d2^n$ real numbers R in the interval $(0, 1)$ using L . For r -interval s , $R \subset \mathbb{R}$, we define the indicator function $\mathbf{i}(s, R) = [R \cap s = \emptyset]$.

$$\mathbf{E}_{R \sim (0,1)^{d2^n}} \mathbf{E}_{s \sim Q} \mathbf{i}(s, R) = \mathbf{E}_{s \sim Q} \mathbf{E}_{R \sim (0,1)^{d2^n}} \mathbf{i}(s, R) = (1 - s)^{d2^n} \leq (1 - 2^{-n})^{d2^n} < e^{-d}.$$

Thus given Q and d , one can determine a set of $d2^n$ rationals $W \subset \mathbb{Q}$, such that $\mathbf{E}_{s \sim Q} \mathbf{i}(s, W) < e^{-d}$. Thus $t(s) = \mathbf{i}(s, W)e^d$ is a Q -test, with $\mathbf{E}_{s \sim Q} [t(s)] < 1$. It must be that $t(v) = 0$, otherwise

$$1.44d \leq \log t(v) <^+ \mathbf{d}(v|Q, d) <^+ d + \mathbf{K}(d),$$

which is a contradiction for large d which one can assume without loss of generality. Thus there exists a rational $r \in W \cap v$ such that, using Lemma 6,

$$\begin{aligned}
\mathbf{K}(r) &<^+ \log |W| + \mathbf{K}(W) \\
&<^+ n + \log d + \mathbf{K}(d, Q) \\
&<^+ n + 3 \log d + \mathbf{K}(Q) \\
&<^+ n + \mathbf{Ks}(v) \\
&<^+ n + \mathbf{I}(\langle v \rangle; \mathcal{H}).
\end{aligned}$$

□

15.1.2 No Synchronization Under Continuous Dynamics

Theorem 59 *Let $(\mathcal{X} \times \mathcal{Y}, \mu \times \nu)$ be a computable product measure space. Let G^t be a transformation group with $(\alpha^t, \beta^t) = G^t(\alpha, \beta)$. If $\mathbf{H}_\mu(\alpha, \beta) > -\infty$ and $\mathbf{I}((\alpha, \beta) : \mathcal{H}) < \infty$ then $\sup_{t \in [0,1]} |\mathbf{H}_\mu(\alpha^t) - \mathbf{H}_\nu(\beta^t)| = \infty$.*

Proof. Let $\bar{\mu} = \mu/\mu(X)$ and $\bar{\nu} = \nu/\nu(Y)$ be computable probability measures due to Claim 1. Assume not and $d = \lceil \sup_{t \in [0,1]} |\mathbf{H}_{\bar{\mu}}(\alpha^t) - \mathbf{H}_{\bar{\nu}}(\beta^t)| \rceil < \infty$. Let $U_n \subset [0, 1]$ be an open set where $U_n = \{t : \mathbf{H}_{\bar{\mu}}(\alpha^t) < -n\}$. By Corollary 18, there is a $c \in \mathbb{N}$ where for Lebesgue measure L , $2^{-n-2 \log n - c} < L(U_n)$.

Given (α, β) and n , one can enumerate an increasing r -interval $v \subseteq U_n$ and stop when $L(v) > L(U_n/2)$. By Lemma 20, there exists a rational $r \in v$, with

$$\mathbf{K}(r) <^{\log} n + \mathbf{I}(r; \mathcal{H}) <^{\log} n + \mathbf{I}((\alpha, \beta) : \mathcal{H}) + \mathbf{K}(n) <^{\log} n + \mathbf{I}((\alpha, \beta) : \mathcal{H}).$$

Let $A_n = \{\gamma : \mathbf{t}_{\bar{\mu}}(\gamma) > 2^n\}$ and $B_n = \{\gamma : \mathbf{t}_{\bar{\nu}}(\gamma) > 2^n\}$. By the definition of tests, $\mu \times \nu(A_n \times B_n) < 2^{-2n}$. This enables us to create the following $\bar{\mu} \times \bar{\nu}$, test

$$\begin{aligned}
t_n(\gamma, \lambda) &= [(\gamma, \lambda) \in G^{-r}(A_n \times B_{n-d})] 2^{2n-d} \\
\mathbf{K}(t_n) &<^+ \mathbf{K}(r, n) <^{\log} n + \mathbf{I}((\alpha, \beta) : \mathcal{H}) \\
&<^{\log} n,
\end{aligned} \tag{15.1}$$

where Equation 15.1 is due to the assumption that $\mathbf{I}((\alpha, \beta) : \mathcal{H}) < \infty$. It must be that $t_n(\alpha, \beta) \neq 0$ because $r \in U_n$. Thus

$$\begin{aligned}
\mathbf{t}_{\bar{\mu} \times \bar{\nu}}(\alpha, \beta) &>^* \sum_n \mathbf{m}(t_n) t_n(\alpha, \beta) \\
&=^* \sum_n 2^{-n-O(\log n)} 2^{2n-d} \\
&= \infty.
\end{aligned}$$

Thus $\mathbf{H}_{\bar{\mu} \times \bar{\nu}}(\alpha, \beta) = -\infty$, which means $\mathbf{H}_{\mu \times \nu}(\alpha, \beta) = -\infty$, causing a contradiction. □

15.2 Ergodic Dynamics

15.2.1 Effective Convergence Time of Ergodic Dynamics

To prove properties of ergodic dynamics, an effective means of determining the convergence of Birkoff averages was needed, and thus the results from [GHR10] were leveraged. Thus the contents in this subsection reviews this referenced work.

Definition 30

Let (\mathcal{X}, μ) be a computable measure space.

- Random variables f_n effectively converge in probability to f if for each $\epsilon > 0$, $\mu\{x : |f_n(x) - f(x)| < \epsilon\}$ converges effectively to 1, uniformly in ϵ . Thus there is a computable function $n(\epsilon, \delta)$ where for all $n \geq n(\epsilon, \delta)$, $\mu\{|f_n - f| \geq \epsilon\} < \delta$.
- Random variables f_n effectively converge almost surely to f if $f'_n = \sup_{k \geq n} |f_k - f|$ effectively converge in probability to 0.
- A simple function $f : X \rightarrow \mathbb{Q}$ is of the form $f(x) = \max_i 1^n q_i \mathbf{1}_{B_n}(x)$, where each q_i is a rational number and B_n is an almost decidable ball.
- The quotient space $L^1(X, \mu)$ can be made into computable metric space. Let $d_1(f, g) = \int_X |f - g| d\mu$ be the distance metric, with the relation $f \sim g$ if $d_1(f, g) = 0$. The ideal points are simple functions. An integral function $f : X \rightarrow \mathbb{R}_{>0}$ is $L^1(X, \mu)$ computable if it is a computable point of the computable metric space $L^1(X, \mu)$. Basic operations $L^1(X, \mu)$ such as addition, multiplication by a scalar, min, max, are computable. If f is computable, then so is $\int f d\mu$.

Proposition 29 If f is a computable element of computable metric space $L^1(X, \mu)$ and if $T : X \rightarrow X$ is computable and μ -measure preserving, then $f \circ T$ is computable.

Proof. Since f is computable, there is a computable fast Cauchy sequences of simple functions (f_1, f_2, \dots) converging to f . Given a simple function $g(x) = q \mathbf{1}_B(x)$, we will show how to compute a simple function h such that $d_1(h, g \circ T)$ can be arbitrarily small. The process can be easily generalized to arbitrary simple functions and thus to each f_i . Since B is almost-decidable $\mu(B)$ can be computed, and thus the effectively open set $T^{-1}(B)$ can be enumerated by ad-sets A such that $\mu(A)$ is arbitrarily close to $\mu(B)$. Thus one can get the simple function $\ell(x) = q \mathbf{1}_A(x)$ such that $d_1(\ell, g)$ is arbitrarily close. \square

Proposition 30 Let (\mathcal{X}, μ) be a computable metric space and $T : X \rightarrow X$ be a computable ergodic function. Let f be a computable element of $L^1(X, \mu)$. The L^1 convergence of Birkhoff averages of f is effective.

Proof. Replacing f with $f - \int f d\mu$, we can assume $\int f d\mu = 0$. Let $A_n = (f + f \circ T + \dots + f \circ T^{n-1})/n$. The sequence $\|A_n\|$ is computable and converges to 0 by Birkoff's Ergodic Theorem. Given $p \in \mathbb{N}$ we have $m \in \mathbb{N}$ where $m = np + k$, with $0 \leq k < p$. So

$$\begin{aligned} A_{np+k} &= \frac{1}{np+k} \left(\sum_{i=0}^{n-1} p A_p \circ T^{pi} + k A_k \circ T^{pn} \right) \\ \|A_{np+k}\| &= \frac{1}{np+k} (np \|A_p\| + k \|A_k\|) \\ &\leq \|A_p\| + \frac{\|A_k\|}{n} \\ &\leq \|A_p\| + \frac{\|f\|}{n} \end{aligned}$$

Let $\epsilon > 0$. We can compute some $p = p(\epsilon)$ such that $\|A_p\| < \epsilon/2$. Then we can compute some $n(\epsilon) \geq \frac{2}{\epsilon} \|f\|$. The function $m(\epsilon) = n(\epsilon)p(\epsilon)$ is computable and for all $m \geq m(\epsilon)$, $\|A_m\| \leq \epsilon$. \square

Lemma 21 (Maximal Ergodic Theorem) For $f \in L^1(X, \mu)$ and $\delta > 0$, $\mu\{x : \sup_n |A_n^f(x)| > \delta\} \leq \frac{1}{\delta} \|f\|_1$.

Theorem 60 Let (\mathcal{X}, μ) be a computable measure space and $T : X \rightarrow X$ be a computable ergodic function. If f is $L^1(X, \mu)$ -computable, then the Birkoff average effectively converge almost surely.

Proof. Let $\epsilon, \delta > 0$. Compute p such that $\|A_p^f\| \leq \delta\epsilon/2$. Applying the maximal ergodic theorem with $g = A_p^f$ has

$$\mu\{x : \sup_n |A_n^g(x)| > \delta/2\} \leq \epsilon. \quad (15.2)$$

One has that

$$A_n^g = A_n^f + \frac{u \circ T^n - u}{np},$$

where $u = (p-1)f + (p-2)f \circ T + \dots + f \circ T^{p-2}$. $\|u\|_\infty \leq \frac{p(p-1)}{2} \|f\|_\infty$ so if $n \geq n_0 \geq 4(p-1)\|f\|_\infty/\delta$ then $\|A_n^g - A_n^f\|_\infty \leq \delta/2$. So if $|A_n^f(x)| > \delta$ for some $n \geq n_0$, then $|A_n^g(x)| > \delta/2$. Using Equation 15.2, we get

$$\mu\{x : \sup_{n \geq n_0} |A_n^f(x)| > \delta\} \leq \epsilon.$$

As n_0 can be computed from δ and ϵ , we prove the theorem. \square

15.2.2 Busy Beaver Functions

We review the material on busy beaver functions, detailed in 13.2. Let $\Omega = \sum \{2^{-\|p\|} : U(p) \text{ halts}\}$ be Chaitin's Omega, $\Omega_n \in \mathbb{Q}_{\geq 0}$ be the rational formed from the first n bits of Ω , and $\Omega^t = \sum \{2^{-\|p\|} : U(p) \text{ halts in time } t\}$. For $n \in \mathbb{N}$, let $\mathbf{bb}(n) = \min\{t : \Omega_n < \Omega^t\}$. $\mathbf{bb}^{-1}(m) = \arg \min_n \{\mathbf{bb}(n-1) < m \leq \mathbf{bb}(n)\}$. Let $\Omega[n] \in \{0, 1\}^*$ be the first n bits of Ω .

Lemma. For $n = \mathbf{bb}^{-1}(m)$, $\mathbf{K}(\Omega[n]|m, n) = O(1)$.

15.2.3 Marginal Entropies of Ergodic Dynamics

Theorem 61 Let $(\mathcal{X} \times \mathcal{Y}, \mu \times \nu)$ be a non-atomic computable product measure space. Let G^t be an ergodic transformation group. Let $(\alpha, \beta) \in X \times Y$, with $(\alpha^t, \beta^t) = G^t(\alpha, \beta)$. If $\mathbf{H}_{\mu \times \nu}(\alpha, \beta) > -\infty$ and $\mathbf{I}((\alpha, \beta) : \mathcal{H}) < \infty$ then $\sup_{t \in \mathbb{N}} |\mathbf{H}_\mu(\alpha^t) - \mathbf{H}_\nu(\beta^t)| = \infty$.

Proof. Let $\bar{\mu} = \mu/\mu(X)$ and $\bar{\nu} = \nu/\nu(Y)$ be computable probability measures due to Claim 1. Assume not. Then there exists $c \in \mathbb{N}$, $c > \max_t |\mathbf{H}_{\bar{\mu}}(\alpha^t) - \mathbf{H}_{\bar{\nu}}(\beta^t)|$. Fix $n \in \mathbb{N}$. Let $U_n = \{(\gamma, \lambda) : \mathbf{H}_{\bar{\mu}}(\gamma) < -n\}$. By Proposition 28, there exists $d \in \mathbb{N}$ where $\bar{\mu}(U_n) > 2^{-n-2\log n-d}$. Given n , one can compute an ad-set $V_n \subset U_n$ with computable $p_n = -\log \mu(V_n)$ and $n+2\log n+d < p_n < n+2\log n+d+1$. Let $B_m^n = (\mathbf{1}_{V_n} + \mathbf{1}_{V_n} \circ T + \dots + \mathbf{1}_{V_n} \circ T^{m-1})/m$, which is computable, as T is measure preserving and due to Proposition 29. By Theorem 60, given $\delta, \epsilon > 0$, there is a computable $m(\delta, \epsilon, n)$ such that $\mu\{(\gamma, \lambda) : \sup_{m > m(\delta, \epsilon, n)} |B_m^n(\gamma, \lambda) - 2^{-p_n}| > \delta\} < \epsilon$. Let $m_n = m(2^{-p_n} - 2^{-1.5p_n}, 2^{-n}, n)$. Let $W_n = \{(\gamma, \lambda) : \sup_{m > m_n} |B_m^n(\gamma, \lambda) - 2^{-p_n}| > 2^{-p_n} - 2^{-1.5p_n}\}$. Either (1) there is an infinite number of n where $(\alpha, \beta) \in W_n$, or (2) there is an infinite number of n where $(\alpha, \beta) \notin W_n$.

Case (1). Each W_n is an effectively open set, computable uniformly in n . Furthermore, $\mu(W_n) < 2^{-n}$. Thus $t(\gamma, \lambda) = \sup_n [(\gamma, \lambda) \in W_n] \mathbf{m}(n) 2^n$ is a $\bar{\mu} \times \bar{\nu}$ test. So $\infty = t(\alpha, \beta) <^* \mathbf{t}_{\bar{\mu} \times \bar{\nu}}(\alpha, \beta)$, which implies $\mathbf{H}_{\mu \times \nu}(\alpha, \beta) = -\infty$, causing a contradiction.

Case (2). Fix one such $n \in \mathbb{N}$, where $(\alpha, \beta) \notin W_n$. Thus $\sup_{m > m_n} |2^{-p_n} - B_m^n(\alpha, \beta)| \leq 2^{-p_n} - 2^{-1.5p_n}$ implies $\sup_{m > m_n} B_m^n(\alpha, \beta) \geq 2^{-1.5p_n}$. Each $T^{-\ell}V_n$ is an effectively open set, uniformly in k and ℓ . So for all $m > m_n$, there are at least $2^{-1.5p_n}m$ indices ℓ , where $(\alpha, \beta) \in T^{-\ell}V_n$. Let $b_n = \mathbf{bb}^{-1}(m_n + 1)$ and N be the smallest power of 2 not less than $\mathbf{bb}(b_n)$. Thus, due to Lemma 16, $\mathbf{K}(N | (\alpha, \beta)) <^+ \mathbf{K}(n, b_n)$. Thus there are at least $2^{-1.5(n+2\log n+d+1)}N$ indices $\ell \in [1, \dots, N]$ where $(\alpha, \beta) \in T^{-\ell}V_n$. Let $D \subseteq \{0, 1\}^{\log N}$, where if $x \in D$ then $(\alpha, \beta) \in T^{-\text{Num}(x)}V_n$ and $|D| \geq 2^{-1.5(n+2\log n+d+1)}N$. The function $\text{Num} : \{0, 1\}^{\log N} \rightarrow \{1, 2, \dots, N\}$ converts strings to numbers in the natural way. Thus $\mathbf{K}(D | (\alpha, \beta)) <^+ \mathbf{K}(n, b_n)$. Let $\text{Uniform}(N)$ be the uniform measure over $\{0, 1\}^{\log N}$. By the EL Theorem (Corollary 41), applied to $\text{Uniform}(N) <^* \mathbf{m}/\mathbf{m}(N)$, and the definition of \mathbf{I} , there exists $x_n \in D$, with

$$\begin{aligned} \mathbf{K}(x_n) &<^{\log} \mathbf{K}(\text{Uniform}(N)) - \log |D| + \mathbf{I}(D; \mathcal{H}) \\ &<^{\log} \mathbf{K}(N) + 1.5n + 3\log n + \mathbf{I}((\alpha, \beta) : \mathcal{H}) + \mathbf{K}(n, b_n) \\ &<^{\log} \mathbf{K}(\Omega[b_n]) + 1.5n + \mathbf{I}((\alpha, \beta) : \mathcal{H}) + \mathbf{K}(b_n). \end{aligned} \quad (15.3)$$

Due to Lemma 16, $\mathbf{K}(\Omega[b_n] | (\alpha, \beta), n, b_n) = O(1)$. Furthermore, it is well known that for bits of Chaitin's Omega, $\mathbf{K}(\Omega[b_n] | \mathcal{H}) <^+ \mathbf{K}(b_n)$ and that $b_n <^+ \mathbf{K}(\Omega[b_n])$. So

$$b_n <^+ \mathbf{K}(\Omega[b_n]) <^{\log} \mathbf{I}(\Omega[b_n]; \mathcal{H}) <^{\log} \mathbf{I}((\alpha, \beta) : \mathcal{H}) + \mathbf{K}(b_n, n) <^{\log} \mathbf{I}((\alpha, \beta) : \mathcal{H}) + \mathbf{K}(n) \quad (15.4)$$

Combining Equations 15.3 and 15.4 together, we get

$$\mathbf{K}(x_n) <^{\log} 1.5n + 2\mathbf{I}((\alpha, \beta) : \mathcal{H}).$$

We define the test

$$\begin{aligned} t_{n,y}(\gamma, \lambda) &= \left[\mathbf{H}_{\bar{\mu}}(T^{\text{Num}(y)}(\gamma, \lambda)_1) < -n \text{ and } \mathbf{H}_{\bar{\nu}}(T^{\text{Num}(y)}(\gamma, \lambda)_2) < -n + c \right] 2^{2n-c}, \\ \mathbf{t}_{\bar{\mu} \times \bar{\nu}}(\gamma, \lambda) &>^* \sum_n \mathbf{m}(t_{n,x_n}) t_{n,x_n}(\gamma, \lambda), \\ &>^* \sum_n \left[\mathbf{H}_{\bar{\mu}}(T^{\text{Num}(x_n)}(\gamma, \lambda)_1) < -n \text{ and } \mathbf{H}_{\bar{\nu}}(T^{\text{Num}(x_n)}(\gamma, \lambda)_2) < -n + c \right] \frac{2^{.5n-2\mathbf{I}((\alpha, \beta) : \mathcal{H})}}{(n + \mathbf{I}((\alpha, \beta) : \mathcal{H}))^{O(1)}}. \end{aligned}$$

In recap, since $(\alpha, \beta) \notin W_n$, $|B_N^n(\alpha, \beta) - 2^{-p_n}| > 2^{-p_n} - 2^{-1.5p_n}$, so $B_N^n > 2^{-1.5p_n} > 2^{-1.5(n+2\log n+d)}$. Thus one can create a large enough set $D \subset \{0, 1\}^N$, and find a simple enough $x_n \in D$ such that $(\alpha, \beta) \in T^{-\text{Num}(x_n)}V_n$. By assumption of the theorem

$$\mathbf{H}_{\bar{\mu}}(T^{\text{Num}(x_n)}(\alpha, \beta)_1) < -n \text{ and } \mathbf{H}_{\bar{\nu}}(T^{\text{Num}(x_n)}(\alpha, \beta)_2) < -n + c.$$

Thus $\mathbf{m}(t_{n,x_n}) t_{n,x_n}(\alpha, \beta) = \frac{2^{.5n-2\mathbf{I}((\alpha, \beta) : \mathcal{H})}}{(n + \mathbf{I}((\alpha, \beta) : \mathcal{H}))^{O(1)}}$. Furthermore, since $\mathbf{I}(\alpha, \beta) < \infty$ and there is an infinite number of n where $(\alpha, \beta) \notin W_n$, $\mathbf{t}_{\bar{\mu} \times \bar{\nu}}(\alpha, \beta) = \infty$, so $\mathbf{H}_{\mu \times \nu}(\alpha, \beta) = -\infty$, causing a contradiction. \square

Corollary 22 (Independent Systems) *Let $(\mathcal{X} \times \mathcal{Y}, \mu \times \nu)$ be a non-atomic computable product measure space. Let G_X^t and G_Y^t be ergodic transformation groups over \mathcal{X} and \mathcal{Y} respectively. Let $(\alpha, \beta) \in X \times Y$. If $\mathbf{H}_{\mu \times \nu}(\alpha, \beta) > -\infty$ and $\mathbf{I}((\alpha, \beta) : \mathcal{H}) < \infty$ then $\sup_{t \in \mathbb{N}} |\mathbf{H}_\mu(G_X^t(\alpha)) - \mathbf{H}_\nu(G_Y^t(\beta))| = \infty$.*

The above theorem has immediate applications to the Cantor space. Let σ be the shift operator and $(\alpha, \beta) = \alpha[1]\beta[1]\alpha[2]\beta[2]\dots$. Let λ be the uniform distribution over $\{0, 1\}^\infty$.

Corollary 23 *If (α, β) is ML Random and $\mathbf{I}((\alpha, \beta) : \mathcal{H}) < \infty$ then $\sup_n |\mathbf{D}(\sigma^{(n)}\alpha|\lambda) - \mathbf{D}(\sigma^{(n)}\beta|\lambda)| = \infty$.*

Chapter 16

Algorithmic Coarse Grained Entropy

16.1 Fundamentals of Coarse Grained Entropy

Coarse grained entropy was introduced in [Gac94] as an update to Boltzmann entropy. The goal was a parameter independent formulation of entropy. It was defined using cells. In this section we define coarse grained entropy with respect to open sets, leveraging Chapter 11. Let $\Pi(\cdot)$ be a set of disjoint uniformly enumerable open sets in the computable metric space \mathcal{X} , as shown in Figure 16.1.

Definition 31 (Algorithmic Coarse Grained Entropy) $\mathbf{H}_\mu(\Pi_i) = \mathbf{K}(i|\mu) + \log \mu(\Pi_i)$.

Remark 2 (Paradox of Typicality) *Discrete dynamics are subject to a paradox of typicality, which is due to a remark by Vladimir Vovk and detailed in [Gac94]. This remark generalizes Example 7. Lets say P is a partition and state ω is a typical state in partition Γ , which has low entropy, thus $\mathbf{H}_\mu(\Gamma)$ is low. Thus $\mathbf{H}_\mu(\alpha) \approx \mathbf{H}(\Gamma)$. Say ω is subject to discrete time dynamics, for some simple $t \in \mathbb{N}$. This new state $\omega' = G^t\omega$ is in a partition Γ' of much greater entropy. Since t is simple, by Proposition 17, $\mathbf{H}_\mu(\omega')$ is still low. However the coarse grain entropy of ω' is $\mathbf{H}_\mu(\Gamma')$ which is high. So the locally typical state ω turns into locally non-typical state ω' . Assuming only the macroscopic variables of ω' are detectable, its low algorithmic fine grain entropy and history is inaccessible to observers.*

Note that this violation of typicality is in effect for a single, simple transform. For continuous transform groups or the averages ergodic discrete ones, this phenomenon is not guaranteed to occur.

Coarse grained entropy is an excellent approximation of fine grained entropy, as shown by Proposition 31 and Lemma 23.

Proposition 31 *Let (\mathcal{X}, μ) be a computable measure space. If $\mu(\Pi_i)$ is uniformly computable and $\alpha \in \Pi_i$ then $\mathbf{H}_\mu(\alpha) <^+ \mathbf{H}_\mu(\Pi_i) + \mathbf{K}(\Pi)$.*

Proof. Let $t(\alpha) = \sum_i [\alpha \in \Pi_i] \mathbf{m}(\Pi_i) / \mu(\Pi_i)$. $t \in \mathcal{F}$ is lower semi-computable and $\int_{\mathcal{X}} t(\alpha) d\mu(\alpha) = \sum_i \int_{\Pi_i} (\mathbf{m}(\Pi_i) / \mu(\Pi_i)) d\mu(\alpha) = \sum_i \mathbf{m}(\Pi_i) \leq 1$. Thus $\mathbf{t}_\mu(\alpha) \stackrel{*}{>} t(\alpha)$.

Example 7 *We detail changes in fine and coarse grained entropy during the course of dynamics, adapted from Example 6.7 from [Gac94], and described in Figure 16.2. Let $(\{0,1\}^{\mathbb{Z}}, \mu)$ be a computable measure space consisting of all bi-infinite sequences. The distance metric between*



Figure 16.1: Partition of phase space into macro sets Γ_ν , with the thermal equilibrium set Γ_{Eq} taking up most of the volume.



Figure 16.2: The following figure shows an example of changes in coarse grained entropy and fine grained entropy. On the first bi-infinite sequence, there are zeros between the position $-n/2$ and $n/2$ are 0, and beyond that are random. Thus both the coarse grained entropy using cells of length n and the fine grained entropy are zero. After dynamics causing the sequence to shift by n positions, the coarse grained entropy is high but the fine grained entropy is still low.

two points $\alpha, \beta \in \{0, 1\}^{\mathbb{Z}}$, is $d(\alpha, \beta) = \exp(-\arg \max_n \alpha[-n/2, n/2] = \beta[-n/2, n/2])$. We use the uniform distribution, where the measure of a cylinder $C = \{0, 1\}^{\mathbb{N}} x \{0, 1\}^{\mathbb{N}}$ is $\mu(C) = 2^{-\|x\|}$. The dynamics is the shift operation, a.k.a. Baker's map. We use the partitions Γ_x , over $x \in \{0, 1\}^n$, where $\omega \in \Gamma_x$ if $\omega[n-2, n+2] = x$. Thus for all $x \in \{0, 1\}^n$, $\mu(\Gamma_x) = 2^{-n}$. Let $\omega \in \{0, 1\}^{\mathbb{Z}}$, where $\omega^n = 0^n$, and all other bits are ML random. Thus the coarse grained entropy of ω is equal to

$$\mathbf{H}_\mu(\Gamma_{0^n}) = {}^+ \mathbf{K}(0^n) + \log \mu(\Gamma_{0^n}) = {}^+ \mathbf{K}(n) - n.$$

Since $\mathbf{K}(\mu(\Gamma_x)) = {}^+ \mathbf{K}(n)$, due to Proposition 31,

$$\mathbf{H}_\mu(\omega) < {}^+ 2\mathbf{K}(n) - n.$$

Now suppose we subject ω to dynamics, by applying the shift operator n times to produce ω' . Thus $\omega' \in \Gamma_x$, for some random string $x \in \{0, 1\}^n$. Thus ω' has algorithmic coarse grained entropy,

$$\mathbf{H}(\Gamma_x) = {}^+ \mathbf{K}(x) + \log \mu(\Gamma_x) = {}^+ 0.$$

However, while the coarse entropy has changed dramatically for ω' , its fine grained entropy has only gradually shifted. Due to Proposition 17,

$$\mathbf{H}_\mu(\omega') < {}^+ \mathbf{K}(n) + \mathbf{H}_\mu(\omega) < {}^+ 3\mathbf{K}(n) - n.$$

Lemma 22 For computable measure space (\mathcal{X}, μ) , for lower computable function $f \in \mathcal{F}$, and enumerable open set U , $\int_U f d\mu$ is lower computable.

Proof. For a finite union of balls $V = \bigcup_{j=1}^n B_{i_j}$ and an enumerable open set $W = \bigcup_{j=1}^\infty B_{k_j}$ and a computable measure μ , the term $\mu(V \cap W)$ is lower computable. Due to Proposition 10, the term $\mu(\bigcup\{B : \exists_{s,t} \text{ such that } B \subseteq B_{i_s} \text{ and } B \subseteq B_{k_t}\}) = \mu(V \cap W)$ is lower computable.

The integral of a finite supremum of step functions over U is lower computable by induction. For the base case $\int_U f_{i,j} d\mu = q_j \mu(B_i \cap U)$ is lower computable by the above reasoning. For the inductive step

$$\int_U \sup\{f_{i_1, j_1}, \dots, f_{i_k, j_k}\} d\mu = q_{j_m} \mu((B_{i_1} \cup \dots \cup B_{i_k}) \cap U) + \int_U \sup\{f_{i_1, j'_1}, \dots, f_{i_k, j'_k}\} d\mu,$$

where q_{j_m} is minimal among $\{q_{j_1}, \dots, q_{j_k}\}$ and $q_{j'_1} = q_{j_1} - q_{i_m}, \dots, q_{j'_k} = q_{j_k} - q_{i_k}$. The first term on the right is lower-computable and by the induction assumption, the last term on the right is lower-computable. \square

The following lemma is an update to the Stability Theorem 5 in [Gac94], using open sets instead of cells.

Lemma 23 For computable measure space (\mathcal{X}, μ) , $\mu\{\alpha \in \Pi_i : \mathbf{H}_\mu(\alpha) < \mathbf{H}_\mu(\Pi_i) - \mathbf{K}(\Pi) - m\} < {}^* 2^{-m} \mu(\Pi_i)$.

Proof. Let $f(i) = \int_{\Pi_i} \mathbf{t}_\mu(\alpha) d\mu(\alpha)$. By Lemma 22, the function $f(i)$ is lower computable, and $\sum_i f(i) \leq 1$. Thus $f(i) < {}^* \mathbf{m}(i)/\mathbf{m}(\Pi)$. So

$$\mu(\Pi_i)^{-1} \int_{\Pi_i} 2^{-\mathbf{H}_\mu(\alpha)} d\mu(\alpha) < {}^* 2^{-\mathbf{H}_\mu(\Pi_i) + \mathbf{K}(\Pi)}.$$

By Markov inequality,

$$\mu\{\alpha \in \Pi_i : \mathbf{H}_\mu(\alpha) < \mathbf{H}_\mu(\Pi_i) - \mathbf{K}(\Pi) - m\} \stackrel{*}{<} 2^{-m} \mu(\Pi_i).$$

Corollary 24 For computable measure space (\mathcal{X}, μ) , $\mu\{\alpha : \mathbf{H}_\mu(\alpha) < \log \mu(\mathcal{X}) - m\} \stackrel{*}{<} 2^{-m} \mu(\mathcal{X})$.

Theorem 62 For non-atomic computable measure space (\mathcal{X}, μ) , with uniformly computable $\mu(\Pi_i)$, $\mu\{\alpha \in \Pi_i : \mathbf{H}_\mu(\alpha) < \mathbf{H}_\mu(\Pi_i) + \mathbf{K}(\Pi) - m\} \stackrel{*}{>} \mu(\Pi_i) 2^{-m}$.

Proof. Similar to that of the proof of Proposition 28, one can compute, uniformly in m , an ad-set $D_m \subset \Pi_i$ such that $\mu(\Pi_i) 2^{-m-2} < \mu(D_m) < \mu(\Pi_i) 2^{-m-1}$ and $D_n \cap D_m = \emptyset$ for $n \neq m$. Suppose this has been done for D_{m-1} . The open set $\Pi_i \setminus \overline{\overline{D_{m-1}}}$ can be enumerated and $\mu(\Pi_i \setminus \overline{\overline{D_{m-1}}})$ can be lower computed. The enumeration of $\Pi_i \setminus \overline{\overline{D_{m-1}}}$ stops when $\mu(\Pi_i \setminus \overline{\overline{D_{m-1}}}) > \mu(\Pi_i) 2^{-m-1}$. Using Proposition 27, one can find an ad-set (see Definition 21) $D_m \subset \mu(\Pi_i \setminus \overline{\overline{D_{m-1}}})$, such that $\mu(\Pi_i) 2^{-m-2} < \mu(D_m) < \mu(\Pi_i) 2^{-m-1}$. We define μ -test t to be $t(\alpha) = \sum_m [\alpha \in D_m] \mathbf{m}(m) 2^m / \mu(\Pi_i)$. Since $\mathbf{m}(i) \mathbf{m}(\Pi) t \stackrel{*}{<} \mathbf{t}_\mu$, for universal lower computable test \mathbf{t}_μ and $\mathbf{H}_\mu = -\log \mathbf{t}_\mu$,

$$\begin{aligned} \mu(\Pi_i) 2^{-m} &< \mu\{\alpha : \mathbf{H}_\mu(\alpha) < \mathbf{K}(i) + \mathbf{K}(\Pi) + \log \mu(\Pi_i) - m\} \\ \mu(\Pi_i) 2^{-m} &< \mu\{\alpha : \mathbf{H}_\mu(\alpha) < \mathbf{H}_\mu(\Pi_i) + \mathbf{K}(\Pi) - m\}. \end{aligned}$$

□

The following Theorem is the coarse grained entropy to the oscillation Theorem 53. As a state travels through different cells Π_i and Π_j , the coarse grained entropy will oscillate, in that its max value $\mathbf{H}_\mu(\Pi_{i_{\max}})$ will become increasingly larger than its min value, $\mathbf{H}_\mu(\Pi_{i_{\min}})$. If the dynamics are ergodic, then by Theorem 55, the state is guaranteed to hit every cell if there are a finite number of them, and if there are an infinite number of cells, the state is guaranteed to hit an unbounded number of them.

Theorem 63 Let (\mathcal{X}, μ) be a computable measure space, G^t be a transformation group, and $\{\Pi_i\}$ a partition of \mathcal{X} . If $i \mapsto \mu(\pi_i)$ is uniformly computable and if a state $\alpha \in \mathcal{X}$, travels through at least 2^n partitions $\{\Pi_i\}_{i=1}^{2^n}$ over $t \in [0, 1]$, then

$$\min_{i \in \{1, \dots, 2^n\}} \mathbf{H}_\mu(\pi_i) <^{\log} \max_{i \in \{1, \dots, 2^n\}} \mathbf{H}_\mu(\pi_i) - n + \mathbf{I}(\alpha : \mathcal{H}).$$

Proof. Let $f(i) = \lceil \log \mu(i) \rceil$. Let $D \subset \mathbb{N}$, $|D| = 2^n$ be a set of partitions that α travels through, so $\mathbf{K}(D|\alpha) <^+ \mathbf{K}(n)$. Theorem 81, on $f : D \rightarrow \mathbb{N}$, produces $x \in D$, where

$$\begin{aligned} f(x) + \mathbf{K}(x) &<^{\log} -\log \sum_{a \in D} \mathbf{m}(a) 2^{-f(a)} + \mathbf{I}(f, D; \mathcal{H}) \\ f(x) + \mathbf{K}(x) + n &<^{\log} \max_{a \in D} f(a) + \mathbf{K}(a) + \mathbf{I}(f, D; \mathcal{H}) \\ \mathbf{H}_\mu(x) + n &<^{\log} \max_{a \in D} \mathbf{H}_\mu(a) + \mathbf{I}(\alpha : \mathcal{H}). \end{aligned}$$

16.2 Entropy Variance in Cells

In this section, we will prove a lower bound of states with low algorithmic fine grained entropy with respect to the uniform distribution over a particular cell, a lower bound. Thus, regardless of what measures are used to define entropy, low entropy states are bound to occur in cells. A result is first proved about infinite sequences, and measures over the Cantor Space, relying on Lemma 13.2.

Theorem 64 *For computable probability measures μ, ρ and nonatomic λ over $\{0, 1\}^\infty$ and $n \in \mathbb{N}$, $\lambda\{\alpha : \mathbf{D}(\alpha|\mu) > n \text{ and } \mathbf{D}(\alpha|\rho) > n\} > 2^{-n-\mathbf{K}(n,\mu,\rho,\lambda)-O(1)}$.*

Proof. We first assume not. For all $c \in \mathbb{N}$, there exist computable nonatomic measures μ, ρ, λ , and there exists n , where $\lambda\{\alpha : \mathbf{D}(\alpha|\mu) > n \text{ and } \mathbf{D}(\alpha|\rho) > n\} \leq 2^{-n-\mathbf{K}(n,\mu,\lambda)-c}$. Sample $2^{n+\mathbf{K}(n,\mu,\rho,\lambda)+c-1}$ elements $D \subset \{0, 1\}^\infty$ according to λ . The probability that all samples $\beta \in D$ has $\mathbf{D}(\beta|\mu) \leq n$ or $\mathbf{D}(\beta|\rho) \leq n$ is

$$\begin{aligned} \prod_{\beta \in D} \lambda\{\mathbf{D}(\beta|\mu) \leq n \text{ or } \mathbf{D}(\beta|\rho) \leq n\} &\geq \\ (1 - |D|2^{-n-\mathbf{K}(n,\mu,\lambda,\rho)-c}) &\geq \\ (1 - 2^{n+\mathbf{K}(n,\mu,\lambda,\rho)+c-1}2^{-n-\mathbf{K}(n,\mu,\lambda,\rho)-c}) &\geq 1/2. \end{aligned}$$

Let $\lambda^{n,c}$ be the probability of an encoding of $2^{n+\mathbf{K}(n,\mu,\lambda)+c-1}$ elements each distributed according to λ . Thus

$$\lambda^{n,c}(\text{Encoding of } 2^{n+\mathbf{K}(n,\mu,\lambda,\rho)+c-1} \text{ elements } \beta, \text{ each having } \mathbf{D}(\beta|\mu) \leq n \text{ or } \mathbf{D}(\beta|\rho) \leq n) \geq 1/2.$$

Let v be a shortest program to compute $\langle n, \mu, \rho, \lambda \rangle$. By Theorem 52, with the universal Turing machine relativized to v ,

$$\lambda^{n,c}(\{\gamma : \mathbf{I}(\gamma : \mathcal{H}|v) > m\}) \stackrel{*}{<} 2^{-m+\mathbf{K}(\lambda^{n,c}|v)} \stackrel{*}{<} 2^{-m+\mathbf{K}(n,\mathbf{K}(n,\mu,\lambda,\rho),c,\lambda|v)} \stackrel{*}{<} 2^{-m+\mathbf{K}(c)}.$$

Therefore,

$$\lambda^{n,c}(\{\gamma : \mathbf{I}(\gamma : \mathcal{H}|v) > \mathbf{K}(c) + O(1)\}) \leq 1/4.$$

Thus, by probabilistic arguments, there exists $\alpha \in \{0, 1\}^\infty$, such that $\alpha = \langle D \rangle$ is an encoding of $2^{n+\mathbf{K}(n,\mu,\rho,\lambda)+c-1}$ elements $\beta \in D \subset \{0, 1\}^\infty$, where each β has $\mathbf{D}(\beta|\mu) \leq n$ or $\mathbf{D}(\beta|\rho) \leq n$ and $\mathbf{I}(\alpha : \mathcal{H}|v) <^+ \mathbf{K}(c)$. By Lemma 17, relativized to v , there are constants $d, f, g \in \mathbb{N}$ where

$$\begin{aligned} m = \log |D| &< \max_{\beta \in D} \min\{\mathbf{D}(\beta|\mu, v), \mathbf{D}(\beta|\rho, v)\} + 2\mathbf{I}(D : \mathcal{H}|v) + d\mathbf{K}(m|v) + f\mathbf{K}(\mu|v) + g\mathbf{K}(\rho|v) \\ m &< \max_{\beta \in D} \min\{\mathbf{D}(\beta|\mu), \mathbf{D}(\beta|\rho)\} + \mathbf{K}(v) + 2\mathbf{I}(D : \mathcal{H}|v) + d\mathbf{K}(m|v) + f\mathbf{K}(\mu|v) + g\mathbf{K}(\rho|v) \\ &<^+ n + \mathbf{K}(n, \mu, \lambda, \rho) + d\mathbf{K}(m|v) + 2\mathbf{K}(c) + (f + g)O(1). \end{aligned} \tag{16.1}$$

Therefore:

$$\begin{aligned} m &= n + \mathbf{K}(n, \mu, \rho, \lambda) + c - 1 \\ \mathbf{K}(m|v) &<^+ \mathbf{K}(c). \end{aligned}$$

Plugging the inequality for $\mathbf{K}(m|v)$ back into Equation 16.1 results in

$$\begin{aligned} n + \mathbf{K}(n, \mu, \lambda, \rho) + c &<^+ n + \mathbf{K}(n, \mu, \lambda, \rho) + 2\mathbf{K}(c) + d\mathbf{K}(c) + (f + g)O(1) \\ c &<^+ (2 + d)\mathbf{K}(c) + (f + g)O(1). \end{aligned}$$

This result is a contradiction for sufficiently large c solely dependent on the universal Turing machine. \square

Theorem 65 *Given computable measures μ, ρ , and computable non-atomic probability measure λ , over a computable metric space \mathcal{X} , there is a $c \in \mathbb{N}$, where for all n , $\lambda(\{\alpha : \mathbf{H}_\mu(\alpha) < \log \mu(X) - n \text{ and } \mathbf{H}_\rho(\alpha) < \log \rho(X) - n\}) > 2^{-n-\mathbf{K}(n)-c}$.*

Proof. We fix the algorithmic descriptions of λ, μ, ρ , and \mathcal{X} . By Theorem 48, easily generalized to three measures, fix a multi representation $(\delta, \lambda_\delta, \mu_\delta, \rho_\delta)$ for computable multi measure space $(\mathcal{X}, \lambda, \mu, \rho)$. Note that δ is a measure-preserving transform, where $\lambda(A) = \lambda_\delta(\delta^{-1}(A))$ for all Borel sets A . Due to Lemma 13, there is a $c \in \mathbb{N}$,

$$\begin{aligned} &\lambda(\{\beta : \mathbf{H}_\mu(\beta) > \log \mu(X) - n \text{ or } \mathbf{H}_\rho(\beta) \geq \log \rho(X) - n\}) \\ &= \lambda(\{\beta : \mathbf{t}_\mu(\beta) \leq 2^n/\mu(X) \text{ or } \mathbf{t}_\rho(\beta) \leq 2^n/\rho(X)\}) \\ &= \lambda_\delta(\delta^{-1}(\{\beta : \mathbf{t}_\mu(\beta) \leq 2^n/\mu(X) \text{ or } \mathbf{t}_\rho(\beta) \leq 2^n/\rho(X)\})) \\ &< \lambda_\delta(\delta^{-1}(\{\beta : \mathbf{t}_{\mu_\delta}(\delta^{-1}(\beta)) < 2^{n+c}/\mu(X) \text{ or } \mathbf{t}_{\rho_\delta}(\delta^{-1}(\beta)) < 2^{n+c}/\rho(X)\})) \\ &= \lambda_\delta(\delta^{-1}(\{\beta \in \delta(\{\alpha : \mathbf{t}_{\mu_\delta}(\alpha) < 2^{n+c}/\mu(X) \text{ or } \mathbf{t}_{\rho_\delta}(\alpha) < 2^{n+c}/\rho(X)\})\})) \\ &= \lambda_\delta(\delta^{-1}(\delta(\{\alpha : \mathbf{t}_{\mu_\delta}(\alpha) < 2^{n+c}/\mu(X) \text{ or } \mathbf{t}_{\rho_\delta}(\alpha) < 2^{n+c}/\rho(X)\}))) \\ &< \lambda_\delta(\{\alpha : \mathbf{t}_{\mu_\delta}(\alpha) < 2^{n+c}/\mu(X) \text{ or } \mathbf{t}_{\rho_\delta}(\alpha) < 2^{n+c}/\rho(X)\}). \end{aligned} \tag{16.2}$$

Let $\bar{\mu}_\delta = \mu_\delta/\mu(X)$ and $\bar{\rho}_\delta = \rho_\delta/\rho(X)$ be two computable probability measures over $\{0, 1\}^\infty$. From Equation 16.3, and Proposition 14, we get,

$$\begin{aligned} X &< \lambda_\delta\{\alpha : \mathbf{D}(\alpha|\mu_\delta) <^+ n + c - \log \mu(X) \text{ or } \mathbf{D}(\alpha|\rho_\delta) <^+ n + c - \log \rho(X)\} \\ &= \lambda_\delta\{\alpha : \mathbf{D}(\alpha|\bar{\mu}_\delta) <^+ n + c \text{ or } \mathbf{D}(\alpha|\bar{\rho}_\delta) <^+ n + c\}. \end{aligned} \tag{16.4}$$

From Theorem 64, we get

$$\begin{aligned} &\lambda(\{\beta : \mathbf{H}_\mu(\beta) > \log \mu(X) - n \text{ or } \mathbf{H}_\rho(\beta) \geq \log \rho(X) - n\}) \\ &< \lambda_\delta\{\alpha : \mathbf{D}(\alpha|\bar{\mu}_\delta) <^+ n + c \text{ or } \mathbf{D}(\alpha|\bar{\rho}_\delta) <^+ n + c\} \\ &< 1 - 2^{-n-c}. \end{aligned}$$

\square

Corollary 25 *Given computable measure μ and computable non-atomic probability measure λ , over a computable metric space \mathcal{X} , there is a $c \in \mathbb{N}$, where for all n , $\lambda(\{\alpha : \mathbf{t}_\mu(\alpha) > \mu(X)2^{-n}\}) > 2^{-n-\mathbf{K}(n)-c}$.*

The following corollary states, given a uniform measure over a cell, there is a lower bound over the measure of the states with low algorithmic entropy with respect to two measures. Note that the corollary can be easily generalized to any finite number of measures.

Corollary 26 *For dual computable measure space (\mathcal{X}, μ, ν) , and with partition Π , let λ be a computable uniform probability measure over cell Π_i . There is a constant $c \in \mathbb{N}$ where $\lambda(\{\alpha : \mathbf{H}_\mu(\alpha) < \log \mu(X) - n \text{ and } \mathbf{H}_\nu(\alpha) < \log \nu(X) - n\}) > 2^{-n-\mathbf{K}(n)-c}$.*

Proof. This follows directly from Theorem 65.

Chapter 17

Computability of Algorithmic Entropy

In this chapter, computability properties of algorithmic entropy are studied. It is easy to see that algorithmic fine grained entropy is uncomputable. In Section 17.1, it is shown that there is no effective method for choosing times where the entropy of a typical state is continually decreasing. In Section 17.2, it is shown if a function can approximate algorithmic fine grained entropy for an infinite amount of typical states, then it has infinite mutual information with the halting sequence.

17.1 Uncomputability of Decreasing Entropy

If a physicist were given the state of a system, (say particles in a box), she could compute times of decreasing entropy (just wait until the particles are in smaller and smaller spaces). However without access to the state, there are no computable method to predict the times in which the state continually decreases.

Theorem 66 *Let (\mathcal{X}, μ) be a computable measure space and $T : X \rightarrow X$ be a computable ergodic function. If $\alpha \in X$, and $\mathbf{H}_\mu(\alpha) > -\infty$, then there does not exist a computable function $f : \mathbb{N} \rightarrow \mathbb{N}$, such that $\mathbf{H}_\mu(T^{f(n)}\alpha) < -n$.*

Proof. Assume not, and such a computable function f exists. Let $U_n = \{\beta : \mathbf{H}_\mu(\beta) < -n\}$, which is an effectively open set, uniformly in n . One can construct a μ -test, where $t(\beta) = \sup_n [\beta \in T^{-f(n)}U_n] \mathbf{m}(n)2^n$. Thus $\infty = t(\alpha) < \mathbf{t}_\mu(\alpha)$, and thus $\mathbf{H}_\mu(\alpha) = -\infty$, causing a contradiction. \square

Corollary 27 *Let (\mathcal{X}, μ) be a computable measure space and G^t and transformation group. If $\alpha \in X$, and $\mathbf{H}_\mu(\alpha) > -\infty$, then there does not exist a computable function $f : \mathbb{N} \rightarrow \mathbb{R}_{>0}$, such that $\mathbf{H}_\mu(T^{f(n)}\alpha) < -n$.*

Proof. The proof follows analogously to that of Theorem 66. \square

17.2 Non-Approximability of Algorithmic Entropy

17.2.1 Kolmogorov Complexity is Exotic

We review the material on busy beaver functions, detailed in 13.2. Let $\Omega = \sum \{2^{-\|p\|} : U(p) \text{ halts}\}$ be Chaitin's Omega, $\Omega_n \in \mathbb{Q}_{\geq 0}$ be the rational formed from the first n bits of Ω , and $\Omega^t = \sum \{2^{-\|p\|} : U(p) \text{ halts in time } t\}$. For $n \in \mathbb{N}$, let $\mathbf{bb}(n) = \min\{t : \Omega_n < \Omega^t\}$. $\mathbf{bb}^{-1}(m) = \arg \min_n \{\mathbf{bb}(n-1) < m \leq \mathbf{bb}(n)\}$. Let $\Omega[n] \in \{0, 1\}^*$ be the first n bits of Ω . For $t \in \mathbb{N}$

define the function $\mathbf{m}^t(x) = \sum \{2^{-\|p\|} : U(p) = x \text{ in } t \text{ steps}\}$ and for $n \in \mathbb{N}$, we have $\mathbf{m}_n(x) = \sum \{2^{-\|p\|} : U(p) = x \text{ in } \mathbf{bb}(n) \text{ steps}\}$.

The following lemma can be strengthened to just $\mathbf{I}(X; \mathcal{H})$, but with an increase to the complexity of the proof.

Lemma 24 *A relation $X = \{(x_i, c_i)\}_{i=1}^{2^n} \subset \{0, 1\}^* \times \mathbb{N}$, $|\mathbf{K}(x_i) - c_i| \leq s$, has $n <^{\log} 2s + 2\mathbf{I}(X; \mathcal{H})$.*

Proof. We relativize the universal Turing machine to (n, s) , which can be done due to the precision of the theorem. Let $T = \min\{t : \lceil -\log \mathbf{m}_t(x_i) \rceil - c_i < s + 1\}$. Let $N = \mathbf{bb}^{-1}(T)$ and $M = \mathbf{bb}(N)$. So for all x_i , $-\log \mathbf{m}_M(x_i) - \mathbf{K}(x_i) <^+ 2s$. Let Q be an elementary probability measure that realizes $\mathbf{Ks}(X)$ and $d = \max\{\mathbf{d}(X|Q), 1\}$. Without loss of generality, the support of Q is restricted to binary relations $B \subset \{0, 1\}^* \times \mathbb{N}$ of size 2^n . Let $B_1 = \bigcup \{y : (y, c) \in B\}$. Let $S = \bigcup \{B_1 : B \in \text{Support}(Q)\}$. We randomly select each string in S to be in a set R independently with probability $d2^{-n}$. Thus $\mathbf{E}[\mathbf{m}_M(R)] \leq d2^{-n}$. For $B \in \text{Support}(Q)$,

$$\mathbf{E}_R \mathbf{E}_{B \sim Q} [[R \cap B_1 = \emptyset]] = \mathbf{E}_{B \sim Q} \Pr(R \cap B_1 = \emptyset) = (1 - d2^{-n})^{2^n} < e^{-d}.$$

Thus there exists a set $R \subseteq S$ such that $\mathbf{m}_M(R) \leq 2 \cdot 2^{-n}$ and $\mathbf{E}_{B \sim Q} [[R \cap B_1 = \emptyset]] < 2e^{-d}$. Let $t(B) = .5[R \cap B_1 = \emptyset]2^d$. t is a Q -test, with $\mathbf{E}_{B \sim Q}[t(B)] \leq 1$. It must be that $t(X) \neq 0$, otherwise,

$$1.44d - 1 < \log t(X) <^+ \mathbf{d}(X|Q) + \mathbf{K}(t|Q) <^+ d + \mathbf{K}(d),$$

which is a contradiction for large enough d , which one can assume without loss of generality. Thus $t(X) \neq 0$ and $R \cap X_1 \neq \emptyset$. Furthermore, if $y \in R$, $\mathbf{K}(y) <^+ -\log \mathbf{m}_M(x) - n + \log d + \mathbf{K}(d, M, R)$. So for $x \in R \cap X_1$,

$$\begin{aligned} \mathbf{K}(x) &<^+ -\log \mathbf{m}_M(x) - n + \log d + \mathbf{K}(d, M, R) \\ \mathbf{K}(x) &<^+ \mathbf{K}(x) + 2s - n + \log d + \mathbf{K}(M) + \mathbf{K}(R, d) \\ n &<^+ 2s + \mathbf{K}(M) + \log d + \mathbf{K}(Q, d) \\ n &<^+ 2s + \mathbf{K}(\Omega[N]) + \mathbf{Ks}(X) \\ n &<^+ 2s + \mathbf{K}(\Omega[N]) + \mathbf{I}(X; \mathcal{H}) \end{aligned} \tag{17.1}$$

From Lemma 16, $\mathbf{K}(\Omega[N]|T, N) =^+ \mathbf{K}(\Omega[N]|X, N) = O(1)$. Furthermore it is well known for the bits of Chaitin's Omega, $N <^+ \mathbf{K}(\Omega[N])$ and $\mathbf{K}(\Omega[N]|\mathcal{H}) <^+ \mathbf{K}(N)$. So, using Lemma 4,

$$N <^+ \mathbf{K}(\Omega[N]) <^{\log} \mathbf{I}(\Omega[N]; \mathcal{H}) <^{\log} \mathbf{I}(X; \mathcal{H}) + \mathbf{K}(N) <^{\log} \mathbf{K}(X; \mathcal{H}). \tag{17.2}$$

So combining Equations 17.1 and 17.2, one gets

$$n <^{\log} 2s + 2\mathbf{I}(X; \mathcal{H}).$$

□

Lemma 25 *Given a computable probability measure μ , over $\{0, 1\}^\infty$, for an infinite set of unique infinite sequences $\{\alpha_i\}_{i=1}^\infty$ where $\mathbf{D}(\alpha_i|\mu) < \infty$ for all i , and an infinite set of numbers $\{c_i\}_{i=1}^\infty$, if $\sup_i |\mathbf{D}(\alpha_i|\mu) - c_i| < \infty$, then $\mathbf{I}(\{(\alpha_i, c_i) : \mathcal{H}\}) = \infty$.*

Proof. Suppose $c = \max_i \mathbf{D}(\alpha_i|\mu) < \infty$. Then let $D_n = \{\alpha_i\}_{i=1}^{2^n}$. So $\mathbf{K}(D_n|\{\alpha_i\}) <^+ \mathbf{K}(n)$. By Lemma 17, $n <^{\log} c + \mathbf{I}(D_n : \mathcal{H}) <^{\log} \mathbf{I}(\{\alpha_i\} : \mathcal{H}) + \mathbf{K}(n)$. Since this holds for all n ,

$$\infty = \mathbf{I}(\{\alpha_i\} : \mathcal{H}) = \mathbf{I}(\{(\alpha_i, c_i)\} : \mathcal{H}).$$

Suppose $\sup_i \mathbf{D}(\alpha_i|\mu) = \infty$ and $r = \max_i |\mathbf{D}(\alpha_i|\mu) - c_i| < \infty$. Given n and $\{(\alpha_i, c_i)\}$, one can find 2^n unique infinite sequences α_i , such that $|c_i - c_j| > 5s$ for $i \neq j$, $i, j \in \{1, \dots, 2^n\}$, and using computable μ , due to the definition of \mathbf{D} , one can compute prefixes $x_i \sqsubset \alpha_i$ where $|\log \mu(x) - \mathbf{K}(x) - c_i| <^+ 2r$. Setting $X_n = \{(x_i, \lceil -\log \mu(x_i) - c_i \rceil)\}$, and $s = 2r + 1$, invoking Lemma 24, gives

$$n <^{\log} 4r + 2\mathbf{I}(X_n; \mathcal{H}) <^{\log} 4r + \mathbf{I}(\{(\alpha_i, c_i)\} : \mathcal{H}) + \mathbf{K}(n),$$

Thus since this hold for arbitrary r , $\mathbf{I}(\{(\alpha_i, c_i)\} : \mathcal{H}) = \infty$. \square

17.2.2 Algorithmic Fine Grain Entropy Cannot be Approximated

It is easy to see that algorithmic fine grained entropy is uncomputable. In this section. we go one step further and show that information about algorithmic fine grained entropy coincides with information about the halting sequence.

Definition 32 (Mutual Information of Infinite Set of Points with \mathcal{H}) *Given an infinite set of infinite sequences $\{\alpha_i\}_{i=1}^\infty$, we encode them into a single infinite sequence $\langle \{\alpha_i\} \rangle$ in the standard way, in that α_1 is encoded into every other bit, α_2 is encoded into every other free space, and so on. Given computable metric space \mathcal{X} , the set of encodings for an infinite set of points in X , $\{\alpha_i\}$, $\alpha_i \in X$, $[\{\alpha_i\}]$, is the set of all encoded $\langle \{\beta_i\} \rangle$, where β_i is an encoding of a fast Cauchy sequence for α . The mutual information that an infinite set $\{\alpha_i\}$ of points has with the halting sequence is $\mathbf{I}(\{\alpha_i\} : \mathcal{H}) = \inf_{\beta \in [\{\alpha_i\}]} \mathbf{I}(\beta : \mathcal{H})$. In a standard way, a number can be appended to the start of each fast Cauchy sequence.*

Theorem 67 *Let (\mathcal{X}, μ) be a computable measure space and $\{\alpha_i\}_{i=1}^\infty$ be an infinite set of unique points in \mathcal{X} where $\mathbf{H}_\mu(\alpha_i) < \infty$, for each α_i . For the infinite set of numbers, $\{c_i\}_{i=1}^\infty$, if $\sup_i |\mathbf{H}_\mu(\alpha_i) - c_i| < \infty$, then $\mathbf{I}(\{(c_i, \alpha_i)\} : \mathcal{H}) = \infty$.*

Proof. Let $\bar{\mu} = \mu/\mu(X)$ be a computable probability measure due to 1. Let $(\delta, \bar{\mu}_\delta)$ be a binary representation for $(\mathcal{X}, \bar{\mu})$. Since (δ, δ^{-1}) is an isomorphism, by Corollary 17 and Proposition 14, and since $\mathbf{H}_\mu(\alpha_i) < \infty$, there is a constant c , where $|\mathbf{D}(\delta^{-1}(\alpha)|\bar{\mu}_\delta) - \mathbf{H}_\mu(\alpha)| < c$. Let β minimize $[\mathbf{I}(\{(c_i, \alpha_i)\} : \mathcal{H})]$. So by applying δ^{-1} to each of the fast Cauchy sequences of α_i encoded in β , one can construct an infinite set $\{(c_i, \gamma_i)\}_{i=1}^\infty$ of infinite sequences where $\sup_i |\mathbf{D}(\gamma_i|\bar{\mu}_\delta) - c_i| < \infty$. So, using Lemma 25,

$$\infty = \mathbf{I}(\{(c_i, \gamma_i)\}_{i=1}^\infty : \mathcal{H}) = \mathbf{I}(\beta : \mathcal{H}) = \mathbf{I}(\{(c_i, \alpha_i)\} : \mathcal{H}).$$

\square

17.3 Computability of Measures

In this section, we provide sufficient conditions for a measure to be uncomputable.

Theorem 68 *Let \mathcal{X} be a computable metric space and μ be a non-atomic measure over \mathcal{X} . If $\mu(\{\alpha : \mathbf{H}_\mu(\alpha) < -n\}) < 2^{-n-\mathbf{K}(n)-\omega(1)}$ then μ is uncomputable.*

Proof. Assume μ is computable. Let $\bar{\mu} = \mu/\mu(X)$ be a computable probability measure due to Claim 1. Let $(\delta, \bar{\mu}_\delta)$ be a binary representation for $(\mathcal{X}, \bar{\mu})$. Thus $\bar{\mu}_\delta$ is computable. Fix n . Let $U_n = \{\alpha \in X : \mathbf{H}_{\bar{\mu}}(\alpha) < -n\}$ and $\delta^{-1}(U_n) = V_n$. Then there is some $c \in \mathbb{R}_{\geq 0}$, independent of n , such that $V_n \supseteq \{\alpha \in \{0, 1\}^\infty : \mathbf{D}(\alpha | \bar{\mu}_\delta) > n - c\}$. By Theorem 64, there is a d , independent of n , where $\bar{\mu}_\delta(V_n) > 2^{-n - \mathbf{K}(n) - d}$. Since δ is a measure-preserving morphism, $\bar{\mu}(U_n) > 2^{-n - \mathbf{K}(n) - d}$ for all n , causing a contradiction.

Definition 33 (Mutual Information of Measures with \mathcal{H}) Let \mathcal{X} be a computable metric space $\mu \in \mathfrak{M}(X)$ be a measure over this space. Its mutual information with the halting sequence is $\mathbf{I}(\mu : \mathcal{H})$, where μ is treated as a point in the computable metric space $\mathfrak{M}(X)$, and Definition 27 is used.

Definition 34 (Neutral Measures) Let \mathcal{X} be a computable measure space. A measure μ is weakly neutral if $\forall \alpha \in X \mathbf{H}_\mu(\alpha) > -\infty$. A measure μ is neutral if $\inf_{\alpha \in X} \mathbf{H}_\mu(\alpha) > -\infty$,

Lemma 26 (Sperner's Lemma) Let p_1, \dots, p_k be points of some finite-dimensional space \mathbb{R}^n . Suppose that there are closed sets F_1, \dots, F_k with the property that for every subset $1 \leq i_1 < \dots < i_j < k$, the simplex $S(p_{i_1}, \dots, p_{i_j})$ spanned by p_{i_1}, \dots, p_{i_j} is covered by the union $F_{i_1} \cup \dots \cup F_{i_j}$. Then the intersection $\cap_i F_i$ is not empty.

Proposition 32 Let \mathcal{X} be a computable measure space. For every closed set $A \subset X$ and probability measure μ , if $\mu(A) = 1$ then there exists $a \in X$, $\mathbf{t}_\mu(x) \leq 1$.

Proof. $\int_X \mathbf{t}_\mu d\mu = \mu^x \mathbf{1}_A(x) \mathbf{t}_\mu(x) \leq 1$. □

Part (1) of the following theorem is due to [Lev76], and conveyed in [G21].

Theorem 69 Let \mathcal{X} be a computable metric space.

1. If \mathcal{X} is compact then it has a neutral measure.
2. If μ is weakly neutral then $\mathbf{I}(\mu : \mathcal{H}) = \infty$.

Proof. (1) For every $x \in X$, let F_x be the set of measures for which $\mathbf{t}_\mu(x) \leq 1$. Since \mathcal{X} is compact, the space of Borel probability measures $\mathcal{M}(X)$ over \mathcal{X} is compact. Therefore, due to compactness, if every finite subset of $\{F_x : x \in X\}$ of closed sets has a nonempty intersection, then $\cap_x F_x \neq \emptyset$. Let $S(x_1, \dots, x_k)$ be the simplex of probability measures concentrated on x_1, \dots, x_k . Proposition 32 implies each such measure belongs to one of the sets F_{x_i} . Thus $S(x_1, \dots, x_k) \subset F_{x_1} \cup \dots \cup F_{x_k}$ and this is true for any subset of the indices $\{1, \dots, k\}$. Lemma 26 implies $F_{x_1} \cap \dots \cap F_{x_k} \neq \emptyset$.

(2) Let $\bar{\mu} = \mu/\mu(X)$, where given a fast Cauchy sequence for μ , one can easily compute a fast Cauchy sequence for $\bar{\mu}$ (just normalize the ideal points). Let $(\delta, \bar{\mu}_\delta)$ be a binary representation of $(\mathcal{X}, \bar{\mu})$. Due to Lemma 13, $\mathbf{t}_{\bar{\mu}_\delta}(\alpha) < \infty$, for all $\alpha \in \{0, 1\}^\infty$, so $\bar{\mu}_\delta$ is a weakly neutral measure. Due to [DJ13], some PA degree is computable from any encoded Cauchy sequence $\langle \bar{\mu}_\delta \rangle \in \{0, 1\}^\infty$ of $\bar{\mu}_\delta$ in $\mathcal{M}(\{0, 1\}^\infty)$ space. Due to [Lev13], $\mathbf{I}(\langle \bar{\mu}_\delta \rangle : \mathcal{H}) = \infty$. So

$$\infty = \inf_{\bar{\mu}_\delta} \mathbf{I}(\langle \bar{\mu}_\delta \rangle : \mathcal{H}) = \mathbf{I}(\bar{\mu}_\delta : \mathcal{H}) <^+ \mathbf{I}(\bar{\mu} : \mathcal{H}) <^+ \mathbf{I}(\mu : \mathcal{H}).$$

□

17.4 Exotic Dynamics

Definition 35 Let \mathcal{X} be a computable metric space. Let $\alpha \in X$ be a point in this space and G^t a (potentially uncomputable) transformation group over \mathcal{X} . For $\beta \in \{0, 1\}^\infty$, let $\text{Num}(\beta) \in [0, 1]$ convert the infinite sequence β into a real in $[0, 1]$ in the standard way. Let $[(G^t, \alpha)]$ consist of all infinite sequences γ such that $U_\beta(\gamma)$ outputs an encoding of a fast Cauchy sequence to $G^{\text{Num}(\beta)}\alpha$. The amount of information that (G^t, α) has with the halting sequence is $\inf_{\gamma \in [(G^t, \alpha)]} \mathbf{I}(\gamma : \mathcal{H})$.

The following theorem states that if entropy does not oscillate enough during the course of dynamics, then the dynamics and starting point are exotic, containing infinite mutual information with the halting sequence. This theorem is a generalization of Theorem 53 in Chapter 13, though the bounds are looser.

Theorem 70 Let L be the Lebesgue measure. Let (\mathcal{X}, μ) be a computable measure space. There is a $c \in \mathbb{N}$ such that for (potentially uncomputable) transformation group G^t and point $\alpha \in X$ if $L\{t : \mathbf{H}_\mu(G^t\alpha) < \log \mu(X) - n\} < 2^{-n-c\mathbf{K}(n)}$, then $\mathbf{I}((G^t, \alpha) : \mathcal{H}) = \infty$.

Proof. Fix n . Let $(\{0, 1\}^\infty, \Gamma)$ be the Cantor space with the uniform measure. The binary representation (see Theorem 48) creates an isomorphism (ϕ, ϕ^{-1}) of computable probability spaces between the spaces $(\{0, 1\}^\infty, \Gamma)$ and $([0, 1], L)$. It is the canonical function $\phi(\gamma) = 0.\gamma$. Thus for all Borel sets $A \subseteq [0, 1]$, $\Gamma(\phi^{-1}(A)) = L(A)$. So

$$\begin{aligned} & L\{t : \mathbf{H}_\mu(G^t\alpha) < \log \mu(X) - n\} \\ &= \Gamma\{\beta : \mathbf{H}_\mu(G^{\phi(\beta)}\alpha) < \log \mu(X) - n\} \\ &= \Gamma\{\beta : \log \mathbf{t}_\mu(G^{\phi(\beta)}\alpha) > n - \log \mu(X)\} \\ &< 2^{-n-c\mathbf{K}(n)}. \end{aligned}$$

Let (δ, μ_δ) be a binary representation for (\mathcal{X}, μ) . Let $\bar{\mu}_\delta = \mu_\delta / \mu_\delta(\{0, 1\}^\infty)$ be a computable probability measure over $\{0, 1\}^\infty$. Thus, due to Lemm 13 and Proposition 14, there is a $d \in \mathbb{N}$ with $\psi(\beta) = \delta^{-1}(G^\beta\alpha)$ and

$$\begin{aligned} & \Gamma\{\beta : \log \mathbf{t}_\mu(G^{\phi(\beta)}\alpha) > n - \log \mu(X)\} \\ &> \Gamma\{\beta : \log \mathbf{t}_{\mu_\delta}(\psi(\beta)) > n - \log \mu(X) + d\} \\ &> \Gamma\{\beta : \mathbf{D}(\psi(\beta) | \mu_\delta) > n - \log \mu(X) + d\} \\ &> \Gamma\{\beta : \mathbf{D}(\psi(\beta) | \bar{\mu}_\delta) > n + d\}. \end{aligned}$$

Let $W \in \{0, 1\}^{\infty 2^{n+c\mathbf{K}(n)-1}}$ be a set of $2^{n+c\mathbf{K}(n)-1}$ infinite sequences with each sequence chosen independently the uniform distribution over $\{0, 1\}^\infty$. The probability that all $\beta \in W$ has $\mathbf{D}(\psi(\beta) | \bar{\mu}_\delta) \leq n + d$ is

$$(1 - 2^{-n-c\mathbf{K}(n)})^{2^{n+c\mathbf{K}(n)-1}} \geq (1 - 2^{n+c\mathbf{K}(n)-1} 2^{-n-c\mathbf{K}(n)}) \geq 1/2.$$

Let \mathcal{U} be a distribution over $\{0, 1\}^\infty$ that is the uniform measure applied independently to $2^{n+c\mathbf{K}(n)-1}$ encoded sequences. Let $\gamma \in [(G^t, \alpha)]$ that minimizes $[\mathbf{I}((G^t, \alpha) : \mathcal{H})] + 1$ and $\lambda = \langle p \rangle \gamma$, where p is a program to compute \mathcal{U} . By Theorem 52, relativized to v ,

$$\Pr_{\beta \sim \mathcal{U}} [\mathbf{I}(\langle \beta, \lambda \rangle : \mathcal{H}) > m] \stackrel{*}{<} 2^{-m+\mathbf{I}(\lambda : \mathcal{H})}.$$

Therefore by probabilistic arguments, there exist a set $W \in \{0, 1\}^{\infty 2^{n+c\mathbf{K}(n)}}$ such that for all $\beta \in W$,

$$\mathbf{D}(\psi(\beta)|\bar{\mu}_\delta) \leq n + d \text{ and } \mathbf{I}(\psi(W) : \mathcal{H}) <^+ \mathbf{I}(W : \mathcal{H}) <^+ \mathbf{I}((G^t, \alpha) : \mathcal{H}).$$

Thus Lemma 17 applied to $\psi(W)$ and $\bar{\mu}_\delta$, results in

$$\begin{aligned} \log |W| &< \max_{\beta \in \psi(W)} \mathbf{D}(\beta|\bar{\mu}_\delta) + 2\mathbf{I}((G^t, \alpha) : \mathcal{H}|v) + O(\mathbf{K}(|W|)). \\ n + c\mathbf{K}(n) &< n + d + 2\mathbf{I}((G^t, \alpha) : \mathcal{H}) + O(\mathbf{K}(n)). \\ c\mathbf{K}(n) &< d + 2\mathbf{I}((G^t, \alpha) : \mathcal{H}) + O(\mathbf{K}(n)). \end{aligned}$$

Thus for proper choice of c , $\mathbf{I}((G^t, \alpha) : \mathcal{H}) = \infty$. □

Corollary 28 *Let L be the Lebesgue measure. Let (\mathcal{X}, μ) be a computable measure space. There is a $c \in \mathbb{N}$ such that for (potentially uncomputable) transformation group G^t and point $\alpha \in X$ if $L\{t : \mathbf{t}_\mu(G^t \alpha) > 2^n / \mu(X)\} < 2^{-n-c\mathbf{K}(n)}$, then $\mathbf{I}((G^t, \alpha) : \mathcal{H}) = \infty$.*

Part IV

Newtonian Gravity

Chapter 18

Typicality in Newtonian Gravity

18.1 Introduction

The universal law of gravitation states that the force between two objects is

$$F = G \frac{m_1 m_2}{r^2},$$

where G is the gravitational constant, m_1 is the mass of the first object, m_2 is the mass of the second object, and r is the distance between the two objects. The gravitational field is a vector field describing the force of several (usually enormous) objects on a receiving object. The force vector applied to an object at position x by n objects at positions $\{x_i\}_{i=1}^n$ with masses $\{m_i\}_{i=1}^n$ is

$$\mathbf{G}_\kappa(x) = -G \sum_{i=1}^n m_i \frac{x - x_i}{\|x - x_i\|^3}.$$

The question is how to determine *typical* points in this vector field. In this chapter we proceed with the following approach to tackle this issue. We treat the magnitude of the force vector as a measure κ and define tests t to be lower computable functions over \mathbb{R}^3 such that $\kappa(t) < 1$. We prove the existence of a universal lower computable test \mathbf{T}_κ and use that as a score of *atypicality* of points in the vector field.

There are several differences in this approach than that of computable measure spaces in Chapter 11. First off, κ is infinite, so the space \mathfrak{M} of finite measures introduced in Chapter 11 cannot be leveraged. However, because the measure can be defined as a finite number of real numbers, there is a standardized method to encode them for the output of programs or as oracles.

Somewhat surprisingly, we show that \mathbf{T}_κ acts like a traditional test over compact spaces with suitable computability properties. Another difference is the assumption of \mathbb{R}^3 space instead of a complete, separable space, which enables theorems to leverage this property. In particular, interesting properties we proven about circular orbits of mass points.

The universal test \mathbf{T}_κ obeys conservation inequalities and can be defined similarly to the randomness deficiency term described in Definition 2. It is infinite over the mass points and decreasing in average as a point travels away from the mass point.

18.2 Preliminaries

Definition 36 L is the Lebesgue measure over \mathbb{R}^3 . L_1 is the Lebesgue measure over \mathbb{R} .

Definition 37 (Rational Rectangles) *The set of all open rectangles with rational coefficients is S . Elements of S are also referred to as rational open rectangles.*

This chapter will leverage computable metric spaces and computable measure spaces, detailed in Chapter 11. It will be particularly useful to define computable measure spaces over closed rational rectangles located in \mathbb{R}^3 . After such a space is defined then results of the previous chapters can be leveraged.

Definition 38 (Computable Measures) *When we say that a measure is R^3 -computable if it is a computable measure of the computable metric space $(R^3, \|\cdot\|, \mathbb{Q}^3)$. Equivalently, due to Propositions 10 and 11, measure μ is \mathbb{R}^3 -computable if $\mu(\mathbb{R}^3)$ is computable and for any finite set of open rational rectangles $V \subset S$, $\mu(\cup V)$ is lower computable.*

Definition 39 (Gravitational Field) *The gravity field κ caused by n point masses $\{x_i\}$ is*

$$\mathbf{G}_\kappa(x) = -G \sum_{i=1}^n M_i \frac{(x - x_i)}{\|x - x_i\|^3}.$$

Each M_i is the mass of mass point x_i and G is the gravitational constant. The magnitude of the gravitational field is $\|\mathbf{G}_\kappa(x)\|$. We denote κ also as a Borel measure, where for a lower semi-continuous function f over \mathbb{R}^3 ,

$$\kappa(f) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x, y, z) \|\mathbf{G}_\kappa(x, y, z)\| dx dy dz.$$

For Borel set S , $\kappa(S) = \kappa(\mathbf{1}_S)$. If κ is computable if $\{(m_i, x_i)\}$ is computable and thus the magnitude of the gravitational field at every point can be computed with a program. If κ is given as an oracle, then $\{(m_i, x_i)\}$ is provided in a standardized fashion and thus for all simple functions f , $\kappa(f)$ can be computed.

Remark 3 (Standard Representation) *A representation $\gamma \in \{0, 1\}^\infty$ of a system κ is an infinite sequence such that $U(\gamma)$ outputs, with increasing precision, the position and mass of each mass point of κ . A representation λ of κ is standard if for every representation γ of κ there is a program that on input γ outputs λ . In this chapter, when κ is given as an oracle or as an output of a program, it is given as a standard representation.*

Definition 40 *A set $W \subset \mathbb{R}^3$ is away from a system κ with mass point $\{x_i\}$ if $\min_{i, x \in W} \|x - x_i\| > 0$.*

Simple functions are defined as follows. They are used as building blocks for lower computable functions and also tests.

Definition 41 (Simple Function) *A simple function is of the form $f((x, y, z)) = [(x, y, z) \in s]q$, where $q \in \mathbb{Q}_{>0}$ and s is a rational rectangle in S . Simple functions can be enumerated as $\{f_n\}$. Given κ , $\kappa(f_n)$ is uniformly computable in n .*

Definition 42 (Tests) *A lower computable function T is of the form $T(x, y, z) = \sup_{n \in N} f_n(x, y, z)$, where N is an enumerable subset of \mathbb{N} . The set of all lower computable κ -tests is $[\kappa]$, where lower computable functions T is in $[\kappa]$ if*

$$\kappa(T) \leq 1.$$

Among all κ -tests there exist universal tests which dominant every member of $[\kappa]$. This universal test is the key metric for determining the level of atypicality of points in a gravitational field. The greater the score, the greater the level of strangeness of the point. Note that equivalently, \mathbf{T}_κ can be defined using lower computable tests over open spheres with rational centers and radii.

Definition 43 (Universal Test) *The universal test is $\mathbf{T}_\kappa(x) = \sum_{T_i \in [\kappa]} \mathbf{m}(i|\kappa)T_i(x)$. $\mathbf{D}_r(x) = \log \mathbf{T}_r(x)$.*

18.3 Properties of Universal Tests

Despite the fact that for a system κ with mass point x_i , $\lim_{x \rightarrow x_i} \mathbf{G}_\kappa(x) = \infty$, $\kappa(B)$ is proportional the radius of balls centered on x_i . Thus, as shown below, as points approach the positions of mass points, they will become more atypical with respect to \mathbf{T}_κ .

Theorem 71 *Let κ be a system with mass points $\{x_i\}$. Given i , there is a $c, d \in \mathbb{R}^+$ such that for all open spheres G_r centered on x_i with radius $r < c$, $\kappa(G_r) < d \cdot r$.*

Proof. Choose c such that $x_j \notin \overline{G_c}$ for all $j \neq i$. Let κ_j be the system containing just mass point x_j . $\kappa(G_r) \leq \sum_{j=1}^n \kappa_j(G_r)$. It is easy to see that since $x_j \notin \overline{G_c}$, for $j \neq i$, we have that $\sum_{j \neq i} \kappa_j(G_r) < fr^3$ for some $f \in \mathbb{R}_{>0}$. $\kappa_i(G_r) = \int_0^r \frac{m_i}{r^2} 4\pi r^2 dr = 4m_i\pi r = gr$, for some $g \in \mathbb{R}_{>0}$. Thus for all $r < c$, $\kappa(G_r) < fr^3 + gr < d \cdot r$.

Corollary 29 *System κ is a locally finite Borel measure.*

Corollary 30 *For system κ , \mathbf{T}_κ is positive over all \mathbb{R}^3 .*

Corollary 31 *For each mass point x_i of a system κ , for each $c > 0$ there is an $r > 0$ such that $\forall x: \|x_i - x\| < r, \mathbf{T}_\kappa(x) > c$.*

Proof. Let G_r be an open ball centered on x_i with radius r . By Theorem 71, there is an r_c where $\kappa(G_{r_c}) < 2^{-n}$. We define the test $t_c(x) = [x \in G_{r_c}]2^n$. Thus for all $x \in G_{r_c}$, $\mathbf{T}_\kappa(x) \stackrel{*}{>} 2^{n-2\log n}$.

Thus each center of mass will have infinite atypicality score.

Corollary 32 *For system κ with mass point x , $\mathbf{T}_\kappa(x) = \infty$.*

Theorem 72 *Let κ be a system and let $w = \max\{\|x_i\| : x_i \text{ is a mass point of } \kappa\}$. U_r be the uniform distribution over the subspace $\{x : r < \|x\| < 2r\}$. Then there is a c where if $r > 2w$ then $U_r(\mathbf{T}_\kappa) < c/r$.*

Proof. Let $\mathbf{T}_\kappa^r(x) = [r < \|x\| < 2r]\mathbf{T}_\kappa(x)$. Let $\theta(z) = \tan^{-1}(w/z)$. Then there is a $c \in \mathbb{N}$ where

$$\begin{aligned}
1 > \kappa(\mathbf{T}_\kappa^r) &= \int_{x,y,z:r < \|(x,y,z)\| < 2r} \mathbf{T}_\kappa(x,y,z) \left\| \sum_{i=1}^n m_i \frac{\widehat{(x,y,z)} - (x_i)}{\|(x,y,z) - (x_i)\|^2} \right\| dx dy dz \\
&\geq \int_{x,y,z:r < \|(x,y,z)\| < 2r} \mathbf{T}_\kappa(x,y,z) \left\| \sum_{i=1}^n m_i \frac{\cos \theta(r)}{4r^2} \right\| dx dy dz \\
&\geq \frac{c}{r^2} \int_{x,y,z:r < \|(x,y,z)\| < 2r} \mathbf{T}_\kappa(x,y,z) dx dy dz \\
&= \frac{c}{r^2} U_r(\mathbf{T}_\kappa) \frac{4}{3} \pi ((2r)^3 - r^3) \\
U_r(\mathbf{T}_\kappa) &< c/r.
\end{aligned}$$

□

Theorem 73 *If point $x \in \mathbb{R}^3$ is a computable distance r from a computable point y , then $\mathbf{T}_\kappa(x) = \infty$.*

Proof. For positive rational $q \in \mathbb{Q}_{>0}$, let $V_q = \{z : r - q < \|z - y\| < r + q\}$. Let W_q consist of all unions Z of a finite number of open rectangles in S such that $V_q \subset Z$. We define the test t_n as follows: enumerate all open sets of W_q for all $q \in \mathbb{Q} \cap (0, 1]$ and stop when an open set $Z \in \cup_{q \in \mathbb{Q} \cap (0, 1]} W_q$ is found such that $\kappa(Z) < 2^{-n}$. Then we define the following test, with $t_n(x) = [x \in Z]2^n$. Since $x \in Z$, $t_n(x) = 2^n$. Thus $\mathbf{T}_\kappa(x) \stackrel{*}{>} \sum_n \mathbf{m}(n)t_n(x) \stackrel{*}{>} \sum_n 2^{n-2 \log n} = \infty$.

Corollary 33 *If a point $x \in \mathbb{R}^3$ is a computable distance from the mass point of a system κ then $\mathbf{T}_\kappa(x) = \infty$.*

Proof. Though the mass points may not be computable, κ is given as an oracle to the universal test \mathbf{T}_κ , and thus the mass points are effectively computable.

18.4 Conservation of Randomness

A key property universally shared between notions of typicality and randomness deficiency is conservation of randomness. The author cannot think of a single example of a robust randomness deficiency property that doesn't have this property. Conservation of randomness says that if the measure and the point are transformed by the same means (usually a total computable function) then its randomness deficiency is constant (or usually changed up to an order of the complexity of the transform). For example Theorem 25 is a conservation of quantum typicality, over quantum operations. In this section we show that random deficiency in magnetic fields is conserved over non-singular linear transformations and systems with a single mass point. It is an open question whether there is conservation of randomness over systems with multiple mass points.

For system κ and invertible matrix A , $A\kappa$ is the system where all the mass points have the matrix A applied to them. For invertible matrix A $A\mathbf{T}_\kappa(x) = \sum_{T_i \in [\kappa]} \mathbf{m}(i|\kappa) T_i(A^{-1}x)$. Similarly, for 3-vector k , $\kappa + k$ displaces all mass points by k and $(\mathbf{T} + k)_\kappa(x) = \sum_{T_i \in [\kappa]} \mathbf{m}(i|\kappa) T_i(x - k)$. $(A\mathbf{D} + k)(x) = \log(A\mathbf{T} + k)(x)$.

Theorem 74 (Conservation of Randomness Deficiency) *For system κ with a single mass point (x_i, y_i, z_i) , for computable invertible 3×3 matrix A and vector k , $(A\mathbf{D} + k)_{A\kappa + k}(x) <^+ \mathbf{D}_\kappa(x)$.*

Proof. $\mathbf{T}_{A\kappa}(y) = \sum_{T_i \in [A\kappa]} \mathbf{m}(i|\kappa) t_i(x)$. Let $\{\lambda_1, \lambda_2, \lambda_3\}$ be the eigenvalues of A , $m = \min |\lambda_i|$ and $M = \max |\lambda_i|$. We use the fact that $m\|x\| \leq \|Ax\| \leq M\|x\|$. We restrict our attention to simple functions, and the generalization to lower computable functions is straightforward. Let simple function f be defined by $f(x, y, z) = [(x, y, z) \in ((x_1, x_2), (y_1, y_2), (z_1, z_2))]v$, with $v \in \mathbb{Q}_{>0}$.

$$\begin{aligned}
A\kappa(f) &= \int \int \int s(x, y, z) \left\| m_i \frac{A(x_i, y_i, z_i) - (x, y, z)}{\|A(x_i, y_i, z_i) - (x, y, z)\|^3} \right\| dx dy dz \\
&= v \int_{x_1}^{x_2} \int_{y_1}^{y_2} \int_{z_1}^{z_2} \left\| m_i \frac{A(x_i, y_i, z_i) - (x, y, z)}{\|A(x_i, y_i, z_i) - (x, y, z)\|^3} \right\| dx dy dz \\
A\kappa(Af) &= v \int_{x_1}^{x_2} \int_{y_1}^{y_2} \int_{z_1}^{z_2} \left\| m_i \frac{A(x_i, y_i, z_i) - A(x, y, z)}{\|A(x_i, y_i, z_i) - A(x, y, z)\|^3} \right\| dx dy dz \\
&\leq \frac{v}{m^3} \int_{x_1}^{x_2} \int_{y_1}^{y_2} \int_{z_1}^{z_2} \left\| m_i \frac{A(x_i, y_i, z_i) - A(x, y, z)}{\|(x_i, y_i, z_i) - (x, y, z)\|^3} \right\| dx dy dz \\
&\leq \frac{vM}{m^3} \int_{x_1}^{x_2} \int_{y_1}^{y_2} \int_{z_1}^{z_2} \left\| m_i \frac{(x_i, y_i, z_i) - (x, y, z)}{\|(x_i, y_i, z_i) - (x, y, z)\|^3} \right\| dx dy dz \\
&\stackrel{*}{\leq} \kappa(f).
\end{aligned}$$

Furthermore, since A is computable, $\mathbf{m}(Af|\kappa) \stackrel{*}{\leq} \mathbf{m}(f|\kappa)/\mathbf{m}(A)$. The k offset argument is self evident. \square

Corollary 34 *For computable matrix A with identical eigenvalues, computable vector k , and system κ . $(\mathbf{AD} + k)_{A\kappa+k}(x) <^+ \mathbf{D}_\kappa(x)$*

Proof. This corollary follows from the reasoning in the proof of Theorem 74. \square

18.5 Comparable Definitions of Typicality

A point $x \in \mathbb{R}^3$ is test random with respect to system κ iff $\mathbf{T}_\kappa(x) < \infty$. A κ -ML test is a series of uniformly enumerable open sets $\{V_n\}$, relative to oracle κ , such that

$$\kappa(V_n) < 2^{-n}.$$

There is a universal ML test U_n such that if $x \in \cap_n V_n$, then $x \in \cap_n U_n$. We say that a point x is ML random if $x \notin \cap_n U_n$.

Proposition 33 *For system κ , $\mathbf{T}_\kappa(x) \neq \infty$ iff x is ML random.*

Proof. We define the ML test $V_n = \{x : \mathbf{T}_\kappa(x) > 2^n\}$. Clearly $\kappa(V_n) < 2^{-n}$, since \mathbf{T}_κ is a test. Thus if $\mathbf{T}_\kappa(x) = \infty$, then $x \in \cap_n V_n \subseteq \cap_n U_n$. Given a ML test V_n , we define the test $T(x) = \sup_n \mathbf{m}(n)[x \in V_n]2^n$. So if $x \in \cap_n V_n$, then $T(x) = \infty$ and thus $\mathbf{T}_\kappa(x) = \infty$. \square

Definition 44 (Randomness Deficiency) *The randomness deficiency of a point $x \in \mathbb{R}^3$ with respect to a system κ is $\mathbf{d}(x|\kappa) = \sup_{x \in s \in S} -\log \kappa(s) - \mathbf{K}(s|\kappa)$.*

Definition 45 (Mixed Rectangles) *Let \hat{S} consist of all “mixed” rectangles with rational boundaries, and each boundary can be open or closed. Thus there are 2^6 types of such rectangles. One example rectangle is $\{(x, y, z) : x \in [x_1, x_2), y \in (y_1, y_2], z \in [z_1, z_2]\}$. Let $\hat{\mathbf{d}}(x|\kappa) = \sup_{x \in \hat{s} \in \hat{S}} -\log \kappa(\hat{s}) - \mathbf{K}(\hat{s}|\kappa)$.*

Proposition 34 *Given a system κ and mixed rectangle $\hat{s} \in \hat{S}$ there exists a rational open rectangle $s \in S$ such that $\hat{s} \subseteq s$, $\kappa(s) < 2\kappa(\hat{s})$, and $\mathbf{K}(s|\kappa) <^+ \mathbf{K}(\hat{s}|\kappa)$.*

Proof. Because every locally finite Borel measure on a separable complete metric space is regular, κ is regular. So there exists an open cover $W \subset \mathbb{R}^3$ such that $\hat{s} \subseteq W$ and $\kappa(W) < 2\kappa(\hat{s})$. So there exists an open rectangle $s \in S$, $s \subseteq W$ that contains \hat{s} . This rectangle can be found using brute force search given \hat{s} and κ .

Note that the following terms hold over the range $\mathbb{R} \cup \{\infty\}$.

Theorem 75 For system κ , $\mathbf{d}(x|\kappa) =^+ \mathbf{D}_\kappa(x)$.

Proof.

(1) $\mathbf{d}(x|\kappa) <^+ \mathbf{D}_\kappa(x)$. For each $s \in S$, one can define a test $T(x) = [x \in s \in S] \mathbf{m}(s|\kappa)/\kappa(s)$. It is easy to see that $\kappa(T) \leq \sum_{s \in S} \mathbf{m}(s|\kappa)(\kappa(s)/\kappa(s)) < 1$, and that T is lower computable. So $\mathbf{d}(x|\kappa) <^+ \log T(x) <^+ \mathbf{D}_\kappa(x)$.

(2) If $\mathbf{D}_\kappa(x) \in \mathbb{R}$, then $\mathbf{D}_\kappa(x) <^+ \mathbf{d}(x|\kappa)$. One can assume the range of \mathbf{T}_κ are 0 and powers of 2. For $n \in \mathbb{Z}$, Let $W_n = \{x : T_r(x) = 2^n\}$. Thus since \mathbf{T}_r is a test, $r(W_n) \leq p(n)2^{-n}$, for some probability p over \mathbb{N} . Let $V_n \subset S$ be the (uniformly in n) enumerable set of disjoint rectangles $\hat{s} \in \hat{S}$ such that $\cup_{\hat{s} \in V_n} \hat{s} = W_n$. Each $\hat{s} \in V_n$ can be identified by a code of size $\mathbf{K}(\hat{s}|\kappa) <^+ -\log \kappa(\hat{s}) - n$. Thus if $n = \mathbf{D}_\kappa(x) <^+ -\log \kappa(\hat{s}) - \mathbf{K}(\hat{s}|\kappa) <^+ \hat{\mathbf{d}}(x|\kappa)$.

Let $\hat{s} \in \hat{S}$ realize $\mathbf{D}_\kappa(x) + O(1)$. Thus by Proposition 34 there is an open rectangle $s \in S$ such that $s \supset \hat{s}$, $\kappa(\hat{s}) < \kappa(s) < 2\kappa(\hat{s})$ and $\mathbf{K}(s|\kappa) <^+ \mathbf{K}(\hat{s}|\kappa)$. Thus $\hat{\mathbf{d}}(x|\kappa) <^+ -\log \kappa(\hat{s}) - \mathbf{K}(\hat{s}|\kappa) <^+ -\log \kappa(s) - \mathbf{K}(s|\kappa) <^+ \mathbf{d}(x|\kappa)$.

(3) If $\mathbf{D}_\kappa(x) = \infty$ then $\mathbf{d}(x|\kappa) = \infty$. Thus there is a ML test $\{V_n\}$ such that $x \in \cap_n V_n$. Fix n . Thus there is a finite or infinite set $W_n \subset \hat{S}$ of disjoint mixed rectangles such that $\cup_{\hat{s} \in W_n} \hat{s} = V_n$. Since $\kappa(V_n) < 2^{-n}$, each $\hat{s} \in W_n$ can be identified by a code of size $\mathbf{K}(\hat{s}|\kappa) <^+ -\log \kappa(s) - n + \mathbf{K}(n|\kappa)$. Thus there is some $\hat{s} \in W_n$ such that $x \in \hat{s}$. So $-\log \kappa(\hat{s}) - \mathbf{K}(\hat{s}|\kappa) >^{\log} n$. Since this occurs for each n , $\hat{\mathbf{d}}(x|\kappa) = \sup_{x \in \hat{s} \in \hat{S}} -\log \kappa(s) - \mathbf{K}(\hat{s}|\kappa) = \infty$. Using the same reasoning as in (2), one can see that $\mathbf{d}(x|\kappa) = \infty$. \square

Corollary 35 For computable invertible 3×3 matrix A and vector k , $\mathbf{d}(Ax+k|\kappa k) <^+ \mathbf{d}(x|\kappa)$.

18.6 Compact Subspaces

In this section, we'll show that the universal test \mathbf{T}_κ behaves like universal uniform test \mathbf{t} defined in Chapter 11 over sufficiently computable compact structures. Over such structures, the \mathbf{T} is shown to be asymptotically balanced.

Definition 46 An open set $W \subseteq \mathbb{R}^3$ is computable if given an open rational rectangle $s \in S$, $\mathbf{1}_W(s) = [s \subseteq W]$ is computable.

Theorem 76 Let V be an open compact computable set, κ be a computable system, and let $\kappa(V)$ be computable. There is a constant $c \in \mathbb{N}$ where $2^{-n-\mathbf{K}(n)-c} < \kappa(\{x : x \in V, \mathbf{T}_\kappa(x) > 2^n\}) < 2^{-n+c}$.

Proof. For Borel set X , let $\kappa'(X) = \kappa(X \cap V)$. So $\mathcal{R} = (\mathbb{R}^3, \|\cdot\|, \mathbb{Q}^3)$ is a computable metric space and (\mathcal{R}, κ') is a computable measure space. Thus κ' is a \mathbb{R}^3 -computable measure since Proposition 11 applies because $\kappa'(V) = \kappa(V)$ is computable and for every $s \in S$,

$$\kappa'(s) = \kappa\left(\bigcup\{s' : s' \in s' \subseteq s \text{ and } s \subseteq V\}\right)$$

which is lower computable because V is computable and κ is computable. Since V is computable, and $\mathbf{t}_{\kappa'}(V) < 1$, $\mathbf{t}_{\kappa'} \stackrel{*}{<} \mathbf{T}_{\kappa}$. Also $\mathbf{t}'_{\kappa'}(x) \stackrel{*}{>} [x \in V] \mathbf{T}_{\kappa}(x)$. So over $x \in V$ $\mathbf{t}_{\kappa'} \stackrel{*}{=} \mathbf{T}_{\kappa}$. By Proposition 20, there is a $c \in \mathbb{N}$ where for all $n \in \mathbb{N}$

$$\kappa'(V)2^{-n-\mathbf{K}(n)-c} < \kappa'\{x : x \in V, \mathbf{t}_{\kappa'}(x) > 2^n / \kappa'(V)\} < \kappa'(V)2^{-n}.$$

The $\kappa'(V)$ term can be folded in the constant c . So,

$$2^{-n-\mathbf{K}(n)-c} < \kappa\{x : x \in V, \mathbf{T}_{\kappa}(x) > 2^n\} < 2^{-n+c}.$$

□

Corollary 36 *Let V be a compact computable open set and let κ be a computable system with computable $L(V)$. Then there is a $c \in \mathbb{N}$ such that for all $n \in \mathbb{N}$, $L\{x : x \in V, \mathbf{T}_{\kappa}(x) > 2^n\} > 2^{-n-\mathbf{K}(n)-c}$.*

Proof. For Borel set W , the function $\lambda(W) = L(W \cap V)/L(V)$ is a \mathbb{R}^3 -computable measure. By Theorem 76, there is a $c \in \mathbb{N}$ where $\lambda\{x : x \in V, \mathbf{T}_{\kappa}(x) > 2^n\} > 2^{-n-\mathbf{K}(n)-c}$. The corollary follows from the fact that in the domain of V , $L(V)\lambda(\cdot) = L(\cdot)$.

Corollary 37 *Let V be an open compact computable set away from a computable system κ and let $\kappa(V)$ be computable. There is a constant $c \in \mathbb{N}$ where $2^{-n-\mathbf{K}(n)-c} < L\{x : x \in V, \mathbf{T}_{\kappa}(x) > 2^n\} < 2^{-n+c}$.*

Proof. Since V is away from κ , there are constants $b_1, b_2 \in \mathbb{R}_{>0}$ such that $b_1 < \mathbf{G}_{\kappa}(x) < b_2$ for all $x \in V$. So for Borel set A , $b_1 \int_A dx dy dz < \kappa(A) < b_2 \int_A dx dy dz$. This means $b_1 L(A) < \kappa(A) < b_2 L(A)$. Applying this equation to Theorem 76 results in the corollary.

Theorem 77 *Let V be a Borel compact set, κ be a computable system κ , and \mathbb{R}^3 -computable non-atomic probability measure λ have support equal to V . Then there is a $c \in \mathbb{N}$ where $\lambda\{x : x \in V, \mathbf{T}_{\kappa}(x) > 2^n\} > 2^{-n-\mathbf{K}(n)-c}$.*

Proof. Let W be a open computable compact set containing V and let $\kappa'(A) = \kappa(A \cap W)$. Thus $\mathcal{R} = (\mathbb{R}^3, \|\cdot\|, \mathbb{Q}^3)$ is a computable metric space and (\mathcal{R}, κ') is a computable measure space. Using the arguments in Theorem 76, for all $x \in W$, $\mathbf{t}_{\kappa'} \stackrel{*}{=} \mathbf{T}_{\kappa}(x)$. Applying Corollary 25 to λ and κ' provides a $c \in \mathbb{N}$ such that

$$\begin{aligned} 2^{-n-\mathbf{K}(n)-c} &< \lambda\{x : \mathbf{t}_{\kappa'}(x) > 2^n\}, \\ 2^{-n-\mathbf{K}(n)-c} &< \lambda\{x : x \in V, \mathbf{T}_{\kappa}(x) > 2^n\}. \end{aligned}$$

□

18.7 Orbits

Notions of typicality can be applied to orbits. Orbits are circular 1 dimensional paths around a mass point. If an orbit is at a computable distance from a mass point, then all its points will have infinite \mathbf{T} -scores. Otherwise, as a point orbits around a mass point, it will have oscillations of typicality, similarly to oscillations of algorithmic fine grained entropy during dynamics.

Definition 47 (Orbit) Let $\hat{\mathbf{x}}$ and $\hat{\mathbf{y}}$ be two perpendicular unit vectors in \mathbb{R}^3 . Let κ be a system. An orbit is defined by $\{r \cos(\theta)\hat{\mathbf{x}} + r \sin(\theta)\hat{\mathbf{y}} + z : \theta \in [0, 2\pi], r \in \mathbb{R}_{>0}\}$, where z is at a mass point. A representation $\alpha \in \text{Rep}(O) \subset \{0, 1\}^\infty$ of an orbit $O = (r, \hat{\mathbf{x}}, \hat{\mathbf{y}}, z)$ is any sequence such that $U(\alpha)$ outputs r , z , $\hat{\mathbf{x}}$, and $\hat{\mathbf{y}}$ to any degree of precision. An orbit is computable if it has a computable representation. The amount of information that an orbit O has with the halting sequence \mathcal{H} is $\mathbf{I}(O : \mathcal{H}) = \inf_{\alpha \in \text{Rep}(O)} \mathbf{I}(\alpha : \mathcal{H})$. We use the notation (r, z) -orbit to specify an orbit with arbitrary $\hat{\mathbf{x}}$ and $\hat{\mathbf{y}}$ vectors.

Corollary 38 If any of the following conditions occur for an orbit O ,

1. O is computable,
2. O is a computable distance from a mass point of system κ ,
3. O is centered at a mass point with a computable axis of rotation,

then $\mathbf{T}_\kappa(x) = \infty$, for all $x \in O$.

Proof. (1) and (2) follow from Theorem 73. Otherwise let $O = (\hat{\mathbf{x}}, \hat{\mathbf{y}}, z)$, where z is a mass point and $(\hat{\mathbf{x}}, \hat{\mathbf{y}})$ is computable. Let $r_1 < r_2$, $r_1, r_2 \in \mathbb{Q}_{>0}$. Let

$$W_q = \{x : r_1 < r_2, x \text{ differs from } (\hat{\mathbf{x}}, \hat{\mathbf{y}}) \text{ axis by } q \text{ radians}\}$$

Given n , one can find a q_n such that $\kappa(W_{q_n}) < 2^{-n}$ and define a test $t_n(x) = [x \in W_{q_n}]2^n$. So for all $x \in O$, $\mathbf{T}_\kappa(x) > \sum_n \mathbf{m}(n|\kappa)t_n(x) > \sum_n 2^{n-2\log n} = \infty$. \square

Corollary 39 Let κ be a system with mass point x_i . For all $c > 0$, there is an $r > 0$ such that for all (r, x_i) orbits O , $\mathbf{T}_\kappa(x) > c$, for all $x \in O$.

Proof. This follows from Corollary 31. \square

Remark 4 We recall that the mutual information of a point $x \in \mathbb{R}^3$ with the halting sequence \mathcal{H} , is $\mathbf{I}(x : \mathcal{H})$, and is introduced in Definition 27.

Claim 2 Given a representation α of an orbit O , there is a computable transformation group G^t around orbit O , such that for all $x \in O$, $G^0x = G^1x = x$. That is, given a fast Cauchy sequence of x and an $t \in \mathbb{R}_{\geq 0}$ one can compute a fast Cauchy sequence for $G^t x$.

Theorem 78 Let κ be a computable system with mass point z and $O = (r, \hat{\mathbf{x}}, \hat{\mathbf{y}}, z)$ be an orbit such that $\mathbf{I}(O : \mathcal{H}) < \infty$. There is a $c \in \mathbb{N}$ where for all $n \in \mathbb{N}$,

$$2^{-n-\mathbf{K}(n)-c} < L_1\{x : x \in O, \mathbf{T}_\kappa(x) > 2^n\}.$$

Furthermore if κ only contains a single mass point z and $(\hat{\mathbf{x}}, \hat{\mathbf{y}})$ is computable, then there is a $c \in \mathbb{N}$ for all $n \in \mathbb{N}$,

$$2^{-n-\mathbf{K}(n)-c} < L_1\{x : x \in O, \mathbf{T}_\kappa(x) > 2^n\} < 2^{-n+c}.$$

Proof. By Claim 2, given O , one can compute a (potentially uncomputable) transformation group G^t such that for all $x \in O$, $G^t x \in O$ for $t \in [0, 1]$, $G^0 x = G^1 x = x$. This transformation group G^t is a rotation centered around mass point z and oriented with the axis (\hat{x}, \hat{y}) . Furthermore there is an x computable from O . Using Definition 35, $\mathbf{I}((G^t, x) : \mathcal{H}) <^+ \mathbf{I}(O : \mathcal{H}) < \infty$.

Let $B = \{y : \|y - z\| < R\}$ where $r < R$, $R \in \mathbb{Q}_{>0}$. For Borel set A , let $\kappa'(A) = \kappa(A \cap B)$. Thus $\mathcal{R} = (\mathbb{R}^3, \|\cdot\|, \mathbb{Q}^3)$ is a computable metric space and (\mathcal{R}, κ') is a computable metric space. In addition B is a compact open computable set. Thus using arguments in the proof of Theorem 76, $\mathbf{t}_{\kappa'}(x) \stackrel{*}{=} \mathbf{T}_{\kappa}(x)$ for all $x \in B$. The application of Corollary 28, applied to κ' and (G^t, x) noting that the transform group need not be measure preserving, with the dynamics starting at x results in a $c \in \mathbb{N}$ where

$$\begin{aligned} 2^{-\mathbf{K}(n)-n-c} &< L_1\{t \in [0, 1], \mathbf{t}_{\kappa'}(G^t x) > 2^n\} \\ 2^{-\mathbf{K}(n)-n-c} &< L_1\{y : y \in O, \mathbf{t}_{\kappa'}(y) > 2^n\}. \end{aligned} \quad (18.1)$$

If the system κ only contains a single mass point z and (\hat{x}, \hat{y}) is computable, then G^t is a computable transformation transform rotating around z . Furthermore G^t is measure preserving over the computable measure space (\mathcal{R}, κ') . Thus applying Theorem 18 results $c \in \mathbb{N}$ where

$$\begin{aligned} 2^{-\mathbf{K}(n)-n-c} &< L_1\{t : t \in [0, 1], \mathbf{t}_{\kappa'}(G^t x) > 2^n\} < 2^{-n} \\ 2^{-\mathbf{K}(n)-n-c} &< L_1\{y : y \in O, \mathbf{t}_{\kappa'}(y) > 2^n\} < 2^{-n}. \end{aligned} \quad (18.2)$$

The fact that $\mathbf{t}_{\kappa'}(x) \stackrel{*}{=} \mathbf{T}_{\kappa}(x)$ for all $x \in O \subset B$ can be applied to Equations 18.1 and 18.2, proving the theorem. \square

Remark 5 (Extended Universal Tests) The set of cylinders is denoted by \mathcal{C} , where $\zeta \in \mathcal{C}$ if there is an $x \in \{0, 1\}^*$ such that $\zeta = \{\alpha : x \sqsubset \alpha \in \{0, 1\}^\infty\}$. Given the space $\mathcal{Z} = \mathbb{R}^3 \times \mathbb{R}^3 \times \{0, 1\}^\infty \times \{0, 1\}^\infty$, a basis extended rectangle $p \in P$ with respect to \mathcal{Z} , are two rational rectangles and two intervals $p = (s_1, s_2, \zeta_1, \zeta_2)$, where $s_1, s_2 \in S$ and $\zeta_1, \zeta_2 \in \mathcal{C}$. A simple function $f : \mathcal{Z} \rightarrow \mathbb{R}_{\geq 0}$ is of the form $f_{s_1, s_2, \zeta_1, \zeta_2}(w, x, y, z) = v[w \in s_1, x \in s_2, y \in \zeta_1, z \in \zeta_2]$. Simple functions can be enumerated producing the list $\{f_n\}$. A lower computable function F is of the form $F(w, x, y, z) = \sup n \in \mathbb{N} f_n(w, x, y, z)$ where N is an enumerable subset of \mathbb{N} . Given computable measure μ over $\{0, 1\}^\infty$ and system κ , a lower computable function F is a (κ, μ) -test, or $f \in [\kappa, \mu]$ if

$$(\kappa, \mu)(F) = \int_{\alpha, \beta \in \{0, 1\}^\infty} \int_{x_1 \in \mathbb{R}^3, x_2 \in \mathbb{R}^3} F(x_1, x_2, \alpha, \beta), d\kappa(x_1) d\kappa(x_2) d\mu(\alpha) d\mu(\beta) \leq 1.$$

Given κ and μ , the set $[\kappa, \mu]$ is enumerable, so there exists a universal lower computable test $\mathbf{T}_{\kappa, \mu}(w, x, y, z) = \sum_{f_i \in [\kappa, \mu]} \mathbf{m}(i | \kappa, \mu) f_i(w, x, y, z)$.

Remark 6 Let $\text{Cauchy}(x)$ be all the fast Cauchy sequences converging to $x \in \mathbb{R}^3$, using the computable metric space $(\mathbb{R}^3, \|\cdot\|, \mathbb{Q}^3)$. For $\zeta \in \text{Cauchy}(x)$, $\langle \zeta \rangle \in \{0, 1\}^\infty$ is an encoding of the fast Cauchy sequence by an infinite encoding of each index of the ideal points converging to x . For points $x_1, x_2 \in \mathbb{R}^3$ and infinite sequences $\alpha, \beta \in \{0, 1\}^\infty$, their multiple information with the halting sequence is

$$\mathbf{I}((x_1, x_2, \alpha, \beta) : \mathcal{H}) = \inf_{(\xi, \zeta) \in (\text{Cauchy}(x_1), \text{Cauchy}(x_2))} \mathbf{I}(\langle \xi \rangle, \langle \zeta \rangle, \alpha, \beta : \mathcal{H}).$$

Remark 7 We recall Definition 29, where an r -interval $v \subseteq (0, 1)$ is a finite collection of open intervals with rational endpoints.

In Theorem 78, it was shown that non-exotic orbits will oscillate in typicality. The following theorem extends this result to two orbits. Given two orbits with two starting points that are not atypical or exotic, the typicality of the points will be out of sync as they orbit the mass points.

Theorem 79 *Let κ be a computable system, μ a computable probability measure over $\{0,1\}^\infty$ and $O_1 \ni x_1$ and $O_2 \ni x_2$ be orbits with $\alpha_1 \in \text{Rep}(O_1)$ and $\alpha_2 \in \text{Rep}(O_2)$. If $\mathbf{T}(x_1, x_2, \alpha_1, \alpha_2) < \infty$ and $\mathbf{I}((x_1, x_2, \alpha_1, \alpha_2) : \mathcal{H}) < \infty$ then for transformation groups (G_1^t, G_2^t) around orbits (O_1, O_2) , we have $\sup_{t \in [0,1]} |\mathbf{T}_\kappa(G_1^t x_1) - \mathbf{T}_\kappa(G_2^t x_2)| = \infty$.*

Proof. Assume not and $d = \lceil \sup_{t \in [0,1]} |\mathbf{T}_\kappa(G_1^t x_1) - \mathbf{T}_\kappa(G_2^t x_2)| \rceil < \infty$. Let $U_n = \{t : \mathbf{T}_\kappa(G^t x_1) > 2^n\}$. Since α_1 is a representation of O_1 , we have $\mathbf{I}(O_1 : \mathcal{H}) <^+ \mathbf{I}((x_1, x_2, \alpha_1, \alpha_2) : \mathcal{H}) < \infty$. Thus by Theorem 78, for all $x \in O$ and $c \in \mathbb{N}$ where

$$2^{-n-2 \log n - c} < L_1\{t \in [0, 1] : \mathbf{T}_\kappa(G^t x) > 2^n\}.$$

So

$$2^{-n-2 \log n - c} < L_1\{t \in [0, 1] : \mathbf{T}_\kappa(G^t x_1) > 2^n\}.$$

Given x_1 and α_1 , one can enumerate an increasing r -interval $v \subseteq U_n$ and stop when $L(v) > L(U_n/2)$. By Lemma 20, there exists a rational $r \in v$, with

$$\mathbf{K}(r) <^{\log} n + \mathbf{I}(r; \mathcal{H}) <^{\log} n + \mathbf{I}((x_1, x_2, \alpha_1, \alpha_2) : \mathcal{H}) + \mathbf{K}(n) <^{\log} n + \mathbf{I}((x_1, x_2, \alpha_1, \alpha_2) : \mathcal{H}).$$

Let $A_n = \{x : \mathbf{T}_\kappa(x) > 2^n\}$. Thus $\kappa(A_n) < 2^{-n}$. Thus one can create the following \mathcal{Z} -test

$$\begin{aligned} t_n(x, y, \alpha, \beta) &= [x \in G_1^{-r} A_n, y \in G_2^{-r} A_{n-d}] 2^{2n-d} \\ \mathbf{K}(t_n) &<^+ \mathbf{K}(r, n) <^{\log} n + \mathbf{I}((x_1, x_2, \alpha_1, \alpha_2) : \mathcal{H}) \\ &<^{\log} n, \end{aligned} \tag{18.3}$$

where Equation 18.3 is due to the fact that $\mathbf{I}((x_1, x_2, \alpha_1, \alpha_2) : \mathcal{H})$ is finite. One can impliment t_n to be a lower computable function in the following manner. As open balls B with center x and radius s are enumerated that are subsets of A_n (or A_{n-d}), the algorithm for t_n rotates them with G_1^t (and also G_2^t), consistently with Claim 2.

As t_n reads the bits of α and β it produces a fast Cauchy sequence $\{x_n\}$ of points that converge to $G^{-r}x$ where $\|x_n - x\| < 2^{-n}$. The algorithm for test t_n creates a sequence of balls $\{B_i\}$, $B_i = \{y : \|y - x_i\| < s - 2^{-i}\}$ such that $B_i \subseteq B_{i+1}$ and $\lim_{i \rightarrow \infty} B_i = G^{-r}B$.

Thus for each $B \supset A_n$, $C \in A_{n-d}$, t_n enumerates two balls B_i and C_i (one for G_1^t and another for G_2^t) and if a point $(y, z) \in B_i \times C_i$, then $t(y, z, \alpha_1, \alpha_2) = 2^{2n-d}$. Note that $\kappa \times \kappa(B_i \times C_i) < 2^{-n}$. The function $t_n \in [\kappa, \mu]$ because

$$\begin{aligned} (\kappa, \mu)(t_n) &= \int_{\alpha, \beta \in \{0,1\}^\infty} \int_{x_1, x_2 \in \mathbb{R}^3} t_n(x_1, x_2, \alpha, \beta) d\kappa(x_1) d\kappa(x_2) d\mu(\alpha) d\mu(\beta) \\ &\leq \sup_{n \rightarrow \infty} \sum_{a, b \in \{0,1\}^n} \mu(a) \mu(b) \int_{x_1, x_2 \in (B^a, C^b)} 2^{2n-d} d\kappa(x_1) d\kappa(x_2) \\ &\leq 2^{2n-d} \sup_{n \rightarrow \infty} \sum_{a, b \in \{0,1\}^n} \mu(a) \mu(b) \int_{x_1, x_2 \in (B^a, C^b)} 2^{2n-d} d\kappa(x_1) d\kappa(x_2) \\ &\leq 2^{2n-d} \sup_{n \rightarrow \infty} \sum_{a, b \in \{0,1\}^n} \mu(a) \mu(b) \kappa(B^a) \kappa(C^b) \\ &\leq 1, \end{aligned}$$

where B^a is the intermediate ball create by th t_n algorithm when just using the bits $a \in \{0, 1\}^*$ (and similarly for C^b). By construction, such balls will have $\kappa(B^a) < 2^{-n}$ and $\kappa(C^b) < 2^{-n+d}$.

It must be that $t_n(x_1, x_2, \alpha_1, \alpha_2) \neq 0$ because $r \in U_n$. Thus

$$\begin{aligned}
\mathbf{T}_{\kappa, \mu}(x_1, x_2, \alpha_1, \alpha_2) &> \sum_n^* \mathbf{m}(t_n | \kappa, \mu) t_n(x_1, x_2, \alpha_1, \alpha_2) \\
&> \sum_n^* \mathbf{m}(n, r_n) t_n(x_1, x_2, \alpha_1, \alpha_2) \\
&> \sum_n^* 2^{-n-O(\log n)} 2^{2n-d} \\
&= \infty,
\end{aligned}$$

causing a contradiction. □

Bibliography

- [BDH⁺12] L. Bienvenu, A. Day, M. Hoyrup, I. Mezhirov, and A. Shen. A constructive version of birkhoff’s ergodic theorem for martin-löf random points. *Information and Computation*, 210:21–30, 2012.
- [BDK⁺05] I Bjelaković, J Deuschel, T Krüger, R Seiler, R Siegmund-Schultze, and A Szkola. A quantum version of sanov’s theorem. *Communications in Mathematical Physics*, 260(3):659–671, 2005.
- [BH96] V. Buzek and M. Hillery. Quantum copying: Beyond the no-cloning theorem. *Phys. Rev. A*, 54:1844–1852, 1996.
- [Bho21] T. Bhojraj. Algorithmic randomness and kolmogorov complexity for qubits. *CoRR*, abs/2106.14280, 2021.
- [BKM⁺06] F. Benatti, T. Krüger, M. Müller, R. Siegmund-Schultze, and A. Szko—a. Entropy and Quantum Kolmogorov Complexity: A Quantum Brudno’s Theorem. *Communications in Mathematical Physics*, 265(2), 2006.
- [BOD14] F. Benatti, S. K. Oskouei, and A. S. Deh Abad. Gacs Quantum Algorithmic Entropy in Infinite Dimensional Hilbert Spaces. *Journal of Mathematical Physics*, 55(8), 2014.
- [Bru78] A. Brudno. The Complexity of the trajectories of a Dynamical System. *Russian Mathematical Surveys*, 33(1):197, 1978.
- [BvL01] A. Berthiaume, W. van Dam, and S. Laplante. Quantum Kolmogorov Complexity. *Journal of Computer and System Sciences*, 63(2), 2001.
- [Deu99] David Deutsch. Quantum theory of probability and decisions. *Proceedings of the Royal Society of London. Series A: Mathematical, Physical and Engineering Sciences*, 455(1988), 1999.
- [DJ13] A. Day and Miller. J. Randomness for non-computable measures. *Transactions of the American Mathematical Society*, 365(7):3575–3591, 2013.
- [Eps19a] S. Epstein. Algorithmic no-cloning theorem. *IEEE Transactions on Information Theory*, 65(9):5925–5930, 2019.
- [Eps19b] S. Epstein. Algorithmic no-cloning theorem. *IEEE Transactions on Information Theory*, 65(9), 2019.
- [Eps19c] S. Epstein. On the algorithmic probability of sets. *CoRR*, abs/1907.04776, 2019.
- [Eps19d] S. Epstein. On the algorithmic probability of sets. *CoRR*, abs/1907.04776, 2019.

- [Eps20] Samuel Epstein. An extended coding theorem with application to quantum complexities. *Information and Computation*, 275, 2020.
- [Eps21a] S. Epstein. On the algorithmic content of quantum measurements. *CoRR*, abs/2102.03905, 2021.
- [Eps21b] Samuel Epstein. All sampling methods produce outliers. *IEEE Transactions on Information Theory*, 67(11):7568–7578, 2021.
- [Eps22] S. Epstein. The outlier theorem revisited. *CoRR*, abs/2203.08733, 2022.
- [Eps23a] S. Epstein. On the algorithmic information between probabilities. *CoRR*, abs/2303.07296, 2023.
- [Eps23b] S. Epstein. A quantum el theorem. *CoRR*, abs/2301.08348, 2023.
- [Eps23c] S. Epstein. A quantum outlier theorem. *CoRR*, abs/2303.06256, 2023.
- [Eve57] Hugh Everett. "relative state" formulation of quantum mechanics. *Rev. Mod. Phys.*, 29, 1957.
- [FMN12] Greenberg N. Franklin, J., J. Miller, and K. Ng. Martin-löf random points satisfy birkhoff's ergodic theorem for effectively closed sets. *Proceedings of the American Mathematical Society*, 140:623–3628, 2012.
- [Gö61] Kurt Gödel. The modern development of the foundations of mathematics in the light of philosophy. In: *Kurt Gödel. Collected Works. Volume III. Oxford University Press.*, 1961.
- [Gó1] P. Gács. Quantum Algorithmic Entropy. *Journal of Physics A Mathematical General*, 34(35), 2001.
- [G21] Peter Gács. Lecture notes on descriptive complexity and randomness. *CoRR*, abs/2105.04704, 2021.
- [Gac94] P. Gacs. The boltzmann entropy and randomness tests. In *Proceedings Workshop on Physics and Computation. PhysComp '94*, pages 209–216, 1994.
- [GHR10] S. Galatolo, M. Hoyrup, and C. Rojas. Computing the speed of convergence of ergodic averages and pseudorandom points in computable dynamical systems. *Electronic Proceedings in Theoretical Computer Science*, 24, 06 2010.
- [GHR11] P. Gács, M. Hoyrup, and C. Rojas. Randomness on computable probability spaces—a dynamical point of view. *Theory of Computing Systems*, 48(3):465–485, 2011.
- [HR09] M. Hoyrup and C. Rojas. Computability of probability measures and martin-löf randomness over metric spaces. *Information and Computation*, 207(7):830–847, 2009.
- [Lev74] L. A. Levin. Laws of Information Conservation (Non-growth) and Aspects of the Foundations of Probability Theory. *Problemy Peredachi Informatsii*, 10(3):206–210, 1974.
- [Lev76] L. A. Levin. Uniform tests of randomness. *Soviet Math. Dokl.*, 17(2):337–340, 1976.
- [Lev84] L. A. Levin. Randomness conservation inequalities; information and independence in mathematical theories. *Information and Control*, 61(1):15–37, 1984.

- [Lev13] L. A. Levin. Forbidden information. *J. ACM*, 60(2), 2013.
- [Lev16] L. A. Levin. Occam bound on lowest complexity of elements. *Annals of Pure and Applied Logic*, 167(10):897–900, 2016. And also: S. Epstein and L.A. Levin, Sets have simple members, arXiv preprint arXiv:1107.1458, 2011.
- [LO97] M. Li and Vitányi O. *An Introduction to Kolmogorov Complexity and Its Applications*. Springer, 1997.
- [Mul08] M. Muller. Strongly Universal Quantum Turing Machines and Invariance of Kolmogorov Complexity. *IEEE Transactions on Information Theory*, 54(2), 2008.
- [NS19] A. Nies and V. Scholz. Martin-Löf Random Quantum States. *Journal of Mathematical Physics*, 60(9), 2019.
- [SBC01] R. Schack, T. Brun, and C. Caves. Quantum bayes rule. *Phys. Rev. A*, 64:014305, 2001.
- [SBKW10] S. Saunders, J. Barrett, A. Kent, and D. Wallace. *Many Worlds?: Everett, Quantum Theory, & Reality*. OUP Oxford, 2010.
- [Sch71] C. Schnorr. A unified approach to the definition of random sequences. *Mathematical systems theory*, 5(3), 1971.
- [Sch95] B. Schumacher. Quantum coding. *Phys. Rev. A*, 51, 1995.
- [She15] M. Shell. *Thermodynamics and Statistical Mechanics: An Integrated Approach*. Cambridge Series in Chemical Engineering. Cambridge University Press, 2015.
- [Vai98] L. Vaidman. On schizophrenic experiences of the neutron or why we should believe in the many-worlds interpretation of quantum theory. *International Studies in the Philosophy of Science*, 12(3):245–261, 1998.
- [Ver21] N. Vereshchagin. Proofs of conservation inequalities for levin’s notion of mutual information of 1974. *Theoretical Computer Science*, 856, 2021.
- [Vit00] P. Vitányi. Three Approaches to the Quantitative Definition of Information in an Individual Pure Quantum State. In *Proceedings of the 15th Annual IEEE Conference on Computational Complexity, COCO ’00*, page 263. IEEE Computer Society, 2000.
- [Vit01] P.M.B. Vitanyi. Quantum kolmogorov complexity based on classical descriptions. *IEEE Transactions on Information Theory*, 47(6):2464–2479, 2001.
- [Wil13] M. Wilde. *Quantum Information Theory*. Cambridge University Press, 2013.
- [WZ82] W. Wootters and W. Zurek. A single quantum cannot be cloned. *Nature*, 299(5886):802–803, 1982.

Part V

Appendix

Appendix A

Information Between Basis States

Special information inequalities and conservation inequalities can be achieved for orthogonal sequences of pure basis states $|1\rangle, |2\rangle, |3\rangle, \dots$. We use Theorem 8 that $\mathbf{K}(i|n) = {}^+ \mathbf{Hg}(|i\rangle)$. Let $S(\rho)$ be the von Neumann entropy of σ .

Theorem 80 *Relativized to an orthogonal sequence of elementary states $|1\rangle, |2\rangle, |3\rangle, \dots$, enumerated by strings $i, j, k \in \{0, 1\}^n$.*

1. $\mathbf{d}(|i\rangle | |j\rangle) = \infty$ for $i \neq j$.
2. $\mathbf{Hg}(\sigma \otimes |i\rangle \langle i|) = {}^+ \mathbf{Hg}(|i\rangle) + \mathbf{Hg}(\sigma | |i\rangle, \mathbf{Hg}(|i\rangle))$.
3. $\mathbf{I}(|k\rangle : |i\rangle) < {}^+ \mathbf{I}(|j\rangle : |i\rangle) + \mathbf{K}(k|j, N)$.
4. $\mathbf{I}(i : j|N) < {}^{\log} \mathbf{I}(|i\rangle : |j\rangle)$.
5. $\mathbf{I}(|i\rangle : |j\rangle) < {}^+ \mathbf{I}(i : j|N) + \mathbf{I}(i, j : \mathcal{H}|N)$.
6. $\mathbf{K}(i|N) < {}^+ \mathbf{I}(|i\rangle : |i\rangle) < {}^+ \mathbf{K}(i|N) + \mathbf{I}(i : \mathcal{H}|N)$.
7. $\mathbf{I}(|i\rangle : |i\rangle) < {}^+ 4n|3$.
8. $\mathbf{I}(|i\rangle : |j\rangle) < {}^+ \mathbf{I}(|i\rangle : |i\rangle) + \mathbf{I}(i, j : \mathcal{H}|n)$.

Proof.

(1) This is due to the fact that $\mathfrak{T}_{|i\rangle \langle i|} >^* \sum_n \mathbf{m}(n)n |j\rangle \langle j|$. Thus $\log \text{Tr} \mathfrak{T}_{|i\rangle \langle i|} |j\rangle \langle j| = \infty$.

(2) We use the lower semicomputable matrix $\rho = \mu_{(|i\rangle, \mathbf{Hg}(|i\rangle))} \otimes |i\rangle \langle i|$. so we have that $\mu_{2n} >^* \mathbf{m}(|i\rangle, \mathbf{Hg}(|i\rangle) | 2^{2n}) \rho \stackrel{*}{=} \mathbf{m}(i, \mathbf{K}(i|n) | 2^{2n}) \rho \stackrel{*}{=} \mathbf{m}(i | 2^n) \rho \stackrel{*}{=} 2^{-\mathbf{Hg}(|i\rangle)} \rho$. So $\mathbf{Hg}(\sigma \otimes |i\rangle \langle i|) < {}^+ \mathbf{Hg}(|i\rangle) - \log \text{Tr} \rho(\sigma \otimes |i\rangle \langle i|) < {}^+ \mathbf{Hg}(|i\rangle) + \mathbf{Hg}(\sigma | |i\rangle, \mathbf{Hg}(|i\rangle))$. The other direction is given by Theorem 12.

(3) We let $T \in \mathcal{T}_{\mu \otimes \mu}$ vary over $\mu \otimes \mu$ tests. Since $\mu >^* \sum_i |i\rangle \mathbf{m}(i|N) \langle i|$ it must be that $1 > \text{Tr} T \mu \otimes \mu >^* \text{Tr} T \sum_{i,j} \mathbf{m}(i|N) \mathbf{m}(j|N) |i\rangle |j\rangle \langle i| \langle j| >^* \sum_{i,j} \mathbf{m}(i|N) \mathbf{m}(j|N) \langle i| \langle j| T |i\rangle |j\rangle$.

From each T , let $T' = \sum_{i,j} (\sum_k \mathbf{m}(k|j, N) \langle i| \langle k| T | i \rangle | k \rangle) | i \rangle | j \rangle \langle i| \langle j|$. Since \mathbf{m} and T are lower computable, then so is T' . In addition, $O(1)T' \in \mathcal{T}_{\mu \otimes \mu}$, because

$$\begin{aligned}
\text{Tr} T' \mu \otimes \mu &= \sum_{i,j} \langle i| \langle j| \mu \otimes \mu | i \rangle | j \rangle \sum_k \mathbf{m}(k|j, N) (\langle i| \langle k| T | i \rangle | k \rangle) \\
&\stackrel{*}{=} \sum_{i,j} \mathbf{m}(i|N) \mathbf{m}(j|N) \sum_k \mathbf{m}(k|j, N) (\langle i| \langle k| T | i \rangle | k \rangle) \\
&< \sum_{i,j} \mathbf{m}(i|N) \sum_k \mathbf{m}(j, k|N) (\langle i| \langle k| T | i \rangle | k \rangle) \\
&\stackrel{*}{=} \sum_{i,j} \mathbf{m}(i|N) \sum_k \mathbf{m}(k|N) \mathbf{m}(j|k, \mathbf{K}(k), N) (\langle i| \langle k| T | i \rangle | k \rangle) \\
&\stackrel{*}{=} \sum_{i,k} \mathbf{m}(i|N) \mathbf{m}(k|N) \sum_j \mathbf{m}(j|k, \mathbf{K}(k), N) \langle i| \langle k| T | i \rangle | k \rangle \\
&< \sum_{i,k} \mathbf{m}(i|N) \mathbf{m}(k|N) (\langle i| \langle k| T | i \rangle | k \rangle) \\
&\stackrel{*}{<} \text{Tr} T \mu \otimes \mu \\
&< O(1).
\end{aligned}$$

So

$$\begin{aligned}
\mathbf{I}(|i\rangle : |j\rangle) &> \log \text{Tr} \sum_T \mathbf{m}(O(1)T' | N^2) O(1)T' | i \rangle | j \rangle \langle i| \langle j| \\
&>^+ \log \text{Tr} \sum_T \mathbf{m}(T | N^2) T' | i \rangle | j \rangle \langle i| \langle j| \\
&>^+ \log \sum_T \mathbf{m}(T | N^2) \sum_k \mathbf{m}(k|j, N) \langle k| \langle i| T | k \rangle | i \rangle \\
&>^+ \log \sum_T \mathbf{m}(T | N^2) \mathbf{m}(k|j, N) \langle k| \langle i| T | k \rangle | i \rangle \\
&=^+ \mathbf{I}(|k\rangle : |i\rangle) - \mathbf{K}(k|j).
\end{aligned}$$

(4) This follow as a special case of Theorem 34.

(5) Let $s(i, j) = \mathbf{m}(i|N) \mathbf{m}(j|N) 2^{\mathbf{I}(|i\rangle : |j\rangle)}$. The function s is lower semicomputable relative to \mathcal{H} because \mathbf{m} and $\mathfrak{T}_{\mu \otimes \mu}$ are lower computable relative to \mathcal{H} . Furthermore we have that

$$\begin{aligned}
\sum_{i,j} s(i, j) &= \sum_{i,j} \mathbf{m}(i|N) \mathbf{m}(j|N) \text{Tr} \mathfrak{T}_{\mu \otimes \mu} | i \rangle \langle i| \otimes | j \rangle \langle j| \\
&= \text{Tr} \mathfrak{T}_{\mu \otimes \mu} \sum_{i,j} \mathbf{m}(i|N) | i \rangle \langle i| \otimes \mathbf{m}(j|N) | j \rangle \langle j| \\
&< O(1) \text{Tr} \mathfrak{T}_{\mu \otimes \mu} \mu \otimes \mu < O(1).
\end{aligned}$$

Therefore $s(i, j) \stackrel{*}{<} \mathbf{m}(i, j|N, \mathcal{H})$ and so $\mathbf{I}(|i\rangle : |j\rangle) <^+ \log \mathbf{m}(i, j|N, \mathcal{H}) | (\mathbf{m}(i|N) \mathbf{m}(j|N)) =^+ \mathbf{I}(i : j|N) + \mathbf{I}(i, j : \mathcal{H}|N)$.

(6) For $\mathbf{K}(i|N) <^+ \mathbf{I}(|i\rangle : |i\rangle)$, we prove the stronger statement: for elementary ρ , $2\mathbf{Hg}(\rho) - \mathbf{K}(\rho, \mathbf{Hg}(\rho)) - 2S(\rho) <^+ \mathbf{I}(\rho : \rho)$. Let $\nu = 2^{2\mathbf{Hg}(\rho)-2}(\rho \otimes \rho)$. The matrix $\nu \in \mathcal{T}_{\mu \otimes \mu}$ because

$\text{Tr}(\boldsymbol{\mu} \otimes \boldsymbol{\mu})\nu \leq 1$. Therefore

$$\begin{aligned}
\mathbf{I}(\rho : \rho) &= \log \text{Tr} \mathfrak{T}_{\boldsymbol{\mu} \otimes \boldsymbol{\mu}}(\rho \otimes \rho) \\
&\geq \log \mathbf{m}(\nu) \text{Tr} \nu(\rho \otimes \rho) \\
&>^+ \log \mathbf{m}(\nu) 2^{2\mathbf{Hg}(\rho)} \text{Tr}(\rho \rho \otimes \rho \rho) \\
&>^+ 2\mathbf{Hg}(\rho) - \mathbf{K}(\rho, \mathbf{Hg}(\rho)) + 2 \log \sum_i \lambda_i^2,
\end{aligned}$$

where λ_i are the eigenvalues of ρ . Due to concavity $-2S(\rho) \leq 2 \log \sum_i \lambda_i^2$. So $\mathbf{I}(\rho : \rho) >^+ 2\mathbf{Hg}(\rho) - \mathbf{K}(\rho, \mathbf{Hg}(\rho)) - 2S(\rho)$. The inequality follows from $\mathbf{Hg}(|i\rangle) =^+ \mathbf{K}(i)$ and $S(|i\rangle \langle i|) = 0$. For $\mathbf{I}(|i\rangle : |i\rangle) <^+ \mathbf{K}(i) + \mathbf{I}(i : \mathcal{H}|n)$, we note that it is a special case of (5).

(7) If $T \in \mathcal{T}_{\boldsymbol{\mu}_n \otimes \boldsymbol{\mu}_n}$, then $\text{Tr} T <^* 2^{2n}$. This is because $1 \geq \text{Tr} T(\boldsymbol{\mu}_n \otimes \boldsymbol{\mu}_n) >^* \text{Tr} T(2^{-n} I \otimes 2^{-n} I)$. Since the set of lower computable matrices of trace not more than 2^{2n} is enumerable, $2^{-2n} \mathfrak{T}_{\boldsymbol{\mu}_n \otimes \boldsymbol{\mu}_n} <^* \boldsymbol{\mu}_{2n}$. Assume $\mathbf{I}(|i\rangle : |i\rangle) = \log \text{Tr} \mathfrak{T}_{\boldsymbol{\mu}_n \otimes \boldsymbol{\mu}_n} |ii\rangle \langle ii| = 2n - c$. Then $-\log \text{Tr} u_{2n} |ii\rangle \langle ii| <^+ c$. This means that $\mathbf{K}(ii|2^{2n}) <^+ c$. So $\mathbf{K}(i|2^n) <^+ c$. So $1 \geq \text{Tr} \mathfrak{T}_{\boldsymbol{\mu}_n \otimes \boldsymbol{\mu}_n} \boldsymbol{\mu} \otimes \boldsymbol{\mu} >^* \mathbf{m}(i|2^n)^2 \text{Tr} \mathfrak{T}_{\boldsymbol{\mu}_n \otimes \boldsymbol{\mu}_n} |ii\rangle \langle ii| >^* 2^{2n-3c}$. Thus $c >^+ 2n/3$. This implies that $\mathbf{I}(|i\rangle : |i\rangle) = 2n - c <^+ 4n/3$.

(8) This follows from (5) and (6).

Corollary 40 For elementary ρ , $2\mathbf{Hg}(\rho) - \mathbf{K}(\rho, \mathbf{Hg}(\rho)) - 2S(\rho) <^+ \mathbf{I}(\rho : \rho)$.

Appendix B

An Extended Coding Theorem

In [Lev16, Eps19c], a new inequality in the field of algorithmic information theory was proven. For a finite set of natural numbers D , it was shown that the size of the smallest description of an element of D , $\min_{x \in D} \mathbf{K}(x)$, is not much smaller than the negative logarithm of the algorithmic probability of the set, $-\log \sum_{x \in D} \mathbf{m}(x)$. This inequality holds for non-exotic sets whose encoding has little mutual information with the halting sequence, $\mathbf{I}(D; \mathcal{H}) = \mathbf{K}(D) - \mathbf{K}(D|\mathcal{H})$.

$$\min_{x \in D} \mathbf{K}(x) <^{\log} -\log \sum_{x \in D} \mathbf{m}(x) + \mathbf{I}(D; \mathcal{H}).$$

Due to algorithmic conservation laws, there are no algorithmic means to produce sets with arbitrary high mutual information with the halting sequence. In this appendix, we introduce an update on the above inequality, proving for non-exotic maps f between whole numbers with a finite domain, $\min_{x \in \text{Dom}(f)} \mathbf{K}(x) + f(x)$ is close to the amount $-\log \sum_{x \in \text{Dom}(f)} \mathbf{m}(x) 2^{-f(x)}$. Exotic maps f have encodings with high mutual information with the halting sequence, $\mathbf{I}(f; \mathcal{H})$, with

$$\min_{x \in \text{Dom}(f)} \mathbf{K}(x) + f(x) <^{\log} -\log \sum_{x \in \text{Dom}(f)} \mathbf{m}(x) 2^{-f(x)} + \mathbf{I}(f; \mathcal{H}).$$

The above inequality can be seen as an extended coding theorem.

B.1 Left-Total Machines

The notion of the “left-total” universal algorithm is needed for the proof of both the mixed state and pure state coding theorems. We say $x \in \{0, 1\}^*$ is total with respect to a machine if the machine halts on all sufficiently long extensions of x . More formally, x is total with respect to T_y for some $y \in \{0, 1\}^{*\infty}$ iff there exists a finite prefix free set of strings $Z \subset \{0, 1\}^*$ where $\sum_{z \in Z} 2^{-\|z\|} = 1$ and $T_y(xz) \neq \perp$ for all $z \in Z$. We say (finite or infinite) string $\alpha \in \{0, 1\}^{*\infty}$ is to the “left” of $\beta \in \{0, 1\}^{*\infty}$, and use the notation $\alpha \triangleleft \beta$, if there exists a $x \in \{0, 1\}^*$ such that $x0 \sqsubseteq \alpha$ and $x1 \sqsubseteq \beta$. A machine T is left-total if for all auxiliary strings $\alpha \in \{0, 1\}^{*\infty}$ and for all $x, y \in \{0, 1\}^*$ with $x \triangleleft y$, one has that $T_\alpha(y) \neq \perp$ implies that x is total with respect to T_α . An example can be seen in Figure B.1.

For the remaining part of this chapter, we can and will change the universal self delimiting machine U into a universal left-total machine U' by the following definition. The algorithm U' enumerates all strings $p \in \{0, 1\}^*$ in order of their convergence time of $U(p)$ and successively assigns them consecutive intervals $i_p \subset [0, 1]$ of width $2^{-\|p\|}$. Then U' outputs $U(p)$ on input p' if the open interval corresponding to p' and not that of $(p')^-$ is strictly contained in i_p . The open interval in




Figure B.1: The above diagram represents the domain of a left total machine T with the 0 bits branching to the left and the 1 bits branching to the right. For $i \in \{1..5\}$, $x_i \triangleleft x_{i+1}$ and $x_i \triangleleft y$. Assuming $T(y)$ halts, each x_i is total. This also implies each x_i^- is total as well.

$[0,1]$ corresponding with p' is $([p']2^{-\|p'\|}, ([p'] + 1)2^{-\|p'\|})$ where $[p]$ is the value of p in binary. For example, the value of both strings 011 and 0011 is 3. The value of 0100 is 4. The same definition applies for the machines U'_α and U_α , over all $\alpha \in \{0,1\}^{*\infty}$. We now set U to equal U' .




Figure B.2: The above diagram represents the domain of the universal left-total algorithm U , with the 0 bits branching to the left and the 1 bits branching to the right. The strings in the above diagram, $0v0$ and $0v1$, are halting inputs to U with $U(0v0) \neq \perp$ and $U(0v1) \neq \perp$. So $0v$ is a total string. The infinite border sequence $\mathcal{B} \in \{0,1\}^\infty$ represents the unique infinite sequence such that all its finite prefixes have total and non total extensions. All finite strings branching to the right of \mathcal{B} will cause U to diverge.

Without loss of generality, the complexity terms of Chapter 1 are defined in this section with respect to the universal left total machine U . The infinite border sequence $\mathcal{B} \in \{0,1\}^\infty$ represents the unique infinite sequence such that all its finite prefixes have total and non total extensions. The term “border” is used because for any string $x \in \{0,1\}^*$, $x \triangleleft \mathcal{B}$ implies that x total with respect to

U and $\mathcal{B} \triangleleft x$ implies that U will never halt when given x as an initial input. Figure B.2 shows the domain of U with respect to \mathcal{B} . The border sequence is computable from \mathcal{H} .

For all total strings $b \in \{0, 1\}^*$, we define the semimeasure $\mathbf{m}_b(x) = \sum \{2^{-\|p\|} : U(p) = x, p \triangleleft b \text{ or } b \sqsubseteq p\}$. If b is not total then $\mathbf{m}_b(x) = \perp$ is undefined. Thus the algorithmic weight \mathbf{m}_b of a string x is approximated using programs that either extend b or are to the left of b .

B.2 Extended Coding Theorem

Lemma 27 *Let f be a elementary map and m be a elementary semi measure. Let $a \in \mathbb{W}$ vary over $\text{Dom}(f)$. Then $\min_a f(a) + \mathbf{K}(a|m) <^{\log} -\log \sum_a m(a)2^{-f(a)} + \mathbf{Ks}(f|m)$.*

Proof. If m is not a proper probability measure, and R is the support of m , we modify m to give an arbitrary $b \in \mathbb{W}$, the value of $1 - m(R)$. So m can be assumed to be an elementary probability measure. Since all terms in the theorem are conditioned on $\langle m \rangle$, we will also condition all complexity terms in the proof on $\langle m \rangle$ and drop its notation. More formally, $U(x)$ is used to denote $U_{\langle m \rangle}(x)$, $\mathbf{K}(x)$ is used to denote $\mathbf{K}(x|m)$, and $\mathbf{Ks}(f)$ is used to denote $\mathbf{Ks}(f|m)$.

For any elementary map g , let $g_n = g^{-1}(n) \cap \text{Supp}(m)$ and let $g_{\leq n} = \cup_{i=0}^n g_i$, for $n \in \mathbb{W} \cup \{\infty\}$. Let $s = \lceil -\log \sum_{a \in f_{\leq \infty}} m(a)2^{-f(a)} \rceil$. Using the reasoning of Markov's inequality,

$$\sum_{a \in f_{\leq \infty}} m(a)2^{-f(a)} \geq 2^{-s}, \quad (\text{B.1})$$

$$\sum_{a \in f_{\leq \infty} \setminus f_{\leq s}} m(a)2^{-f(a)} \leq \sum_{a \in f_{\leq \infty} \setminus f_{\leq s}} m(a)2^{-s-1} \leq 2^{-s-1}, \quad (\text{B.2})$$

$$\sum_{a \in f_{\leq s}} m(a)2^{-f(a)} \geq 2^{-s-1}. \quad (\text{B.3})$$

Equation (B.1) follows from the definition of s and Equation (B.3) follows from Equations (B.1) and (B.2). We now turn our attention to creating an elementary probability measure Q with the following properties:

1. f is typical of Q and Q is simple, i.e. there is a $v \in \{0, 1\}^*$ with $U(v) = \langle Q \rangle$ and $\|v\| + 3 \log \max\{\mathbf{d}(f|Q, v), 1\}$ is not much larger than $\mathbf{Ks}(f)$.
2. All strings in the support of Q encode elementary functions g whose range contain a lot of values that are not greater than s , with $\sum_{a \in g_{\leq s}} m(a)2^{-g(a)} \geq 2^{-s-1}$.

To accomplish this goal, we start with the program $v' \in \{0, 1\}^*$ and elementary probability measure Q' that realizes the stochasticity of f , with $U(v') = \langle Q' \rangle$, and also with the relation $\mathbf{Ks}(f) = \|v'\| + 3 \log \max\{\mathbf{d}(f|Q', v'), 1\}$. Note that this implies $\langle f \rangle \in \text{Supp}(Q')$. Let Q be the elementary probability measure equal to Q' conditioned on the set of (encoded) elementary maps g such that $\sum_{a \in g_{\leq s}} m(a)2^{-g(a)} \geq 2^{-s-1}$. Thus $Q(\langle g \rangle) = [g \in S]Q'(g)/Q'(S)$, where $S \subset \{0, 1\}^*$, the support of Q , is defined as $S = \{\langle g \rangle : g \in \text{Supp}(Q'), \sum_{a \in g_{\leq s}} m(a)2^{-g(a)} \geq 2^{-s-1}\}$. This Q is computable from v' and s . Using this fact, define the Q program $v \in \{0, 1\}^*$, to be of the form $v = v_0 v_s v'$, where $v_0 \in \{0, 1\}^*$ is helper code of size $O(1)$, and $v_s \in \{0, 1\}^*$ is a shortest U -program

for s . So $\|v\| <^+ \|v'\| + \mathbf{K}(s)$. We define $d = \max\{\mathbf{d}(f|Q, v), 1\}$ and we have that

$$\begin{aligned}
& \|v\| <^+ \|v'\| + \mathbf{K}(s), \\
& \|v\| + 3 \log d <^+ \|v'\| + \mathbf{K}(s) + 3 \log d \\
& \quad <^+ \|v'\| + \mathbf{K}(s) + 3 \log(\max\{-\log Q(f) - \mathbf{K}(f|v), 1\}) \\
& \quad <^+ \|v'\| + \mathbf{K}(s) + 3 \log(\max\{-\log Q'(f) - \mathbf{K}(f|v), 1\}) \tag{B.4} \\
& \quad <^+ \|v'\| + \mathbf{K}(s) + 3 \log(\max\{-\log Q'(f) - \mathbf{K}(f|v') + \mathbf{K}(v|v'), 1\}) \tag{B.5} \\
& \quad <^+ \|v'\| + \mathbf{K}(s) + 3 \log(\max\{-\log Q'(f) - \mathbf{K}(f|v') + \mathbf{K}(s), 1\}) \tag{B.6} \\
& \quad <^{\log} \|v'\| + \mathbf{K}(s) + 3 \log(\max\{-\log Q'(f) - \mathbf{K}(f|v'), 1\}), \\
& \|v\| + 3 \log d <^{\log} \mathbf{K}s(f) + \mathbf{K}(s). \tag{B.7}
\end{aligned}$$

Equation (B.4) follows from $Q(f) = Q'(f)/Q'(\text{Supp}(Q))$, and thus $-\log Q(f) \leq -\log Q'(f)$. Equation (B.5) follows from the inequality $\mathbf{K}(f|v') <^+ \mathbf{K}(f|v) + \mathbf{K}(v|v')$. Equation (B.6) follows from v being computable from v' and v_s , and thus $\mathbf{K}(v|v') <^+ \mathbf{K}(s)$.

We now create a small set of lists of numbers A that will intersect with the range of a large percentage of the support of Q . We do so by using the probabilistic method. Let $c \in \mathbb{N}$ be a constant solely dependent on the universal Turing machine U to be determined later. We use an elementary measure w_n over lists A^n of (possibly repeating) whole numbers of size $cd2^{s+1-n}$ where $w_n(A^n) = \prod_{i=1}^{cd2^{s+1-n}} m(A_i^n)$. For a set of $s+1$ lists $A = \{A^n\}_{n=0}^s$, we a measure w over A , where $w(A) = \prod_{n=0}^s w_n(A^n)$.

For a set of lists A and elementary function g , let $\mathbf{1}(g, A) = 1$ if $g_n \cap A^n = \emptyset$ for all $n \in [0, s]$, and $\mathbf{1}(g, A) = 0$, otherwise. Thus

$$\begin{aligned}
\mathbf{E}_{g \sim Q} \mathbf{E}_{A \sim w} [\mathbf{1}(g, A)] &= \sum_g Q(g) \prod_{n=0}^s (1 - m(g_n))^{|A^n|} \\
&\leq \sum_g Q(g) \prod_{n=0}^s \exp\{-|A^n| m(g_n)\} \tag{B.8}
\end{aligned}$$

$$\begin{aligned}
&= \sum_g Q(g) \exp\left\{-\sum_{n=0}^s |A^n| m(g_n)\right\} \\
&= \sum_g Q(g) \exp\left\{-\sum_{n=0}^s cd2^{s+1-n} m(g_n)\right\} \\
&= \sum_g Q(g) \exp\left\{-cd2^{s+1} \sum_{n=0}^s m(g_n) 2^{-n}\right\} \\
\mathbf{E}_{g \sim Q} \mathbf{E}_{A \sim \lambda} [\mathbf{1}(g, A)] &\leq \sum_g Q(g) \exp\{-cd\} = \exp\{-cd\}. \tag{B.9}
\end{aligned}$$

Equation (B.8) follows from the inequality $(1-a) \leq e^{-a}$ over $a \in [0, 1]$. Equation (B.9) follows from the definition of the support of Q , where $g \in \text{Supp}(Q)$ iff $\sum_{a \in g \leq s} m(a) 2^{-g(a)} \geq 2^{-s-1}$. By the probability argument, there exists a set of lists $A = \{A^n\}_{n=0}^s$ such that $|A^n| = cd2^{s+1-n}$ and

$$\mathbf{E}_{g \sim Q} [\mathbf{1}(g, A)] \leq \exp\{-cd\}.$$

There exists a brute force search algorithm that on input c, d, v , outputs A . Note that the strings s and $\langle Q \rangle$ are computable from v . This algorithm computes all possible sets of lists $A' = \{A'^n\}_{n=0}^s$, $|A'^n| = cd2^{s+1-n}$, $A'^n \subseteq \text{Supp}(Q)$ and outputs the first A' such that $\mathbf{E}_{g \sim Q}[\mathbf{1}(g, A')] \leq \exp\{-cd\}$. The existence of such an A' is guaranteed by Equation (B.9). So

$$\mathbf{K}(A) <^+ \mathbf{K}(c, d, v). \quad (\text{B.10})$$

We now show that there is an n where $f_n \cap A^n \neq \emptyset$. To do so, we show that any function g in the support of Q whose range does not intersect with A , i.e. $\mathbf{1}(g, A) = 1$ will have a very high deficiency of randomness with respect to Q and v . For all such g and proper choice of c solely dependent on U ,

$$\begin{aligned} \mathbf{d}(g|Q, v) &= \lfloor -\log Q(g) \rfloor - \mathbf{K}(g|v) \\ &> -\log Q(g) - (-\log \mathbf{1}(g, A) \lfloor e^{cd} \rfloor Q(g) + \mathbf{K}(\mathbf{1}(\cdot, A) \lfloor e^{cd} \rfloor Q(\cdot)|v)) - O(1) \end{aligned} \quad (\text{B.11})$$

$$\begin{aligned} &> cd \log e - \mathbf{K}(\mathbf{1}(\cdot, A) \lfloor e^{cd} \rfloor Q(\cdot)|v) - O(1) \\ &> cd \log -\mathbf{K}(A, c, d|v) - O(1) \\ &> cd \log e - \mathbf{K}(c, d) > d. \end{aligned} \quad (\text{B.12})$$

With c being chosen, it is removed from consideration for the rest of the proof, with $c \in \emptyset(1)$. Equation B.11 is due to the fact that for any elementary semimeasure P , $\mathbf{K}(x) <^+ \mathbf{K}(P) - \log P(x)$. Equation B.12 is due to Equation B.10. So $\mathbf{1}(f, A) = 0$, otherwise by the above equation, $\mathbf{d}(f|Q, v) > d$, causing a contradiction. So there exists $n \in [0, s]$ with $a \in f_n \cap A^n$ and

$$\begin{aligned} \mathbf{K}(a) &<^+ \log |A^n| + \mathbf{K}(A^n) \\ &<^+ \log |A^n| + \mathbf{K}(A) + \mathbf{K}(A^n|A) \\ &<^+ (\log d + s - n) + \mathbf{K}(d, v) + \mathbf{K}(n) \\ &=^+ \log d + s - f(a) + \mathbf{K}(d, v) + \mathbf{K}(f(a)) \end{aligned} \quad (\text{B.13})$$

$$\begin{aligned} \mathbf{K}(a) + f(a) &<^+ \log d + s + \mathbf{K}(v) + \mathbf{K}(d) + \mathbf{K}(f(a)) \\ \mathbf{K}(a) + f(a) &<^{\log} s + \|v\| + 3 \log d \end{aligned} \quad (\text{B.14})$$

$$\mathbf{K}(a) + f(a) <^{\log} s + \mathcal{H}(f) \quad (\text{B.15})$$

$$\min_{a \in f_{\leq \infty}} \mathbf{K}(a) + f(a) <^{\log} -\log \sum_{a \in f_{\leq \infty}} m(a) 2^{-f(a)} + \mathcal{H}(f). \quad (\text{B.16})$$

Equation (B.13) follows from Equation (B.10), and from $c \in O(1)$. Equation (B.14) follows from $\mathbf{K}(x) <^{\log} \|x\|$ for $x \in \{0, 1\}^* \cup \mathbb{W}$. Equation (B.15) follows directly from Equation (B.7). Equation (B.16) follows from the definition of s and its form proves the theorem.

Proposition 35 For border prefix $b \sqsubseteq \mathcal{B}$, $\mathbf{K}(b|\mathcal{H}) <^+ \mathbf{K}(\|b\|)$ and $\|b\| <^+ \mathbf{K}(b)$.

Proof. The border \mathcal{B} is computable from the halting sequence \mathcal{H} , so it follows easily $\mathbf{K}(b|\mathcal{H}) <^+ \mathbf{K}(\|b\|)$. We recall that $\Omega = \sum_x \mathbf{m}(x)$ is Chaitin's Omega, the probability that U will halt. It is well known that the binary expansion $\Omega' \in \{0, 1\}^\infty$ of Ω is Martin L f random. Given $b \sqsubset \mathcal{B}$, $\|b\| \in \{0, 1\}^n$, one can compute $\hat{\Omega} = \sum \{2^{-\|y\|} [U(y) \neq \perp] : y \triangleleft b\}$ with differs from Ω in the summation of programs which branch from \mathcal{B} at positions $n+1$ or higher. Thus $\Omega - \hat{\Omega} \leq 2^{-n}$. So $n <^+ \mathbf{K}(\Omega'[0..n-1]) <^+ \mathbf{K}(\Omega'[0..n-1], b) <^+ \mathbf{K}(\Omega'[0..n-1]|b) + \mathbf{K}(b) <^+ \mathbf{K}(b)$.

Proposition 36 If $b \in \{0, 1\}^*$ is total and b^- is not total, then b^- is a border prefix, with $b^- \sqsubset \mathcal{B}$.

\mathcal{B} (see Figure B.3). In addition, Q is computable from v . Therefore

$$\begin{aligned} \mathbf{K}(x|\mathcal{H}) &<^+ \mathbf{K}(x|Q) + \mathbf{K}(Q|\mathcal{H}) \\ &<^+ \mathbf{K}(x|Q) + \mathbf{K}(v|\mathcal{H}) \\ &<^+ -\log Q(x) + \mathbf{K}(\|v\|) \end{aligned} \tag{B.18}$$

$$\begin{aligned} &<^+ \mathbf{K}(x) - \|v\| + \mathbf{K}(\|v\|), \\ \|v\| &<^+ \mathbf{K}(x) - \mathbf{K}(x|\mathcal{H}) + \mathbf{K}(\|v\|), \\ \|v\| &<^{\log} \mathbf{I}(x; \mathcal{H}). \end{aligned} \tag{B.19}$$

Equation (B.18) is due to Proposition (35). Since Q is computable from v , one gets $\mathbf{Ks}(x) <^+ \mathbf{K}(v) + 3 \log(\max\{\mathbf{d}(x|Q, v), 1\}) <^+ \|v\| + \mathbf{K}(\|v\|) + 3 \log(\max\{\mathbf{d}(x|Q, v), 1\})$. Due to Equation B.17, one gets $\mathbf{Ks}(x) \leq \|v\| + O(\mathbf{K}(\|v\|)) <^{\log} \|v\|$. Due to Equation B.19, one gets $\mathbf{Ks}(x) <^{\log} \mathbf{I}(x; \mathcal{H})$.

Theorem 81 For elementary map f , $\min_{a \in \text{Dom}(f)} f(a) + \mathbf{K}(a) <^{\log} -\log \sum_{a \in \text{Dom}(f)} \mathbf{m}(a) 2^{-f(a)} + \mathbf{I}(\langle f \rangle; \mathcal{H})$.

Proof. Let $s = \lceil 1 - \log \sum_{a \in \text{Dom}(f)} \mathbf{m}(a) 2^{-f(a)} \rceil$ and let $S(z) = \lceil -\log \sum_{a \in \text{Dom}(f)} \mathbf{m}_z(a) 2^{-f(a)} \rceil$ be a partial recursive function from strings to rational numbers. S is defined solely on total strings, where $S(z) \neq \perp$ iff z is total. For total strings z, z^- , one has that $\mathbf{m}_{z^-}(x) \geq \mathbf{m}_z(x)$ and therefore $S(z^-) \leq S(z)$. Let b be the shortest total string with the property that $S(b) < s$. This implies $S(b^-) = \perp$ and thus b^- is not total. So by proposition (36), $b^- \sqsubseteq \mathcal{B}$ is a prefix of border. Lemma 27, with U containing b on an auxilliary tape, with $m(a) = \mathbf{m}_b(a)$, provides $a \in \mathbb{W}$ such that $\mathbf{K}(a|m, b) + f(a) <^{\log} s + \mathbf{Ks}(f|m, b)$. Since $\mathbf{K}(m|b) = O(1)$, we have Equation (B.20). Lemma (28), conditional on b , results in Equation (B.21), with

$$\mathbf{K}(a|b) + f(a) <^{\log} s + \mathbf{Ks}(f|b), \tag{B.20}$$

$$\mathbf{K}(a|b) + f(a) <^{\log} s + \mathbf{I}(f; \mathcal{H}|b), \tag{B.21}$$

$$\mathbf{K}(a|b) + f(a) <^{\log} s + \mathbf{K}(f|b) - \mathbf{K}(f|b, \mathcal{H}). \tag{B.22}$$

Using the fact that $\mathbf{K}(a) <^+ \mathbf{K}(a|b) + \mathbf{K}(b)$, we get $\mathbf{K}(a) - \mathbf{K}(b) <^+ \mathbf{K}(a|b)$, and combined with Equation (B.22), we get Equation (B.23). Equation (B.24) is due to the chain rule $\mathbf{K}(b) + \mathbf{K}(f|b) <^{\log} \mathbf{K}(f) + \mathbf{K}(b|f)$. Equation (B.25) follows from the inequality $\mathbf{K}(f|\mathcal{H}) <^+ \mathbf{K}(f|b, \mathcal{H}) + \mathbf{K}(b|\mathcal{H})$.

$$\mathbf{K}(a) + f(a) <^{\log} s + \mathbf{K}(b) + \mathbf{K}(f|b) - \mathbf{K}(f|b, \mathcal{H}), \tag{B.23}$$

$$\mathbf{K}(a) + f(a) <^{\log} s + \mathbf{K}(f) + \mathbf{K}(b|f) - \mathbf{K}(f|b, \mathcal{H}), \tag{B.24}$$

$$\mathbf{K}(a) + f(a) <^{\log} s + \mathbf{K}(f) + \mathbf{K}(b|f) - \mathbf{K}(f|\mathcal{H}) + \mathbf{K}(b|\mathcal{H}), \tag{B.25}$$

$$\mathbf{K}(a) + f(a) <^{\log} s + \mathbf{I}(f; \mathcal{H}) + (\mathbf{K}(b|f) + \mathbf{K}(b|\mathcal{H})). \tag{B.26}$$

The remaining part of the proof shows that $\mathbf{K}(b|f) + \mathbf{K}(b|\mathcal{H}) = O(\log(s + \mathbf{K}(b)))$. This is sufficient to proof the theorem due to its logarithmic precision and by the right hand side of the inequality of Equation (B.23) being larger than $s + \mathbf{K}(b)$ (up to a logarithmic factor). Since b is a prefix of border, due to proposition (35), one gets that $\mathbf{K}(b|\mathcal{H}) < O(\mathbf{K}(\|b\|)) < O(\log \|b\|) < O(\log \mathbf{K}(b))$. Thus combined with Equation (B.26) and also Equation (B.23), one gets

$$\mathbf{K}(a) + f(a) <^{\log} s + \mathbf{I}(f; \mathcal{H}) + \mathbf{K}(b|f). \tag{B.27}$$

We now prove $\mathbf{K}(b|f) <^+ \mathbf{K}(s, \|b\|)$. This follows from the existence of an algorithm, that when given f , s , and $\|b\|$, computes $S(b')$ for all $b' \in \{0, 1\}^{\|b\|}$ ordered by \triangleleft , and then outputs the first b' such that $S(b') < s$. This output is b otherwise there exists total $b' \triangleleft b$, with $\|b'\| = \|b\|$, and $S(b') < s$. This implies the existence of total string b'^- such that $S(b'^-) < s$. This contradicts the definition of b being the shortest total string with $S(b) < s$. So $\mathbf{K}(b|f) <^+ \mathbf{K}(s, \|b\|)$ and thus one gets the final form of the theorem, as shown below. Equation (B.28) is again due to the right hand side of Equation (B.23).

$$\begin{aligned}
\mathbf{K}(a) + f(a) &<^{\log} s + \mathbf{I}(f; \mathcal{H}) + \mathbf{K}(s, \|b\|), \\
\mathbf{K}(a) + f(a) &<^{\log} s + \mathbf{I}(f; \mathcal{H}), \\
\min_{a \in \text{Dom}(f)} \mathbf{K}(a) + f(a) &<^{\log} -\log \sum_{a \in \text{Dom}(f)} \mathbf{m}(a) 2^{-f(a)} + \mathbf{I}(f; \mathcal{H}).
\end{aligned} \tag{B.28}$$

Corollary 41 (EL Theorem) *For finite $D \subset \{0, 1\}^*$, $\min_{x \in D} \mathbf{K}(x) <^{\log} -\log \sum_{x \in D} \mathbf{m}(x) + \mathbf{I}(x; \mathcal{H})$.*