A Level Maths - M2 Sam Robbins 13SE

# Centres of mass

# 1 Centre if mass of a discrete mass distribution

$$\bar{x} = \frac{\sum m_i x_i}{\sum m_i}$$
$$\bar{y} = \frac{\sum m_i y_i}{\sum m_i}$$

#### Example

| Mass | 2 | 3 | 2  |
|------|---|---|----|
| x    | 2 | 3 | -3 |
| y    | 3 | 6 | 2  |

$$\bar{x} = \frac{2 \times 2 + 3 \times 3 + 2 \times -3}{2 + 3 + 2} = 1$$

$$\bar{y} = \frac{2 \times 3 + 3 \times 6 + 2 \times 2}{2 + 3 + 2} = 4$$

### 2 Uniform laminae

For a triangular lamina the centre of mass is  $\frac{2}{3}$  along the line from the vertex to the middle of the line opposite. For a sector of a circle, radius r, where the angle at the centre is  $2\alpha$  the centre of mass is  $\frac{2r\sin\alpha}{2\alpha}$ 

#### 3 Rods

In a circular arc, radius r, where the angle at the centre is  $2\alpha$ , the centre of mass is  $\frac{r \sin \alpha}{\alpha}$  away from the centre.

## 4 Equilibrium

To avoid tipping, the lime of action of the weight must be within the side of the lamina in contact with the plane. If a lamina is suspended from a fixed point, the centre of mass will be vertically below the point of suspension. Assumptions made in equilibrium calculations:

- No friction at the point of suspension
- The mass of each area is uniform
- The mass is uniform at the join