


Radioactivity


1 Rutherford Scattering

1.1 The plum pudding model



The plum pudding model was the initial model of the atom, stating a sphere of positive charge with electrons embedded into it.


1.2 Rutherford's experiment



Rutherford's experiment involved firing a beam of alpha particles at gold foil and measuring the paths of particles from the foil.

- Gold was used as it was expected to have a large nucleus
- The screen fluoresces when collided with
- This showed the atom was mostly empty space with a positive nucleus


1.2.1 Results



| Observation | Explanation |
|--|--|
| Most electrons pass all the way through | Atoms are mostly empty space |
| Some are deflected | The atom has a positive centre |
| Some are deflected by significant angles | The positive charge is condensed in a small area |

1.3 Estimating the size of the nucleus

1.3.1 Closest approach method




KE=EPE

$$8.0 \times 10^{-13} = \frac{1}{4\pi\epsilon_0} \times \frac{Q_{Au}}{r} \times Q_{\alpha}$$

$$r = 4.55 \times 10^{-14}$$

1.3.2 Estimate from scattering data

- About $\frac{1}{10,000}$ deflected through more than 90°
- Foil had n layers of atoms



$$\frac{\frac{1}{4}\pi d^2}{\frac{1}{4}\pi D^2} = \frac{d^2}{D^2} = \frac{1}{10,000n}$$

$n = 10^4$ layers

$$\frac{d^2}{D^2} = \frac{1}{10,000 \times 1 \times 10^4}$$


$$d = \frac{D}{10,000}$$

2 Radioactive materials

2.1 Sources of background radiation by most common

1. Air (e.g. radon gas)
2. Medical
3. Ground and buildings
4. Food and drink
5. Cosmic rays
6. Nuclear weapons
7. Air travel
8. Nuclear power

2.2 Geiger Müller tube



When a particle of ionising radiation enters the tube, the particle ionises the gas atoms along its track. The negative ions are attracted to the rod and the positive ions to the wall. These ions cause further ionisation, creating enough ions for a current to flow. A pulse of charge passes round the circuit through resistor R , causing the voltage pulse across R which is recorded as a single count by the pulse counter

The dead time of the tube, the time taken to regain its non conducting state after an ionising particle enters it, is typically of the order of 0.2ms.

| | Alpha | Beta | Gamma |
|--------------------------------|-----------------------------|--|-----------------------------|
| Nature | 2 Protons+2 Neutrons | High speed electron or positron | High energy photon |
| Range | Up to 10cm | Up to 1m | Infinite |
| Deflection in a magnetic field | Deflected | Opposite direction to α particles and more easily deflected | Not deflected |
| Absorption | Paper | Aluminium | Lead |
| Ionisation | 10^4 ions per mm | 100 ions per mm | Very weak ionising effect |
| Energy of each particle | Constant for a given source | Varies up to a maximum for a given source | Constant for a given source |