

# Maclaurin and Taylor Series

## 1 Maclaurin's expansion

$$f(x) = f(0) + f'(0)x + f''(0)\frac{x^2}{2!} + f'''(0)\frac{x^3}{3!} + \dots + f^{(r)}(0)\frac{x^r}{r!} \dots$$

For the continuous function,  $f$ , given by  $f : x \Rightarrow f(x)$  (where  $x$  is real), then providing  $f(0), f'(0), f''(0)$  etc all have finite values. This is an infinite series.

### 1.1 Example

Given that  $f(x) = e^x$  can be written as an infinite series in the form:

$$f(x) = e^x = a_0 + a_1x + a_2x^2 + a_3x^3 + \dots + a_rx^4 + \dots$$

And that it is valid to differentiate an infinite series term by term, show that:

$$e^2 = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots + \frac{x^r}{r!} + \dots$$

Find up to the third differential of  $f(x)$  and the value of zero for each

$$\begin{array}{ll} f(x) = e^x & f(0) = 1 \\ f'(x) = e^x & f''(0) = 1 \\ f''(x) = e^x & f'''(0) = 1 \\ f'''(x) = e^x & f^{(4)}(0) = 1 \end{array}$$

$$\begin{aligned} f(x) &= 1 + 1 \times x + 1 \times \frac{x^2}{2!} + 1 \times \frac{x^3}{3!} \\ f(x) &= 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots + \frac{x^r}{r!} + \dots \end{aligned}$$

### 1.2 Standard results

Standard results are given on the data sheet, these can then be used for adapted forms of the results also. Remember to consider the limits where appropriate.

## 2 Taylor expansion

The conditions of the Maclaurin expansion mean that some functions, such as  $\ln x$  cannot be expanded as a series in ascending powers of  $x$ .

The construction of the Maclaurin expansion focuses on  $x = 0$  and values of  $x$  very close to zero. The Taylor expansion focuses on  $x = a$ .

Considering the functions  $f$  and  $g$ , where  $f(x + a) \equiv g(x)$  then:

$$f^r(a) = g^r(0)$$

Turning the Maclaurin expansion for  $g$  from:

$$g(x) = g(0) + g'(0)x + \frac{g''(0)}{2!}x^2 \dots$$

Into

$$f(x + a) = f(a) + f'(a)x + \frac{f''(a)}{2!}x^2 + \dots + \frac{f^r(a)}{r!}x^r$$

Replacing  $x$  by  $x-a$  gives

$$f(x) = f(a) + f'(a)(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \dots + \frac{f^r(a)}{r!}(x-a)^r$$

These are the two forms of the Taylor expansion, when  $a=0$ , they both become the Maclaurin expansion.

### 2.1 Example

Find the Taylor expansion of  $\cos 2x$ , in ascending powers of  $(x - \frac{\pi}{4})$  up to  $(x - \frac{\pi}{4})^5$  Find the differentials of  $f(x)$  up to the fifth derivative, and the associated values when  $x = \frac{\pi}{4}$

$$\begin{array}{ll} f(x) = \cos 2x & f(\frac{\pi}{4}) = 0 \\ f'(x) = -2 \sin 2x & f'(\frac{\pi}{4}) = -2 \\ f''(x) = -4 \cos 2x & f''(\frac{\pi}{4}) = 0 \\ f'''(x) = 8 \sin 2x & f'''(\frac{\pi}{4}) = 8 \\ f^{(4)}(x) = 16 \cos 2x & f^{(4)}(\frac{\pi}{4}) = 0 \\ f^{(5)}(x) = -32 \sin 2x & f^{(5)}(\frac{\pi}{4}) = -32 \end{array}$$

Substitute in the associated values into the formula

$$\cos 2x = 0 - 2\left(x - \frac{\pi}{4}\right) + 0 + \frac{8}{3!}\left(x - \frac{\pi}{4}\right)^3 + 0 - \frac{32}{5!}\left(x - \frac{\pi}{4}\right)^5$$

$$\cos 2x = -2\left(x - \frac{\pi}{4}\right) + \frac{4}{5}\left(x - \frac{\pi}{4}\right)^3 - \frac{4}{15}\left(x - \frac{\pi}{4}\right)^5$$

### 3 Finding the solution, in the form of a series to a differential equation using the Taylor series method

Suppose you have a first order differential equation of the form  $\frac{dy}{dx} = f(x, y)$  and you know the initial condition that at  $x = x_0, y = y_0$ , then you can calculate  $\left(\frac{dy}{dx}\right)_{x_0}$  by substituting  $x_0$  and  $y_0$  into the original differential equation.

By successive differentiation of the original differential equation, the values of  $\left(\frac{d^2y}{dx^2}\right)_{x_0}$  and  $\left(\frac{d^3y}{dx^3}\right)_{x_0}$  and so on can be found by substituting previous results into the derived equations.

The series solution to the differential equation is found using the Taylor series in the form:

$$y = y_0 + (x - x_0)\left(\frac{dy}{dx}\right)_{x_0} + \frac{(x - x_0)^2}{2!}\left(\frac{d^2y}{dx^2}\right)_{x_0} + \frac{(x - x_0)^3}{3!}\left(\frac{d^3y}{dx^3}\right)_{x_0} + \dots$$

In the common situation where  $x_0 = 0$  then this reduces to the Maclaurin series

$$y = y_0 + x\left(\frac{dy}{dx}\right)_{x_0} + \frac{x^2}{2!}\left(\frac{d^2y}{dx^2}\right)_{x_0} + \frac{x^3}{3!}\left(\frac{d^3y}{dx^3}\right)_{x_0} + \dots$$

#### 3.1 Example

Using the Taylor method to find a series solution, in ascending powers of  $x$  up to and including the term in  $x^3$ , of:

$$\frac{d^2y}{dx^2} = y - \sin x$$

Given that when  $x = 0, y = 1$  and  $\frac{dy}{dx} = 2$

Use the formula to find  $\frac{d^2y}{dx^2}$

$$\frac{d^2y}{dx^2} = 1 - \sin(0) = 1$$

$$\frac{d^2y}{dx^2} = 1$$

Differentiate the formula to obtain a formula for  $\frac{d^3y}{dx^3}$

$$\frac{d^3y}{dx^3} = \frac{dy}{dx} - \cos x = 2 - \cos(0) = 1$$

$$\frac{d^3y}{dx^3} = 1$$

Substitute the known values into the Maclaurin formula

$$y = 1 + x \times 2 + \frac{x^2}{2!} \times 1 + \frac{x^3}{3!} \times 1$$

$$y = 1 + 2x + \frac{x^2}{2} + \frac{x^3}{6}$$