

HEALTH CARE REFORM

Home Blood Pressure Management and Improved Blood Pressure Control

Results From a Randomized Controlled Trial

Hayden B. Bosworth, PhD; Benjamin J. Powers, MD; Maren K. Olsen, PhD; Felicia McCant, MSSW; Janet Grubber, MSPH; Valerie Smith, MS; Pamela W. Gentry, RN, BSN; Cynthia Rose, RN, BSN; Courtney Van Houtven, PhD; Virginia Wang, PhD; Mary K. Goldstein, MD; Eugene Z. Oddone, MD, MHS

Background: To determine which of 3 interventions was most effective in improving blood pressure (BP) control, we performed a 4-arm randomized trial with 18-month follow-up at the primary care clinics at a Veterans Affairs Medical Center.

Methods: Eligible patients were randomized to either usual care or 1 of 3 telephone-based intervention groups: (1) nurse-administered behavioral management, (2) nurse- and physician-administered medication management, or (3) a combination of both. Of the 1551 eligible patients, 593 individuals were randomized; 48% were African American. The intervention telephone calls were triggered based on home BP values transmitted via telemonitoring devices. Behavioral management involved promotion of health behaviors. Medication management involved adjustment of medications by a study physician and nurse based on hypertension treatment guidelines.

Results: The primary outcome was change in BP control measured at 6-month intervals over 18 months. Both the

behavioral management and medication management alone showed significant improvements at 12 months—12.8% (95% confidence interval [CI], 1.6%-24.1%) and 12.5% (95% CI, 1.3%-23.6%), respectively—but not at 18 months. In subgroup analyses, among those with poor baseline BP control, systolic BP decreased in the combined intervention group by 14.8 mm Hg (95% CI, -21.8 to -7.8 mm Hg) at 12 months and 8.0 mm Hg (95% CI, -15.5 to -0.5 mm Hg) at 18 months, relative to usual care.

Conclusions: Overall intervention effects were moderate, but among individuals with poor BP control at baseline, the effects were larger. This study indicates the importance of identifying individuals most likely to benefit from potentially resource intensive programs.

Trial Registration: clinicaltrials.gov Identifier: NCT00237692

Arch Intern Med. 2011;171(13):1173-1180

Author Affiliations: Center for Health Services Research in Primary Care, Durham Veterans Affairs Medical Center, Durham, North Carolina (Drs Bosworth, Powers, Olsen, Van Houtven, Wang, and Oddone and Mss McCant, Grubber, Smith, Gentry, and Rose); Departments of Medicine, Division of General Internal Medicine (Drs Bosworth, Powers, Van Houtven, Wang, and Oddone and Mss Grubber and Smith) and Biostatistics and Bioinformatics (Dr Olsen), Duke University, Durham; and Veterans Affairs Palo Alto Health Care System and Center for Primary Care and Outcomes Research, Department of Medicine, Stanford University, Palo Alto, California (Dr Goldstein).

IMPROVING THE RATE OF BLOOD pressure (BP) control may require expanding the model of care delivery beyond the traditional clinic-based physician visits. Interventions that use home BP telemonitoring linked with patient feedback and medication titration may enhance access and improve outcomes for adults with hypertension. Prior interventions have shown a benefit using this model up to 12 months in highly selected patients, with few racial minorities, and with most patients prescribed fewer than 2 antihypertensive medications at enrollment.^{1,2} It is not known whether these results apply to a racially diverse primary care population with limited resources and more aggressive medication regimens or whether treatment effects are sustained beyond 12 months.

Home BP telemonitoring may improve the quality of decision making and provide the ongoing surveillance required for timely interventions for poor BP control, although it may be more resource intensive than traditional clinic visits.^{3,4} Home BP measurements are predictive of target organ damage and may reduce unnecessary treatment of “white-coat hypertension” (high BP in office and normal BP out of the office).⁵⁻⁷ Home BP telemonitoring is also consistent with the patient-centered medical home (PCMH)

See Invited Commentary at end of article

model of health care.⁸ The PCMH seeks to retain the primacy of an ongoing relationship between patients and primary care providers but also to support this relation-

ship by using nonphysician team members in the systematic processes for non-clinic-based care. Improving current hypertension control rates may require new methods of care delivery that can be applied to primary care populations and are consistent with the PCMH. However, few elements of PCMH have been tested, and there have been calls for more rigorous evaluation.⁹ Furthermore, it is becoming increasingly important to identify individuals who are most likely to benefit from these programs given increasingly scarce resources.

This randomized clinical trial evaluates novel hypertension treatment delivery methods based on home telemonitoring of BP and tests to determine which of 3 interventions delivered via telephone was most effective in improving BP control.

METHODS

DESIGN, SETTING, AND RECRUITMENT

The Hypertension Intervention Nurse Telemedicine Study (HINTS) trial evaluated 3 telephone-based interventions in a 4-group design: (1) nurse-administered, behavioral management intervention; (2) nurse-administered, physician-directed medication management intervention using a validated clinical decision support system; (3) combined behavioral management and medication management intervention; and (4) usual care. Potentially eligible individuals were selected from among primary care patients in Durham, North Carolina, Veterans Affairs (VA) general internal medicine clinics (VAMC). Eligible patients had a diagnosis of hypertension, were using a BP-lowering medication, and had inadequate BP control ($>140/90$ mm Hg for all patients) based on the average of the prior 12 months of clinic BP recordings obtained from electronic medical records. We excluded patients who received hemodialysis; had a serum creatinine level greater than 2.5 mg/dL (221 μ mol/L) or no documentation of renal function; had ever had an organ transplant; had been hospitalized for stroke, myocardial infarction, or coronary artery revascularization within 3 months of contact; had a diagnosis of metastatic cancer or dementia; did not have a home telephone; resided in a nursing home; received home health care; or had severely impaired hearing or speech.

A research assistant sent recruitment letters to eligible patients, contacted patients by telephone, and arranged an in-person meeting at the patient's next primary care provider visit to obtain informed consent and conduct a baseline interview. Consenting patients were then randomized to 1 of the 4 arms using consecutively numbered envelopes; randomization was stratified by diabetic status. Participants were reimbursed \$10 for the baseline visit and the 3 subsequent 6-month BP measurements (\$40 total). The Durham VAMC institutional review board approved this study.

INTERVENTIONS

Telemedicine and Home BP Monitoring

All patients randomized to an intervention arm were provided a wireless home BP monitor (model UA-767PC; A&D Medical Digital Blood Pressure, Tokyo, Japan) and telemedicine device (model 102; Carematix Inc, Chicago, Illinois) and advised to measure their BP once every other day. The telemedicine device connected to a telephone line like an answering machine. Patients' BP measurements were transmitted automati-

cally to a secure server.¹⁰ Because BP measured at home averages 6 to 8 mm Hg and 5 to 6 mm Hg lower than values obtained during a routine clinic visit for systolic and diastolic BP, respectively,¹¹ we designed intervention trigger alerts based on a 2-week average home BP of 135/85 mm Hg mm for those without diabetes mellitus (DM).¹² However, because an internal expert panel raised concerns about reducing diastolic BP too low with home BP values for those without DM, intervention trigger alerts were based on a 2-week home BP average of at least 135/80 mm Hg for patients with DM. If an intervention was activated, a patient's home BP was reassessed again at 6 weeks before triggering the intervention again. If 3 BP values were not transmitted over a 2-week period, a research assistant contacted patients to remind them to use the equipment and to answer any potential technical questions regarding the equipment. A mean home BP measurement that exceeded the threshold over this interval triggered 1 of 3 assigned interventions (eFigure; <http://www.archinternmed.com>). Patients who maintained adequate BP control did not activate the intervention but triggered a contact every 6 months to reinforce their positive behavior. A safety alert was activated if 2 consecutive home BP measurements recorded at least 12 hours apart were greater than 175/105 mm Hg. In this instance, a nurse contacted the patient and initiated an approved safety protocol.

Behavioral Management Intervention

The behavioral management intervention consisted of 11 tailored health behavior modules focused on improving hypertension self-management as noted in eTable 1.¹³ The behaviors addressed in the intervention included hypertension knowledge, including medication adverse effects and medication memory; resources to improve health care access; and the patient-health care provider relationship. Patients were also provided evidence-based recommendations regarding hypertension-related behaviors, including salt intake, weight, stress reduction, smoking cessation, and alcohol use. Verbal information was reinforced with mailed handouts. The nurse used an intervention software application that contained predetermined scripts and patient-specific tailored algorithms for the modules. Each encounter consisted of 3 or 4 modules and lasted 12 to 14 minutes.

Medication Management Intervention

On triggers in the medication management intervention, a nurse notified a study physician and provided the physician with a medication change recommendation based on the decision support software.¹⁴ The study physician reviewed the patient's BP, medication, and adherence (based on patient report and prescription refill data) with the nurse and decided whether to change hypertension medication. The nurse communicated recommended changes to the patient, while the study physician electronically prescribed the medication and generated a note in the patient's medical record that was cosigned by the patient's primary care physician. The nurse called the patient 3 weeks after the medication change to obtain reports of adverse effects and address patient questions.

Combined Intervention

In this arm, patients received the full dose of each intervention. The nurse initially addressed recommended medication adjustments using the established method described in the previous subsection, followed by the tailored behavioral intervention.

Usual Care

Patients in all 4 study arms received primary care and management of hypertension according to the discretion of their primary care provider. Patients treated with usual care received no contact with the intervention nurses and did not receive home telemonitoring equipment.

Study Measures

Study recruitment started in May 2006, and the final 18-month follow-up appointment was completed in July 2009. The primary outcome of the study was BP control measured at baseline and at 6, 12, and 18 months using a standardized research protocol.¹⁵ At each measurement point, a research assistant masked to intervention arms asked patients to rest for 5 minutes before obtaining 2 BP measurements using a digital sphygmomanometer.¹⁶ Inadequate BP control was defined as standardized research protocol measurement of systolic BP of at least 140 mm Hg or a diastolic BP of at least 90 mm Hg for individuals without DM, and systolic BP of at least 130 mm Hg or a diastolic BP of at least 80 mm Hg for individuals with DM.^{17,18}

We conducted the cost analysis from the perspective of the VA health care system, including both direct and indirect costs of the intervention. Resource utilization costs were derived from the VA's DSS National Data Extract inpatient and outpatient files.¹⁹ We examined outpatient utilization counts (primary and specialty care) and aggregated inpatient and outpatient costs to obtain total VA resource utilization costs.

Sample size estimation was based on the primary hypothesis that patients randomized to an intervention arm would have improved BP control at 18 months compared with the usual care group. A common intercept and a linear change in BP control were assumed, so the comparison was a difference in slopes (the treatment by time interaction in a logistic mixed-effects regression model). Sample size and power estimates were generated empirically via a simulation study using PROC NLMIXED statistical software (version 9.2; SAS Institute Inc, Cary, North Carolina). From previous studies^{13,20} baseline BP control at a single measurement was estimated as 30% and the 18-month dropout rate as 15%. The random intercept variance component was assumed to be 1.7 (equivalent to a patient intraclass correlation of 0.34).²¹ To detect a difference in slopes resulting in a 15% improvement in BP control at 18 months relative to the control group (ie, 45% vs 30%) with 80% power and a type-I error rate of 5%, 600 patients were needed.

For the primary analyses, a logistic mixed-effects regression model was used to estimate differences in BP control at each postbaseline time point for each of the intervention groups relative to usual care. Model parameters were estimated using maximum likelihood via an adaptive quadrature method (using PROC NLMIXED software) and included a common intercept,²² baseline DM status, dummy coded time, and intervention arms interacted with each follow-up time point. A patient-level random effect was included in the model to account for the correlation of patients' repeated measurements over time. Marginalized estimates and corresponding confidence intervals for the proportion in BP control for the usual care and each intervention group at 6, 12, and 18 months were calculated to estimate the relative improvement in proportion of patients with BP control.²¹

For the secondary analyses, we examined the intervention group effect on mean systolic and diastolic BPs over time. We used longitudinal data analysis models with an unstructured covariance to account for the correlation of patients' repeated measurements over time (using PROC MIXED software). The outcome variable in these models was the study BP at base-

line, 6, 12, and 18 months²³; model parameters included a common intercept, baseline DM status, and dummy coded time, and intervention arms interacted with each follow-up time point. Mean differences between each intervention group and the usual care group at 6, 12, and 18 months were calculated, along with corresponding 95% confidence intervals (CIs), using SAS ESTIMATE statements. Finally, in post hoc analyses, we created 2 subgroups of patients: (1) those whose BP was in control at baseline and (2) those whose BP was out of control at baseline. We then examined the intervention group effect on mean systolic and diastolic BPs over time within each of these subgroups using a similar model to that described in the previous subsection.

To determine whether the interventions were more resource intensive than usual care in the clinic setting, we also examined differences in patients' median medical resource use and costs in the 18 months after initiating the interventions. In this analysis, we used *t* test and Wilcoxon test to assess whether there were intervention cost or health care use differences between the intervention groups.

Patients were evaluated based on the initial randomization group; 591 patients were included in the analyses (**Figure 1**). All available data, including data from participants who subsequently discontinued the study, were used for analyses. All analyses were conducted with full likelihood methods, which are valid under a missing data framework in which the missing values are dependent on intervention group, baseline DM status, and any observed BP values.²¹

RESULTS

PATIENTS

Of the 591 study patients, 48% were African American, 92% were male, and 59% of participants had their BP under control at baseline using standardized measurements (**Table 1**). No crossover among study groups occurred. No study-related adverse events occurred in any intervention group. Overall, BP measurements were available for 503 patients (85%) at 18 months of follow-up (**Figure 1**). Completion rates of the 18-month follow-up were similar across groups (83%-89%).

PRIMARY OUTCOME: BP CONTROL

The improvement in BP control relative to usual care at 12 months was statistically significant in the behavioral management group and medication management group at 12.8% (95% CI, 1.6% to 24.1%; *P* = .03) and 12.5% (95% CI, 1.3% to 23.6%; *P* = .03), respectively. The benefit in the combined intervention group was 8.3% (95% CI, -3.3% to 19.9%; *P* = .16) and was not statistically significant (**Table 2** and **Figure 2**). At 18 months, only the combined intervention group showed evidence of improved BP control relative to usual care, 7.7% (95% CI, -4.1% to 19.5%; *P* = .20); however, the difference was not statistically significant.

SYSTOLIC AND DIASTOLIC BP

The largest sustained improvement for systolic BP was observed in the combined intervention group (**Table 2** and **Figure 3**). Relative to usual care at 12 months, the mean systolic BP was 2.1 mm Hg lower (95% CI, -6.2 to




Figure 1. Hypertension Intervention Nurse Telemedicine Study recruitment flowchart. BP indicates blood pressure. *Patients no longer receiving care at Durham, North Carolina, Veterans Affairs clinics; residing in a nursing home; receiving hemodialysis; having creatinine levels higher than 2.5 mg/dL (221 μ mol/L); or diagnosed as having dementia.

2.1 mm Hg; $P = .33$) in the behavioral management group, 2.4 mm Hg lower (95% CI, -6.5 to 1.7 mm Hg; $P = .25$) in the medication management group, and 4.3 mm Hg lower (95% CI, -8.5 to -0.2 mm Hg; $P = .04$) in the combined intervention group compared with the usual care group. However, by 18 months, the mean systolic BP was lower only in the medication management and combined intervention groups compared with the usual care group, but these differences were not statistically significant (-1.2 mm Hg [95% CI, -5.7 to 3.2 mm Hg], $P = .58$; and -3.6 mm Hg [95% CI, -8.1 to 0.9 mm Hg], $P = .11$). There were differences between intervention groups and usual care for diastolic BP, but they were not statistically significant (Table 2).

Over half of patients enrolled, 348, had adequate control at baseline (mean systolic BP, 117.9 mm Hg; mean diastolic BP, 71.6 mm Hg) compared with 243 patients with inadequate control (mean systolic BP, 145.0 mm Hg; mean diastolic BP, 85.7 mm Hg). In post hoc analyses of these subgroups, we found that individuals with adequate BP control at baseline continued to remain in control over the 18 months of the study (Table 3). For individuals with poor BP control, systolic BP decreased in the behavioral arm by 8.3 mm Hg at 12 months (95% CI, -15.1 to -1.6 mm Hg; $P = .02$) compared with the decrease in the usual care group but was no longer significantly lower at 18 months. The mean systolic BP was significantly lower in the medication management arm at 12 months, -7.9 mm Hg (95% CI, -14.5 to -1.4 mm Hg;

$P = .02$) relative to usual care. In the combined intervention arm, systolic BP decreased by 14.8 mm Hg (95% CI, -21.8 to -7.8 mm Hg; $P < .001$) at 12 months and 8.0 mm Hg (95% CI, -15.5 to -0.5 mm Hg; $P = .04$) at 18 months relative to usual care (Table 3 and Figure 4). Among those with inadequate baseline BP control, diastolic BP decreased significantly in the medication management arm by 4.2 mm Hg (95% CI, -8.3 to -0.2 mm Hg; $P = .04$) at 18 months compared with usual care. In the combined intervention arm, diastolic BP decreased by 5.3 mm Hg (95% CI, -9.5 to -1.2 mm Hg; $P = .01$) and 5.5 mm Hg (95% CI, -9.7 to -1.2 mm Hg; $P = .01$) at 12 and 18 months, respectively, relative to usual care.

INTERVENTION ACTIVATION

Over the 18-month intervention period, 389 of the 444 intervention patients (88%) triggered an alert for the nurse to conduct an intervention call as a result of poor BP control (eTable 2). These 389 patients activated a total number of 1945 intervention alerts. Seventy-nine (18%) activated a safety alert, while 65 (37%) activated praise calls as a result of continued BP control. In total, we referred 43 individuals 90 times for care owing to triggering safety levels. The number of alerts activated and the number of patients who generated alerts were fairly similar across the 3 intervention arms (eTable 2).

Across the 3 intervention groups, 13.2 minutes was the mean time spent by nurses on each completed en-

Table 1. HINTS Baseline Sample Characteristics

Baseline Characteristic ^a	Patient Groups				
	Total (n=591)	Usual Care (n=147)	Behavioral Management Intervention (n=148)	Medication Management Intervention (n=149)	Combined Intervention (n=147)
Demographics					
Age, mean (SD)	64 (10)	64 (10)	63 (11)	64 (10)	63 (11)
Race/ethnicity					
White	49	50	53	49	44
African American	48	48	45	48	52
Other	3	1	3	3	5
Male	92	96	92	93	86
Married	66	65	71	65	61
Completed <12 y of school	13	10	14	12	18
Low literacy level (<ninth grade; REALM score ≤60)	38	45	39	32	37
Employed	35	35	34	34	35
Current smoker	21	22	19	20	22
BMI, mean (SD)	30.3 (5.3)	30.0 (5.5)	30.6 (5.6)	30.2 (5.0)	30.6 (5.1)
Medical history					
>10 y history of high BP	75	71	76	77	74
DM	43	44	44	43	40
Hypertension medications, mean (SD), No. ^b	2.4 (1.2)	2.5 (1.2)	2.4 (1.2)	2.3 (1.1)	2.4 (1.1)
BP ^c					
Controlled BP ^d	59	61	58	52	65
Systolic BP, mean (SD), mm Hg	129 (20)	128 (17)	129 (19)	132 (21)	127 (21)
Diastolic BP, mean (SD), mm Hg	77 (13)	78 (14)	77 (12)	78 (14)	77 (13)

Abbreviation: BMI, body mass index (calculated as weight in kilograms divided by height in meters squared); BP, blood pressure; DM, diabetes mellitus; HINTS, Hypertension Intervention Nurse Telemedicine Study; REALM, Rapid Estimate of Adult Literacy in Medicine²⁴; VA, Veterans Affairs.

^aAll data except BP, age, BMI, DM status, sex/ethnicity, and number of hypertension medications (national VA database) were patient-reported and are given as percentages except where noted. When missing values existed, they were included in the calculation of percentages.

^bMeans are for the 507 VA patients whose prescribed hypertension medication fills, as recorded in the national VA database, overlapped with their baseline interview date.

^cWhen patients had multiple BP readings during their baseline visit, means of their systolic and diastolic BP readings were used as the baseline BP values.

^dAccording to the Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure control¹⁷: less than 140/90 mm Hg for those without DM; less than 130/80 mm Hg for those with DM.

counter: 12.0, 13.9, and 13.7 minutes in behavioral management, medication management, and combined intervention, respectively.

COSTS OF INTERVENTION AND MEDICAL RESOURCE USE

Intervention costs over 18 months were \$947 for behavioral management, \$1275 for medication management, and \$1153 for the combined intervention arm. The number of primary care and specialty care encounters over 18 months was similar for the 4 groups; the medians were 4 primary care visits for all groups ($P=.21$) and 5 to 6 specialty care visits ($P=.12$). There was no significant difference in median 18-month total VA medical cost across groups, which ranged from \$5180 per patient for the medication management arm to \$6910 for the behavioral intervention ($P=.28$).

COMMENT

We examined the effects of a patient behavioral management intervention, medication management, and a combination of the 2 interventions delivered by telephone and activated by home BP monitoring among adults with hypertension treated in primary care. The behavioral man-

agement and medication management groups alone showed significantly improved BP control at 12 months compared with usual care. The combined intervention group demonstrated statistically significant improvement in systolic BP at 12 months relative to usual care, but this was no longer significant at 18 months.^{13,23} In post hoc analyses, stratified by baseline BP control, there were significant and clinically meaningful BP improvements in all 3 arms at 12 months and in the combined intervention at 18 months compared with usual care, with the largest effects in the combined intervention arm (systolic BP decrease of 8 mm Hg and diastolic BP decrease of 5.5 mm Hg at 18 months).

In a review of nurse-led interventions for hypertension, nurse prescribing showed greater reductions in BP.²⁵ In a recent Cochrane review,²⁶ self-monitoring was associated with moderate net reduction in systolic BP and diastolic BP. Randomized controlled trials (RCTs) of educational interventions directed at patients were unlikely to be associated with large net reductions in BP by themselves. However, nurse- or pharmacist-led care may be a promising way forward, with most RCTs being associated with improved hypertension outcome.²⁶⁻²⁸ Our intervention, similar to prior work,^{13,23} demonstrated the potential for a nurse-led intervention to improve hypertension management in the short term and, among a care-

Table 2. Estimated Differences in BP Control, Systolic BP, and Diastolic BP Between HINTS Intervention and Usual Care Groups at 6, 12, and 18 Months

Outcome Time Period	Behavioral Management vs Usual Care	Medication Management vs Usual Care	Combined Intervention vs Usual Care
Primary Outcome, % (95% CI)			
BP in control, % ^a			
6 mo	3.3 (−8.7 to 15.3)	6.7 (−5.1 to 18.5)	5.4 (−6.5 to 17.3)
12 mo	12.8 (1.6 to 24.1)	12.5 (1.3 to 23.6)	8.3 (−3.3 to 19.9)
18 mo	−2.9 (−15.0 to 9.3)	−0.3 (−12.4 to 11.9)	7.7 (−4.1 to 19.5)
Secondary Outcomes, mm Hg (95% CI)^b			
Systolic BP			
6 mo	1.3 (−2.9 to 5.4)	−0.4 (−4.6 to 3.7)	−0.9 (−5.1 to 3.2)
12 mo	−2.1 (−6.2 to 2.1)	−2.4 (−6.5 to 1.7)	−4.3 (−8.5 to −0.2)
18 mo	2.2 (−2.2 to 6.6)	−1.2 (−5.7 to 3.2)	−3.6 (−8.1 to 0.9)
Diastolic BP			
6 mo	0.5 (−1.9 to 2.9)	−1.2 (−3.5 to 1.2)	−0.2 (−2.5 to 2.2)
12 mo	−0.7 (−3.2 to 1.9)	−0.9 (−3.5 to 1.6)	−0.01 (−2.6 to 2.6)
18 months	0.6 (−2.0 to 3.3)	−0.5 (−3.2 to 2.1)	−1.4 (−4.0 to 1.3)

Abbreviations: BP, blood pressure; CI, confidence interval; HINTS, Hypertension Intervention Nurse Telemedicine Study.

^aThe BP control estimates are marginalized probabilities with corresponding 95% CIs derived from a logistic mixed-effects regression model.²¹ Positive differences reflect improvement compared with usual care, and negative differences reflect worsening compared with usual care. The model-estimated intrapatient correlation was 0.39. See the Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure.¹⁷

^bSystolic and diastolic BP estimates are based on a longitudinal data model with an unstructured covariance matrix. Negative differences reflect improvement compared with usual care, and positive differences reflect worsening compared with usual care. The model-estimated systolic BP correlation between time points ranged from 0.24 (at baseline and 18 months) to 0.41 (at 6 and 18 months), and the estimated diastolic BP correlation between time points ranged from 0.40 (at baseline and 18 months) to 0.52 (at 6 and 18 months).




Figure 2. Estimated proportion in blood pressure (BP) control from baseline to 18 months, by intervention group. Bars represent 95% confidence intervals (CIs). The estimates and 95% CIs are marginalized probabilities from a logistic mixed effects model and were calculated using predict statements in PROC NLMIXED statistical software (version 9.2; SAS Institute Inc, Cary, North Carolina). The model estimated intrapatient correlation was 0.39.

fully selected group—patients with inadequate BP at baseline—over the longer term.

Overall, the behavioral and medication management intervention groups had significant improvements in BP control relative to the usual care group at 12 months; however, these findings were not sustained at 18 months. One possible explanation for these effects was that patients were receiving more aggressive medication regimens at baseline and may therefore have been less likely to benefit from medication management interventions than those prescribed fewer medications.

Recent studies examining self-monitoring and telemedicine have typically involved higher socioeconomic status, lower level of intensity of baseline treatment (≤ 2 hypertensive drugs, whereas we had individuals pre-




Figure 3. Estimated systolic blood pressure (BP) from baseline to 18 months, by intervention group. Bars represent 95% confidence intervals. Estimates are based on a longitudinal data model with unstructured covariance matrix. The model-estimated systolic BP correlation between time points ranged from 0.24 (at baseline and 18 months) to 0.41 (at 6 and 18 months).

scribed as many as 4),^{1,2} and shorter interventions (< 6 months).^{29–32} Thus, our study was more diverse in literacy level, education, and race/ethnicity and may be more generalizable to the larger population of male Americans with hypertension.

The study has potential limitations. First, despite our inclusion criteria that patients were required to have inadequate BP control based on the average of the prior 12 months of clinic BP recordings obtained from the electronic medical record, 59% at baseline were in control using a more accurate method of assessing BP (ie, systematic assessment in a quiet room, by a trained research assistant). Thus, the discrepancy between BP val-

Table 3. Estimated Differences (95% CI) in Systolic and Diastolic BP Between HINTS Intervention and Usual Care Groups at 12 and 18 Months for BP Control Subgroups

Outcome Subgroup ^a	mm Hg (95% CI)		
	Behavioral Management vs Usual Care	Medication Management vs Usual Care	Combined Intervention vs Usual Care
Systolic BP			
Adequate baseline BP control ^b			
12 mo	1.6 (−3.5 to 6.8)	0.1 (−5.1 to 5.3)	1.3 (−3.7 to 6.3)
18 mo	2.8 (−2.7 to 8.4)	1.8 (−3.9 to 7.4)	−1.7 (−7.2 to 3.8)
Inadequate baseline BP control ^b			
12 mo	−8.3 (−15.1 to −1.6)	−7.9 (−14.5 to −1.4)	−14.8 (−21.8 to −7.8)
18 mo	0.3 (−6.9 to 7.5)	−6.6 (−13.8 to 0.53)	−8.0 (−15.5 to −0.5)
Diastolic BP			
Adequate baseline BP control ^b			
12 mo	0.4 (−2.9 to 3.7)	0.7 (−2.6 to 4.0)	3.3 (0.08 to 6.5)
18 mo	0.07 (−3.4 to 3.5)	2.0 (−1.5 to 5.5)	1.0 (−2.4 to 4.4)
Inadequate baseline BP control ^d			
12 mo	−2.1 (−6.2 to 1.9)	−3.4 (−7.3 to 0.5)	−5.3 (−9.5 to −1.2)
18 mo	1.1 (−3.0 to 5.1)	−4.2 (−8.3 to −0.2)	−5.5 (−9.7 to −1.2)

Abbreviations: BP, blood pressure; CI, confidence interval; DM, diabetes mellitus; HINTS, Hypertension Intervention Nurse Telemedicine Study.

^aInadequate baseline BP control was defined as systolic BP 140 mm/Hg or higher or diastolic BP 90 mm/Hg or higher for individuals without DM and systolic BP 130 mm/Hg or higher or a diastolic BP 80 mm/Hg or higher for individuals with DM. A total of 348 patients had adequate baseline BP control, and 243 had inadequate baseline BP control. Systolic and diastolic BP estimates are based on a longitudinal data model with an unstructured covariance matrix. Negative differences reflect improvement compared with usual care, and positive differences reflect worsening compared with usual care.

^bThe numbers of patients with adequate baseline BP control in treatment groups were as follows: usual care, 89; behavioral management, 86; medication management, 78; combined intervention, 95.

^cThe numbers of patients with inadequate baseline BP control in treatment groups were as follows: usual care, 58; behavioral management, 62; medication management, 71; combined intervention, 52.

ues obtained in the clinic and standardized BP measurement has implications for quality indicators. We know from prior work that BP values obtained in the research setting are, on average, 8 mm Hg lower than those obtained in the clinic.^{33,34} It is also important to acknowledge that the home BP values, and those obtained by research staff, were lower, on average, than the values obtained in the clinic. It is difficult to tell to what degree this is a white-coat effect or some other factor. The fact that our study BPs were lower than anticipated may have hampered our ability to detect larger improvements in BP control in the overall analyses. Last, one should exercise caution in interpreting our post hoc analyses given that these analyses were not initially planned.

Receiving all of one's treatment in primary care office settings may not be the optimal care strategy owing to competing issues that arise during routine office visits or due to lack of accessibility to primary care. This study addresses several of the core features of the PCMH model³⁵ and is one of the first to examine a tailored, home-based BP monitoring used to initiate a behavioral intervention to improve hypertension adherence with a home-based hypertension medication management strategy. The intervention included nurses to facilitate non-face-to-face medication management outside the clinic setting, as well as systematic reporting back to the primary care physician, and proactively calling patients—not just relying on the clinic visit to make adjustments. None of the intervention groups increased health care use, but the cost of the interventions alone was approximately \$1100 per patient over 18 months.

Each intervention demonstrated improvements in BP control or systolic BP at 12 months; none of these im-




Figure 4. Estimated systolic blood pressure (BP) from baseline to 18 months, by intervention group, for 241 patients with inadequate baseline BP Control. Bars represent 95% confidence intervals. Estimates are based on a longitudinal data model with unstructured covariance matrix. The model-estimated systolic BP correlation between time points ranged from 0.28 (at baseline and 18 months) to 0.43 (at 6 and 18 months).

provements were sustained at 18 months and did not result in lower medical care costs. However, among those individuals with poor baseline BP control, the combined intervention significantly decreased systolic BP and diastolic BP at 12 and 18 months. Further work is required to identify individuals who will most benefit from this more intensive model of BP management interventions as well as to identify methods to increase or sustain initial effects obtained beyond 12 months.

Accepted for Publication: April 8, 2011.

Correspondence: Hayden B. Bosworth, PhD, Health Services Research and Development, Durham Veterans Affairs Medical Center (152), 508 Fulton St, Durham, NC 27705 (hayden.bosworth@duke.edu).

Author Contributions: All authors had full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis. **Study concept and design:** Bosworth, Olsen, Van Houtven, Goldstein, and Oddone. **Acquisition of data:** Bosworth, Powers, McCant, Gentry, Rose, and Wang. **Analysis and interpretation of data:** Bosworth, Powers, Olsen, McCant, Grubber, Smith, Van Houtven, Wang, and Oddone. **Drafting of the manuscript:** Bosworth, Olsen, and Van Houtven. **Critical revision of the manuscript for important intellectual content:** Bosworth, Powers, Olsen, McCant, Grubber, Smith, Gentry, Rose, Wang, Goldstein, and Oddone. **Statistical analysis:** Olsen, Grubber, and Smith. **Obtained funding:** Bosworth and Olsen. **Administrative, technical, and material support:** Bosworth, Powers, McCant, Gentry, Rose, Wang, and Oddone. **Study supervision:** Bosworth, Powers, and Oddone. **Cost analysis:** Wang and Van Houtven. **Analysis of data:** Bosworth, Olsen, Grubber, Smith, Wang, and Van Houtven. **Providing and maintaining medical management system:** Goldstein.

Financial Disclosure: None reported.

Funding/Support: This research is supported by a grant from Veterans Affairs, Health Services Research and Development (No. IIR 04-426), a career scientist award (No. RCS 08-027) and an Established Investigator Award from American Heart Association to Dr Bosworth. ATHENA-HTN development and testing was supported in part by VA Health Services Research and Development (Nos. CPI 99-275 and IMV-04-062).

Disclaimer: The views expressed in this manuscript are those of the authors and do not necessarily represent the views of the Department of Veterans Affairs.

Online-Only Material: The eFigure and eTables are available at <http://www.archinternmed.com>.

REFERENCES

- Green BB, Cook AJ, Ralston JD, et al. Effectiveness of home blood pressure monitoring, Web communication, and pharmacist care on hypertension control: a randomized controlled trial. *JAMA*. 2008;299(24):2857-2867.
- McManus RJ, Mant J, Bray EP, et al. Telemonitoring and self-management in the control of hypertension (TASMINH2): a randomised controlled trial. *Lancet*. 2010;376(9736):163-172.
- Patient-Centered Medical Home. National Committee for Quality Assurance Web site. <http://www.ncqa.org/tabid/631/Default.aspx>. Accessed May 12, 2011.
- Crabtree BF, Nutting PA, Miller WL, Stange KC, Stewart EE, Jaén CR. Summary of the National Demonstration Project and recommendations for the patient-centered medical home. *Ann Fam Med*. 2010;8(suppl 1):S80-S90, S92.
- Bobrie G, Chatellier G, Genes N, et al. Cardiovascular prognosis of "masked hypertension" detected by blood pressure self-measurement in elderly treated hypertensive patients. *JAMA*. 2004;291(11):1342-1349.
- Staessen JA, Den Hond E, Celis H, et al. Treatment of Hypertension Based on Home or Office Blood Pressure (THOP) Trial Investigators. Antihypertensive treatment based on blood pressure measurement at home or in the physician's office: a randomized controlled trial. *JAMA*. 2004;291(8):955-964.
- Verberk WJ, Kroon AA, Lenders JW, et al; Home Versus Office Measurement, Reduction of Unnecessary Treatment Study Investigators. Self-measurement of blood pressure at home reduces the need for antihypertensive drugs: a randomized, controlled trial. *Hypertension*. 2007;50(6):1019-1025.
- Currell R, Urquhart C, Wainwright P, Lewis R. Telemedicine versus face to face patient care: effects on professional practice and health care outcomes. *Cochrane Database Syst Rev*. 2000;(2):CD002098.
- Barr MS. The need to test the patient-centered medical home. *JAMA*. 2008;300(7):834-835.
- Bosworth HB, Olsen MK, McCant F, et al. Hypertension Intervention Nurse Telemedicine Study (HINTS): testing a multifactorial tailored behavioral/educational and a medication management intervention for blood pressure control. *Am Heart J*. 2007;153(6):918-924.
- Weisser B, Mengden T, Dusing R, Vetter H, Vetter W. Normal values of blood pressure self-measurement in view of the 1999 World Health Organization-International Society of Hypertension guidelines. *Am J Hypertens*. 2000;13(8):940-943.
- Staessen JA, Thijs L; First International Consensus Conference on Blood Pressure Self-Measurement. Development of diagnostic thresholds for automated self-measurement of blood pressure in adults. *Blood Press Monit*. 2000;5(2):101-109.
- Bosworth HB, Olsen MK, Dudley T, et al. Patient education and provider decision support to control blood pressure in primary care: a cluster randomized trial. *Am Heart J*. 2009;157(3):450-456.
- Goldstein MK, Coleman RW, Tu SW, et al. Translating research into practice: organizational issues in implementing automated decision support for hypertension in three medical centers. *J Am Med Inform Assoc*. 2004;11(5):368-376.
- Bosworth HB, Olsen MK, Dudley TK, et al. The Take Control of Your Blood pressure (TCYB) study: study design and methodology. *Contemp Clin Trials*. 2007;28(1):33-47.
- Wright BM, Dore CF. A random-zero sphygmomanometer. *Lancet*. 1970;1(7642):337-338.
- Chobanian AV, Bakris GL, Black HR, et al; National Heart, Lung, and Blood Institute Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure; National High Blood Pressure Education Program Coordinating Committee. The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure: the JNC 7 report. *JAMA*. 2003;289(19):2560-2572.
- Joint National Committee on Detection, Evaluation, and Treatment of High Blood Pressure. *The Sixth Report of the Joint National Committee on Detection, Evaluation, and Treatment of High Blood Pressure (JNC VI)*. Bethesda, MD: US Department of Health and Human Services, National Institutes of Health; 1997.
- VIREC Research User Guide. VHA Decision Support System Clinical National Data Extracts. 2nd ed. US Department of Veterans Affairs, Health Services Research and Development Service, VA Information Resource Center, Web site. <http://www.virec.research.va.gov/References/RUG/RUG-DSS-2nd-Ed-er.pdf>. Accessed May 12, 2011.
- Bosworth HB, Olsen MK, Goldstein MK, et al. The veterans' study to improve the control of hypertension (V-STITCH): design and methodology. *Contemp Clin Trials*. 2005;26(2):155-168.
- Hedeker D, Gibbons RD. *Longitudinal Data Analysis*. New York, NY: John Wiley & Sons; 2006.
- Fitzmaurice G, Laird NM, Ware JH. *Applied Longitudinal Analysis*. New York, NY: Wiley Series in Probability and Statistics; 2004.
- Bosworth HB, Olsen MK, Grubber JM, et al. Two self-management interventions to improve hypertension control: a randomized trial. *Ann Intern Med*. 2009;151(10):687-695.
- Parker RM, Baker DW, Williams MV, Nurss JR. The test of functional health literacy in adults: a new instrument for measuring patients' literacy skills. *J Gen Intern Med*. 1995;10(10):537-541.
- Clark CE, Smith LF, Taylor RS, Campbell JL. Nurse led interventions to improve control of blood pressure in people with hypertension: systematic review and meta-analysis. *BMJ*. 2010;341:c3995.
- Glynn LG, Murphy AW, Smith SM, Schroeder K, Fahey T. Interventions used to improve control of blood pressure in patients with hypertension. *Cochrane Database Syst Rev*. 2010;(3):CD005182.
- Carter BL, Foppe van Mil JW. Comparative effectiveness research: evaluating pharmacist interventions and strategies to improve medication adherence. *Am J Hypertens*. 2010;23(9):949-955.
- Walsh JM, McDonald KM, Shojania KG, et al. Quality improvement strategies for hypertension management: a systematic review. *Med Care*. 2006;44(7):646-657.
- Friedman RH, Kazis LE, Jette A, et al. A telecommunications system for monitoring and counseling patients with hypertension. Impact on medication adherence and blood pressure control. *Am J Hypertens*. 1996;9(4, pt 1):285-292.
- Rogers MA, Small D, Buchan DA, et al. Home monitoring service improves mean arterial pressure in patients with essential hypertension: a randomized, controlled trial. *Ann Intern Med*. 2001;134(11):1024-1032.
- Mehos BM, Saseen JJ, MacLaughlin EJ. Effect of pharmacist intervention and initiation of home blood pressure monitoring in patients with uncontrolled hypertension. *Pharmacotherapy*. 2000;20(11):1384-1389.
- Artinian NT, Washington OG, Templin TN. Effects of home telemonitoring and community-based monitoring on blood pressure control in urban African Americans: a pilot study. *Heart Lung*. 2001;30(3):191-199.
- Kim JW, Bosworth HB, Voils CI, et al. How well do clinic-based blood pressure measurements agree with the mercury standard? *J Gen Intern Med*. 2005;20(7):647-649.
- Ishikawa J, Carroll DJ, Kuruvilla S, Schwartz JE, Pickering TG. Changes in home versus clinic blood pressure with antihypertensive treatments: a meta-analysis. *Hypertension*. 2008;52(5):856-864.
- Iglehart JK. No place like home: testing a new model of care delivery. *N Engl J Med*. 2008;359(12):1200-1202.