



## Blood pressure control by home monitoring: meta-analysis of randomised trials

Francesco P Cappuccio, Sally M Kerry, Lindsay Forbes and Anna Donald

*BMJ* 2004;329:145; originally published online 11 Jun 2004;  
doi:10.1136/bmj.38121.684410.AE

---

Updated information and services can be found at:  
<http://bmj.com/cgi/content/full/329/7458/145>

---

*These include:*

**Data supplement**

"Appendices A - D"  
<http://bmj.com/cgi/content/full/bmj.38121.684410.AE/DC1>

**References**

This article cites 41 articles, 14 of which can be accessed free at:  
<http://bmj.com/cgi/content/full/329/7458/145#BIBL>

3 online articles that cite this article can be accessed at:  
<http://bmj.com/cgi/content/full/329/7458/145#otherarticles>

**Rapid responses**

11 rapid responses have been posted to this article, which you can access for free at:  
<http://bmj.com/cgi/content/full/329/7458/145#responses>

You can respond to this article at:  
<http://bmj.com/cgi/eletter-submit/329/7458/145>

**Email alerting service**

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article

---

**Topic collections**

Articles on similar topics can be found in the following collections

[Clinical Research](#) (497 articles)  
[Systematic reviews \(incl meta-analyses\): examples](#) (185 articles)  
[Health Services Research](#) (440 articles)  
[Hypertension](#) (352 articles)

---

**Correction**

A correction has been published for this article. The contents of the [correction](#) have been appended to the original article in this reprint. The correction is also available online at:  
<http://bmj.com/cgi/content/full/329/7464/499-b>

---

**Notes**

---

To order reprints of this article go to:  
<http://bmj.bmjournals.com/cgi/reprintform>

To subscribe to *BMJ* go to:  
<http://www.bmjournals.com/subscriptions>

Cite this article as: BMJ, doi:10.1136/bmj.38121.684410.AE (published 11 June 2004)

# Primary care

## Blood pressure control by home monitoring: meta-analysis of randomised trials

Francesco P Cappuccio, Sally M Kerry, Lindsay Forbes, Anna Donald

### Abstract

**Objective** To determine the effect of home blood pressure monitoring on blood pressure levels and proportion of people with essential hypertension achieving targets.

**Design** Meta-analysis of 18 randomised controlled trials.

**Participants** 1359 people with essential hypertension allocated to home blood pressure monitoring and 1355 allocated to the "control" group seen in the healthcare system for 2-36 months.

**Main outcome measures** Differences in systolic (13 studies), diastolic (16 studies), or mean (3 studies) blood pressures, and proportion of patients achieving targets (6 studies), between intervention and control groups.

**Results** Systolic blood pressure was lower in people with hypertension who had home blood pressure monitoring than in those who had standard blood pressure monitoring in the healthcare system (standardised mean difference 4.2 (95% confidence interval 1.5 to 6.9) mm Hg), diastolic blood pressure was lower by 2.4 (1.2 to 3.5) mm Hg, and mean blood pressure was lower by 4.4 (2.0 to 6.8) mm Hg. The relative risk of blood pressure above predetermined targets was lower in people with home blood pressure monitoring (risk ratio 0.90, 0.80 to 1.00). When publication bias was allowed for, the differences were attenuated: 2.2 (–0.9 to 5.3) mm Hg for systolic blood pressure and 1.9 (0.6 to 3.2) mm Hg for diastolic blood pressure.

**Conclusions** Blood pressure control in people with hypertension (assessed in the clinic) and the proportion achieving targets are increased when home blood pressure monitoring is used rather than standard blood pressure monitoring in the healthcare system. The reasons for this are not clear. The difference in blood pressure control between the two methods is small but likely to contribute to an important reduction in vascular complications in the hypertensive population.

### Introduction

High blood pressure is one of the most readily preventable causes of stroke and other cardiovascular complications.<sup>1-4</sup> It can be easily detected, and most cases have no underlying detectable cause; the most effective way to reduce the associated risk is to reduce the blood pressure. Unlike many other common, chronic conditions, we have very effective ways of treating high blood pressure and we have clear evidence of the benefits of such interventions.<sup>1</sup> However, despite a great deal of time and effort, hypertension is still underdiagnosed and undertreated.<sup>5</sup> Furthermore, losses to follow up are high and are responsible for avoidable vascular deaths.<sup>6</sup>

Blood pressure is usually measured and monitored in the healthcare system by doctors or nurses in hospital outpatient departments and, increasingly, in primary care settings. New electronic devices have been introduced and validated in the clinical setting to replace the mercury sphygmomanometer and to overcome the large variations in measurement due to variability between observers. Ambulatory blood pressure monitoring is also being used more often to assess individuals' blood pressures outside the clinical setting.

Measuring blood pressure at home is becoming increasingly popular with both doctors and patients.<sup>7,8</sup> Some national and international guidelines also recommend home monitoring in certain circumstances.<sup>9</sup> A recent qualitative review of the role of home blood pressure measurement in managing hypertension concluded that no evidence exists as to whether home monitoring leads to better control of high blood pressure.<sup>10</sup>

We reviewed the literature on home blood pressure monitoring and did a meta-analysis of the effect of home monitoring on blood pressure levels and the control of hypertension in randomised trials that compared home or "self" blood pressure monitoring and usual blood pressure monitoring in the healthcare system.

### Methods

#### Identification and selection of trials


To identify published trials that met the inclusion criteria we searched Medline (1966 to January 2003) and Embase (1980 to January 2003) for randomised controlled trials of home or self blood pressure monitoring in people with high blood pressure (see appendix A on [bmj.com](http://bmj.com) for strategy). We also searched the Cochrane Database of Systematic Reviews, the Database of Abstracts of Clinical Effectiveness, the Health Technology Assessment Database, the NHS Economic Evaluation Database, the TRIP database, and the websites of the Centre for Reviews and Dissemination and the Agency for Healthcare Research and Quality for reviews of blood pressure monitoring studies. Finally, we examined reference lists of the relevant reviews and all identified studies and reviewed the cited literature. We extended the search to all languages.

We included studies in which the intervention under test was at least one measurement of blood pressure at home by study participants or their family members, whether the result was recorded by the participant or transmitted to a healthcare provider.<sup>11-31</sup> We excluded studies that were not randomised con-



Appendices A-D are on [bmj.com](http://bmj.com)

## Primary care



**Fig 1** QUORUM statement flow diagram. RTC=randomised controlled trial

trolled trials and those that used “ambulatory” blood pressure monitoring rather than “home” or “self” blood pressure monitoring. When several publications reported aspects of the same study, we chose only one paper to represent the trial data on blood pressure control or on achievement of hypertension targets. Where endpoints were presented at different time points—for example, Earp et al and Stahl et al with endpoints at one and two years of follow up<sup>17 19</sup>—we repeated analyses with the alternative time point.

We extracted data from text, tables, and graphs. Two reviewers (SMK and LF) independently examined the data. Differences about inclusion of studies and interpretation of data were resolved by arbitration (FPC), and consensus was reached after discussion. We found 253 references. Twenty one studies met the inclusion criteria.<sup>11–31</sup> We excluded three of these studies because they did not use blood pressure as an outcome measure<sup>11–13</sup> (fig 1).

### Outcome measures

We assessed change in blood pressure (systolic, diastolic, and mean) between intervention and control arms as mean (SD) and change in the proportion of people with blood pressure above target (see appendix B on [bmj.com](http://bmj.com) for methods of assessment of outcome). We used target blood pressure as defined in each paper (see appendix C on [bmj.com](http://bmj.com) for targets used in each study); older studies used diastolic blood pressure only (90 or 95 mm Hg), and others used systolic pressure of 140 mm Hg and diastolic pressure of 90 mm Hg (see appendices D1–D3 on [bmj.com](http://bmj.com) for detailed blood pressure values and effects in each study).

### Statistical analysis

We used a random effects model (StataCorp, College Station, TX, USA) for the meta-analysis of the difference in change in systolic blood pressure, diastolic blood pressure, or mean arterial pressure. Where standard deviation of the change was not reported or could not be calculated from the 95% confidence interval, we estimated it. As the standard deviation of the change was approximately the same as the standard deviations of the initial and follow up blood pressures in studies in which these

were reported, we estimated the standard deviation of the change as the average of the standard deviations of the initial and follow up pressures where only the standard deviation of the change was missing. If no standard deviations were reported then we used the average standard deviation for all the remaining studies. We used relative risk to estimate the effect of intervention on the percentage of patients with blood pressure above target at follow up.

We assessed potential publication bias by using a funnel plot and Egger’s test.<sup>32</sup> Publication bias is due to small negative studies failing to be accepted for publication, which then causes the funnel plot to display asymmetry. We recalculated the combined estimate after estimating from the asymmetry of the funnel plot the number of “missing” studies and their effect sizes and standard errors, a method known as “trim and fill.”<sup>33 34</sup> We assessed heterogeneity between trials by using the  $\chi^2$  test.

## Results

We identified 18 randomised controlled trials that compared blood pressure control or the proportion of people with blood pressure above target. The table shows the characteristics of the analysed trials. Six were based in hospital outpatient clinics,<sup>14 19 21 22 25 31</sup> eight in communities and general practices,<sup>16 18 23 24 26–29</sup> and four in mixed settings.<sup>15 17 20 30</sup> Treatment in the “control” group was mainly “usual” or “standard” care,<sup>15–19 21 22 24–29 31</sup> but some trials had nurse clinics,<sup>14 30</sup> educational interventions,<sup>20</sup> or flagged medical records.<sup>23</sup> Trials used different methods of home or self blood pressure monitoring. In total, 1359 people were randomised to home or self blood pressure monitoring and 1355 to a control group of blood pressure monitoring by health professionals in clinical settings. Two trials used a factorial design,<sup>16 18</sup> four had more than two randomised groups,<sup>17 19 20 29</sup> and one was randomised in clusters.<sup>23</sup> Only in eight trials was outcome assessment stated to have been blind,<sup>14–16 24 25 29 31</sup> and only in nine was randomisation concealed.<sup>15–18 20 21 24 29 31</sup> The duration of the intervention varied between two months<sup>31</sup> and 36 months.<sup>19</sup>

### Systolic blood pressure

Thirteen studies reported systolic blood pressure at follow up and baseline or the change from baseline (see appendix D1 on [bmj.com](http://bmj.com)), but only five of these studies reported full data on means and the standard deviation of the difference. For the remaining seven studies we estimated standard deviations. The overall effect of intervention was 4.2 (95% confidence interval 1.5 to 6.9) mm Hg, with highly significant heterogeneity between studies ( $P < 0.001$ ) (fig 2, top panel). The funnel plot showed some asymmetry, and Egger’s test for publication bias was significant ( $P = 0.038$ ) (fig 3, top panel). The trim and fill method estimated three missing studies and gave a revised estimate of 2.2 (–0.9 to 5.3) mm Hg.

### Diastolic blood pressure

Sixteen studies reported diastolic blood pressure at follow up and baseline or the change from baseline (see appendix D2 on [bmj.com](http://bmj.com)), but only eight of these studies reported full data on means and the standard deviation of the difference. For the remaining eight studies we estimated standard deviations. One study had multiple endpoints.<sup>19</sup> We included results from the one year endpoint. Use of the two year endpoint did not make an important difference to the results (2.2 (1.0 to 3.3) mm Hg). The overall effect of intervention was 2.4 (1.2 to 3.5) mm Hg, with significant heterogeneity between studies ( $P = 0.014$ ) (fig 2, middle panel). The funnel plot showed some asymmetry (fig 3, bottom

## Characteristics of trials included in meta-analysis of home or self blood pressure monitoring

Author and country	Setting	Age group (years)	Definition of hypertension	Length of intervention (months)	Intervention group	Control group
Carnahan 1975, USA <sup>14</sup>	Hospital outpatient	Not stated	DBP >90 mm Hg	6	Home BP self recorded twice daily	Nurse clinic
Haynes 1976, Canada <sup>15</sup>	Workplace	Not stated	DBP ≥90 mm Hg	6	Daily BP self recorded on chart	"Usual care"
Johnson 1978, Canada <sup>16</sup>	Community	35-65	On BP treatment at baseline and DBP ≥95 mm Hg	6	Daily BP self recorded on chart	Home visits or "usual care"
Earp 1982 <sup>17</sup>	Hospital outpatient and general practice	Not stated	Not stated	18	Daily or twice weekly BP by family member	Home visits
Pierce 1984, Australia <sup>18</sup>	General practice	<70	SBP >160 mm Hg or DBP >95 mm Hg	6	Daily BP self recorded on chart	"Usual care" or health education programme
Stahl 1984, USA <sup>19</sup>	Hospital outpatient	16-70	3xDBP ≥95 mm Hg if >30 years; DBP >90 mm Hg if 16-30 years; 2xDBP >100 mm Hg; DBP >120 mm Hg	36	Daily BP self recorded at home	"Standard care"
Binstock 1988, USA <sup>20</sup>	Not clear	Not stated	"Documented" hypertension on treatment	9	Home BP monitoring	Educational intervention
Midanik 1991, USA <sup>21</sup>	Hospital outpatient	Not stated	SBP <180 mm Hg and DBP 90-99 mm Hg	12	Twice weekly BP self recorded at home	Standard care (not measuring BP at home)
Soghikian 1992, USA <sup>22</sup>	Hospital outpatient	Not stated	Not stated	12	Twice weekly BP self recorded at home	"Usual care"
Muhlhauser 1993, Germany <sup>23</sup>	General practice	30-60	2xBP >160 or >95 mm Hg	18	Twice daily BP self recorded at home	"Usual care" with flagged notes
Friedman 1996, USA <sup>24</sup>	Community	≥60	On BP treatment at baseline and SBP ≥160 or DBP ≥90 mm Hg	6	Weekly BP self recorded at home	Regular medical care
Zarke 1997, Canada <sup>25</sup>	Hospital outpatient	18-80	Not stated	2	Twice daily BP self recorded at home	Standard office based care
Bailey 1999, Australia <sup>26</sup>	General practice	Not stated	Not stated	2	Twice daily BP self recorded at home	"Usual care"
Mehos 2000, USA <sup>27</sup>	General practice	≥35	SBP 140-179 mm Hg or DBP 90-109 mm Hg	6	Daily BP self recorded at home	No home monitoring
Vetter 2000, Switzerland <sup>28</sup>	General practice	18-85	SBP 160-200 and DBP 95-115 mm Hg and losartan	2	Twice daily BP self recorded at home before and 12 hours after treatment	Doctor's office
Artinian 2001, USA <sup>29</sup>	Community	Not stated	SBP ≥140 or DBP ≥90 mm Hg; if diabetes or myocardial infarction, SBP ≥130 or DBP ≥85 mm Hg	3	BP self recorded at home three times a week	"Usual care"
Broege 2001, USA <sup>30</sup>	Community and hospital outpatient	≥65	SBP >150 and DBP <90 (on treatment) or >90 mm Hg (not on treatment)	3	BP self recorded at home every other day (also monthly clinic visits)	Fortnightly nurse clinic
Rogers 2001, USA <sup>31</sup>	Hospital outpatient	≥18	Between ≥130 or ≥85 mm Hg and ≥180 or ≥110 mm Hg, depending on complications	2-7	BP self recorded at home three times a week	Usual outpatient care

BP=blood pressure; DBP=diastolic blood pressure; SBP=systolic blood pressure.

panel) (Egger's test for publication bias,  $P = 0.095$ ). The trim and fill method estimated two missing studies and gave a revised estimate of 1.9 (0.6 to 3.2) mm Hg.

### Mean arterial pressure

Three studies reported mean arterial pressure, one of which did not report either systolic or diastolic blood pressure.<sup>25</sup> All studies reported change from baseline (see appendix D3 on [bmj.com](http://bmj.com)) with standard deviation of the difference. The overall effect was 4.4 (2.0 to 6.8) mm Hg, with no significant heterogeneity ( $P = 0.319$ ) (fig 2, bottom panel).

### Blood pressure above target


Six studies reported the number of patients whose blood pressure was controlled at follow up. Different definitions of blood pressure control were used (see appendix C on [bmj.com](http://bmj.com)). Two studies reported the outcome at more than one time point. The analysis reported here is for the one year outcome in both studies. The overall relative risk was 0.90 (0.80 to 1.00), with no significant heterogeneity between studies ( $P = 0.34$ ) (fig 4). Inclusion of the two year outcomes for Earp<sup>17</sup> and Stahl<sup>19</sup> slightly reduced the effect—relative risk 0.92 (0.83 to 1.04).

## Discussion

### Main findings

The meta-analysis of 18 randomised controlled clinical trials found that "self" blood pressure monitoring at home results in better blood pressure control and greater achievement of blood pressure targets than "usual" blood pressure monitoring in the healthcare system. The size of the difference is rather small from the clinical viewpoint: 2.2/1.9 mm Hg (when allowing for publication bias), with 10% greater proportion on target. However, this may represent an adjunctive useful improvement in management of hypertension likely to contribute to a better outlook for cardiovascular events. The main inclusion criterion in the study was that participants had undertaken blood pressure monitoring at home either by themselves or with the aid of a family member. As this is the likely scenario for implementation in a population setting, the results of our meta-analysis could be applicable to the general population of people with mild to moderate essential hypertension.

## Primary care



**Fig 2** Standardised mean differences (95% confidence interval) in systolic (top), diastolic (middle), and mean (bottom) blood pressures achieved in people monitoring blood pressure at home compared with people whose blood pressure was monitored by health professionals in clinical settings

### Limitations of the study

The studies included in the quantitative review were done in a variety of settings, with different methods, using different criteria and different comparative groups. Any potentially consistent effect might have been underestimated. Furthermore, despite our adjustments with statistical methods, the likelihood of publi-

cation bias cannot be excluded. The analysis of hypertension targets may not be easily extrapolated to today's recommended targets of national and international guidelines, because different thresholds were used in different studies.






Fig 3 Funnel plots for systolic and diastolic blood pressure

### Implications

Home blood pressure monitoring has been shown to be feasible; acceptable to patients, nurses, and doctors in general practice; and more suitable for the screening of “white coat” hypertension than ambulatory blood pressure monitoring.<sup>35 36</sup> The white coat effect is important in the diagnosis and treatment of hypertension, even in a primary care setting, and is not a research artefact.<sup>37</sup> Either repeated measurements by health professionals or ambulatory or home measurements may substantially improve estimates of blood pressure and management and control of hypertension. Home blood pressure measurements are the most acceptable method to patients and are preferred to either readings in the surgery or ambulatory monitoring.<sup>38 39</sup> They provide accurate blood pressure measurements in most patients, although some patients of low educational level may have poor reporting accuracy.<sup>39</sup> Finally, blood pressure monitoring at home might help to improve awareness and concordance, and thus overall effective management.

### What is already known on this topic

Blood pressure is usually measured and monitored in the healthcare system by health professionals

With the introduction and validation of new electronic devices, self blood pressure monitoring at home is becoming increasingly popular

No evidence exists as to whether use of home monitoring is associated with better control of high blood pressure

### What this study adds

Patients who monitor their blood pressure at home have a lower “clinic” blood pressure than those whose blood pressure is monitored in the healthcare system

A greater proportion of them also achieve blood pressure targets when assessed in the clinic

After we submitted our manuscript, a multicentre randomised trial was published that compared the use of blood pressure measurements taken in the physician’s office and at home and the potential impact on the management of hypertension.<sup>40</sup> After a year, home blood pressure levels were lower than office blood pressures. Adjustment of antihypertensive treatment on the basis of home blood pressure instead of office blood pressure led to less intensive drug treatment and lower costs. Less good blood pressure control as judged by office blood pressure targets was obviously recorded. At variance with this trial, our results indicate that the practice of monitoring blood pressure “at home” leads to a better control of blood pressure “in the clinic.” Nevertheless, the results of our systematic review and of the latest trial highlight the need for further evidence from prospective studies of outcome to inform potential modifications of treatment guidelines.

### Conclusions

We conclude that blood pressure monitoring by patients at home is associated with better blood pressure values and improved control of hypertension than usual blood pressure monitoring in the healthcare system. As home blood pressure monitoring is now feasible, acceptable to patients, and reliable for most of them,<sup>41</sup> it could be considered as a useful, though adjunctive, practice to involve patients more closely in the management of their own blood pressure and help to manage their hypertension more effectively.




Fig 4 Standardised relative risk of blood pressure above target in people monitoring blood pressure at home compared with people whose blood pressure was monitored by health professionals in clinical settings

## Primary care

FPC is a member of the St George's Cardiovascular Research Group.

Contributors: FPC, LE, and AD conceived the idea, set the objectives, and contributed to design and interpretation. LF ran the searches. SMK and LF abstracted the data, consulting FPC when necessary. SMK did the statistical analysis. FPC drafted the paper, and all authors reviewed it. FPC is the guarantor.

Funding: Bazian Ltd was supported, in part, by an educational grant from Bristol-Myers Squibb.

Competing interests: Bazian Ltd is an independent company that specialises in evidence based reviews and training. It has a policy of strict scientific integrity and does not accept contracts that threaten impartiality when assessing and reviewing research. Bazian Ltd provides managing editorship to publications of the BMJ Publishing Group.

Ethical approval: Not needed.

- Collins R, Peto R, MacMahon S, Hebert P, Fiebach NH, Eberlein KA, et al. Blood pressure, stroke and coronary heart disease. II. Short term reductions in blood pressure: overview of randomised drug trials in their epidemiological context. *Lancet* 1990;335:827-38.
- MacMahon S, Peto R, Cutler J, Collins R, Sorlie P, Neaton J, et al. Blood pressure, stroke and coronary heart disease. I. Effect of prolonged differences in blood pressure: evidence from nine prospective observational studies corrected for the regression dilution bias. *Lancet* 1990;335:765-74.
- Prospective Studies Collaboration. Age-specific relevance of usual blood pressure to vascular mortality: a meta-analysis of individual data for one million adults in 61 prospective studies. *Lancet* 2002;360:1903-13.
- Prospective Studies Collaboration. Cholesterol, diastolic blood pressure, and stroke: 13,000 strokes in 450,000 people in 45 prospective cohorts. *Lancet* 1995;346:1647-53.
- Primates P, Brookes M, Poulter NR. Improved hypertension management and control: results from the health survey for England 1998. *Hypertension* 2001;38:827-32.
- Du X, Cruickshank K, McNamee R, Sarace M, Sourbutts J, Summers A, et al. Case-control study of stroke and the quality of hypertension control in north west England. *BMJ* 1997;314:272-6.
- Yarows SA, Staessen JA. How to use home blood pressure monitors in clinical practice. *Am J Hypertens* 2002;15:93-6.
- Yarows SA, Julius S, Pickering TG. Home blood pressure monitoring. *Arch Intern Med* 2000;160:1251-7.
- Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL Jr, et al. The seventh report of the joint national committee on prevention, detection, evaluation, and treatment of high blood pressure: the JNC 7 report. *JAMA* 2003;289:2560-71.
- Rickerby J. The role of home blood pressure measurement in managing hypertension: an evidence-based review. *J Hum Hypertens* 2002;16:469-72.
- Cartwright W, Dalton KJ, Swindells H, Rushant S, Mooney P. Objective measurement of anxiety in hypertensive pregnant women managed in hospital and in the community. *Br J Obstet Gynaecol* 1992;99:182-5.
- Glanz K, Kirscht JP, Rosenstock IM. Linking research and practice in patient education for hypertension: patient responses to four educational interventions. *Med Care* 1981;19:141-52.
- Ross-McGill H, Hewison J, Hirst J, Dowswell T, Holt A, Brunskill P, et al. Antenatal home blood pressure monitoring: a pilot randomised controlled trial. *Br J Obstet Gynaecol* 2000;107:217-21.
- Carnahan JE, Nugent CA. The effects of self monitoring by patients on the control of hypertension. *Am J Med Sci* 1975;269:69-73.
- Haynes RB, Sackett DL, Gibson ES, Taylor DW, Hackett BC, Roberts RS, et al. Improvement of medication compliance in uncontrolled hypertension. *Lancet* 1976;i:1265-8.
- Johnson AL, Taylor DW, Sackett DL, Dunnett CW, Shimizu AG. Self-recording of blood pressure in the management of hypertension. *CMAJ* 1978;119:1034-9.
- Earp JA, Ory MG, Strogatz DS. The effects of family involvement and practitioner home visits on the control of hypertension. *Am J Public Health* 1982;72:1146-54.
- Pierce JP, Watson DS, Knights S, Gliddon T, Williams S, Watson R. A controlled trial of health education in the physician's office. *Prev Med* 1984;13:185-94.
- Stahl SM, Kelley CR, Neill PJ, Grim CE, Mamlin J. Effects of home blood pressure measurement on long-term BP control. *Am J Public Health* 1984;74:704-9.
- Binstock ML, Franklin KL. A comparison of compliance techniques on the control of high blood pressure. *Am J Hypertens* 1988;1:192-4S.

- Midanik LT, Resnick B, Hurley LB, Smith EJ, McCarthy M. Home blood pressure monitoring for mild hypertensives. *Public Health Rep* 1991;106:85-9.
- Soghikian K, Casper SM, Fireman BH, Hunkeler EM, Hurley LB, Tekawa IS, et al. Home blood pressure monitoring: effect on use of medical services and medical care costs. *Med Care* 1992;30:855-65.
- Muhlhauser I, Sawicki PT, Didjurgeit U, Jorgens V, Trampisch HJ, Berger M. Evaluation of a structured treatment and teaching programme on hypertension in general practice. *Clin Exp Hypertens* 1993;15:125-42.
- Friedman RH, Kazis LE, Jette A, Smith MB, Stollerman J, Torgerson J, et al. A telecommunications system for monitoring and counseling patients with hypertension: impact on medication adherence and blood pressure control. *Am J Hypertens* 1996;9:285-92.
- Zarnke KB, Feagan BG, Mahon JL, Feldman RD. A randomized study comparing a patient-directed hypertension management strategy with usual office-based care. *Am J Hypertens* 1997;10:58-67.
- Bailey B, Carney SL, Gillies AA, Smith AJ. Antihypertensive drug treatment: a comparison of usual care with self blood pressure measurement. *J Hum Hypertens* 1999;13:147-50.
- Mehos BM, Saseen JJ, MacLaughlin EJ. Effect of pharmacist intervention and initiation of home blood pressure monitoring in patients with uncontrolled hypertension. *Pharmacotherapy* 2000;20:1384-9.
- Vetter W, Hess L, Brignoli R. Influence of self-measurement of blood pressure on the responder rate in hypertensive patients treated with losartan: results of the SVATCH study. *J Hum Hypertens* 2000;14:235-41.
- Artinian NT, Washington OG, Templin TN. Effects of home telemonitoring and community-based monitoring on blood pressure control in urban African Americans: a pilot study. *Heart Lung* 2001;30:191-9.
- Broege PA, James GD, Pickering TG. Management of hypertension in the elderly using home blood pressures. *Blood Press Monit* 2001;6:139-44.
- Rogers MA, Small D, Buchan DA, Butch CA, Stewart CM, Krenzer BE, et al. Home monitoring service improves mean arterial pressure in patients with essential hypertension: a randomized, controlled trial. *Ann Intern Med* 2001;134:1024-32.
- Egger M, Davey SG, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997;315:629-34.
- Sutton AJ, Song F, Gilbody SM, Abrams KR. Modelling publication bias in meta-analysis: a review. *Stat Methods Med Res* 2000;9:421-45.
- Sutton AJ, Duval SJ, Tweedie RL, Abrams KR, Jones DR. Empirical assessment of effect of publication bias on meta-analyses. *BMJ* 2000;320:1574-7.
- Aylett M, Marples G, Jones K. Home blood pressure monitoring: its effect on the management of hypertension in general practice. *Br J Gen Pract* 1999;49:725-8.
- Brueren MM, Schouten HJA, de Leeuw PW, van Montfrans GA, van Ree JW. A series of self-measurements by the patient is a reliable alternative to ambulatory blood pressure measurement. *Br J Gen Pract* 1998;48:1585-9.
- Nordmann A, Frach B, Walker T, Martina B, Battagay E. Reliability of patients measuring blood pressure at home: prospective observational study. *BMJ* 1999;319:1172.
- Little P, Barnett J, Barnsley L, Marjoram J, Fitzgerald-Barron A, Mant D. Comparison of agreement between different measures of blood pressure in primary care and daytime ambulatory blood pressure. *BMJ* 2002;325:254.
- Little P, Barnett J, Barnsley L, Marjoram J, Fitzgerald-Barron A, Mant D. Comparison of acceptability of and preferences for different methods of measuring blood pressure in primary care. *BMJ* 2002;325:258-9.
- Staessen JA, Den Hond E, Celis H, Fagard R, Keary L, Vandenhoven G, et al. Antihypertensive treatment based on blood pressure measurement at home or in the physician's office. *JAMA* 2004;291:955-64.
- Williams B, Poulter NR, Brown MJ, Davis M, McInnes GT, Potter JF, et al. Guidelines for management of hypertension: report of the fourth working party of the British Hypertension Society, 2004—BHS IV. *J Hum Hypertens* 2004;18:139-85.

(Accepted 1 February 2004)

doi: 10.1136/bmj.38121.684410.AE

Department of Community Health Sciences, St George's Hospital Medical School, London SW17 0RE

Francesco P Cappuccio *professor of clinical epidemiology and primary care*

Sally M Kerry *senior lecturer in medical statistics*

Bazian Ltd, London N1 1QP

Lindsay Forbes *public health consultant*

Anna Donald *managing director*

Correspondence to: F P Cappuccio [f.cappuccio@sghms.ac.uk](mailto:f.cappuccio@sghms.ac.uk)

evaluated. We think that trauma units with access to a helical, multiplane, computed tomography scanner should routinely image the entire cervical spine at high resolution since the number needed to treat to detect a further injury beyond directed scanning may be only eight to 22 patients,<sup>20 21 36</sup> and this is the standard recommended in figure 2. This is similarly a level 2-3 recommendation; an urgent need remains for an adequately powered, prospective comparison of these modalities to justify any further recommendations.

We propose the removal of cervical collars and patients' mobilisation as a priority for management. Therefore if plain radiographs and computed tomography imaging do not show evidence of traumatic abnormality and the patient is not expected to be conscious within 48-72 hours, current evidence supports the declaration "cervical spine cleared" without further delay.

We draw attention to the routine inclusion of thoracolumbar plain radiography, where unconscious patients with multiple injuries have a compatible mechanism of injury.

Contributors: CGM had the original idea, searched the literature, and wrote the main text. CGM and GGL managed the cases. EPM is guarantor.

Funding: None.

Competing interests: None declared.

- MacDonald RL, Schwartz ML, Mirich D. Diagnosis of cervical spine injury in motor vehicle crash victims: how many X-rays are enough? *J Trauma* 1990;30:392-7.
- Chiu WC, Haan JH, Cushing BM, Kramer ME, Scalea TM. Ligamentous injuries of the cervical spine in unreliable blunt trauma patients: incidence, evaluation and outcome. *J Trauma* 2001;50:457-63.
- Demetriades D, Charalambides K, Chahwan S, Hampeter D, Alo K, Velmahos G, et al. Nonskeletal cervical spine injuries: epidemiology and diagnostic pitfalls. *J Trauma* 2000;48:724-7.
- Holly LT, Kelly DF, Counelis GJ, Blinman T, McArthur DL, Cryer HG. Cervical spine trauma associated with moderate and severe head injury: incidence risk factors and injury characteristics. *J Neurosurg* 2002;96:S285-91.
- Iida H, Tachibana S, Kithara T, Horiike S, Ohwada T, Fujii K, et al. Association of head trauma with cervical spine injury, spinal cord injury or both. *J Trauma* 1999;46:450-2.
- Blackmore CC, Emerson SS, Mann FA, Koepsell TD. Cervical spine imaging in patients with trauma: determination of fracture risk to optimize use. *Radiology* 1999;211:759-65.
- American College of Surgeons Committee on Trauma. *Advanced trauma life support for doctors student course manual*. Chicago: American College of Surgeons, 1997.
- Reid DC, Hernderson R, Saboe L, Miller JD. Etiology and clinical course of missed spine fractures. *J Trauma* 1987;27:980-6.
- Velmahos GC, Theodorou D, Tatevossian R, Belzbergh H, Cornwell EE 3rd, Berne TV, et al. Radiographic cervical evaluation in the alert, asymptomatic blunt trauma victim: much ado about nothing? *J Trauma* 1996;40:768-74.
- Bachulis BL, Long WB, Hynes GD, Johnson MC. Clinical indications for cervical radiographs in the traumatized patient. *Am J Surg* 1987;153:473-8.
- Ersoy G, Karcioğlu O, Enginbas Y, Eray O, Ayrik C. Are cervical spine X-rays mandatory in all blunt trauma patients? *Eur J Med* 1995;2:191-5.
- Brohi K, Wilson-MacDonald J. Evaluation of unstable cervical injury: a 6 year experience. *J Trauma* 2000;49:76-80.
- Davis JW, Parks SN, CL Detlefs MD, Williams GG, Williams JL, Smith RW. Clearing the cervical spine in obtunded patients: the use of dynamic fluoroscopy. *J Trauma* 1995;39:435-8.
- Morris CG, Mullan B. Clearing the cervical spine after polytrauma, instituting unified management for unconscious victims in the intensive care unit. *Anaesthesia* 2004;59:755-61.
- Nawoczenski DA. Pressure sores: prevention and management. In: Buchanan LE, Nawoczenski DA, eds. *Spinal cord injury—concepts and management approaches*. Baltimore, Williams and Wilkins, 1987:99-121.
- Lieberman IH, Webb JK. Cervical injuries in the elderly. *J Bone Joint Surg* 1994;76:877-81.
- McKinley WO, Jackson AB, Cardenas DD, DeVivo MJ. Long-term medical complications after traumatic spinal cord injury: a regional model systems analysis. *Arch Phys Med Rehabil* 1999;80:1402-10.
- Bauer TM, Öfner E, Just HM, Just H, Daschner FD. An epidemiological study assessing the relative importance of airborne and direct contact transmission of microorganisms in a medical intensive care unit. *J Hosp Infect* 1990;15:301-9.
- Huebener J, Frank U, Kappstein I, et al. Influence of architectural design on nosocomial infection in intensive care units—a prospective 2 year analysis. *Intensive Care Med* 1989;15:179-83.
- Berne JD, Velmahos GC, El-Tawil Q, Demetriades D, Asenso JA, Murray JA, et al. Value of complete cervical computed tomographic scanning in identifying cervical injury in the unevaluable blunt trauma patient with multiple injuries: a prospective study. *J Trauma* 1999;47:896-902.
- Nunez DB, Zuluaga A, Fuentes-Bernardo DA, Rivas LA, Becerra JL. Cervical spine trauma: how much more do we learn by routinely using helical CT? *Radiographics* 1996;16:1307-18.
- Rehabilitation of persons with traumatic brain injury. NIH Consensus Statement 1998 Oct 26-28;16:1-41.
- Grossman MD, Reilly PM, Gillett T, Gillett D. National survey of the incidence of cervical spine injury and approach to cervical spine clearance in US trauma centers. *J Trauma* 1999;47:684-90.
- Gupta KJ, Clancy M. Discontinuation of cervical spine immobilisation in unconscious patients with trauma in intensive care units—telephone survey of practice in south and west region. *BMJ* 1997;314:1652-5.
- Porter KM, Allison KP. The UK emergency department practice for spinal board unloading. Is there conformity? *Resuscitation* 2003;58:117-20.
- Morris CG, McCoy E. Cervical immobilisation collars in ICU: friend or foe? *Anaesthesia* 2003;58:1051-3.
- Ajani AE, Cooper DJ, Scheinkestel CD, Laidlaw J, Tuxen DV. Optimal assessment of cervical spine trauma in severe nonpenetrating closed head injury: a prospective study. *Anaesth Intensive Care* 1998;26:487-91.
- Mawson AR, Biundo JJ Jr, Neville P, Linares HA, Winchester Y, Lopez A. Risk factors for early occurring pressure ulcers following spinal injury. *Am J Phys Med Rehabil* 1998;67:123-7.
- Kato H, Inoue T, Torii S. A new postoperative management scheme for preventing sacral pressure sores in patients with spinal cord injuries. *Ann Plast Surg* 1998;40:39-43.
- O'Sullivan KL, Engrav LH, Maier RV, Pilcher SL, Isik FF, Copass MK. Pressure sores in the acute trauma patient: incidence and causes. *J Trauma* 1997;42:276-8.
- Linares HA, Mawson AR, Suarez E, Biundo JJ. Association between pressure sores and immobilisation in the immediate post-injury period. *Orthopedics* 1987;10:571-3.
- Curry K, Cassidy L. The relationship between extended periods of immobility and decubitus ulcer formation in the acutely spinal cord-injured patient. *J Neurosci Nurs* 1992;24:185-9.
- Mawson AR, Siddiqui FH, Connolly BJ, Sharp CJ, Summer WR, Biundo JJ Jr. Sacral transcutaneous oxygen tensions levels in the spinal cord injured: risk factors for pressure ulcers? *Arch Phys Med Rehabil* 1993;74:745-51.
- Cooke MW. Use of spinal board within the accident and emergency department. *J Accid Emerg Med* 1998;15:108-9.
- Hauswald M, McNally T. Confusing extrication with immobilisation: the inappropriate use of hard spine boards for interhospital transfers. *Air Med J* 2000;19:126-7.
- Morris CG, McCoy E. Clearing the cervical spine in unconscious polytrauma victims: balancing risks and effective screening. *Anaesthesia* 2004;59:464-82.
- EAST Ad Hoc Committee on Practice Management Guideline Development. Practice management guidelines for trauma from the Eastern Association for the Surgery of Trauma. *J Trauma* 1998;44:941-6.
- Eastern Association for the Surgery of Trauma. *Determination of cervical spine stability in trauma patients*. Winston-Salem: EAST, 2000. [www.east.org/tpg/chap3u.pdf](http://www.east.org/tpg/chap3u.pdf) (accessed 22 Jul 2004).

(Accepted 29 June 2004)

## Corrections and clarifications

### New treatments for colon cancer

Some errors occurred in this editorial by Maurice Slevin and Sarah Payne (17 July, pp 124-6). The misspelling of a drug persisted to publication—cetuximab was spelt wrongly throughout the article. The drug's manufacturer, Merck, has pointed out that the drug is licensed in many countries, not just Switzerland and the United States, as the article implied, and marketing authorisation was granted in the European Union at the end of June 2004.

### Health experts warn of disease as worst flooding in years hits Bangladesh

In this news article by Peter Moszynski (31 July, p 247) we misspelt Dhaka (in Bangladesh). We had added an "r" to the end of the word; if we had also omitted the "h" we would have successfully spelt Dakar, the capital of Senegal.

### Blood pressure control by home monitoring: meta-analysis of randomised trials

We let the wrong spelling of QUOROM (Quality of Reporting of Meta-analyses) slip through in this Primary Care paper by Francesco P Cappuccio and colleagues (17 July, pp 145-8).