

SELECTED DRUG COMPLICATIONS AND TREATMENT CONFLICTS IN THE PRESENCE OF COEXISTENT DISEASES

Dimitrios Vassilopoulos, MD, Charles Camisa, MD,
and Robert M. Strauss, MD

Rheumatologists treating patients with various rheumatic diseases are often faced with challenging clinical situations in which the available treatment options can adversely influence the course of a coexistent disease. The propensity of certain rheumatic diseases to involve multiple organs and the potential of various antirheumatic drugs to adversely affect different organs further complicates the selection of the appropriate treatment for the individual patient.

This article focuses on the treatment conflicts that are commonly encountered in the setting of underlying gastrointestinal (GI), liver, skin, kidney, or malignant diseases.

INTESTINAL DISEASES

Among the various medications that rheumatologists prescribe in their daily practice, nonsteroidal anti-inflammatory drugs (NSAIDs)

From the Departments of Rheumatic and Immunologic Diseases (DV) and Dermatology (CC), Section of Clinical Dermatology (CC), Cleveland Clinic Foundation (DV, CC), Cleveland, Ohio; Department of Internal Medicine, Ohio State University College of Medicine (CC), Columbus, Ohio; and Department of Gastrointestinal Oncology, Emory University School of Medicine (RMS), Atlanta, Georgia

have been predominantly associated with a number of complications affecting the upper and lower GI tract and the hepatobiliary and pancreatic systems.⁴⁵ The role of NSAIDs in the development of dyspepsia and peptic ulcer disease is well recognized among rheumatologists and is discussed elsewhere in this article. On the contrary, NSAID-induced, small and large intestinal toxicity is usually unrecognized and probably underreported.⁴⁹

NSAID-Related Small Intestine Disease

NSAID use has been associated with various forms of small intestinal toxicity that varies from ulcer and stricture formation to perforation and bleeding (Table 1).^{21, 45, 124} The most commonly described condition is the NSAID-induced enteropathy, which is characterized by diffuse intestinal inflammation and increased mucosal permeability.¹²⁴ Although it has been reported to occur in up to 70% of patients taking NSAIDs,⁴⁵

Table 1. LOWER GASROINTESTINAL DISEASES ATTRIBUTABLE TO THE USE OF NONSTEROIDAL ANTI-INFLAMMATORY DRUGS

| GI Location | Lesion | Comments |
|--|--|--|
| Small intestine ^{21, 45, 124} | Ulcers Perforation/bleeding Strictures/diaphragms Enteropathy characterized by: Occult blood loss Protein-losing enteropathy Iron deficiency anemia Malabsorption | 8% in a large autopsy study ⁴ Rare Rare Presence of small intestine diaphragms is pathognomonic for NSAID-induced damage Usually mild without significant clinical findings |
| Colon ^{21, 45, 49, 75} | | |
| No pre-existing colonic disease (de novo injury) | Ulcers Strictures Perforation Bleeding Colitis Proctitis caused by suppositories | Rare Cases of NSAID-induced colitis due to a hypersensitivity reaction have been reported (reviewed in ref ⁴⁹) Mainly from indomethacin |
| Pre-existing colonic disease | Exacerbation of quiescent IBD Complications from underlying diverticular disease (bleeding/perforation/peritonitis) | See text for details See text for details |

NSAIDs = nonsteroidal anti-inflammatory drugs; IBD = inflammatory bowel disease

it is usually mild, without significant clinical manifestations. Its pathogenesis is unclear, but it has been proposed that direct NSAID-induced uncoupling of the oxidative metabolism in enterocyte mitochondria could lead to disruption of the small intestine mucosal barrier causing increased intestinal permeability.²⁰¹ NSAID-induced cyclooxygenase inhibition impairs the prostaglandin-dependent regenerative process, leading to prolongation of the increased intestinal permeability. Bacteria, various enzymes, and bile subsequently can penetrate easily the intestinal mucosa, perpetuating the mucosal injury. It has been shown that NSAIDs, such as indomethacin and piroxicam with extensive enterohepatic circulation, are causing the most injury compared with other agents.¹⁹⁷

In patients with underlying rheumatic diseases who develop signs and symptoms of small intestinal injury, discontinuing the offending NSAID is the preferable approach. If this option is not possible or if the injury is mild, NSAIDs with limited enterohepatic circulation such as nabumetone may be tried.¹³⁹ In small trials, sulfasalazine^{23, 90} and metronidazole²² have been shown to decrease NSAID-induced intestinal inflammation and blood loss, but large scale prospective studies supporting this approach have not been reported.

NSAID-Related Large Intestine Disease

Most of the published information on NSAID-induced large intestinal damage has been from case reports or small retrospective studies, hence the true frequency of the NSAID-induced colonic toxicity is unknown (see Table 1). Overall, it appears to be less frequent compared with NSAID-induced small intestine toxicity.^{21, 45} Most of the cases of *de novo* colitis induced by NSAIDs were caused by sustained release NSAIDs, fenemates or diclofenac.^{49, 75} Its pathogenesis remains unclear.

Inflammatory Bowel Disease

NSAIDs have been associated with relapses of quiescent inflammatory bowel disease (IBD).¹¹² This seems to occur more often in patients with ulcerative colitis as compared with patients with Crohn's disease.⁷⁵ The frequency of exacerbation of IBD attributed to NSAIDs is unknown because large retrospective or prospective studies have not been done. In a recent case-control study, Evans et al reported an increased risk of admission to the hospital because of colitis from inflammatory bowel disease, in patients with current (less than 45 days) or recent (up to 6 months) exposure to NSAIDs (overall odds ratio 1.77–1.93).⁵⁶

Different NSAIDs have been implicated in the reactivation of IBD, which histologically can range from mild sigmoid edema to pancolitis.⁷⁵ The time from initiation of NSAID therapy to IBD exacerbation ranges from a few hours to 8 weeks (median less than a week).⁷⁵ Each of the eight patients that has been rechallenged with NSAIDs showed evidence

of disease relapse, indicating a causal relationship between NSAID use and IBD exacerbation.⁷⁵ Most of the patients responded well to discontinuation of the responsible NSAID or to steroid treatment.

Diverticular Disease

Several case reports,⁴⁹ case-control studies,^{33, 40, 96, 128} and a prospective study²²⁴ have implicated NSAID intake with serious complications of diverticular disease, such as perforation and lower gastrointestinal bleeding. In a prospective study, Wilson et al found a statistically increased risk of perforation among patients with diverticular disease taking NSAIDs compared to age- and sex-matched controls from a general practice and patients with colorectal cancer taking NSAIDs.²²⁴ In a recent retrospective case-control study from the USA, Holt et al observed a two-fold increased risk of lower GI bleeding in patients consuming NSAIDs at the time of admission compared with age- or sex-matched controls without evidence of GI disease.⁹⁶

Further substantiating the above observations, a recent retrospective study from Finland found that NSAID-induced perforation or bleeding from the lower GI accounted for 36% of the NSAID-attributable deaths in a large population of rheumatoid arthritis (RA) patients ($n = 1666$).¹⁴⁹

Although large epidemiologic studies linking NSAID use with exacerbation of IBD or with increased risk of complications from diverticular disease are still missing, the authors believe that NSAIDs should not be used in patients with known IBD and used very cautiously (if at all) in patients with known diverticulosis. An increased awareness of their potential life threatening complications is necessary and prompt discontinuation of the medication is indicated in patients who develop these complications.

LIVER DISEASES

Understanding the potential side effects of the antirheumatic medications in patients with compromised liver function is complicated by their occasional direct hepatotoxic effects on the "normal" liver^{25, 210} and the concomitant involvement of the liver by certain autoimmune diseases.

NSAIDs⁶³ and methotrexate (MTX)²²¹ have been most commonly implicated in various forms of liver injury. Other antirheumatic drugs that may cause hepatotoxicity include gold,⁹⁹ D-penicillamine,¹⁸¹ allopurinol,³ azathioprine (AZA),¹¹⁷ and cyclosporine (CsA)¹¹⁷ (Table 2).

RA and systemic lupus erythematosus (SLE) may affect the liver. Elevated alkaline phosphatase levels have been reported in 25% to 50% of patients with RA,²¹⁹ although significant abnormalities in liver biopsies rarely have been noted.²²¹ Liver enzyme abnormalities have been found in 23% to 60% of patients with SLE during the course of their disease^{148, 184} and tend to correlate with disease activity.¹²³ Severe involve-

Table 2. HEPATOTOXICITY OF ANTIRHEUMATIC MEDICATIONS

| Medication | Type of Hepatic Injury | Frequency/Comments |
|-----------------|---|--|
| NSAIDs | | |
| Phenylbutazone* | Hepatocellular | Minor AST/ALT elevation = 1%–15% Severe liver injury = Rare (<0.1%) |
| Diclofenac* | | |
| Etodolac | | |
| Ibuprofen | | |
| Indomethacin | | |
| Fenoprofen | | |
| Piroxicam | | |
| Phenylbutazone* | Cholestatic | |
| Sulindac* | | |
| Naproxen | | |
| Nabumetone | | |
| Fenoprofen | | |
| Diflunisal | | |
| Sulindac* | Mixed | |
| Diflunisal | | |
| Ibuprofen | | |
| Phenylbutazone | Granulomatous | |
| Methotrexate | | Frequency of fibrosis in RA patients: Mild (Roenigk grade IIIA) = 14.3% Moderate (Roenigk grade IIIB) = 0.83% Cirrhosis (Roenigk grade IV) = 0.28% Occasionally manifesting as a hypersensitivity syndrome |
| | Fibrosis | |
| Sulfasalazine | Hepatocellular | Rare, occasionally in older patients with underlying renal failure or on diuretic therapy |
| Allopurinol | Hepatocellular | Rare (overall <0.1%) Mainly in the post-transplant setting |
| | Cholestatic | |
| | Granulomatous | |
| Azathioprine | Cholestatic (most common) | |
| | Hepatocellular | Rare |
| | Vascular liver disease (peliosis hepatis/venooclusive disease/nodular regenerative hyperplasia) | |
| Cyclosporine | Cholestatic (most common) | Rare Mainly in the post-transplant setting (4%–7%), dose dependent |
| | Hepatocellular | |
| Gold | Cholestatic | Rare case reports |
| D-Penicillamine | Cholestatic | Rare case reports |

*Most commonly reported.

AST = aspartate aminotransferase; ALT = alanine aminotransferase; RA = rheumatoid arthritis.

Data from References 3, 57, 62, 63, 99, 117, 120, 181, 210

ment of the liver causing chronic hepatitis, fibrosis, or cirrhosis in SLE is rare (0.8%–2.4% in a recent autopsy series).¹⁴³ Rarely, Sjögren's syndrome, systemic sclerosis, Felty's syndrome, polymyalgia rheumatica/giant cell arteritis, and Still's disease can show evidence of liver involvement.²¹⁹

An overview of the potential interactions of medications commonly used in the clinical practice of rheumatology in patients with underlying liver disease is presented in this article.

Nonsteroidal Anti-inflammatory Drugs

NSAIDs have been associated with minor increases in liver enzymes in 1% to 15% of patients treated with these medications.⁶³ The pathogenesis of the liver injury has been attributed to immunologic or metabolic idiosyncratic mechanisms. Usually asymptomatic elevations in transaminase levels occur during the first few months of therapy and they usually subside after discontinuation of the offending NSAID. Based on the results of large retrospective studies, it seems that the risk of severe liver disease because of NSAIDs is low.^{34, 62, 69}

Among the different NSAIDs, sulindac, diclofenac, and phenylbutazone have been associated with an increased reported incidence of liver injury.^{144, 210} In a recent review of the reported cases of sulindac-induced liver injury, Tarazi et al showed that in 78% of the cases, liver damage occurred during the first 2 months of therapy and in almost half of the cases it was caused by a hypersensitivity type of reaction.²⁰⁷ Similar results have been reported in association with diclofenac or phenylbutazone use.⁶³

Although there are adequate scientific data to suggest a role of NSAID in direct liver injury, our knowledge of their effect in patients with pre-existing liver disease is limited. NSAIDs should be avoided in patients with cirrhosis of the liver and ascites regardless of its origin, because of the increased risk of NSAID-induced renal insufficiency.⁹ Renal failure in these patients is caused by NSAID-induced inhibition of the vasodilating prostaglandins, which results in decreased renal blood flow and glomerular filtration rate (GFR).^{18, 125} Although some studies have shown that misoprostol can minimize the renal dysfunction caused by NSAIDs,^{7, 152} other studies did not reproduce these findings.^{65, 100}

Patients with advanced liver disease characterized by decreased synthesis of coagulation factors or coexistent thrombocytopenia are prone to bleeding that can be exaggerated by the concomitant use of NSAIDs. In these patients NSAIDs should also be avoided.

Whether or not NSAIDs enhance liver damage in patients with underlying chronic viral hepatitis caused by hepatitis C virus (HCV) or hepatitis B virus (HBV) is unclear. There have been no studies examining their direct effect on liver function in these patients. Preliminary data have suggested an enhanced antiviral activity of interferon-alpha (IFN- α) when used in combination with NSAIDs in patients with chronic hepatitis.⁶ Clinical studies have failed to show any additional efficacy in patients with chronic hepatitis C treated with a combination of IFN- α and NSAIDs in comparison with patients treated with IFN- α alone.^{5, 230} In the study by Zarski et al,²³⁰ administration of IFN- α and tenoxicam for 6 months did not result in worsened liver function (biochemical or histological) or in increased HCV RNA levels compared with patients treated with IFN- α alone.

Although it is difficult to extrapolate from these limited data, it seems that NSAIDs can be safely used in patients with chronic hepatitis (without evidence of cirrhosis) with close monitoring of the transaminase

levels, especially during the first 6 months of therapy.⁶³ NSAIDs with an increased risk for hepatotoxicity (e.g., phenylbutazone, diclofenac, Sulindac, see Table 2) probably should be avoided. NSAIDs should be discontinued if significant elevation of transaminase levels (> 3 times normal value) occurs. If other causes of elevated transaminase levels have been ruled out, the offending NSAID should not be used again.

Corticosteroids

Corticosteroids (CS) are used in the treatment of various liver diseases including autoimmune hepatitis,⁴⁷ granulomatous hepatitis,²³² severe alcoholic hepatitis,¹⁷⁴ and after liver transplantation. Although CS have been associated with the development of fatty liver, significant liver dysfunction from their use is rare.¹¹⁷ CS should be used cautiously in patients with advanced liver failure with or without associated portal hypertension because of increased risk of infection and possibly gastrointestinal bleeding (especially in patients taking NSAIDs at the same time).^{67, 164}

CS treatment leads to enhancement of HBV-antigen expression in hepatocytes,^{129, 212} an increase in viral replication in animals,⁶⁴ an increase in serum hepatitis B surface antigen (HBsAg) and DNA polymerase levels in humans,¹⁹³ and clinical deterioration of patients with chronic HBV infection.^{97, 126} Although liver inflammation decreases during treatment, withdrawal of steroids is usually followed by a rebound increase in transaminase levels because of an enhanced cellular immune response against the HBV-infected hepatocytes.⁹⁷ In some cases, this rebound phenomenon can lead to viral clearance²²⁰ but can also cause fulminant hepatitis and death.⁹⁷ Antiviral therapy (IFN- α) following pretreatment with CS, has not been proven to be of additional benefit for the treatment of chronic hepatitis B,³⁸ but this regimen was not associated with an increased rate of liver decompensation.¹⁶¹ These data emphasize the need for cautious use of CS in patients with chronic HBV infection, with close clinical and laboratory monitoring, especially after withdrawal of therapy. Antiviral therapy should be considered for each patient receiving steroids and consultation with a hepatologist is recommended, particularly in patients with HBV-associated polyarteritis nodosa.¹⁴¹

Taking into account that chronic hepatitis C affects 1% to 2% of the general population⁹³ and is frequently associated with various rheumatological/autoimmune manifestations,⁸⁶ rheumatologists are increasingly encountering patients requiring CS or other forms of immunosuppressive therapy for control of HCV-associated conditions (vasculitis, polyarthritis, renal disease, and so forth) or for the treatment of coexisting rheumatic diseases.

Short courses of CS treatment (1–6 months) increase viremia in HCV infected patients, whereas transaminase levels decrease during therapy.^{60, 138, 209} Fong et al have found that discontinuation of prednisone therapy led to a return of viral RNA and transaminase levels to their pretreat-

ment values in patients with chronic HCV infection.⁶⁰ In contrast to hepatitis B, no cases of fulminant hepatitis were observed.⁶⁰ Furthermore, in HCV-positive patients who underwent kidney transplantation and were treated with different immunosuppressive regimens containing CS for 1 to 7 years, although elevations of transaminases and viral titers were seen, severe compromise of the liver function was rare (0%–4.3%).^{114, 183} Addition of IFN to methylprednisolone in a long-term randomized controlled study of patients with HCV-associated mixed cryoglobulinemia prevented the rise of HCV RNA levels.⁴⁸

These data suggest that although treatment with steroids leads to increased viremia in patients with hepatitis C, fulminant hepatitis or severe decompensation of the liver function is rare. Similarly to patients with hepatitis B, close monitoring of liver function and concomitant antiviral therapy (IFN \pm ribavirin) should be considered in every patient with HCV infection receiving CS treatment.

Cyclosporine

CsA, an immunosuppressive agent that has been used successfully in the organ transplantation field for the last 20 years, has been recently included in the armamentarium of agents used for the treatment of various autoimmune and rheumatologic conditions.¹²⁷ Hepatotoxicity related to CsA use is rare¹¹⁷; it is usually manifested as a cholestatic syndrome with or without associated mild elevation of transaminases. A possible association with the formation of gallstones and biliary sludge also has been suggested. Most of the cases of CsA-induced hepatotoxicity have been reported in the post-transplant setting (ranging from 4%–7%), especially when a high dose of the medication was used.¹¹⁷ The liver abnormalities usually subside after reduction of the dose of CsA.

CsA acts mainly by inhibiting interleukin-2 production by T lymphocytes, an essential cytokine for an effective immune response. Its potent immunosuppressive action can increase the risk of infection in all patients. Given its known nephrotoxic effect,¹⁹⁵ extreme caution should be paid to the renal function of patients with severe liver dysfunction who may have abnormal renal function.

The effect of CsA in patients with chronic liver disease has not been studied extensively. According to recent guidelines, CsA use is not recommended in patients with elevated liver enzymes (> 2 times the normal values).¹⁵⁷ Nevertheless, CsA has shown beneficial effects in patients with autoimmune hepatitis resistant to CS or AZA^{54, 131, 162} and has been used with moderate success in patients with primary biliary cirrhosis (PBC) although an increased incidence of nephrotoxicity was noted.¹³³

The impact of CsA in patients with chronic hepatitis B is controversial. CsA does not cause increased HBV antigen expression in hepatocyte cultures from patients with chronic hepatitis B¹²⁹ and does not lead to

increased rates of chronicity in an animal model of hepatitis B.⁴² Most of the available data in humans are derived from the study of patients with HBV infection receiving immunosuppressive regimens containing CsA after renal transplantation. Unfortunately, the interpretation of these data is complicated by the concomitant use of CS or AZA by these patients. Huang et al observed a 50% incidence of chronic hepatitis in renal allograft recipients treated with CsA and prednisone but the mortality rate was low (5%).¹⁰¹ These results were not different from patients treated with prednisone and AZA. Similarly, Stempel et al found no increased mortality in HBsAg-positive renal transplant patients treated with CsA and prednisone over a 4-year period.²⁰⁴ On the other hand, Kliem et al followed renal transplant patients for almost 7 years and found that 60% of HBsAg-positive patients died from liver failure compared with 2% of HBsAg-positive patients.¹¹⁴ Furthermore, Bang et al reported a 25% mortality in HBsAg-positive patients treated with CsA/prednisone (45% because of liver-related diseases) compared with a mortality of 7.8% in a similar group of HCV positive patients (0% from liver diseases).¹³

A milder course of hepatitis C after kidney transplantation in patients receiving different immunosuppressive regimens containing CsA has been suggested from different studies,^{13, 114, 183} although in some studies an increased incidence of sepsis was found.^{114, 159} Elevated levels of CsA metabolites in patients with chronic hepatitis C receiving CsA have been reported, but the clinical significance of these findings is unclear.^{98, 114} In the nontransplant setting, CsA given for 3 months at a dose of 1.5 to 4.0 mg/kg/day, did not have any effect in the HCV RNA levels in patients with chronic hepatitis C.¹⁰⁸ CsA also has been used successfully for the treatment of HCV-associated cryoglobulinemia without any reported worsening of liver function.¹²

In conclusion, CsA appears to have a milder effect on the clinical course of patients with chronic viral hepatitis. Close monitoring of liver and kidney function is mandatory in these patients as is a high alert state for the development of infections.

Methotrexate

MTX is currently one of the most commonly used medications for the treatment of RA. Its hepatotoxicity has been well documented, although significant controversy exists regarding its frequency and severity.²²¹

The American College of Rheumatology (ACR) recently has published guidelines regarding the monitoring for hepatotoxicity in patients with RA receiving MTX.¹²⁰ In patients with history of excessive alcohol consumption, persistently elevated aspartate aminotransferase (AST) levels or with underlying chronic hepatitis B or C infection, consultation with a gastroenterologist for consideration for a pretreatment liver biopsy is recommended.¹²⁰ If there is evidence of persistent AST abnormal-

ities over a year, a repeat liver biopsy is recommended. Presence of advanced histologic changes, such as moderate to severe fibrosis or cirrhosis (Roenigk grade IIIB or IV, respectively),¹⁸⁰ in the liver biopsy is an indication for discontinuation of therapy.¹²⁰

When MTX is used for treatment of psoriasis or psoriatic arthritis, a higher incidence of liver toxicity has been observed.¹⁷⁹ Based on these findings, a recent consensus conference proposed that patients without risk factors for liver disease should undergo a liver biopsy when 1 to 1.5 gm of cumulative MTX has been reached and thereafter at intervals of approximately 1.5 gm of cumulative MTX dose.¹⁷⁹ In patients with risk factors for liver disease, such as excessive alcohol consumption, persistent abnormal liver chemistry studies, family history of inheritable liver diseases, history of liver diseases like chronic hepatitis B or C, diabetes mellitus, obesity, or history of significant exposure to hepatotoxic drugs or chemicals, an initial liver biopsy is indicated during the first 2 to 4 months of therapy, with repeat biopsies at 1.0 to 1.5, 3.0, and 4.0 gm of cumulative MTX dose. Similarly to the ACR guidelines, discontinuation of MTX is recommended if severe fibrosis or cirrhosis (Roenigk grade IIIB or IV) are present.

Although these guidelines emphasize the importance of close monitoring in patients with pre-existing liver diseases, the mere presence of an underlying liver disease should not be a contraindication for treatment. MTX should be avoided in patients with advanced liver disease or cirrhosis regardless of its cause because of concerns of increased risk of infection and drug-related toxicities (renal, pulmonary, bone marrow, liver, and so forth). MTX also should be avoided in patients with underlying alcoholic liver disease because that represents a significant risk factor for MTX-induced hepatotoxicity.²²¹ On the other hand, MTX has been used over the last decade in patients with various inflammatory liver diseases including PBC,¹¹⁰ primary sclerosing cholangitis,²²² and idiopathic granulomatous hepatitis¹¹⁵ with promising results.

Although a case report of fulminant hepatitis caused by reactivation of hepatitis B infection after discontinuation of low-dose MTX has been reported,⁵⁹ the short and long term effects of MTX treatment in patients with underlying chronic hepatitis B or C are unknown. As mentioned previously, a baseline liver biopsy is indicated and then a decision should be made, balancing the risk to benefit ratio for each individual patient. Close monitoring of liver function and virologic parameters, as discussed with the other immunosuppressive agents, is mandatory.

Azathioprine

Liver disease caused by AZA is rare (< 0.1%).⁵⁷ AZA-induced hepatotoxicity has been more commonly reported in the post-transplant setting and it can present with a wide spectrum of manifestations, including asymptomatic elevation of liver tests, cholestasis, cholestatic hepatitis, and vascular liver disease (peliosis hepatis, veno-occlusive

disease and nodular regenerative hyperplasia).¹¹⁷ Despite its potential hepatotoxic effect, AZA is being used successfully for the treatment of autoimmune hepatitis⁴⁶ particularly, as a steroid-sparing agent.

In patients with advanced liver disease, the same precautions that have been mentioned previously for the other immunosuppressive agents apply to AZA. A recent report by Pol et al¹⁶⁶ suggested that AZA-induced hepatitis in renal transplant patients was facilitated by an underlying chronic viral hepatitis B or C. Similarly, Hestin et al found a correlation between AZA use and histologically proven chronic active hepatitis C in renal transplant patients.⁹⁴ Although data from patients with chronic viral hepatitis treated with AZA in the nontransplant setting are lacking, caution is recommended in patients with chronic viral hepatitis receiving AZA therapy. Differentiating between AZA-induced liver injury and exacerbation of the underlying viral process may require liver biopsy in addition to drug discontinuation.

PSORIASIS

The skin is frequently involved in different rheumatologic or autoimmune diseases including SLE, scleroderma, dermatomyositis, psoriatic arthritis (PsA), Reiter's syndrome, Still's disease, sarcoidosis, Behçet's disease, and various vasculitides.⁷⁰ Successful treatment of the underlying rheumatologic condition usually leads to improvement of the coexistent skin lesions. In some cases, treatment of the underlying rheumatic disease has been associated with exacerbations of the skin manifestations. Typically, this has been described in patients with psoriasis undergoing treatment for associated PsA or coexistent rheumatic diseases (e.g., SLE).^{1, 30, 83}

Several medications have been associated with induction or exacerbation of psoriatic lesions. These medications are listed below:

Definite association

- Lithium²
- β -blockers²
- Corticosteroid withdrawal¹⁸⁵
- Antimalarials¹

Possible association

- NSAIDs¹
- Interferons:
 - α ^{19, 66, 68, 74, 107, 134, 140, 227}
 - β ^{116, 218}
 - γ ⁵⁸
- ACE-inhibitors^{43, 76, 87, 102, 206, 226}
- Potassium iodide¹
- Amiodarone¹
- Digoxin¹
- Antibiotics (tetracycline,¹ penicillin,¹ sulfonamides,¹ terbinafine⁸⁴)
- Alcohol¹⁶⁵
- Calcium channel blockers¹¹³

Most of the drug-induced psoriasis cases have been reported as sporadic case reports or as small retrospective studies.^{1, 30} The absence of well-designed, prospective placebo-controlled studies makes the interpretation of the available data problematic.³⁰

Among the medications that rheumatologists frequently prescribe, antimalarials, CS, NSAIDs, and gold have been frequently implicated in the exacerbation of psoriasis.

ANTIMALARIALS

Antimalarials are prescribed to patients with psoriasis for treatment of associated PsA, coexistent SLE or RA, or for prophylaxis against malaria in patients traveling to endemic areas.¹ Although there have been no controlled trials indicating efficacy in PsA, recent reports have suggested a potential role for antimalarials in the treatment of PsA.^{77, 109} Concerns about their potential toxicity have limited their use. In a study by Gladman et al only 5% of patients with PsA were receiving antimalarials.⁷⁸

Earlier reports suggested an increased incidence of psoriasis exacerbation, especially in patients taking quinacrine⁴¹ or chloroquine.^{11, 20, 41, 121, 153, 155, 231} Similarly, there have been several case reports in the literature describing exacerbation of psoriasis, resistance to treatment, and erythroderma, following treatment with hydroxychloroquine (HCQ).^{10, 61, 82, 91, 137, 199, 216} Some *in vitro* studies suggested a potential role of these agents in the induction of epidermal injury in normal and psoriatic skin.^{50, 192, 225} Chloroquine enhances mitogen-induced lymphocyte proliferation¹⁹² and the allostimulatory properties of fresh and cultured epidermal antigen presenting cells from patients with psoriasis.⁵⁰ HCQ has been shown to inhibit the epidermal transglutaminase in normal cultured human skin, leading to epidermal injury and subsequent irregular keratinization of the upper and lower epidermis.²²⁵ Chloroquine, however, in contrast to lithium and propranolol, had no enhancing effect in tyrosine phosphorylation of mitogen-stimulated T cells from psoriatic patients.¹⁵⁴

Clinical studies have portrayed a safer profile for antimalarials in the treatment of psoriatic arthritis. Gladman et al⁷⁷ found no differences in psoriasis exacerbation between a group of patients with PsA treated with chloroquine for 6 months compared to a control group (18% and 25%, respectively). Only 1 out of 32 patients had to discontinue the medication because of psoriasis exacerbation.⁷⁷ Two retrospective studies in patients with PsA treated with HCQ did not show also any significant incidence of skin exacerbations.^{109, 190} Specifically, Kammer et al reported no cases of psoriasis exacerbation in 50 trials of HCQ, although drug-induced cutaneous eruptions developed in 4 cases (8%).¹⁰⁹ Wilke and Sayers, reviewing the experience at the Cleveland Clinic, showed that only 3 of 32 patients (9%) treated with HCQ for 4 to 6 years had a "possible flare" of their psoriatic lesions.²²³

Collectively, these limited data suggest that the incidence of antima-

larial-induced exacerbation of psoriasis is low. It also seems that HCQ is safer compared with chloroquine and quinacrine. Nevertheless, these medications should be used cautiously in psoriatic patients with close monitoring of their skin lesions.³⁰

CORTICOSTEROIDS

The impression that systemic CS aggravate psoriasis is derived from the seminal observations of Ryan and Baker¹⁸⁵ in the treatment of 73 patients with generalized pustular psoriasis. Prednisolone alone was used in 24 patients in doses ranging from 30 to 300 mg/day. Triamcinolone was used in 17 patients in doses of 8 to 60 mg daily. Smaller numbers of patients received betamethasone, ACTH, dexamethasone, or hydrocortisone alone, and 21 patients received varying combinations together or in sequence. Six months or longer seemed to be a typical treatment course. The authors concluded that CS-treated patients tended to experience increasingly severe generalized pustular relapses, often precipitated by attempting to withdraw the steroid. Satisfactory control with acceptable side effects was achieved in only 12 of 73 (17%) patients.

They later expanded their experience to 155 patients and reported that 37 (24%) endured their first attack of generalized pustular psoriasis within a month of first receiving CS or within a month of their withdrawal; 25 (16%) had common plaque-type psoriasis before the pustular phase.¹⁸⁶ They concluded that systemic CS used for plaque or erythrodermic psoriasis may precipitate and perpetuate the severe phase of the disease. Moreover, MTX seemed to be less efficacious in patients previously treated with steroids. MTX was valuable during steroid withdrawal in patients who became refractory to steroids; however, rebound flares after steroid withdrawal were not always controllable with MTX.

The least common form of psoriasis is exfoliative dermatitis or psoriatic erythroderma, representing only 1% to 2% of all forms. Erythrodermic psoriasis usually develops gradually or abruptly during the course of ordinary chronic psoriasis, but it may be the initial manifestation. The most important precipitating factor is thought to be the preceding use of systemic steroids or excessive potent topical steroids (>60 gm/week). Of 50 patients admitted to Baylor University for psoriatic erythroderma, 25 (50%) received excessive topical steroids or systemic steroids (oral or intramuscular).²⁴ Some patients received both oral and intramuscular steroids.

Because of the foregoing indirect clinical evidence, whenever possible, the use of high-dose systemic CS therapy should be eschewed in patients with psoriasis. Their use is associated with the majority of extremely unstable (inflammatory) psoriasis as well as the conversion of ordinary cases into generalized pustular or erythrodermic psoriasis.³¹ Despite these concerns, systemic CS are still used in approximately 20% of patients with PsA.^{78, 211} Short courses of low-dose CS (such as prednisone at 10–20 mg/d) and local intra-articular CS injections may be

justifiable for severe flares of PsA but always in combination with disease modifying agents like MTX, sulfasalazine, AZA, CsA, and so forth. A prior consultation with a dermatologist is recommended for better planning and coordination of the therapeutic strategy for these patients.

Nonsteroidal Anti-Inflammatory Drugs

NSAIDs are the most commonly used medications in the treatment of PsA.^{78, 81} Although there have been anecdotal case reports of psoriasis exacerbations in patients receiving various NSAIDs (including indomethacin,^{111, 168, 194} phenylbutazone,¹⁷⁵ ibuprofen,¹⁵ and oxyphenbutazone¹⁷⁵) two double-blind placebo controlled trials using NSAIDs for treatment of psoriasis did not substantiate these findings.^{55, 118} Kragballe et al compared the efficacy of benoxaprofen (n=20) with placebo (n=20) in an 8-week double-blind randomized trial.¹¹⁸ There was a significant improvement of the psoriatic lesions in the benoxaprofen group compared with placebo, without any cases of worsening psoriasis in the NSAID group. A higher incidence of photosensitivity was observed in patients receiving benoxaprofen (11 of 20) compared with placebo (3 of 19). In a similar study, Ellis et al randomized patients to receive either meclofenamate (n=47) or placebo (n=56) in a double-blind fashion for 4 weeks followed by an open trial of the medication for 4 more weeks.⁵⁵ Although there was no evidence of improvement with meclofenamate, the incidence of worsening psoriatic lesions did not differ between the meclofenamate and the placebo group (13% versus 13% at 4 weeks and 8% versus 10% at 8 weeks, respectively).

Overall, it appears that NSAIDs are well tolerated in patients with psoriasis without significant skin side effects.

Gold

Gold therapy has been shown to have modest efficacy in patients with PsA. Both oral (auranofin) and intramuscular (IM) (gold sodium thiomalate) gold have been used.^{81, 223} In a recent double-blind trial, Palit et al found a marginal improvement of PsA in patients treated with IM gold compared to auranofin.¹⁵⁶ The incidence of psoriasis exacerbation is low for both routes of administration, but it seems to be higher in patients treated with IM gold. In a recent review of published studies, Bruckle et al reported an overall incidence of 3.4% of psoriasis exacerbation in patients receiving auranofin.²⁶ In contrast, in patients treated with IM gold, the incidence ranges from 0% to 18% (7.8% from 4 published studies).^{26, 52, 156, 177}

Gold therapy is well tolerated in patients with PsA, although its efficacy is modest. Auranofin has a lower incidence of psoriasis exacerbations but is less effective compared with IM gold.¹⁰⁵

Treatment Conflicts in the Presence of Coexistent SLE

Based on the prevalence of psoriasis in the US and UK population (1%–2%),³² the coexistence of all forms of lupus with psoriasis seems to be less than expected. Dubois reported that 0.6% of 520 patients with SLE had concurrent psoriasis.⁵³ The coexistence of psoriasis and lupus raises issues of diagnosis and treatment.⁹¹ Subacute cutaneous lupus erythematosus (SCLE) lesions may be either annular or psoriasiform. Biopsy, direct immunofluorescence, and serologic testing are necessary to distinguish some cases of SCLE from psoriasis.

Photochemotherapy with longwave ultraviolet (UV) light and oral 8-methoxypsoralen (PUVA) is being used for the treatment of psoriasis and has shown some benefit in the treatment of patients with peripheral PsA.¹⁶⁰ UV light exposure (UVA or UVB) can induce significant flares of the skin disease in patients with lupus, especially in those with SCLE who are positive for anti-Ro/SSA antibodies.^{89, 132} Hence, UV therapy should be avoided in patients with coexistent psoriasis and SLE.

The control of serious SLE manifestations (renal, CNS, hematologic, and so forth) often requires systemic CS therapy, but a rebound flare of psoriasis is possible upon withdrawal of the CS. Antimetabolites such as AZA and MTX used as steroid-sparing agents may prevent this as well as be beneficial for the psoriasis.³¹

KIDNEY DISEASES

Kidney involvement is a common finding in different rheumatic diseases and represents an important determinant of morbidity and mortality.²⁹ SLE, systemic sclerosis, mixed connective tissue disease (MCTD), and vasculitides like Wegener's granulomatosis, microscopic polyarteritis, polyarteritis nodosa, mixed cryoglobulinemia, and Henoch-Schönlein purpura, are the diseases that are most commonly associated with renal injury. Kidney involvement also can be secondary to amyloidosis because of chronic inflammatory conditions like RA and ankylosing spondylitis or more rarely caused by complications from chronic gout.²⁹

In addition to kidney abnormalities caused by the underlying rheumatologic conditions, a number of antirheumatic drugs have been shown to possess nephrotoxic properties. The most commonly used potentially nephrotoxic medications are listed in Table 3.

Renal insufficiency either caused by the underlying rheumatic condition or by unrelated causes (e.g., hypertension, diabetes mellitus, hereditary causes, and so forth) complicates the appropriate treatment of the affected patients. For most antirheumatic drugs, renal insufficiency can lead to inappropriately high levels in the blood and prolonged half-life of the medications, causing increased toxicity (see Table 4). Certain medications, like NSAIDs, MTX, gold (oral or intramuscular), D-penicillamine, and probenecid should be avoided in the setting of advanced

Table 3. RENAL TOXICITY OF ANTIRHEUMATIC MEDICATIONS

| Drug | Renal Toxicity | Risk Factors | Comments |
|--|---|--|---|
| NSAIDs ¹⁸ | Acute renal failure | Plasma volume contraction Congestive heart failure | |
| | Hyperkalemia | Cirrhosis of the liver with ascites Concomitant drug use, e.g., β -blockers, α -agonists, K^+ -sparing diuretics, ACE-inhibitors | |
| | | Insulin-dependent diabetes mellitus Moderate-severe renal failure (GFR <30 mL/min) | Transient effect Usually reversible |
| | Sodium retention Water retention Acute interstitial nephritis \pm minimal- change glomerulopathy/nephrotic syndrome Chronic renal failure Renal papillary necrosis \uparrow protein catabolic rate, azotemia \uparrow GFR (high-dose) | Concomitant use of OTC analgesics | Rare "False" evidence of improvement in renal function |
| Corticosteroids ^{19, 24, 37} | Hypokalemic alkalosis Proteinuria | | Mild/transient = 10%-20% Severe = 1%-3% Occurs in 1%-2% Usually reversible (70%) Rare |
| Gold ²⁰ (mainly with IM route) | Hematuria Nephrotic syndrome caused by membranous glomerulopathy Renal insufficiency | HLA-B8/DR3 | |

| | | |
|--|---|---|
| D-Penicillamine ⁷⁹ | Proteinuria Nephrotic syndrome Membranous glomerulonephritis Microscopic hematuria Rapidly progressive glomerulonephritis Goodpasture-like syndrome | Common (10%–20%) Rare Rare Usually benign Rare Rare Granular deposits in basement membrane Anti-GBM = (–) Dose-related Usually reversible |
| Cyclosporine ^{178, 195} | Acute nephrotoxicity Chronic nephrotoxicity | <p>↑ Creatinine HTN ↑ K⁺ Mild distal RTA Slowly progressive renal failure Proteinuria HTN</p> <p>Increasing age Dose >5 mg/kg/d ↑ of pretreatment creatinine by >50% CsA-induced HTN</p> <p>10% in RA patients treated for ~3 yr</p> |
| FK-506/Tacrolimus ^{145, 167, 202, 208, 215} | Hemolytic-uremic syndrome Acute/chronic toxicity similar to cyclosporine | <p>Renal transplant patients Most of the data derived from renal/ liver transplant recipients Usually reversible after dose reduction ? autoimmune diseases</p> |

GFR = glomerular filtration rate; OTC = over the counter; RTA = renal tubular acidosis; CsA = cyclosporine; GBM = glomerular basement membrane; RA = rheumatoid arthritis; HTN = hypertension; ACE = angiotensin converting enzyme.

Table 4. TOXICITY OF ANTIRHEUMATIC DRUGS IN THE PRESENCE OF RENAL INSUFFICIENCY

| Drug | Major Toxicity | Comments |
|---|---|--|
| Allopurinol ^{88, 198} | Allopurinol hypersensitivity syndrome manifested by: Skin rash (erythematous desquamative) Fever Hepatitis Eosinophilia Worsening renal function | Renal insufficiency present in 81% of reported cases Mortality = 26% Dose adjustment required (see Table 5) for patients with renal failure |
| Colchicine ^{72, 122, 170, 230} | Proximal myopathy with ↑ creatinine kinase Neuropathy (mild, axonal) | Usually reversible after drug discontinuation ↓ dose by 50% for GFR <10 mL/min Special precautions for posttransplant patients receiving CsA |
| Methotrexate ^{85, 95, 176} | Pancytopenia ↑ Overall severe toxicity | 54% of all reported cases of bone marrow suppression had evidence of renal insufficiency Odds ratio for overall severe toxicity ranging from 3 to 5.7 Probably avoid for patients with creatinine >2.0 mg/dL |
| NSAIDs ¹⁸ | Hyperkalemia ↑ Risk of GI bleeding (?) | NSAIDs should be probably avoided for patients with GFR <50 mL/min and used cautiously in hemodialysis patients |
| Cyclophosphamide/ azathioprine | Possible potentiation of bone marrow suppressive effects | Close monitoring of cell counts is mandatory |

GFR = glomerular filtration rate; CsA = cyclosporine; GI = gastrointestinal; NSAIDs = nonsteroidal anti-inflammatory drugs.

renal failure. For other medications, appropriate adjustment of their dose is imperative in the presence of mild or severe renal failure. An overview of the pharmacokinetics of different antirheumatic medications and their dosage adjustment in patients with end-stage renal disease (ESRD) requiring hemodialysis or peritoneal dialysis is presented in Table 5.

Special precautions also should be taken in order to avoid hazardous interactions between different medications, especially in the setting of underlying renal failure. A complete review of the multiple interactions between different antirheumatic medications is beyond the scope of this review. Most antirheumatic drugs should be used very cautiously in patients with renal insufficiency, with close laboratory monitoring and increased alertness for any signs of clinical toxicity.

Table 5. PHARMACOKINETICS AND DOSAGE ADJUSTMENT OF ANTIRHEUMATIC DRUGS

| Drug | Clearance/ Metabolism | Protein Binding (%) | Half-life Normal (t 1/2) | Half-life ESRD (t 1/2) | Adjustment for Renal Failure by GFR (mL/min) | | | Supple- mentation After Hemodialysis | Supple- mentation After Peritoneal Dialysis |
|--|--|---|--|---|---|---|--|---|---|
| | | | | | > 50 (%) | 10-50 (%) | < 10 (%) | | |
| Acetylsalicylic acid NSAIDs | Hepatic metabolism Hepatic metabolism | 80-90 95-99 | 15 min Variable (1-100 h) | Unchanged Unchanged except Diflunisal (5-20 h → 62 h) Ketorolac (4-6 h → 10 h) | 100 100 | ? ↓ by 50 for Diflunisal Ketorolac | Avoid ↓ by 50 for Diflunisal Ketorolac Ketoprofen Oxaprozin Sulindac | Yes None | None None |
| Prednisone Methylprednisolone Cyclophosphamide | Hepatic metabolism Hepatic metabolism Hepatic metabolism | 75 40-60 14 (metabolites ~ 60) | 2.5-3.5 h 1.9-6 h 4-7.5 h | Unchanged Unchanged 10 h | 100 100 100 | 100 100 100 | 100 100 75 | None Yes Half dose | Unknown Unknown Unknown |
| Sulfasalazine Azathioprine | Colon → SP/5-ASA Hepatic metabolism/ renal excretion of metabolites | 99 20 | 4-14 h 15 min (metabolites ~ 5 h) | ↑ ↑ | 100 100 | ? 75 | ? 50 | Unknown Yes | Unknown Unknown |
| Methotrexate Cyclosporine FK-506 Auranofin | Renal excretion Hepatic metabolism Hepatic metabolism Renal excretion | 45-50 96-99 75-99 60 | 7 h 5-18 h 4-40 h Plasma = 17-26 d Whole body = 70-80 d | ↑ Unchanged Unchanged Unknown | 100 100 100 50 | 50 100 100 Avoid | Avoid 50 ? | None None Unknown None | None None Unknown None |
| Gold sodium thiuronate Penicillamine | Renal excretion Hepatic metabolism Renal excretion | 95 80 | Variable (7-250 d) 1.5-3 h | Unknown ↑ | 50 100 | Avoid Avoid | Avoid Avoid | None One-third dose | None Unknown |
| Allopurinol | Oxidation to oxipurinol Renal excretion | <5 | Allopurinol = 2-3 h Oxypurinol = 15 h | Unchanged | 75 | 50 | 25 | Half dose | Unknown |
| Colchicine Probenecid Chloroquine | Renal excretion Renal excretion Renal excretion | 31 85-95 50-65 | 19 h 5-8 h 2-4 d | 40 h Unchanged 5-50 d | 100 100 100 | 100 Avoid 100 | 50 Avoid 50 | None Unknown None | Unknown None None |

SP = sulfapyridine; 5-ASA = 5-aminosalicylic acid; ESRD = end stage renal disease; GFR = glomerular filtration rate.
Data from references 16, 17, 39, 163, 173, 202.

MALIGNANT DISEASES

Treating patients with various rheumatologic conditions in the setting of an underlying malignancy is a challenging task. Rheumatologists are faced with this situation in patients with:

- De novo appearing rheumatologic conditions caused by underlying malignancies or as a complication of cancer treatment (chemotherapy, radiation therapy, etc)
- Exacerbation of pre-existing rheumatic diseases caused by malignancy or cancer therapy
- Development of malignancies in the course of a chronic rheumatic disease

The choice of the appropriate therapeutic regimen should be influenced by several factors. Obviously, the nature and the severity of the coexistent malignant condition, which usually determines the expected survival for the individual patient, is the most important factor. In patients with projected short life expectancy, therapy aiming primarily at relief of symptoms should be offered. Because most of these patients have coexistent conditions such as cytopenias, infections, or evidence of different organ dysfunction (liver, lung, gastrointestinal, etc.) selecting medications with the least possible toxicity is appropriate.

In patients with projected longer life expectancy after treatment of their primary malignancy, several issues must be taken into account before designing the appropriate therapy. Differentiating between a primary rheumatologic condition versus a malignancy or drug-associated musculoskeletal syndrome is an important first step.

Various musculoskeletal syndromes have been described in association with malignant diseases.^{27, 35, 150} An overview of the most commonly described conditions is summarized in Table 6. Treatment of the underlying malignant disease occasionally leads to improvement of the associated rheumatic conditions like carcinoma-associated polyarthritis,²⁸ malignancy-associated polymyositis (PM)/dermatomyositis (DM),³⁵ and so forth. In other conditions, conventional antirheumatic therapy is required for symptom control and improvement of the patient's quality of life. Certain conditions like myelodysplastic syndrome (MDS)-associated polyarthritis, polymyalgia rheumatica, malignancy-associated PM/DM, and various malignancy-associated vasculitides may respond to steroid therapy.³⁵ George and Newman observed a good response to steroids in 5 out of 8 patients with seronegative inflammatory arthritis associated with MDS.⁷¹ Similar responses in various MDS-associated rheumatologic conditions have been reported in a study from the Mayo Clinic.³⁶ When steroids are used for the treatment of these disorders, increased vigilance for early diagnosis and aggressive treatment of emerging infections is required. The use of cytotoxic immunosuppressive agents should be avoided because of concerns about recurrence of the underlying malignant process or transformation of a premalignant condition to frank malignancy (e.g., MDS to leukemia). A similar approach should be taken in patients with pre-existing rheumatic diseases that are exacerbated

Table 6. MUSCULOSKELETAL SYNDROMES ASSOCIATED WITH MALIGNANCIES

| Musculoskeletal Syndrome | Malignancy | Comments |
|--------------------------------------|---|---|
| Inflammatory polyarthritis (RA-like) | Carcinomas* (most common) MDS Lymphoma (especially NHL) Paraproteinemias (MM/ WM/MGUS) Leukemias (rare/children) | Predominantly asymmetric with lower extremity involvement ↑ acute phase reactants/RF(-) Occasionally good response to steroids |
| Mono/oligo-articular involvement | Leukemia Lymphoma Metastatic cancer with local invasion Primary bone tumors | Common knee involvement Nonspecific synovial biopsy changes Pain a prominent feature |
| Hypertrophic osteoarthropathy | Mainly lung neoplasms Intrathoracic carcinomas | Knee/ankle/MTP/wrist involvement/+ clubbing Bone scan (+)/periosteal thickening in radiographs Occasional improvement with steroids |
| Polymyalgia rheumatica | Various malignancies (ovary, prostate, kidney)/ Questionable relationship | Accompanied by polyarthritis |
| Palmar fasciitis | Ovarian tumors | |
| Relapsing polychondritis | MDS (rare) Lymphoma Leukemia | |
| Dermatomyositis/polymyositis | Carcinomas (breast, ovary, lung, GI) | Most common with DM Variable response to steroid treatment |
| Necrotizing myopathy | Carcinomas (lung/breast/GI/bladder) | Shoulder girdle>>lower extremities Steroid-resistant |
| Systemic sclerosis | Carcinoma of the stomach/melanoma | |
| Eosinophilic fasciitis | Hematopoietic malignancies | |
| Panniculitis | Pancreatic cancer | Accompanied by ankle arthritis |
| Vasculitides | Hematopoietic malignancies (hairy cell leukemia/lymphomas/MDS, etc.) Solid tumors (Lung/colon/renal cell, etc.) | Usually leukocytoclastic but any size vessel can be affected |

*Typically seronegative, explosive onset arthritis.

MDS = myelodysplastic syndrome; MM = multiple myeloma; WM = Waldenström macroglobulinemia; MGUS = monoclonal gammopathy of undetermined significance; NHL = non-Hodgkin's lymphoma; MTP = metatarsophalangeal; RF = rheumatoid factor; DM = dermatomyositis.

Data from references 28, 35, 150.

because of the underlying malignancy or as a consequence of the cancer treatment (see below).

Rheumatologists should also be aware of the potential of anticancer drugs to induce various rheumatologic syndromes or to exacerbate underlying rheumatic diseases. In Table 7 the most commonly reported drug-induced musculoskeletal conditions are presented. Identifying and discontinuing the offending medication, after appropriate consultation

Table 7. ANTICANCER DRUG INDUCED RHEUMATOLOGICAL SYNDROMES

| Drug | Musculoskeletal Syndrome | Comments |
|---|---|---|
| Intravesical bacille Calmette-Guérin (BCG) ⁵¹ 80, 104, 169, 189, 213, 228 | Arthralgias/reactive arthritis | Usually symmetric polyarthritis involving small joints HLA-B27 (+) = 44% Good response to NSAIDs |
| Interleukin-2 ^{130, 142} | Symmetric polyarthritis (RA-like) Reactivation of quiescent RA or Reiter's syndrome | Usually few days to weeks after IV infusion Occasional improvement with NSAIDs |
| G-CSF/GM-CSF ^{103, 146, 188, 203} | Cutaneous vasculitis Sweet's syndrome Pyoderma gangrenosum Psoriasis exacerbation Gout/pseudogout exacerbation (G-CSF) | |
| Interferons (- α , - β , - γ) ^{146, 151} | Induction/exacerbation of various autoimmune diseases (SLE/RA/psoriasis/vasculitis) Arthritis (IFN- α) Raynaud's (rare) Leukocytoclastic vasculitis (rare) | Polyarthritis (50%) with RF(+) = 34% Arthritis usually improves with drug discontinuation and/or addition of NSAIDs \pm steroids |
| Tamoxifen ^{44, 217} | Induction/exacerbation of inflammatory polyarthritis | |
| Various chemotherapeutic agents ^{44, 135, 147, 171, 182, 196, 200, 217} | "Post-chemotherapy rheumatism" | See text for details |

NSAIDs = non-steroidal anti-inflammatory drugs; RA = rheumatoid arthritis; G-CSF = granulocyte colony-stimulating factor; GM-CSF = granulocyte-macrophage colony-stimulating factor; RF = rheumatoid factor; SLE = systemic lupus erythematosus; IFN = interferon.

with the patient's hematologist/oncologist, could lead to improvement of patient's symptoms and avoidance of an extensive diagnostic work-up. In some cases, addition of NSAIDs or low-dose CS is indicated.

Loprinzi et al have recently described a syndrome of widespread arthralgias associated with mild inflammation in a few cases, occurring 1 to 3 months after completion of chemotherapy for breast cancer.¹³⁶ They called this syndrome *postchemotherapy rheumatism*. Since then, 24 more cases have been described.^{135, 147, 171, 182, 196, 200, 217} Most patients were women (31 of 32) receiving chemotherapy for breast cancer (27 of 31). Symmetric polyarthralgias/myalgias, morning stiffness, and absence of laboratory findings suggestive of an inflammatory process characterize this syndrome. In a few cases objective findings suggestive of synovitis were present. Treatment with NSAIDs has been ineffective in the major-

ity of patients, whereas CS provided little additional benefit. The symptoms usually subside spontaneously 4 to 12 months later. The pathogenesis of this condition is unknown. It is interesting that all reported patients had received cyclophosphamide as part of their chemotherapy regimen. Still, though, it remains unclear if this condition represents a syndrome with a common origin or a heterogeneous collection of musculoskeletal conditions caused by chemotherapy withdrawal, chemotherapy-induced menopause, steroid-withdrawal, tamoxifen-induced polyarthritides, or fibromyalgia.

Certain rheumatic conditions have been associated with an increased risk of malignancy. These include RA (hematopoietic malignancies), Sjögren's syndrome (lymphomas), Felty's syndrome (non-Hodgkin lymphomas), scleroderma (lung cancer), SLE (non-Hodgkin's lymphomas), discoid lupus erythematosus (squamous cell epithelioma), Paget's disease (osteogenic sarcoma), and lymphomatoid granulomatosis (lymphoma).³⁵ Cytotoxic agents that are commonly used for the treatment of various rheumatic diseases, such as AZA,^{106, 197} cyclophosphamide,^{172, 205} chlorambucil,^{92, 158} and CsA⁸ have been shown to further increase this risk. A highly controversial issue is the possible oncogenic potential (mainly for lymphomas) of MTX when it is used in low doses for the treatment of RA.^{73, 119} Regardless of the exact contribution of cytotoxic agents in the development of certain malignant disorders, recent evidence suggests that occasionally discontinuation of these agents can lead to reversal of the malignant process.^{187, 214} It is prudent to discontinue all cytotoxic agents when a malignancy is identified in patients with a chronic rheumatic disease. Whether or not these agents should be used for life-threatening complications of the underlying rheumatic disease or for symptomatic relief of these patients is a question that has to be addressed on an individual basis, taking into account all the factors that have been mentioned previously.

References

1. Abel EA: Diagnosis of drug-induced psoriasis. *Seminars in Dermatology* 11:269-274, 1992
2. Abel EA, DiCicco LM, Orenberg EK, et al: Drugs in exacerbation of psoriasis. *J Am Acad Dermatol* 15:1007-1022, 1986
3. Al-Kawas FH, Seeff LB, Berendson RA, et al: Allopurinol hepatotoxicity. Report of two cases and review of the literature. *Ann Intern Med* 95:588-590, 1981
4. Allison MC, Howatson AG, Torrance CJ, et al: Gastrointestinal damage associated with the use of nonsteroidal antiinflammatory drugs. *N Engl J Med* 327:749-754, 1992
5. Anderson FH, Zeng L, Yoshida EM, et al: Failure of ketoprofen and interferon combination therapy to improve interferon-resistant chronic hepatitis C. *Can J Gastroenterol* 11:294-297, 1997
6. Andreone P, Cursaro C, Gramenzi A, et al: Indomethacin enhances serum 2'5'-oligoadenylate synthetase in patients with hepatitis B and C virus chronic active hepatitis. *J Hepatol* 21:984-988, 1994
7. Antillon M, Cominelli F, Lo S, et al: Effects of oral prostaglandins on indomethacin-induced renal failure in patients with cirrhosis and ascites. *J Rheumatol Suppl* 20:46-49, 1990

8. Arellano F, Krupp P: Malignancies in rheumatoid arthritis patients treated with cyclosporin A. *Br J Rheumatol* 32(suppl 1):72-75, 1993
9. Arroyo V, Gines P, Rimola A, et al: Renal function abnormalities, prostaglandins, and effects of nonsteroidal anti-inflammatory drugs in cirrhosis with ascites. An overview with emphasis on pathogenesis. *Am J Med* 81:104-122, 1986
10. Assier H, Auffret N, Beltzer-Garely E, et al: Pustulose exanthematique aigue generalisee a l'hydroxychloroquine. *Ann Dermatol Venerol* 120:848-850, 1993
11. Baker H: The influence of chloroquine and related drugs on psoriasis and keratoderma blennorrhagicum. *Br J Dermatol* 78:161-166, 1966
12. Ballare M, Bobbio F, Poggi S, et al: A pilot study on the effectiveness of cyclosporine in type II mixed cryoglobulinemia. *Clin Exp Rheumatol* 13(suppl 13):S201-S203, 1995
13. Bang BK, Yang CW, Yoon SA, et al: Prevalence and clinical course of hepatitis B and hepatitis C liver disease in ciclosporin-treated renal allograft recipients. *Nephron* 70:397-401, 1995
14. Baylis C, Handa RK, Sorkin M: Glucocorticoids and control of glomerular filtration rate. *Semin Nephrol* 10:320-329, 1990
15. Ben-Chetrit E, Rubinow A: Exacerbation of psoriasis by ibuprofen. *Cutis* 38:45, 1986
16. Benet LZ, Oie S, Schwartz JB: Design and optimization of dosage regimens; pharmacokinetic data. In Hardman JG, Gilman AG, Limbird LE (eds): *Goodman & Gilman's The Pharmacological basis of therapeutics*, ed 9. New York, McGraw-Hill, 1996, pp 1702-1792
17. Bennett WM, Avonoff GR, Golper TA, et al: Drug prescribing in renal failure: Dosing guidelines for adults, ed 3. Philadelphia, American College of Physicians, 1994
18. Bennett WM, Henrich WL, Stoff JS: The renal effects of nonsteroidal anti-inflammatory drugs: Summary and recommendations. *Am J Kidney Dis* 28(suppl 1):S56-S62, 1996
19. Berge E, Laurent-Puig P, Clerc D, et al: Rheumatism proviasique au cours d'un traitement par l'interferon alpha [letter]. *Revue du Rhumatisme. Edition Francaise* 60:77, 1993
20. Bielicky T: Light-sensitive psoriasis and antimalarial drugs. *J Invest Dermatol* 41:1-2, 1963
21. Bjarnason I, Hayllar J, MacPherson AJ, et al: Side effects of nonsteroidal anti-inflammatory drugs on the small and large intestine in humans. *Gastroenterology* 104:1832-1847, 1993
22. Bjarnason I, Hayllar J, Smethurst P, et al: Metronidazole reduces intestinal inflammation and blood loss in non-steroidal anti-inflammatory drug induced enteropathy. *Gut* 33:1204-1208, 1992
23. Bjarnason I, Hopkinson N, Zanelli G, et al: Treatment of non-steroidal anti-inflammatory drug induced enteropathy. *Gut* 31:777-780, 1990
24. Boyd AS, Menter A: Erythrodermic psoriasis. Precipitating factors, course, and prognosis in 50 patients. *J Am Acad Dermatol* 21:985-991, 1989
25. Brass EP: Hepatic toxicity of antirheumatic drugs. *Cleve Clin J Med* 60:466-472, 1993
26. Bruckle W, Dixel T, Grasedyck K, et al: Treatment of psoriatic arthritis with auranofin and gold sodium thiomalate. *Clin Rheumatol* 13:209-216, 1994
27. Caldwell DS: Musculoskeletal syndromes associated with malignancy. In Kelley WM, Harris ED Jr, Ruddy S, Sledge CB (eds): *Textbook of Rheumatology*, ed 5. Philadelphia, WB Saunders, 1997, pp 1521-1533
28. Caldwell DS, McCallum RM: Rheumatologic manifestations of cancer. *Med Clin North Am* 70:385-417, 1986
29. Cameron JS: The kidney and rheumatic disease. In Klippel JH, Dieppe PA (eds): *Rheumatology*, ed 2. London, Mosby, 1998, pp 2.9.1-2.9.10
30. Camisa C: Concurrent diseases, associations, and drugs. In Camisa C (ed): *Psoriasis*, ed 1. Boston, Blackwell, 1994, pp 25-39
31. Camisa C: Corticosteroids. In Camisa C (ed): *Psoriasis*, ed 1. Boston, Blackwell, 1994, pp 177-196
32. Camisa C: Overview of psoriasis. In Camisa C (ed): *Handbook of Psoriasis*, Malden, Blackwell, 1998, pp 1-6
33. Campbell K, Steele RJ: Non-steroidal anti-inflammatory drugs and complicated diverticular disease: A case-control study. *Br J Surg* 78:190-191, 1991

34. Carson JL, Strom BL, Duff A, et al: Safety of nonsteroidal anti-inflammatory drugs with respect to acute liver disease. *Arch Intern Med* 153:1331-1336, 1993
35. Carsons S: The association of malignancy with rheumatic and connective tissue diseases. *Semin Oncol* 24:360-372, 1997
36. Castro M, Conn DL, Su WP, et al: Rheumatic manifestations in myelodysplastic syndromes. *J Rheumatol* 18:721-727, 1991
37. Cogan MG, Sargent JA, Yarbrough SG, et al: Prevention of prednisone-induced negative nitrogen balance. Effect of dietary modification on urea generation rate in patients on hemodialysis receiving high-dose glucocorticoids. *Ann Intern Med* 95:158-161, 1981
38. Cohard M, Poynard T, Mathurin P, et al: Prednisone-interferon combination in the treatment of chronic hepatitis B: Direct and indirect metanalysis. *Hepatology* 20:1390-1398, 1994
39. Cohen R, Berns JS, Rudnick M: End-stage renal disease in autoimmune diseases. *In* Mandell BF (ed): *Acute Rheumatic and Immunological Diseases: Management of the critically ill patient*, ed 1. New York, Marcel Dekker, 1994, pp 365-397
40. Corder A: Steroids, non-steroidal anti-inflammatory drugs, and serious septic complications of diverticular disease. *BMJ* 295:1238, 1987
41. Cornbleet T: Preliminary and short report: Action of synthetic antimalarial drugs in psoriasis. *J Invest Dermatol* 26:435-436, 1956
42. Cote PJ, Korba BE, Baldwin B, et al: Immunosuppression with cyclosporine during the incubation period of experimental woodchuck hepatitis virus infection increases the frequency of chronic infection in adult woodchucks. *J Infect Dis* 166:628-631, 1992
43. Coulter DM, Pillans PI: Angiotensin-converting enzyme inhibitors and psoriasis [letter]. *NZ Med J* 106:392-393, 1993
44. Creamer P, Lim K, George E, et al: Acute inflammatory polyarthritis in association with tamoxifen. *Br J Rheumatol* 33:583-585, 1994
45. Cryer B: Nonsteroidal anti-inflammatory drugs and gastrointestinal disease. *In* Feldman M, Scharschmidt BF, Sleisenger MH (eds): *Sleisenger and Fordtran's Gastrointestinal and Liver Disease: Pathophysiology/Diagnosis/Management*, ed 6. Philadelphia, WB Saunders, 1998, pp 343-357
46. Czaja AJ: Diagnosis and therapy of autoimmune liver disease. *Med Clin North Am* 80:973-994, 1996
47. Czaja AJ: The variant forms of autoimmune hepatitis. *Ann Intern Med* 125:588-598, 1996
48. Dammacco F, Sansonno D, Han JH, et al: Natural interferon-alpha versus its combination with 6-methyl-prednisolone in the therapy of type II mixed cryoglobulinemia: A long-term, randomized, controlled study. *Blood* 84:3336-3343, 1994
49. Davies NM: Toxicity of nonsteroidal anti-inflammatory drugs in the large intestine. *Dis Colon Rectum* 38:1311-1321, 1995
50. Demidem A, Taylor JR, Grammer SF, et al: Effects of chloroquine on antigen-presenting functions of epidermal cells from normal and psoriatic skin. *J Invest Dermatol* 98:181-186, 1992
51. Devlin M, Deodhar A, Davis M: Arthritis as a complication of intravesical BCG vaccine [letter]. *BMJ* 308:1638, 1994
52. Dorwart BB, Gall EP, Schumacher HR, et al: Chrysotherapy in psoriatic arthritis. Efficacy and toxicity compared to rheumatoid arthritis. *Arthritis Rheum* 21:513-515, 1978
53. Dubois EL: *Lupus Erythematosus*, ed 2. Los Angeles, University of Southern California Press, 1974
54. Duclos-Vallee JC, Hadengue A, Ganne-Carrie N, et al: Primary biliary cirrhosis-autoimmune hepatitis overlap syndrome. Corticoreistance and effective treatment by cyclosporine A. *Dig Dis Sci* 40:1069-1073, 1995
55. Ellis CN, Goldfarb MT, Roenigk HHJ, et al: Effects of oral meclofenamate therapy in psoriasis. *J Am Acad Dermatol* 14:49-52, 1986
56. Evans JM, McMahon AD, Murray FE, et al: Non-steroidal anti-inflammatory drugs are associated with emergency admission to hospital for colitis due to inflammatory bowel disease. *Gut* 40:619-622, 1997

57. Farrell GC: Liver disease caused by drugs, anesthetics and toxins. In Feldman M, Scharschmidt BF, Sleisenger MH (eds): *Sleisenger and Fordtran's Gastrointestinal and Liver Disease: Pathophysiology/Diagnosis/Management*, ed 6. Philadelphia, WB Saunders Company, 1998, pp 1221–1253
58. Fierlbeck G, Rassner G, Muller C: Psoriasis induced at the injection site of recombinant interferon gamma. Results of immunohistologic investigations. *Arch Dermatol* 126:351–355, 1990
59. Flowers MA, Heathcote J, Wanless IR, et al: Fulminant hepatitis as a consequence of reactivation of hepatitis B virus infection after discontinuation of low-dose methotrexate therapy. *Ann Intern Med* 112:381–382, 1990
60. Fong TL, Valinluck B, Govindarajan S, et al: Short-term prednisone therapy affects aminotransferase activity and hepatitis C virus RNA levels in chronic hepatitis C. *Gastroenterology* 107:196–199, 1994
61. Friedman SJ: Pustular psoriasis associated with hydroxychloroquine [letter]. *J Am Acad Dermatol* 16:1256–1257, 1987
62. Friis H, Andreasen PB: Drug-induced hepatic injury: An analysis of 1100 cases reported to the Danish Committee on Adverse Drug Reactions between 1978 and 1987. *J Intern Med* 232:133–138, 1992
63. Fry SW, Seeff LB: Hepatotoxicity of analgesics and anti-inflammatory agents. *Gastroenterol Clin North Am* 24:875–905, 1995
64. Fukuda R, Okinaga S, Akagi S, et al: Alteration of infection pattern of duck hepatitis B virus by immunomodulatory drugs. *J Med Virol* 26:387–396, 1988
65. Fullerton T, Sica DA, Blum RA: Evaluation of the renal protective effect of misoprostol in elderly, osteoarthritic patients at risk for nonsteroidal anti-inflammatory drug-induced renal dysfunction. *J Clin Pharmacol* 33:1225–1232, 1993
66. Funk J, Langeland T, Schrupf E, et al: Psoriasis induced by interferon-alpha. *Br J Dermatol* 125:463–465, 1991
67. Gabriel SE, Jaakkimainen L, Bombardier C: Risk for serious gastrointestinal complications related to use of nonsteroidal anti-inflammatory drugs. A meta-analysis. *Ann Intern Med* 115:787–796, 1991
68. Garcia-Lora E, Tercedor J, Massare E, et al: Interferon-induced psoriasis in a patient with chronic hepatitis C [letter]. *Dermatology* 187:280, 1993
69. Garcia RL, Williams R, Derby LE, et al: Acute liver injury associated with nonsteroidal anti-inflammatory drugs and the role of risk factors. *Arch Intern Med* 154:311–316, 1994
70. George DL, Wiener SG: Skin and rheumatic disease. In Klippel JH, Dieppe PA (eds): *Rheumatology*, ed 2. London, Mosby, 1998, pp 2.5.1–2.5.12
71. George SW, Newman ED: Seronegative inflammatory arthritis in the myelodysplastic syndromes. *Semin Arthritis Rheum* 21:345–354, 1992
72. George T, Mandell BF: Gout in the transplant patient. *Journal of Clinical Rheumatology* 1:328–334, 1995
73. Georgescu L, Quinn GC, Schwartzman S, et al: Lymphoma in patients with rheumatoid arthritis: Association with the disease state or methotrexate treatment. *Semin Arthritis Rheum* 26:794–804, 1997
74. Georgetson MJ, Yarze JC, Losos AT, et al: Exacerbation of psoriasis due to interferon-alpha treatment of chronic active hepatitis. *Am J Gastroenterol* 88:1756–1758, 1993
75. Gibson GR, Whitacre EB, Ricotti CA: Colitis induced by nonsteroidal anti-inflammatory drugs. Report of four cases and review of the literature. *Arch Intern Med* 152:625–632, 1992
76. Gilleaudeau P, Vallat VP, Carter DM, et al: Angiotensin-converting enzyme inhibitors as possible exacerbating drugs in psoriasis. *J Am Acad Dermatol* 28:490–492, 1993
77. Gladman DD, Blake R, Brubacher B, et al: Chloroquine therapy in psoriatic arthritis. *J Rheumatol* 19:1724–1726, 1992
78. Gladman DD, Shuckett R, Russell ML, et al: Psoriatic arthritis (PSA)—an analysis of 220 patients. *QJM* 62:127–141, 1987
79. Gordon DA: Gold compounds and penicillamine in the rheumatic diseases. In Kelley WM, Harris ED Jr, Ruddy S, et al (eds): *Textbook of Rheumatology*, ed 5. Philadelphia, WB Saunders, 1997, pp 759–769

80. Goupille P, Poet JL, Jattiot F, et al: Three cases of arthritis after BCG therapy for bladder cancer. *Clin Exp Rheumatol* 12:195-197, 1994
81. Goupille P, Soutif D, Valat JP: Treatment of psoriatic arthropathy. *Semin Arthritis Rheum* 21:355-367, 1992
82. Gray RG: Hydroxychloroquine provocation of psoriasis [letter]. *J Rheumatol* 12:391, 1985
83. Griffiths CE: Therapy for psoriatic arthritis: Sometimes a conflict for psoriasis [editorial]. *Br J Rheumatol* 36:409-410, 1997
84. Gupta AK, Sibbald RG, Knowles SR, et al: Terbinafine therapy may be associated with the development of psoriasis de novo or its exacerbation: Four case reports and a review of drug-induced psoriasis. *J Am Acad Dermatol* 36:858-862, 1997
85. Gutierrez-Urena S, Molina JE, Garcia CO, et al: Pancytopenia secondary to methotrexate therapy in rheumatoid arthritis. *Arthritis Rheum* 39:272-276, 1996
86. Hadziyannis SJ: The spectrum of extrahepatic manifestations in hepatitis C virus infection. *J Viral Hepat* 4:9-28, 1997
87. Hamlet NW, Keefe M, Kerr RE: Does captopril exacerbate psoriasis? [letter]. *BMJ* 295:1352, 1987
88. Hande KR, Noone RM, Stone WJ: Severe allopurinol toxicity. Description and guidelines for prevention in patients with renal insufficiency. *Am J Med* 76:47-56, 1984
89. Hasan T, Nyberg F, Stephansson E, et al: Photosensitivity in lupus erythematosus, UV photoprovision results compared with history of photosensitivity and clinical findings. *Br J Dermatol* 136:699-705, 1997
90. Hayllar J, Smith T, Macpherson A, et al: Nonsteroidal antiinflammatory drug-induced small intestinal inflammation and blood loss. Effects of sulfasalazine and other disease-modifying antirheumatic drugs. *Arthritis Rheum* 37:1146-1150, 1994
91. Hays SB, Camisa C, Luzar MJ: The coexistence of systemic lupus erythematosus and psoriasis. *J Am Acad Dermatol* 10:619-622, 1984
92. Hazleman B: Incidence of neoplasms in patients with rheumatoid arthritis exposed to different treatment regimens. *Am J Med* 78:39-43, 1985
93. Heintges T, Wands JR: Hepatitis C virus: Epidemiology and transmission. *Hepatology* 26:521-526, 1997
94. Hestin D, Hussenet F, Frimat L, et al: Hepatitis C after renal transplantation: Histopathological correlations. *Nephrol Dial Transplant* 11 (suppl 4):52-53, 1996
95. Hillson JL, Furst DE: Pharmacology and pharmacokinetics of methotrexate in rheumatic disease. Practical issues in treatment and design. *Rheum Dis Clin North Am* 23:757-778, 1997
96. Holt S, Rigoglioso V, Sidhu M, et al: Nonsteroidal antiinflammatory drugs and lower gastrointestinal bleeding. *Dig Dis Sci* 38:1619-1623, 1993
97. Hoofnagle JH, Davis GL, Pappas SC, et al: A short course of prednisolone in chronic type B hepatitis. Report of a randomized, double-blind, placebo-controlled trial. *Ann Intern Med* 104:12-17, 1986
98. Horina JH, Wirnsberger GH, Kenner L, et al: Increased susceptibility for CsA-induced hepatotoxicity in kidney graft recipients with chronic viral hepatitis C. *Transplantation* 56:1091-1094, 1993
99. Howrie DL, Gartner JCJ: Gold-induced hepatotoxicity: Case report and review of the literature. *J Rheumatol* 9:727-729, 1982
100. Hsia HC, Lin HC, Tsai YT, et al: The effects of chronic administration of indomethacin and misoprostol on renal function in cirrhotic patients with and without ascites. *Scand J Gastroenterol* 30:1194-1199, 1995
101. Huang CC, Lai MK, Fong MT: Hepatitis B liver disease in cyclosporine-treated renal allograft recipients. *Transplantation* 49:540-544, 1990
102. Ikai K: Exacerbation and induction of psoriasis by angiotensin-converting enzyme inhibitors [letter]. *J Am Acad Dermatol* 32:819, 1995
103. Jain KK: Sweet's syndrome associated with granulocyte colony-stimulating factor. *Cutis* 57:107-110, 1996
104. Jawad AS, Kahn L, Copland RF, et al: Reactive arthritis associated with Bacillus Calmette-Guerin immunotherapy for carcinoma of the bladder: A report of two cases. *Br J Rheumatol* 32:1018-1020, 1993

105. Jones G, Crotty M, Brooks P: Psoriatic arthritis: A quantitative overview of therapeutic options. The Psoriatic Arthritis Meta-Analysis Study Group. *Br J Rheumatol* 36:95-99, 1997
106. Jones M, Symmons D, Finn J, et al: Does exposure to immunosuppressive therapy increase the 10 year malignancy and mortality risks in rheumatoid arthritis? A matched cohort study. *Br J Rheumatol* 35:738-745, 1996
107. Jucgla A, Marcoval J, Curco N, et al: Psoriasis with articular involvement induced by interferon alfa [letter]. *Arch Dermatol* 127:910-911, 1991
108. Kakumu S, Takayanagi M, Iwata K, et al: Cyclosporine therapy affects aminotransferase activity but not hepatitis C virus RNA levels in chronic hepatitis C. *J Gastroenterol Hepatol* 12:62-66, 1997
109. Kammer GM, Soter NA, Gibson DJ, et al: Psoriatic arthritis: A clinical, immunologic and HLA study of 100 patients. *Semin Arthritis Rheum* 9:75-97, 1979
110. Kaplan MM: The use of methotrexate, colchicine, and other immunomodulatory drugs in the treatment of primary biliary cirrhosis. *Semin Liver Dis* 17:129-136, 1997
111. Katayama H, Kawada A: Exacerbation of psoriasis induced by indomethacin. *J Dermatol* 8:323-327, 1981
112. Kaufmann HJ, Taubin HL: Nonsteroidal anti-inflammatory drugs activate quiescent inflammatory bowel disease. *Ann Intern Med* 107:513-516, 1987
113. Kitamura K, Kanasashi M, Suga C, et al: Cutaneous reactions induced by calcium channel blocker: High frequency of psoriasiform eruptions. *J Dermatol* 20:279-286, 1993
114. Kliem V, van den Hoff U, Brunkhorst R, et al: The long-term course of hepatitis C after kidney transplantation. *Transplantation* 62:1417-1421, 1996
115. Knox TA, Kaplan MM, Gelfand JA, et al: Methotrexate treatment of idiopathic granulomatous hepatitis. *Ann Intern Med* 122:592-595, 1995
116. Kowalick L: Psoriasis flare caused by recombinant interferon beta injections [letter]. *J Am Acad Dermatol* 36:501, 1997
117. Kowdley KV, Keefe EB: Hepatotoxicity of transplant immunosuppressive agents. *Gastroenterol Clin North Am* 24:991-1001, 1995
118. Kragballe K, Herlin T: Benoxaprofen improves psoriasis. A double-blind study. *Arch Dermatol* 119:548-552, 1983
119. Kremer JM: Is methotrexate oncogenic in patients with rheumatoid arthritis? *Semin Arthritis Rheum* 26:785-787, 1997
120. Kremer JM, Alarcon GS, Lightfoot RWJ, et al: Methotrexate for rheumatoid arthritis. Suggested guidelines for monitoring liver toxicity. American College of Rheumatology. *Arthritis Rheum* 37:316-328, 1994
121. Kuflik EG: Effect of antimalarial drugs on psoriasis. *Cutis* 26:153-155, 1980
122. Kunkl RW, Duncan G, Watson D, et al: Colchicine myopathy and neuropathy. *N Engl J Med* 316:1562-1568, 1987
123. Kushimoto K, Nagasawa K, Ueda A, et al: Liver abnormalities and liver membrane autoantibodies in systemic lupus erythematosus. *Ann Rheum Dis* 48:946-952, 1989
124. Kwo PY, Tremaine WJ: Nonsteroidal anti-inflammatory drug-induced enteropathy: Case discussion and review of the literature. *Mayo Clin Proc* 70:55-61, 1995
125. Laffi G, La Villa G, Pinzani M, et al: Arachidonic acid derivatives and renal function in liver cirrhosis. *Semin Nephrol* 17:530-548, 1997
126. Lam KC, Lai CL, Trepo C, et al: Deleterious effect of prednisolone in HBsAg-positive chronic active hepatitis. *N Engl J Med* 304:380-386, 1981
127. Langford CA, Klippel JH, Balow JE, et al: Use of cytotoxic agents and cyclosporine in the treatment of autoimmune disease. Part 1: Rheumatologic and renal diseases. *Ann Intern Med* 128:1021-1028, 1998
128. Langman MJ, Morgan L, Worrall A: Use of anti-inflammatory drugs by patients admitted with small or large bowel perforations and haemorrhage. *BMJ* 290:347-349, 1985
129. Lau JY, Bain VG, Smith HM, et al: Modulation of hepatitis B viral antigen expression by immunosuppressive drugs in primary hepatocyte culture. *Transplantation* 53:894-898, 1992
130. Lavelle-Jones M, al-Hadrani A, Spiers EM, et al: Reactivation of rheumatoid arthritis

- during continuous infusion of interleukin 2: Evidence of lymphocytic control of rheumatoid disease. *BMJ* 301:97, 1990
131. Lawrence SP, Sherman KE, Lawson JM, et al: A 39 year old man with chronic hepatitis. *Semin Liver Dis* 14:97-105, 1994
 132. Lehmann P, Holzle E, Kind P, et al: Experimental reproduction of skin lesions in lupus erythematosus by UVA and UVB radiation. *J Am Acad Dermatol* 22:181-187, 1990
 133. Lombard M, Portmann B, Neuberger J, et al: Cyclosporin A treatment in primary biliary cirrhosis: Results of a long-term placebo controlled trial. *Gastroenterology* 104:519-526, 1993
 134. Lombardini F, Taglione E, Riente L, et al: Psoriatic arthritis with spinal involvement in a patient receiving alpha-interferon for chronic hepatitis C. *Scand J Rheumatol* 26:58-60, 1997
 135. Loprinzi CL, Duffy J, Ingle JN: More postchemotherapy rheumatism [letter]. *J Clin Oncol* 11:2052, 1993
 136. Loprinzi CL, Duffy J, Ingle JN: Postchemotherapy rheumatism. *J Clin Oncol* 11:768-770, 1993
 137. Luzar MJ: Hydroxychloroquine in psoriatic arthropathy: Exacerbations of psoriatic skin lesions. *J Rheumatol* 9:462-464, 1982
 138. Magrin S, Craxi A, Fabiano C, et al: Hepatitis C viremia in chronic liver disease: Relationship to interferon-alpha or corticosteroid treatment. *Hepatology* 19:273-279, 1994
 139. Mahmud T, Rafi SS, Scott DL, et al: Nonsteroidal antiinflammatory drugs and uncoupling of mitochondrial oxidative phosphorylation. *Arthritis Rheum* 39:1998-2003, 1996
 140. Makino Y, Tanaka H, Nakamura K, et al: Arthritis in a patient with psoriasis after interferon-alpha therapy for chronic hepatitis C. *J Rheumatol* 21:1771-1772, 1994
 141. Mandell BF, Calabrese LH: Infections and systemic vasculitis. *Curr Opin Rheumatol* 10:51-57, 1998
 142. Massarotti EM, Liu NY, Mier J, et al: Chronic inflammatory arthritis after treatment with high-dose interleukin-2 for malignancy. *Am J Med* 92:693-697, 1992
 143. Matsumoto T, Yoshimine T, Shimouchi K, et al: The liver in systemic lupus erythematosus: Pathologic analysis of 52 cases and review of Japanese Autopsy Registry Data. *Hum Pathol* 23:1151-1158, 1992
 144. Matzke GR: Nonrenal toxicities of acetaminophen, aspirin, and nonsteroidal anti-inflammatory agents. *Am J Kidney Dis* 28(suppl 1):S63-S70, 1996
 145. McDiarmid SV, Colonna JO, Shaked A, et al: A comparison of renal function in cyclosporine- and FK-506-treated patients after primary orthotopic liver transplantation. *Transplantation* 56:847-853, 1993
 146. Merkel PA: Drugs associated with vasculitis. *Curr Opin Rheumatol* 10:45-50, 1998
 147. Michl I, Zielinski CC: More postchemotherapy rheumatism [letter]. *J Clin Oncol* 11:2051-2052, 1993
 148. Miller MH, Urowitz MB, Gladman DD, et al: The liver in systemic lupus erythematosus. *QJM* 53:401-409, 1984
 149. Myllykangas-Luosujarvi R, Aho K, Isomaki H: Death attributed to antirheumatic medication in a nationwide series of 1666 patients with rheumatoid arthritis who have died. *J Rheumatol* 22:2214-2217, 1995
 150. Naschitz JE, Rosner I, Rozenbaum M, et al: Cancer-associated rheumatic disorders: Clues to occult neoplasia. *Semin Arthritis Rheum* 24:231-241, 1995
 151. Neshar G, Ruchlemer R: Alpha-interferon-induced arthritis: Clinical presentation, treatment, and prevention. *Semin Arthritis Rheum* 27:360-365, 1998
 152. Neshar G, Sonnenblick M, Dwolatzky T: Protective effect of misoprostol on indomethacin-induced renal dysfunction in elderly patients. *J Rheumatol* 22:713-716, 1995
 153. O'Quinn SE, Kennedy CB, Naylor LZ: Psoriasis, ultraviolet light and chloroquine. *Arch Dermatol* 90:211-216, 1964
 154. Ockenfels HM, Nussbaum G, Schultewolter T, et al: Tyrosine phosphorylation in psoriatic T cells is modulated by drugs that induce or improve psoriasis. *Dermatology* 191:217-225, 1995
 155. Olsen TG: Chloroquine and psoriasis [letter]. *Ann Intern Med* 94:546-547, 1981
 156. Palit J, Hill J, Capell HA, et al: A multicentre double-blind comparison of auranofin,

- intramuscular gold thiomalate and placebo in patients with psoriatic arthritis. *Br J Rheumatol* 29:280-283, 1990
157. Panayi GS, Tugwell P: The use of cyclosporin A microemulsion in rheumatoid arthritis: Conclusions of an international review. *Br J Rheumatol* 36:808-811, 1997
 158. Patapanian H, Graham S, Sambrook PN, et al: The oncogenicity of chlorambucil in rheumatoid arthritis. *Br J Rheumatol* 27:44-47, 1988
 159. Periera BJ, Wright TL, Schmid CH, et al: The impact of pretransplantation hepatitis C infection on the outcome of renal transplantation. *Transplantation* 60:799-805, 1995
 160. Perlman SG, Gerber LH, Roberts RM, et al: Photochemotherapy and psoriatic arthritis. A prospective study. *Ann Intern Med* 91:717-722, 1979
 161. Perrillo RP, Schiff ER, Davis GL, et al: A randomized, controlled trial of interferon alfa-2b alone and after prednisone withdrawal for the treatment of chronic hepatitis B. The Hepatitis Interventional Therapy Group. *N Engl J Med* 323:295-301, 1990
 162. Person JL, McHutchison JG, Fong TL, et al: A case of cyclosporine-sensitive, steroid-resistant, autoimmune chronic active hepatitis. *J Clin Gastroenterol* 17:317-320, 1993
 163. Physicians Desk Reference, ed 52. Montvale, Medical Economics Company, 1998
 164. Piper JM, Ray WA, Daugherty JR, et al: Corticosteroid use and peptic ulcer disease: Role of nonsteroidal anti-inflammatory drugs. *Ann Intern Med* 114:735-740, 1991
 165. Poikolainen K, Reunala T, Karvonen J: Smoking, alcohol and life events related to psoriasis among women. *Br J Dermatol* 130:473-477, 1994
 166. Pol S, Cavalcanti R, Carnot F, et al: Azathioprine hepatitis in kidney transplant recipients. A predisposing role of chronic viral hepatitis. *Transplantation* 61:1774-1776, 1996
 167. Porayko MK, Textor SC, Krom RA, et al: Nephrotoxic effects of primary immunosuppression with FK-506 and cyclosporine regimens after liver transplantation. *Mayo Clin Proc* 69:105-111, 1994
 168. Powles AV, Griffiths CE, Seifert MH, et al: Exacerbation of psoriasis by indomethacin [letter]. *Br J Dermatol* 117:799-800, 1987
 169. Price GE: Arthritis and iritis after BCG therapy for bladder cancer. *J Rheumatol* 21:564-565, 1994
 170. Puttermann C, Ben-Chetrit E, Caraco Y, et al: Colchicine intoxication: clinical pharmacology, risk factors, features, and management. *Semin Arthritis Rheum* 21:143-155, 1991
 171. Raderer M, Scheithauer W: Postchemotherapy rheumatism following adjuvant therapy for ovarian cancer. *Scand J Rheumatol* 23:291-292, 1994
 172. Radis CD, Kahl LE, Baker GL, et al: Effects of cyclophosphamide on the development of malignancy and on long-term survival of patients with rheumatoid arthritis. A 20-year follow-up study. *Arthritis Rheum* 38:1120-1127, 1995
 173. Rains CP, Noble S, Faulds D: Sulfasalazine. A review of its pharmacological properties and therapeutic efficacy in the treatment of rheumatoid arthritis [published erratum appears in *Drugs* 50, 625, 1995]. *Drugs* 50:137-156, 1995
 174. Ramond MJ, Poynard T, Rueff B, et al: A randomized trial of prednisolone in patients with severe alcoholic hepatitis. *N Engl J Med* 326:507-512, 1992
 175. Reshad H, Hargreaves GK, Vickers CF: Generalized pustular psoriasis precipitated by phenylbutazone and oxyphenbutazone. *Br J Dermatol* 109:111-113, 1983
 176. Rheumatoid Arthritis Clinical Trial Archive Group: The effect of age and renal function on the efficacy and toxicity of methotrexate in rheumatoid arthritis. *J Rheumatol* 22:218-223, 1995
 177. Richter MB, Kinsella P, Corbett M: Gold in psoriatic arthropathy. *Ann Rheum Dis* 39:279-280, 1980
 178. Rodriguez F, Krayenbuhl JC, Harrison WB, et al: Renal biopsy findings and follow-up of renal function in rheumatoid arthritis patients treated with cyclosporin A. An update from the International Kidney Biopsy Registry. *Arthritis Rheum* 39:1491-1498, 1996
 179. Roenigk HHJ, Auerbach R, Maibach H, et al: Methotrexate in psoriasis: Consensus conference. *J Am Acad Dermatol* 38:478-485, 1998
 180. Roenigk HHJ, Auerbach R, Maibach HI, et al: Methotrexate guidelines—revised. *J Am Acad Dermatol* 6:145-155, 1982

181. Rosenbaum J, Katz WA, Schumacher HR: Hepatotoxicity associated with use of D-penicillamine in rheumatoid arthritis. *Ann Rheum Dis* 39:152-154, 1980
182. Rosenstein ED, Kramer N, Leitner SP, et al: Exacerbation of rheumatoid arthritis after termination of chemotherapy for breast carcinoma. *J Rheumatol* 23:1988-1990, 1996
183. Roth D, Zucker K, Cirocco R, et al: A prospective study of hepatitis C virus infection in renal allograft recipients. *Transplantation* 61:886-889, 1996
184. Runyon BA, LaBrecque DR, Anuras S: The spectrum of liver disease in systemic lupus erythematosus. Report of 33 histologically-proved cases and review of the literature. *Am J Med* 69:187-194, 1980
185. Ryan TJ, Baker H: Systemic corticosteroids and folic acid antagonists in the treatment of generalized pustular psoriasis. Evaluation and prognosis based on the study of 104 cases. *Br J Dermatol* 81:134-145, 1969
186. Ryan TJ, Baker H: The prognosis of generalized pustular psoriasis. *Br J Dermatol* 85:407-411, 1971
187. Salloum E, Cooper DL, Howe G, et al: Spontaneous regression of lymphoproliferative disorders in patients treated with methotrexate for rheumatoid arthritis and other rheumatic diseases. *J Clin Oncol* 14:1943-1949, 1996
188. Sandor V, Hassan R, Kohn E: Exacerbation of pseudogout by granulocyte colony-stimulating factor [letter]. *Ann Intern Med* 125:781, 1996
189. Saporta L, Gumus E, Karadag H, et al: Reiter's syndrome following intracavitary BCG administration. *Scand J Urol Nephrol* 31:211-212, 1997
190. Sayers ME, Mazanec DJ: Use of antimalarial drugs for the treatment of psoriatic arthritis. *Am J Med* 93:474-475, 1992
191. Scheiman JM: NSAIDs, gastrointestinal injury, and cytoprotection. *Gastroenterol Clin North Am* 25:279-298, 1996
192. Schopf RE, Ockenfels HM, Schultewolter T, et al: Chloroquine stimulates the mitogen-driven lymphocyte proliferation in patients with psoriasis. *Dermatology* 187:100-103, 1993
193. Scullard GH, Smith CI, Merigan TC, et al: Effects of immunosuppressive therapy on viral markers in chronic active hepatitis B. *Gastroenterology* 81:987-991, 1981
194. Sendagorta E, Allegue F, Rocamora A, et al: Generalized pustular psoriasis precipitated by diclofenac and indomethacin [letter]. *Dermatologica* 175:300-301, 1987
195. Shihab FS: Cyclosporine nephropathy: Pathophysiology and clinical impact. *Semin Nephrol* 16:536-547, 1996
196. Siegel JE: Postchemotherapy rheumatism: Is this a menopausal symptom? [letter]. *J Clin Oncol* 11:2051, 1993
197. Silman AJ, Petrie J, Hazleman B, et al: Lymphoproliferative cancer and other malignancy in patients with rheumatoid arthritis treated with azathioprine: A 20 year follow up study. *Ann Rheum Dis* 47:988-992, 1988
198. Singer JZ, Wallace SL: The allopurinol hypersensitivity syndrome. Unnecessary morbidity and mortality. *Arthritis Rheum* 29:82-87, 1986
199. Slagel GA, James WD: Plaquenil-induced erythroderma. *J Am Acad Dermatol* 12:857-862, 1985
200. Smith DE: Additional cases of postchemotherapy rheumatism [letter]. *J Clin Oncol* 11:1625-1626, 1993
201. Somasundaram S, Rafi S, Hayllar J, et al: Mitochondrial damage: A possible mechanism of the "topical" phase of NSAID induced injury to the rat intestine. *Gut* 41:344-353, 1997
202. Spencer CM, Goa KL, Gillis JC: Tacrolimus. An update of its pharmacology and clinical efficacy in the management of organ transplantation. *Drugs* 54:925-975, 1997
203. Spitzer T, McAfee S, Poliquin C, et al: Acute gouty arthritis following recombinant human granulocyte colony-stimulating factor therapy in an allogeneic blood stem cell donor [letter]. *Bone Marrow Transplant* 21:966-967, 1998
204. Stempel C, Lake J, Ferrell L, et al: Effect of cyclosporine on the clinical course of HBsAg-positive renal transplant patients. *Transplant Proc* 23:1251-1252, 1991
205. Talar-Williams C, Hijazi YM, Walther MM, et al: Cyclophosphamide-induced cystitis and bladder cancer in patients with Wegener granulomatosis. *Ann Intern Med* 124:477-484, 1996

206. Tamir A, Wolf R, Brenner S: Exacerbation and induction of psoriasis by angiotensin-converting enzyme inhibitors [letter]. *J Am Acad Dermatol* 30:1045, 1994
207. Tarazi EM, Harter JG, Zimmerman HJ, et al: Sulindac-associated hepatic injury: Analysis of 91 cases reported to the Food and Drug Administration. *Gastroenterology* 104:569-574, 1993
208. The U.S. Multicenter FK 506 Liver Study Group: A comparison of tacrolimus (FK 506) and cyclosporine for immunosuppression in liver transplantation. *N Engl J Med* 331:1110-1115, 1994
209. Thiele DL, DuCharme L, Cunningham MR, et al: Steroid therapy of chronic hepatitis: Characteristics associated with response in anti-hepatitis C virus-positive and -negative patients. *Am J Gastroenterol* 91:300-308, 1996
210. Tolman KG: Hepatotoxicity of antirheumatic drugs. *J Rheumatol Suppl* 22:6-11, 1990
211. Torre AJ, Rodriguez PA, Arribas CJ, et al: Psoriatic arthritis (PA): A clinical, immunological and radiological study of 180 patients. *Br J Rheumatol* 30:245-250, 1991
212. Tur-Kaspa R, Laub O: Corticosteroids stimulate hepatitis B virus DNA, mRNA and protein production in a stable expression system. *J Hepatol* 11:34-36, 1990
213. Van Thiel RJ, Nossent GD, Tjon PGN: Articular complications of intravesical BCG treatment for bladder carcinoma. *Br J Urol* 70:446-447, 1992
214. Vassilopoulos D, Waltuck J, Clarke G, et al: Reversible Epstein-Barr virus associated T cell non-Hodgkin's lymphoma in a patient with rheumatoid arthritis undergoing chronic immunosuppressive therapy [letter]. *J Rheumatol* 25:389-391, 1998
215. Vincenti F, Laskow DA, Neylan JE, et al: One-year follow-up of an open-label trial of FK 506 for primary kidney transplantation. A report of the U.S. Multicenter FK 506 Kidney Transplant Group. *Transplantation* 61:1576-1581, 1996
216. Vine JE, Hymes SR, Warner NB, et al: Pustular psoriasis induced by hydroxychloroquine: A case report and review of the literature. *J Dermatol* 23:357-361, 1996
217. Warner E, Keshavjee A, Shupak R, et al: Rheumatic symptoms following adjuvant therapy for breast cancer. *Am J Clin Oncol* 20:322-326, 1997
218. Webster GF, Knobler RL, Lublin FD, et al: Cutaneous ulcerations and pustular psoriasis flare caused by recombinant interferon beta injections in patients with multiple sclerosis [published erratum appears in *J Am Acad Dermatol* 35:587, 1996]. *J Am Acad Dermatol* 34:365-367, 1996
219. Weinblatt ME, Tesser JR, Gilliam JH: The liver in rheumatic diseases. *Semin Arthritis Rheum* 11:399-405, 1982
220. Weller IV, Bassendine MF, Murray AK, et al: Effects of prednisolone/azathioprine in chronic hepatitis B viral infection. *Gut* 23:650-655, 1982
221. West SG: Methotrexate hepatotoxicity. *Rheum Dis Clin North Am* 23:883-915, 1997
222. Wilke WS: Methotrexate use in miscellaneous inflammatory diseases. *Rheum Dis Clin North Am* 23:855-882, 1997
223. Wilke WS, Sayers ME: Psoriatic arthritis. In Camisa C (ed): *Psoriasis*, ed 1. Boston, Blackwell, 1994, pp 109-163
224. Wilson RG, Smith AN, Macintyre IM: Complications of diverticular disease and non-steroidal anti-inflammatory drugs: A prospective study. *Br J Surg* 77:1103-1104, 1990
225. Wolf R, Lo SA, Lombardi ML, et al: The in vitro effect of hydroxychloroquine on skin morphology and transglutaminase. *Int J Dermatol* 36:704-707, 1997
226. Wolf R, Tamir A, Brenner S: Psoriasis related to angiotensin-converting enzyme inhibitors. *Dermatologica* 181:51-53, 1990
227. Wolfe JT, Singh A, Lessin SR, et al: De novo development of psoriatic plaques in patients receiving interferon alfa for treatment of erythrodermic cutaneous T-cell lymphoma. *J Am Acad Dermatol* 32:887-893, 1995
228. Xerri B, Chretien Y, Le Parc JM: Reactive polyarthritis induced by intravesical BCG therapy for carcinoma of the bladder [letter]. *Eur J Med* 2:503-505, 1993
229. Yussim A, Bar-Nathan N, Lustig S, et al: Gastrointestinal, hepatorenal, and neuromuscular toxicity caused by cyclosporine-colchicine interaction in renal transplantation. *Transplant Proc* 26:2825-2826, 1994
230. Zarski JP, Maynard-Muet M, Chousterman S, et al: Tenoxicam, a non-steroid anti-inflammatory drug, is unable to increase the response rate in patients with chronic hepatitis C treated by alpha interferon. *Hepatology* 27:862-867, 1998

- 231. Ziprowski L, Haim S, Bank H: Atabrine in psoriasis. *Acta Med Orient* 13:45-52, 1954
- 232. Zoutman DE, Ralph ED, Frei JV: Granulomatous hepatitis and fever of unknown origin. An 11-year experience of 23 cases with three years' follow-up. *J Clin Gastroenterol* 13:69-75, 1991

Address reprint requests to

Dimitrios Vassilopoulos, MD
Department of Rheumatic and Immunologic Disease
Cleveland Clinic Foundation
9500 Euclid Avenue, A50
Cleveland, OH 44915