

# PBM-MAP Clinical Practice Guideline for the Pharmacologic Management of **Chronic Heart Failure** in Primary Care Practice

*Pharmacy Benefits Management  
Strategic Healthcare Group and  
Medical Advisory Panel*

Veterans Health Administration  
Department of Veterans Affairs  
Publication No. 00-0015  
September 2007

Version: Final

Updates may be found at [www.pbm.va.gov](http://www.pbm.va.gov) or <http://vaww.pbm.va.gov>

## EXECUTIVE SUMMARY

1. Treatment of chronic heart failure (HF) is based upon the four-stage classification system developed by the American College of Cardiology/American Heart Association (ACC/AHA) Task Force on Practice Guidelines: Stage A includes patients who are at high risk for developing HF, but do not have structural heart disease; Stage B are patients who do have structural damage to the heart, but have not developed symptoms; Stage C refers to patients with past or current HF symptoms and evidence of structural heart damage; and Stage D includes patients with end-stage disease, requiring special interventions. It is the intent of the ACC/AHA recommendations to be used in conjunction with the New York Heart Association (NYHA) functional classification that estimates the severity of disease based on patient symptoms.
2. Goals of therapy for HF include improving symptoms, increasing functional capacity, improving quality of life, slowing disease progression, decreasing need for hospitalization, and prolonging survival.
3. Nonpharmacologic therapy includes abstaining from alcohol and tobacco, limiting dietary sodium, reducing weight if appropriate, exercising regularly, and influenza and pneumococcal vaccinations. Other nonpharmacologic therapies such as automatic implantable defibrillators or cardiac resynchronization therapy should be considered in appropriate patients but are beyond the scope of this document.
4. Risk factor modification and treatment of concomitant cardiac conditions and underlying causes should be implemented in patients in Stage A to potentially reduce the development of HF.
5. In addition to risk factor modification, patients in Stage B should receive post-myocardial infarction (MI) treatment with an angiotensin-converting enzyme inhibitor (ACEI) and beta-adrenergic blocker, regardless of the presence of left ventricular systolic dysfunction, to prevent future development of HF and improve overall survival (Grade A Recommendation, Good Overall Quality of Evidence). It is also recommended that patients with evidence of left ventricular systolic dysfunction who are without symptoms should be treated with an ACEI (Grade A Recommendation, Good Overall Quality of Evidence) and beta-adrenergic blocker (Grade B Recommendation, Fair Overall Quality of Evidence). An angiotensin II receptor antagonist may be prescribed in patients with a history of MI who have a reduced left ventricular ejection fraction without symptoms of HF if they are ACEI intolerant (Grade A Recommendation, Good Overall Quality of Evidence).
6. Patients with HF in Stage C should also be educated on risk factor modification. Pharmacotherapy recommendations for these patients include:
  - A diuretic should be used in the treatment of patients with signs of fluid overload (Grade B Recommendation, Fair Overall Quality of Evidence).
  - All patients should be treated with an ACEI unless contraindicated or not tolerated (Grade A Recommendation, Good Overall Quality of Evidence). These agents improve HF symptoms, functional status, and quality of life, while decreasing frequency of hospitalization and mortality. An angiotensin II receptor antagonist may be considered as an alternative to an ACEI (in patients who are on standard therapy for HF) and are unable to tolerate an ACEI (Grade A Recommendation, Good Overall Quality of Evidence).
  - A beta-adrenergic blocker that has proven to reduce mortality (i.e., bisoprolol, carvedilol, sustained release metoprolol succinate) should be used in conjunction with an ACEI in all patients who are considered stable (i.e., minimal or no signs of fluid overload or volume depletion and not in an intensive care unit), unless contraindicated or not tolerated. These agents have been shown to reduce mortality and decrease the symptoms of HF (Grade A Recommendation, Good Overall Quality of Evidence).
  - Low dose of an aldosterone antagonist should be considered in patients with recent New York Heart Association (NYHA) Class IV HF and current Class III or IV symptoms and left ventricular ejection fraction (LVEF)  $\leq 35\%$ , provided the patient has preserved renal function and normal potassium levels. This therapy improves symptoms (as assessed by change in NYHA functional class), decreases hospitalizations for worsening HF, and decreases mortality (Grade A Recommendation, Good Overall Quality of Evidence). An aldosterone antagonist may also be considered in patients with LVEF  $\leq 40\%$  in patients early post-MI on standard therapy for HF.

- The combination of hydralazine and a nitrate should be considered, especially in African American patients with NYHA Class III or IV HF, who continue to have symptoms despite therapy with an ACEI and beta-adrenergic blocker (Grade B Recommendation, Good Overall Quality of Evidence). The combination of hydralazine and a nitrate may be considered as an alternative to an ACEI in patients who are unable to tolerate an ACEI (or angiotensin II receptor antagonist) due to hypotension, renal insufficiency, hyperkalemia, or possibly, angioedema (Grade C Recommendation, Fair Overall Quality of Evidence).
  - Addition of an angiotensin II receptor antagonist to standard therapy (i.e., an ACEI and beta-adrenergic blocker) in patients with systolic HF may be considered to decrease cardiovascular death or HF hospitalizations (Grade B Recommendation, Fair Overall Quality of Evidence); although routine use of an angiotensin II receptor antagonist, ACEI, and aldosterone antagonist is not recommended.
  - Digoxin can be used in patients whose symptoms persist despite treatment with an ACEI (or an angiotensin II receptor antagonist if an ACEI is not tolerated), a beta-blocker, and a diuretic. Digoxin reduces symptoms associated with HF and decreases the risk for hospitalizations due to HF but does not improve mortality (Grade B Recommendation, Fair Overall Quality of Evidence).
  - Patients should receive regular follow-up in order to provide the most effective care. At each encounter, an inquiry should be made as to the patient's adherence to the medication regimen, nonpharmacologic measures, and adverse effects to therapy. Patients should be scheduled for routine laboratory monitoring. The patient should also be assessed for any change in functional status or frequency of hospitalizations, and medication therapy should be optimized.
7. Patients with HF in Stage D may require special treatment interventions including mechanical circulatory support, continuous therapy with positive inotropic agents, consideration for cardiac transplantation, or hospice care. In patients where therapeutic interventions may no longer be appropriate, discussions regarding end-of-life care should be considered. Specific recommendations are beyond the scope of this document, and these patients should be referred to a HF management program that includes experts on the management of patients with refractory HF.

## **The Medical Advisory Panel for the Pharmacy Benefits Management Strategic Healthcare Group**

### **Mission**

The mission of the Medical Advisory Panel (MAP) for Pharmacy Benefits Management (PBM) includes the development of evidence-based pharmacologic management guidelines for improving quality and providing best-value patient care.

The MAP comprises practicing VA physicians from facilities across the nation:

Thomas Craig, MD  
Chief Quality and Performance Officer  
Office of Quality and Performance Management  
Department of Veterans Affairs  
Washington DC

Gregory W. Dalack, MD  
Staff Psychiatrist  
VA Ann Arbor Healthcare System  
Vice Chair and Associate Chair for Education and Academic Affairs  
Department of Psychiatry  
University of Michigan

Sylvain DeLisle, MD, MBA  
Associate Chief of Staff, Performance Improvement  
VA Maryland HCS  
Associate Professor of Medicine & Physiology  
University of Maryland, School of Medicine

Thomas H. Dickinson, MD  
Co-Service Line Manager, Primary Care  
Brockton Division, VA Boston HCS  
Clinical Instructor in Medicine  
Harvard Medical School

John R. Downs, MD, FACP  
Ambulatory Care  
AMLD, South Texas Veterans HCS  
Associate Professor of Medicine  
University of Texas Health Science Center  
Col, USAF (retired)

Peter A. Glassman, MBBS, MSc  
Staff Physician  
West Los Angeles VAMC  
VA Greater Los Angeles HCS  
Associate Professor  
University of California, Los Angeles (UCLA)

Matthew Goetz, MD  
Chief, Infectious Diseases  
West Los Angeles VAMC  
VA Greater Los Angeles HCS  
Professor of Clinical Medicine  
David Geffen School of Medicine at UCLA

Chester B. Good, MD, MPH, FACP  
Chairman, Medical Advisory Panel  
Staff Physician, Department of Medicine  
VA Pittsburgh HCS  
Associate Professor of Medicine  
University of Pittsburgh

Robert C. Goodhope, MD, MBA  
Chief Medical Officer  
VA Outpatient Clinic  
Tallahassee, FL

Robert J. Hariman, MD  
Director, Cardiac Electrophysiology  
Hines VA Hospital  
Professor of Medicine  
Loyola University School of Medicine

William Korchik, MD  
Medical Director, Extended Care Center  
VAMC Minneapolis  
Assistant Professor of Medicine  
University of Minnesota

Alexander M. Shepherd, MD, PhD, FAHA  
Professor and Chief, Division of Clinical Pharmacology  
Departments of Medicine and Pharmacology  
University of Texas Health Science Center at San Antonio

## **Pharmacy Benefits Management (PBM) Strategic Healthcare Group (SHG)**

VHA's PBM-SHG has been directed by the Under Secretary for Health to coordinate the development of recommendations for the pharmacologic management of common diseases treated within the VA, establish a national level VA formulary, and to manage pharmaceutical costs, utilization, and measure outcomes as they apply to patient care. The MAP provides support and direction to the PBM staff, located in Hines, Illinois.

Michael Valentino, RPh, MHSA  
Chief Consultant, PBM-SHG

Janet Dailey, PharmD  
Clinical Pharmacy Specialist

Virginia S. Torrise, PharmD  
Deputy Chief Consultant, PBM-SHG

Elaine Furmaga, PharmD  
Clinical Pharmacy Specialist

Joseph Canzolino, RPh  
Associate Chief Consultant, PBM-SHG

Mark Geraci, PharmD, BCOP  
Clinical Pharmacy Specialist

Vincent Calabrese, PharmD  
Assistant Chief Consultant, PBM-SHG

Francine Goodman, PharmD, BCPS  
Clinical Pharmacy Specialist

Fran Cunningham, PharmD  
Director, Center for Medication Safety PSCI and  
Program Manager Pharmacoepidemiologic/  
Outcomes Research

Cathy Kelley, PharmD  
Clinical Pharmacy Specialist

Deborah Khachikian, PharmD  
Clinical Pharmacy Specialist

Muriel Burk, PharmD  
Clinical Pharmacy Specialist, Pharmacoepidemiologic  
and Outcomes Research

Lisa Longo, PharmD, BCPS  
Clinical Pharmacy Specialist

Marie Sales, PharmD  
Clinical Pharmacy Specialist, Pharmacoepidemiologic  
and Outcomes Research

Todd Semla, PharmD, BCPS  
Clinical Pharmacy Specialist

Kathy Tortorice, PharmD, BCPS  
Clinical Pharmacy Specialist

Suzanne Lenz, RPh  
Pharmacist Specialist/Contract Liaison

Lucy Szudarski  
Program Specialist

Lisa Torphy  
Program Specialist

## Acknowledgements

This document was developed in consultation with members of the PBM-MAP and subject matter experts in cardiology. A draft was then forwarded to the field through the VISN Formulary Leaders for peer review. The final version was forwarded to the Director of Performance Management and the National Advisory Council for the Adoption, Development and Implementation of Clinical Practice Guidelines for approval. The following clinicians provided comments on drafts of this report. The final document incorporates reviewers' comments; however, the PBM-SHG takes full responsibility for the content of this guideline.

Alan Cohen, MD, FACP  
Chief of Primary Care,  
VA Central California Healthcare System

Gregg C. Fonarow, MD  
Member, VA CHF QUERI Executive Committee  
Professor of Cardiovascular Medicine and Science,  
David Geffen School of Medicine at UCLA

Charles Hoesch, MD  
Medical Director, Perry Point VAMC  
VA Maryland Healthcare System

David Parra, PharmD, BCPS  
Clinical Pharmacy Specialist in Cardiology,  
West Palm Beach VAMC  
Clinical Assistant Professor,  
Department of Experimental and Clinical Pharmacology  
University of Minnesota College of Pharmacy

Eugene S. Smith, III, MD  
Acting Chief of Cardiology,  
Central Arkansas Veterans Healthcare System  
Associate Professor of Medicine, Division of Cardiology  
University of Arkansas for Medical Sciences

---

\* This list does not represent all the clinicians who reviewed the document, only those who agreed to be acknowledged. Reviewers of prior versions have been previously acknowledged.

## TABLE OF CONTENTS

EXECUTIVE SUMMARY.....	ii
ABBREVIATIONS.....	viii
INTRODUCTION.....	1
TREATMENT ALGORITHM.....	6
ANNOTATIONS.....	8
A. Diagnose and Evaluate a Patient at Risk for or Suspected of Having Heart Failure (HF).....	9
B. Management of Concomitant Cardiac Conditions and Risk Factors, Nonpharmacologic Interventions, and Treatment of Underlying Causes .....	12
C. Pharmacologic Management of HF with Preserved Left Ventricular Ejection Fraction.....	15
D. Interventions in Patients With Asymptomatic Left Ventricular Systolic Dysfunction .....	17
E. Systolic Dysfunction and Assessment for Symptoms of Volume Overload .....	19
F. Diuretic Therapy .....	20
G. Angiotensin-Converting Enzyme Inhibitors .....	21
H. Beta-Adrenergic Blockers.....	24
I. Angiotensin II Receptor Antagonists .....	27
J. Hydralazine in Combination with a Nitrate .....	31
K. Digitalis.....	32
L. Aldosterone Antagonists.....	34
M. Continue Present Management and Schedule Regular Follow-up.....	36
REFERENCES.....	38

## TABLES

Table 1. Quality of Evidence.....	3
Table 2. Overall Quality.....	3
Table 3. Net Effect of Intervention.....	3
Table 4. Strength of Recommendation.....	4
Table 5. Grade for Strength of Recommendation .....	4
Table 6. ACC/AHA Class of Recommendation.....	4
Table 7. ACC/AHA Level of Evidence.....	4
Table 8. ACC/AHA Guidelines for the Evaluation and Management of HF.....	11
Table 9. NYHA Functional Classification and Objective Assessment of HF.....	12

## FIGURE

Figure. VA Utilization of Medications for HF by Drug Class .....	50
--	----

## APPENDICES

Appendix A. Medications Commonly Used in the Management of Heart Failure.....	51
Appendix B. Common Drug Interactions with Agents Used in Heart Failure.....	53
Appendix C: Long-term, Randomized, Controlled, Outcome Trials in Systolic HF by Drug Class.....	56

## ABBREVIATIONS

---

AIIRA (also ARB)	Angiotensin II receptor antagonist (also referred to as angiotensin receptor blocker)
ACC/AHA	American College of Cardiology/American Heart Association
ACEI	Angiotensin-converting enzyme inhibitor
ARR	Absolute risk reduction
AV	Atrioventricular
BNP	Brain natriuretic peptide
BPH	Benign prostatic hyperplasia
BUN	Blood urea nitrogen
CCB	Calcium channel blocker
CI	95% confidence interval
Cr	Creatinine
CrCl	Creatinine clearance
DM	Diabetes mellitus
DOE	Dyspnea on exertion
HCTZ	Hydrochlorothiazide
HF	Heart failure
HFSA	Heart Failure Society of America
HTN	Hypertension
INR	International normalization ratio
ISDN	Isosorbide dinitrate
JVD	Jugular venous distention
K <sup>+</sup>	Potassium
LV	Left ventricular
LVEF	Left ventricular ejection fraction
LVEDP	Left ventricular end diastolic pressure
LVH	Left ventricular hypertrophy
MI	Myocardial infarction
NNT	Number needed to treat
NSAID	Non-steroidal anti-inflammatory drug
NYHA	New York Heart Association
PND	Paroxysmal nocturnal dyspnea
RR	Relative risk
SNS	Sympathetic nervous system
SOB	Shortness of breath
TSH	Thyroid-stimulating hormone (thyrotropin)
TZD	Thiazolidinedione

---





# INTRODUCTION

## THE PHARMACOLOGIC MANAGEMENT OF CHRONIC HEART FAILURE

Approximately 5 million patients in the United States have heart failure (HF), with 550,000 new cases diagnosed each year.<sup>1,2</sup> The prevalence of HF increases with age, with nearly 5% of patients seen at Veterans Affairs (VA) Medical Centers having a primary diagnosis of HF. According to a recent report from the American Heart Association, 80% of men and 70% of women with a diagnosis of HF who are less than 65 years of age will die within 8 years. In addition, the one year mortality rate was reported as 20%.<sup>2</sup> Hospitalizations for HF have increased,<sup>1,2</sup> accounting for 6.5 million hospital days annually.<sup>1</sup> Heart failure was also reported to be the main reason for 12 to 15 million clinic visits each year.<sup>1</sup> The VA provides care for approximately 240,000 veterans with HF, with over 42,000 of these patients being hospitalized during Fiscal Year (FY) 2005 with a primary diagnosis of HF. It has been estimated that the cost for HF in 2006 is \$29.6 billion in the United States alone.<sup>1</sup>

The pharmacologic treatment of HF has advanced significantly over the years, with clinical research establishing the benefit of drug therapy in preventing morbidity and mortality in patients with this condition.<sup>1</sup> The clinical outcomes and resulting economic benefits of drug therapy have also been documented in the clinical practice setting.<sup>3</sup> With HF being such a prevalent disease, especially among the older patient population, and with it a high rate of morbidity and mortality, it is prudent that evidence-based therapy and associated clinical practice guidelines be utilized to improve patient outcomes.<sup>4,5</sup>

Since the beginning of fiscal year (FY) 2003, the VA Office of Quality and Performance has implemented performance measures evaluating the treatment of HF with the angiotensin-converting enzyme inhibitors (ACEIs), including data on the angiotensin II receptor antagonists as part of the measure in FY2005. After annual chart review of approximately 3,000 veterans, the External Peer Review Program reported that nearly 90% of these veterans with HF were prescribed an ACEI or angiotensin II receptor antagonist (refer to <http://vaww.pdw.med.va.gov/pdwframe.asp> for details on the indicators and updated results).

Utilization of beta-adrenergic blockers and renin-angiotensin-aldosterone inhibitors have continued to increase in veteran patients with HF (refer to the Figure); although use of these drug classes in combination is less than optimal. Continued efforts to optimize evidence-based therapy should be encouraged.

These clinical practice guidelines for the management of patients with HF focus on the pharmacologic treatment of the disease. The clinician is referred to the American College of Cardiology/American Heart Association (ACC/AHA) Guideline Update for the Diagnosis and Management of Chronic Heart Failure in the Adult<sup>1</sup> and other medical literature and cardiology experts for the overall management of patients with HF.

### Summary

This consensus and evidence-based document on the pharmacologic management of patients with chronic HF is intended to update the August 2003 publication of the PBM-MAP The Pharmacologic Management of Chronic Heart Failure. Whenever possible, the PBM and MAP rely upon evidence-based, multidisciplinary, nationally recognized consensus statements for the basis of VA guidance. Relevant literature is reviewed and assessed with consideration given to the VA population. Draft documents are sent to the field for comments prior to being finalized.

### Development Process and Sources of Information

Development of the recommendations included reference to the following consensus document: *Hunt SA, Abraham WT, Chin MH, et al. ACC/AHA 2005 guideline update for the diagnosis and management of chronic heart failure in the adult: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Update the 2001 Guidelines for the Evaluation and Management of Heart Failure). American College of Cardiology Web site. Available at: <http://www.acc.org/clinical/guidelines/failure/index.pdf>.*

To update the August 2003 PBM-MAP guideline “The Pharmacologic Management of Chronic Heart Failure”, a literature search of the National Library of Medicine’s MEDLINE/PubMed database and Evidence Based Medicine reviews available on OVID was conducted. Preference was given to randomized controlled trials, meta-analyses, and systematic reviews. The following search terms were used: heart failure, angiotensin-converting enzyme inhibitor, beta-adrenergic blocker, digoxin, spironolactone, angiotensin receptor blocker, aldosterone antagonist, hydralazine, isosorbide dinitrate, diastolic dysfunction, clinical trial, review, meta-analysis. The literature was limited to adult human subjects and articles published in the English language. The bibliographies of articles and consensus documents were reviewed for additional relevant literature. In updating the December 2006 document, 208 abstracts and 144 articles were reviewed. One hundred thirty-seven articles were added to the update of the 2006 document, 49 of which were randomized controlled trials. In addition to randomized controlled trials of patients with a diagnosis of chronic HF, the references added to the annotations discussing recommendations for specific pharmacologic classes or HF in general included 67 pertinent subgroup or retrospective evaluations, 9 meta-analyses or systematic reviews of controlled trials relevant to the recommendations in the document, 5 case reports and 7 review articles, some that provided a comprehensive inclusion of information and others that discussed patient care considerations not addressed by clinical trials. Literature known to the PBM-MAP on medical history, physical examination, diagnosis and evaluation, consensus statements and clinical practice guidelines were also included in the document. Major changes to the 2006 update include addition of pertinent medical evidence published since the last iteration of the document including data with the angiotensin II receptor antagonists, isosorbide dinitrate and hydralazine, the aldosterone antagonists, and a comparison of an ACEI versus a beta-adrenergic blocker as initial therapy. The treatment algorithm has been revised to reflect the recommended place in therapy of these drug classes based on this information. The document has been reformatted as per other VA/DoD clinical practice guidelines. In addition, relevant data from long-term outcome trials in patients with chronic HF due to systolic dysfunction have been compiled in the Appendix.

## Methods to Formulate Recommendations

The literature was critically analyzed and evidence was graded using a standardized format. The evidence rating system for this document is based on the system used by the U.S. Preventative Services Task Force and also references the grading system used in the ACC/AHA Practice Guidelines for the Evaluation and Management of HF. The rating scale of the U.S. Preventive Services Task Force is summarized in Tables 1 to 5.<sup>6</sup>

<b>Table 1    Quality of Evidence (QE)</b>	
<b>I</b>	Evidence obtained from at least one properly randomized controlled trial
<b>II-1</b>	Evidence obtained from well-designed controlled trials without randomization
<b>II-2</b>	Evidence obtained from well-designed cohort or case-control analytic studies
<b>II-3</b>	Evidence obtained from multiple time series studies; dramatic results in uncontrolled experiments
<b>III</b>	Opinions of respected authorities; descriptive studies and case reports; reports of expert committees

<b>Table 2    Overall Quality (OQ)</b>	
<b>I    Good</b>	High grade evidence (I or II-1) directly linked to health outcome
<b>II   Fair</b>	High grade evidence (I or II-1) linked to intermediate outcome or Moderate grade evidence (II-2 or II-3) directly linked to health outcome
<b>III   Poor</b>	Level III evidence or no linkage of evidence to health outcome

<b>Table 3    Net Effect of Intervention</b>	
<b>Substantial</b>	More than a small relative impact on a frequent condition with a substantial burden of suffering, or A large impact on an infrequent condition with a significant impact on the individual patient level
<b>Moderate</b>	A small relative impact on a frequent condition with a substantial burden of suffering, or A moderate impact on an infrequent condition with a significant impact on the individual patient level
<b>Small</b>	A negligible relative impact on a frequent condition with a substantial burden of suffering, or A small impact on an infrequent condition with a significant impact on the individual patient level
<b>Zero or Negative</b>	Negative impact on patients, or No relative impact on either a frequent condition with a substantial burden of suffering, or An infrequent condition with a significant impact on the individual patient level

<b>Table 4 Strength of Recommendation</b>	
<b>A</b>	A strong recommendation that the intervention is always indicated and acceptable
<b>B</b>	A recommendation that the intervention may be useful/effective
<b>C</b>	A recommendation that the intervention be considered
<b>D</b>	A recommendation that an intervention may be considered not useful/effective, or may be harmful
<b>I</b>	Insufficient evidence to recommend for or against; clinical judgment should be used

<b>Table 5 Grade for Strength of Recommendation</b>				
<b>Overall Quality of Evidence</b>	<b>Net benefit of intervention</b>			
	<b>Substantial</b>	<b>Moderate</b>	<b>Small</b>	<b>Zero or Negative</b>
<b>I</b>	A	B	C	D
<b>II</b>	B	B	C	D
<b>III</b>	I	I	I	D

The evidence rating system used in the ACC/AHA Practice Guidelines on the Evaluation and Management of HF are included below.<sup>1</sup> As this is used by ACC/AHA guidelines, this format will also be included in the recommendations in the text to assist in the application of the recommendations to clinical practice.

<b>Table 6 ACC/AHA Class of Recommendation</b>	
<b>Class</b>	<b>Recommendation</b>
<b>I</b>	Conditions for which there is evidence and/or general agreement that a given procedure/therapy is useful and effective
<b>II</b>	Conditions for which there is conflicting evidence and/or a divergence of opinion about usefulness/efficacy of performing the procedure/therapy
<b>Ila</b>	Weight of evidence/opinion is in favor of usefulness/efficacy
<b>Ilb</b>	Usefulness/efficacy is less well established by evidence/opinion
<b>III</b>	Conditions for which there is evidence and/or general agreement that a procedure/therapy is not useful/effective and in some cases may be harmful

<b>Table 7 ACC/AHA Level of Evidence</b>	
<b>A</b>	Data is derived from multiple randomized clinical trials or meta-analyses
<b>B</b>	Data is derived from a single randomized trial or nonrandomized studies
<b>C</b>	Consensus opinion of experts is the primary source of recommendation

Recommendations were based on evidence published in the medical literature. Critical literature review focused on pharmacologic management of HF. The annotations that include discussion on medical history, physical examination, diagnosis and evaluation, nonpharmacologic intervention, management of concomitant cardiac conditions, and treatment of underlying causes were based on consensus and did not undergo critical literature review. Where evidence was not available, expert opinion of the MAP and cardiology expert reviewers were used. After review and discussion by the PBM-MAP, the draft guideline was sent to experts in the field of Cardiology for review. After the Cardiologist reviewers' comments were considered and incorporated into the document where appropriate, the draft was then circulated to practicing clinicians (primarily cardiologists and primary care providers) for input on clarity and applicability.

## Use of the Document

The document is divided into four sections: Executive Summary, Algorithm, Annotations, and Appendices. The algorithm is intended to provide a systematic approach to the pharmacologic management of patients with HF. The letters within the boxes in the algorithm refer to the corresponding annotation. The annotation is further discussion of the evidence for making each recommendation. Details on drug therapy are provided to encourage the safe and effective implementation of the pharmacotherapy recommendations made in this guideline. Recommendations discussed in the annotations on pharmacotherapy are referenced and graded according to the grading system outlined above. The appendices provide additional information for the clinician when considering treatment options.

The recommendations are meant to focus on the pharmacologic management of patients with HF. Other sections have been included that highlight areas such as physical examination, diagnosis,

nonpharmacologic management, etc. Practitioners should refer to comprehensive clinical practice guideline on HF, cardiology texts or local experts for the finer points of diagnosis and these other areas.

The purpose of the recommendations is to assist practitioners in clinical decision-making, to standardize and improve the quality of patient care, and to promote cost-effective drug prescribing. This document attempts to define principles of practice that should produce high quality patient care. They are attuned to the needs of a primary care practice but are directed to providers at all levels. Care of patients with HF may occur in several clinical settings including primary care, cardiology, or by multidisciplinary HF treatment teams. Regardless of the setting in which patients with HF are cared for, the clinician is encouraged to follow these and other HF guidelines and to use clinical judgment of when to refer to a specialist. This will depend on the skill and experience of managing patients with HF, and also the resources available to the practitioner. The recommendations also serve as a basis for monitoring local, regional and national patterns of pharmacologic care.

The recommendations in this document should not be considered inclusive of all proper methods of care or exclusive of other methods of care reasonably directed at obtaining the same results. The ultimate judgment regarding the propriety of any course of conduct must be made by the clinician in light of individual patient situations.

## **Plan for Implementation**

The document will be available on the PBM home page at [www.pbm.va.gov](http://www.pbm.va.gov) or <http://vaww.pbm.va.gov> as well as the VA Office of Quality and Performance at [www.oqp.med.va.gov](http://www.oqp.med.va.gov). It is recommended that a hard copy be kept on file in the medical libraries. Distribution to all clinicians who manage patients with HF is strongly recommended. Clinicians are encouraged to have a copy of the document or a summary of key points available for reference when treating patients with HF.

Continuing education programs will be developed.

Departmental and individual education at the facility is also encouraged.

## **Referencing the Document**

This document should be referenced as follows:


*The Pharmacologic Management of Chronic Heart Failure*. Washington, DC: Pharmacy Benefits Management Strategic Healthcare Group and the Medical Advisory Panel, Veterans Health Administration, Department of Veterans Affairs. September 2007 Update. PBM-MAP Publication No. 00-0015.

## **Updating the Guideline**

The PBM will review the guideline routinely. Updating will occur as new information is made available from well-designed, scientifically valid studies and as outcome data may direct. Any member of the VA community is encouraged to recommend changes based on such evidence.

A current copy of the pharmacologic management guideline can be obtained from the PBM home page at [www.pbm.va.gov](http://www.pbm.va.gov) or <http://vaww.pbm.va.gov>.

# **TREATMENT ALGORITHM**





# ANNOTATIONS

# THE PHARMACOLOGIC MANAGEMENT OF CHRONIC HEART FAILURE

## Annotations

### A. Diagnose and Evaluate a Patient at Risk for or Suspected of Having Heart Failure (HF)

#### OBJECTIVES

- To identify patient factors associated with HF
- To distinguish between the diagnosis of HF and other conditions, such as pulmonary, hepatic, renal, hematopoietic diseases that can produce symptoms or signs suggestive of HF
- To distinguish systolic from diastolic dysfunction
- To evaluate the patient's functional status

#### BACKGROUND

As previously discussed in the Introduction, HF is a prevalent condition, especially in the older patient population, with a high rate of morbidity and mortality.<sup>1</sup> This treatment guideline focuses on the pharmacologic management of HF; the clinician is referred to other resources including comprehensive treatment guidelines or clinical expertise in the diagnosis and evaluation of patients with HF. The recommendations below are to be used as a general guide and are a summary of Class I Recommendations where there is evidence/consensus that the treatment/procedure is of benefit as per the ACC/AHA Practice Guidelines on initial and follow-up assessments of patients with HF.<sup>1</sup>

#### RECOMMENDATIONS

The following are Class I recommendations by the ACC/AHA (i.e., there is evidence and/or general agreement that a given procedure/therapy is useful and effective).<sup>1</sup>

##### Initial Assessment

- Obtain a thorough medical history and perform a comprehensive physical examination in patients suspected of having HF to identify risk factors or conditions that may lead to the development or progression of HF
- Obtain a comprehensive medication history (including past and current treatments for HF, alternative therapies, and antineoplastic agents) and past or current alcohol or illicit drug use
- Perform an assessment of the patient's ability to perform activities of daily living
- Document the patient's height, weight, and body mass index; assess the patient's volume status and evaluate for orthostatic blood pressure changes
- Obtain baseline laboratory parameters including a complete blood count, serum electrolytes with calcium and magnesium, blood urea nitrogen (BUN), serum creatinine (Cr), fasting blood glucose and glycosylated hemoglobin, lipid profile, liver function tests, and thyroid-stimulating hormone (TSH)
- Perform a twelve-lead electrocardiogram and chest radiograph
- Perform a two-dimensional echocardiography with Doppler to assess left ventricular ejection fraction (LVEF), left ventricular (LV) size, wall thickness, and valve function; radionuclide ventriculography to assess LVEF and volumes can also be performed
- Perform a coronary arteriography in patients with HF and angina or significant ischemia (unless not eligible candidate for revascularization)

##### Follow-Up Evaluations

- Perform an assessment of the patient's ability to perform activities of daily living at each clinic visit
- Assess the patient's volume status and weight at each clinic visit
- Inquire as to the patient's current use of alcohol, illicit drugs, alternative therapies, or antineoplastic agents; evaluate the patient's diet and intake of sodium

## DISCUSSION

Heart failure is defined as a “complex clinical syndrome that can result from any structural or functional cardiac disorder that impairs the ability of the ventricle to fill with or eject blood.”<sup>1</sup> The leading causes of HF are coronary artery disease, hypertension (HTN), and dilated cardiomyopathy (i.e., a structural abnormality).<sup>1</sup> In addition, identification of other conditions or risk factors contributing to the development or progression of HF is important as some of these may be treated or avoided.<sup>1</sup>

Patients with heart failure typically present with symptoms including dyspnea and fatigue, as well as edema and rales on physical examination.<sup>1</sup> Signs and symptoms of HF are nonspecific and must be distinguished from those of other conditions such as pulmonary disease, liver failure, and/or nephrotic syndrome. Heart failure due to myocardial muscle dysfunction may be characterized by systolic dysfunction, diastolic dysfunction, or both. Systolic dysfunction is generally defined as a LVEF of < 40%.<sup>1</sup> Patients with diastolic dysfunction often have impaired ventricular relaxation and distensibility resulting in increased ventricular filling pressure (LVEDP). The ejection fraction in these patients may be normal or increased.

### Medical history<sup>1</sup>

- Coronary artery disease
- Hypertension
- Valvular heart disease
- Diabetes
- Peripheral vascular disease
- Dyslipidemia
- Myopathy
- Rheumatic fever
- Mediastinal irradiation
- Sleep-disordered breathing
- Exposure to cardiotoxic agents (e.g., anthracyclines, trastuzumab, ephedra, high-dose cyclophosphamide)
- Alcohol or illicit drug use
- Smoking
- Exposure to sexually transmitted diseases
- Thyroid disorder
- Pheochromocytoma
- Obesity
- Family history of atherosclerotic disease, cardiomyopathy, sudden death, conduction system disease, skeletal myopathies, or tachyarrhythmias

Patient presentation: Patients with LV dysfunction generally present in one of the following manners:<sup>1</sup>

- Decreased exercise tolerance
- Fluid retention
- Cardiac enlargement or dysfunction noted during evaluation for a condition other than HF

Patient symptoms of HF:<sup>1,7,8</sup> Most patients will present with complaints of exercise intolerance due to dyspnea and/or fatigue. However, no symptom is sufficiently sensitive or specific for the diagnosis of HF to allow ruling in or out disease. Patients with at least one of the following symptoms are at somewhat higher likelihood of having HF. Some patients with HF may have only signs of the condition without any active symptoms.

- Shortness of breath (SOB)
- Fatigue

- Orthopnea
- Paroxysmal nocturnal dyspnea (PND)
- Dyspnea on exertion (DOE)
- Cough
- Edema
- Weight gain (anorexia may be seen in advanced HF)

Physical examination findings of HF:<sup>7,8</sup> No single finding is sufficiently sensitive or specific for use alone in the diagnosis of HF. However, patients with at least one of the following signs are more likely to have HF. Some patients with HF may only have symptoms of the condition without any of the physical signs listed below.

- Tachycardia
- Increasing weight
- Jugular venous distention (JVD) or hepatjugular reflux
- Presence of S<sub>3</sub> (third heart sound)
- Laterally displaced apical impulse
- Pulmonary crackles or wheezes
- Hepatomegaly
- Peripheral (dependent) edema
- Abdominal distention or ascites

Laboratory Assessment:<sup>1</sup> Laboratory parameters are recommended to evaluate the patient for conditions that may contribute to the development or exacerbation of HF. The initial assessment should include:

- Complete blood count
- Serum electrolytes with calcium and magnesium
- Blood urea nitrogen, serum Cr
- Fasting blood glucose and glycosylated hemoglobin
- Lipid profile
- Liver function tests
- Thyroid-stimulating hormone
- B-type natriuretic peptide (BNP): elevated levels may be helpful in diagnosing a patient suspected of having HF or used to consider a diagnosis of HF when the diagnosis is unknown. The ACC/AHA recommends (i.e., weight of evidence/opinion is in favor of usefulness/efficacy) that the measurement of BNP may be useful in evaluating patients who present short of breath to the urgent care setting where the diagnosis of HF may be uncertain.<sup>1,9</sup> The ACC/AHA also states that the value of serial BNP measurements to guide therapy is not well established<sup>1</sup>

### Classification of HF

Different classification systems help characterize HF based on cardiac cycle (systolic, diastolic or both), and/or ventricular involvement (right, left or both). The following recommendations of the ACC/AHA are for staging patients with HF based on the progression of disease.<sup>1</sup>

<b>Table 8 ACC/AHA Guidelines for the Evaluation and Management of HF<sup>1</sup></b>	
<b>Stage</b>	<b>Disease Progression</b>
<b>A</b>	Patients who are high risk for developing HF, but do not have structural heart disease
<b>B</b>	Patients who have structural damage to the heart, but have not developed symptoms
<b>C</b>	Patients with past or current HF symptoms and evidence of structural heart damage
<b>D</b>	Patients with end-stage disease, requiring special interventions

It is the intent of the ACC/AHA recommendations to be used in conjunction with the New York Heart Association (NYHA) functional classification<sup>10</sup> that estimates the severity of disease based on patient

symptoms.<sup>1</sup> According to the above classification system, once a patient develops symptoms they should be treated according to the recommendations for patients with Stage C (even if NYHA Class I, see below), and do not return to Stage B.<sup>1</sup>

<b>Table 9 NYHA Functional Classification and Objective Assessment of HF<sup>10</sup></b>	
<b>Class</b>	<b>Disease Progression</b>
<b>I</b>	No limitation of physical activity. Ordinary physical activity does not cause undue fatigue, palpitation, dyspnea, or angina
<b>II</b>	Slight limitation of physical activity. Ordinary physical activity results in fatigue, palpitation, dyspnea, or angina
<b>III</b>	Marked limitation of physical activity. Comfortable at rest, but less than ordinary physical activity results in fatigue, palpitation, dyspnea, or angina
<b>IV</b>	Unable to carry on any physical activity without discomfort. Symptoms are present at rest. With any physical activity, symptoms increase

## **B. Management of Concomitant Cardiac Conditions and Risk Factors, Nonpharmacologic Interventions, and Treatment of Underlying Causes**

### **OBJECTIVES**

- To provide general interventions to be recommended in patients at risk for developing HF or who have a diagnosis of HF

### **BACKGROUND**

Identification and treatment of chronic medical conditions or risk factors that may impact the development or progression of HF is important in the overall management of patients at risk for HF or who have been diagnosed with the condition. Many of these diseases have clinical practice guidelines that have been reviewed and approved by the VA/DoD Evidence-Based Practice Work Group and can be found on the VA Office of Quality and Performance Web site at <http://www.oqp.med.va.gov/cpg/cpg/htm>.

### **RECOMMENDATIONS**

#### Control risk factors

The following are Class I recommendations by the ACC/AHA (i.e., there is evidence and/or general agreement that a given procedure/therapy is useful and effective)<sup>1</sup>

- Control HTN (<http://www.oqp.med.va.gov/cpg/cpg/htm>)
- Treat hyperlipidemia (<http://www.oqp.med.va.gov/cpg/cpg/htm>)
- Treat DM (<http://www.oqp.med.va.gov/cpg/cpg/htm>)
- Encourage smoking cessation (<http://www.oqp.med.va.gov/cpg/cpg/htm>) and discourage alcohol consumption and illicit drug use
- Control ventricular rate or restore sinus rhythm in patients with supraventricular tachyarrhythmias
- Treat thyroid disorders
- Conduct periodic evaluations for signs and symptoms of HF
- Manage atherosclerotic vascular disease
- For those with a family history of cardiomyopathy or who are receiving cardiotoxic medications, perform a noninvasive evaluation of LV

### **DISCUSSION**

#### Treatment of conditions that may lead to HF

Patients with hypertension, diabetes mellitus, or atherosclerotic vascular disease, or those who smoke tobacco are at an increased risk for the development of HF.<sup>1,11,12</sup> Treatment of these conditions and other

risk factors can contribute to an improvement in patient outcomes,<sup>1,13-19</sup> and it is recommended that patients be treated according to the corresponding VA/DoD clinical practice guidelines (available at <http://www.oqp.med.va.gov/cpg/cpg/htm>).

### Recommendations in selected patients

*Atrial fibrillation:* In patients with HF due to systolic dysfunction and atrial fibrillation requiring rate control, a beta-adrenergic blocker is preferred due to its favorable effect on patients with HF (in patients that are hemodynamically and otherwise stable).<sup>1</sup> Digoxin is also commonly used. Some patients may require combination therapy with digoxin and a beta-adrenergic blocker.<sup>20</sup> Although the long-term use of diltiazem and verapamil [calcium channel blockers (CCBs) with atrioventricular (AV) nodal blocking activity] have been associated with worsening LV dysfunction in patients with HF, patients with atrial fibrillation and rapid ventricular response resistant to combinations of digoxin and a beta-adrenergic blocker have responded with better rate control by adding diltiazem or verapamil. The decision to add diltiazem or verapamil in such patients should be based on weighing the benefit of better rate control against the deleterious long-term effects of these drugs. If additional rate control is needed, referral should be made to a cardiologist with expertise in electrophysiology.

*Anticoagulation:* Warfarin anticoagulation [with a target international normalization ratio (INR) of 2.5; range 2.0 to 3.0] is recommended in patients with HF and atrial fibrillation, previous systemic or pulmonary thromboembolism, or mobile LV thrombus.<sup>21,22</sup> The routine use of warfarin anticoagulation for HF has not been confirmed by controlled clinical trials.<sup>1,23</sup> The Warfarin and Antiplatelet Therapy in Chronic Heart Failure (WATCH) trial, was unable to demonstrate a significant difference between warfarin, aspirin, and clopidogrel in patients with HF. Another trial comparing warfarin to aspirin in patients with HF is currently underway.<sup>24</sup> Arterial thromboembolism may occur in patients with HF due to systolic dysfunction as a result of the low cardiac output and poor contractility. Analysis of cohorts in the Studies of Left Ventricular Dysfunction (SOLVD) who received warfarin, compared to those who did not, suggests a 25% risk reduction in all-cause mortality.<sup>25</sup> However, a post-hoc analysis of a single study is not evidence enough to recommend anticoagulation in patients with systolic dysfunction. Patients with atrial fibrillation and HF with contraindications to warfarin (e.g., increased risk of bleeding, difficulty adhering to the medication regimen or regular INR monitoring, current alcohol abuse or frequent falls) should receive aspirin unless contraindicated.<sup>26,27</sup> For patients with HF who do not have coronary disease, additional information is needed as to the risk vs. benefit of recommending aspirin therapy.<sup>28,29</sup>

*Concomitant HTN and/or angina:* Patients with systolic HF and concomitant HTN should be maximized on therapy with agents such as diuretics, ACEIs, and beta-adrenergic blockers, or beta-adrenergic blockers and nitrates in patients with concomitant angina, before adding other agents. However, in patients who are not adequately controlled on these agents, treatment with a long-acting dihydropyridine (felodipine or amlodipine) may be considered based on the following information.

The negative inotropic properties of the CCBs may cause deleterious effects in patients with HF due to systolic dysfunction. Studies have looked at the use of the long-acting dihydropyridines, felodipine and amlodipine, in patients with systolic dysfunction (Note: neither amlodipine nor felodipine have approval by the Food and Drug Administration for use in patients with HF and should be used with caution in patients with this diagnosis).

The Prospective Randomized Amlodipine Survival Evaluation (PRAISE) evaluated patients with NYHA class IIIB or IV with a LVEF of < 30%, who remained symptomatic despite treatment with digoxin, diuretics, and an ACEI. There were 571 patients who received amlodipine up to 10mg once daily compared to 582 patients on placebo. The average follow-up was 13.8 months (range 6-33). There was no significant difference in the primary endpoint (combined risk of death and major cardiovascular hospitalizations) between groups. There was a trend toward a decrease in all-cause mortality with amlodipine (p=0.07). Subgroup analysis showed that amlodipine significantly decreased the risk of death from all causes in patients with HF due to nonischemic dilated cardiomyopathy, without a difference in patients with ischemic dilated cardiomyopathy.<sup>30</sup> This result was not considered *a priori* endpoint. The

survival benefit of amlodipine in patients with nonischemic dilated cardiomyopathy found in the original PRAISE trial was not confirmed in PRAISE-2.<sup>31</sup>

The third Vasodilator Heart Failure Trial (V-HeFT III) included patients with NYHA class II or III HF with a LVEF of 18-42% who remained symptomatic despite treatment with digoxin, diuretics, and an ACEI. There were 224 patients who received felodipine at a maximum dose of 5mg twice daily compared to 226 patients on placebo. The average follow-up was 18 months (range 3-39). The primary endpoint of the study was the effect of treatment on exercise tolerance. There was no significant difference between groups in death from all causes, worsening HF, or number of hospitalizations. This study was not sufficiently powered to demonstrate that felodipine did not alter mortality, however. Exercise tolerance and quality of life significantly improved with felodipine at 27 months.<sup>32</sup>

Clinical practice guidelines have stated that only trials with amlodipine and felodipine have provided long-term safety data in patients with HF.<sup>22</sup> The evidence with amlodipine suggests that this agent does not adversely affect survival in patients with systolic HF. Felodipine or amlodipine may be considered in patients with HF due to systolic dysfunction for the treatment of hypertension for those who have been maximized on pharmacotherapy including diuretics, ACEIs, and beta-adrenergic blockers, and an angiotensin II receptor antagonist, hydralazine/nitrate, or aldosterone antagonist, as indicated; or beta-adrenergic blockers and long-acting nitrates in patients with concomitant angina.

*Cardiac amyloidosis:* If cardiac amyloidosis is known or suspected from echocardiography or clinical grounds, further work-up and referral to a cardiologist is warranted for appropriate treatment.

Conventional wisdom has been that digoxin and CCBs should be avoided in patients with amyloid cardiomyopathy.<sup>33,34</sup> However, this point is controversial<sup>35</sup> and supported by only weak published evidence. Several case reports suggest a sensitivity to digoxin,<sup>36-41</sup> however one prospective autopsy study found no association.<sup>42</sup> Digoxin can be useful in controlling rapid ventricular response to atrial fibrillation and might be useful, especially in early stages of systolic dysfunction caused by amyloid cardiomyopathy.<sup>35</sup> The data supporting a CCB sensitivity is based on case reports for nifedipine<sup>43,44</sup> and verapamil.<sup>45</sup> Both these drugs can exacerbate chronic systolic dysfunction independent of etiology. No case reports of other CCBs have been found to suggest sensitivity to them. The following recommendations are based on review of available evidence:

- Avoid verapamil and diltiazem (except in patients with atrial fibrillation and rapid ventricular rate that do not achieve rate control on a beta-adrenergic blocker and digoxin), and nifedipine in systolic dysfunction of all etiologies
- If digoxin is necessary in a patient with known or suspected amyloid cardiomyopathy (e.g., to control ventricular response to atrial fibrillation), it should be used very cautiously with careful monitoring for evidence of cardiac toxicity
- Use digoxin in severe cases of known or suspected amyloid cardiomyopathy only in close consultation with a cardiologist and after carefully weighing the potential risks and benefits
- Use felodipine or amlodipine only according to prescribing guidelines; monitor patients with known or suspected amyloid cardiomyopathy very closely when using any CCB
- Consider using other agents for diastolic dysfunction before resorting to a CCB in patients with known or suspected amyloid cardiomyopathy

#### Medications to avoid or to be used with caution

- **Anti-arrhythmic agents:** Anti-arrhythmic agents, other than beta-adrenergic blockers, are not recommended to suppress asymptomatic ventricular arrhythmia or ectopy. Class I antiarrhythmic agents have been shown to increase the risk of sudden death in patients with HF. Of the class III agents, treatment with amiodarone or dofetilide does not appear to increase the risk of death in patients with HF.<sup>46-49</sup> Patients with ventricular arrhythmias should be referred to a cardiologist for individualized treatment



- **Calcium channel blockers:** Most CCBs (except felodipine and amlodipine; refer to above discussion) should not be used in patients with systolic dysfunction
- **Non-steroidal anti-inflammatory drugs (NSAIDs):** NSAIDs may cause fluid retention and should be avoided;<sup>1,50,51</sup> alternative anti-inflammatory agents may be used (e.g., non-acetylated salicylates)
- **Antineoplastic agents:** Antineoplastic agents such as anthracyclines or trastuzumab may lead to the development of HF and should be avoided, if possible
- **Thiazolidinediones (TZDs):** TZDs, including rosiglitazone and pioglitazone, are used in the management of patients with DM and have been found to cause edema, an adverse effect that is more common when a TZD is used in combination with other oral hypoglycemic agents or insulin. In addition, clinical trials with the TZDs generally did not include patients with NYHA class III or IV HF and an increased risk of HF in patients prescribed the TZDs has been reported, although the risk appears to be low. Current recommendations include evaluation of the patient's cardiac and fluid status prior to prescribing a TZD and upon follow-up. If a TZD is prescribed in patients without HF but who have one or more risk factors for HF,<sup>52</sup> or in patients who have NYHA class I or II HF, a low dose should be started and the patient should be closely monitored for signs and symptoms of HF including shortness of breath, edema, or excessive or rapid weight gain, as treatment with a TZD has been associated with worsening of HF, and a higher rate of hospitalization and cardiovascular related events. Treatment with a TZD should be reconsidered in patients who develop HF after initiation of the drug. Clinician discretion may be used in patients receiving a TZD who are stabilized and without evidence of volume overload. The use of a TZD is contraindicated in patients with NYHA Class III or IV HF.
- **Cilostazol:** Cilostazol, a phosphodiesterase (PDE) inhibitor used in the management of intermittent claudication and as antiplatelet therapy, is contraindicated in patients with HF due to the decreased survival seen in patients with NYHA class III or IV HF receiving other PDE type 3 inhibitors
- **Metformin:** Metformin should not be used in patients with unstable or acute congestive HF due to the propensity for hypoperfusion or hypoxemia and resultant increased risk of developing lactic acidosis

#### Additional recommendations<sup>22,53,54</sup>

- Patients and their families or caregivers should receive education on HF, dietary restrictions including reducing salt intake and fluid control (especially in advanced HF), weight monitoring to assess fluid status, moderation of alcohol intake, weight loss if obese, regular physical activity or exercise training if appropriate, smoking cessation, drug therapy and importance of adherence to the medication regimen, symptoms associated with worsening HF and what to do if they occur, and prognosis
- Unless contraindicated, influenza vaccination should be offered every fall
- Pneumococcal immunizations should be provided at diagnosis if not previously vaccinated; if initial vaccination was at age less than 65 years, revaccinate at age 65 or 5 years after initial immunization, whichever is later
- An implantable cardioverter-defibrillator or cardiac resynchronization therapy should be considered in appropriate patients but are beyond the scope of this document.
- Patients should be followed closely by a clinician competent in caring for patients with HF. Care of patients with HF may occur in several clinical settings including primary care, cardiology, or by multidisciplinary HF treatment teams. Regardless of the setting in which patients with HF are cared for, the clinician is encouraged to follow these and other HF guidelines and to use clinical judgment of when to refer to a specialist. This will depend on the skill and experience of managing patients with HF, and also the resources available to the practitioner

### **C. Pharmacologic Management of HF with Preserved Left Ventricular Ejection Fraction**

#### **OBJECTIVE**

- To review the pharmacologic recommendations for patients with HF and preserved LVEF



## BACKGROUND

In patients with HF and normal LVEF the systolic function of the left ventricle is preserved. The defect of ventricular function lies in the reduced LV compliance and difficulty in passive filling. Increased LVEDP can result in pulmonary congestion indistinguishable clinically from LV systolic dysfunction. As patients with HF and normal LVEF are often symptomatic, these patients may also be categorized as Stage C HF according to the ACC/AHA.<sup>1</sup> In addition, there is a high rate of morbidity and mortality seen in these patients.<sup>1,55</sup>

Compared to HF due to systolic dysfunction, there is a paucity of data from randomized trials about the pharmacologic management of patients with preserved LVEF,<sup>1,56-71</sup> despite the estimate that 20-60% of patients with HF can be considered to have normal LVEF (depending on the definition for reduced LVEF).<sup>1</sup> Since questions remain regarding the optimal treatment of patients with HF and normal LVEF, it is recommended that these patients be treated in conjunction with a cardiologist if the patient does not adequately respond to initial interventions.

## RECOMMENDATIONS

### Drug Therapy

The following are Class I recommendations by the ACC/AHA (i.e., there is evidence and/or general agreement that a given procedure/therapy is useful and effective)<sup>1</sup>

- Control HTN
- Control ventricular rate in patients with atrial fibrillation
- Use diuretics in patients with symptoms of volume overload (e.g., pulmonary congestion or peripheral edema)

The following are Class IIb recommendations by the ACC/AHA (i.e., the usefulness/efficacy is less well established by evidence/opinion)<sup>1</sup>

- Consider use of beta-adrenergic blockers, ACEIs, non-dihydropyridine calcium channel blockers, or angiotensin II receptor antagonists in patients with controlled HTN who continue to have symptoms
- Use of digoxin to improve symptoms is not well established
- Restoring and maintaining sinus rhythm in patients with atrial fibrillation may be useful to improve symptoms

## DISCUSSION

General principles of lowering systolic and diastolic blood pressure, treating myocardial ischemia, controlling heart rate and blood volume, and providing anticoagulation for patients with atrial fibrillation apply to these patients as well as to patients with systolic dysfunction. Conditions that can lead to HF with a normal LVEF (e.g., HTN, coronary artery disease, aortic stenosis) should also be treated.<sup>1</sup>

The main goal of therapy is to improve symptoms by lowering the filling pressures of the left ventricle without significantly reducing cardiac output. Agents that decrease heart rate can be helpful by increasing diastolic filling time.

The majority of clinical trials in patients with HF and preserved LVEF have been in a limited number of patients. The CHARM (Candesartan in Heart failure Assessment of Reduction in Mortality and morbidity) Preserved trial<sup>71</sup> enrolled 3023 patients with HF and LVEF > 40% and compared the addition of an angiotensin II receptor antagonist (candesartan) or placebo to current therapy. Median follow-up was 36.6 months. The primary endpoint of cardiovascular mortality or HF hospitalizations occurred in 22% of

patients receiving candesartan compared to 24% of patients in the placebo group; a difference that was not statistically significant.

## **D. Interventions in Patients With Asymptomatic Left Ventricular Systolic Dysfunction**

### **OBJECTIVE**

- To provide recommendations for patients with asymptomatic LV systolic dysfunction (Stage B)

### **BACKGROUND**

The management goals for patients with asymptomatic systolic dysfunction is to prevent the development of HF.<sup>1</sup> These recommendations are divided into the following patient groups.

### **RECOMMENDATIONS**

#### Drug Therapy

- Use an ACEI and beta-adrenergic blocker in patients with a recent or remote history of MI, regardless of LVEF
- Use an ACEI in patients with a reduced LVEF who do not have symptoms of HF
- A beta-adrenergic blocker is indicated in patients without a history of MI who have reduced LVEF and do not have symptoms of HF
- An angiotensin II receptor antagonist may be given to patients with a history of MI who have a reduced LVEF and do not have symptoms of HF if they are intolerant of ACEIs

### **DISCUSSION**

#### Patients with a Recent or Remote History of Myocardial Infarction

Prescribing an ACEI in patients with an acute or recent MI<sup>72,73</sup> and evidence of left ventricular systolic dysfunction (generally defined as LVEF < 40%) may reduce mortality and slow the progression to symptomatic heart failure.<sup>74-77</sup> In the Survival and Ventricular Enlargement (SAVE),<sup>74</sup> Acute Infarction Ramipril Efficacy (AIRE),<sup>75,76</sup> and Trandolapril Cardiac Evaluation (TRACE)<sup>77,78</sup> trials, patients with a recent MI and evidence of HF experienced a significant decrease in all-cause mortality and risk of developing severe HF when treated with an ACEI compared to placebo. Treatment with an ACEI in patients recently recovered from an MI can decrease the risk of reinfarction and death in patients with evidence of HF at the time of the infarction.<sup>75</sup> Patients with a history of MI without reduced LVEF may also benefit from treatment with an ACEI.<sup>16,79</sup>

The use of a beta-adrenergic blocker in patients with asymptomatic left ventricular systolic dysfunction post-MI reduces the risk of cardiovascular morbidity and mortality.<sup>80-83</sup> In the Carvedilol Post-Infarct Survival Control in LV Dysfunction (CAPRICORN) trial that randomized 1959 patients with a LVEF ≤ 40% post-MI to carvedilol or placebo, there was not a statistically significant difference in the primary endpoint of all-cause mortality or cardiovascular hospitalizations (originally a prespecified secondary endpoint). The original primary endpoint of all-cause mortality (changed to co-primary endpoint due to inadequate sample size and power) was lower, but not statistically significant in patients on carvedilol compared to placebo.<sup>84</sup> Although the results of this study did not achieve statistical significance, the endpoints were numerically lower in patients treated with carvedilol. Taking this into account with results of other trials, there still appears to be a benefit of using a beta-adrenergic blocker in patients with asymptomatic left ventricular systolic dysfunction post-MI.

Combination therapy with a beta-adrenergic blocker and an ACEI may also be beneficial in patients with left ventricular systolic dysfunction post-MI.<sup>85-87</sup>

In the Optimal Trial in Myocardial Infarction with the Angiotensin II Antagonist Losartan (OPTIMAAL), losartan (target dose 50mg once daily) was compared to captopril (target dose 50mg three times daily) in 5477 high-risk (i.e., signs and symptoms of HF or Q-wave MI) patients with acute MI. After a mean follow-up of 2.7 years, the primary endpoint of all-cause mortality occurred in 18% of patients on losartan and 16% of patients on captopril, a difference that was not statistically significant.<sup>88</sup> In another trial comparing an angiotensin II receptor antagonist with an ACEI in patients with MI complicated by HF, LV dysfunction, or both, the Valsartan in Acute Myocardial Infarction Trial (VALIANT) randomized 14,703 patients to treatment with valsartan (target dose 160mg twice daily), captopril (target dose 50mg three times daily), or the combination of valsartan (target dose 80mg twice daily) and captopril (target dose 50mg three times daily). During a median follow-up of 24.7 months, the primary endpoint of all-cause mortality was similar in the treatment groups and occurred in 19.9% of patients randomized to valsartan, 19.5% treated with captopril, and 19.3% receiving the combination. Treatment with valsartan was found to be noninferior to captopril for the endpoint of all-cause mortality.<sup>89</sup> Based on these results, an angiotensin II receptor antagonist can be considered in patients who are intolerant to an ACEI (refer to Annotation G) in high risk patients following MI.<sup>1</sup>

#### Patients with Asymptomatic Left Ventricular Dysfunction

In the Studies of Left Ventricular Dysfunction (SOLVD) Prevention trial, patients with asymptomatic left ventricular dysfunction treated with an ACEI experienced a significant reduction in the combined risk of death and hospitalization for HF by 20% compared to placebo. However, there was no significant decrease in all-cause mortality alone in the ACEI group.<sup>90</sup> In the Prevention trial component of the 12 year follow-up of SOLVD (median duration of follow-up for the Prevention trial 11.2 months), there was a significant reduction in all-cause mortality in patients treated with enalapril (median duration during trial 3.2 years) compared to those receiving placebo (50.9% vs. 56.4%).<sup>91</sup>

The benefit of an ACEI in men compared to women with HF was recently evaluated. According to a subgroup analysis of trials including treatment of patients with asymptomatic LV dysfunction, there did not appear to be a clear benefit of ACEI in women, with a relative risk of 0.96 (95% CI 0.75-1.22). It was concluded that further investigation is warranted before making a definitive recommendation on the use of ACEIs in women with asymptomatic left ventricular dysfunction.<sup>92</sup> While the benefit, to the extent that one exists, remains to be quantified, an ACEI should still be considered standard therapy given the current level of data overall.

Although the benefit of beta-adrenergic blockers in patients with asymptomatic HF (not in the post-MI setting) has not been critically evaluated, current recommendations include use of a beta-adrenergic blocker in this patient population.<sup>1,83-85</sup>

Digoxin is currently recommended in patients with symptomatic HF to improve clinical status and decrease the risk of hospitalization due to HF,<sup>93</sup> after optimization of standard therapy. Since there is not a significant reduction in disease progression or mortality, digoxin is not recommended in patients with asymptomatic left ventricular dysfunction.<sup>1</sup>

#### **EVIDENCE**

Grading System		USPSTF <sup>b</sup>			ACC/AHA <sup>1</sup>	
Intervention	References	QE	OQ	SR	CR	LE

Use an ACEI in patients with a recent or remote history of MI, regardless of LVEF	SAVE (1992) AIRE (1993) GISSI-3 (1994) ISIS-4 (1995) TRACE (1995) AIREX (1997) HOPE (2000) TRACE 12yr (2005) PREAMI (2006) ACC/AHA (2005) HFSA (2006)	I I I I I II-1 I II-1 I III III	I I A	I A
Use a beta-adrenergic blocker in patients with a recent or remote history of MI, regardless of LVEF	Norwegian trial (1981) BHAT (1982) Chadda (1986) Gottlieb (1998) CAPRICORN (2001) ACC/AHA (2005) HFSA (2006)	I I II-2 II-2 I III III	I I A	I A
Use an ACEI in patients with a reduced LVEF who do not have symptoms of HF	SOLVD (1992) SOLVD 12yr (2003) ACC/AHA (2005) HFSA (2006)	I II-1 III III	I I A	I A
A beta-adrenergic blocker is indicated in patients without a history of MI who have reduced LVEF and do not have symptoms of HF	Chadda (1986) CAPRICORN (2001) SAVE (1997) ACC/AHA (2005) HFSA (2006)	II-2 I II-2 III III	II B	I C
An angiotensin II receptor antagonist may be given to patients with a history of MI who have a reduced LVEF and do not have symptoms of HF if they are intolerant of ACEIs	OPTIMAAL (2002) VALIANT (2003) ACC/AHA (2005)	I I III	I I A	I B

## E. Systolic Dysfunction and Assessment for Symptoms of Volume Overload

### OBJECTIVE

- To provide recommendations for initial assessment and therapy in patients with a diagnosis of systolic HF who exhibit symptoms of volume overload

### BACKGROUND

The goals of treating patients with HF due to systolic dysfunction are to improve the patient's symptoms and quality of life, and to reduce the risk of morbidity and mortality by slowing the progression of disease. Patient's symptoms are often related to volume overload.

### RECOMMENDATION

#### Initial Assessment of Volume Status

The following is a Class I recommendation by the ACC/AHA (i.e., there is evidence and/or general agreement that a given procedure/therapy is useful and effective)<sup>1</sup>

- Patients presenting with HF should receive an assessment of volume status

### DISCUSSION

Symptoms of volume overload include ankle swelling, weight gain, fatigue, orthopnea, PND, DOE, SOB at rest and nocturnal cough. The physical signs of volume overload are pulmonary crackles, third heart sound, cardiomegaly, JVD, hepatjugular reflux, hepatomegaly, ascites, dependent edema (presacral, flank, lower extremity), tachypnea, tachycardia, and pulmonary edema.

Chest radiography is useful to identify signs of volume overload (pleural effusion, pulmonary edema, cardiomegaly).

A diuretic is recommended in patients with HF who exhibit signs or symptoms of volume overload (refer to Annotation F).<sup>1</sup>

## **F. Diuretic Therapy**

### **OBJECTIVE**

- To provide recommendations for the appropriate use of diuretics in patients with a diagnosis of systolic HF (for a discussion on the use of aldosterone antagonists in HF, refer to Annotation L)

### **BACKGROUND**

Many patients with HF will present with symptoms of fluid retention and require treatment with a diuretic. It is recommended that the diuretic should be continued (particularly in patients with NYHA III or IV failure) even if symptoms resolve to prevent recurrence of volume overload. Patients should notice symptomatic improvement early on with diuretic therapy; however, a diuretic should not be prescribed alone in patients with Stage C HF but combined with drug classes that have been shown to reduce morbidity and mortality.<sup>1</sup>

### **RECOMMENDATION**

#### Diuretic Therapy in Stage C HF

- A diuretic is indicated in patients with current or previous symptoms of HF with evidence of fluid retention

### **DISCUSSION**

Diuretics act by inhibiting sodium or chloride reabsorption in the renal tubules. The loop diuretics exert their effects more proximally and are therefore the most potent of the diuretics. The diuretics primarily differ in their duration of action (e.g., furosemide 6-8 hours, hydrochlorothiazide 6-12 hours, metolazone 12-24 hours) and in their ability to cause sodium excretion ('low ceiling' diuretics like hydrochlorothiazide or 'high ceiling' diuretics like furosemide). As HF progresses, a delay in absorption and failure to filter the drug in the tubular fluid may be contributing factors to the need for increasing diuretic doses in some patients.<sup>1,94-96</sup>

There have been no long-term properly blinded, randomized controlled clinical outcome trials evaluating the effectiveness of loop or thiazide diuretic therapy in patients with HF.<sup>1</sup> Short-term and intermediate length studies have demonstrated that diuretics can decrease the signs and symptoms of fluid retention, and improve cardiac conduction and exercise tolerance.<sup>1,97-100</sup> The majority of patients enrolled in long-term trials demonstrating a decreased morbidity or mortality with ACEI or beta-adrenergic blocker therapy, were also receiving a diuretic.

Some patients with HF may experience a recurrence of symptoms if diuretic therapy is withdrawn.<sup>101</sup> In one trial the risk of requiring reinstitution of diuretic therapy was 36% in patients in the withdrawal group compared with controls.<sup>102</sup> A LVEF  $\leq$  27%, diuretic dose greater than 40mg of furosemide daily, or a history of HTN were independent risk factors for early reinstitution of diuretic therapy.<sup>103</sup>

Patients with HF may have symptoms that interfere with their daily activities and, therefore, impact on their quality of life. A diuretic should be used for preload reduction in patients with HF and current or previous signs or symptoms of volume overload (e.g., orthopnea, PND, DOE, or edema).<sup>1,100</sup> Patients with

symptoms of fluid overload benefit from treatment with a diuretic in conjunction with an ACEI and beta-adrenergic blocker,<sup>1</sup> and possibly digoxin.<sup>104</sup>

Loop diuretics are most commonly used for patients with HF and volume overload. They are effective in patients with renal insufficiency or creatinine clearance (CrCl) < 30 mL/min, whereas the effectiveness of thiazides are diminished in patients with CrCl < 40 mL/min.<sup>1</sup> Edema resistant to large doses of loop diuretics may intermittently require combined diuretic therapy (e.g., adding metolazone or thiazide at low doses two to three times per week or more frequently if needed, one hour prior to a loop diuretic), consideration of change to another loop diuretic, or intravenous diuretics.<sup>1, 94-96,104-111</sup> The use of combination diuretics increases the risk of electrolyte imbalances and overdiuresis leading to prerenal azotemia. Therefore, combination diuretic therapy requires close monitoring.

Monitoring parameters with diuretics include the following:<sup>1</sup>

1. Weight: (initially 1 - 2 pound weight loss per day until “ideal weight” achieved); weight loss may be greater during the first few days when significant edema is present; obtain daily weights
2. Signs or symptoms of volume depletion: weakness, dizziness, decreased urine output, symptomatic hypotension, orthostatic hypotension
3. Serum potassium (K<sup>+</sup>), BUN or Cr (and serum BUN/Cr ratio) within 1 to 2 weeks after initiating therapy; consider checking serum levels of magnesium (especially if high doses diuretic used), sodium, calcium, bicarbonate, uric acid, glucose as indicated. Use of an ACEI (or angiotensin II receptor antagonist) and/or spironolactone may offset potential diuretic-induced hypokalemia, minimizing the need for potassium or potassium-sparing diuretics
4. Diuretic dosage may require adjustment if hypotension or decrease in renal function occurs. Avoid excessive diuresis, which could also limit ACEI dosage due to hypotension or renal dysfunction

## EVIDENCE

Grading System Intervention	References	USPSTF <sup>6</sup>			ACC/AHA <sup>1</sup>	
		QE	OQ	SR	CR	LE
Use a diuretic in patients with evidence of fluid retention	Wilson (1981) Richardson (1987) Parker (1993) Patterson (1994) ACC/AHA (2005) HFSA (2006)	I I I I III III	II	B	I	C

## G. Angiotensin-Converting Enzyme Inhibitors

### OBJECTIVE

- To provide recommendations for the appropriate use of ACEIs in patients with a diagnosis of systolic HF

### BACKGROUND

Due to their beneficial effects on morbidity and mortality, an ACEI should be prescribed in patients with Stage C HF, unless contraindicated. Therapy should be initiated at low doses and titrated to target doses or patient tolerability.

### RECOMMENDATION

#### ACEI Therapy in Stage C HF

- An ACEI is recommended in all patients with current or prior symptoms of HF and reduced LVEF, unless contraindicated

## DISCUSSION

Angiotensin-converting enzyme (ACE) is responsible for converting angiotensin I to angiotensin II. Angiotensin II is a potent vasoconstrictor and it stimulates aldosterone secretion, which leads to increased sodium and water retention. By inhibiting this enzyme, ACEIs ultimately reduce the vasoconstriction associated with angiotensin II and decrease the sodium and water retention associated with aldosterone. ACE is structurally similar to kininase II, so it may also inhibit the breakdown of bradykinin, a vasodilator. The importance of ACE's effect on kinin-mediated prostaglandin synthesis in the management of patients with HF is not yet known, but it may be as important as angiotensin suppression.<sup>1</sup>

In addition to improving HF symptoms and functional status,<sup>1,112-118</sup> treatment with an ACEI has been shown to decrease the frequency of hospitalization and the mortality rate<sup>91,119-123</sup> (Appendix C).

In the Captopril-Digoxin Multicenter Trial, patients with mild to moderate HF were randomized to placebo, captopril, or digoxin in addition to treatment with diuretics for 6 months. Compared with placebo, patients on captopril experienced significant improvement in exercise tolerance and decreased frequency of hospital or emergency care for worsening HF. Similar results were not seen with digoxin.<sup>119</sup> Patients with mild to moderate HF who received enalapril for an average of 41 months in the SOLVD Treatment Trial had a significant decrease of 16% in all-cause mortality and a 26% decreased risk of death or hospitalizations for HF compared to patients on placebo.<sup>120</sup> The Vasodilator Heart Failure Trial (V-HeFT) II showed that patients with mild to moderate HF who received enalapril for an average of 2.5 years experienced a significant decrease of 28% in the risk of death at 2 years compared to patients on the combination of isosorbide dinitrate (ISDN) and hydralazine.<sup>121</sup> The Cooperative North Scandinavian Enalapril Survival Study (CONSENSUS) evaluated treatment with enalapril for 6 months compared to placebo in patients with NYHA class IV HF. Treatment with enalapril significantly decreased all-cause mortality at 6 months by 40%.<sup>122</sup>

The possibility of racial differences in response to therapy has been seen in a subanalysis of V-HeFT and V-HeFT II. In V-HeFT I, white patients did not experience the same mortality benefit as black patients on ISDN and hydralazine (compared to placebo). In V-HeFT II, white patients on an ACEI experienced a decrease in mortality compared to treatment with ISDN and hydralazine. There was not a statistically significant difference in mortality between treatments in black patients.<sup>124</sup> When matched cohorts of white patients were compared to black patients on an ACEI enrolled in the SOLVD Treatment Trial, white patients experienced a decreased risk for hospitalizations due to HF which was not seen in the cohort of black patients.<sup>125</sup> Based on a pooled relative risk analysis, there was no evidence that mortality differed substantially with an estimate for white patients of 0.89 (95% CI 0.82-0.97) and 0.89 (85% CI 0.74-1.06) for black patients.<sup>92</sup> Results of a trial comparing treatment with the combination of ISDN and hydralazine vs. placebo in self-identified black patients (the majority being treated with an ACEI or angiotensin II receptor antagonist and a beta-adrenergic blocker) are discussed in Annotation J. Further trials will need to be conducted to determine if therapy with an ACEI for HF should be modified based on patient demographics.

It is recommended that an ACEI be offered to all patients with reduced left ventricular systolic dysfunction unless the patient has specific contraindications:<sup>1,126</sup>

1. A history of angioedema, anuric renal failure, or other documented hypersensitivity to an ACEI
2. Bilateral renal artery stenosis or renal artery stenosis in a solitary kidney
3. Pregnancy
4. Serum potassium > 5.5 mEq/L that cannot be reduced
5. Hypotension in patients at risk of cardiogenic shock

Prior to initiating ACEIs, obtain baseline serum potassium, Cr, and BUN; ACEIs should be used cautiously in patients with serum Cr > 3mg/dL. Before initiating therapy, patients should first be assessed for adequate volume status. In patients taking diuretics, symptomatic hypotension may occur following



initiation of an ACEI; if the diuretic cannot be reduced or discontinued, consider a lower starting dose of an ACEI. If the patient is on a potassium-sparing diuretic when an ACEI is begun, close monitoring of potassium is recommended. Alternatively, if the patient is hypokalemic or normokalemic, the potassium-sparing diuretic may be stopped while titrating the ACEI and re-started later, with subsequent close monitoring of potassium. An ACEI should also be used with caution with an aldosterone antagonist (refer to Annotation L). Concomitant use with an NSAID should be avoided whenever possible as NSAIDs used in conjunction with an ACEI may worsen renal function and contribute to hyperkalemia (refer to Appendix B for common drug interactions). Patients started on an ACEI should be evaluated within 1 to 2 weeks to monitor blood pressure, serum potassium and creatinine; more frequent monitoring may be warranted depending on the severity of the patient's condition.

Doses should initially be low and then titrated upward over several weeks to the maximum dose tolerated, with the target doses based on those used in large scale clinical trials (refer to Appendices A and C).<sup>1</sup> Despite the overwhelming evidence in favor of treating HF patients with ACEIs and that a large majority of patients are able to tolerate high doses, these agents are often underutilized, and frequently at low doses,<sup>127-129</sup> although this may depend on the clinical setting.<sup>130</sup>

There appears to be a dose response benefit as shown in the Assessment of Treatment with Lisinopril and Survival (ATLAS) study. In this study, patients with NYHA class II-IV HF on maximal doses of lisinopril (average of  $33.2 \pm 5.4$  mg daily) experienced a significant 12% decrease in the risk of death or hospitalization for any reason and 24% fewer hospitalizations for HF, compared to patients receiving lower doses (average of  $4.5 \pm 1.1$  mg daily). There was also a nonsignificant 8% lower risk of death in the high dose compared to the low dose treatment group. The authors observed that the decrease in risk with the high dose compared to the low dose group in the ATLAS study was approximately half that seen with target doses of an ACEI compared to placebo in other trials. This suggests that even patients on suboptimal doses will derive benefit, although not as great as patients receiving higher doses.<sup>131,132</sup> This is important since other factors may preclude a patient from achieving target doses. In another trial, patients on high doses of an ACEI (enalapril 20mg/d) had a decreased risk of HF hospitalizations compared to patients on medium and lower doses (enalapril 10mg/d and 5mg/d, respectively). There was no difference between doses in symptoms or mortality.<sup>133</sup> There was also no difference in NYHA class, LVEF, or mortality in a trial of patients on standard ( $17.9 \pm 4.3$  mg/d) compared to high ( $42 \pm 19.3$  mg/d) doses of enalapril.<sup>134</sup>

Every effort should be made to titrate patients to the doses used in clinical trials, although if this is not feasible, patients should be maintained on the maximum tolerated dose.<sup>1</sup> Patients prescribed an ACEI prior to discharge from the hospital are more likely to be on an ACEI long-term compared to those not discharged on an ACEI.<sup>135</sup> In addition, initiation of a beta-adrenergic blocker should not be delayed due to an inability to achieve target doses of an ACEI,<sup>1</sup> as patients treated with a beta-adrenergic blocker derived benefit regardless if the patient were receiving low or high doses of an ACEI.<sup>136</sup>

Due to the strong evidence for the beneficial effects of ACEIs in patients with HF, every effort should be made to adjust the dosage before a patient is documented as intolerant.<sup>137</sup> Dosage should be modified if the patient develops any of the following:<sup>128</sup>

1. While creatinine often increases (usually < 25%) after initiation of an ACEI, clinically significant decline in renal function (suggested by a change in serum Cr concentration of at least 0.5 mg/dL) should be investigated. Consultation with a nephrologist should be considered for persistent deteriorations in renal function that cannot be explained or corrected.
2. Hyperkalemia (potassium > 5.5 mEq/L), after other causes have been excluded
3. If patient cannot tolerate ACEI due to symptomatic hypotension, consider referral to a cardiologist for assistance in titrating the ACEI dosage
4. The cough associated with an ACEI has been described as dry, nonproductive, persistent, beginning with a tickling sensation, and often worse at night. The onset is usually within the first week of ACEI therapy and continues throughout treatment, resolving within a few days to 4 weeks after the ACEI is discontinued. The cough is not usually dose-dependent, although in some instances it may be eliminated with a reduction in dose. In addition, fosinopril may be considered in patients who



experience cough on another ACEI.<sup>138-140</sup> Since therapy with an ACEI has proven valuable, it is important to consider alternative diagnoses (e.g., asthma, chronic obstructive pulmonary disease, allergic rhinitis, upper respiratory tract infection, heart failure, gastroesophageal reflux disease) before a diagnosis of ACEI-induced cough is made. If the cough is not bothersome, the benefits of continuing the ACEI should be discussed with the patient.

There is some controversy as to whether use of aspirin decreases the cardiovascular benefit of an ACEI when used concomitantly. Some of the beneficial effects of ACEIs are thought to be due to inhibiting the breakdown of bradykinin, which in turn, increases the production of vasodilatory prostaglandins. Aspirin, which blocks cyclooxygenase, may therefore interfere with the full benefit of an ACEI by inhibiting vasodilatory prostaglandin synthesis.<sup>141,142</sup> Much of the discussion was prompted from the publication of retrospective analyses of data from large trials evaluating the benefits of treatment with an ACEI.<sup>24,143</sup> A cohort analysis of SOLVD found that treatment with an antiplatelet agent (e.g., aspirin or dipyridamole) was associated with a reduction in all-cause mortality and a decrease in the risk of death or hospital admission for HF. In contrast, this association was not apparent in patients treated with an ACEI who were on an antiplatelet agent at baseline, and patients on an ACEI did not experience a reduction in all-cause mortality as did patients randomized to enalapril who were not on an antiplatelet agent. There was a reduction in the combined risk of death or hospital admission for HF in patients on an ACEI and antiplatelet agent.<sup>24</sup> In an analysis of CONSENSUS II in patients with acute MI, those in the ACEI treatment group who were taking aspirin at baseline experienced a lower mortality benefit than patients who were on an ACEI without aspirin.<sup>143</sup> In a retrospective analysis of over 22,000 patients from six long-term randomized controlled trials, treatment with an ACEI decreased the risk of major clinical outcomes (composite death, MI, stroke, HF hospitalization, or revascularization) by 20% in patients also receiving aspirin, and by 29% in patients not on concomitant aspirin therapy (interaction test not statistically significant).<sup>144</sup> Two additional evaluations of patients prescribed an ACEI in conjunction with aspirin compared to an ACEI without aspirin, did not find an association between outcome and concomitant aspirin use.<sup>145,146</sup> A dose-related adverse effect of aspirin was reported in one retrospective evaluation where there was an increase in mortality in the patients receiving high dose aspirin ( $\geq 325\text{mg}$ ) compared to those on low dose or no aspirin.<sup>147</sup> It is difficult to determine the clinical significance of these results given the retrospective nature of the analyses and the potential contribution of differences in the groups at baseline.<sup>24,141-147</sup> A prospective evaluation of patients with systolic HF receiving warfarin, aspirin, or clopidogrel reported no significant difference in the primary outcome of death, MI, or stroke; although there was a significant reduction in HF hospitalizations with warfarin compared to aspirin.<sup>148,149</sup> This trial was terminated early due to poor enrollment.<sup>148</sup> Therefore, given the benefit of aspirin in patients with coronary artery disease, there is insufficient evidence to warrant a change in the current recommendations in patients with coronary artery disease and HF.

## EVIDENCE

Grading System		USPSTF <sup>6</sup>			ACC/AHA <sup>1</sup>	
Intervention	References	QE	OQ	SR	CR	LE
An ACEI is recommended in all patients with Stage C HF, unless contraindicated	CONSENSUS (1987)	I	I	A	I	C
	SOLVD (1991)	I				
	V-HeFT II (1991)	I				
	SOLVD 12yr (2003)	II-1				
	ACC/AHA (2005)	III				
	HFSA (2006)	III				

## H. Beta-Adrenergic Blockers

### OBJECTIVE

- To provide recommendations for the appropriate use of beta-adrenergic blockers in patients with a diagnosis of systolic HF

## BACKGROUND

Due to their beneficial effects on morbidity and mortality, a beta-adrenergic blocker should be used in patients with Stage C HF, unless contraindicated. Therapy should be initiated at low doses and titrated to target doses or based on patient tolerability. The majority of clinical trials evaluating efficacy of a beta-adrenergic blocker in patients with HF were conducted in patients receiving an ACEI and a diuretic. Patient factors may be taken into consideration when determining whether to initiate therapy first with an ACEI or beta-adrenergic blocker; with subsequent addition of the alternate drug class.

## RECOMMENDATION

### Beta-Adrenergic Blocker Therapy in Stage C HF

- Stable patients with current or prior symptoms of HF due to systolic dysfunction should receive therapy with a beta-adrenergic blocker that has proven to reduce mortality (i.e., bisoprolol, carvedilol, sustained release metoprolol succinate) unless contraindicated

## DISCUSSION

Activation of the sympathetic nervous system (SNS) is one of the proposed compensatory mechanisms to maintain circulation in the presence of left ventricular dysfunction. However, activation of the SNS can result in beta-receptor down-regulation, LVH, cardiotoxic effects, and arrhythmia. It is thought that one or more of these effects may contribute to HF progression.<sup>1,150</sup> Therefore, using a beta-adrenergic blocker in a patient with HF due to systolic dysfunction could potentially negate some of these adverse effects on the heart.

Numerous trials have been conducted that demonstrate the beneficial effects of beta-adrenergic blockers in reducing symptoms, hospitalization, and progression of disease in patients with HF due to systolic dysfunction.<sup>1,151-165</sup> However, more recent evidence has demonstrated a mortality benefit with the use of beta-adrenergic blockers in this patient population<sup>159-167</sup> (Appendix C). The beta-adrenergic blockers that have been studied for chronic HF and have demonstrated a reduction in mortality include bisoprolol, carvedilol, and sustained release metoprolol succinate. Bisoprolol, titrated to 10 mg once daily, was compared to placebo in patients with primarily NYHA class III HF receiving standard therapy in the second Cardiac Insufficiency Bisoprolol Study (CIBIS II). The primary endpoint of all-cause mortality was reduced with bisoprolol, occurring in 11.8% of patients, compared to 17.3% of patients on placebo.<sup>160</sup> Carvedilol was studied in patients with NYHA class II and III HF (U.S. Carvedilol Heart Failure Study),<sup>161</sup> as well as in patients with more severe HF as in the Carvedilol Prospective Randomized Cumulative Survival Study (COPERNICUS).<sup>162</sup> After a median of 6.5 months, the primary endpoint of death was reported in 3.2% of patients in the U.S. Carvedilol Study receiving carvedilol (target dose 25 mg twice daily) compared to 7.8% of patients on placebo.<sup>161</sup> In COPERNICUS, the primary endpoint of all-cause mortality occurred in 11.3% of patients randomized to carvedilol compared to 16.8% of patients receiving placebo.<sup>162</sup> In the Carvedilol Or Metoprolol European Trial (COMET), carvedilol at a target dose of 25 mg twice daily was compared to the immediate-release formulation of metoprolol tartrate, at target doses of 50 mg twice daily. All-cause mortality was reported to be lower in patients on carvedilol (33.9%) compared to patients receiving metoprolol (39.5%) in this study (additional discussion below).<sup>167</sup> All-cause mortality was also a primary endpoint in the Metoprolol CR/XL Randomized Intervention Trial in Congestive Heart Failure (MERIT-HF), and was reported in 7.3% of patients randomized to the extended-release formulation of metoprolol succinate (metoprolol XL at a target dose 200mg once daily) compared to 10.9% of patients receiving placebo.<sup>159</sup>

It is unknown if other beta-adrenergic blockers have a similar benefit, as not all beta-adrenergic blockers studied have shown a clear reduction in mortality.<sup>151,167,168</sup>

These agents have also demonstrated efficacy in patients with advanced HF.<sup>160,162,169</sup> In a subgroup analysis of MERIT-HF, 795 patients with NYHA class III or IV HF with a LVEF < 25% who received placebo or metoprolol XL were compared. Similar to COPERNICUS with carvedilol,<sup>162</sup> the mean baseline LVEF was 19.1% and the annual mortality for patients in the placebo group was 19%. Patients randomized to metoprolol XL experienced a significant decrease in risk of total mortality (39%), death due to worsening HF (55%), hospitalization due to worsening HF (45%), and combined all-cause mortality or all-cause hospitalization (29%) compared to placebo.<sup>169</sup>

In another post hoc analysis of MERIT-HF, the beneficial effects on morbidity and mortality with metoprolol XL were also seen in the subgroup of 898 women, including 183 women with stable severe HF.<sup>170</sup>

The difference in response to treatment with a beta-adrenergic blocker based on race has also been evaluated. After subgroup analysis in the Beta-Blocker Evaluation of Survival Trial (BEST), there was a significant survival benefit in nonblack patients but not in black patients.<sup>168</sup> These results are contrary to findings from a retrospective comparison of patients enrolled in the U.S. Carvedilol Study where the benefit of carvedilol was not statistically significantly different between black and nonblack patients.<sup>171</sup> A meta-analysis by the U.S. Department of Health and Human Services reported the estimate of pooled random-effects of the relative risk for mortality in black patients to be 0.67 (95% CI 0.39-1.16) compared to 0.63 (95% CI 0.52-0.77) for white patients. Results were similar for the pooled estimates from the hazard ratio analysis. The evidence report to address the potential difference in mortality of beta-adrenergic blockers depending on race concluded that black patients should derive the same benefits as white patients when treated with bisoprolol, carvedilol, or metoprolol succinate (the results of BEST were not included in the pooled analysis).<sup>92</sup>

The question of whether to use a selective beta-adrenergic blocker (e.g., bisoprolol or metoprolol) versus a non-selective agent with alpha-adrenergic blocking and antioxidant effects (e.g., carvedilol) remains controversial.<sup>167,172-174</sup> Although COMET demonstrated a statistically significant improvement in survival with carvedilol compared to immediate-release metoprolol (tartrate), it is unknown whether there is a difference between carvedilol and immediate-release metoprolol tartrate or metoprolol succinate (metoprolol XL) when prescribed at the recommended target doses. Since metoprolol succinate was not available at the time of enrollment in COMET, immediate-release metoprolol tartrate was selected as the comparator to carvedilol, at doses that were expected to result in comparable beta-blockade. Much of the discussion about the results of COMET includes the difference in target dose and effect on resting heart rate.<sup>175</sup> The dose of carvedilol used in COMET achieved a similar reduction in heart rate as seen in U.S. Carvedilol (i.e., 13 beats per minute).<sup>161,167</sup> The mean dose of metoprolol tartrate used in COMET was less than the mean dose in the Metoprolol in Dilated Cardiomyopathy (MDC) trial (i.e., 85 vs. 108mg/d), and resulted in less of a decrease in heart rate (i.e., 11.7 vs. 15 beats per minute).<sup>151,167</sup> The mean dose in MERIT-HF was 159mg/d and achieved a reduction in heart rate of 14 beats per minute.<sup>159</sup> Whether these factors had an influence on the results is unknown. Very few trials with beta-adrenergic blockers that are available in the U.S. other than bisoprolol, carvedilol, or metoprolol succinate have been published. It is therefore unknown if treatment with other beta-adrenergic blockers would provide the same benefits as seen with the agents that have demonstrated a reduction in mortality in patients with heart failure.<sup>176,177</sup>

The majority of patients included in the beta-adrenergic blocker trials received therapy with an ACEI. Survival benefit in the ACEI trials ranged from 12 to 33%, which was mainly a result of reduction in deaths from worsening HF. Meta-analyses of the beta-adrenergic blocker trials show a reduction in mortality of approximately 30 to 35%.<sup>85,177-179</sup> It is felt that the use of an ACEI and beta-adrenergic blocker in patients with HF is synergistic<sup>180</sup> and should be used in combination whenever possible.<sup>1</sup> Whether to begin treatment naïve patients with a beta-adrenergic blocker or an ACEI has yet to be resolved; according to the results from CIBIS III, it appears that initial therapy with bisoprolol may be as safe and efficacious as starting treatment with enalapril (refer to Appendix C for detailed results).<sup>166</sup> In patients with HF, utilization of the beta-adrenergic blockers is typically not as high as that seen with the ACEIs,<sup>4</sup> even though patient tolerability appears to be similar.<sup>181</sup> As the majority of clinical trials evaluating efficacy of a beta-adrenergic blocker in patients with HF were conducted in patients receiving an ACEI and a diuretic,

clinicians may choose to initiate therapy with a beta-adrenergic blocker once the patient has been stabilized on treatment with an ACEI.<sup>54</sup> Initiation of therapy with a beta-adrenergic blocker may be considered prior to achieving a target dose of the ACEI, with concomitant titration;<sup>53</sup> as benefit with combination therapy, even at lower doses of an ACEI, has been demonstrated.<sup>1,131,133</sup> Every effort should be made to achieve target doses of both an ACEI and beta-adrenergic blocker as tolerated by the patient. Implementation of treatment guideline recommendations should be emphasized in order to provide patients with the opportunity for optimal drug therapy benefit.<sup>4</sup>

Caution should be exercised when initiating a beta-adrenergic blocker in patients with HF. Initial dosages should be low and titrated upward slowly and as tolerated. Patients can become transiently worse with each dosage increase. Since patients may experience fluid retention during initiation, daily weights are recommended with corresponding adjustments in diuretic dose. Some patients may also experience fatigue or weakness that may resolve after several weeks or require dosage adjustments. Selection of a different beta-adrenergic blocker may also be considered.<sup>182</sup> Another factor that may contribute to a need for a delay in titration is a low heart rate;<sup>79</sup> although, the absolute increase in risk for hypotension, dizziness, and bradycardia is small and should be weighed against the overall benefit of beta-adrenergic blockers seen in clinical trials.<sup>182</sup> Clinicians who do not have experience with beta-adrenergic blockers in patients with HF should consult with a cardiologist or healthcare provider specializing in the management of HF. Another opportunity to initiate therapy may be predischARGE, provided the patient is stable and their condition does not necessitate use of intravenous therapy for HF.<sup>1,183</sup> It is important that patients with HF on a beta-adrenergic blocker are titrated carefully to a target dose as used in clinical trials or as tolerated (refer to Appendices A and C).<sup>184-191</sup> Common drug interactions are listed in Appendix B.

Factors that appear to contribute to a beneficial response are selection of patients who are clinically stable (i.e., not hospitalized in intensive care, no or minimal evidence of volume overload or depletion, no recent treatment with intravenous positive inotropic agents) when therapy starts, a low initial dosage, a gradual increase in the dosage (2 week intervals; with optimal doses achieved in 8 to 12 weeks<sup>53</sup>), and an adequate duration of treatment (3-12 months before effects are seen).

Beta-adrenergic blockers should not be used in patients with bronchospastic disease, symptomatic bradycardia, or advanced heart block without a pacemaker. Caution should be used in patients with asymptomatic bradycardia with a HR of less than 60 beats per minute.<sup>1</sup> If the patient is on digoxin with a HR of less than 60 bpm, reconsider digoxin in favor of the benefits of a beta-adrenergic blocker, or consider referral to a cardiologist for adjustment in therapy. It should be noted that patients with DM or chronic obstructive pulmonary disease were not excluded from the clinical trials.<sup>1,159-161,192</sup>

## EVIDENCE

Grading System Intervention	References	USPSTF <sup>6</sup>			ACC/AHA <sup>1</sup>	
		QE	OQ	SR	CR	LE
Use a beta-adrenergic blocker with proven mortality benefit in patients with stable Stage C HF	US Carvedilol (1996) MERIT-HF (1999) CIBIS-II (1999) COPERNICUS (2001) Shibata (2001) CIBIS I & II (2002) COMET (2003) ACC/AHA (2005) HFSA (2006)	I I I I I I I III III	I         	A         	I         	A         

## I. Angiotensin II Receptor Antagonists

### OBJECTIVE

- To provide recommendations for the appropriate use of the angiotensin II receptor antagonists in patients with a diagnosis of systolic HF

## BACKGROUND

Due to the established beneficial effects of the ACEIs and beta-adrenergic blockers in treating patients with HF, long-term outcome trials with the angiotensin II antagonists have been conducted in patients already receiving standard therapy for HF or in patients who are unable to tolerate an ACEI. Treatment with an angiotensin II receptor antagonist has shown a beneficial effect in reducing cardiovascular death and HF hospitalizations in patients unable to tolerate an ACEI, as well as in addition to standard therapy; although data are conflicting as to the benefit of adding an angiotensin II receptor antagonist to patients receiving an ACEI and a beta-adrenergic blocker. The effect of treatment with an angiotensin II receptor antagonist on all-cause mortality has not yet been established.

## RECOMMENDATIONS

### Angiotensin II Receptor Antagonist in Stage C HF

- An angiotensin II receptor antagonist with demonstrated efficacy (i.e., candesartan and valsartan) in the treatment of HF is recommended in patients with Stage C HF who are unable to tolerate therapy with an ACEI
- Addition of an angiotensin II receptor antagonist to standard therapy (i.e., an ACEI and beta-adrenergic blocker) may be considered to decrease cardiovascular death or HF hospitalizations in patients with persistent symptoms (see discussion below); although it should also be noted that routine use of an ACEI, angiotensin II receptor antagonist, and aldosterone antagonist is not recommended

## DISCUSSION

Angiotensin-converting enzyme inhibitors reduce levels of angiotensin II, a potent vasoconstrictor, and inhibit the breakdown of bradykinin, a vasodilator. Production of angiotensin II also occurs through alternative pathways. The angiotensin II receptor antagonists selectively block the angiotensin II type 1 receptor so that the effects of angiotensin II are blocked regardless of how it is produced. The contribution of bradykinin to the favorable results of the ACEI trials in HF patients is unknown, but may be as important as suppression of angiotensin.<sup>1</sup>

Trials have been conducted evaluating the majority of the angiotensin II receptor antagonists, demonstrating a favorable effect on patient symptoms or NYHA functional class compared to placebo,<sup>191-195</sup> with comparable benefits to an ACEI.<sup>196-201</sup> Results of long-term effects on morbidity and mortality have been published in patients treated with the angiotensin II receptor antagonists, candesartan, losartan, and valsartan. The results of these trials are briefly discussed with details found in Appendix C.

One of the first trials to evaluate long-term morbidity and mortality outcomes with an angiotensin II receptor antagonist in patients with HF was ELITE II (Evaluation of Losartan in the Elderly); which evaluated the effects of losartan 50mg once daily compared to captopril 50mg three times daily on overall mortality and cardiac events (sudden cardiac death or resuscitated cardiac arrest).<sup>202</sup> This trial was a follow-up to the original ELITE Study, that although not hypothesized *a priori*, reported a favorable mortality rate with losartan.<sup>203</sup> In ELITE II, there was no significant difference in all-cause mortality between the treatment groups (17.7% on losartan vs. 15.9% on captopril). There was no difference between the groups in sudden death or resuscitated cardiac arrest, or hospital admissions. However, this was a superiority trial not designed to detect equivalence between groups.<sup>202</sup>

The Val-HeFT (Valsartan Heart Failure Treatment) study was a placebo-controlled trial that evaluated the addition of valsartan 160 mg twice daily to patients with HF on standard therapy (93% ACEI, 35% beta-adrenergic blockers). Overall mortality (a primary endpoint) was similar, occurring in 19.7% of patients in the valsartan group and 19.4% of patients on placebo. The combined primary endpoint of mortality and morbidity (i.e., cardiac arrest with resuscitation, HF hospitalization, or intravenous inotropic agents or



vasodilators for over 4 hours) occurred in 28.8% and 32.1% of patients on valsartan and placebo, respectively. This included a reduction in hospitalizations for HF (13.8% valsartan vs. 18.2% placebo). However, death from any cause (as first event) was higher in patients on valsartan compared to patients receiving placebo (14.2% vs. 12.6%, respectively). According to a subgroup analysis, there was an increased risk of mortality ( $p=0.009$ ) and a trend toward an increased risk of combined morbidity and mortality in patients receiving valsartan in conjunction with an ACEI and beta-adrenergic blocker. Patients who were not on an ACEI or beta-adrenergic blocker experienced a significant reduction in mortality ( $p=0.012$ ). Patients on valsartan but not on an ACEI (with or without a beta-adrenergic blocker) had a lower risk of death and a lower risk of the combined endpoint.<sup>204</sup> A subanalysis of the 366 patients in Val-HeFT who were not on an ACEI reported a 33% decrease in all-cause mortality ( $p=0.017$ ) and a 53% decrease in combined morbidity and mortality ( $p<0.001$ ).<sup>205</sup>

The CHARM (Candesartan in Heart failure Assessment of Reduction in Mortality and morbidity) Overall program combined the results of three placebo-controlled trials evaluating therapy with candesartan titrated to a target dose of 32mg once daily:<sup>206</sup> CHARM-Added evaluated patients with systolic HF on standard therapy (100% ACEI; 55% beta-adrenergic blockers);<sup>207</sup> CHARM-Alternative studied patients with systolic HF and previous ACEI intolerance;<sup>208</sup> and CHARM-Preserved included patients with HF and LVEF > 40%.<sup>70</sup> In CHARM-Overall, the primary outcome of all-cause mortality was numerically reduced with candesartan, although the result did not achieve statistical significance. The secondary endpoint of combined CV death or HF hospitalization was significantly reduced by 16% compared to placebo.<sup>206</sup> In a pooled analysis of patients with LVEF  $\leq 40\%$ , there was a significant 12% reduction in mortality.<sup>209</sup>

In the CHARM-Added trial, the combined primary endpoint of CV mortality or HF hospitalization was significantly reduced by 15% compared to placebo in patients on candesartan in addition to standard therapy including an ACEI. The difference in all-cause mortality was not statistically significant. In the subgroup of patients on therapy with candesartan in combination with an ACEI and beta-adrenergic blocker, there was also a significant reduction in the risk of CV death or HF hospitalization compared to patients on placebo; the difference in all-cause mortality in this subgroup was not statistically significant.<sup>207</sup> This is in conflict with the increase in mortality seen in the subgroup analysis of Val-HeFT in patients on combination ACEI, beta-adrenergic blockers, and angiotensin II antagonist.<sup>204</sup> Regarding the addition of an angiotensin II receptor antagonist to standard therapy for HF, the ACC/AHA HF clinical practice guidelines recommend (Class IIb: i.e., usefulness/efficacy is less well established by evidence/opinion) that this may be considered in patients with persistent symptoms despite standard therapy for HF (Level of Evidence B).<sup>1</sup> It is also important to note that the routine use of an ACEI, angiotensin II receptor antagonist, and aldosterone antagonist is not recommended.<sup>1</sup>

The combined primary endpoint of CV mortality or HF hospitalization was reduced by 23% in patients with a history of ACEI intolerance randomized to candesartan compared to those on placebo in the CHARM-Alternative trial. There was not a statistically significant reduction in all-cause mortality.<sup>208</sup>

The angiotensin II receptor antagonists have yet to be shown to be equivalent or superior to the ACEIs in reducing long-term outcomes of morbidity and mortality in randomized controlled trials of patients with HF. A meta-analysis of 38,080 patients reported that use of an angiotensin II receptor antagonist in patients with HF reduced all-cause mortality [OR (odds ratio) 0.83; 95% CI 0.69-1.00] compared to placebo, although this was influenced largely by data from CHARM-Alternative. There was also a statistically significant reduction in HF hospitalizations (OR 0.64; 95% CI 0.53-0.78) with an angiotensin II receptor antagonist compared to placebo. There was not a significant difference in all-cause mortality or HF hospitalizations when data with an angiotensin II receptor antagonist were compared to results with an ACEI. The analysis also compared an angiotensin II receptor antagonist in combination with an ACEI vs. an ACEI alone, without a significant difference in all-cause mortality; although there was a statistically significant reduction in HF hospitalizations (OR 0.77; 95% CI 0.69-0.87) favoring combination therapy.<sup>210</sup> These results are similar to a previous meta-analysis of 12,469 patients that reported a trend toward improved mortality and hospitalizations with an angiotensin II receptor antagonist compared to placebo in patients not on an ACEI, and the combination of an angiotensin II receptor antagonist and ACEI significantly reduced the risk of hospitalizations compared to patients on an ACEI alone.<sup>211</sup> Another meta-

analysis reported a reduction in morbidity and mortality, but not mortality alone, in patients receiving an angiotensin II receptor antagonist in combination with an ACEI, regardless of a beta-adrenergic blocker, or when a beta-adrenergic blocker was not part of therapy. Combined morbidity and mortality, or the endpoint of mortality alone, was not reduced in patients receiving all three classes of medications.<sup>212</sup>

Use of an angiotensin II receptor antagonist can be considered in patients who are unable to tolerate treatment with an ACEI due to cough, although there is a slight chance that patients may develop a cough with an angiotensin II receptor antagonist.<sup>213</sup>

An angiotensin II receptor antagonist should be used with extreme caution in a patient who has previously experienced angioedema on an ACEI.<sup>1</sup> The incidence of angioedema in patients taking ACEIs is approximately 0.1-1.2 %.<sup>214</sup> It has been reported that black American patients have an increased relative risk of 4.5 of angioedema associated with use of an ACEI compared to white patients.<sup>215</sup> Angioedema has been reported with the angiotensin II receptor antagonists but to a much lesser degree than ACEIs. The exact mechanism is unknown; in ACEIs, it is thought to be related to bradykinin accumulation. In the CHARM-Alternative trial with candesartan in patients with HF and a history of ACEI intolerance, 3 of 1013 patients randomized to candesartan experienced angioedema. One of these patients required discontinuation of the drug (0.1%). All 3 cases occurred out of the 39 patients who previously experienced angioedema or anaphylaxis on an ACEI (7.7%). None of the 1015 patients who received placebo experienced angioedema.<sup>208</sup> There have been a number of published case reports of angioedema in patients treated with an angiotensin II receptor antagonist.<sup>214,216-230</sup> In approximately one-third of these cases, the patients previously experienced angioedema with an ACEI. Therefore, extreme caution is warranted in patients who have previously experienced angioedema.<sup>216,224,226,227,231</sup>

The angiotensin II receptor antagonists, like the ACEIs, decrease release of aldosterone from the adrenal cortex, which can lead to decreased potassium excretion. It is unclear at this time if treatment with an angiotensin II receptor antagonist would be an appropriate alternative in patients who develop hyperkalemia on an ACEI.<sup>120,201,203,232</sup> In the CHARM-Overall programme, hyperkalemia resulted in discontinuation of study drug in 2.2% of patients on candesartan compared to 0.6% patients on placebo (p<0.0001). In the overall analysis, 41% of patients received concomitant treatment with an ACEI and approximately 17% were on spironolactone.<sup>206</sup> As with the ACEIs, it is recommended that patients on an angiotensin II receptor antagonist have their blood pressure, renal function, and potassium reevaluated within one to two weeks after initiating therapy, and monitored after dose adjustments.<sup>1</sup> Patients receiving an angiotensin II receptor antagonist in conjunction with an ACEI, potassium supplements, or potassium-sparing diuretics (including spironolactone) should be monitored closely as combination therapy may result in an increased potassium level. Other clinically significant drug interactions with the angiotensin II receptor antagonists are listed in Appendix B.

## EVIDENCE

Grading System		USPSTF <sup>6</sup>			ACC/AHA <sup>1</sup>	
Intervention	References	QE	OQ	SR	CR	LE
Use an angiotensin II receptor antagonist in patients on standard therapy for HF who cannot tolerate an ACEI	Val-HeFT (2001) Maggioni (2002) Jong (2002) CHARM-Alternative (2003) CHARM Overall (2003) Lee (2004) ACC/AHA (2005) HFSA (2006)	I II-2 I I I I III III	I	A	I	A
Consider an angiotensin II receptor antagonist in addition to standard therapy (i.e., an ACEI and beta-blocker) in patients with persistent symptoms	Val-HeFT (2001) Jong (2002) CHARM-Added (2003) CHARM Overall (2003) Lee (2004) Dimopoulos (2004) ACC/AHA (2005) HFSA (2006)	I I I I I I III III	II	B	IIB	B

## **J.      Hydralazine in Combination with a Nitrate**

### **OBJECTIVE**

- To provide recommendations for the appropriate use of hydralazine and a nitrate in patients with a diagnosis of systolic HF

### **BACKGROUND**

An earlier trial compared the combination of hydralazine and isosorbide dinitrate (ISDN) to therapy with an ACEI and based on these results, the combination of hydralazine and ISDN was considered a therapeutic option in patients unable to tolerate an ACEI. The combination of hydralazine and ISDN had not been previously studied in addition to standard therapy (i.e., an ACEI and beta-adrenergic blocker); however, results of a recent trial in self-identified black patients reported a significant reduction in mortality and HF hospitalization with the combination of hydralazine and ISDN. It is not clear whether these results can be extrapolated to the general patient population.

### **RECOMMENDATION**

#### Therapy with Hydralazine and a Nitrate in Stage C HF

- The combination of hydralazine and a nitrate should be considered, especially in African American patients with NYHA Class III or IV HF, who continue to have symptoms despite therapy with an ACEI (or an angiotensin II receptor antagonist if an ACEI is not tolerated) and beta-adrenergic blocker
- The combination of hydralazine and a nitrate may be considered as an alternative to an ACEI in patients who are unable to tolerate an ACEI (or angiotensin II receptor antagonist) due to hypotension, renal insufficiency, hyperkalemia, or possibly, angioedema

### **DISCUSSION**

Peripheral vasodilators such as ISDN (venodilator) and hydralazine (arterial vasodilator) can produce favorable hemodynamic effects in patients with HF.<sup>1</sup> Earlier trials evaluated the combination of hydralazine and ISDN in patients receiving standard therapy for HF (at the time, digoxin and a diuretic). In the first of these two VA trials, the Vasodilator-Heart Failure Trial I (V-HeFT I), treatment with ISDN and hydralazine was reported to significantly reduce mortality by two years compared to placebo (25.6% vs. 34.3%, respectively).<sup>233</sup> The second trial, V-HeFT II, compared treatment with ISDN and hydralazine to that of an ACEI in HF patients (majority with NYHA class II or III HF). Mortality with ISDN and hydralazine was similar to that seen in V-HeFT I (25%), although mortality after two years was lower in patients treated with an ACEI (18%) compared to patients on hydralazine and ISDN.<sup>121</sup> The authors concluded that the similar reduction in mortality seen with the combination of hydralazine and ISDN in V-HeFT I and V-HeFT II, compared with the mortality in the placebo group, along with the reduction in mortality seen with an ACEI, suggested that there is benefit in using a vasodilator as part of the treatment regimen in patients with HF, and that there may be an additional benefit of using the two treatments together.<sup>121</sup>

As discussed previously, there may be racial differences in response to therapy with the ACEIs where black patients may not derive as much benefit as seen in white patients.<sup>125</sup> Different results have been found with hydralazine and ISDN, where there has been a greater benefit in black patients compared to white patients.<sup>124</sup> Racial differences in response to therapy have been reported in subanalyses of the V-HeFT I and V-HeFT II trials. The annual mortality rate was significantly lower in black patients receiving ISDN and hydralazine in V-HeFT I compared to black patients receiving placebo (9.7% vs. 17.3%, respectively); a similar effect was not seen in white patients on hydralazine and ISDN vs. placebo (annual mortality rate 16.9% vs. 18.8%, respectively).<sup>124</sup> In V-HeFT II, white patients on enalapril experienced a significant decrease in mortality compared to treatment with hydralazine and ISDN (annual mortality rate 11.0% vs.



14.9%, respectively), whereas black patients did not have a similar benefit (annual mortality rate with enalapril 12.8% vs. 12.9% with hydralazine and ISDN).<sup>124</sup>

More recently, the African-American Heart Failure Trial (A-HeFT), a long-term morbidity and mortality trial in self-identified black patients with NYHA class III to IV HF, evaluated the fixed-dose combination of hydralazine and ISDN compared to placebo. The majority of patients enrolled in the trial were also receiving treatment with an ACEI (or angiotensin II receptor antagonist), a beta-adrenergic blocker, diuretic, and digoxin; over one-third of patients were also receiving an aldosterone antagonist. The trial was planned for 18 months of follow-up but was terminated early (mean follow-up 10 months) due to a significant reduction in mortality in patients receiving treatment (6.2%) compared to those on placebo (10.2%). Treatment was associated with a 43% improvement in survival. There was also a significant 33% reduction in first hospitalization for HF (another pre-specified component of the primary endpoint) with treatment compared to placebo (16.4% vs. 24.4%, respectively). The primary endpoint was a composite score (possible range -6 to +2, with a higher score representing a better outcome) with weighted values based on mortality, survival to the end of the trial, first hospitalization for HF, no hospitalizations, and change in quality of life. It was reported that patients receiving the combination hydralazine and ISDN had a primary composite score of  $-0.1 \pm 1.9$  compared to  $-0.5 \pm 2.0$  in the placebo group ( $p=0.01$ ), indicating a benefit with hydralazine and ISDN in addition to standard drug therapy.<sup>234</sup>

Side-effects such as headache, tachycardia, flushing, hypotension, and edema, as well as dosing frequency, preclude the use of this regimen in as many as one third of patients. Other adverse effects reported with hydralazine include rash, arthralgia, and other lupus-like symptoms. Common drug interactions are listed in Appendix B.

## EVIDENCE

Grading System Intervention	References	USPSTF <sup>6</sup>			ACC/AHA <sup>1</sup>	
		QE	OQ	SR	CR	LE
Consider combination of hydralazine and a nitrate, especially in African American patients in NYHA Class III or IV, who continue to have symptoms despite therapy with an ACEI (or angiotensin II receptor antagonist if an ACEI is not tolerated) and beta-adrenergic blocker	A-HeFT (2004) V-HeFT I (1986) V-HeFT II (1991) ACC/AHA (2005) HFSA (2006)	I I I III III	I	B	Ila	A
Consider combination of hydralazine and a nitrate in patients unable to take an ACEI or angiotensin II receptor antagonist due to hypotension, renal insufficiency, or drug intolerance	V-HeFT I (1986) V-HeFT II (1991) ACC/AHA (2005) HFSA (2006)	I I III III	II	C	Ilb	C

## K. Digitalis

### OBJECTIVE

- To provide recommendations for the appropriate use of digoxin in patients with a diagnosis of systolic HF

### BACKGROUND

Trials with digoxin have shown it to be beneficial in reducing HF associated symptoms and hospitalizations in patients on standard therapy at trial enrollment (i.e., diuretic and an ACEI), but not in improving survival.

### RECOMMENDATION

#### Digoxin Therapy in Stage C HF

- Digoxin may be useful in decreasing hospitalizations in patients with current or previous HF symptoms

## DISCUSSION

Digoxin is thought to be beneficial in patients with systolic HF through inhibition of sodium-potassium adenosine triphosphatase resulting in reduced activation of the neurohormonal system and increased contractility of the heart.<sup>1</sup> The use of agents with positive inotropic activity as the mainstay of therapy for HF has decreased over the years. This has primarily been due to the increased mortality associated with some of the agents in this class. Digoxin appears to continue to have a role in the treatment of patients with HF by improving patient symptoms and decreasing hospitalizations, without adversely affecting survival.<sup>56,119,132,235</sup>

According to a meta-analysis, treatment with digoxin in patients with HF due to systolic dysfunction can reduce the incidence of clinical deterioration<sup>235,236</sup> by 12% compared to patients on placebo.<sup>235</sup> The Randomized Assessment of (the effect of) Digoxin on Inhibitors of the Angiotensin-Converting Enzyme (RADIANCE) Study evaluated 178 patients with NYHA class II or III HF stabilized on digoxin, diuretics, and an ACEI. Patients were randomized to continuation of treatment or withdrawal of digoxin therapy for 12 weeks. Patients who were withdrawn from digoxin experienced worsening HF and a decreased exercise tolerance, worsening NYHA class, decreased quality of life and LVEF.<sup>237</sup> The Prospective Randomized Study of Ventricular Failure and the Efficacy of Digoxin (PROVED) trial was a study evaluating 88 patients with NYHA class II or III HF on digoxin and diuretics and the effect of digoxin withdrawal or continuation of therapy. Patients who had digoxin withdrawn experienced a worsening of maximum exercise performance, a higher percentage of treatment failures, and a decreased time to treatment failure.<sup>238</sup>

These trials demonstrate the benefit of digoxin in reducing symptoms associated with mild to moderate HF. The Digitalis Investigators Group (DIG) trial evaluated the benefit of digoxin on survival. This trial enrolled 6,800 patients on diuretics and an ACEI who were randomized to receive digoxin or placebo for a mean of 37 months. The results showed that treatment with digoxin significantly decreased the risk for hospitalizations due to HF by 28%, although there was no significant reduction in mortality with digoxin treatment.<sup>56</sup> In a post hoc analysis of the DIG trial, a decrease in the rate of cardiovascular deaths and deaths from worsening HF was found in the men, but not in the women who were treated with digoxin. The death rate in women on digoxin was higher than women randomized to placebo (33.1% vs. 28.9%, respectively;  $p=0.078$ ). There was a decrease in hospitalizations for worsening HF in women on digoxin compared to women on placebo (30.2% vs. 34.4%, respectively;  $p=0.079$ ). The median serum digoxin concentration was significantly higher in women compared to men (0.9ng/ml based on 475 randomly selected women vs. 0.8ng/ml in 1653 randomly selected men at one month after randomization;  $p=0.007$ ); although, there was not a statistically significant difference at 12 months (0.6ng/ml in randomly selected men and women, respectively). Due to these findings, the authors suggest that the role of digoxin in women be reevaluated.<sup>239</sup> Others suggest that a lower dose with a resultant serum concentration < 1ng/ml be used.<sup>239,240</sup>

More recently, the relationship between serum digoxin concentrations and morbidity and mortality in women in the DIG trial were evaluated. This retrospective analysis demonstrated a reduction in death or HF hospitalization and no increase in mortality in women with a digoxin concentration 0.5-0.9ng/ml. A serum digoxin concentration of 1.2-2.0ng/ml was associated with an increase in risk for death in women.<sup>241</sup> Another retrospective analysis of the DIG trial did not find a relationship on outcomes based on race.<sup>242</sup>

Digoxin is recommended in patients with symptomatic HF, without bradycardia, to improve clinical status and thereby decrease the risk of hospitalization due to HF. Treatment is usually initiated in conjunction with a diuretic, ACEI, and beta-adrenergic blocker since these latter two classes of agents have been shown to improve survival in patients with HF.<sup>1</sup> If there is no symptomatic improvement after one to two months of therapy, the risk vs. benefit of continued digoxin therapy should be considered. Digoxin is particularly useful to control rapid ventricular response in patients with systolic dysfunction and atrial fibrillation.<sup>1</sup>

Loading doses are not necessary for patients in normal sinus rhythm. The most commonly prescribed dose of digoxin is 0.125-0.25mg/day. Initial dosing should be conservative (e.g., 0.125mg once daily or every other day) especially for patients with reduced CrCl, decreased weight and/or decreased muscle mass. The utility of monitoring serum digoxin levels to assess efficacy has not been established.<sup>1</sup> Subgroup analysis from the DIG trial as well as in the Prospective Randomized Milrinone Survival Evaluation (PROMISE) trials showed that higher concentrations (even within the therapeutic range) were associated with an increased risk of mortality.<sup>243-246</sup> In both the RADIANCE and PROVED trials, the mean digoxin serum concentration was 1.2ng/ml and in the DIG trial, the mean serum digoxin level was 0.8 ng/ml at 12 months.<sup>237,238,244,246</sup> In a meta-analysis of the PROVED and RADIANCE trials, the clinical efficacy (e.g., worsening HF, change in LVEF, treadmill time) of low (0.5-0.9ng/ml), moderate (0.9-1.2ng/ml), and high (>1.2ng/ml) serum digoxin concentrations were compared. There was no relationship between the endpoints and the three groups.<sup>247</sup> A post hoc analysis of the DIG trial showed a linear relationship for mortality and increasing serum digoxin concentrations with a lower mortality seen in patients with a digoxin serum concentration of 0.5-0.8ng/ml, no reduction in mortality at 0.9-1.1ng/ml, and an increase in mortality at levels  $\geq 1.2$ ng/ml. The analysis was limited to men.<sup>248</sup> The authors concluded that lower levels may provide optimal benefit without the risk of detrimental effects seen with higher levels,<sup>246,247</sup> although levels are not typically drawn unless monitoring for toxicity.

In general, trough (or a minimum of 6 hours post dose due to distribution) serum digoxin levels should be monitored if any of the following occurs:

1. HF worsens or renal function deteriorates
2. Signs of toxicity develop (e.g., confusion, nausea, vomiting, abdominal pain, diarrhea, anorexia, fatigue, arrhythmias, visual disturbances)
3. Dose adjustments are made
4. Medications are added that affect the serum digoxin concentration (e.g., quinidine, verapamil, amiodarone, antibiotics, anticholinergics) (refer to Appendix B), or the sensitivity to digoxin by altering potassium levels

## EVIDENCE

Grading System	USPSTF <sup>6</sup>			ACC/AHA <sup>1</sup>	
Intervention	References	QE	OQ	SR	CR LE
Consider digoxin to improve functional status and reduce frequency of hospitalizations if continued symptoms	Captopril-Digoxin (1988)	I	II	B	Ila B
	Jaeschke (1990)	I			
	RADIANCE (1993)	I			
	PROVED (1993)	I			
	DIG (1999)	I			
	Hood (2004)	I			
	ACC/AHA (2005)	III			

## L. Aldosterone Antagonists

### OBJECTIVE

- To provide recommendations for the appropriate use of aldosterone antagonists in patients with a diagnosis of systolic HF

### BACKGROUND

Aldosterone antagonists (e.g., spironolactone, eplerenone) competitively inhibit the effects of aldosterone. One of the proposed mechanisms for benefit of using ACEIs in patients with HF is that of suppression of production of aldosterone. Additional therapy with an aldosterone antagonist was originally felt not to be necessary, with concern for an increase in the risk of hyperkalemia due to potential for potassium retention if aldosterone is decreased. Evidence has shown that addition of an aldosterone antagonist may be beneficial in patients with severe HF (recent NYHA class IV HF and current class III or IV symptoms and LVEF  $\leq 35\%$ ), even in patients already receiving an ACEI.<sup>1,249</sup> This suggests that therapy with an ACEI

may not achieve long-term suppression of aldosterone production. There is insufficient evidence to make a recommendation as to the use of aldosterone antagonists in patients with mild to moderate HF.

## RECOMMENDATION

### Aldosterone Antagonist Therapy in Stage C HF

- An aldosterone antagonist is beneficial in selected patients (e.g., moderately severe to severe HF symptoms with reduced LVEF, or patients with  $\text{LVEF} \leq 40\%$  early post-MI, and with adequate kidney function and no hyperkalemia) who can be monitored for hyperkalemia or renal dysfunction

## DISCUSSION

The above recommendations are based on the Randomized Aldactone Evaluation Study (RALES), a study that enrolled 1663 patients with severe class IV HF within the last 6 months (and class III or IV at time of enrollment), a  $\text{LVEF} \leq 35\%$  within the last 6 months, and treated with conventional therapy (95% ACEI, 100% loop diuretic, 75% digoxin). In addition, 11% of patients were on a beta-adrenergic blocker. Patients were randomized to spironolactone 25mg once daily or placebo. The primary endpoint was to evaluate all-cause mortality. After a mean follow-up of 24 months, the trial was discontinued early due to a 30% reduction in the risk of death due to progressive HF and sudden death of a cardiac cause in patients in the spironolactone group (45.9% on placebo vs. 34.6% on spironolactone). There was also a significant 35% decrease in hospitalizations due to worsening HF in patients on spironolactone; these patients also experienced significant improvement in symptoms.<sup>249</sup>

The Eplerenone Post-Acute Myocardial Infarction Heart Failure Efficacy Survival Study (EPHESUS) compared an aldosterone antagonist, eplerenone (mean dose 42.6mg per day), to placebo in 6642 patients with acute MI complicated by LV dysfunction, and symptoms of HF (patients with DM could be enrolled without having HF symptoms), with the majority on standard therapy for this indication. Ninety percent of patients had symptomatic HF. Treatment with eplerenone significantly reduced the primary endpoints of death from any cause (14.4% of patients on eplerenone vs. 16.7% of patients in the placebo group) and death from cardiovascular causes or first hospitalization for a cardiovascular event, including HF, recurrent acute MI, stroke, or ventricular arrhythmia (eplerenone 26.7% vs. placebo 30.0%).<sup>250</sup> Based on these data, an aldosterone antagonist may also be considered early after an acute MI in patients with LV dysfunction and HF.<sup>1,53</sup>

These are highly complex patients with a high mortality rate and should be cared for by a multidisciplinary HF team including a primary care provider in consultation with a cardiologist.<sup>251</sup> The risk vs. benefit of using an aldosterone antagonist in these patients needs to be determined. An aldosterone antagonist may contribute to serious hyperkalemia if not used properly in patients with HF.<sup>251,252</sup>

In addition to hyperkalemia, aldosterone antagonists can cause gynecomastia, gastrointestinal side effects, and menstrual irregularities. In RALES, gynecomastia or breast pain was reported in 10% of male patients in the spironolactone group. The incidence of hyperkalemia was not significant. However, it should be noted that in both the RALES and EPHESUS trials, patients with serum creatinine  $> 2.5$  mg/dL and serum potassium  $> 5.0$  mmol/L were excluded and patients were not taking other potassium-sparing diuretics. In EPHESUS, the mean serum Cr concentration was 1.1 mg/dl at baseline.<sup>250</sup> In clinical practice, reports of discontinuations due to hyperkalemia appear to be higher than seen in the clinical trial.<sup>253-255</sup> Hyperkalemia occurs more frequently in patients receiving potassium supplements and in patients with renal insufficiency. Use of potassium supplements with an aldosterone antagonist should be avoided unless hypokalemia develops. The aldosterone antagonists should be used with caution in patients with renal insufficiency; patients should be scheduled for follow-up electrolytes and renal function after initiation and dose adjustments.<sup>256</sup> An aldosterone antagonist should also be used with caution in patients receiving ACEIs or angiotensin II receptor antagonists due to the potential for hyperkalemia; potassium should be monitored closely in these patients.<sup>257,258</sup> In general, potassium supplements should be discontinued when therapy with an aldosterone antagonist is initiated.<sup>1</sup> Serum potassium should be monitored within 3 days

and at 1 week, and every 4 weeks for the first 3 months, then every 3 months thereafter.<sup>249,252</sup> More frequent monitoring may be indicated in patients on concomitant medications that may increase potassium levels, with renal insufficiency or DM, who are of advanced age, experiencing worsening HF or conditions that may contribute to dehydration.<sup>257-260</sup> If the potassium increases to > 5.5 mEq/L, the aldosterone antagonist should be discontinued or the dose reduced. If serious hyperkalemia develops, therapy with the aldosterone antagonist should be discontinued.<sup>1</sup>

The initial dose of spironolactone used in RALES was 25mg once daily. The dose was decreased to 25mg every other day in patients exhibiting hyperkalemia. The dose was increased to 50mg once daily at 8 weeks in patients who had signs or symptoms of worsening HF and did not have hyperkalemia. Patients receiving 50mg spironolactone should have their serum potassium measured one week after the dose was increased, and then follow-up as described above.<sup>252</sup> Refer to Appendix B for common drug interactions.

## EVIDENCE

Grading System Intervention	References	USPSTF <sup>b</sup>			ACC/AHA <sup>1</sup>	
		QE	OQ	SR	CR	LE
Consider low dose of an aldosterone antagonist (e.g., 12.5 to 25mg/d spironolactone) in patients with severe HF (recent NYHA class IV HF and current class III or IV symptoms), provided the potassium is normal (< 5 mEq/L) and kidney function is adequate (serum Cr ≤ 2.5 mg/dL in men; ≤ 2.0 mg/dL in women) and in whom potassium and renal function can be carefully monitored	RALES (1999) ACC/AHA (2005) HFSA (2006)	I III III	I	A	I	B
Consider addition of an aldosterone antagonist in patients on standard therapy for HF with LVEF ≤ 40% early post-MI, who have a normal potassium and adequate kidney function; patients should be monitored for changes in potassium and kidney function	EPHESUS (2003) ACC/AHA (2005) HFSA (2006)	I III III	I	A	NA	NA

## M. Continue Present Management and Schedule Regular Follow-up

### OBJECTIVE

- To provide recommendations for appropriate follow-up of patients with a diagnosis of systolic HF

### BACKGROUND

Patients should receive regular follow-up in order to provide the most effective care. At each encounter, an inquiry should be made as to the patient's adherence to the medication regimen, nonpharmacologic measures, and adverse effects to therapy. Patients should be scheduled for routine laboratory monitoring. The patient should also be assessed for any change in functional status or frequency of hospitalizations, and medication therapy should be optimized. A multidisciplinary approach to care and follow-up should be utilized if appropriate to potentially improve care and outcomes.<sup>1,22,53</sup>

### RECOMMENDATIONS

#### General Recommendation

- Patients should receive regular follow-up

#### Multidisciplinary Disease Management Programs

The following is a Class I recommendation by the ACC/AHA (i.e., there is evidence and/or general agreement that a given procedure/therapy is useful and effective)<sup>1</sup>

- Recommended for patients at high risk for hospital admission or clinical deterioration to facilitate implementation of clinical practice guidelines, address barriers to behavioral change, and to decrease the risk of HF hospitalization

The following is a Class IIb recommendation by the ACC/AHA (i.e., the usefulness/efficacy is less well established by evidence/opinion)<sup>1</sup>

- Consider for patients at low risk for hospital admission or clinical deterioration to facilitate implementation of clinical practice guidelines

#### Performance Measures (refer to discussion in Introduction)

The following is a Class IIa recommendation by the ACC/AHA (i.e., the weight of evidence/opinion is in favor of usefulness/efficacy)<sup>1</sup>

- Performance measures based on clinical practice guidelines may improve quality of patient care

### **DISCUSSION**

Routine follow-up is an essential component of the overall management of patients with HF.<sup>1,261</sup> At this time the patient's functional status can be evaluated and any adjustments made to the medication regimen. The presence of any adverse events should also be determined. Evaluation of the patient's serum potassium is important due to the influence of medications on this parameter. There is the potential for hypokalemia with diuretics that may lead to toxicity in a patient receiving digoxin. The ACEIs, angiotensin II receptor antagonists, and aldosterone antagonists may all increase potassium, leading to potential toxicity.<sup>1</sup>

Adherence to the medication regimen is often not optimal<sup>262,263</sup> and may lead to clinical deterioration in patients with HF.<sup>5</sup> Patients need to be educated on the importance of adherence to the medication regimen in order to derive the benefits of decreased morbidity and mortality. The reason for not taking a medication as prescribed should be investigated. If it is a result of an adverse effect, the dosage of the medication can be adjusted or another class of medication considered.

Some facilities may have interdisciplinary HF disease management clinics or specialized programs to provide continuity of care and improve treatment outcomes for patients with HF.<sup>1,186,265-282</sup> Heart failure disease management clinics have improved patient outcomes including improved function status,<sup>186,272,274</sup> fewer hospitalizations,<sup>272-275,282</sup> a reduction in mortality,<sup>186,266,275-278</sup> increased utilization of ACEI and/or beta-adrenergic blockers or their doses.<sup>186,266,267,278</sup> In addition, reports suggest the use of these disease management programs may be cost-effective.<sup>186,273</sup>

Proper education of patients and their family is imperative so that they may have an understanding of the cause of HF, prognosis, therapy, dietary restrictions, activity, adherence, and the signs and symptoms of recurrent HF.<sup>1,279</sup> If patients and/or caregivers are cognizant of the signs and symptoms of recurrent HF, they may have the opportunity to present to the healthcare practitioner before the patient's condition deteriorates. Patients and caregivers should also be educated on the patient's prognosis for function and survival. Treatment options, a living will, and advanced directives should be discussed with the patient and caregiver in response to different events that may occur. The availability of hospice care should also be discussed. Continuity of care is important for the patient's overall care and for the implementation of the patient's request for end of life care.<sup>1</sup>

## REFERENCES



1. Hunt SA, Abraham WT, Chin MH, et al. ACC/AHA 2005 guideline update for the diagnosis and management of chronic heart failure in the adult: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Update the 2001 Guidelines for the Evaluation and Management of Heart Failure). American College of Cardiology Web site. Available at: <http://www.acc.org/clinical/guidelines/failure/index.pdf>.
2. Thom T, Haase N, Rosamond W, et al. Heart disease and stroke statistics-2006 update. A report from the American Heart Association statistics committee and stroke statistics subcommittee. 2006. *Circulation*. Available at <http://circ.ahajournals.org/cgi/content/short/113/6/e85>.
3. Skrepnek GH, Abarca J, Malone DC, et al. Incremental effects of concurrent pharmacotherapeutic regimens for heart failure on hospitalizations and costs. *Ann Pharmacother* 2005;39:1785-91.
4. Komajda M, Lapuerta P, Hermans N, et al. Adherence to guidelines is a predictor of outcome in chronic heart failure: the MAHLER survey. *Eur Heart J* 2005;26:1653-9.
5. Lee DS, Mamdani MM, Austin PC, et al. Trends in heart failure outcomes and pharmacotherapy: 1992 to 2000. *Am J Med* 2004;116:581-9.
6. Harris RP, Helfand M, Woolf SH, et al. for the Methods Work Group, Third U.S. Preventive Services Task Force. Current methods of the U.S. Preventive Services Task Force. A review of the process. *Am J Prev Med* 2001;20(3S):21-35.
7. Badgett RG, Lucey CR, Mulrow CD. Can the clinical examination diagnose left-sided heart failure in adults? *JAMA* 1997;277:1712-9.
8. Thomas JT, Kelly RF, Thomas SJ, et al. Utility of history, physical examination, electrocardiogram, and chest radiograph for differentiating normal from decreased systolic function in patients with heart failure. *Am J Med* 2002;112:437-45.
9. Mueller C, Scholer A, Laule-Kilian K, et al. Use of B-type natriuretic peptide in the evaluation and management of acute dyspnea. *N Engl J Med* 2004;350:647-54.
10. Criteria Committee of the American Heart Association. 1994 revisions to the classification of functional capacity and objective assessment of patients with disease of the heart. *Circulation* 1994;90:644-5.
11. Levy D, Larson MG, Vasan RS, Kannel WB, Ho KK. The progression from hypertension to congestive heart failure. *JAMA* 1996;275:1557-62.
12. He J, Ogden LG, Bazzano LA, et al. Risk factors for congestive heart failure in US men and women: NHANES I epidemiologic follow-up study. *Arch Intern Med* 2001;161:996-1002.
13. Chobanian AV, Bakris GL, Black HR, et al. The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure: the JNC 7 report. *JAMA* 2003;289:2560-72.
14. ALLHAT Officers and Coordinators for the ALLHAT Collaborative Research Group. The Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial. Major outcomes in high-risk hypertensive patients randomized to angiotensin-converting enzyme inhibitor or calcium channel blocker vs diuretic: The Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial (ALLHAT). *JAMA* 2002;288:2981-97.
15. Heart Outcomes Prevention Evaluation (HOPE) Study Investigators. Effects of ramipril on cardiovascular and microvascular outcomes in people with diabetes mellitus: results of the HOPE study and MICRO-HOPE substudy. *Lancet* 2000;355:253-59.
16. Yusuf S, Sleight P, Pogue J, et al. for the Heart Outcomes Prevention Evaluation Study Investigators. Effects of an angiotensin-converting-enzyme inhibitor, ramipril, on cardiovascular events in high-risk patients. *N Engl J Med* 2000;342:145-53.
17. Fox KM. Efficacy of perindopril in reduction of cardiovascular events among patients with stable coronary artery disease: randomized, double-blind, placebo-controlled, multicenter trial (the EUROPA study). *Lancet* 2003;362:782-8.
18. Braunwald E, Domanski MJ, Fowler SE, et al. for the PEACE Trial Investigators. Angiotensin-converting enzyme inhibition in stable coronary artery disease. *N Engl J Med* 2004;351:2058-68.
19. National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III) final report. *Circulation* 2002;106:3143-421.
20. Waldo AL, Prystowsky EN. Drug treatment of atrial fibrillation in the managed care era. *Am J Cardiol* 1998;81(5A):23C-9C.
21. Singer DE, Albers GW, Dalen JE, Go AS, Halperin JL, Manning WJ. Antithrombotic therapy in atrial fibrillation: the Seventh ACCP Conference on Antithrombotic and Thrombolytic Therapy. *Chest* 2004;126(3 Suppl):429S-56S.
22. The Task Force for the Diagnosis and Treatment of Chronic Heart failure of the European Society of Cardiology. Guidelines for the diagnosis and treatment of chronic heart failure: executive summary (update 2005). *Eur Heart J* 2005;26:1115-40.



23. Baker DW, Wright RF. Management of heart failure IV: Anticoagulation for patients with heart failure due to left ventricular systolic dysfunction. *JAMA* 1994;272:1614-18.
24. Thatai D, Ahooja V, Pullicino PM. Pharmacological prevention of thromboembolism in patients with left ventricular dysfunction. *Am J Cardiovasc Drugs* 2006;6:41-9.
25. Al-Khadra AS, Salem DN, Rand WM, et al. Warfarin anticoagulation and survival: a cohort analysis from the Studies of Left Ventricular Dysfunction. *J Am Coll Cardiol* 1998;31:749-53.
26. The Stroke Prevention in Atrial Fibrillation Investigators. Stroke prevention in atrial fibrillation study. Final results. *Circulation* 1991;84:527-39.
27. Hirsch J, Dalen JE, Fuster V, et al. Aspirin and other platelet-active drugs: the relationship among dose, effectiveness, and side effects. *Chest* 1995;108(Suppl):247S-57S.
28. Aspirin use and outcomes in a community-based cohort of 7352 patients discharged after first hospitalization for heart failure. *Circulation* 2006;113:2572-8.
29. Cleland JG, Findlay I, Jafri S, et al. The Warfarin/Aspirin Study in Heart failure (WASH): a randomized trial comparing antithrombotic strategies for patients with heart failure. *Am Heart J* 2004;148:157-64.
30. Packer M, O'Connor M, Ghali JK et al for The Prospective Randomized Amlodipine Survival Evaluation Study Group. Effect of amlodipine on morbidity and mortality in severe chronic heart failure. *N Engl J Med* 1996;335:1107-14.
31. Jafary F. Late-breaking clinical trial results – Session III. PRAISE-2: Prospective Randomized Amlodipine Survival Evaluation. Presented by: Packer M. ACC Scientific Session 2000 <http://www.medscape.com/>
32. Cohn J, Ziesche S, Smith R et al for the V-HeFT Study Group. Effect of the calcium antagonist felodipine as supplementary vasodilator therapy in patients with chronic heart failure treated with enalapril: V-HeFT III. *Circulation* 1997;96:856-63.
33. Alexander RW, Schlant RC, Fuster V, eds. Hurst's the heart, arteries and veins. 9<sup>th</sup> ed. New York:McGraw Hill; 1998:2050.
34. Braunwald E, ed. Heart disease: a textbook of cardiovascular medicine. 5<sup>th</sup> ed. Philadelphia: W. B. Saunders;1997:1429.
35. Willerson JT, Chen PC, Hartwell EA, Buja LM. Congestive heart failure in a 70-year-old man [clinical conference]. *Circulation* 1993;88:1336-47.
36. Brandt K, Cathcart ES, Cohen AS. A clinical analysis of the course and prognosis of forty-two patients with amyloidosis. *Am J Med* 1968;44:955-69.
37. Buja LM, Khoi NB, Roberts WC. Clinically significant cardiac amyloidosis: clinicopathologic findings in 15 patients. *Am J Cardiol* 1970;26:394-405.
38. Jacobson DR, Ittmann M, Buxbaum JN, Wiecezorek R, Gorevic PD. Cardiac amyloidosis resulting from transthyretin Ile 122 deposition in african-americans: two case reports. *Texas Heart Inst J* 1997;24:45-52.
39. Cassidy JT. Cardiac amyloidosis: two cases with digitalis sensitivity. *Ann Intern Med* 1961;55:989.
40. Pomerance A. Senile cardiac amyloidosis. *Br Heart J* 1965;27:711-8.
41. Ridolfi RL, Bulkley BH, Hutchins GM. The conduction system in cardiac amyloidosis. Clinical and pathologic features of 23 patients. *Am J Med* 1977;62:677-86.
42. Hodkinson HM, Pomerance A. The clinical significance of senile cardiac amyloidosis: a prospective clinicopathological study. *Q J Med* 1977;46:381-7.
43. Gertz MA, Falk RH, Skinner M, Cohen AS, Kyle RA. Worsening of congestive heart failure in amyloid heart disease treated by calcium channel-blocking agents. *Am J Cardiol* 1985;55:1645.
44. Griffiths BE, Hughes P, Dowdle R, Stephens MR. Cardiac amyloidosis with asymmetrical septal hypertrophy and deterioration after nifedipine. *Thorax* 1982;37:711-2.
45. Pollak A, Falk RH. Left ventricular systolic dysfunction precipitated by verapamil in cardiac amyloidosis. *Chest* 1993;104:618-20.
46. Doval HC, Nul DR, Grancelli HO, Perrone SV, Bortman GR, Curiel R, for the Grupo de Estudio de la Sobrevida' en la Insuficiencia Cardiaca en Argentina (GESICA). Randomized trial of low-dose amiodarone in severe congestive heart failure. *Lancet* 1994;344:493-8.
47. Singh SN, Fletcher RD, Fisher SG, et al., for the Survival Trial of Antiarrhythmic Therapy in Congestive Heart Failure. Amiodarone in patients with congestive heart failure and asymptomatic ventricular arrhythmias. *N Engl J Med* 1995;333:77-82.
48. Massie BM, Fisher SG, Radford M, et al. Effect of amiodarone on clinical status and left ventricular function in patients with congestive heart failure. CHF-STAT Investigators. *Circulation* 1996;93:2128-34.
49. Torp-Pederson C, Møller M, Bloch-Thomsen PE, et al., for the Danish Investigations of Arrhythmia and Mortality on Dofetilide Study Group. Dofetilide in patients with congestive heart failure and left ventricular dysfunction. *N Engl J Med* 1999;341:857-65.
50. Page J, Henry D. Consumption of NSAIDs and the development of congestive heart failure in elderly patients: an underrecognized public health problem. *Arch Intern Med* 2000;160:777-84.

51. Feenstra J, Heerdink ER, Grobbee DE, Stricker BHC. Association of nonsteroidal anti-inflammatory drugs with first occurrence of heart failure and with relapsing heart failure: the Rotterdam study. *Arch Intern Med* 2002;162:265-70.
52. Nesto RW, Bell D, Bonow RO, et al. Thiazolidinedione use, fluid retention, and congestive heart failure. A Consensus Statement From the American Heart Association and American Diabetes Association. *Circulation* 2003;108:2941-8.
53. Adams KF, Lindenfeld J, Arnold JMO, et al. HFSA 2006 Comprehensive Heart Failure Practice Guideline. *J Cardiac Failure* 2006;12:e1-e122.
54. Arnold JMO, Liu P, Demers P, et al. Canadian Cardiovascular Society consensus conference recommendations on heart failure 2006: Diagnosis and management. *Can J Cardiol* 2006;22:23-45.
55. Ahmed A, Aronow WS, Fleg JL. Higher New York Heart Association classes and increased mortality and hospitalization in patients with heart failure and preserved left ventricular function. *Am Heart J* 2006;151:444-50.
56. Massie BM, Abdalla I. Heart failure in patients with preserved left ventricular systolic function: do digitalis glycosides have a role? *Prog Cardiovasc Dis* 1998;40:357-69.
57. The Digitalis Investigators Group: The effect of digoxin on mortality and morbidity in patients with heart failure. *N Engl J Med* 1997;336:525-33.
58. Lenihan DJ, Gerson MC, Hoit BD, Walsh RA. Mechanisms, diagnosis and treatment of diastolic heart failure. *Am Heart J* 1995;130:153-66.
59. Goldsmith SR, Dick C. Differentiating systolic from diastolic heart failure: pathophysiologic and therapeutic considerations. *Am J Med* 1993;95:645-54.
60. Bonow RO, Edelson JE. Left ventricular diastolic dysfunction as a cause of congestive heart failure: mechanisms and management. *Ann Intern Med* 1992;117:502-10.
61. Vasan RS, Benjamin EJ, Levy D. Congestive heart failure with normal left ventricular systolic function. *Arch Intern Med* 1996;156:146-57.
62. Weinberger HD. Diagnosis and treatment of diastolic heart failure. *Hosp Pract* 1999;34:115-8, 121-2, 125-6.
63. Setaro J, Zaret BL, Schulman DS, et al. Usefulness of verapamil for congestive heart failure associated with abnormal left ventricular diastolic performance. *Am J Cardiol* 1990;66:981-6.
64. Dahlöf B, Pennert K, Hansson L. Reversal of left ventricular hypertrophy in hypertensive patients: a meta-analysis of 109 treatment studies. *Am J Hypertens* 1992;5:95-110.
65. Gottdiener JS, Reda DJ, Massie BM, et al. Effect of single-drug therapy on reduction of left ventricular mass in mild to moderate hypertension: comparison of six antihypertensive agents. *Circulation* 1997;95:2007-14.
66. Warner JG, Jr., Metzger DC, Kitzman DW, Wesley DJ, Little WC. Losartan improves exercise tolerance in patients with diastolic dysfunction and a hypertensive response to exercise. *J Am Coll Cardiol* 1999;33:1567-72.
67. Zile MR, Brutsaert DL. New concepts in diastolic dysfunction and diastolic heart failure: part II: causal mechanisms and treatment. *Circulation* 2002;105:1503-8.
68. Aronow WS, Kronzon I. Effect of enalapril on congestive heart failure treated with diuretics in elderly patients with prior myocardial infarction and normal left ventricular ejection fraction. *Am J Cardiol* 1993;71:602-4.
69. Aronow WS, Ahn C, Kronzon I. Effect of propranolol versus no propranolol on total mortality plus nonfatal myocardial infarction in older patients with prior myocardial infarction, congestive heart failure, and left ventricular ejection fraction  $\geq 40\%$  treated with diuretics plus angiotensin-converting enzyme inhibitors. *Am J Cardiol* 1997;80:207-9.
70. Little WC, Wesley-Farrington DJ, Hoyle J, et al. Effect of candesartan and verapamil on exercise tolerance in diastolic dysfunction. *J Cardiovasc Pharmacol* 2004;43:288-93.
71. Yusuf S, Pfeffer MA, Swedberg K, et al. for the CHARM Investigators and Committees. Effects of candesartan in patients with chronic heart failure and preserved left-ventricular ejection fraction: the CHARM-Preserved Trial. *Lancet* 2003;777-81.
72. ISIS-4 (Fourth International Study of Infarct Survival) Collaborative Group. ISIS-4: A randomized factorial trial assessing early oral captopril, oral mononitrate, and intravenous magnesium sulphate in 58 050 patients with suspected acute myocardial infarction. *Lancet* 1995;345:669-85.
73. Gruppo Italiano per lo Studio della Sopravvivenza nell'Infarto Miocardico. GISSI-3: effects of lisinopril and transdermal glyceryl trinitrate singly and together on 6-week mortality and ventricular function after acute myocardial infarction. *Lancet* 1994;343:1115-22.
74. Pfeffer MA, Braunwald E, Moye LA, et al. on Behalf of the SAVE Investigators. Effect of captopril on mortality and morbidity in patients with left ventricular dysfunction after myocardial infarction. *N Engl J Med* 1992;327:669-77.
75. The Acute Infarction Ramipril (AIRE) Study Investigators. Effect of ramipril on mortality and morbidity of survivors of acute myocardial infarction with clinical evidence of heart failure. *Lancet* 1993;342:821-8.
76. Hall AS, Murray GD, Ball SG, on behalf of the AIREX Investigators. Follow-up study of patients randomly allocated ramipril or placebo for heart failure after acute myocardial infarction: AIRE Extension Study (AIREX). *Lancet* 1997;349:1493-7.

77. Kober L, Torp-Pedersen C, Carlsen JE, et al. for the Trandolapril Cardiac Evaluation (TRACE) Study Group. A clinical trial of the angiotensin-converting-enzyme inhibitor trandolapril in patients with left ventricular dysfunction after myocardial infarction. *N Engl J Med* 1995;333:1670-6.
78. Buch P, Rasmussen S, Abildstrom SZ, et al. for the TRACE investigators. The long-term impact of the angiotensin-converting enzyme inhibitor trandolapril on mortality and hospital admissions in patients with left ventricular dysfunction after a myocardial infarction: follow-up to 12 years. *Eur Heart J* 2005;26:145-52.
79. Ferrari R, for the Perindopril and Remodeling in Elderly with Acute Myocardial Infarction Investigators. Effects of angiotensin-converting enzyme inhibition with perindopril on left ventricular remodeling and clinical outcome: results of the randomized Perindopril and Remodeling in Elderly with Acute Myocardial Infarction (PREAMI) Study. *Arch Intern Med* 2006;166:659-66.
80. Gottlieb SS, McCarter RJ, Vogel RA. Effect of beta-blockade on mortality among high-risk and low-risk patients after myocardial infarction. *N Engl J Med* 1998;339:489-97.
81. The Norwegian Multicenter Study Group. Timolol-induced reduction in mortality and reinfarction in patients surviving acute myocardial infarction. *N Engl J Med* 1981;304:801-7.
82. Beta-Blocker Heart Attack Trial Research Group. A randomized trial of propranolol in patients with acute myocardial infarction. I. Mortality results. *JAMA* 1982;247:1707-14.
83. Chadda K, Goldstein S, Byington R, Curb JD. Effect of propranolol after acute myocardial infarction in patients with congestive heart failure. *Circulation* 1986;73:503-10.
84. The CAPRICORN Investigators. Effect of carvedilol on outcome after myocardial infarction in patients with left-ventricular dysfunction: the CAPRICORN randomized trial. *Lancet* 2001;357:1385-90.
85. Vantrimpont P, Rouleau JL, Wun CC, et al., for the SAVE Investigators. Additional beneficial effects of beta-blockers to angiotensin-converting enzyme inhibitors in the Survival and Ventricular Enlargement (SAVE) Study. *J Am Coll Cardiol* 1997;29:229-36.
86. Exner DV, Dries DL, Waclawiw MA, Shelton B, Domanski MJ. Beta-adrenergic blocking agent use and mortality in patients with asymptomatic and symptomatic left ventricular systolic dysfunction: a post hoc analysis of the studies of left ventricular dysfunction. *J Am Coll Cardiol* 1999;33:916-923.
87. Shlipak MG, Browner WS, Noguchi H, et al. Comparison of the effects of angiotensin converting-enzyme inhibitors and beta blockers on survival in elderly patients with reduced left ventricular function after myocardial infarction. *Am J Med* 2001;110:425-33.
88. Dickstein K, Kjekshus J, and the OPTIMAAL Steering Committee, for the OPTIMAAL Study Group. Effects of losartan and captopril on mortality and morbidity in high-risk patients after acute myocardial infarction: the OPTIMAAL randomized trial. *Lancet* 2002;360:752-60.
89. Pfeffer MA, McMurray J, Leizorovicz A, et al. Valsartan, captopril, or both in myocardial infarction complicated by heart failure, left ventricular dysfunction, or both. (VALIANT). *N Engl J Med* 2003;349:1893-906.
90. The SOLVD Investigators. Effect of enalapril on mortality and the development of heart failure in asymptomatic patients with reduced left ventricular ejection fractions. *N Engl J Med* 1992;327:685-91.
91. Jong P, Yusuf S, Rousseau MF, Ahn SA, Bangdiwala SI. Effect of enalapril on 12-year survival and life expectancy in patients with left ventricular systolic dysfunction: a follow-up study. *Lancet* 2003;361:1843-8.
92. Pharmacologic management of heart failure and left ventricular systolic dysfunction: effect in female, black, and diabetic patients, and cost-effectiveness. U.S. Department of Health and Human Services, 2003; Evidence Report/Technology Assessment No. 82.
93. The Digitalis Investigation Group. The effect of digoxin on mortality and morbidity in patients with heart failure. *N Engl J Med* 1997;336:525-33.
94. Vargo DL, Kramer WG, Black PK, et al. Bioavailability, pharmacokinetics, and pharmacodynamics of torsemide and furosemide in patients with congestive heart failure. *Clin Pharmacol Ther* 1995;57:601-9.
95. Brater DC, Day B, Burdette A, Anderson S. Bumetanide and furosemide in heart failure. *Kidney Int* 1984;26:183-9.
96. Murray MD, Deer MM, Ferguson JA, et al. Open-label randomized trial of torsemide compared with furosemide therapy for patients with heart failure. *Am J Med* 2001;111:513-20.
97. Patterson JH, Adams KF Jr, Applefeld MM, Corder CN, Masse BR, for the Torsemide Investigators Group. Oral torsemide in patients with chronic congestive heart failure: effects on body weight, edema, and electrolyte excretion. *Pharmacotherapy* 1994;14:514-21.
98. Wilson JR, Reichek N, Dunkman WB, Goldberg S. Effect of diuresis on the performance of the failing left ventricle in man. *Am J Med* 1981;70:234-9.
99. Parker JQ. The effects of oral ibopamine in patients with mild heart failure: a double-blind placebo controlled comparison to furosemide. *Int J Cardiol* 1993;40:221-7.
100. Richardson A, Bayliss J, Scriven AJ, et al. Double-blind comparison of captopril alone against frusemide plus amiloride in mild heart failure. *Lancet* 1987;2:709-11.
101. Cleland JGF, Swedberg K, Poole-Wilson PA. Successes and failures of current treatment of heart failure. *Lancet* 1998;352(1S):19S1-28S1.

102. Walma EP, Hoes AW, van Dooren C, Prins A, van der Does E. Withdrawal of long term diuretic medication in elderly patients: a double blind randomised trial. *Br Med J* 1997;315:464-8.
103. Grindstead WC, Francis MJ, Marks GF et al. Discontinuation of chronic diuretic therapy in stable congestive heart failure secondary to coronary artery disease or to idiopathic dilated cardiomyopathy. *Am J Cardiol* 1994;73:881-6.
104. Young JB, Gheorghiade M, Uretsky BF, Patterson JH, Adams KF. Superiority of "triple" drug therapy in heart failure: insights from PROVED and RADIANCE trials. *J Am Coll Cardiol* 1998;32:686-92.
105. Ellison DH. The physiologic basis of diuretic synergism: its role in treating diuretic resistance. *Ann Intern Med* 1991;114:886-94.
106. Brater DC. Diuretic resistance: mechanisms and therapeutic strategies. *Cardiology* 1994;84(suppl 2):57-67.
107. Channer KS, McLean KA, Lawson-Matthew P, Richardson M. Combination diuretic treatment in severe heart failure: a randomized controlled trial. *Br Heart J* 1994;71:146-50.
108. Oster JR, Epstein M, Smoler S. Combined therapy with thiazide-type and loop diuretic agents for resistant sodium retention. *Ann Intern Med* 1983;99:405-6.
109. Sica DA, Gehr TWB. Diuretic combinations in refractory oedema states: pharmacokinetic-pharmacodynamic relationships. *Clin Pharmacokinet* 1996;30:229-49.
110. Brater DC. Pharmacology of diuretics. *Am J Med Sci* 2000;319:38-50.
111. Kramer BK, Schweda F, Riegger GAJ. Diuretic treatment and diuretic resistance in heart failure. *Am J Med* 1999;106:90-6.
112. Captopril Multicenter Research Group. A placebo-controlled trial of captopril in refractory chronic congestive heart failure. *J Am Coll Cardiol* 1983;2:755-63.
113. Sharpe DN, Murphy J, Coxon R, Hannan SF. Enalapril in patients with chronic heart failure: a placebo-controlled, randomized, double-blind study. *Circulation* 1984;70:271-8.
114. Chalmers JP, West MJ, Cyran J et al. Placebo-controlled study of lisinopril in congestive heart failure: a multi-centre study. *J Cardiovasc Pharmacol* 1987;9(suppl III):S89-S97.
115. Pflugfelder PW, Baird MG, Tonkon MH, DiBianco R, Pitt B, for the Quinapril Heart Failure Trial Investigators. Clinical consequences of angiotensin-converting enzyme inhibitor withdrawal in chronic heart failure: a double-blind, placebo-controlled study of quinapril. *J Am Coll Cardiol* 1993;22:1557-63.
116. Gunderson T, Swedberg K, Amtorp O, Remes J, Milsson B. Absence of an effect on exercise capacity of 12-weeks treatment with ramipril in patients with moderate congestive heart failure. *Eur Heart J* 1994;15:1659-65.
117. Erhardt L, MacLean A, Ilgenfritz JU, Gelperin K, Blumenthal M, for the Fosinopril Efficacy/Safety Trial (FEST) Study Group. Fosinopril attenuates clinical deterioration and improves exercise tolerance in patients with heart failure. *Eur Heart J* 1995;16:1892-9.
118. Lechat P, Garnham SP, Desche P, Bounhoure JP. Efficacy and acceptability of perindopril in mild to moderate chronic congestive heart failure. *Am Heart J* 1993;126(3 Pt 2):798-806.
119. The Captopril-Digoxin Multicenter Research Group. Comparative effects of therapy with captopril and digoxin in patients with mild to moderate heart failure. *JAMA* 1988;259:539-44.
120. The SOLVD Investigators. Effect of enalapril on survival in patients with reduced left ventricular ejection fractions and congestive heart failure. *N Engl J Med* 1991;325:293-302.
121. Cohn JN, Johnson G, Ziesche S et al. A comparison of enalapril with hydralazine-isosorbide dinitrate in the treatment of chronic congestive heart failure. *N Engl J Med* 1991;325:303-10.
122. The CONSENSUS Trial Study Group. Effects of enalapril on mortality in severe congestive heart failure: results of the Cooperative North Scandinavian Enalapril Survival Study (CONSENSUS). *N Engl J Med* 1987;316:1429-35.
123. Garg R, Yusuf S. Collaborative Group on ACE Inhibitor Trials. Overview of randomized trials of angiotensin-converting enzyme inhibitors on mortality and morbidity in patients with heart failure. *JAMA* 1995;273:1450-56.
124. Carson P, Ziesche S, Johnson G, Cohn JN, for the Vasodilator-Heart Failure Trial Study Group. Racial differences in response to therapy for heart failure: analysis of the Vasodilator-Heart Failure Trials. *J Cardiac Fail* 1999;5:178-87.
125. Exner DV, Dries DL, Domanski MJ, Cohn JN. Lesser response to angiotensin-converting-enzyme inhibitor therapy in black as compared to white patients with left ventricular dysfunction. *N Engl J Med* 2001;344:1351-7.
126. Israili ZH, Hall WD. Cough and angioneurotic edema associated with angiotensin-converting enzyme inhibitor therapy. *Ann Intern Med* 1992;117:234-42.
127. Masoudi FA, Rathore SS, Wang Y, et al. National patterns of use and effectiveness of angiotensin-converting enzyme inhibitors in older patients with heart failure and left ventricular systolic dysfunction. *Circulation* 2004;110:724-31.
128. Bungard TJ, McAlister FA, Johnson JA, Tsuyuki RT. Underutilisation of ACEI inhibitors in patients with congestive heart failure. *Drugs* 2001;61:2021-33.
129. Roe CM, Motheral BR, Teitelbaum F, Rich MW. Angiotensin-converting enzyme inhibitor compliance and dosing among patients with heart failure. *Am Heart J* 1999;138:818-25.
130. Massie BM, Armstrong PW, Cleland JGF, et al. Tolerant of high doses of angiotensin-converting enzyme inhibitors in patients with chronic heart failure: results from the ATLAS trial. *Arch Intern Med* 2001;161:165-71.



131. Packer M, Poole-Wilson PA, Armstrong PW et al on behalf of the ATLAS Study Group. Comparative effects of low and high doses of the angiotensin-converting enzyme inhibitor, lisinopril, on morbidity and mortality in chronic heart failure. *Circulation* 1999;100:2312-18.
132. Majumdar SR, McAlister FA, Cree M, et al. for the Group. Do evidence-based treatments provide incremental benefits to patients with congestive heart failure already receiving angiotensin-converting enzyme inhibitors? A secondary analysis of one-year outcomes from the Assessment of Treatment with Lisinopril and Survival (ATLAS) study. *Clin Ther*. 2004 May;26(5):694-703.
133. The NETWORK investigators. Clinical outcome with enalapril in symptomatic chronic heart failure; a dose comparison. *Eur Heart J* 1998;19:481-9.
134. Nanas JN, Alexopoulos G, Anastasiou-Nana MI, et al., for the High Enalapril Dose Study Group. Outcome of patients with congestive heart failure treated with standard versus high doses of enalapril: a multicenter study. *J Am Coll Cardiol* 2000;36:2090-5.
135. Butler J, Arbogast PG, Daugherty J, et al. Outpatient utilization of angiotensin-converting enzyme inhibitors among heart failure patients after hospital discharge. *J Am Coll Cardiol* 2004;43:2036-43.
136. Ghali JK, Dunselman P, Waagstein F, et al. Consistency of the beneficial effect of metoprolol succinate extended release across a wide range dose of angiotensin-converting enzyme inhibitors and digitalis. *J Card Fail* 2004;10:452-9.
137. Ahmed A, Centor RM, Weaver MT, Perry GJ. A propensity score analysis of the impact of angiotensin-converting enzyme inhibitors on long-term survival of older adults with heart failure and perceived contraindications. *Am Heart J* 2005;149:737-43.
138. Punzi HA. Safety update: focus on cough. *Am J Cardiol* 1993;72:45H-8H.
139. Sharif MN, Evans BL, Pylypchuk GB. Cough induced by quinapril with resolution after changing to fosinopril. *Ann Pharmacother* 1994;28:720-1.
140. Germino FW, Lastra J, Pool P et al. Evaluation of the cough profile of fosinopril in hypertensive patients with ACE inhibitor-associated cough: a pilot study. *Curr Ther Res* 1993;54:469-75.
141. Stys T, Lawson WE, Smaldone GC, Stys A. Does aspirin attenuate the beneficial effects of angiotensin-converting enzyme inhibition in heart failure? *Arch Intern Med* 2000;160:1409-13.
142. Olson KL. Combined aspirin/ACE inhibitor treatment for CHF. *Ann Pharmacother* 2001;35:1653-8.
143. Nguyen KN, Aursnes I, Kjekshus J. Interaction between enalapril and aspirin on mortality after acute myocardial infarction: subgroup analysis of the Cooperative New Scandinavian Enalapril Survival Study II (CONSENSUS II). *Am J Cardiol* 1997;79:115-19.
144. Teo KK, Yusuf S, Pfeffer M, et al. for the ACE Inhibitors Collaborative Group. Effects of long-term treatment with angiotensin-converting-enzyme inhibitors in the presence or absence of aspirin: a systematic review. *Lancet* 2002;360:1037-43.
145. Harjai KJ, Solis S, Prasad A, Loupe J. Use of aspirin in conjunction with angiotensin-converting enzyme inhibitors does not worsen long-term survival in heart failure. *Int J Cardiol* 2003;88:207-14.
146. Aumegeat V, Lamblin N, de Groote P, et al. Aspirin does not adversely affect survival in patients with stable congestive heart failure treated with angiotensin-converting enzyme inhibitors. *Chest* 2003;124:1250-8.
147. Guazzi M, Brambilla R, Reina G, Tumminello G, Guazzi MD. Aspirin-angiotensin-converting enzyme inhibitor coadministration and mortality in patients with heart failure: a dose-related adverse effect of aspirin. *Arch Intern Med* 2003;163:1574-9.
148. Massie BM, Krol WF, Ammon SE, et al. The Warfarin and Antiplatelet Therapy in Heart Failure trial (WATCH): rationale, design, and baseline patient characteristics. *J Card Fail* 2004;10:101-12.
149. Thatai D, Ahooja V, Pullicino PM. Pharmacological prevention of thromboembolism in patients with left ventricular dysfunction. *Am J Cardiovasc Drugs* 2006;6:41-9.
150. Pepper GS, Lee RW. Sympathetic activation in heart failure and its treatment with  $\beta$ -blockade. *Arch Intern Med* 1999;159:225-34.
151. Waagstein F, Bristow MR, Swedberg K et al for the Metoprolol in Dilated Cardiomyopathy (MDC) Trial Study Group. Beneficial effects of metoprolol in idiopathic dilated cardiomyopathy. *Lancet* 1993;342:1441-6.
152. White M, Rouleau JL, Pericak D et al, on behalf of the RESOLVD Study Group. Effects of metoprolol-CR in patients with ischaemic and dilated cardiomyopathy: the RESOLVD pilot study (phase II). (Abstract) *Eur Heart J* 1998;19(suppl):308.
153. CIBIS Investigators and Committees. A randomized trial of  $\beta$ -blockade in heart failure: The Cardiac Insufficiency Bisoprolol Study (CIBIS). *Circulation* 1994;90:1765-73.
154. Colucci WS, Packer M, Bristow MR et al for the US Carvedilol Heart Failure Study Group. Carvedilol inhibits clinical progression in patients with mild symptoms of heart failure. *Circulation* 1996;94:2800-6.
155. Australia/New Zealand Heart Failure Research Collaborative Group. Randomised, placebo-controlled trial of carvedilol in patients with congestive heart failure due to ischaemic heart disease. *Lancet* 1997;349:375-80.
156. Packer M, Colucci WS, Sackner-Bernstein JD et al for the PRECISE Study Group. Double-blind, placebo-controlled study of the effects of carvedilol in patients with moderate to severe heart failure: the PRECISE Trial. *Circulation* 1996;94:2793-9.

157. Bristow MR, Gilbert EM, Abraham WT et al for the MOCHA Investigators. Carvedilol produces dose-related improvements in left ventricular function and survival in subjects with chronic heart failure. *Circulation* 1996;94:2807-16.
158. Cohn JN, Fowler MB, Bristow MR et al for the U.S. Carvedilol Heart Failure Study Group. Safety and efficacy of carvedilol in severe heart failure. *J Card Failure* 1997;3:173-9.
159. MERIT-HF Study Group. Effect of metoprolol CR/XL in chronic heart failure: Metoprolol CR/XL Randomized Intervention Trial in Congestive Heart Failure (MERIT-HF). *Lancet* 1999;353:2001-7.
160. CIBIS-II Investigators and Committees. The Cardiac Insufficiency Bisoprolol Study (CIBIS-II): a randomised trial. *Lancet* 1999;353:9-13.
161. Packer M, Bristow MR, Cohn JN et al for the U.S. Carvedilol Heart Failure Study Group. The effect of carvedilol on morbidity and mortality in patients with chronic heart failure. *N Engl J Med* 1996;334:1349-55.
162. Packer M, Coats AJS, Fowler MB et al. for the Carvedilol Prospective Randomized Cumulative Survival Study Group. Effect of carvedilol on survival in severe chronic heart failure. *N Engl J Med* 2001;344:1651-8.
163. Packer M, Fowler MB, Roecker EB, et al. for the Carvedilol Prospective Randomized Cumulative Survival Study Group. Effect of carvedilol on the morbidity of patients with severe chronic heart failure: results of the Carvedilol Prospective Randomized Cumulative Survival (COPERNICUS) Study. *Circulation* 2002;106:2194-9.
164. Leizorovicz A, Lechat P, Cucherat M, Bugnard F. Bisoprolol for the treatment of chronic heart failure: a meta-analysis on individual data of two placebo-controlled studies-CIBIS and CIBIS II. *Am Heart J* 2002;143:301-7.
165. Shibata MC, Flather MD, Wang D. Systematic review of the impact of beta blockers on mortality and hospital admissions in heart failure. *Eur J Heart Fail* 2001;3:351-7.
166. Willenheimer R, van Veldhuisen DJ, Silke B, et al on behalf of the CIBIS III Investigators. Effect on survival and hospitalization of initiating treatment for chronic heart failure with bisoprolol followed by enalapril, as compared with the opposite consequence: results of the randomized Cardiac Insufficiency Bisoprolol Study (CIBIS) III. *Circulation* 2005;112:2426-35.
167. Poole-Wilson PA, Swedberg K, Cleland JGF, et al. Comparison of carvedilol and metoprolol on clinical outcomes in patients with chronic heart failure in the Carvedilol Or Metoprolol European Trial (COMET): randomized controlled trial. *Lancet* 2003;362:7-13.
168. The Beta-Blocker Evaluation of Survival Trial Investigators. A trial of the beta-blocker bucindolol in patients with advanced chronic heart failure. *N Engl J Med* 2001;344:1659-67.
169. Goldstein S, Fagerberg B, Hjalmarson A et al for the MERIT-HF Study Group. Metoprolol controlled release/extended release in patients with severe heart failure: analysis of the experience in the MERIT-HF study. *J Am Coll Cardiol* 2001;38:932-8.
170. Ghali JK, Piña IL, Gottlieb SS, Deedwania PC, Wikstrand JC, on behalf of the MERIT-HF Study Group. Metoprolol CR/XL in female patients with heart failure: analysis of the experience in metoprolol extended-release randomized intervention trial in heart failure (MERIT-HF). *Circulation* 2002;105:1585-91.
171. Yancy CW, Fowler MB, Colucci WS et al. for the U.S. Carvedilol Heart Failure Study Group. Race and response to adrenergic blockade with carvedilol in patients with chronic heart failure. *N Engl J Med* 2001;344:1358-65.
172. Packer M, Antonopoulos GV, Berlin JA et al. Comparative effects of carvedilol and metoprolol on left ventricular ejection fraction in heart failure: results of a meta-analysis. *Am Heart J* 2001;141:899-907.
173. Adams KF Jr. Which  $\beta$ -blocker for heart failure? *Am Heart J* 2001;141:884-8.
174. Yancy CW. Clinical trials of  $\beta$ -blockers in heart failure: a class review. *Am J Med* 2001;110:7S-10S.
175. Dargie HJ. Beta-blockers in heart failure. *Lancet* 2003;362:2-3.
176. Doughty RN, Rodgers A, Sharpe N, MacMahon S. Effects of beta-blocker therapy on mortality in patients with heart failure: a systematic overview of randomized controlled trials. *Eur Heart J* 1997;18:560-5.
177. Heidenreich PA, Lee TT, Massie BM. Effect of beta-blockade on mortality in patients with heart failure: a meta-analysis of randomized clinical trials. *J Am Coll Cardiol* 1997;30:27-34.
178. Lechat P, Packer M, Chalon S et al. Clinical effects of  $\beta$ -adrenergic blockade in chronic heart failure: a meta-analysis of double-blind, placebo-controlled, randomized trials. *Circulation* 1998; 98: 1184-91.
179. Brophy JM, Joseph L, Rouleau JL.  $\beta$ -blockers in congestive heart failure: a Bayesian meta-analysis. *Ann Intern Med* 2001;134:550-60.
180. Gottlieb SS, Fisher ML, Kjekshus J, et al., on behalf of the MERIT-HF Investigators. Tolerability of  $\beta$ -blocker titration in the metoprolol CR/XL randomized intervention trial in congestive heart failure (MERIT-HF). *Circulation* 2002;105:1182-88.
181. Komajda M, Lutiger B, Madeira H, et al. for the CARMEN investigators and co-ordinators. Tolerability of carvedilol and ACE-Inhibition in mild heart failure. Results of CARMEN (Carvedilol ACE-Inhibitor Remodelling Mild CHF EvaluationN). *Eur J Heart Fail* 2004;6:467-75.
182. Butler J, Khadim G, Belue R, et al. Tolerability to beta-blocker therapy among heart failure patients in clinical practice. *J Card Fail* 2003;9:203-9.
183. Gattis WA, O'Connor CM, Gallup DS, Hasselblad V, Gheorghiade M, IMPACT-HF Investigators and Coordinators. PredischARGE initiation of carvedilol in patients hospitalized for decompensated heart failure: results

- of the Initiation Management Predischarge: Process for Assessment of Carvedilol Therapy in Heart Failure (IMPACT-HF) trial. *J Am Coll Cardiol* 2004;43:1534-41.
184. Ko DT, Hebert PR, Coffey CS, et al. Adverse effects of beta-blocker therapy for patients with heart failure: a quantitative overview of randomized trials. *Arch Intern Med* 2004;164:1389-94.
  185. Nul D, Zambrano C, Diaz A, et al. for the Grupo de Estudio de la Sobrevida en la Insuficiencia Cardiaca en Argentina. Impact of a standardized titration protocol with carvedilol in heart failure: safety, tolerability, and efficacy-a report from the GESICA registry. *Cardiovasc Drugs Ther* 2005;19:125-34.
  186. Jain A, Mills P, Nunn LM, et al. Success of a multidisciplinary heart failure clinic for initiation and up-titration of key therapeutic agents. *Eur J Heart Fail* 2005;7:405-10.
  187. Gullestad L, Wikstrand J, Deedwania P, et al. for the MERIT-HF Study Group. What resting heart rate should one aim for when treating patients with heart failure with a beta-blocker? Experiences from the Metoprolol Controlled Release/Extended Release Randomized Intervention Trial in Chronic Heart Failure (MERIT-HF). *J Am Coll Cardiol* 2005;45:252-9.
  188. Simon T, Mary-Krause M, Funck-Brentano C, Lechat P, Jaillon P. Bisoprolol dose-response relationship in patients with congestive heart failure: a subgroup analysis in the cardiac insufficiency bisoprolol study (CIBIS II). *Eur Heart J* 2003;24:552-9.
  189. Maggioni AP, Sinagra G, Opasich C, et al. Beta blockers in patients with congestive heart failure: guided use in clinical practice Investigators. Treatment of chronic heart failure with beta adrenergic blockade beyond controlled clinical trials: the BRING-UP experience. *Heart* 2003;89:299-305.
  190. Deedwania PC, Giles TD, Klibaner M, et al. for the MERIT-HF Study Group. Efficacy, safety and tolerability of metoprolol CR/XL in patients with diabetes and chronic heart failure: experiences from MERIT-HF. *Am Heart J* 2005;149:159-67.
  191. Riegger GAJ, Bouzo H, Petr P, et al. Improvement in exercise tolerance and symptoms of congestive heart failure during treatment with candesartan cilexetil. *Circulation* 1999;100:2224-30.
  192. Hamroff G, Katz SD, Mancini D, et al. Addition of angiotensin II receptor blockade to maximal angiotensin-converting enzyme inhibition improves exercise capacity in patients with severe congestive heart failure. *Circulation* 1999;99:990-2.
  193. Warner JG Jr., Metzger DC, Kitzman DW, et al. Losartan improves exercise tolerance in patients with diastolic dysfunction and a hypertensive response to exercise. *J Am Coll Cardiol* 1999;33:1567-72.
  194. Blanchet M, Sheppard R, Racine N, et al. Effects of angiotensin-converting enzyme inhibitor plus irbesartan on maximal and submaximal exercise capacity and neurohumoral activation in patients with congestive heart failure. *Am Heart J* 2005;149:e1-7.
  195. O'Meara E, Solomon S, McMurray J, et al. Effect of candesartan on New York Heart Association functional class. *Eur Heart J* 2004;25:1920-6.
  196. Houghton AR, Harrison M, Cowley AJ. Haemodynamic, neurohumoral and exercise effects of losartan vs. captopril in chronic heart failure: Results of an ELITE trial substudy. *Eur J Heart Fail* 1999;1:385-93.
  197. Lang RM, Elkayam U, Yellen LG, et al. Comparative effects of losartan and enalapril on exercise capacity and clinical status in patients with heart failure. *J Am Coll Cardiol* 1997;30:983-91.
  198. Dickstein K, Chang P, Willenheimer R, et al. Comparison of the effects of losartan and enalapril on clinical status and exercise performance in patients with moderate or severe chronic heart failure. *J Am Coll Cardiol* 1995;26:438-45.
  199. Dunselman PH, Replacement of Angiotensin Converting Enzyme Inhibition I. Effects of the replacement of the angiotensin converting enzyme inhibitor enalapril by the angiotensin II receptor blocker telmisartan in patients with congestive heart failure. The replacement of angiotensin converting enzyme inhibition (REPLACE) investigators. *Int J Cardiol* 2001;77:131-8.
  200. Willenheimer R, Helmers C, Pantev E, et al. Safety and efficacy of valsartan versus enalapril in heart failure patients. *Int J Cardiol* 2002;85:261-70.
  201. McKelvie RS, Yusuf S, Pericak D, et al., for the RESOLVD Pilot Study Investigators. Comparison of candesartan, enalapril, and their combination in congestive heart failure: Randomized Evaluation of Strategies for Left Ventricular Dysfunction (RESOLVD) Pilot Study. *Circulation* 1999;100:1056-64.
  202. Pitt B, Poole-Wilson PA, Segal R, et al. Effect of losartan compared with captopril on mortality in patients with symptomatic heart failure: randomized trial: the Losartan Heart Failure Survival Study ELITE II. *Lancet* 2000;355:1582-7.
  203. Pitt B, Segal R, Martinez FA, et al. Randomized trial of losartan versus captopril in patients over 65 with heart failure: Evaluation of Losartan in Elderly Study (ELITE). *Lancet* 1997;349:747-52.
  204. Cohn JN, Tognoni G, for the Valsartan Heart Failure Trial Investigators. A randomized trial of the angiotensin-receptor blocker valsartan in chronic heart failure. *N Engl J Med* 2001;345:1667-75.
  205. Maggioni AP, Anand I, Gottlieb SO, et al on behalf of the Val-HeFT investigators. Effects of valsartan on morbidity and mortality in patients with heart failure not receiving angiotensin-converting enzyme inhibitors. *J Am Coll Cardiol* 2002;40:1414-21.


206. Pfeffer MA, Swedberg K, Granger CB, et al. for the CHARM Investigators and Committees. Effects of candesartan on mortality and morbidity in patients with chronic heart failure: the CHARM-Overall programme. *Lancet* 2003;362:759-66.
207. McMurray JJV, Östergren J, Swedberg K, et al. for the CHARM Investigators and Committees. Effects of candesartan in patients with chronic heart failure and reduced left-ventricular systolic function taking angiotensin converting-enzyme inhibitors: the CHARM-Added trial. *Lancet* 2003;362:767-71.
208. Granger CB, McMurray JJV, Yusuf S, et al. for the CHARM Investigators and Committees. Effects of candesartan in patients with chronic heart failure and reduced left-ventricular systolic function intolerant to angiotensin converting-enzyme inhibitors: the CHARM-Alternative trial. *Lancet* 2003;362:772-6.
209. Young JB, Dunlap ME, Pfeffer MA, et al. Mortality and morbidity reduction with candesartan in patients with chronic heart failure and left ventricular systolic dysfunction: results of the CHARM low-left ventricular ejection fraction trials. *Circulation* 2004;110:2618-26.
210. Lee VC, Rhew DC, Dylan M, et al. Meta-analysis: angiotensin-receptor blockers in chronic heart failure and high-risk acute myocardial infarction. *Ann Intern Med* 2004;141(9):693-704.
211. Jong P, Demers C, McKelvie RS, Liu PP. Angiotensin receptor blockers in heart failure: meta-analysis of randomized controlled trials. *J Am Coll Cardiol* 2002;39:463-70.
212. Dimopoulos K, Salukhe TV, Coats AJ, et al. Meta-analyses of mortality and morbidity effects of an angiotensin receptor blocker in patients with chronic heart failure already receiving an ACE inhibitor (alone or with a beta-blocker). *Int J Cardiol* 2004;93:105-11.
213. Conigliaro RL, Gleason PP. Losartan-induced cough after lisinopril therapy. *Am J Health-Syst Pharm* 1999;56:914-5. Letter.
214. Pylypchuk GB. ACE inhibitor-versus angiotensin II blocker-induced cough and angioedema. *Ann Pharmacother* 1998;32:1060-6.
215. Brown NJ, Ray WA, Snowden M, Griffin MR. Black Americans have an increased rate of angiotensin converting enzyme inhibitor-associated angioedema. *Clin Pharmacol Ther* 1996;60:8-13.
216. van Rijnsoever EW, Kwee-Zuiderwijk WJ, Feenstra J. Angioneurotic edema attributed to the use of losartan. *Arch Intern Med* 1998;158:2063-5.
217. Boxer M. Accupril- and Cozaar-induced angioedema in the same patient (letter). *J Allergy Clin Immunol* 1996;98:471.
218. Acker CG, Greenberg A. Angioedema induced by the angiotensin II blocker losartan (letter). *N Engl J Med* 1995;333:1572.
219. Sharma PK, Yium JJ. Angioedema associated with angiotensin II receptor antagonist losartan. *South Med J* 1997;90:552-3.
220. Frye CB, Pettigrew TJ. Angioedema and photosensitive rash induced by valsartan. *Pharmacotherapy* 1998;18:866-8.
221. Rivera JO. Losartan-induced angioedema. *Ann Pharmacother* 1999;33:933-5.
222. Cha YJ, Pearson VE. Angioedema due to losartan. *Ann Pharmacother* 1999;33:936-8.
223. Rupprecht R, Vente C, Grafe A, Fuchs T. Angioedema due to losartan. *Allergy* 1999;54:81-2.
224. Warner KK, Visconti JA, Tschampel MM. Angiotensin II receptor blockers in patients with ACE inhibitor-induced angioedema. *Ann Pharmacother* 2000;34:526-8.
225. Kymrakis DE, Papadakis CE, Liolios AD, et al. Angiotensin-converting enzyme inhibitors and angiotensin II receptor antagonists. *Arch Otolaryngol Head Neck Surg* 2004;130:1416-9.
226. Cicardi M, Zingale LC, Bergamaschini L, Agostoni A. Angioedema associated with angiotensin-converting enzyme inhibitor use: outcome after switching to a different treatment. *Arch Intern Med* 2004;164:910-3.
227. Abdi R, Dong VM, Lee CJ, Ntoso KA. Angiotensin II receptor blocker-associated angioedema: on the heels of ACE inhibitor angioedema. *Pharmacotherapy*. 2002;22:1173-5.
228. Chiu AG, Krowiak EJ, Deeb ZE. Angioedema associated with angiotensin II receptor antagonists: challenging our knowledge of angioedema and its etiology. *Laryngoscope* 2001;111:1729-31.
229. Irons BK, Kumar A. Valsartan-induced angioedema. *Ann Pharmacother* 2003;37:1024-7.
230. Lo K. Angioedema associated with candesartan. *Pharmacotherapy* 2002; 22:1176-9.
231. Howes LG, Tran D. Can angiotensin receptor antagonists be used safely in patients with previous ACE inhibitor-induced angioedema? *Drug Saf* 2002;25:73-6.
232. Bakris GL, Siomos M, Richardson D, et al. ACE inhibition or angiotensin receptor blockade: impact on potassium in renal failure. VAL-K Study Group. *Kidney Int* 2000;58:2084-92.
233. Cohn JN, Archibald DG, Ziesche S et al. Effect of vasodilator therapy on mortality in chronic congestive heart failure: Results of a Veterans Administration Cooperative Study. *N Engl J Med* 1986;314:1547-52.
234. Taylor AL, Ziesche S, Yancy C, et al. for the African-American Heart Failure Trial Investigators. Combination of isosorbide dinitrate and hydralazine in blacks with heart failure. *N Engl J Med* 2004;351:2049-57.
235. Hood WB Jr, Dans AL, Guyatt GH, Jaeschke R, McMurray JJ. Digitalis for treatment of congestive heart failure in patients in sinus rhythm: a systematic review and meta-analysis. *J Card Fail* 2004;10:155-64.



236. Jaeschke R, Oxman AD, Guyatt GH. To what extent do congestive heart failure patients in sinus rhythm benefit from digoxin therapy? A systematic overview and meta-analysis. *Am J Med* 1990;88:279-86.
237. Packer M, Gheorghiade M, Young JB et al for the RADIANCE Study. Withdrawal of digoxin from patients with chronic heart failure treated with angiotensin-converting enzyme inhibitors. *N Engl J Med* 1993;329:1-7.
238. Uretsky BF, Young JB, Shahidi E et al. Randomized study assessing the effect of digoxin withdrawal in patients with mild to moderate chronic congestive heart failure: results of the PROVED trial. *J Am Coll Cardiol* 1993;22:955-62.
239. Rathore SS, Wang Y, Krumholz HM. Sex-based differences in the effect of digoxin for the treatment of heart failure. *N Engl J Med* 2002;347:1403-11.
240. Eichhorn EJ, Gheorghiade M. Digoxin-new perspective on an old drug. *N Engl J Med* 2002;347:1394-5.
241. Adams KF Jr, Patterson JH, Gattis WA, et al. Relationship of serum digoxin concentration to mortality and morbidity in women in the digitalis investigation group trial: a retrospective analysis. *J Am Coll Cardiol* 2005;46:497-504.
242. Mathew J, Wittes J, McSherry F, et al., Digitalis Investigation Group. Racial differences in outcome and treatment effect in congestive heart failure. *Am Heart J* 2005;150:968-76.
243. Riaz K, Forker AD. Digoxin use in congestive heart failure: current status. *Drugs* 1998;55:747-58.
244. Gheorghiade M, Pitt B. Digitalis Investigation Group (DIG) trial: a stimulus for further research. *Am Heart J* 1997;134:3-12.
245. Mancini DM, Benotti JR, Elkayam U et al, and the PROMISE Investigators & Coordinators. Antiarrhythmic drug use and high serum levels of digoxin are independent adverse prognostic factors in patients with chronic heart failure [abstr]. *Circulation* 1991;84:II-243.
246. Terra SG, Washam JB, Dunham GD, Gattis WA. Therapeutic range of digoxin's efficacy in heart failure: what is the evidence? *Pharmacotherapy* 1999;19:1123-6.
247. Adams KF, Gheorghiade M, Uretsky BF, et al. Clinical benefits of low serum digoxin concentrations in heart failure. *J Am Coll Cardiol* 2002;39:946-953.
248. Rathore SS, Curtis JP, Wang Y, Bristow MR, Krumholz HM. Association of serum digoxin concentration and outcomes in patients with heart failure. *JAMA* 2003;289:871-8.
249. Pitt B, Zannad F, Remme WJ, et al. The effect of spironolactone on morbidity and mortality in patients with severe heart failure: Randomized Aldactone Evaluation Study Investigators. *N Engl J Med* 1999;341:709-17.
250. Pitt B, Remme W, Zannad F, et al, for the Eplerenone Post-Myocardial Infarction Heart Failure Efficacy and Survival Study Investigators. Eplerenone, a selective aldosterone blocker, in patients with left ventricular dysfunction after myocardial infarction. *N Engl J Med* 2003;348:1309-21.
251. Bozkurt B, Agoston I, Knowlton AA. Complications of inappropriate use of spironolactone in heart failure: when an old medicine spirals out of new guidelines. *J Am Coll Cardiol* 2003;41:211-4.
252. The VA Chronic Heart Failure Quality Enhancement Research Initiative. Notice: Safety of spironolactone for heart failure patients. *CHF QUERI News*. February 2000.
253. Tamirisa KP, Aaronson KD, Koelling TM. Spironolactone-induced renal insufficiency and hyperkalemia in patients with heart failure. *Am Heart J* 2004;148:971-8.
254. Sligl W, McAlister FA, Ezekowitz J, Armstrong PW. Usefulness of spironolactone in a specialized heart failure clinic. *Am J Cardiol* 2004;94:443-7.
255. Svensson M, Gustafsson F, Galatius S, Hildebrandt PR, Atar D. How prevalent is hyperkalemia and renal dysfunction during treatment with spironolactone in patients with congestive heart failure? *J Card Fail* 2004;10:297-303.
256. Shah KB, Rao K, Sawyer R, Gottlieb SS. The adequacy of laboratory monitoring in patients treated with spironolactone for congestive heart failure. *J Am Coll Cardiol* 2005;46:845-9.
257. Saito M, Takada M, Hirooka K, Isobe F, Yasumura Y. Serum concentration of potassium in chronic heart failure patients administered spironolactone plus furosemide and either enalapril maleate, losartan potassium or candesartan cilexetil. *J Clin Pharm Ther* 2005;30: 603-10.
258. Juurlink DN, Mamdani MM, Lee DS, et al. Rates of hyperkalemia after publication of the Randomized Aldactone Evaluation Study. *N Engl J Med* 2004;351:543-51.
259. Schepkens H, Vanholder R, Billioud JM, Lameire N. Life-threatening hyperkalemia during combined therapy with angiotensin-converting enzyme inhibitors and spironolactone: an analysis of 25 cases. *Am J Med* 2001;110:438-41.
260. Berry C, McMurray JJV. Serious adverse events experienced by patients with chronic heart failure taking spironolactone. *Heart* 2001;85:e8-e9.
261. Oddone EZ, Weinberger M, Horner M, et al. Classifying general medicine readmissions. Are they preventable? Veterans Affairs Cooperative Studies in Health Services Group on Primary Care and Hospital Readmissions. *J Gen Intern Med* 1996;11:597-607.
262. Roe CM, Motheral BR, Teitelbaum F, Rich MW. Angiotensin-converting enzyme inhibitor compliance and dosing among patients with heart failure. *Am Heart J* 1999;138:818-25.

263. Cline CM, Bjorck-Linne BY, Willenheimer RB, Erhardt LR. Non-compliance and knowledge of prescribed medication in elderly patients with heart failure. *Eur J Heart Fail* 1999;1:145-9.
264. Ghali JK, Kadakia S, Cooper R, et al. Precipitating factors leading to decompensation of heart failure. *Arch Intern Med* 1988;148:2013-6.
265. Rich MW. Multidisciplinary interventions for the management of heart failure: where do we stand? *Am Heart J* 1999;138:599-601.
266. Gattis WA, Hasselblad V, Whellan DJ, O'Connor CM. Reduction in heart failure events by the addition of a clinical pharmacist to the heart failure management team: results of the Pharmacist in Heart Failure Assessment Recommendation and Monitoring (PHARM) Study. *Arch Intern Med* 1999;159:1939-45.
267. Whellan DJ, Gauden L, Gattis WA, et al. The benefit of implementing a heart failure disease management program. *Arch Intern Med* 2001;161:2223-8.
268. West JA, Miller NH, Parker KM, et al. A comprehensive management system for heart failure improves clinical outcomes and reduces medical resource utilization. *Am J Cardiol* 1997;79:58-63.
269. Shah NB, Der E, Ruggerio C, Heidenreich PA, Massie BM. Prevention of hospitalizations for heart failure with an interactive home monitoring program. *Am Heart J* 1998;135:373-8.
270. Rich MW, Nease RF. Cost-effectiveness analysis in clinical practice: the case of heart failure. *Arch Intern Med* 1999;159:1690-1700.
271. Ramahi TM, Longo MD, Rohlf K, Sheynberg N. Effect of heart failure program on cardiovascular utilization and dosage in patients with chronic heart failure. *Clin Cardiol* 2000;23:909-14.
272. Philbin EF. Comprehensive multidisciplinary programs for the management of patients with congestive heart failure. *J Gen Intern Med* 1999;14:130-5.
273. Rich MW, Beckham V, Wittenberg C, et al. A multidisciplinary intervention to prevent the readmission of elderly patients with congestive heart failure. *N Engl J Med* 1995;333:1190-5.
274. Rich MW. Heart failure disease management: a critical review. *J Card Fail* 1999;5:64-75.
275. McAlister FA, Stewart S, Ferrua S, McMurray JJ. Multidisciplinary strategies for the management of heart failure patients at high risk for admission: a systematic review of randomized trials. *J Am Coll Cardiol* 2004;44:810-9.
276. Hebert KA, Horswell RL, Dy S, et al. Mortality benefit of a comprehensive heart failure disease management program in indigent patients. *Am Heart J* 2006;151:478-83.
277. Akosah KO, Schaper AM, Haus LM, et al. Improving outcomes in heart failure in the community: long-term survival benefit of a disease-management program. *Chest* 2005;127:2042-8.
278. Hershberger RE, Nauman DJ, Byrkit J, et al. Prospective evaluation of an outpatient heart failure disease management program designed for primary care: the Oregon model. *J Card Fail* 2005;11:293-8.
279. Franciosa JA, Massie BM, Lukas MA, et al., COHERE Participant Physicians. Beta-blocker therapy for heart failure outside the clinical trial setting: findings of a community-based registry. *Am Heart J* 2004;148:718-26.
280. Ledwidge M, Barry M, Cahill J, et al. Is multidisciplinary care of heart failure cost-beneficial when combined with optimal medical care? *Eur J Heart Fail* 2003;5:381-9.
281. Ansari M, Alexander M, Tutar A, Bello D, Massie BM. Cardiology participation improves outcomes in patients with new-onset heart failure in the outpatient setting. *J Am Coll Cardiol* 2003;41:62-8.
282. Gonseth J, Guallar-Castillon P, Banegas JR, Rodriguez-Artalejo F. The effectiveness of disease management programmes in reducing hospital re-admission in older patients with heart failure: a systematic review and meta-analysis of published reports. *Eur Heart J* 2004;25:1570-95.

**Figure: VA Utilization of Medications for HF by Drug Class**



Denominator=patients with at least one diagnosis of heart failure (i.e., at least one inpatient primary diagnosis or any outpatient diagnosis within 24 months prior to the end date of the fiscal year) and at least one active prescription during the fiscal year  
 Active prescription=at least 90 days of therapy during the given year  
 Combination therapy=prescriptions for both medications overlapping  $\geq 50\%$   
 AA=aldosterone antagonist; ACEI=angiotensin-converting enzyme inhibitor; ARB=angiotensin II receptor antagonist; BB=beta-adrenergic blocker; ACEI/ARB+BB=ACEI or ARB in combination with BB; H-N=hydralazine in combination with a nitrate

## Appendix A: Medications Commonly Used for the Management of HF<sup>a,c</sup>

In general, patients should be titrated to target doses as used in randomized controlled trials (refer to Appendix C), or highest tolerated dose

DRUG	INITIAL DOSE	MAXIMUM DOSE <sup>d</sup>	COMMENTS/CAUTIONS
<i>Diuretics</i>			
Furosemide	20 to 40 mg given once or twice daily	600 mg in divided doses	Monitor serum K <sup>+</sup> at 1 to 2 weeks after initiating therapy or changing dose, then every few months; more frequently if patient is also on digoxin or has demonstrated hypokalemia Add potassium supplement or low dose potassium-sparing diuretic <sup>e</sup> if the patient becomes hypokalemic (serum K <sup>+</sup> < 4.0 mEq/L) Use cautiously in poorly controlled DM, symptomatic BPH, or in patients with increased risk of volume depletion Furosemide usually administered once daily unless higher doses (e.g., > 160mg/d) are needed, then more frequent daily dosing should be considered Ethacrynic acid may be used in patients with sulfonamide sensitivity Thiazides lose effectiveness in patients with CrCl < 40 mL/min Reserve indapamide for patients with CrCl < 25 mL/min Reserve metolazone for intermittent use as an adjunct to loop diuretics for diuresis in patients with HF or in patients with CrCl < 25 mL/min; thiazide/loop combinations are also effective and are less expensive
Bumetanide	0.5 to 1.0 mg given once or twice daily	10mg in divided doses	
Ethacrynic acid	50 mg once or twice daily	400mg in divided doses	
Torsemide	10 to 20 mg once daily	200 mg once daily	
Hydrochlorothiazide	25 mg given once or twice daily	200 mg in divided doses	
Chlorthalidone	12.5 to 25 mg once daily	100 mg once daily	
Indapamide	2.5 mg once daily	5 mg once daily	
Metolazone <sup>f,g</sup> Zaroxolyn®	2.5 to 5 mg once daily	20 mg once daily	
Mykrox®	0.5 mg once daily	1 mg once daily	
<i>Angiotensin Converting Enzyme Inhibitor</i>			
Captopril <sup>h</sup>	6.25 to 12.5 mg three times daily	50 mg three times daily <sup>d</sup>	Start with lower or less frequent doses in patients with renal insufficiency; use with caution in patients with renal artery stenosis Should not be used if K <sup>+</sup> > 5.5 mEq/L that cannot be reduced Due to the potential risk for fetal abnormalities in patients taking ACEIs during pregnancy, it is recommended that therapy be discontinued as soon as a woman becomes pregnant. Alternate therapy should be considered. ACEIs should only be prescribed in pregnant women when the benefit clearly outweighs the potential risk for fetal abnormalities
Enalapril	2.5 mg twice daily	10 to 20mg twice daily <sup>d</sup>	
Fosinopril	5 to 10 mg once daily	20 to 40 mg once daily <sup>d</sup>	
Lisinopril	2.5 to 5 mg once daily	20 to 40 mg once daily <sup>d</sup>	
<i>Beta-adrenergic blockers</i>			
Metoprolol succinate (XL)	12.5 to 25 mg once daily; double dose every 2 weeks to target dose	200 mg once daily <sup>d</sup> (or highest dose tolerated)	Low initial doses should be implemented; use slow gradual increases in the dosage Effects are generally seen in 3-12 months Carvedilol should be given with food to reduce the incidence of orthostatic hypotension Consider separating the ACEI, adjusting dose of diuretic, or temporary ACEI dose reduction if dizziness occurs Should not be abruptly discontinued
Bisoprolol	1.25 mg once daily; increase by 1.25 mg every week until 5 mg once daily, then increase by 2.5 mg every 4 weeks to target dose	10 mg once daily <sup>d</sup>	
Carvedilol (alpha & beta antagonist)	3.125 mg twice daily; titrate at minimum of every 2 weeks to target dose	25 mg twice daily <sup>d</sup> (should be titrated as tolerated to 50mg twice daily if ≥ 85kg)	
<i>Angiotensin II Receptor Antagonists</i>			
Candesartan	4 to 8 mg once daily	32 mg once daily <sup>d</sup>	Contraindicated in 2 <sup>nd</sup> and 3 <sup>rd</sup> trimesters pregnancy due to potential neonatal/fetal

Valsartan	40 to 80 mg divided twice daily	320 mg divided twice daily <sup>d</sup>	morbidity and death Use with caution in patients with renal artery stenosis Should not be used if K <sup>+</sup> > 5.5 mEq/L that cannot be reduced Consider lower doses in patients with intravascular volume depletion
<i>Hydralazine in Combination with a Nitrate</i>			
Hydralazine	112.5 to 150 mg divided three times daily	225 to 300 mg divided three times daily <sup>d</sup>	Hydralazine: Adverse effects include dizziness, headache, lupus-like syndrome, nausea, tachycardia, postural hypotension Advise patient to take with food
Isosorbide dinitrate	30 to 60 mg divided three times daily	120 to 160 mg divided three times daily <sup>d</sup>	ISDN: Adverse effects include flushing, headache, postural hypotension, rash May cause an increase in ocular pressure; caution with presence of glaucoma
<i>Digoxin</i>			
Digoxin	0.125 to 0.25 micrograms once daily	Usually 0.25 micrograms once daily (0.375 to 0.5 micrograms once daily may be used rarely)	Initiate therapy with 0.125 micrograms once daily (or every other day) in patients > 70, with impaired kidney function, or with a low lean body mass Lower trough serum digoxin concentrations may be preferable (i.e., 0.5-0.9ng/ml), and should not exceed 1.1ng/ml Signs of toxicity include confusion, nausea, vomiting, abdominal pain, diarrhea, anorexia, fatigue, arrhythmias, visual disturbances
<i>Aldosterone Antagonists</i>			
Spironolactone	12.5 to 25 mg once daily	25 mg once daily <sup>d</sup>	If CrCl < 50 ml/min, initial dose should be 12.5 mg once daily or 25 mg every other day for spironolactone and 25 mg once daily of eplerenone; not recommended if CrCl < 30 ml/min Monitor closely for hyperkalemia (should not be used if baseline K <sup>+</sup> > 5.0 mEq/L) or renal dysfunction; patients may also experience gynecomastia, especially with spironolactone
Eplerenone	25 mg once daily	50 mg once daily <sup>d</sup>	

<sup>a</sup> Adapted from McEvoy GK, ed. American Hospital Formulary Service Drug Information, Bethesda, MD: American Society of Health-System Pharmacists, Inc., 2006.

<sup>b</sup> Adapted from Hebel SK ed. Drug Facts and Comparisons, St. Louis, Missouri: Facts and Comparisons Inc., May 2006.

<sup>c</sup> Hunt SA, Abraham WT, Chin MH, et al. ACC/AHA 2005 guideline update for the diagnosis and management of chronic heart failure in the adult: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Update the 2001 Guidelines for the Evaluation and Management of Heart Failure). American College of Cardiology Web site. Available at: <http://www.acc.org/clinical/guidelines/failure/index.pdf>.

<sup>d</sup> Target doses (also refer to Appendix C for target doses in randomized controlled trials)

<sup>e</sup> Unless patients have persistent hypokalemia or are being treated with low dose spironolactone for severe HF (refer to Annotation M), potassium-sparing diuretics should not be used in combination with ACEI (refer to Appendix B for common diuretic drug interactions)

<sup>f</sup> The brand names of metolazone are not bioequivalent, therefore doses vary

<sup>g</sup> Intermittent use recommended once the response of the patient is stabilized

<sup>h</sup> One hour before meals, on an empty stomach

<sup>i</sup> Also available as a fixed-dose combination product ISDN 20mg/hydralazine 37.5 mg, one to two tablets three times daily; ISMN once daily has also been used in place of ISDN; hydralazine dosing variable, total daily dose may also be divided two to four times daily

## Appendix B. Drug Interactions with Agents Used in HF<sup>a,b</sup>

DRUG CLASS	INTERACTING DRUG	DESCRIPTION
<b>DIURETICS</b>		
	ACEI	↑ hypotensive effect in the presence of intensive diuretic therapy due to sodium depletion and hypovolemia; effects of loop diuretics may be ↓ by inhibition of angiotensin II production [significance=3]
	Bile Acid Resins	↓ absorption of all diuretics; bile acid resin should be taken at least 2 hours after diuretic [significance=2]
	Digoxin	Loop and thiazide diuretics may induce hypokalemia which may ↑ risk of digitalis toxicity [significance=2]
	Dofetilide	Risk of torsade de pointes increased with hypokalemia [significance=1]
	Lithium	With thiazides, a compensatory ↑ in proximal tubule reabsorption of sodium occurs, which results in ↑ lithium reabsorption (reduce lithium dose by 50%); ↑ plasma lithium concentrations may also occur with loop diuretics [significance=2]
	Oral hypoglycemics	Thiazides may ↓ hypoglycemic effects of sulfonylureas possibly due to ↓ insulin sensitivity, ↓ insulin secretion or ↓ in K <sup>+</sup> [significance=2]
	K <sup>+</sup> preparations, ACEI, ARBs	K <sup>+</sup> -sparing diuretics used concomitantly may ↑ K <sup>+</sup> serum levels [significance=3]
<b>ACEIs</b>		
	Allopurinol	Isolated case reports with allopurinol and captopril or enalapril may have caused predisposition to hypersensitivity reactions (e.g., Stevens Johnson Syndrome, anaphylaxis, skin eruptions, fever, arthralgias) [significance=4]
	Lithium	Potential for ↑ serum lithium levels and resultant toxicity [significance=2]
	NSAIDs	NSAIDs ↓ antihypertensive effects due to inhibition of PG synthesis resulting in ↓ GFR, ↓ sodium and water excretion, and vasoconstriction [significance=2]
	K <sup>+</sup> preparations K <sup>+</sup> -sparing diuretics	Concomitant therapy may ↑ K <sup>+</sup> serum levels [significance=1]
<b>ANGIOTENSIN II RECEPTOR ANTAGONISTS</b>		
	Digoxin	See digoxin for description of drug interaction [significance=4]
	Lithium	Angiotensin II receptor antagonists may ↓ lithium renal secretion and ↑ serum lithium levels [significance=2]
	K <sup>+</sup> preparations K <sup>+</sup> -sparing diuretics	Concomitant therapy may ↑ K <sup>+</sup> serum levels [significance=1]

BETA-ADRENERGIC BLOCKERS		
	Cimetidine	Hypotension and bradycardia have been reported with propranolol and metoprolol when used with cimetidine due to ↑ serum levels of beta-blockers that undergo hepatic metabolism [significance=2]
	Diltiazem <b>Verapamil</b> <b>Epinephrine</b>	Combination may potentiate the pharmacologic effects of beta-blockers; additive effects on cardiac conduction [significance=2] Noncardioselective agents may ↑ the pressor response resulting in ↑ in HTN/bradycardia [significance=1]
	Lidocaine	↑ toxicity due to reduced hepatic metabolism of lidocaine [significance=2]
	NSAIDs	NSAIDs ↓ antihypertensive effect due to inhibition of PG synthesis resulting in ↓ GFR, ↓ sodium and water excretion, and vasoconstriction [significance=2]
	<b>Neuroleptics</b>	Some beta-blockers and neuroleptics (chlorpromazine/ thioridazine) may ↑ the plasma concentrations of one another; monitor for enhanced effects of both drugs; concomitant use of thioridazine and propranolol or pindolol is contraindicated [significance=1]
	Oral hypoglycemics	↓ hypoglycemic action may occur, may also mask symptoms of hypoglycemia (more likely with nonselective beta-blocker); clinical significance is unclear [significance=5]
	Prazosin	Potential for ↑ postural hypotension [significance=2]
	Propafenone	↑ hypotensive effect has been seen with propranolol and metoprolol due to inhibition of metabolic clearance; HF and nightmares have been reported [significance=2]
	Rifampin	May enhance the hepatic metabolism of propranolol and metoprolol; enzyme induction effect may resolve after a 3-4 week washout period [significance=2]
	Theophylline	↑ serum concentration in a dose-dependent manner has been seen with propranolol [significance=2]
CALCIUM CHANNEL BLOCKERS		
	Beta-blockers	See beta-blockers for description of drug interaction [significance=2]
	Carbamazepine	↑ toxicity has been noted with verapamil and diltiazem due to ↓ metabolism of carbamazepine; felodipine bioavailability may be ↓, making it difficult to achieve therapeutic felodipine concentrations [significance=2]
	Cyclosporine	Blood concentrations have ↑ with verapamil, diltiazem and nifedipine; renal toxicity has been reported [significance=2]
	<b>Digoxin</b>	Verapamil, diltiazem, bepridil, and nisoldipine have ↑ digoxin levels by 20-70% [significance=1]
	Lithium	Combination with verapamil or diltiazem may result in neurotoxicity that may occur without attendant ↑ in serum level [significance=4]
	HMG-CoA reductase inhibitors	Diltiazem produces marked ↑ lovastatin and simvastatin concentrations through inhibition of CYP3A4, therefore potential for ↑ toxicity (rhabdomyolysis reported with atorvastatin and simvastatin in combination with diltiazem); ↑ concentration of simvastatin seen with concomitant verapamil [significance=2]
	<b>Quinidine</b>	Verapamil inhibits metabolism of quinidine leading to ↑ toxicity [significance=1]; nifedipine appears to ↓ blood concentrations [significance=4]
	Theophylline	Inhibition of hepatic metabolism with verapamil and diltiazem may lead to ↑ serum levels [significance=4]



DIGOXIN		
	<b>Amiodarone</b>	↑ serum digoxin concentrations; may need to decrease digoxin dose by ~ 50%; monitor for digoxin toxicity (i.e., anorexia, nausea, fatigue, vomiting, diarrhea, visual disturbances, confusion, ventricular tachycardia) [significance=1]
	Beta-blockers	Carvedilol may ↑ serum digoxin concentrations; potential for synergistic bradycardia with propranolol [significance=2]
	<b>Cyclosporine</b>	↑ serum digoxin concentrations; may need to discontinue digoxin or ↓ dose when treatment resumed; monitor for toxicity (i.e. anorexia, nausea, vomiting, diarrhea, fatigue, visual disturbances, confusion, and ventricular tachycardia) [significance=1]
	<b>Diuretics</b>	↑ risk of digitalis toxicity due to diuretic induced hypokalemia [significance=1]
	<b>Quinidine</b>	↑ serum digoxin concentrations; may need dose ↓ ~ 50%; monitor for toxicity (i.e. anorexia, nausea, vomiting, diarrhea, fatigue, visual disturbances, confusion, and ventricular tachycardia) [significance=1]
	Spironolactone	Renal excretion of digoxin may be reduced; false increases in plasma digoxin concentrations may occur depending on the assay method used [significance=2]
	Telmisartan	May increase digoxin peak plasma concentrations (49%) and in trough concentrations (20%); monitor digoxin levels when starting, adjusting, or discontinuing therapy with telmisartan [significance=4]
	<b>Verapamil</b> (see also Calcium Channel Blockers)	↑ digoxin serum concentrations on average ~70%; dose related; may need to ↓ dose be at least 50%; monitor for toxicity (i.e. anorexia, nausea, vomiting, diarrhea, fatigue, visual disturbances, confusion, and ventricular tachycardia) [significance=1]
SPIRONOLACTONE		
	Digoxin	See digoxin for description of drug interaction [significance=2]
	<b>Mitotane</b>	Spironolactone may antagonize the activity of mitotane; avoid concomitant use [significance=4]
	<b>Potassium, other potassium-sparing diuretics, ACEIs, ARBs</b>	Coadministration may result in hyperkalemia [significance=1]
VASODILATORS		
Hydralazine	Beta-blockers	Serum levels of propranolol or metoprolol may be ↑ with hydralazine use; clinical effects may be enhanced [significance=2]
Nitrates	<b>PDE 5 inhibitors</b>	Sildenafil, tadalafil, and vardenafil potentiate the hypotensive effects of nitrates, severe hypotension may occur; concomitant use is contraindicated [significance=1]

<sup>a</sup>Wickersham RM, et al. eds. Drug Facts and Comparisons, St. Louis: Wolters Kluwer Health, Inc., 2006.

<sup>b</sup> Online Facts and Comparisons 4.0 <http://online.factsandcomparisons.com/> Accessed August 2006.

**Significance:** 1=potentially severe or life-threatening; interaction suspected, established or probable in well controlled studies; 2=may cause deterioration in patient's clinical status; interaction suspected, established or probable in well controlled studies; 3=causes minor effects; interaction suspected, established or probable in well controlled studies; 4=may cause moderate to major effects, very limited data; 5=minor to major effects, interaction is unlikely or not good evidence of an altered clinical effect; **Bold**=major drug interaction

AUC=area under the curve; CV=cardiovascular; CYP=cytochrome P-450 enzyme system; GFR=glomerular filtration rate; PDE=phosphodiesterase; PG=prostaglandin

## Appendix C: Long-term, Randomized, Controlled, Outcome Trials in Systolic HF by Drug Class

Angiotensin Converting Enzyme Inhibitors																			
Trial	Patient Population	N	Treatment	Duration	Results	Study Conclusions	Quality Rating												
<b>SOLVD<sup>120</sup></b> <b>1991</b> MC, R, DB (vs. placebo) U.S., Canada, Belgium  Supported by NHLBI; Merck Sharpe and Dohme provided drug/placebo	NYHA I (11%), II (57%), III (30%), IV (2%) Mean EF 24.8%	2569	Enalapril 10 mg twice daily (target dose) vs. Placebo  <u>HF therapy</u> BB: 8% Digitalis: 67% Diuretics: 86% Vasodilators (any): 51%	Average 41.4 months	<b>Primary Endpoint:</b> Total mortality (16% ↓ with enalapril 95% CI 0.05-0.26; ARR 4.6%, NNT 22) <table><tr><th>Endpoint</th><th>Enalapril (N=1285)</th><th>Placebo (N=1284)</th><th>p value</th></tr><tr><td>Primary</td><td>452 (35.2%)</td><td>510 (39.7%)</td><td>&lt;0.0036</td></tr><tr><td>Death or HF hosp</td><td>613 (47.7%)</td><td>736 (57.3%)</td><td>&lt;0.0001</td></tr></table> <b>Target dose on monotherapy:</b> Bisoprolol (65%) vs. enalapril (84%) <b>Mean dose:</b> 16.6 mg per day (patients taking study drug); 11.2 mg per day (all randomized patients)	Endpoint	Enalapril (N=1285)	Placebo (N=1284)	p value	Primary	452 (35.2%)	510 (39.7%)	<0.0036	Death or HF hosp	613 (47.7%)	736 (57.3%)	<0.0001	Enalapril significantly reduced mortality and HF hospitalizations in patients with HF	Good
Endpoint	Enalapril (N=1285)	Placebo (N=1284)	p value																
Primary	452 (35.2%)	510 (39.7%)	<0.0036																
Death or HF hosp	613 (47.7%)	736 (57.3%)	<0.0001																
<b>CONSENSUS<sup>122</sup></b> <b>1987</b> MC, R, DB, PG (vs. placebo) Scandinavia  Supported by Merck Sharpe and Dohme	All patients with NYHA IV at randomization	253	Enalapril 20 mg twice daily (maximum dose) vs. Placebo  <u>HF therapy</u> BB: 3% Digitalis: 93% Furosemide: 98% Hydralazine: 2% ISDN: 46% Spironolactone: 53%	Average 188 days	<b>Primary Endpoint:</b> All-cause mortality at 6 months (40% ↓ with enalapril; ARR 17.7%, NNT 6) <table><tr><th>Endpoint</th><th>Enalapril (N=127)</th><th>Placebo (N=126)</th><th>p value</th></tr><tr><td>Primary</td><td>33 (26%)</td><td>55 (44%)</td><td>0.002</td></tr><tr><td>Total mortality</td><td>50 (39%)</td><td>68 (54%)</td><td>0.003</td></tr></table> <b>Maximum dose:</b> Enalapril (28 patients) vs. placebo (57 patients) <b>Mean dose:</b> 18.4 mg per day	Endpoint	Enalapril (N=127)	Placebo (N=126)	p value	Primary	33 (26%)	55 (44%)	0.002	Total mortality	50 (39%)	68 (54%)	0.003	Enalapril reduced mortality in patients with severe HF on conventional therapy (i.e., diuretics and digitalis)	Good
Endpoint	Enalapril (N=127)	Placebo (N=126)	p value																
Primary	33 (26%)	55 (44%)	0.002																
Total mortality	50 (39%)	68 (54%)	0.003																

ARR=absolute risk reduction; CI=confidence interval; DB=double-blind; EF=ejection fraction; HF=heart failure; hosp=hospitalizations; ISDN: isosorbide dinitrate; N=number of patients; NNT=number needed to treat; NYHA=New York Heart Association; PG=parallel group; R=randomized

Beta-Adrenergic Blockers																							
Trial	Patient Population	N	Treatment	Duration	Results	Study Conclusions	Quality Rating																
<b>CIBIS III<sup>166</sup></b> <b>2005</b> MC, PROBE (BB 1 <sup>st</sup> vs. ACEI 1 <sup>st</sup> ) Europe, Australia, Tunisia  Supported by Merck KGaA	NYHA II (49%), III (51%) Mean EF 28.8%	1010	(Initial monotherapy X 6 months) Bisoprolol 10 mg once daily (target dose) vs. (Initial monotherapy X 6 months) Enalapril 10 mg twice daily (target dose) Followed by combination therapy X 6 to 24 months  <u>HF therapy</u> Cardiac glycoside: 32% Diuretics: 84% Aldosterone antagonist: 13%	Mean 1.22 yrs	<b>Primary Endpoints:</b> Combined all-cause mortality or all-cause hosp (per protocol analysis bisoprolol 1 <sup>st</sup> vs. enalapril 1 <sup>st</sup> HR 0.97 95% CI 0.78-1.21; ITT 178 (35.2%) vs. 186 (36.8%) HR 0.94 95% CI 0.77-1.16; p=0.019 <sup>a</sup> ) <table><tr><th>Endpoint</th><th>Bisoprolol 1st (N=505<sup>b</sup>)</th><th>Enalapril 1st (N=505<sup>b</sup>)</th><th>p value</th></tr><tr><td>Primary</td><td>163 (32.4%)</td><td>165 (33.1%)</td><td>0.046<sup>a</sup></td></tr><tr><td>CV death<sup>c</sup></td><td>55 (NR)</td><td>56 (NR)</td><td>0.86</td></tr><tr><td>HF hosp<sup>d</sup></td><td>63 (NR)</td><td>51 (NR)</td><td>0.23</td></tr></table> <sup>a</sup> test for noninferiority; <sup>b</sup> N for ITT; <sup>c</sup> IHR 0.97; <sup>d</sup> HR 1.25 <b>Target dose on monotherapy:</b> Bisoprolol (65%) vs. enalapril (84%)	Endpoint	Bisoprolol 1st (N=505 <sup>b</sup> )	Enalapril 1st (N=505 <sup>b</sup> )	p value	Primary	163 (32.4%)	165 (33.1%)	0.046 <sup>a</sup>	CV death <sup>c</sup>	55 (NR)	56 (NR)	0.86	HF hosp <sup>d</sup>	63 (NR)	51 (NR)	0.23	Bisoprolol 1 <sup>st</sup> noninferior to enalapril 1 <sup>st</sup> in ITT analysis, but not by per-protocol analysis; initial therapy with bisoprolol may be as safe and efficacious as starting with enalapril	Fair
Endpoint	Bisoprolol 1st (N=505 <sup>b</sup> )	Enalapril 1st (N=505 <sup>b</sup> )	p value																				
Primary	163 (32.4%)	165 (33.1%)	0.046 <sup>a</sup>																				
CV death <sup>c</sup>	55 (NR)	56 (NR)	0.86																				
HF hosp <sup>d</sup>	63 (NR)	51 (NR)	0.23																				
<b>COMET<sup>167</sup></b> <b>2003</b> MC, R, DB, PG (BB vs. BB) Europe  Supported by F Hoffmann La Roche and GlaxoSmithKline	NYHA II (48%), III (48%), IV (4%) Mean EF 26%	3029	Carvedilol 25 mg twice daily (target dose) vs. Metoprolol IR 50 mg twice daily (target dose)  <u>HF therapy</u> ACEI: 91% ARB: 7% Digoxin: 59% Diuretics: 99% Aldosterone antagonist: 11%	Mean 58 months	<b>Primary Endpoints:</b> 1) All-cause mortality (↓ with carvedilol vs. metoprolol; HR 0.83 95% CI 0.74-0.93; ARR 5.6%, NNT 18); and 2) Composite all-cause mortality or all-cause admission (HR 0.94 95% CI 0.86-1.02) <table><tr><th>Endpoint</th><th>Carvedilol (N=1511)</th><th>Metoprolol (N=1518)</th><th>p value</th></tr><tr><td>Primary1</td><td>512 (33.9%)</td><td>600 (39.5%)</td><td>0.017</td></tr><tr><td>Primary2</td><td>1116 (73.9%)</td><td>1160 (76.4%)</td><td>0.122</td></tr></table> <b>Target dose:</b> Carvedilol (75%) vs. metoprolol IR (78%) <b>Mean dose:</b> Carvedilol (41.8 ± 14.6 mg per day); metoprolol IR (85 ± 28.9 mg per day)	Endpoint	Carvedilol (N=1511)	Metoprolol (N=1518)	p value	Primary1	512 (33.9%)	600 (39.5%)	0.017	Primary2	1116 (73.9%)	1160 (76.4%)	0.122	Carvedilol had a greater benefit on survival compared to metoprolol IR in patients with chronic HF on standard therapy (i.e., diuretics plus ACEI)	Fair				
Endpoint	Carvedilol (N=1511)	Metoprolol (N=1518)	p value																				
Primary1	512 (33.9%)	600 (39.5%)	0.017																				
Primary2	1116 (73.9%)	1160 (76.4%)	0.122																				
<b>COPERNICUS<sup>163</sup></b> <b>2001</b> MC, R, DB (vs. placebo) U.S., Canada, Mexico, Europe, S. America, Israel, S. Africa, Australia  Supported by SmithKline Beecham and Boehringer-Mannheim	Severe HF (≥ 2 months dyspnea or fatigue at rest or minimal exertion, EF < 25%) Mean EF 19.9%	2289	Carvedilol 25 mg twice daily (target dose) vs. Placebo  <u>HF therapy</u> ACEI or ARB: 97% Digoxin: 66% Diuretics: 99% Spironolactone: 20%	Mean 10.4 months (stopped early due to improved survival)	<b>Primary Endpoint:</b> All-cause mortality (35% ↓ with carvedilol; 95% CI 0.19-0.48; ARR 5.5%, NNT 18) <table><tr><th>Endpoint</th><th>Carvedilol (N=1156)</th><th>Placebo (N=1133)</th><th>p value</th></tr><tr><td>Primary</td><td>130 (11.3%)</td><td>190 (16.8%)</td><td>0.0014</td></tr><tr><td>Death or hosp</td><td>425 (36.8%)</td><td>507 (44.8%)</td><td>&lt;0.001</td></tr></table> <b>Target dose:</b> Carvedilol (65.1%) vs. placebo (78.2%) at 4 months <b>Mean dose:</b> 37 mg per day	Endpoint	Carvedilol (N=1156)	Placebo (N=1133)	p value	Primary	130 (11.3%)	190 (16.8%)	0.0014	Death or hosp	425 (36.8%)	507 (44.8%)	<0.001	Carvedilol reduced the rate of death in patients with severe HF on conventional therapy (i.e., diuretics plus ACEI or ARB)	Good				
Endpoint	Carvedilol (N=1156)	Placebo (N=1133)	p value																				
Primary	130 (11.3%)	190 (16.8%)	0.0014																				
Death or hosp	425 (36.8%)	507 (44.8%)	<0.001																				

Beta-Adrenergic Blockers (continued)												
Trial	Patient Population	N	Treatment	Duration	Results	Study Conclusions	Quality Rating					
<b>MERIT-HF<sup>159</sup> 1999</b> MC, R, DB (vs. placebo) U.S., Europe   Supported by Astra Hässle AB	NYHA II (41%), III (56%), IV (3.4%) HF Mean EF 28%	3991	Metoprolol XL 200 mg once daily (target dose) vs. Placebo  <u>HF therapy</u> ACEI: 90% ARB: 7% Digoxin: 64% Diuretics: 90%	Mean 1 yr (terminated early due to survival benefit)	<b>Primary Endpoints:</b> 1) All-cause mortality (↓ with metoprolol XL; RR 0.66 95% CI 0.53-0.81; ARR 3.6%, NNT 28); and 2) Combined all-cause mortality and all-cause hosp admissions (NR)	Metoprolol XL significantly improved survival in patients with symptomatic HF on standard therapy for HF (i.e., diuretics plus ACEI)	Good					
					<table><tr><th>Endpoint</th><th>Metoprolol XL (N=1990)</th><th>Placebo (N=2001)</th><th>p value</th></tr><tr><td>Primary<sup>1</sup></td><td>145 (7.3%)</td><td>217 (10.9%)</td><td>0.00009</td></tr><tr><td>CV death</td><td>128 (6.4%)</td><td>203 (10.2%)</td><td>0.00003</td></tr></table> <b>Target dose:</b> Metoprolol XL (64%) vs. placebo (82%) <b>Mean dose:</b> 159 mg			Endpoint	Metoprolol XL (N=1990)	Placebo (N=2001)	p value	Primary <sup>1</sup>
Endpoint	Metoprolol XL (N=1990)	Placebo (N=2001)	p value									
Primary <sup>1</sup>	145 (7.3%)	217 (10.9%)	0.00009									
CV death	128 (6.4%)	203 (10.2%)	0.00003									
<b>CIBIS II<sup>160</sup> 1999</b> MC, R, DB (vs. placebo) Europe   Supported by E Merck	NYHA III (83%), IV (17%) Mean EF 27.5%	2647	Bisoprolol 10 mg once daily (target dose) vs. Placebo  <u>HF therapy</u> ACEI: 96% Digoxin: 52% Diuretics: 99%	Mean 1.3 yrs (stopped early due to improved survival)	<b>Primary Endpoint:</b> All-cause mortality (↓ with bisoprolol; HR 0.66 95% CI 0.54-0.81; ARR 5.5%, NNT 18)	Bisoprolol significantly improved survival in patients with stable symptomatic HF (NYHA class III to IV) on standard therapy (i.e., diuretics plus ACEI)	Good					
					<table><tr><th>Endpoint</th><th>Bisoprolol (N=1327)</th><th>Placebo (N=1320)</th><th>p value</th></tr><tr><td>Primary</td><td>156 (11.8%)</td><td>228 (17.3%)</td><td>&lt;0.0001</td></tr><tr><td>CV death</td><td>119 (9%)</td><td>161 (12.2%)</td><td>0.0049</td></tr></table> <b>Target dose:</b> Bisoprolol 10 mg (564 patients); 7.5 mg (152 patients); 5 mg (176 patients) <b>Most common dose:</b> 10 mg			Endpoint	Bisoprolol (N=1327)	Placebo (N=1320)	p value	Primary
Endpoint	Bisoprolol (N=1327)	Placebo (N=1320)	p value									
Primary	156 (11.8%)	228 (17.3%)	<0.0001									
CV death	119 (9%)	161 (12.2%)	0.0049									
<b>US Carvedilol<sup>161</sup> 1996</b> MC, R, DB (vs. placebo) U.S.   Supported by Roche and GlaxoSmithKline	NYHA II (53%), III (44%), IV (3%) Mean EF 23%	1094	Carvedilol 25 to 50 mg twice daily (target dose) or 6.25, 12.5, or 25 mg twice daily (dose-ranging protocol) vs. Placebo  <u>HF therapy</u> ACEI: 95% Digoxin: 91% Diuretics: 95%	Median 6.5 months (stopped early due to improved survival)	<b>Primary Endpoint:</b> Death (65% ↓ with carvedilol; 95% CI 0.39-0.80; ARR 4.6%, NNT 22)	Carvedilol reduced the risk of death in patients with symptomatic HF on standard therapy (i.e., diuretics plus ACEI); individual protocols designed to assess nonfatal endpoints, with mortality prespecified to evaluate safety and benefit in overall trial	Fair					
					<table><tr><th>Endpoint</th><th>Carvedilol (N=696)</th><th>Placebo (N=398)</th><th>p value</th></tr><tr><td>Primary</td><td>22 (3.2%)</td><td>31 (7.8%)</td><td>&lt;0.001</td></tr><tr><td>CV hosp</td><td>98 (14.1%)</td><td>78 (19.6%)</td><td>0.036</td></tr></table> <b>Target dose:</b> Achieved in 80% of patients <b>Mean dose:</b> 45 ± 27 mg per day			Endpoint	Carvedilol (N=696)	Placebo (N=398)	p value	Primary
Endpoint	Carvedilol (N=696)	Placebo (N=398)	p value									
Primary	22 (3.2%)	31 (7.8%)	<0.001									
CV hosp	98 (14.1%)	78 (19.6%)	0.036									

ACEI=angiotensin-converting enzyme inhibitor; ARB=angiotensin II receptor blocker; ARR=absolute risk reduction; CI=confidence interval; CV=cardiovascular; DB=double-blind; EF=ejection fraction; HF=heart failure; hosp=hospitalizations; HR=hazard ratio; IR=immediate-release; ITT=intention-to-treat analysis; N=number of patients; NNT=number needed to treat; NR=not reported; NYHA=New York Heart Association; PROBE=prospective, randomized, open-label, blinded endpoint evaluation; R=randomized; RR=relative risk; XL=extended-release; yrs=years

Angiotensin II Receptor Antagonists																							
Trial	Patient Population	N	Treatment	Duration	Results	Study Conclusions	Quality Rating																
<b>CHARM-Overall<sup>206</sup> 2003</b> MC, R, DB, PG (vs. placebo) U.S., Canada, Europe, S. Africa, Australia, Malaysia  Supported by AstraZeneca R&D	NYHA II (45%), III (52%), IV (3%) HF EF < 40% (57%); EF ≥ 40% (43%)	7601	Candesartan 32 mg once daily (target dose) vs. Placebo  <u>HF therapy</u> ACEI: 41% BB: 55% Digoxin: 43% Diuretics: 83% Spironolactone: 17%	Median 37.7 months	<b>Primary Endpoint:</b> All-cause mortality; no statistically significant difference vs. placebo (unadjusted HR 0.91 95% CI 0.83-1.00)	Candesartan significantly reduced CV deaths and HF hospitalizations	Good																
					<table><tr><th>Endpoint</th><th>Candesartan (N=3803)</th><th>Placebo (N=3796)</th><th>p value</th></tr><tr><td>Primary</td><td>886 (23%)</td><td>945 (25%)</td><td>0.055</td></tr><tr><td>CV death/HF hosp</td><td>1150 (30.2%)</td><td>1310 (34.5%)</td><td>&lt;0.0001</td></tr></table>			Endpoint	Candesartan (N=3803)	Placebo (N=3796)	p value	Primary	886 (23%)	945 (25%)	0.055	CV death/HF hosp	1150 (30.2%)	1310 (34.5%)	<0.0001				
					Endpoint			Candesartan (N=3803)	Placebo (N=3796)	p value													
					Primary			886 (23%)	945 (25%)	0.055													
					CV death/HF hosp			1150 (30.2%)	1310 (34.5%)	<0.0001													
<b>Target dose:</b> Candesartan (63%) vs. placebo (75%) at 6 months																							
<b>Mean dose:</b> 24 mg at 6 months																							
<b>CHARM-Alternative<sup>208</sup> 2003</b> MC, R, DB (vs. placebo) U.S., Canada, Europe, S. Africa, Australia, Malaysia  Supported by AstraZeneca R&D	NYHA II (48%), III (49%), IV (4%) HF Mean EF 30% ACEI intolerant	2028	Candesartan 32 mg once daily (target dose) vs. Placebo  <u>HF therapy</u> BB: 55% Digoxin: 45% Diuretics: 85% Spironolactone: 25%	Median 33.7 months	<b>Primary Endpoint:</b> Composite CV death or HF hospitalizations; ↓ with candesartan (unadjusted HR 0.77 95% CI 0.67-0.89; ARR 7.0%, NNT 14)	Candesartan significantly reduced CV deaths and HF hospitalizations in patients with symptomatic HF who are ACEI intolerant	Good																
					<table><tr><th>Endpoint</th><th>Candesartan (N=1013)</th><th>Placebo (N=1015)</th><th>p value</th></tr><tr><td>Primary</td><td>334 (33%)</td><td>406 (40%)</td><td>0.0004</td></tr><tr><td>CV death</td><td>219 (21.6%)</td><td>252 (24.8%)</td><td>0.072</td></tr><tr><td>HF hosp</td><td>207 (20.4%)</td><td>286 (28.2%)</td><td>&lt;0.0001</td></tr></table>			Endpoint	Candesartan (N=1013)	Placebo (N=1015)	p value	Primary	334 (33%)	406 (40%)	0.0004	CV death	219 (21.6%)	252 (24.8%)	0.072	HF hosp	207 (20.4%)	286 (28.2%)	<0.0001
					Endpoint			Candesartan (N=1013)	Placebo (N=1015)	p value													
					Primary			334 (33%)	406 (40%)	0.0004													
					CV death			219 (21.6%)	252 (24.8%)	0.072													
HF hosp	207 (20.4%)	286 (28.2%)	<0.0001																				
No significant difference (p=0.11) in all-cause death (not a pre-specified endpoint)																							
<b>Target dose:</b> Candesartan (59%) vs. placebo (73%) at 6 months																							
<b>Mean dose:</b> 23 mg at 6 months																							
<b>CHARM-Added<sup>207</sup> 2003</b> MC, R, DB (vs. placebo) U.S., Canada, Europe, S. Africa, Australia, Malaysia  Supported by AstraZeneca R&D	NYHA II (24%), III (73%), IV (3%) Mean EF 28%	2548	Candesartan 32 mg once daily (target dose) vs. Placebo  <u>HF therapy</u> ACEI: 100% BB: 55% Digoxin: 58% Diuretics: 90% Spironolactone: 17%	Median 41 months	<b>Primary Endpoint:</b> Composite CV death or HF hospitalizations; ↓ with candesartan (unadjusted HR 0.85 95% CI 0.75-0.96; ARR 4.4%, NNT 23)	The addition of candesartan to treatment with an ACEI and other standard therapy for HF significantly reduced CV deaths and HF hospitalizations in patients with symptomatic HF	Good																
					<table><tr><th>Endpoint</th><th>Candesartan (N=1276)</th><th>Placebo (N=1272)</th><th>p value</th></tr><tr><td>Primary</td><td>483 (37.9%)</td><td>538 (42.3%)</td><td>0.011</td></tr><tr><td>CV death</td><td>302 (23.7%)</td><td>347 (27.3%)</td><td>0.029</td></tr><tr><td>HF hosp</td><td>309 (24.2%)</td><td>356 (28.0%)</td><td>0.014</td></tr></table>			Endpoint	Candesartan (N=1276)	Placebo (N=1272)	p value	Primary	483 (37.9%)	538 (42.3%)	0.011	CV death	302 (23.7%)	347 (27.3%)	0.029	HF hosp	309 (24.2%)	356 (28.0%)	0.014
					Endpoint			Candesartan (N=1276)	Placebo (N=1272)	p value													
					Primary			483 (37.9%)	538 (42.3%)	0.011													
					CV death			302 (23.7%)	347 (27.3%)	0.029													
HF hosp	309 (24.2%)	356 (28.0%)	0.014																				
No significant difference (p=0.086) in all-cause death (not a pre-specified endpoint)																							
<b>Target dose:</b> Candesartan (61%) vs. placebo (73%) at 6 months																							
<b>Mean dose:</b> 24 mq at 6 months																							

Angiotensin II Receptor Antagonists (continued)																							
Trial	Patient Population	N	Treatment	Duration	Results	Study Conclusions	Quality Rating																
<b>Val-HeFT<sup>204</sup> 2001</b> MC, R, DB (vs. placebo) U.S., Europe, S. Africa, Australia       Supported by Novartis Pharma	NYHA II (62%), III (36%), IV (2%) Mean EF 27%	5010	Valsartan 160 mg twice daily (target dose) vs. Placebo  <u>HF therapy</u> ACEI: 93% BB: 35% Digoxin: 67% Diuretics: 85% Spironolactone: 5%	Mean 23 months	<b>Primary Endpoint:</b> 1) Overall mortality (no statistically significant difference vs. placebo); and 2) Combined morbidity and mortality (↓ with valsartan; RR 0.87 97.5% CI 0.77-0.97; ARR 3.3%, NNT 30) <table><tr><th>Endpoint</th><th>Valsartan (N=2511)</th><th>Placebo (N=2499)</th><th>p value</th></tr><tr><td>Primary1</td><td>495 (19.7%)</td><td>484 (19.4%)</td><td>0.80</td></tr><tr><td>Primary2</td><td>723 (28.8%)</td><td>801 (32.1%)</td><td>0.009</td></tr><tr><td>HF hosp*</td><td>346 (13.8%)</td><td>455 (18.2%)</td><td>&lt;0.001</td></tr></table> *adjudicated hosp for worsening HF as 1 <sup>st</sup> event <b>Target dose:</b> Valsartan (84%) vs. placebo (93%) <b>Mean dose:</b> 254 mg	Endpoint	Valsartan (N=2511)	Placebo (N=2499)	p value	Primary1	495 (19.7%)	484 (19.4%)	0.80	Primary2	723 (28.8%)	801 (32.1%)	0.009	HF hosp*	346 (13.8%)	455 (18.2%)	<0.001	Valsartan significantly reduced the combined morbidity and mortality endpoint when given to patients with HF currently on therapy; a post hoc analysis noted an increase in mortality in patients receiving combination of valsartan with an ACEI and BB	Good
Endpoint	Valsartan (N=2511)	Placebo (N=2499)	p value																				
Primary1	495 (19.7%)	484 (19.4%)	0.80																				
Primary2	723 (28.8%)	801 (32.1%)	0.009																				
HF hosp*	346 (13.8%)	455 (18.2%)	<0.001																				
<b>ELITE II<sup>202</sup> 2000</b> MC, R, DB (vs. ACEI) U.S., Canada, Europe, S. Africa, S. America       Supported by Merck Research Laboratories	NYHA II (52%), III (43%), IV (5%) Mean EF 31%	3152	Losartan 50 mg once daily (target dose) vs. Captopril 50 mg three times daily (target dose)  <u>HF therapy</u> BB: 22% Digoxin: 50% Diuretics: 78%	Median 1.5 yrs	<b>Primary Endpoint:</b> All-cause mortality; no statistically significant difference vs. captopril (HR 1.13 95.7% CI 0.95-1.35) <table><tr><th>Endpoint</th><th>Losartan (N=1578)</th><th>Captopril (N=1574)</th><th>p value</th></tr><tr><td>Primary</td><td>280 (17.7%)</td><td>250 (15.9%)</td><td>0.16</td></tr><tr><td>Secondary*</td><td>142 (9.0%)</td><td>115 (7.3%)</td><td>0.08</td></tr><tr><td>HF hosp*</td><td>270 (17.1%)</td><td>293 (18.6%)</td><td>0.32</td></tr></table> *composite sudden cardiac death or resuscitated cardiac arrest <b>Mean dose:</b> NA <b>Target dose:</b> NA	Endpoint	Losartan (N=1578)	Captopril (N=1574)	p value	Primary	280 (17.7%)	250 (15.9%)	0.16	Secondary*	142 (9.0%)	115 (7.3%)	0.08	HF hosp*	270 (17.1%)	293 (18.6%)	0.32	Losartan was not found to be superior to captopril in reducing all-cause mortality in patients with HF currently on therapy; designed as a superiority trial, unable to determine equivalence between losartan and captopril	Good
Endpoint	Losartan (N=1578)	Captopril (N=1574)	p value																				
Primary	280 (17.7%)	250 (15.9%)	0.16																				
Secondary*	142 (9.0%)	115 (7.3%)	0.08																				
HF hosp*	270 (17.1%)	293 (18.6%)	0.32																				

ACEI=angiotensin-converting enzyme inhibitor; ARR=absolute risk reduction; BB=beta-blockers; CI=confidence interval; CV=cardiovascular; DB=double-blind; EF=ejection fraction; HF=heart failure; hosp=hospitalizations; HR=hazard ratio; N=number of patients; NA=not available; NNT=number needed to treat; PG=parallel-group; R=randomized; RR=relative risk; Sx=symptoms; yrs=years

Hydralazine/Isosorbide Dinitrate																											
Trial	Patient Population	N	Treatment	Duration	Results	Study Conclusions	Quality Rating																				
<b>A-HeFT<sup>234</sup> 2004</b> MC, DB, RCT (vs. placebo) U.S.  Supported by NitroMed	Self-identified black with (African descent) NYHA III or IV HF ≥ 3 months Mean EF ~24%	1050	ISDN/HYD 40 mg/75 mg three times daily (120 mg/225 mg total daily target dose) vs. Placebo  <u>HF therapy</u> ACEI: 69% ARB: 17% BB: 74% Carvedilol: 55% Digoxin: 60% Diuretics: 90% Spironolactone: 39%	10 months (terminated early due to difference in mortality)	<b>Primary endpoint:</b> Composite score (weighted values for all-cause mortality, 1 <sup>st</sup> HF hosp during 18 months, change in QOL by MLHF at 6 months); possible score -6 to +2 <table><tr><th>Endpoint</th><th>ISDN/HYD (N=518)</th><th>Placebo (N=532)</th><th>p value</th></tr><tr><td>Primary (score)</td><td>-0.1+1.9</td><td>-0.5+2.0</td><td>0.01</td></tr><tr><td>All-cause death</td><td>32 (6.2%)</td><td>54 (10.2%)</td><td>0.02</td></tr><tr><td>1<sup>st</sup> HF hosp</td><td>85 (16.4%)</td><td>130 (24.4%)</td><td>0.001</td></tr><tr><td>Change QOL*</td><td>-5.6+20.6</td><td>-2.7+21.2</td><td>0.02</td></tr></table> *lower score reflects better QOL <b>Survival:</b> 43% improvement with ISDN/HYD (HR 0.57; p=0.01) <b>Target dose:</b> ISDN/HYD (68%) vs. placebo (88.9%); p<0.001 <b>Mean # tabs/d:</b> ISDN/HYD (3.8±2.5) vs. placebo (4.7±2.2); p<0.001	Endpoint	ISDN/HYD (N=518)	Placebo (N=532)	p value	Primary (score)	-0.1+1.9	-0.5+2.0	0.01	All-cause death	32 (6.2%)	54 (10.2%)	0.02	1 <sup>st</sup> HF hosp	85 (16.4%)	130 (24.4%)	0.001	Change QOL*	-5.6+20.6	-2.7+21.2	0.02	Combination ISDN/HYD, in addition to standard therapy for HF, improved survival and decreased rate of first hospitalizations for HF, in self-identified black patients with NYHA class III to IV HF	Fair
Endpoint	ISDN/HYD (N=518)	Placebo (N=532)	p value																								
Primary (score)	-0.1+1.9	-0.5+2.0	0.01																								
All-cause death	32 (6.2%)	54 (10.2%)	0.02																								
1 <sup>st</sup> HF hosp	85 (16.4%)	130 (24.4%)	0.001																								
Change QOL*	-5.6+20.6	-2.7+21.2	0.02																								
<b>V-HeFT II<sup>121</sup> 1991</b> MC (VA), DB, RCT (vs. ACEI) U.S. Supported by VA Cooperative Studies	Males with primarily NYHA II (51%) or III (43%) HF Mean EF ~29%	804	ISDN/HYD 40 mg/75 mg four times daily (160 mg/300 mg total daily target dose) vs. Enalapril 10 mg twice daily (20 mg total daily target dose)  <u>HF therapy</u> Digoxin Diuretics	Ave 2.5 yrs	<b>Primary Endpoint:</b> Overall and 2-yr mortality; 2-yr mortality ↓ with ACEI vs. ISDN/HYD (risk reduction 28.2%; ARR 7.0%, NNT 14) <table><tr><th>Mortality</th><th>ISDN/HYD (N=401)</th><th>ACEI (N=403)</th><th>p value</th></tr><tr><td>Overall</td><td>153 (38.2%)</td><td>132 (32.8%)</td><td>0.08</td></tr><tr><td>2-yr</td><td>NR (25%)</td><td>NR (18%)</td><td>0.016</td></tr><tr><td>HF hosp</td><td>78 (18.4%)</td><td>76 (18.9%)</td><td>NR</td></tr></table> <b>Ave daily dose:</b> ISDN/HYD (100/199 mg); enalapril (15 mg)	Mortality	ISDN/HYD (N=401)	ACEI (N=403)	p value	Overall	153 (38.2%)	132 (32.8%)	0.08	2-yr	NR (25%)	NR (18%)	0.016	HF hosp	78 (18.4%)	76 (18.9%)	NR	Mortality was lower with enalapril compared to ISDN/HYD, a difference that was statistically significant at 2 yrs	Fair				
Mortality	ISDN/HYD (N=401)	ACEI (N=403)	p value																								
Overall	153 (38.2%)	132 (32.8%)	0.08																								
2-yr	NR (25%)	NR (18%)	0.016																								
HF hosp	78 (18.4%)	76 (18.9%)	NR																								
<b>V-HeFT I<sup>123</sup> 1986</b> MC (VA), DB, RCT (vs. prazosin or placebo) U.S.  Supported by VA Cooperative Studies	Males with chronic congestive HF Mean EF ~30%	642	ISDN/HYD 40 mg/75 mg four times daily (160 mg/300 mg total daily target dose) vs. Prazosin 5 mg four times daily (20 mg total daily target dose) vs. Placebo  <u>HF therapy</u> Digoxin Diuretics	Ave 2.3 yrs	<b>Primary Endpoint:</b> Overall and 2-yr mortality; ↓ in 2-yr mortality with ISDN/HYD (risk reduction 34%, CI 0.04 to 0.54; ARR 8.7%, NNT=12) <table><tr><th>Mortality</th><th>ISDN/HYD (N=186)</th><th>Placebo (N=273)</th><th>p value</th></tr><tr><td>Overall*</td><td>NR (36.2%)</td><td>NR (46.9%)</td><td>NR</td></tr><tr><td>2-yr</td><td>NR (25.6%)</td><td>NR (34.3%)</td><td>&lt;0.028</td></tr></table> *by 3 yrs, mortality risk reduction 36% with ISDN/HYD vs. placebo; small sample size > 3yrs <b>Ave daily dose:</b> ISDN/HYD (136/270 mg); prazosin (18.6 mg)	Mortality	ISDN/HYD (N=186)	Placebo (N=273)	p value	Overall*	NR (36.2%)	NR (46.9%)	NR	2-yr	NR (25.6%)	NR (34.3%)	<0.028	Mortality reduced with ISDN/HYD compared to placebo up to 3 yrs; unable to determine benefit beyond this point	Fair								
Mortality	ISDN/HYD (N=186)	Placebo (N=273)	p value																								
Overall*	NR (36.2%)	NR (46.9%)	NR																								
2-yr	NR (25.6%)	NR (34.3%)	<0.028																								

ACEI=angiotensin-converting enzyme inhibitor; AE=adverse event; ARB=angiotensin II receptor blocker; ARR=absolute risk reduction; Ave=average; BB=beta-adrenergic blocker; DB=double-blind; EF=ejection fraction; HF=heart failure; hosp=hospitalizations; ISDN/HYD=isosorbide dinitrate and hydralazine; MC=multicenter; n=number of patients; MLHF=Minnesota Living with Heart Failure questionnaire; NNT=number needed to treat; NR=not reported; NYHA=New York Heart Association; QOL=quality of life; RCT=randomized controlled trial; VA=Veterans Affairs Medical Center; yrs=years



Digitalis										
Trial	Patient Population	N	Treatment	Duration	Results	Study Conclusions	Quality Rating			
DIG <sup>93</sup> 1997 MC, R, DB (vs. placebo) U.S., Canada   NHLBI and VA Cooperative study; drug and placebo provided by Glaxo Wellcome	NYHA I (13%); II (54%); III (31%), IV (2%) HF Mean EF 29%	6800	Digoxin dosed per algorithm (based on age, gender, weight, and kidney function) vs. Placebo  <u>HF therapy</u> ACEI: 95% Diuretics: 82%	Mean 37 months	Primary Endpoint: All-cause mortality; no statistically significant difference vs. placebo (RR 0.99 95% CI 0.91-1.07)	All-cause mortality was not significantly reduced with digoxin; there was a significant decrease in HF hospitalization in patients with HF receiving treatment with digoxin	Good			
					Endpoint			Digoxin (N=3397)	Placebo (N=3403)	p value
					Primary			1181 (34.8%)	1194 (35.1%)	0.8
					CV death			1016 (29.9%)	1004 (29.5%)	0.78
					HF hosp			910 (26.8%)	1180 (34.7%)	<0.001
					Median dose (at randomization): 0.25 mg once daily Mean serum digoxin concentration (steady state at 12 months): 0.80 ng/ml					

ACEI=angiotensin-converting enzyme inhibitor; CI=confidence interval; CV=cardiovascular; DB=double-blind; EF=ejection fraction; HF=heart failure; hosp=hospitalizations; N=number of patients; NYHA=New York Heart Association; R=randomized; RR=relative risk

Aldosterone Antagonists																							
Trial	Patient Population	N	Treatment	Duration	Results	Study Conclusions	Quality Rating																
<b>EPHESUS<sup>250</sup></b> <b>2003</b> MC, R, DB (vs. placebo) U.S., Canada, Europe, Latin America  Supported by Pharmacia	Days from MI to R (7.3) HF symptoms (90%) Mean EF 33%	6632	Eplerenone 25 mg once daily (50 mg once daily target dose) vs. Placebo  <u>Drug therapy</u> ACEI or ARB: 87% BB: 75% Diuretics: 61%	Mean 16 months	<b>Primary Endpoints:</b> 1) Time to death (any cause); ↓ with eplerenone (RR 0.85 95% CI 0.75-0.96; ARR 2.3%; NNT 43); 2) Time to death from CV causes or 1 <sup>st</sup> CV hosp; ↓ with eplerenone (RR 0.87 95% CI 0.79-0.95; ARR 3.3%; NNT 30) <table><tr><th>Endpoint</th><th>Eplerenone (N=3319)</th><th>Placebo (N=3313)</th><th>p value</th></tr><tr><td>Primary1</td><td>478 (14.4%)</td><td>554 (16.7%)</td><td>0.008</td></tr><tr><td>CV death or hosp</td><td>885 (26.7%)</td><td>993 (30.0%)</td><td>0.002</td></tr></table> <b>Mean dose:</b> 42.6 mg once daily	Endpoint	Eplerenone (N=3319)	Placebo (N=3313)	p value	Primary1	478 (14.4%)	554 (16.7%)	0.008	CV death or hosp	885 (26.7%)	993 (30.0%)	0.002	Eplerenone significantly reduced combined death from CV cause or CV hospitalization in patients with acute MI complicated by LVD and HF	Good				
Endpoint	Eplerenone (N=3319)	Placebo (N=3313)	p value																				
Primary1	478 (14.4%)	554 (16.7%)	0.008																				
CV death or hosp	885 (26.7%)	993 (30.0%)	0.002																				
<b>RALES<sup>249</sup></b> <b>1999</b> MC, R, DB (vs. placebo) U.S., Canada, Mexico, Europe, S. America, S. Africa, New Zealand, Japan  Supported by Searle	NYHA III (71%), IV (29%) HF Mean EF 25%	1663	Spironolactone 25 mg once daily (increased to 50 mg once daily if signs or symptoms of HF progression without hyperkalemia) vs. Placebo  <u>HF therapy</u> ACEI: 95% BB: 11% Digoxin: 74% Diuretics: 100%	Mean 24 months (terminated early due to survival benefit)	<b>Primary Endpoint:</b> All-cause mortality; ↓ with spironolactone (RR 0.70 95% CI 0.60-0.82; ARR 11.4%; NNT=9) <table><tr><th>Endpoint</th><th>Spironolactone (N=841)</th><th>Placebo (N=822)</th><th>p value</th></tr><tr><td>Primary</td><td>284 (34.6%)</td><td>386 (45.9%)</td><td>&lt;0.001</td></tr><tr><td>CV death</td><td>226 (27.5%)</td><td>314 (37.3%)</td><td>&lt;0.001</td></tr><tr><td>HF hosp</td><td>215 (26.2%)</td><td>300 (35.7%)</td><td>&lt;0.001</td></tr></table> <b>Mean dose:</b> 26 mg once daily	Endpoint	Spironolactone (N=841)	Placebo (N=822)	p value	Primary	284 (34.6%)	386 (45.9%)	<0.001	CV death	226 (27.5%)	314 (37.3%)	<0.001	HF hosp	215 (26.2%)	300 (35.7%)	<0.001	Spironolactone significantly reduced all-cause mortality in patients with severe HF	Good
Endpoint	Spironolactone (N=841)	Placebo (N=822)	p value																				
Primary	284 (34.6%)	386 (45.9%)	<0.001																				
CV death	226 (27.5%)	314 (37.3%)	<0.001																				
HF hosp	215 (26.2%)	300 (35.7%)	<0.001																				

ACEI=angiotensin-converting enzyme inhibitor; ARB=angiotensin II receptor blocker; ARR=absolute risk reduction; BB=beta-blockers; CI=confidence interval; CV=cardiovascular; DB=double-blind; EF=ejection fraction; HF=heart failure; hosp=hospitalizations; LVD=left ventricular dysfunction; MI=myocardial infarction; N=number of patients; NYHA=New York Heart Association; NNT=number needed to treat; R=randomized; RR=relative risk