

MSiA-401-hw7-q4

2022-11-29

```
setwd("~/Desktop/MSiA-401-hw7")
desert = read.csv("desert.csv",header = T)
# omit NA
desert2 = na.omit(desert)
head(desert2,10)
```

	FIPS	newsPub	age	pop	BAhigher	income	raceBlack	race_Hisp	digDistress
## 1	1003	4	42.6	203360	30.7	52562	9.5	4.4	28.58287
## 2	1005	1	39.7	26201	12.0	33368	47.8	4.2	50.65492
## 3	1007	1	39.8	22580	13.2	43404	22.0	2.4	52.60406
## 4	1009	1	40.9	57667	13.1	47412	1.5	9.0	39.09295
## 5	1011	1	40.8	10478	13.4	29655	75.6	0.3	62.23687
## 6	1013	1	40.7	20126	16.1	36326	44.7	0.3	48.72265
## 7	1015	1	39.1	115527	17.9	43686	20.4	3.6	32.97889
## 8	1017	2	43.0	33895	13.3	37342	39.3	2.2	46.54169
## 9	1019	1	46.1	25855	12.5	40041	5.0	1.6	43.37013
## 10	1021	2	38.9	43805	15.1	43501	9.5	7.7	51.71853

```
fit1 = glm(newsPub ~ . -FIPS, poisson, data = desert2)
summary(fit1)
```

```
##
## Call:
## glm(formula = newsPub ~ . - FIPS, family = poisson, data = desert2)
##
## Deviance Residuals:
##      Min       1Q   Median       3Q      Max
## -8.9405  -0.6614  -0.2605   0.3281   9.8664
##
## Coefficients:
##              Estimate Std. Error z value Pr(>|z|)
## (Intercept)  6.058e-01  1.570e-01   3.859 0.000114 ***
## age         -1.876e-03  2.654e-03  -0.707 0.479574
## pop          3.890e-07  9.183e-09  42.360 < 2e-16 ***
## BAhhigher    1.416e-02  1.944e-03   7.283 3.27e-13 ***
## income       3.734e-06  1.360e-06   2.745 0.006049 **
## raceBlack    4.300e-03  1.102e-03   3.902 9.54e-05 ***
## race_Hisp     3.278e-03  1.069e-03   3.068 0.002159 **
## digDistress -1.627e-02  1.977e-03  -8.229 < 2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## (Dispersion parameter for poisson family taken to be 1)
```

```
##
##      Null deviance: 6123.8   on 3141   degrees of freedom
## Residual deviance: 3693.6   on 3134   degrees of freedom
## AIC: 10792
##
## Number of Fisher Scoring iterations: 5
```

First we fit a Poisson model `fit1` by simply use all variables except `FIPS`, we observe the deviance for the model is 3693.6 and AIC is 10792.

```
library(Hmisc)
```

```
## Loading required package: lattice
```

```
## Loading required package: survival
```

```
## Loading required package: Formula
```

```
## Loading required package: ggplot2
```

```
##
```


```
## Attaching package: 'Hmisc'
```

```
## The following objects are masked from 'package:base':
```

```
##
```

```
##      format.pval, units
```

```
hist.data.frame(desert2)
```



By observing the data distribution for all variables, we find variables `pop`, `raceBlack`, `race_Hisp` have very non-symmetric distribution. In that case, we will transform these three variables by taking log in the new model.

Besides, since `digDistress` is considered as a pip (determined by `pop` and `income`), we will not include `digDistress` in the new model.

```
fit2 = glm(newsPub ~ age + log(pop) + BAhhigher + income + log(raceBlack+1) + log(race_Hisp+1), poisson,
summary(fit2))
```

```
##
## Call:
## glm(formula = newsPub ~ age + log(pop) + BAhhigher + income +
##      log(raceBlack + 1) + log(race_Hisp + 1), family = poisson,
##      data = desert2)
##
## Deviance Residuals:
##      Min       1Q   Median       3Q      Max
## -4.4815  -0.6708  -0.1023   0.3972  12.5225
##
## Coefficients:
##              Estimate Std. Error z value Pr(>|z|)
## (Intercept)  -5.133e+00  1.746e-01 -29.395  < 2e-16 ***
## age           2.099e-02  2.810e-03   7.469 8.11e-14 ***
## log(pop)      4.548e-01  1.134e-02  40.106 < 2e-16 ***
## BAhhigher     2.727e-03  1.940e-03   1.405  0.1599
## income        2.613e-06  1.290e-06   2.025  0.0428 *
```

```
## log(raceBlack + 1) -1.109e-01 1.320e-02 -8.407 < 2e-16 ***
## log(race_Hisp + 1) 3.060e-02 1.588e-02 1.927 0.0540 .
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## (Dispersion parameter for poisson family taken to be 1)
##
## Null deviance: 6123.8 on 3141 degrees of freedom
## Residual deviance: 2957.6 on 3135 degrees of freedom
## AIC: 10054
##
## Number of Fisher Scoring iterations: 5
```

We observe the new model fit2 improves significantly because deviance and AIC of fit2 is 2957.6 and 10054, which is less than 3693.6 and 10792.

We will then use drop1 to see if dropping any other variables could improve the model.

```
drop1(fit2)
```

```
## Single term deletions
##
## Model:
## newsPub ~ age + log(pop) + BAhhigher + income + log(raceBlack +
## 1) + log(race_Hisp + 1)
##
```

	Df	Deviance	AIC
<none>		2957.6	10054
age	1	3012.9	10107
log(pop)	1	4693.6	11788
BAhhhigher	1	2959.6	10054
income	1	2961.7	10056
log(raceBlack + 1)	1	3029.8	10124
log(race_Hisp + 1)	1	2961.3	10055

```
summary(fit2)
```

```
##
## Call:
## glm(formula = newsPub ~ age + log(pop) + BAhhigher + income +
## log(raceBlack + 1) + log(race_Hisp + 1), family = poisson,
## data = desert2)
##
## Deviance Residuals:
##      Min       1Q   Median       3Q      Max
## -4.4815  -0.6708  -0.1023   0.3972  12.5225
##
## Coefficients:
##              Estimate Std. Error z value Pr(>|z|)
## (Intercept)  -5.133e+00  1.746e-01 -29.395 < 2e-16 ***
## age           2.099e-02  2.810e-03   7.469 8.11e-14 ***
## log(pop)      4.548e-01  1.134e-02  40.106 < 2e-16 ***
## BAhhigher     2.727e-03  1.940e-03   1.405  0.1599
## income        2.613e-06  1.290e-06   2.025  0.0428 *
```

```
## log(raceBlack + 1) -1.109e-01  1.320e-02  -8.407  < 2e-16 ***
## log(race_Hisp + 1)  3.060e-02  1.588e-02   1.927   0.0540 .
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## (Dispersion parameter for poisson family taken to be 1)
##
##      Null deviance: 6123.8  on 3141  degrees of freedom
## Residual deviance: 2957.6  on 3135  degrees of freedom
## AIC: 10054
##
## Number of Fisher Scoring iterations: 5
```

We observe dropping any of variables would not improve the model because dropping any variables would increase deviance of the model. But we notice variables BAhiger and $\log(\text{race_Hisp} + 1)$ is non-significant (have p-value greater than 0.05), dropping them also barely affect deviance and AIC. In that case, we drop variables BAhiger and $\log(\text{race_Hisp} + 1)$.

```
fit3 = glm(newsPub ~ age + log(pop) + income + log(raceBlack+1), poisson, data = desert2)
summary(fit3)
```

```
##
## Call:
## glm(formula = newsPub ~ age + log(pop) + income + log(raceBlack +
##      1), family = poisson, data = desert2)
##
## Deviance Residuals:
##      Min       1Q   Median       3Q      Max
## -4.4642  -0.6701  -0.0997   0.4025  12.5988
##
## Coefficients:
##              Estimate Std. Error z value Pr(>|z|)
## (Intercept)    -5.136e+00  1.703e-01 -30.155  < 2e-16 ***
## age              1.942e-02  2.719e-03   7.143 9.11e-13 ***
## log(pop)         4.661e-01  1.020e-02  45.689  < 2e-16 ***
## income           3.915e-06  9.378e-07   4.175 2.98e-05 ***
## log(raceBlack + 1) -1.125e-01  1.310e-02  -8.590  < 2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## (Dispersion parameter for poisson family taken to be 1)
##
##      Null deviance: 6123.8  on 3141  degrees of freedom
## Residual deviance: 2962.8  on 3137  degrees of freedom
## AIC: 10055
##
## Number of Fisher Scoring iterations: 5
```

From the model above, we can see predictor variables age, $\log(\text{pop})$, income have positive effect on dependent variable newsPub, whereas predictor variable $\log(\text{raceBlack} + 1)$ has negative effect on newsPub. This makes common sense since older people tend to read more newspaper; people read newspaper will increase as the population increase; people with higher income can afford buying newspaper than people with less income.