# FRACTAL AND MULTIFRACTAL ANALYSIS AS TOOLS TO CHARACTERIZE SUPERNOVAE AND MOLECULAR CLOUDS: SUBTITLE

Samuel Brenner<sup>1</sup>
Draft version July 18, 2013

#### ABSTRACT

Keywords: supernovae, fractals, multifractals

#### 1. INTRODUCTION

Many physical phenomena cannot be characterized by the idealizations of Euclidean geometry alone; they exhibit "roughness". That is, they have a detailed structure at any arbitrarily small size scale (Falconer 2003). The development of fractal geometry allows a mathematical treatment of the "roughness" inherent in the non-idealized phenomena of the real world. Multifractal analysis permits us to examine how those fractal characteristics themselves change with scale; in fact, they may even be fractal themselves. In this paper, we detail the application of fractal geometry to flame fronts in white dwarfs and then analyze the multifractal characteristics of the cool molecular clouds involved in star formation.

### 1.1. Fractal analysis of type Ia supernova flame fronts

The accepted model for a type Ia supernova is a white dwarf that accumulates mass from a binary companion until it reaches a mass so large that the compression of the heavier elements in the core causes them to ignite and fuse, releasing more energy in the process. This deflagration results in a flame front that spreads rapidly throughout the star and causes the visible effects of supernova<sup>2</sup>.

The surface of the expanding flame front can be dramatically affected by multiple types of turbulence. Landau & Lifshitz (1959) showed that the flame surface is subject to Landau-Darrieus instability in which the front is convoluted due to thermal expansion across the flame front. It is also widely established (see Kull (1991)) that Rayleigh-Taylor instability, in which a denser fluid pushes through a less-dense one under the influence of a gravitational field, can contort the deflagration front in a supernova.

These wrinkles on the surface of the flame front increase the front's effective surface area, and because the flame speed is proportional to the flame's surface area, the convolution of the surface can influence the rate at which the flame consumes the star. Thus, characterizing the shape of the flame front plays a crucial part in characterizing the supernova explosion process as a whole.

These wrinkles on the surface of a turbulent front can be viewed within the framework of fractal geometry, as was proposed by Mandelbrot (1975) and further developed by Timmes, F. X. (1994). Furthermore, Timmes, F. X. (1994) showed that the final effective speed of the flame front is a function of its density. In  $\S 2.1$ , a method is implemented to perform fractal analysis of a supernova flame front. The results are discussed in  $\S 3.1$ , and the implications of the calculated fractal dimension are discussed in  $\S 4.1$ .

- 1.2. Mutlfractal analysis of dense molecular clouds
  - 1.3. Physical phenomena studied
    - 2. METHODS
    - 2.1. Fractal analysis methods
  - 2.2. Multifractal analysis methods
    - 3. RESULTS
    - 3.1. Fractal analysis methods
  - $3.2.\ Multifractal\ analysis\ methods$ 
    - 4. DISCUSSION
    - 4.1. Fractal analysis methods
  - 4.2. Multifractal analysis methods

Timmes eqn: The final effective speed  $v_{eff}$  of the flame front on a macroscopic scale is given by

$$v_{eff} = v_{lam} \left(\frac{l_{min}}{l_{max}}\right)^{2-D} \tag{1}$$

where  $v_{lam}$  is the laminar speed of the front through the material (that is, the speed of the flame in the absence of turbulence),  $l_{min}$  is the minimum length scale at which the turbulence takes place,  $l_{max}$  is the maximum length scale at which the turbulence takes place, and D is the fractal dimension of the flame front.

#### 5. ACKNOWLEDGMENTS

The work of SB has been supported by Tomasz Plewa and Tim Handy at Florida State University under the Young Scholars Program.

## REFERENCES

Falconer, K. 2003, Fractal Geometry: Mathematical Foundations and Applications, Vol. 46 (Wiley), 366
Kull, H. J. 1991, Physics Reports, 206, 197
Landau, L. D., & Lifshitz, E. M. 1959, Course of Theoretical

Physics, Vol. 6, Image Rochester NY (Pergamon Press), 539 Mandelbrot, B. B. 1975, Journal of Fluid Mechanics, 72, 401 Timmes, F. X. 1994, Astrophysical Journal, 423, L131

Electronic address: samuel.e.brenner@gmail.com  $^{1}$  Young Scholars Program, Florida State University  $^{2}$  it does, right?