

Soil carbon and nitrogen mineralization following deposition of insect frass and greenfall from forests under elevated CO₂ and O₃

Michael Hillstrom · Timothy D. Meehan ·
Kristine Kelly · Richard L. Lindroth

Received: 18 February 2010 / Accepted: 21 May 2010 / Published online: 5 June 2010
© Springer Science+Business Media B.V. 2010

Abstract Elevated CO₂ and O₃ alter tree quality and the quality of herbivore inputs, such as frass, to forest soil. Altered quality or quantity of herbivore inputs to the forest floor can have large impacts on below-ground processes. We collected green leaves and frass from whitemarked tussock moth caterpillars from aspen-birch stands at the Aspen Free Air CO₂ Enrichment (FACE) site near Rhinelander, WI, USA. Small or large quantities of frass, greenfall, or a 1:1 ratio of frass and greenfall were added to microcosms for each FACE treatment (control, +CO₂, +O₃, +CO₂+O₃). We measured initial frass and greenfall quality, and recorded microbial respiration, and nitrate leaching over 40 days. Elevated carbon dioxide (eCO₂) and tropospheric ozone (eO₃) significantly altered the carbon, nitrogen, and condensed tannin content of insect frass and green leaves. Although FACE treatments affected input quality, they had minimal effect on microbial respiration and no effect on nitrogen leaching. In contrast, input quantity substantially influenced microbial respiration and nitrate leaching. Respiratory carbon loss and nitrate immobilization were nearly double in microcosms

receiving large amounts of herbivore inputs than those receiving no herbivore inputs. Small amounts of herbivore inputs, however, did not significantly alter microbial respiration or immobilization, suggesting that effects of herbivore inputs on soil processes will be detected only at moderate to high herbivory/input levels. These results suggest that subtle changes in frass and greenfall quality may not affect soil nutrient cycling. In contrast, environmental change induced increases in insect population size or frass and greenfall inputs to the soil may substantially impact nutrient cycling.

Keywords Carbon and nitrogen mineralization · Aspen FACE · *Orgyia leucostigma* · Canopy herbivore inputs · Soil respiration · Carbon dioxide · Ozone

Introduction

Forest ecosystems of the 21st century face a multitude of global change factors that will alter their structure and function, as well as the services they provide. Fragmentation, invasion of alien species and changes in the atmosphere will alter forests directly through effects on plants and indirectly through impacts on higher trophic levels (e.g., herbivores). Changes in atmospheric concentrations of two gases, carbon dioxide and tropospheric ozone, are expected to have significant impacts on forest ecosystems (Karnosky et al. 2003). Annual human-caused CO₂ emissions have increased more

Responsible Editor: Per Ambus.

M. Hillstrom (✉) · T. D. Meehan · K. Kelly ·
R. L. Lindroth
Department of Entomology, University of Wisconsin,
237 Russell Labs, 1630 Linden Drive,
Madison, WI 53706-1598, USA
e-mail: mlhillstrom@wisc.edu

than 80% since 1970 (IPCC 2007) and tropospheric ozone concentrations are expected to increase from an average ambient background concentration of 40 ppb today to 68 ppb by 2050 and 85 ppb by 2100 (Akimoto 2003; Wittig et al. 2009). Numerous studies have evaluated the effects of both elevated CO₂ (eCO₂) and elevated O₃ (eO₃) on tree health, nutrient status and defense characteristics. Generally, forest trees respond to eCO₂ with increased photosynthesis and growth above- and below-ground (Saxe et al. 1998; Karnosky et al. 2003), decreased foliar nitrogen and increased foliar carbohydrates and phenolics, resulting in increased carbon to nitrogen ratios (Zvereva and Kozlov 2006; Stiling and Cornelissen 2007; Lindroth 2010). Unlike the fertilizing effects of CO₂, tropospheric O₃ causes significant damage to plants, inducing foliar injury (Karnosky 1976; Karnosky et al. 2003) and decreasing photosynthesis and growth (Karnosky et al. 2003). Changes in phytochemistry of trees exposed to eO₃ can shift in magnitude or direction depending on the timing of the sampling (season), and the species and genotypes (O₃ sensitive vs. O₃ resistant) involved (Lindroth et al. 2001; Holton et al. 2003; Kopper and Lindroth 2003; Lindroth 2010). Although the effects of eCO₂ and eO₃ on plant quality vary, the implications are clear: eCO₂ and eO₃ alter the quality of forest trees for herbivores.

Changes in tree quality due to eCO₂ and eO₃ have been shown to affect herbivore growth, survivorship and abundance (Bezemer and Jones 1998; Zvereva and Kozlov 2006; Valkama et al. 2007; Lindroth 2010), but the implications for herbivore-mediated ecosystem processes have received little attention. Canopy herbivory can strongly affect nutrient cycling (Belovsky and Slade 2000; Schowalter 2000; Hunter 2001). Canopy insects may influence soil nutrient dynamics through four types of soil surface inputs: frass deposition, turnover of insect cadavers, changes in throughfall chemistry, or changes in quality or quantity of leaf litter that falls to the forest floor (e.g., greenfall deposition) (Hunter 2001). Frass and greenfall inputs, in particular, are important because they occur during the growing season and frass inputs have been shown to influence tree nutrient levels (Christenson et al. 2002; Frost and Hunter 2007) and soil carbon and nitrogen cycling within the same growing season (Frost and Hunter 2004, 2007). This fast processing of organic matter is possible because, unlike senesced leaves that have much of their

nutrient content resorbed, frass and greenfall inputs are “higher quality” inputs that may contain 74% more available nitrogen than senescent leaves (Risely and Crossley 1993; Fonte and Schowalter 2004; Madritch et al. 2007a). Further, soil nitrogen inputs from frass can exceed those from leaf litter during severe outbreaks (Fogal and Slansky 1984; Grace 1986; Hollinger 1986). Frass and greenfall inputs to the soil may result in increased nitrogen mineralization (Constantinides and Fownes 1994; Lightfoot and Whitford 1990; Reynolds et al. 2000) leading to enhanced microbial immobilization (Lovett and Ruesink 1995; Christenson et al. 2002; Lovett et al. 2002) or export from the system (Swank and Crossley 1988; Eshleman et al. 1998). Nitrogen export may occur rapidly (e.g., within 1 week) if timed with a rainfall event, but in most situations the nitrogen is immobilized by microbes or fungi near the soil surface (top 0–30 cm) and is not available for plant uptake (Christenson et al. 2002; Lovett et al. 2002; Frost and Hunter 2007).

The effects of herbivore inputs on soil carbon and nitrogen dynamics depend on both input quality and quantity. Input quality may change due to the direct effects of atmospheric chemistry on foliar chemistry, which, in turn, affects the chemistry of both frass and greenfall. Knepp et al. (2007) demonstrated that under eCO₂ *Polyphemus* caterpillars have altered carbon and nitrogen concentrations and caterpillar frass has altered elemental composition, phenolic content and carbon to nitrogen ratios. Input quantity may change due to the effects of foliar chemistry on the ingestion and egestion rates of individual herbivores (Lindroth 1996), as well as on herbivore population density (Liebhold and Elkinton 1988). Of these, changes in population densities are likely particularly important, as research has documented increased effects of herbivore inputs at higher population densities (Hunter et al. 2003).

Given that atmospheres of the future are likely to alter the quality and quantity of frass and greenfall inputs, we investigated how eCO₂ and eO₃ would affect carbon respiration of soil microorganisms and soil nitrate leaching. We employed an experimental microcosm approach, using different quantities of whitemarked tussock moth (*Orgyia leucostigma* Fitch) frass and greenfall, both of which varied in quality due to atmospheric chemistry treatments administered at the Aspen Free Air CO₂ Enrichment (FACE) facility near Rhinelander, WI, USA.

Methods

Aspen FACE facility

The Aspen FACE site consists of 12 circular plots (rings), each 30 m in diameter. During the growing season, each ring receives one of two levels of CO₂ (ambient ~350 ppm, aCO₂; elevated ~550 ppm) and O₃ (ambient ~35 ppb, aO₃; elevated ~50 ppb). Thus, at the level of atmospheric gas treatment, this is a two-way factorial experiment with three replicates of each of the four treatment combinations. Fumigation of trees began in 1998 with eCO₂ and eO₃ dispensed at concentrations predicted for 2060. Elevated O₃ is fumigated at approximately 1.5 times that of monitored, daily, ambient levels (Dickson et al. 2000).

Frass and greenfall

In August 2006, we reared whitemarked tussock moth larvae from eggs obtained from the Canadian Forest Service (Sault St. Marie, ON, Canada), at 22°C in a Percival® growth chamber. The whitemarked tussock moth was chosen because it is a common generalist, native herbivore in this system. This tussock moth is not considered an outbreak species, but has similar feeding behavior and gut and frass chemistry to the gypsy moth (*Lymantria dispar* L.) (Kopper et al. 2002; Lovett et al. 1998; Frost and Hunter 2007). Larvae were fed paper birch foliage from trees at Aspen FACE that were not exposed to the fumigation treatments. Once larvae had reached the third instar, 10 individuals were put into a mesh enclosure on each of four aspen (*Populus tremuloides* Michx., clone 216) and four birch (*Betula papyrifera* Marsh.) trees in each of the 12 rings. Larvae were then allowed to feed until pupation. Frass was collected from fifth-instar caterpillars, air-dried, and weighed. Equal amounts (1:1 mix by dry mass) of frass from aspen- and birch-fed caterpillars were then combined into a single aggregate pool for each FACE ring (three replicate FACE rings per fumigation treatment). “Greenfall” was artificially generated for this study using green leaves collected from each FACE ring in August 2006. To preserve chemical characteristics, field collected leaves were stored on crushed ice while in the field, flash frozen in liquid nitrogen, and freeze-dried. Freeze-dried leaves easily crumbled into small pieces. We selected pieces roughly 0.1 to

0.5 cm² for this experiment. Equal amounts of aspen and birch foliage were weighed and pooled (1:1 mix by dry mass) for each FACE ring.

Subsamples of frass and composite greenfall were measured for carbon and nitrogen concentration (% dry mass) with a Thermo Finnigan (San Jose, CA, USA) Flash 1112 elemental analyzer. Condensed tannins of frass and greenfall were extracted with 70% acetone at 4°C and quantified by the acid butanol method of Porter et al. (1986), with a purified 1:1 mix of aspen and birch tannin standards. We do not report lignin concentrations in this study because the fine particulate nature of frass caused too much loss of material during the standard lignin digestion procedure.

Microcosms

We set up seven microcosms for each of the 12 FACE rings. Microcosms consisted of pieces of clear acrylic plastic (18 cm in height and 6 cm in diameter) enclosed at the bottom by 1 mm nylon mesh and a clear plastic funnel. Each microcosm was filled to 10 cm depth with 150 g (dry mass) of field moist soil mix. This mix contained 100 g of topsoil from a non-fumigated portion of the Aspen FACE site and 50 g of sand, added to improve drainage. On top of the soil, we placed 1.5 g of chopped, air-dried leaf litter to simulate natural soil conditions at Aspen FACE. This litter was also collected from a non-fumigated portion of the FACE facility during August 2006, and was a mixture of partially decomposed aspen and birch leaves from previous growing seasons. Soil and old leaf litter were defaunated by repeated (three times) freezing and thawing. Sand was sterilized using an autoclave.

After adding soil and old leaf litter, each microcosm was flushed with 30 ml of double distilled water to rehydrate the soil and litter. After four days, either a low or high quantity of frass, greenfall, or a mix of frass and greenfall (1:1 mix by dry mass) was added to each microcosm (Table 1). Input quantities were based on estimates of gypsy moth herbivory in mature aspen stands calculated from aspen specific leaf area, leaf area index, litterfall and gypsy moth digestion efficiencies (Raich and Nadelhoffer 1989; Lindroth and Hwang 1996; Steele et al. 1997; Roth et al. 1998; Burrows et al. 2002; Davidson et al. 2002; Madritch et al. 2007a). The 100 mg treatments represent frass and greenfall inputs from background canopy herbivory of less than 20% leaf area consumed while the

Table 1 Frass and greenfall treatments applied to the soil microcosms. Each of the seven treatments was applied to one microcosm for each of the 12 FACE rings for a total of 84 microcosms

	Treatment	Abbreviation	Quantity (mg)
1	Control	C	0
2	Frass (low)	FL	100
3	Greenfall (low)	GL	100
4	Frass + Greenfall (low)	FGL	50+50
5	Frass (high)	FH	700
6	Greenfall (high)	GH	700
7	Frass + Greenfall (high)	FGH	350+350

700 mg treatments represent frass and greenfall inputs from outbreak herbivory levels of greater than 70% leaf area consumed. After substrates were added to the microcosms, we added 20 ml of distilled water to rehydrate the inputs. Microcosms were incubated at room temperature (19°C) for 40 days.

Soil carbon and nitrogen mineralization

On days 1–5, 7, 9, 11, 16, 18, 21, 24, 30, and 37 we measured CO₂ efflux from each of the 84 microcosms (12 rings × 7 microcosms) for two minutes using a PP Systems (Amesbury, MA) infrared gas analyzer. Five milliliters of water were added to each microcosm approximately 24 h before each CO₂ measurement and as necessary between measurements to maintain soil moisture levels. The amount of water added to each microcosm over the entirety of the study approximated the average summer monthly rainfall per 37 days in northern Wisconsin. We report CO₂ respired in conventional units of g/m²/hr.

After the 40 day incubation period, microcosms were flushed with 50 ml distilled water. Leachate was collected by drawing a vacuum through each microcosm and was analyzed for combined NO₂ and NO₃ concentration using a modified version of the vanadium chloride method (Miranda et al. 2001; Doane and Horwath 2003; Allison et al. 2008). Leachate was analyzed for NH₄ concentration using a modified version of the Berthelot-salicylate method (Weatherburn 1967; Madritch et al. 2007a; Allison et al. 2008). Ammonium concentration is typically much lower than nitrate concentration in leachate due to lower mobility. Nitrogen concentra-

tion of leachate is reported as NO₃-N and NH₄-N loss (mg/L) per liter of leachate.

Statistical analysis

Average carbon, nitrogen, and condensed tannins in frass and greenfall pools from the 12 FACE rings were analyzed with a blocked, two-way ANOVA, where effects were block, CO₂, O₃, and CO₂ × O₃ interaction. CO₂ respiration and nitrogen leaching results were analyzed using a blocked, split-plot ANOVA. Whole plot effects were related to input quality and included experimental block, CO₂, O₃, and CO₂ × O₃ interaction. At the sub-plot level, effects were microcosm treatment (Table 1), CO₂ × microcosm treatment, O₃ × microcosm treatment, and CO₂ × O₃ × microcosm treatment. Correlation analysis was used to reveal the pattern of relationship between nitrate leaching and soil respiration. Analyses were conducted using JMP® 8.0.1 software (SAS Institute, Inc, Cary, NC, USA). P-values are considered statistically significant at $\alpha \leq 0.05$ and marginally significant for $0.05 < \alpha < 0.1$ (Filion et al. 2000).

Results

Input quality

FACE treatments significantly affected frass quality (Table 2). Frass carbon concentration was 1% lower, nitrogen concentration was unaltered, and condensed tannin concentration was 41% higher in eCO₂ plots compared with aCO₂ plots. Frass carbon concentration was not affected by eO₃, but nitrogen concentration was reduced 12% and condensed tannin concentration was increased 30% at eO₃ compared with aO₃. However, the 30% increase in tannins at eO₃ was not statistically significant.

Elevated CO₂ and O₃ also affected green leaf quality (Table 2). Greenfall carbon concentration was 1% lower in eCO₂ plots compared with aCO₂ plots. Greenfall nitrogen and condensed tannin concentrations were not altered by eCO₂. Elevated O₃ increased greenfall carbon concentration 1%. Greenfall nitrogen concentration decreased 9% and condensed tannin concentration increased 13% compared with aO₃ but the decrease in tannins was not significant. For comparison, greenfall, frass, and leaf litter collected

Table 2 Mean (± 1 SE) percent carbon, nitrogen, and condensed tannins of frass and greenfall added to microcosms. Statistics represent the results of two-way crossed ANOVA tests (df 1,6). Significant P-values (<0.10) are indicated in bold

	Control	+CO ₂	+O ₃	+CO ₂ +O ₃	CO ₂		O ₃		CO ₂ \times O ₃	
					F	P	F	P	F	P
Frass carbon	50.21 \pm 0.25	49.61 \pm 0.16	50.24 \pm 0.09	49.61 \pm 0.21	10.52	0.02	0.01	0.94	0.01	0.95
Frass nitrogen	1.18 \pm 0.07	1.12 \pm 0.06	0.99 \pm 0.04	1.05 \pm 0.05	0.00	0.99	5.34	0.06	1.12	0.33
Frass tannins	6.83 \pm 0.62	11.32 \pm 1.53	10.49 \pm 2.20	12.93 \pm 1.96	4.00	0.09	2.33	0.17	0.38	0.56
Greenfall carbon	49.95 \pm 0.11	49.5 \pm 0.11	50.36 \pm 0.07	50.1 \pm 0.23	4.91	0.07	10.23	0.02	0.39	0.56
Greenfall nitrogen	1.87 \pm 0.02	1.88 \pm 0.04	1.59 \pm 0.04	1.83 \pm 0.13	2.77	0.15	5.51	0.06	2.48	0.17
Greenfall tannins	17.32 \pm 2.37	20.01 \pm 3.34	23.44 \pm 4.43	18.82 \pm 2.53	0.08	0.79	0.5	0.51	1.09	0.34

from the FACE rings had carbon concentrations of 50%, 50%, and 49%, respectively, and nitrogen concentrations of 1.8%, 1.0%, and 0.9%, respectively.

Soil processes

Consistent with the minimal effects of FACE CO₂ and O₃ treatments on input quality, we found only minor effects on microbial respiration (Table 3; Fig. 1a, c). We found no CO₂ or O₃ main effects on microbial respiration averaged across microcosm treatments.

Table 3 Two-way crossed split-plot ANOVA of microbial respiration and nitrate leaching from microcosms. P-values in bold are significant at $\alpha \leq 0.05$. Microcosm trt. (= treatment) includes the seven microcosm treatments from Table 1

Source	F	df	P
Microbial respiration			
CO ₂	1.92	1,6	0.21
O ₃	1.04	1,6	0.35
CO ₂ \times O ₃	0.99	1,6	0.36
Microcosm trt.	21.57	6,48	<0.01
CO ₂ \times microcosm trt.	2.05	6,48	0.08
O ₃ \times microcosm trt.	0.30	6,48	0.93
CO ₂ \times O ₃ \times microcosm trt.	1.91	6,48	0.10
Nitrate-N leaching			
CO ₂	0.53	1,6	0.50
O ₃	0.14	1,6	0.72
CO ₂ \times O ₃	0.10	1,6	0.77
Microcosm trt.	6.40	6,48	<0.01
CO ₂ \times microcosm trt.	0.92	6,48	0.49
O ₃ \times microcosm trt.	1.60	6,48	0.17
CO ₂ \times O ₃ \times microcosm trt.	0.90	6,48	0.50

Microbial respiration was marginally affected by the interaction of CO₂ and microcosm treatment ($F_{6,48} = 2.05$, $P=0.08$). We found a trend for larger differences in CO₂ respiration between low and high inputs at eCO₂ compared with aCO₂. Microcosms with the FGL treatment had greater carbon respiration than control microcosms at aCO₂ but similar carbon respiration to FGL microcosms at eCO₂. Microbial respiration was not significantly different between greenfall, frass, and combinations of the two at the same quantity of input (Fig. 1b, c) even though the substrates differed slightly in terms of carbon and nitrogen concentrations.

In contrast to input quality, input quantity had a substantial positive effect on microbial respiration (Table 3, Fig. 1b, c). Losses from microcosms receiving small amounts of herbivore inputs were larger (0.079 ± 0.006 g CO₂/m²/hr) but not significantly different from those receiving no herbivore inputs (0.066 ± 0.004 g CO₂/m²/hr; Fig. 1b). However, respiratory carbon losses from microcosms receiving large amounts of herbivore inputs were nearly double (0.126 ± 0.007 g CO₂/m²/hr) those receiving no herbivore inputs (Table 3, Fig. 1b).

Nitrate loss from microcosms was unaffected by the CO₂ or O₃ treatments (Table 3, Fig. 2b, c) but was negatively related to input quantity. Nitrate concentrations of leachate from microcosms receiving small amounts of herbivore inputs was lower, but not significantly different from those receiving no herbivore inputs (72.5 ± 8.9 vs. 80.3 ± 6.9 mg/L; Fig. 2b). Nitrate concentrations of leachate from microcosms receiving large amounts of herbivore inputs were approximately half (38.5 ± 8.0 mg/L) those receiving low herbivore inputs. However, only GH and FGH




Fig. 1 Effects of FACE fumigation treatments (CO₂, O₃) and organic inputs (frass, greenfall) on soil respiration (± 1 SE). ANOVA results are presented in Table 3. **a** effects of fumigation treatments, pooled across microcosm treatments. **b** effects of microcosm treatment, pooled across fumigation treatments. **c** effects of individual fumigation and microcosm treatments. Abbreviations for microcosm treatments are provided in Table 1

inputs (not FH) were significantly different from no input (Table 3, Fig. 2b). We found a trend for a greater reduction in nitrate leaching in microcosms with greenfall compared with frass organic inputs across fumigation treatments (Fig. 2b, c). We also found a clear negative correlation between respiratory carbon loss and nitrate leaching across microcosm treatments (Fig. 3).

Relatively little ammonium nitrogen leached from the soil microcosms. The average concentration of NH₄-N in leachate across all microcosms was 0.058 ± 0.003 mg/L, only 1% of the concentration of nitrate. Neither input quality nor quantity influenced rates of ammonium leaching.




Fig. 2 Effects of FACE fumigation treatments (CO₂, O₃) and organic inputs (frass, greenfall) on nitrate concentration of leachate (± 1 SE). ANOVA results are presented in Table 3. **a** effects of fumigation treatments, pooled across microcosm treatments. **b** effects of microcosm treatment, pooled across fumigation treatments. **c** effects of individual fumigation and microcosm treatments. Abbreviations for microcosm treatments are provided in Table 1

Discussion

Effects of input quality and quantity on soil processes

Changes in frass and greenfall quality due to changes in atmospheric chemistry were sufficiently subtle that they did not significantly affect carbon and nitrogen mineralization. Differences in input quality, in terms of whether it was frass or greenfall, also had minor effects on soil carbon and nitrogen mineralization. In contrast, variation in the quantity of herbivore inputs strongly influenced carbon and nitrogen mineralization rates.




Fig. 3 Correlation of microbial respiration and nitrate leached for the 84 soil microcosms. Each point represents one microcosm

Elevated CO_2 and O_3 had relatively minor effects on frass and greenfall carbon and nitrogen concentrations. Condensed tannin levels increased in leaves exposed to eCO_2 and eO_3 . Caterpillar frass also had increased amounts of tannins at eCO_2 and eO_3 . Previous work with gypsy moth and forest tent caterpillars found that frass carbon to nitrogen ratios and condensed tannin levels reflected the carbon to nitrogen ratio and condensed tannin levels of the aspen foliage on which they fed (Madritch et al. 2007a). Frass of whitemarked tussock moth caterpillars fed birch condensed tannins also reflected the concentration of tannins in their food source (Kopper et al. 2002). These results suggest that if eCO_2 or eO_3 alter host plant chemical composition, frass chemical composition will be similarly altered. Altered chemical composition of frass, especially altered condensed tannin levels, could significantly alter soil processes because condensed tannins have been shown to reduce nutrient availability and slow leaf litter decomposition (Schweitzer et al. 2008) and alter soil respiration (Madritch et al. 2007b). Elevated CO_2 has been shown to reduce leaf litter quality (e.g., increased tannin concentrations and carbon to nitrogen ratios) at Aspen FACE, resulting in slower decomposition (Parsons et al. 2004, 2008). Similarly, lower frass or greenfall quality could impact soil processes, although the mostly weak effects of eCO_2 and eO_3 on input quality in this study were insufficient to do so.

Our results show that low levels of herbivore inputs (inputs from <20% leaf area removed) had a minor effect on soil microbial respiration, but high levels of herbivore inputs (inputs from >70% leaf area removed) nearly doubled microbial respiration. Pre-

vious studies suggest that small input levels can significantly alter soil processes, but do so less often than when inputs are large (e.g., during outbreaks) (Hunter et al. 2003). Only moderate or high input levels may provide enough labile carbon to allow detection of increased soil respiration. Both our results and those of Lovett and Ruesink (1995) support the concept that soil microbes do not respond unless frass inputs reach some carbon threshold. However, the response at even high input levels is likely to be short-lived (Frost and Hunter 2004). Soil temperature and precipitation events likely set the stage for the strength of the effect of herbivore inputs on soil microbe respiration rates. For example, high humidity or precipitation leaches frass carbon and nitrogen into the soil, while high temperature will increase microbial activity once inputs become available (Reynolds and Hunter 2001). Although we focused on microbial respiration, we recognize that part of the respiration we measured could be due to the frass microbial community and not the soil microbial community. We also recognize that the inclusion of leaf litter in our microcosms may have resulted in an early flush of CO_2 that masked smaller fumigation treatment effects on respiration.

Variation in input quantity also strongly influenced nitrate leaching rates. Nitrate leaching was relatively high at low-input levels, while microbial immobilization occurred at high-input levels. Frost and Hunter (2007) suggest that significant amounts of frass nitrogen may be leached if frass inputs are timed with precipitation, but otherwise the nitrogen will be incorporated into soil organic matter. Indeed, frass nitrogen may rapidly move into nonexchangeable soil pools near the soil surface (top 30 cm) (Christenson et al. 2002). It appears, therefore, that leaching losses likely occur shortly after deposition but most frass nitrogen is rapidly immobilized (Frost and Hunter 2007). Our results add that the quantity of organic input may alter whether nitrate is leached from, or immobilized in, the soil. The clear negative correlation we found between respiratory carbon loss and nitrate leaching across microcosm treatments strongly supports the concept that only moderate to high carbon inputs supply enough labile carbon for soil microbial populations to grow and increase nitrate immobilization (Fig. 3). Regardless, frass nitrogen inputs change the timing of nitrogen input to the forest floor and can alter the ultimate distribution of

nitrogen, even within the season of defoliation (Frost and Hunter 2004). For example, Frost and Hunter (2007) found that nitrogen from frass deposited early in the summer was taken up by plants within the same growing season. Approximately 1.2% of the frass nitrogen ended up in senescent leaves of the same trees at the end of the growing season, linking the fast (frass) and slow (senescent leaves) nutrient cycles. However, plants may acquire senesced leaf nitrogen more easily than frass nitrogen (Christenson et al. 2002).

Greenfall, respiration, and nitrate leaching

Our results suggest that, similar to large frass inputs, microbial respiration is enhanced by large inputs of greenfall. In another study, Reynolds and Hunter (2001) found reduced microbial respiration in forest soils where greenfall was excluded. While soil respiration was similar after frass and greenfall inputs, nitrate immobilization appeared to be slightly, although not significantly greater, for greenfall than for frass. The 65% higher nitrogen concentration of greenfall compared with frass in our study may have afforded greater microbial population growth and, thereby, increased immobilization. Reynolds and Hunter (2001) found no leaching of nitrate in their greenfall exclusion plots and suggested this may have been due to immobilization by roots or microbes. ^{15}N isotope studies have been useful for tracking frass N cycling and should be performed for greenfall given its high resource quality for microbes. However, as for frass, greenfall effects in the soil may be short-lived and dependent on temperature and precipitation.

Limitations

Microcosm studies are useful because they reduce environmental complexity and facilitate interpretation of results, but they are also less ecologically realistic (Carpenter 1996; Verhoef 1996). For example, our microcosms did not include detritivores or plant roots, and both will have important effects on soil respiration and nitrogen dynamics. Also, we did not consider other herbivore inputs, such as throughfall, which can influence soil processes (Stadler et al. 2001, 2006). We also recognize that some nutrients may have leached from our frass in the rain storm that occurred the day we collected frass. However, no other rain

events occurred during caterpillar rearing so very little leaching should have occurred. The nitrogen concentration of frass used in our study was approximately 1.0%, which is in line with values reported by Frost and Hunter (2007), but lower than the 2.4% documented by Lovett and Ruesink (1995) and the 1.8% reported by Madritch et al. (2007a). Our numbers may differ because results vary depending on the herbivore species used (Madritch et al. 2007a) and the time of sampling within a year. Insects feeding on early season foliage with greater concentrations of nitrogen and water and lower concentrations of defensive compounds (e.g., expanding leaves) could produce different results.

Implications of changes in herbivore input quantity

We found that soil processes were much more sensitive to the quantity of herbivore inputs than to the qualitative changes due to altered atmospheric chemistry. Liu et al. (2005) found that eCO_2 increased leaf litter inputs to the forest floor, and more recently that the increased inputs were more important to soil carbon and nitrogen cycles than were changes in litter chemistry due to eCO_2 (Liu et al. 2009). Larger inputs from the canopy also resulted in greater microbial respiration (King et al. 2004) and soil nitrogen concentration (Liu et al. 2009), in parallel with our large frass and greenfall inputs. These results suggest that altered atmospheric chemistry may have indirect effects on soil processes via changes in input quantity. Canopy damage rates at Aspen FACE have increased over 300% under eCO_2 , resulting in a more than 40% increase in frass and greenfall (Couture, Meehan, and Lindroth unpublished data). Thus, continued increases in eCO_2 in the future may increase the quantity of herbivore inputs, which may substantially impact nutrient cycling in forest soils.

Summary

Elevated CO_2 and eO_3 affected frass and greenfall quality, but the changes were minimal, producing little effect on short-term microbial respiration and no effect on nitrogen leaching. In contrast, large quantities of herbivore inputs nearly doubled microbial respiration and nitrogen immobilization compared with no inputs. Although microbial response to herbivore inputs may be relatively short-lived, herbi-

vore inputs are very important for carbon and nitrogen cycling, at least within the season of deposition. Changes in the abundance or feeding habits of canopy herbivores due to environmental change could lead to changes in soil carbon sequestration and nitrogen balance in forests of the future.

Acknowledgments Aspen FACE is principally supported by the Office of Science (BER), U.S. Department of Energy, Grant No. DE-FG02-95ER62125 to Michigan Technological University, and Contract No. DE-AC02-98CH10886 to Brookhaven National Laboratory, the U.S. Forest Service Northern Global Change Program and North Central Research Station, Michigan Technological University, and Natural Resources Canada—Canadian Forest Service. We thank the members of the Lindroth Lab, especially Mike Madritch, for input into study design and lab techniques, Adam Gusse and Andy Vogelzang for help raising whitemarked tussock moth larvae, and Leanne Vigue for collecting senesced leaves from the Aspen FACE site. We also thank the two anonymous reviewers for their helpful comments on the manuscript. This work was supported by U.S. Department of Energy (Office of Science, BER) grant DE-FG02-06ER64232, and University of Wisconsin Hatch grant WIS04898 to R.L. Lindroth.

References

- Akimoto H (2003) Global air quality and pollution. *Science* 302:1716–1719
- Allison SD, Czimczik CI, Treseder KK (2008) Microbial activity and soil respiration under nitrogen addition in Alaskan boreal forest. *Glob Chang Biol* 14:1156–1168
- Belovsky GE, Slade JB (2000) Insect herbivory accelerates nutrient cycling and increases plant production. *Proc Natl Acad Sci* 97:14414–14417
- Bezemer TM, Jones TH (1998) Plant-insect herbivore interactions in elevated atmospheric CO₂: quantitative analyses and guild effects. *Oikos* 82:212–222
- Burrows SN, Gower ST, Clayton MK, Mackay DS, Ahl DE, Norman JM, Diak G (2002) Application of geostatistics to characterize leaf area index (LAI) from flux tower to landscape scales using a cyclic sampling design. *Ecosystems* 5:667–679
- Carpenter SR (1996) Microcosm experiments have limited relevance for community and ecosystem ecology. *Ecology* 77:677–690
- Christenson LM, Lovett GM, Mitchell MJ, Groffman PM (2002) The fate of nitrogen in gypsy moth frass deposited to an oak forest floor. *Oecologia* 131:444–452
- Constantinides M, Fownes JH (1994) Nitrogen mineralization from leaves and litter of tropical plants: relationship to nitrogen, lignin, and soluble polyphenol concentrations. *Soil Biol Biochem* 26:49–55
- Davidson EA, Savage K, Bolstad P, Clark DA, Curtis PS, Ellsworth DS, Hanson PJ, Luo Y, Pregitzer KS, Randolph JC, Zak DR (2002) Belowground carbon allocation in forests estimated from litterfall and IRGA-based soil respiration measurements. *Agric For Meteorol* 113:39–51
- Dickson RE, Lewin KF, Isebrands JG, Coleman MD, Heilman WE, Riemenschneider DE, Sober J, Host GE, Zak DR, Hendry GR, Pregitzer KS, Karnosky DF (2000) Forest atmosphere carbon transfer storage-II (FACTS II)—the aspen free-air CO₂ and O₃ enrichment (FACE) project: an overview, General Technical Report NC-214, USDA Forest Service, North Central Research Station, Rhineland, WI
- Doane TA, Horwath WR (2003) Spectrophotometric determination of nitrate with a single reagent. *Analytical Letters* 36:2713–2722
- Eshleman KN, Morgan RP II, Webb JR, Deviney FA, Galloway JN (1998) Temporal patterns of nitrogen leakage from mid-Appalachian forested watersheds: role of insect defoliation. *Water Resour Res* 34:2005–2016
- Filion M, Dutilleul P, Potvin C (2000) Optimum experimental design for Free-Air Carbon Dioxide Enrichment (FACE) studies. *Glob Change Biol* 6:843–854
- Fogal WH, Slansky F Jr (1984) Contribution of feeding by European pine sawfly larvae to litter production and element flux in Scots pine plantations. *Can J For Res* 15:484–487
- Fonte SJ, Schowalter TD (2004) Decomposition of greenfall vs. senescent foliage in a tropical forest ecosystem in Puerto Rico. *Biotropica* 36:474–482
- Frost CJ, Hunter MD (2004) Insect canopy herbivory and frass deposition affect soil nutrient dynamics and export in oak mesocosms. *Ecology* 85:3335–3347
- Frost CJ, Hunter MD (2007) Recycling of nitrogen in herbivore feces: plant recovery, herbivore assimilation, soil retention, and leaching losses. *Oecologia* 151:42–53
- Grace JR (1986) The influence of gypsy moth on the composition and nutrient content of litter fall in a Pennsylvania oak forest. *For Sci* 32:855–870
- Hollinger DY (1986) Herbivory and the cycling of nitrogen and phosphorus in isolated California oak trees. *Oecologia* 70:291–297
- Holton MK, Lindroth RL, Nordheim EV (2003) Foliar quality influences tree-herbivore-parasitoid interactions: effects of elevated CO₂, O₃, and genotype. *Oecologia* 137:233–244
- Hunter MD (2001) Insect population dynamics meets ecosystem ecology: effects of herbivory on soil nutrient dynamics. *Agric For Entomol* 3:77–84
- Hunter MD, Linnen CR, Reynolds BC (2003) Effects of endemic densities of canopy herbivores on nutrient dynamics along a gradient in elevation in the southern Appalachians. *Pedobiologia* 47:231–244
- Intergovernmental Panel on Climate Change (2007) Climate Change 2007: The Physical Scientific Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge
- Karnosky DF (1976) Threshold levels for foliar injury to *Populus tremuloides* Michx. by sulfur dioxide and ozone. *Can J For Res* 6:166–169
- Karnosky DF, Zak DR, Pregitzer KS, Awmack CS, Bockheim JG, Dickson RE, Hendrey GR, Host GE, King JS, Kopper BJ, Kruger EL, Kubiske ME, Lindroth RL, Mattson WJ, McDonald EP, Noormets A, Oksanen E, Parsons WFJ,

- Percy KE, Podila GK, Riemenschneider DE, Sharma P, Thakur RC, Sober A, Sober J, Jones WS, Anttonen S, Vapaavuori E, Mankovska B, Heilman WE, Isebrands JG (2003) Tropospheric O₃ moderates responses of temperate hardwood forests to elevated CO₂: a synthesis of molecular to ecosystem results from the Aspen FACE project. *Funct Ecol* 17:289–304
- King JS, Hanson PJ, Bernhardt E, DeAngelis P, Norby RJ, Pregitzer KS (2004) A multiyear synthesis of soil respiration responses to elevated atmospheric CO₂ from four forest FACE experiments. *Glob Change Biol* 10:1027–1042
- Knepp RG, Hamilton JG, Zangerl AR, Berenbaum MR, DeLucia EH (2007) Foliage of oaks grown under elevated CO₂ reduced performance of *Antheraea polyphemus* (Lepidoptera: Saturniidae). *Environ Entomol* 36:609–617
- Kopper BJ, Lindroth RL (2003) Effects of elevated carbon dioxide and ozone on the phytochemistry of aspen and performance of an herbivore. *Oecologia* 134:95–103
- Kopper BJ, Jakobi VN, Osier TL, Lindroth RL (2002) Effects of paper birch condensed tannin on whitemarked tussock moth (Lepidoptera: Lymantriidae) performance. *Environ Entomol* 31:10–14
- Liebold AM, Elkinton JS (1988) Techniques for estimating the density of late-instar gypsy moth, *Lymantria dispar* (Lepidoptera: Lymantriidae), populations using frass drop and frass production measurements. *Environ Entomol* 17:381–384
- Lightfoot DC, Whitford WG (1990) Phytophagous insects enhance nitrogen flux in a desert creosote bush community. *Oecologia* 82:18–25
- Lindroth RL (1996) Consequences of elevated atmospheric CO₂ for forest insects. In: Korner C, Bazzaz FA (eds) Carbon dioxide, populations, and communities. Physiological Ecology Series, Academic Press, San Diego, pp 347–361
- Lindroth RL (2010) Impacts of elevated atmospheric CO₂ and O₃ on forests: phytochemistry, trophic interactions, and ecosystem dynamics. *J Chem Ecol* 36:2–21
- Lindroth RL, Hwang SY (1996) Clonal variation in foliar chemistry of quaking aspen (*Populus tremuloides* Michx). *Biochem Syst Ecol* 24:357–364
- Lindroth RL, Roth S, Nordheim EV (2001) Genotypic variation in response of quaking aspen (*Populus tremuloides*) to atmospheric CO₂ enrichment. *Oecologia* 126:371–379
- Liu LL, King JS, Giardina CP (2005) Effects of elevated concentrations of atmospheric CO₂ and tropospheric O₃ on leaf litter production and chemistry in trembling aspen and paper birch communities. *Tree Physiol* 25:1511–1522
- Liu LL, King JS, Booker FL, Giardina CP, Allen HL, Hu S (2009) Enhanced litter input rather than changes in litter chemistry drive soil carbon and nitrogen cycles under elevated CO₂: a microcosm study. *Glob Change Biol* 15:441–453
- Lovett GM, Ruesink AE (1995) Carbon and nitrogen mineralization from decomposing gypsy moth frass. *Oecologia* 104:133–138
- Lovett GM, Hart JE, Christenson LM, Jones CG (1998) Caterpillar gut and ammonia volatilization: retention of nitrogen by gypsy moth larvae consuming oak foliage. *Oecologia* 117:513–516
- Lovett GM, Christenson LM, Groffman PM, Jones CG, Hart JE, Mitchell JM (2002) Insect defoliation and nitrogen cycling in forests. *Bioscience* 52:335–341
- Madritch MD, Donaldson JR, Lindroth RL (2007a) Canopy herbivory can mediate the influence of plant genotype on soil processes through frass deposition. *Soil Biol Biochem* 29:1192–1201
- Madritch MD, Jordan LM, Lindroth RL (2007b) Interactive effects of condensed tannin and cellulose additions on soil respiration. *Can J For Res* 37:2063–2067
- Miranda KM, Espey MG, Wink DA (2001) A rapid, simple spectrophotometric method for simultaneous detection of nitrate and nitrite. *Nitric Oxide-Biol Chem* 5:62–71
- Parsons WFJ, Lindroth RL, Bockheim JG (2004) Decomposition of *Betula papyrifera* leaf litter under the independent and interactive effects of elevated CO₂ and O₃. *Glob Change Biol* 10:1666–1677
- Parsons WFJ, Bockheim JG, Lindroth RL (2008) Independent, interactive, and species-specific responses of leaf litter decomposition to elevated CO₂ and O₃ in a northern hardwood forest. *Ecosystems* 11:505–519
- Porter LJ, Hrstich LN, Chan BG (1986) The conversion of procyanidins and prodelphinidins to cyaniding and delphinidin. *Phytochemistry* 25:223–230
- Raich JW, Nadelhoffer KJ (1989) Belowground carbon allocation in forest ecosystems: global trends. *Ecology* 70:1346–1354
- Reynolds BC, Hunter MD (2001) Responses of soil respiration, soil nutrients, and litter decomposition to inputs from canopy herbivores. *Soil Biol Biochem* 33:1641–1652
- Reynolds BC, Hunter MD, Crossley DA Jr (2000) Effects of canopy herbivory on nutrient cycling in a northern hardwood forest in western North Carolina. *Selbyana* 21:74–78
- Risely LS, Crossley DA Jr (1993) Contribution of herbivore-caused greenfall to litterfall nitrogen flux in several southern Appalachian forested watersheds. *Am Midl Nat* 129:67–74
- Roth S, Lindroth RL, Volin JC, Kruger EL (1998) Enriched atmospheric CO₂ and defoliation: effects on tree chemistry and insect performance. *Glob Change Biol* 4:419–430
- Saxe H, Ellsworth DS, Heath J (1998) Tree and forest functioning in an enriched CO₂ atmosphere. *New Phytol* 139:395–436
- Schowalter TD (2000) Insect ecology: an ecosystem approach. Academic Press, New York
- Schweitzer JA, Bailey JK, Fischer DG, LeRoy CJ, Lonsdorf EV, Whitham TG, Hart SC (2008) Plant-soil-microorganism interactions: heritable relationship between plant genotype and associate soil microorganisms. *Ecology* 89:773–781
- Stadler B, Solinger S, Michalzik B (2001) Insect herbivores and the nutrient flow from the canopy to the soil in coniferous and deciduous forests. *Oecologia* 126:104–113
- Stadler B, Muller T, Orwig D (2006) The ecology of energy and nutrient fluxes in hemlock forests invaded by hemlock woolly adelgid. *Ecology* 87:1792–1804
- Steele SJ, Gower ST, Vogel JG, Norman JM (1997) Root mass, net primary production and turnover in aspen, jack pine and black spruce forest in Saskatchewan and Manitoba, Canada. *Tree Physiol* 17:577–587
- Stiling P, Cornelissen T (2007) How does elevated carbon dioxide (CO₂) affect plant-herbivore interactions? A field

- experiment and meta-analysis of CO₂-mediated changes on plant chemistry and herbivore performance. *Glob Change Biol* 13:1–20
- Swank WT, Crossley DA Jr (1988) Forest ecology and hydrology at Coweeta. Springer, New York
- Valkama E, Koricheva J, Oksanen E (2007) Effects of elevated O₃, alone and in combination with elevated CO₂, on tree leaf chemistry and insect herbivore performance: a meta-analysis. *Glob Change Biol* 13:184–201
- Verhoef HA (1996) The role of soil microcosms in the study of ecosystem processes. *Ecology* 77:685–690
- Weatherburn MW (1967) Phenol-hypochlorite reaction for determination of ammonia. *Anal Chem* 39:971–974
- Wittig VE, Ainsworth EA, Naidu SL, Karnosky DF, Long SP (2009) Quantifying the impact of current and future tropospheric ozone on tree biomass, growth, physiology and biochemistry: a quantitative meta-analysis. *Glob Change Biol* 15:396–424
- Zvereva EL, Kozlov MV (2006) Consequences of simultaneous elevation of carbon dioxide and temperature for plant-herbivore interactions: a metaanalysis. *Glob Change Biol* 12:27–41