

3/10 Cyclotomic Extensions §5.8


Our goal now is to study finite extensions of \mathbb{Q} of specific forms, leading up to a treatment of the problem which motivated the development of Galois theory: the solubility of polynomials by radicals.

Def'n Recall that a number $u \in \mathbb{C}$ is called an n^{th} root of unity, for some $n \geq 1$, if $u^n = 1$, i.e., if u is a root of $X^n - 1 \in \mathbb{Q}[X]$. If u is an n^{th} root of unity, it is also a $(mn)^{\text{th}}$ root of unity for any $m \geq 1$. We say u is a primitive n^{th} root of unity if it is an n^{th} root of unity but not a k^{th} root of unity for any $k < n$.

Prop. The n^{th} roots of unity are $e^{\frac{2\pi i}{n}j}$ for $j=0,1,\dots,n-1$.

The primitive n^{th} roots of unity are those $e^{\frac{2\pi i}{n}j}$ with $\gcd(j,n)=1$.

- (*) E.g. We've seen before how the n^{th} roots of unity are equally spaced on the unit circle, for instance for $n=12$ we get



\Leftarrow the primitive 12^{th} roots of unity are circled; they are $e^{\frac{2\pi i}{12}j}$ for $j=1,5,7,11$, the integers coprime to 12.

Pf sketch of prop: That the $e^{\frac{2\pi i}{n}j}$ for $j=0,1,2,\dots,n-1$ are the n^{th} roots of unity follows from the fact that $e^{\frac{2\pi i}{n}j} \cdot e^{\frac{2\pi i}{n}k} = e^{\frac{2\pi i}{n}(j+k \bmod n)}$ (phases of complex #'s add when multiplied).

That the primitive ones are the coprime j 's then follows from $e^{\frac{2\pi i}{n}j}$ is a primitive n^{th} root of unity \Leftrightarrow

j is a generator of $(\mathbb{Z}/n\mathbb{Z}, +) \Leftrightarrow$

j is a unit in the ring $\mathbb{Z}/n\mathbb{Z} \Leftrightarrow$

j is coprime to n . You will flesh out this argument on your next HW assignment. \square

Notice: $\zeta_n = e^{\frac{2\pi i}{n}}$ is always a primitive n^{th} root of unity, and all n^{th} roots of unity are powers of this ζ_n .

Def'n Let $n \geq 1$. The n^{th} cyclotomic polynomial $\Phi_n(x) \in \mathbb{C}[x]$ is $\Phi_n(x) = \prod_{\omega \text{ a primitive } n^{\text{th}} \text{ root of unity}} (x - \omega)$ (The book uses $g_n(x)$.)

E.g.: The primitive 3rd roots of unity are $\omega = e^{\frac{2\pi i}{3}} = -\frac{1}{2} + \frac{\sqrt{3}}{2}i$ and $\omega^2 = -\frac{1}{2} - \frac{\sqrt{3}}{2}i$; so $\Phi_3(x) = (x - \omega)(x - \omega^2) = x^2 + x + 1$.

In fact, the first 6 cyclotomic polynomials are:

$$\Phi_1(x) = x - 1, \quad \Phi_2(x) = x + 1, \quad \Phi_3(x) = x^2 + x + 1, \quad \Phi_4(x) = x^2 + 1, \\ \Phi_5(x) = x^4 + x^3 + x^2 + x + 1, \quad \Phi_6(x) = x^2 - x + 1.$$

Thm $x^n - 1 = \prod_{d|n} \Phi_d(x)$

Pf: Every root of $x^n - 1$ is an n^{th} root of unity, which is a primitive d^{th} root of unity for some $d|n$. \square

Note: Even though $\Phi_d(x)$ is a priori defined as an element of $\mathbb{C}[x]$, books give it belongs to $\mathbb{Q}[x]$. This is true and we'll prove it!

In fact the coefficients are integers, which can get arbitrarily big, but take a while ($\Phi_{105}(x)$ is first with a coeff. not in $\{1, -1\}$).

The way we will show cyclotomic polynomials are rational is by studying the extensions of \mathbb{Q} we get by adjoining their roots.

Def'n The n^{th} cyclotomic extension of \mathbb{Q} is the splitting field of $x^n - 1$. Equivalently, ...

Thm The n^{th} cyclotomic extension is $\mathbb{Q}(\zeta_n)$, where ζ_n is a primitive n^{th} root of unity.

Pf: Since ζ_n is an n^{th} root of unity, it belongs to splitting field of $x^n - 1$.
But on other hand, every root of unity is a power of ζ_n , hence in $\mathbb{Q}(\zeta_n)$. \square

Thm Let $\Psi_k: \mathbb{Q}(\zeta_n) \rightarrow \mathbb{Q}(\zeta_n)$ be defined by $\Psi_k(\zeta_n) = \zeta_n^k$.

Then $\text{Aut}_{\mathbb{Q}}(\mathbb{Q}(\zeta_n)) \subseteq \{ \Psi_k : 1 \leq k \leq n, \gcd(n, k) = 1 \}$.

Pf: Any $\sigma \in \text{Aut}_{\mathbb{Q}}(\mathbb{Q}(\zeta_n))$ is determined by where it sends ζ_n , which must be to some ζ_n^k since these are roots of $x^n - 1$. But it cannot be sent to a non-primitive n^{th} root of unity, since it's not a root of any $x^m - 1$ (with $m < n$). \square

Cor The cyclotomic polynomial $\Phi_n(x) \in \mathbb{Q}[x]$.

Pf: $\mathbb{Q}(\zeta_n)$ is a Galois extension, since it's a splitting field, and every $\sigma \in \text{Aut}_{\mathbb{Q}}(\mathbb{Q}(\zeta_n))$ fixes $\Phi_n(x)$ since just permutes roots, so in fact coefficients of $\Phi_n(x)$ are rational. \square

Thm (Gauss) $\Phi_n(x)$ is irreducible over \mathbb{Q} .

Pf: This is non-trivial but I skip it - see the book. \square

Cor $\Phi_n(x)$ is the minimal polynomial of ζ_n , and every Ψ_k for $\gcd(n, k) = 1$ is indeed an element of $G = \text{Aut}_{\mathbb{Q}}(\mathbb{Q}(\zeta_n))$.
Hence $G \cong (\mathbb{Z}/n\mathbb{Z})^\times$, the multiplicative group mod n , via the isomorphism $\Psi_k \mapsto k \in (\mathbb{Z}/n\mathbb{Z})^\times$.

Remark: This shows $G \cong (\mathbb{Z}/n\mathbb{Z})^\times$ is an abelian group of order $\varphi(n)$ where $\varphi(n) = \# \{ 1 \leq k \leq n : \gcd(n, k) = 1 \}$

i) Euler's totient function. When $n = p$ is prime we have seen that $(\mathbb{Z}/p\mathbb{Z})^\times$ is in fact cyclic (of order $p-1$), but in general it need not be:

e.g. $(\mathbb{Z}/8\mathbb{Z})^\times \cong \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}$.

//

3/12

Cyclic Extensions § 5.7

We are almost ready to study the solvability of polynomials by radicals. We just need one more preparatory result...

Def'n An ^{algebraic, Galois} extension L/K is called abelian if $\text{Aut}_K(L)$ is abelian, it is called cyclic if $\text{Aut}_K(L)$ is cyclic, and it is called cyclic of degree n if $\text{Aut}_K(L) \cong \mathbb{Z}/n\mathbb{Z}$.

Remark: We have seen that the cyclotomic extension $\mathbb{Q}(\zeta_n)$ of \mathbb{Q} is always abelian, and sometimes cyclic (e.g. if n is prime) although not always.

In general it is hard to classify cyclic extensions, but there is a nice situation where we can do this.

Def'n For an arbitrary field K , $u \in K$ is called an n^{th} root of unity if $u^n = 1 \in K$ and is called a primitive n^{th} root of unity if u^0, u^1, \dots, u^{n-1} are all distinct (hence all the ^{are} n^{th} roots of unity).

For subfields of \mathbb{C} , this agrees with our previous definition. Then let K be a field containing a primitive n^{th} root of unity ζ_n for some $n \geq 1$. Then the following are equivalent for L/K :

- 1) L/K is cyclic of degree d , for some $d \mid n$.
- 2) L/K is the splitting field of a polynomial of form $f(x) = x^n - a \in K[x]$, in which case $L = K(u)$ for u a root of $f(x)$.
- 3) L/K is splitting field of irreducible polynomial of form $f(x) = x^d - a$ for some $d \mid n$, in which case $L = K(u)$ for u root of $f(x)$.

E.g. Any degree 2 extension of \mathbb{Q} is the splitting field of a polynomial of the form $x^2 - d$ where d is not a square in \mathbb{Q} , and this extension has Galois group $\mathbb{Z}/2\mathbb{Z}$.

E.g. On a previous homework you showed that if $L = \mathbb{Q}(\omega, \sqrt[3]{2})$ ^{prim. 3rd root of unity} is the splitting field of $x^3 - 2$ over \mathbb{Q} , then $\text{Aut}_K(L) \cong S_3$, which is not cyclic (not even abelian!). But \mathbb{Q} does not have a prim. 3rd root of unity! If we instead take $K = \mathbb{Q}(\omega)$, then $\text{Aut}_K(L) = \mathbb{Z}/3\mathbb{Z}$.

In the theorem, 2) and 3) are easily seen to be equivalent, just having to do with whether $x^n - a$ is irreducible, equivalently, whether a has a d th root in K for some $d \mid n$. The main point is showing $3) \Leftrightarrow 1)$. In fact we will mostly come about $3) \Rightarrow 1)$, which we will prove now ~~and~~ we just need:

Lemma If K is a field with a primitive n th root of unity ζ , then for any $d \mid n$, $\eta = \zeta^{n/d}$ is a primitive d th root of 1.

And if L is an extension of K such that $u \in L$ is a root of $x^d - a \in K[x]$, then all the roots of $x^d - a$ are $u, \eta u, \eta^2 u, \dots, \eta^{d-1} u$ (all distinct).

Pf. Straight forward exercise. P3

Pf of $3) \Rightarrow 1)$ in thm: By the lemma, the roots of $x^d - a$ in L are $u, \eta u, \dots, \eta^{d-1} u$ where u is any root and $\eta = \zeta^{n/d}$ as above. So any $\sigma \in \text{Aut}_K(L)$ is determined by where it sends u (since $\eta \in K$ is fixed by σ). Since $x^d - a$ is irreducible, there must be some σ with $\sigma(u) = \eta u$, and this σ generates all of $\text{Aut}_K(L)$. Since $\sigma^k(u) = \eta^k u$, which give all the possible automorphisms in the Galois group by the previous sentence. QED

We will only sketch the ideas that go into the pt of 1) \Rightarrow 3):

Def'n Let L/K be a finite ^{Galois} extension, and suppose that $\text{Aut}_K(L) = \{\sigma_1, \dots, \sigma_n\}$. For any $u \in L$, the norm of u is $N(u) = \sigma_1(u) \cdot \sigma_2(u) \cdot \dots \cdot \sigma_n(u)$.

Eg. Let $K = \mathbb{R}$ and $L = \mathbb{C}$. Recall that $\text{Aut}_{\mathbb{R}}(\mathbb{C}) = \{1, \sigma\}$ where $\sigma: \mathbb{C} \mapsto \bar{\mathbb{C}}$ is complex conjugation. So the norm of $z = a + bi \in \mathbb{C}$ is $N(z) = z \cdot \bar{z} = a^2 + b^2$, usual complex norm.

Eg. For $K = \mathbb{Q}$ and $L = \mathbb{Q}(i)$, the same is true: the norm of $a + bi$ is $(a + bi)(a - bi) = a^2 + b^2 \in \mathbb{Q}$.

Prop. If L/K is a finite Galois extension, then the norm $N(u)$ of any $u \in L$ is an element of the base field K .

Pf. For any $\tau \in \text{Aut}_K(L)$, $\tau(N(u)) = \tau(\sigma_1(u) \cdot \sigma_2(u) \cdot \dots \cdot \sigma_n(u))$
 $= \tau(\sigma_1(u)) \cdot \dots \cdot \tau(\sigma_n(u)) = \sigma_{i_1}(u) \cdot \dots \cdot \sigma_{i_n}(u) = N(u)$

(where i_1, \dots, i_n is some permutation of $1, \dots, n$), so because L/K is Galois, $N(u) \in K$ as claimed. \square

Remark: We can define the norm for non-Galois extensions too, and it remains true that it belongs to the ground field, but it's a little more technical.

Another important property of the norm is multiplicativity:

Prop. We have $N(u) \cdot N(v) = N(uv)$ for $u, v \in L$.

Pf. Straight forward exercise. \square

The norm is particularly useful for cyclic extensions...

Thm (Hilbert Theorem 90) Let L/K be a finite cyclic ^(Galois) extension and let $\sigma \in \text{Aut}_K(L)$ be a generator of the Galois group. Then for $u \in L$, $N(u) = 1 \Leftrightarrow u = v/\sigma(v)$ for some $v \in L$.

Pf: One direction is easy: if $u = \frac{v}{\sigma(v)}$ then $N(u) = \frac{\sigma_1(v) \cdots \sigma_n(v)}{\sigma_1(v) \cdots \sigma_n(v)} = 1$

The other direction is nontrivial - see the book for a proof. \square

Ex. Consider $L = \mathbb{Q}(i)$ over $K = \mathbb{Q}$. The elements in $\mathbb{Q}(i)$ of norm 1 are $\frac{p}{r} + \frac{q}{r}i$ with $\frac{p^2}{r^2} + \frac{q^2}{r^2} = 1$, i.e., $p^2 + q^2 = r^2$, $p, q, r \in \mathbb{Z}$.

These are Pythagorean triples. Hilbert's Thm 90 says they can all be written in form $\frac{a+bi}{a-bi} = \frac{a^2-b^2}{a^2+b^2} + \frac{2ab}{a^2+b^2}i$, $a, b \in \mathbb{Z}$

It is a classic fact going back to Euclid that (primitive) Pythagorean triples can be parameterized in this way.

With Hilbert's thm 90 we can complete the pf of main thm.

Pf of 1) \Rightarrow 3): Let $\sigma \in \text{Aut}_K(L)$ be a generator, and let $\eta = \zeta^{1/d}$ be a primitive d th root of unity.

Then $N(\eta) = \eta \cdot \sigma(\eta) \cdots \sigma^{d-1}(\eta) = \eta^d$ (since $\eta \in K$)

so by Hilbert 90 we can write $\eta = v/\sigma(v)$

for some $v \in L$. Notice $\sigma(v^d) = (\sigma(v))^d = \left(\frac{v}{\eta}\right)^d = \frac{v^d}{\eta^d} = v^d$

(since $\eta^d = 1$), so because the extension L/K is Galois,

this means that $v^d \in K$. Then v is a root of

the polynomial $x^d - v^d \in K[x]$, and it can be

shown that this polynomial is in fact irreducible

over K and that $L = K(v)$ is the splitting field. \square