


2/23

Arc lengths of curves § 8.1

Having studied techniques for integration, we return to applications of integrals. We've already used integrals to compute areas (2D measures), and volumes (3D measures), what about lengths (1D measures)?


Suppose we have a curve $y = f(x)$ from $x=a$ to $x=b$: what is the length of this curve? Of course, if the curve were a line $y = mx + c$ we could compute its length using the Pythagorean Theorem:

$$y = mb + c +$$



Notice: length depends on slope of line

But what if $y = f(x)$ is not a line? As usual, we break it into smaller parts where we treat it as approximately linear:



Thus, length of $y = f(x)$ from $x=a$ to $x=b$ is

$$\begin{aligned} &= \lim_{n \rightarrow \infty} \sum_{i=1}^n \sqrt{1 + \left(\frac{\Delta y_i}{\Delta x}\right)^2} \Delta x \\ &= \int_a^b \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx = \boxed{\int_a^b \sqrt{1 + (f'(x))^2} dx} \end{aligned}$$

In limit, $\frac{\Delta y_i}{\Delta x}$ becomes the derivative $\frac{dy}{dx}$

E.g.: If $f(x) = mx + c$ is a line, then $f'(x) = m$
 length from $x=a$ to $x=b$ $\Rightarrow \int_a^b \sqrt{1+(f'(x))^2} dx = \int_a^b \sqrt{1+m^2} dx = (b-a)\sqrt{1+m^2}$

E.g.: Consider the curve $y = x^{3/2}$ from $x=0$ to $x=1$.

$$\text{Length} = \int_0^1 \sqrt{1+(\frac{d}{dx} x^{3/2})^2} dx = \int_0^1 \sqrt{1+(3/2 x^{1/2})^2} dx \\ = \int_0^1 \sqrt{1+\frac{9}{4}x} dx \leftarrow \text{can solve w/ a u-sub.}$$

① indef. integral: $\int \sqrt{1+\frac{9}{4}x} dx = \int \sqrt{u} \frac{4}{9} du = \frac{4}{9} \cdot \frac{2}{3} u^{3/2} \\ u = 1 + \frac{9}{4}x \quad = \frac{8}{27} (1 + \frac{9}{4}x)^{3/2}$

② plug in: $\int_0^1 \sqrt{1+\frac{9}{4}x} dx = \left[\frac{8}{27} (1 + \frac{9}{4}x)^{3/2} \right]_0^1 = \frac{8}{27} ((\frac{13}{4})^{3/2} - 1)$

E.g.: Even for curve $y=x^2$ from $x=0$ to $x=1$, integral is nasty:

$$\text{Length} = \int_0^1 \sqrt{1+(\frac{d}{dx} x^2)^2} dx = \int_0^1 \sqrt{1+(2x)^2} dx = \int_0^1 \sqrt{1+4x^2} dx$$

① indef. integral: $\int \sqrt{1+4x^2} dx$ good idea: trig sub! $x = \frac{1}{2} \tan \theta$
 $= \int \sqrt{1+\tan^2 \theta} \frac{1}{2} \sec^2 \theta d\theta = \frac{1}{2} \int \sec^3 \theta d\theta$

But... $\int \sec^3 \theta d\theta$ is not easy! Int. by parts helps, but even then you still need to know $\int \sec \theta d\theta = \ln(\sec \theta + \tan \theta)$.

E.g.: Sometimes $(1+(f'(x))^2)$ has a square root:

$$\text{If } f(x) = \frac{1}{4}x^2 - \frac{1}{2}\ln(x) \text{ then } f'(x) = \frac{1}{2}x - \frac{1}{2x} = \frac{x^2 - 1}{2x}$$

$$\text{So } 1+(f'(x))^2 = 1 + \left(\frac{x^2-1}{2x}\right)^2 = 1 + \frac{x^4 - 2x^2 + 1}{4x^2} = \frac{x^4 + 2x^2 + 1}{4x^2} = \frac{(x^2+1)^2}{(2x)^2}$$

Thus, $\int \sqrt{1+(f'(x))^2} dx = \int \sqrt{\frac{(x^2+1)^2}{(2x)^2}} dx = \int \frac{x^2+1}{2x} dx = \int \frac{x}{2} dx + \int \frac{1}{2x} dx \\ = \frac{x^2}{4} + \frac{1}{2}\ln(x) + C$, so we get...


$$\int_1^e \sqrt{1+(f'(x))^2} dx = \left[\frac{x^2}{4} + \frac{1}{2}\ln(x) \right]_1^e = \frac{e^2}{4} + \frac{1}{4} \leftarrow \begin{array}{l} \text{length of} \\ y=f(x) \text{ from } x=1 \text{ to } x=e \end{array}$$

2/26

Area of Surface of Revolution § 8.2

Intuitively, the surface area of a solid is the amount of "wrapping paper" you would need to wrap it.


As usual, we start our discussion of surface area with simple shapes. Consider a cylinder of length l and radius r :



(Note: we do not consider area of left/right ends of cylinder, it is "open")


By cutting the cylinder and unwinding it into a rectangle we see that it has surface area = $2\pi r \cdot l$.

Similarly, if we take a cone of slant length l and base radius r :



a simple calculation shows surface area = $\pi r l$

More generally still, if we consider a cone slice:




then its surface area = $2\pi r l$ where $l = \text{slant length}$

and $r = (r_1 + r_2)/2$ is average of radii of the bases.


Cylinders, cones, and cone slices are all examples of surfaces of revolution, and we can use Calculus to obtain an integral formula for Surface area of any surface of revolution ...

Consider a curve $y = f(x)$ from $x=a$ to $x=b$. By rotating this curve around the x -axis, we get a surface of revolution:



So a surface of revolution is just the (lateral) boundary of the corresponding solid of revolution.

As usual, to find the area of a surface of revolution, we break the curve into short intervals where we approximate it by a linear function, giving cone segments:



We explained last class when talking about arc lengths that the slant length of the i^{th} cone segment $= \sqrt{1 + (\frac{dy_i}{dx})^2} \Delta x$. Meanwhile, the circumference $= 2\pi f(x_i^*)$ for some $x_i^* \in [x_{i-1}, x_i]$.

So the area of the i^{th} segment $= 2\pi f(x_i^*) \cdot \sqrt{1 + (\frac{dy_i}{dx})^2} \Delta x$.

and the total area of surface $\approx \sum_{i=1}^n 2\pi f(x_i^*) \sqrt{1 + (\frac{dy_i}{dx})^2} \Delta x$.

Taking the limit as $n \rightarrow \infty$, we get:

$$\begin{aligned} & \text{Area of surface} \\ & \text{of revolution} \\ & \text{rotating } y = f(x) \\ & \text{from } x=a \text{ to } x=b \\ & \text{about } x\text{-axis} \end{aligned} = \boxed{\int_a^b 2\pi f(x) \sqrt{1 + (f'(x))^2} dx}$$

To remember this formula, think:

$$\frac{\text{circumference}}{2\pi f(x)} \cdot \frac{\text{length}}{\sqrt{1 + (\frac{dy}{dx})^2} dx}$$

2/28

E.g.: Consider $y = \sqrt{x}$ from $x=0$ to $x=1$ rotated about x-axis.

Area of surface of revolution = $\int_0^1 2\pi f(x) \sqrt{1+(f'(x))^2} dx$ where $f(x) = x^{1/2}$
 $f'(x) = \frac{1}{2}x^{-1/2}$

$$= \int_0^1 2\pi \sqrt{x} \sqrt{1+(\frac{1}{2}\frac{1}{\sqrt{x}})^2} dx = \int_0^1 2\pi \sqrt{x} \sqrt{1+\frac{1}{4x}} dx$$


$$= 2\pi \int_0^1 \sqrt{x} \cdot (1+\frac{1}{4x}) dx = 2\pi \int_0^1 \sqrt{x+\frac{1}{4x}} dx$$

(1) Indef. integral: $\int \sqrt{x+\frac{1}{4x}} dx = \int \sqrt{u} du = \frac{2}{3} u^{3/2}$

$$u = x + \frac{1}{4x} \quad u = \frac{2}{3}(x + \frac{1}{4x})^{3/2}$$

(2) Plug in FTC: $\Rightarrow 2\pi \int_0^1 \sqrt{x+\frac{1}{4x}} dx = 2\pi \left[\frac{2}{3} \left(x + \frac{1}{4x} \right)^{3/2} \right]_0^1 = \frac{4\pi}{3} \left((\frac{5}{4})^{3/2} - (\frac{1}{4})^{3/2} \right)$

E.g.: Let's compute the surface area of a sphere of radius r .



recall sphere is surface of revolution
for $y = f(x) = \sqrt{r^2 - x^2}$ for $x = -r$ to $x = r$
we compute $f'(x) = 2x \cdot \frac{1}{2} (r^2 - x^2)^{-1/2} = \frac{-x}{\sqrt{r^2 - x^2}}$

So area = $\int_{-r}^r 2\pi f(x) \sqrt{1+(f'(x))^2} dx$

$$= \int_{-r}^r 2\pi \sqrt{r^2 - x^2} \sqrt{1 + \left(\frac{-x}{\sqrt{r^2 - x^2}} \right)^2} dx = \int_{-r}^r 2\pi \sqrt{r^2 - x^2} \sqrt{1 + \frac{x^2}{r^2 - x^2}} dx$$
$$= 2\pi \int_{-r}^r \sqrt{(r^2 - x^2)(1 + \frac{x^2}{r^2 - x^2})} dx = 2\pi \int_{-r}^r \sqrt{(r^2 - x^2) + x^2} dx$$
$$= 2\pi \int_{-r}^r \sqrt{r^2} dx = 2\pi r \int_{-r}^r dx = [4\pi r^2]$$

Note! If we did $\int_a^b 2\pi f(x) \sqrt{1+(f'(x))^2} dx$ here instead


we would get $\int_a^b 2\pi r dx = 2\pi r(b-a)$,

which gives the surface area of sphere segment

from $x=a$ to $x=b$
(result of Archimedes!)

It is also possible to compute surface area by integrating w.r.t. y .

Suppose that $x = g(y)$ for $y=c$ to $y=d$, and we rotate this curve about the x -axis:



↙ Same surface of revolution
but given x in terms of y

A similar computation shows that

$$\text{Area} = \int_c^d 2\pi y \cdot \sqrt{1 + (g'(y))^2} dy$$

Think: circumference \times length
 $\sqrt{1 + (\frac{dx}{dy})^2} dy = \sqrt{1 + (\frac{dy}{dx})^2} dx$

E.g. Consider curve $x = \frac{2}{3} y^{3/2}$ from $y=0$ to $y=3$.

Compute surface area of surface get by rotating about x -axis.

Since we already have x in terms of y , it is easiest here to use the y -integral formula:

$$\text{Area} = \int_c^d 2\pi y \sqrt{1 + (g'(y))^2} dy \quad \text{where } g(y) = \frac{2}{3} y^{3/2} \\ \text{so } g'(y) = y^{1/2}$$

$$= \int_0^3 2\pi y \sqrt{1 + (y^{1/2})^2} dy = \int_0^3 2\pi y \sqrt{1+y} dy$$

① indef. integral: $\int y \sqrt{1+y} dy = \int (u-1) \sqrt{u} du$
 $u = 1+y \Rightarrow y = u-1 \Rightarrow = \int u^{3/2} - u^{1/2} du$
 $du = dy \quad = \frac{2}{5} u^{5/2} - \frac{2}{3} u^{3/2} = \frac{2}{5} (1+y)^{5/2} - \frac{2}{3} (1+y)^{3/2}$

② Plug in FTC: $\Rightarrow 2\pi \int_0^3 y \sqrt{1+y} dy = 2\pi \left[\frac{2}{5} (1+y)^{5/2} - \frac{2}{3} (1+y)^{3/2} \right]_0^3$
 $= 2\pi \left(\left(\frac{2}{5} 4^{5/2} - \frac{2}{3} 4^{3/2} \right) - \left(\frac{2}{5} - \frac{2}{3} \right) \right) = \dots = \frac{232\pi}{15} \cdot //$