


The mystery of plethysm coefficients

Anne Schilling

Department of Mathematics, UC Davis

based on joint work with Rosa Orellana (Dartmouth), Franco Saliola (UQAM), Mike Zabrocki (York), Algebraic Combinatorics (2022), to appear OSZ, Laura Colmenarejo (NCSU) in progress



OPAC
University of Minnesota
May 20, 2022

FPSAC 2023 at UC Davis: July 17-21, 2023



fpsac23.math.ucdavis.edu

Outline

- 1 The plethysm problem
- 2 Diagram algebras
- 3 Uniform block permutation algebra
- 4 Symmetric chain decompositions

Why work on a combinatorial interpretation?

- **Inspiration**/excuse to learn a lot more mathematics

Why work on a combinatorial interpretation?

- **Inspiration**/excuse to learn a lot more mathematics
- Develop a better **understanding of the underlying structure** (representation theory, geometry,)

Why work on a combinatorial interpretation?

- **Inspiration**/excuse to learn a lot more mathematics
- Develop a better **understanding of the underlying structure** (representation theory, geometry,)
- Research is a little like a **random walk**, you bump into a lot of cool stuff on the way, even if you do not return necessarily to the original question.

Representations

G group, V vector space

Representations

G group, V vector space

- **Representation** $\rho: G \rightarrow \text{End}(V)$ homomorphism

Representations

G group, V vector space

- **Representation** $\rho: G \rightarrow \text{End}(V)$ homomorphism
- **Character** $\text{char}(g) = \text{trace} \rho(g)$

Representations

G group, V vector space

- **Representation** $\rho: G \rightarrow \text{End}(V)$ homomorphism
- **Character** $\text{char}(g) = \text{trace} \rho(g)$

Remark

Characters are **class functions**, that is, they are constant on conjugacy classes $\text{char}(hgh^{-1}) = \text{char}(g)$.

Plethysm via representations of GL_n

Definition

$GL_n(\mathbb{C}) =$ invertible $n \times n$ matrices

Plethysm via representations of GL_n

Definition

$GL_n(\mathbb{C}) =$ invertible $n \times n$ matrices

- GL_n -representation $\rho: GL_n \rightarrow GL_m$
- GL_m -representation $\tau: GL_m \rightarrow GL_r$

Plethysm via representations of GL_n

Definition

$GL_n(\mathbb{C}) =$ invertible $n \times n$ matrices

- GL_n -representation $\rho: GL_n \rightarrow GL_m$
- GL_m -representation $\tau: GL_m \rightarrow GL_r$
- Composition is GL_n -representation

$$\tau \circ \rho: GL_n \rightarrow GL_r$$

Plethysm via representations of GL_n

Definition

$GL_n(\mathbb{C}) =$ invertible $n \times n$ matrices

- GL_n -representation $\rho: GL_n \rightarrow GL_m$
- GL_m -representation $\tau: GL_m \rightarrow GL_r$
- Composition is GL_n -representation

$$\tau \circ \rho: GL_n \rightarrow GL_r$$

Definition

Character of composition is **plethysm**:

$$\text{char}(\tau \circ \rho) = \text{char}(\tau)[\text{char}(\rho)]$$

Frobenius map

R^n space of class functions of GL_n

Λ^n ring of symmetric functions of degree n

Frobenius map

R^n space of class functions of GL_n

Λ^n ring of symmetric functions of degree n

Power sum symmetric function p_λ

$$p_\lambda = p_{\lambda_1} p_{\lambda_2} \cdots p_{\lambda_\ell}$$

$$p_r = x_1^r + x_2^r + \cdots$$

Frobenius map

R^n space of class functions of GL_n

Λ^n ring of symmetric functions of degree n

Power sum symmetric function p_λ

$$p_\lambda = p_{\lambda_1} p_{\lambda_2} \cdots p_{\lambda_\ell}$$

$$p_r = x_1^r + x_2^r + \cdots$$

Schur function s_λ

$$s_\lambda = \sum_{T \in \text{SSYT}(\lambda)} x^{\text{wt}(T)}$$

Frobenius map – continued

Definition

The **Frobenius characteristic map** is $\text{ch}^n: R^n \rightarrow \Lambda^n$

$$\text{ch}^n(\chi) = \sum_{\mu \vdash n} \frac{1}{z_\mu} \chi_\mu p_\mu$$

where $z_\mu = 1^{a_1} a_1! 2^{a_2} a_2! \dots$ for $\mu = 1^{a_1} 2^{a_2} \dots$

Frobenius map – continued

Definition

The **Frobenius characteristic map** is $\text{ch}^n: R^n \rightarrow \Lambda^n$

$$\text{ch}^n(\chi) = \sum_{\mu \vdash n} \frac{1}{z_\mu} \chi_\mu p_\mu$$

where $z_\mu = 1^{a_1} a_1! 2^{a_2} a_2! \dots$ for $\mu = 1^{a_1} 2^{a_2} \dots$

Remark

The **irreducible character** χ^λ indexed by λ under the Frobenius map is

$$\text{ch}^n(\chi^\lambda) = s_\lambda$$

by the identity

$$s_\lambda = \sum_{\mu} \frac{1}{z_\mu} \chi_\mu^\lambda p_\mu$$

Plethysm for symmetric functions

Definition

$f, g \in \Lambda$ symmetric functions

Monomial expansion $f = \sum_{i \geq 1} x^{a_i}$

Plethysm for symmetric functions

Definition

$f, g \in \Lambda$ symmetric functions

Monomial expansion $f = \sum_{i \geq 1} x^{a^i}$

The **plethysm** $g[f]$ is defined by

$$g[f] = g(x^{a^1}, x^{a^2}, \dots)$$

Plethysm for symmetric functions

Definition

$f, g \in \Lambda$ symmetric functions

Monomial expansion $f = \sum_{i \geq 1} x^{a^i}$

The **plethysm** $g[f]$ is defined by

$$g[f] = g(x^{a^1}, x^{a^2}, \dots)$$

Remark

Plethysm: Greek for multiplication

Plethysm for symmetric functions

Definition

$f, g \in \Lambda$ symmetric functions

Monomial expansion $f = \sum_{i \geq 1} x^{a_i}$

The **plethysm** $g[f]$ is defined by

$$g[f] = g(x^{a^1}, x^{a^2}, \dots)$$

Remark

Plethysm: Greek for multiplication

Example

$$s_1 = x_1 + x_2 + \dots \quad \Rightarrow \quad g[s_1] = g(x_1, x_2, \dots) = g$$

Plethysm for symmetric functions

Definition

$f, g \in \Lambda$ symmetric functions

Monomial expansion $f = \sum_{i \geq 1} x^{a^i}$

The **plethysm** $g[f]$ is defined by

$$g[f] = g(x^{a^1}, x^{a^2}, \dots)$$

Remark

Plethysm: Greek for multiplication

Example

$$s_1 = x_1 + x_2 + \dots \quad \Rightarrow \quad g[s_1] = g(x_1, x_2, \dots) = g$$

$$p_n = x_1^n + x_2^n + \dots \quad \Rightarrow \quad f[p_n] = f(x_1^n, x_2^n, \dots) = \sum_{i \geq 1} x^{a^i n} = p_n[f]$$

Plethysm for symmetric functions – example

Example

$$s_2[x_1, x_2] = x_1^2 + x_1x_2 + x_2^2$$

11

12

22

Plethysm for symmetric functions – example

Example

$$s_2[x_1, x_2] = x_1^2 + x_1 x_2 + x_2^2$$

$$\boxed{1\,1} \quad \boxed{1\,2} \quad \boxed{2\,2}$$

Plethysm

$$s_2[s_2[x_1, x_2]] = s_2[x_1^2, x_1 x_2, x_2^2]$$

$$= x_1^4 + x_1^3 x_2 + x_1^2 x_2^2 + x_1^2 x_2^2 + x_1 x_2^3 + x_2^4$$

$$\boxed{1\,1} \quad \boxed{1\,2} \quad \boxed{1\,3} \quad \boxed{2\,2} \quad \boxed{2\,3} \quad \boxed{3\,3}$$

$$\boxed{1111} \quad \boxed{1112} \quad \boxed{1122} \quad \boxed{1212} \quad \boxed{1222} \quad \boxed{2222}$$

$$= s_4[x_1, x_2] + s_{2,2}[x_1, x_2]$$

Plethysm problem

Problem

Find a *combinatorial interpretation* for the coefficients $a_{\lambda\mu}^\nu \in \mathbb{N}$ in the expansion

$$s_\lambda[s_\mu] = \sum_{\nu} a_{\lambda\mu}^\nu s_\nu$$

Plethysm problem


Problem

Find a *combinatorial interpretation* for the coefficients $a_{\lambda\mu}^\nu \in \mathbb{N}$ in the expansion

$$s_\lambda[s_\mu] = \sum_{\nu} a_{\lambda\mu}^\nu s_\nu$$


Problem

Find a *crystal on tableaux of tableaux* which explains $a_{\lambda\mu}^\nu$.



Plethysm problem – special cases

Partition λ is **even** if all columns have even length




Plethysm problem – special cases

Partition λ is **even** if all columns have even length



Partition λ is **threshold** if $\lambda'_i = \lambda_i + 1$ for all $1 \leq i \leq d(\lambda)$




Plethysm problem – special cases

Partition λ is **even** if all columns have even length



Partition λ is **threshold** if $\lambda'_i = \lambda_i + 1$ for all $1 \leq i \leq d(\lambda)$



Theorem

We have

$$s_h[s_2] = \sum_{\substack{\lambda \vdash 2h \\ \lambda \text{ even}}} s_{\lambda'}$$


$$s_1 s_h[s_2] = \sum_{\substack{\lambda \vdash 2h \\ \lambda \text{ threshold}}} s_{\lambda'}$$

$$s_h[s_1^2] = \sum_{\substack{\lambda \vdash 2h \\ \lambda \text{ even}}} s_{\lambda}$$


$$s_1 s_h[s_1^2] = \sum_{\substack{\lambda \vdash 2h \\ \lambda \text{ threshold}}} s_{\lambda}$$

Plethysm problem – special cases

Partition λ is **even** if all columns have even length



Partition λ is **threshold** if $\lambda'_i = \lambda_i + 1$ for all $1 \leq i \leq d(\lambda)$



Theorem

We have

$$s_h[s_2] = \sum_{\substack{\lambda \vdash 2h \\ \lambda \text{ even}}} s_{\lambda'}$$

$$s_{1^h}[s_2] = \sum_{\substack{\lambda \vdash 2h \\ \lambda \text{ threshold}}} s_{\lambda'}$$

$$s_h[s_{1^2}] = \sum_{\substack{\lambda \vdash 2h \\ \lambda \text{ even}}} s_{\lambda}$$

$$s_{1^h}[s_{1^2}] = \sum_{\substack{\lambda \vdash 2h \\ \lambda \text{ threshold}}} s_{\lambda}$$

Appeared in **Littlewood 1950**, **Macdonald 1998** (pg 138)

Littlewood and Macdonald



Easy proof – s -perp trick

Action of s_λ^\perp on $f \in \Lambda$

$$s_\lambda^\perp f = \sum_{\mu} \langle f, s_\lambda s_\mu \rangle s_\mu$$

Easy proof – s -perp trick

Action of s_λ^\perp on $f \in \Lambda$

$$s_\lambda^\perp f = \sum_{\mu} \langle f, s_\lambda s_\mu \rangle s_\mu$$

Proposition (The s -perp trick)

Let f and g be two symmetric functions of homogeneous degree d . If

$$s_r^\perp f = s_r^\perp g \quad \text{for all } 1 \leq r \leq d,$$

then $f = g$. Same statement is true if s_r^\perp is replaced by $s_{1^r}^\perp$.

Easy proof – s -perp trick

Action of s_λ^\perp on $f \in \Lambda$

$$s_\lambda^\perp f = \sum_{\mu} \langle f, s_\lambda s_\mu \rangle s_\mu$$

Proposition (The s -perp trick)

Let f and g be two symmetric functions of homogeneous degree d . If

$$s_r^\perp f = s_r^\perp g \quad \text{for all } 1 \leq r \leq d,$$

then $f = g$. Same statement is true if s_r^\perp is replaced by $s_{1^r}^\perp$.

The following hold:

$$\begin{aligned} s_r^\perp s_{1^h}[s_{1^w}] &= s_{1^{h-r}}[s_{1^w}] s_{1^r}[s_{1^{w-1}}] & s_{1^r}^\perp s_h[s_w] &= s_{h-r}[s_w] s_{1^r}[s_{w-1}] \\ s_r^\perp s_h[s_{1^w}] &= s_{h-r}[s_{1^w}] s_r[s_{1^{w-1}}] & s_{1^r}^\perp s_{1^h}[s_w] &= s_{1^{h-r}}[s_w] s_r[s_{w-1}] \end{aligned}$$

Easy proof – s -perp trick

Action of s_λ^\perp on $f \in \Lambda$

$$s_\lambda^\perp f = \sum_{\mu} \langle f, s_\lambda s_\mu \rangle s_\mu$$

Proposition (The s -perp trick)

Let f and g be two symmetric functions of homogeneous degree d . If

$$s_r^\perp f = s_r^\perp g \quad \text{for all } 1 \leq r \leq d,$$

then $f = g$. Same statement is true if s_r^\perp is replaced by $s_{1^r}^\perp$.

Remark

Benefit: Fast computational algorithm to compute plethysm of Schur functions!

Relationship between restriction problem and plethysm

Restriction: λ partition with at most n parts

$$\text{Res}_{S_n}^{GL_n} V_{GL_n}^\lambda = \bigoplus (V_{S_n}^\mu)^{r_{\lambda\mu}}$$

Relationship between restriction problem and plethysm

Restriction: λ partition with at most n parts

$$\operatorname{Res}_{S_n}^{GL_n} V_{GL_n}^\lambda = \bigoplus (V_{S_n}^\mu)^{r_{\lambda\mu}}$$

$r_{\lambda\mu}$ = coefficient of s_μ in the plethysm $s_{(n-|\lambda|,\lambda)}[s_{(1)} + s_{(2)} + \cdots]$

Outline

- 1 The plethysm problem
- 2 Diagram algebras
- 3 Uniform block permutation algebra
- 4 Symmetric chain decompositions

Diagram algebras

- **Restrict** diagonal action of GL_n on $V^{\otimes k}$ to $S_n \subseteq GL_n$: for $\sigma \in S_n$

$$\sigma(v_{i_1} \otimes v_{i_2} \otimes \cdots \otimes v_{i_k}) = \sigma v_{i_1} \otimes \cdots \otimes \sigma v_{i_k}$$

Diagram algebras

- **Restrict** diagonal action of GL_n on $V^{\otimes k}$ to $S_n \subseteq GL_n$: for $\sigma \in S_n$

$$\sigma(v_{i_1} \otimes v_{i_2} \otimes \cdots \otimes v_{i_k}) = \sigma v_{i_1} \otimes \cdots \otimes \sigma v_{i_k}$$

- What **commutes** with this action?

Diagram algebras

- **Restrict** diagonal action of GL_n on $V^{\otimes k}$ to $S_n \subseteq GL_n$: for $\sigma \in S_n$

$$\sigma(v_{i_1} \otimes v_{i_2} \otimes \cdots \otimes v_{i_k}) = \sigma v_{i_1} \otimes \cdots \otimes \sigma v_{i_k}$$

- What **commutes** with this action?

Answer: **Partition algebra** $P_k(n)$

Martin, Jones 1990s

Diagram algebras

- **Restrict** diagonal action of GL_n on $V^{\otimes k}$ to $S_n \subseteq GL_n$: for $\sigma \in S_n$

$$\sigma(v_{i_1} \otimes v_{i_2} \otimes \cdots \otimes v_{i_k}) = \sigma v_{i_1} \otimes \cdots \otimes \sigma v_{i_k}$$

- What **commutes** with this action?

Answer: **Partition algebra** $P_k(n)$ Martin, Jones 1990s

- **Basis:** set partitions of $\{1, 2, \dots, k\} \cup \{\bar{1}, \bar{2}, \dots, \bar{k}\}$

Diagram algebras

- **Restrict** diagonal action of GL_n on $V^{\otimes k}$ to $S_n \subseteq GL_n$: for $\sigma \in S_n$

$$\sigma(v_{i_1} \otimes v_{i_2} \otimes \cdots \otimes v_{i_k}) = \sigma v_{i_1} \otimes \cdots \otimes \sigma v_{i_k}$$


- What **commutes** with this action?

Answer: **Partition algebra** $P_k(n)$ Martin, Jones 1990s

- **Basis:** set partitions of $\{1, 2, \dots, k\} \cup \{\bar{1}, \bar{2}, \dots, \bar{k}\}$

Example

The set partition $\pi = \{\{1, 2, 4, \bar{2}, \bar{5}\}, \{3\}, \{5, 6, 7, \bar{3}, \bar{4}, \bar{6}, \bar{7}\}, \{8, \bar{8}\}, \{\bar{1}\}\}$ is represented by the following diagram:



Martin and Jones



Centralizer pair

$V_{P_k(n)}^{(n-|\lambda|, \lambda)} =$ **simple module** indexed by partitions λ such that
 $\lambda_1 + \lambda_2 + \cdots \leq k$

Centralizer pair

$V_{P_k(n)}^{(n-|\lambda|, \lambda)} = \text{simple module indexed by partitions } \lambda \text{ such that}$
 $\lambda_1 + \lambda_2 + \cdots \leq k$

Example

$$V_{P_3(6)}^{(4,2)} = \text{span} \left\{ \begin{array}{|c|c|c|c|} \hline & & & 3 \\ \hline 1 & 2 & & \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline & & & 2 \\ \hline 1 & 3 & & \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline & & & 1 \\ \hline 2 & 3 & & \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline & & & \\ \hline 1 & 2 & 3 & \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline & & & \\ \hline 1 & 2 & 3 & \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline & & & \\ \hline 2 & 1 & 3 & \\ \hline \end{array} \right\}$$

Centralizer pair

$$V_{P_k(n)}^{(n-|\lambda|, \lambda)} = \text{simple module indexed by partitions } \lambda \text{ such that } \lambda_1 + \lambda_2 + \cdots \leq k$$

Example

$$V_{P_3(6)}^{(4,2)} = \text{span} \left\{ \begin{array}{|c|c|c|c|} \hline & & & 3 \\ \hline 1 & 2 & & \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline & & & 2 \\ \hline 1 & 3 & & \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline & & & 1 \\ \hline 2 & 3 & & \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline & & & \\ \hline 1 & 2 & 3 & \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline & & & \\ \hline 1 & 2 & 3 & \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline & & & \\ \hline 2 & 1 & 3 & \\ \hline \end{array} \right\}$$

Dimension is number of **set valued tableaux**

Centralizer pair

$$V_{P_k(n)}^{(n-|\lambda|,\lambda)} = \text{simple module indexed by partitions } \lambda \text{ such that } \lambda_1 + \lambda_2 + \cdots \leq k$$

Example

$$V_{P_3(6)}^{(4,2)} = \text{span} \left\{ \begin{array}{|c|c|c|c|} \hline & & & 3 \\ \hline 1 & 2 & & \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline & & & 2 \\ \hline 1 & 3 & & \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline & & & 1 \\ \hline 2 & 3 & & \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline & & & \\ \hline 12 & 3 & & \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline & & & \\ \hline 1 & 23 & & \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline & & & \\ \hline 2 & 13 & & \\ \hline \end{array} \right\}$$

Dimension is number of **set valued tableaux**

Theorem (Jones 1994)

$$V^{\otimes k} \cong \bigoplus_{\lambda, \lambda_1 + \lambda_2 + \cdots \leq k} V_{P_k(n)}^{(n-|\lambda|,\lambda)} \otimes V_{S_n}^{(n-|\lambda|,\lambda)}$$

Centralizer pair

$V_{P_k(n)}^{(n-|\lambda|, \lambda)}$ = **simple module** indexed by partitions λ such that
 $\lambda_1 + \lambda_2 + \cdots \leq k$

Example

$$V_{P_3(6)}^{(4,2)} = \text{span} \left\{ \begin{array}{|c|c|c|c|} \hline & & & 3 \\ \hline 1 & 2 & & \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline & & & 2 \\ \hline 1 & 3 & & \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline & & & 1 \\ \hline 2 & 3 & & \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline & & & \\ \hline 1 & 2 & 3 & \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline & & & \\ \hline 1 & 2 & 3 & \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline & & & \\ \hline 2 & 1 & 3 & \\ \hline \end{array} \right\}$$

Dimension is number of **set valued tableaux**


Theorem (Jones 1994)

$$V^{\otimes k} \cong \bigoplus_{\lambda, \lambda_1 + \lambda_2 + \cdots \leq k} V_{P_k(n)}^{(n-|\lambda|, \lambda)} \otimes V_{S_n}^{(n-|\lambda|, \lambda)}$$

Remark

- S_k and GL_n form a **centralizer pair**
- $P_k(n)$ and S_n form a **centralizer pair**

See-Saw pairs



(See book by **Goodman, Wallach**)

See-Saw pairs


$A \hookrightarrow B$ algebra embedding

$$\operatorname{Res}_A^B V_B^\lambda = \bigoplus_{\mu} (V_A^\mu)^{\oplus c_{\lambda\mu}}$$

See-Saw pairs

$A \hookrightarrow B$ algebra embedding

$$\operatorname{Res}_A^B V_B^\lambda = \bigoplus_{\mu} (V_A^\mu)^{\oplus c_{\lambda\mu}}$$




- B and C centralizer pair
- A and D centralizer pair

See-Saw pairs

$A \hookrightarrow B$ algebra embedding

$$\text{Res}_A^B V_B^\lambda = \bigoplus_{\mu} (V_A^\mu)^{\oplus c_{\lambda\mu}}$$




- B and C centralizer pair
- A and D centralizer pair

- 1 Indices for the simple modules for B and C are the same.
- 2 Indices for the simple modules for A and D are the same.

See-Saw pairs

$A \hookrightarrow B$ algebra embedding

$$\text{Res}_A^B V_B^\lambda = \bigoplus_{\mu} (V_A^\mu)^{\oplus c_{\lambda\mu}}$$




- B and C centralizer pair
- A and D centralizer pair


- 1 Indices for the simple modules for B and C are the same.
- 2 Indices for the simple modules for A and D are the same.

$$\text{Res}_C^D V_D^\mu = \bigoplus_{\lambda} (V_C^\lambda)^{\oplus c_{\lambda\mu}}$$

Our See-Saw pair




Our See-Saw pair



$$\text{Res}_{S_n}^{GL_n} V_{GL_n}^\lambda = \bigoplus_{\mu} (V_{S_n}^\mu)^{\oplus r_{\lambda\mu}}$$

$$\text{Res}_{S_k}^{P_k(n)} V_{P_k(n)}^\mu = \bigoplus_{\lambda} (V_{S_k}^\lambda)^{\oplus r_{\lambda\mu}}$$

Our See-Saw pair



$$\text{Res}_{S_n}^{GL_n} V_{GL_n}^\lambda = \bigoplus_{\mu} (V_{S_n}^\mu)^{\oplus r_{\lambda\mu}}$$

$$\text{Res}_{S_k}^{P_k(n)} V_{P_k(n)}^\mu = \bigoplus_{\lambda} (V_{S_k}^\lambda)^{\oplus r_{\lambda\mu}}$$

Idea: Restrict representations of $P_k(n)$ to S_k

The approach

\mathcal{U}_k uniform block permutation algebra

$$\underbrace{S_k \hookrightarrow}_{\text{special cases of plethysm}} \mathcal{U}_k \hookrightarrow \underbrace{P_k(n)}_{\text{generalized LR coefficients}}$$

The approach

\mathcal{U}_k uniform block permutation algebra

$$\underbrace{S_k \hookrightarrow}_{\text{special cases of plethysm}} \mathcal{U}_k \hookrightarrow \underbrace{P_k(n)}_{\text{generalized LR coefficients}}$$

Goal: Combinatorial model for the representation theory of \mathcal{U}_k

Outline

- 1 The plethysm problem
- 2 Diagram algebras
- 3 Uniform block permutation algebra**
- 4 Symmetric chain decompositions

Uniform block permutations

Tanabe, Kosuda

Party algebra, centralizer algebra for complex reflection groups

Uniform block permutations

Tanabe, Kosuda

Party algebra, centralizer algebra for complex reflection groups

Definition

The set partition $d = \{d_1, d_2, \dots, d_\ell\}$ of $[k] \cup [\bar{k}]$ is **uniform** if $|d_i \cap [k]| = |d_i \cap [\bar{k}]|$ for all $1 \leq i \leq \ell$. Let

$$\mathcal{U}_k = \{d \vdash [k] \cup [\bar{k}] : d \text{ uniform}\}.$$

Uniform block permutations

Tanabe, Kosuda

Party algebra, centralizer algebra for complex reflection groups

Definition

The set partition $d = \{d_1, d_2, \dots, d_\ell\}$ of $[k] \cup [\bar{k}]$ is **uniform** if $|d_i \cap [k]| = |d_i \cap [\bar{k}]|$ for all $1 \leq i \leq \ell$. Let

$$\mathcal{U}_k = \{d \vdash [k] \cup [\bar{k}] : d \text{ uniform}\}.$$

Example

$$d = \{\{2, \bar{4}\}, \{5, \bar{7}\}, \{1, 3, \bar{1}, \bar{2}\}, \{4, 6, \bar{3}, \bar{6}\}, \{7, 8, 9, \bar{5}, \bar{8}, \bar{9}\}\}$$

Uniform block permutations

Tanabe, Kosuda

Party algebra, centralizer algebra for complex reflection groups

Definition

The set partition $d = \{d_1, d_2, \dots, d_\ell\}$ of $[k] \cup [\bar{k}]$ is **uniform** if $|d_i \cap [k]| = |d_i \cap [\bar{k}]|$ for all $1 \leq i \leq \ell$. Let

$$\mathcal{U}_k = \{d \vdash [k] \cup [\bar{k}] : d \text{ uniform}\}.$$

Example

$$d = \{\{2, \bar{4}\}, \{5, \bar{7}\}, \{1, 3, \bar{1}, \bar{2}\}, \{4, 6, \bar{3}, \bar{6}\}, \{7, 8, 9, \bar{5}, \bar{8}, \bar{9}\}\}$$

Think of d as a **size-preserving bijection**

$$\left(\begin{array}{ccccc} \{2\} & \{5\} & \{1, 3\} & \{4, 6\} & \{7, 8, 9\} \\ \{4\} & \{7\} & \{1, 2\} & \{3, 6\} & \{5, 8, 9\} \end{array} \right)$$

Uniform block permutations

Tanabe, Kosuda

Party algebra, centralizer algebra for complex reflection groups

Definition

The set partition $d = \{d_1, d_2, \dots, d_\ell\}$ of $[k] \cup [\bar{k}]$ is **uniform** if $|d_i \cap [k]| = |d_i \cap [\bar{k}]|$ for all $1 \leq i \leq \ell$. Let

$$\mathcal{U}_k = \{d \vdash [k] \cup [\bar{k}] : d \text{ uniform}\}.$$

Example

$$d = \{\{2, \bar{4}\}, \{5, \bar{7}\}, \{1, 3, \bar{1}, \bar{2}\}, \{4, 6, \bar{3}, \bar{6}\}, \{7, 8, 9, \bar{5}, \bar{8}, \bar{9}\}\}$$

Think of d as a **size-preserving bijection**


$$\left(\begin{array}{ccccc} \{2\} & \{5\} & \{1, 3\} & \{4, 6\} & \{7, 8, 9\} \\ \{4\} & \{7\} & \{1, 2\} & \{3, 6\} & \{5, 8, 9\} \end{array} \right)$$

\Rightarrow Elements of \mathcal{U}_k are called **uniform block permutations**

Uniform block permutations – continued

Example


Diagram for $\{\{1, 3, \bar{1}, \bar{2}\}, \{2, \bar{4}\}, \{4, 6, \bar{3}, \bar{6}\}, \{5, \bar{7}\}, \{7, 8, 9, \bar{5}, \bar{8}, \bar{9}\}\}$




Uniform block permutations – continued

Example


Diagram for $\{\{1, 3, \bar{1}, \bar{2}\}, \{2, \bar{4}\}, \{4, 6, \bar{3}, \bar{6}\}, \{5, \bar{7}\}, \{7, 8, 9, \bar{5}, \bar{8}, \bar{9}\}\}$




The product of




and



is obtained by stacking the diagrams of d and d' :



$=$



Idempotents

For every set partition π of $[k]$ we define:

$$e_\pi = \{A \cup \bar{A} : A \in \pi\} \in \mathcal{U}_k$$

where $\bar{A} = \{\bar{i} : i \in A\}$.

Idempotents

For every set partition π of $[k]$ we define:

$$e_\pi = \{A \cup \bar{A} : A \in \pi\} \in \mathcal{U}_k$$

where $\bar{A} = \{\bar{i} : i \in A\}$. For example,

$$e_{2|7|14|36|589} =$$

Idempotents

For every set partition π of $[k]$ we define:

$$e_\pi = \{A \cup \bar{A} : A \in \pi\} \in \mathcal{U}_k$$


where $\bar{A} = \{\bar{i} : i \in A\}$. For example,

$$e_{2|7|14|36|589} =$$


Lemma

The set $E(\mathcal{U}_k) = \{e_\pi : \pi \vdash [k]\}$ is a *complete set of idempotents* in \mathcal{U}_k .

Factorizable monoid



Factorizable monoid



Proposition



For every $d \in \mathcal{U}_k$ and every $\sigma \in S_k$ satisfying $\sigma(B \cap [k]) = \overline{B} \cap [k]$, we have

$$d = e_{\text{top}(d)} \sigma = \sigma e_{\text{bot}(d)}.$$

Consequently, \mathcal{U}_k is a **factorizable monoid**

$$\mathcal{U}_k = E(\mathcal{U}_k) S_k = S_k E(\mathcal{U}_k).$$

Factorizable monoid



(See book by **Steinberg** 2016)

Maximal subgroups

Definition

M finite monoid, e idempotent

Maximal subgroup: $G_e =$ unique largest subgroup of M containing e

Maximal subgroups

Definition

M finite monoid, e idempotent

Maximal subgroup: $G_e =$ unique largest subgroup of M containing e

Lemma

The maximal subgroup of \mathcal{U}_k at the idempotent e_π is

$$G_{e_\pi} = \{d \in \mathcal{U}_k : \text{top}(d) = \text{bot}(d) = \pi\}$$

Maximal subgroups

Definition

M finite monoid, e idempotent

Maximal subgroup: $G_e =$ unique largest subgroup of M containing e

Lemma

The maximal subgroup of \mathcal{U}_k at the idempotent e_π is

$$G_{e_\pi} = \{d \in \mathcal{U}_k : \text{top}(d) = \text{bot}(d) = \pi\}$$

Example

For $\pi = \{\{1\}, \{2\}, \{3, 4\}, \{5, 6\}\}$

$$G_{e_\pi} = \left\{ \begin{array}{c} \text{Diagram 1} \end{array}, \begin{array}{c} \text{Diagram 2} \end{array}, \begin{array}{c} \text{Diagram 3} \end{array}, \begin{array}{c} \text{Diagram 4} \end{array} \right\}$$

Maximal subgroups – continued

Example

For $\pi = \{\{1\}, \{2\}, \{3, 4\}, \{5, 6\}\}$ with $\text{type}(\pi) = (1^2 2^2)$

$$G_{e_\pi} = \left\{ \begin{array}{c} \text{Diagram 1: } \begin{array}{cc} \text{Two vertical lines} \end{array}, \quad \begin{array}{c} \text{Diagram 2: } \begin{array}{cc} \text{Two vertical lines} \end{array}, \quad \begin{array}{c} \text{Diagram 3: } \begin{array}{cc} \text{Two vertical lines} \end{array}, \quad \begin{array}{c} \text{Diagram 4: } \begin{array}{cc} \text{Two vertical lines} \end{array}, \end{array} \right.$$

Theorem

For $\pi \vdash [k]$ with $\text{type}(\pi) = (1^{a_1} 2^{a_2} \dots k^{a_k})$

$$G_{e_\pi} \simeq S_{a_1} \times S_{a_2} \times \dots \times S_{a_k}$$

Representation theory of \mathcal{U}_k

Indexing set of simple modules

$$I_k = \left\{ \left(\lambda^{(1)}, \lambda^{(2)}, \dots, \lambda^{(k)} \right) : \lambda^{(i)} \text{ are partitions such that } \sum_{i=1}^k i|\lambda^{(i)}| = k \right\}$$

Representation theory of \mathcal{U}_k

Indexing set of simple modules

$$I_k = \left\{ \left(\lambda^{(1)}, \lambda^{(2)}, \dots, \lambda^{(k)} \right) : \lambda^{(i)} \text{ are partitions such that } \sum_{i=1}^k i|\lambda^{(i)}| = k \right\}$$

Example

$$I_3 = \{((3), \emptyset, \emptyset), ((2, 1), \emptyset, \emptyset), ((1, 1, 1), \emptyset, \emptyset), ((1), (1), \emptyset), (\emptyset, \emptyset, (1))\}$$

Representation theory of \mathcal{U}_k – continued

Definition

A **uniform tableau** $\mathbf{S} = (S^{(1)}, \dots, S^{(k)})$ of shape $\vec{\lambda} \in I_k$ satisfies:

- 1 $S^{(i)}$ is a tableau of shape $\lambda^{(i)}$ filled with subsets of $[k]$ of size i ;
- 2 $S^{(i)}$ is standard;
- 3 the subsets appearing in \mathbf{S} form a set partition of $[k]$.

We define $\mathcal{T}_{\vec{\lambda}}$ to be the set of uniform tableaux of shape $\vec{\lambda}$.

Representation theory of \mathcal{U}_k – continued

Definition

A **uniform tableau** $\mathbf{S} = (S^{(1)}, \dots, S^{(k)})$ of shape $\vec{\lambda} \in I_k$ satisfies:

- 1 $S^{(i)}$ is a tableau of shape $\lambda^{(i)}$ filled with subsets of $[k]$ of size i ;
- 2 $S^{(i)}$ is standard;
- 3 the subsets appearing in \mathbf{S} form a set partition of $[k]$.

We define $\mathcal{T}_{\vec{\lambda}}$ to be the set of uniform tableaux of shape $\vec{\lambda}$.

Irreducible representations: $V_{\mathcal{U}_k}^{\vec{\lambda}} = \text{span}\{\mathbf{S} \in \mathcal{T}_{\vec{\lambda}}\}$

Representation theory of \mathcal{U}_k – continued

Definition

A **uniform tableau** $\mathbf{S} = (S^{(1)}, \dots, S^{(k)})$ of shape $\vec{\lambda} \in I_k$ satisfies:

- ① $S^{(i)}$ is a tableau of shape $\lambda^{(i)}$ filled with subsets of $[k]$ of size i ;
- ② $S^{(i)}$ is standard;
- ③ the subsets appearing in \mathbf{S} form a set partition of $[k]$.

We define $\mathcal{T}_{\vec{\lambda}}$ to be the set of uniform tableaux of shape $\vec{\lambda}$.

Irreducible representations: $V_{\mathcal{U}_k}^{\vec{\lambda}} = \text{span}\{\mathbf{S} \in \mathcal{T}_{\vec{\lambda}}\}$

Example

$$V_{\mathcal{U}_3}^{((1),(1),\emptyset)} = \text{span}\left\{\left(\boxed{1}, \boxed{23}\right), \left(\boxed{2}, \boxed{13}\right), \left(\boxed{3}, \boxed{12}\right)\right\}$$

Characters of \mathcal{U}_k

Definition

M be a finite monoid.

- Subsemigroup of M generated by $m \in M$ contains a unique idempotent m^ω
- $m, n \in M$ are **conjugate** if there exist $x, x' \in M$ such that $xx'x = x$, $x'xx' = x'$, $x'x = m^\omega$, $xx' = n^\omega$ and $xm^{\omega+1}x' = n^{\omega+1}$

Characters of \mathcal{U}_k

Definition

M be a finite monoid.

- Subsemigroup of M generated by $m \in M$ contains a unique idempotent m^ω
- $m, n \in M$ are **conjugate** if there exist $x, x' \in M$ such that $xx'x = x$, $x'xx' = x'$, $x'x = m^\omega$, $xx' = n^\omega$ and $xm^{\omega+1}x' = n^{\omega+1}$

Definition

- $d \in G_{e_\pi}$: **cycletype**(d) = $(\mu^{(1)}, \mu^{(2)}, \dots, \mu^{(k)})$
where $\mu^{(i)}$ is the cycle type of the permutation $d^{(i)}$
- $x \in \mathcal{U}_k$: **cycletype**(x) = **cycletype**($x^{\omega+1}$)

Characters of \mathcal{U}_k

Definition

M be a finite monoid.

- Subsemigroup of M generated by $m \in M$ contains a unique idempotent m^ω
- $m, n \in M$ are **conjugate** if there exist $x, x' \in M$ such that $xx'x = x$, $x'xx' = x'$, $x'x = m^\omega$, $xx' = n^\omega$ and $xm^{\omega+1}x' = n^{\omega+1}$

Definition

- $d \in G_{e_\pi}$: **cycletype**(d) = $(\mu^{(1)}, \mu^{(2)}, \dots, \mu^{(k)})$
where $\mu^{(i)}$ is the cycle type of the permutation $d^{(i)}$
- $x \in \mathcal{U}_k$: **cycletype**(x) = **cycletype**($x^{\omega+1}$)

$d_{\vec{\mu}}$ representative for generalized conjugacy class of cycle type $\vec{\mu}$

Characters of \mathcal{U}_k – continued

Theorem (OSSZ 2022)

$\vec{\lambda}, \vec{\mu} \in I_k$, $a_i = |\lambda^{(i)}|$, $\lambda = (1^{a_1} 2^{a_2} \dots k^{a_k})$

$$\chi_{\mathcal{U}_k}^{\vec{\lambda}}(d_{\vec{\mu}}) = \sum_{\substack{\vec{\nu} \in I_k \\ |\nu^{(i)}| = a_i}} b_{\vec{\mu}}^{\vec{\nu}} \chi_{G_\lambda}^{\vec{\lambda}}(d_{\vec{\nu}})$$

Characters of \mathcal{U}_k – continued

Theorem (OSSZ 2022)

$\vec{\lambda}, \vec{\mu} \in I_k$, $a_i = |\lambda^{(i)}|$, $\lambda = (1^{a_1} 2^{a_2} \dots k^{a_k})$

$$\chi_{\mathcal{U}_k}^{\vec{\lambda}}(d_{\vec{\mu}}) = \sum_{\substack{\vec{\nu} \in I_k \\ |\nu^{(i)}| = a_i}} b_{\vec{\mu}}^{\vec{\nu}} \chi_{G_\lambda}^{\vec{\lambda}}(d_{\vec{\nu}})$$

Example

Let $\vec{\lambda} = (\emptyset, (1, 1), \emptyset, \emptyset)$, so that $\lambda = (2, 2)$:

$$\chi_{\mathcal{U}_4}^{\vec{\lambda}} \left(\begin{array}{cc} \circ & \circ \\ \diagdown & \diagup \\ \circ & \circ \end{array} \quad \begin{array}{cc} \circ & \circ \\ \diagup & \diagdown \\ \circ & \circ \end{array} \right) = \chi_{G_\lambda}^{\vec{\lambda}} \left(\begin{array}{cc} \circ & \circ \\ \text{---} & \text{---} \\ \circ & \circ \end{array} \quad \begin{array}{cc} \circ & \circ \\ \text{---} & \text{---} \\ \circ & \circ \end{array} \right) + 2 \chi_{G_\lambda}^{\vec{\lambda}} \left(\begin{array}{cccc} \circ & \circ & \circ & \circ \\ \diagdown & \diagup & \diagdown & \diagup \\ \circ & \circ & \circ & \circ \end{array} \right) = -1$$

Coefficients in characters

$$z_\lambda = 1^{a_1} a_1! 2^{a_2} a_2! \cdots k^{a_k} a_k! \quad \text{for } \lambda = (1^{a_1} 2^{a_2} \cdots k^{a_k})$$

$$z_{\vec{\lambda}} = z_{\lambda(1)} z_{\lambda(2)} \cdots z_{\lambda(k)}$$

Coefficients in characters

$$z_\lambda = 1^{a_1} a_1! 2^{a_2} a_2! \cdots k^{a_k} a_k! \quad \text{for } \lambda = (1^{a_1} 2^{a_2} \cdots k^{a_k})$$

$$\mathbf{z}_{\vec{\lambda}} = z_{\lambda(1)} z_{\lambda(2)} \cdots z_{\lambda(k)}$$

Theorem (OSSZ 2022)

$$\vec{\mu}, \vec{\nu} \in I_k$$

$$b_{\vec{\mu}}^{\vec{\nu}} = \frac{1}{z_{\vec{\nu}}} \sum_{\vec{\tau}(\bullet, \bullet)} \frac{z_{\vec{\mu}}}{\prod_{i,j} z_{\vec{\tau}(i,j)}}$$

where sum is over all $\vec{\tau}(\bullet, \bullet)$ with $\vec{\tau}(i, j) \in I_j$ and $\vec{\mu} = \biguplus_{i,j} \nu_i^{(j)} \vec{\tau}(i, j)$.

Connections to symmetric functions

Symmetric functions on multiple variables: $\mathbf{X} = X_1, X_2, \dots$

Connections to symmetric functions

Symmetric functions on multiple variables: $\mathbf{X} = X_1, X_2, \dots$

We define the polynomial ring

$$\mathrm{Sym}_{\mathbf{X}}^* := \mathbb{C}[p_i[X_j] \mid i, j \geq 1]$$

Connections to symmetric functions

Symmetric functions on multiple variables: $\mathbf{X} = X_1, X_2, \dots$

We define the polynomial ring

$$\mathrm{Sym}_{\mathbf{X}}^* := \mathbb{C}[p_i[X_j] \mid i, j \geq 1]$$

Power sum symmetric functions:

$$p_{\mu}[X_j] := p_1[X_j]^{a_1} p_2[X_j]^{a_2} \cdots p_r[X_j]^{a_r} \quad \mu = (1^{a_1} 2^{a_2} \cdots k^{a_k})$$

$$\mathbf{p}_{\vec{\mu}}[\mathbf{X}] := p_{\mu(1)}[X_1] p_{\mu(2)}[X_2] \cdots p_{\mu(k)}[X_k] \quad \vec{\mu} \in I_k$$

Connections to symmetric functions

Symmetric functions on multiple variables: $\mathbf{X} = X_1, X_2, \dots$

We define the polynomial ring

$$\mathrm{Sym}_{\mathbf{X}}^* := \mathbb{C}[p_i[X_j] \mid i, j \geq 1]$$

Power sum symmetric functions:

$$\begin{aligned} p_{\mu}[X_j] &:= p_1[X_j]^{a_1} p_2[X_j]^{a_2} \cdots p_r[X_j]^{a_r} & \mu &= (1^{a_1} 2^{a_2} \cdots k^{a_k}) \\ \mathbf{p}_{\vec{\mu}}[\mathbf{X}] &:= p_{\mu^{(1)}}[X_1] p_{\mu^{(2)}}[X_2] \cdots p_{\mu^{(k)}}[X_k] & \vec{\mu} &\in I_k \end{aligned}$$

Schur functions

$$\mathbf{s}_{\vec{\mu}}[\mathbf{X}] := s_{\mu^{(1)}}[X_1] s_{\mu^{(2)}}[X_2] \cdots s_{\mu^{(k)}}[X_k] \quad \vec{\mu} \in I_k$$

Connections to symmetric functions

Symmetric functions on multiple variables: $\mathbf{X} = X_1, X_2, \dots$

We define the polynomial ring

$$\text{Sym}_{\mathbf{X}}^* := \mathbb{C}[p_i[X_j] \mid i, j \geq 1]$$

Power sum symmetric functions:

$$\begin{aligned} p_{\mu}[X_j] &:= p_1[X_j]^{a_1} p_2[X_j]^{a_2} \cdots p_r[X_j]^{a_r} & \mu &= (1^{a_1} 2^{a_2} \cdots k^{a_k}) \\ \mathbf{p}_{\vec{\mu}}[\mathbf{X}] &:= p_{\mu^{(1)}}[X_1] p_{\mu^{(2)}}[X_2] \cdots p_{\mu^{(k)}}[X_k] & \vec{\mu} &\in I_k \end{aligned}$$

Schur functions

$$\mathbf{s}_{\vec{\mu}}[\mathbf{X}] := s_{\mu^{(1)}}[X_1] s_{\mu^{(2)}}[X_2] \cdots s_{\mu^{(k)}}[X_k] \quad \vec{\mu} \in I_k$$

Scalar product

$$\langle \mathbf{p}_{\vec{\lambda}}[\mathbf{X}], \mathbf{p}_{\vec{\mu}}[\mathbf{X}] \rangle = \begin{cases} z_{\vec{\mu}} & \text{if } \vec{\lambda} = \vec{\mu} \\ 0 & \text{else} \end{cases}$$

Connections to symmetric functions – continued

Frobenius characteristic of trivial representation of \mathcal{U}_k

$$\begin{aligned} E_r &:= \sum_{\vec{\mu} \in I_r} \frac{\mathbf{p}_{\vec{\mu}}[\mathbf{X}]}{\mathbf{z}_{\vec{\mu}}} \\ &= \sum_{(1^{a_1} 2^{a_2} \dots r^{a_r}) \vdash r} s_{a_1}[X_1] s_{a_2}[X_2] \cdots s_{a_r}[X_r] \end{aligned}$$

Connections to symmetric functions – continued

Frobenius characteristic of trivial representation of \mathcal{U}_k

$$\begin{aligned} E_r &:= \sum_{\vec{\mu} \in I_r} \frac{\mathbf{p}_{\vec{\mu}}[\mathbf{X}]}{\mathbf{z}_{\vec{\mu}}} \\ &= \sum_{(1^{a_1} 2^{a_2} \dots r^{a_r}) \vdash r} s_{a_1}[X_1] s_{a_2}[X_2] \cdots s_{a_r}[X_r] \end{aligned}$$

Proposition (OSSZ 2022)

$$b_{\vec{\mu}}^{\vec{\nu}} = \frac{1}{\mathbf{z}_{\vec{\nu}}} \langle \mathbf{p}_{\vec{\nu}}[\mathbf{E}], \mathbf{p}_{\vec{\mu}}[\mathbf{X}] \rangle$$

Characters, symmetric functions, and plethysm

Theorem (OSSZ 2022)

$$\chi_{\mathcal{U}_k}^{\vec{\lambda}}(d_{\vec{\mu}}) = \langle \mathbf{s}_{\vec{\lambda}}[\mathbf{E}], \mathbf{p}_{\vec{\mu}}[\mathbf{X}] \rangle$$

Characters, symmetric functions, and plethysm

Theorem (OSSZ 2022)

$$\chi_{\mathcal{U}_k}^{\vec{\lambda}}(d_{\vec{\mu}}) = \langle \mathbf{s}_{\vec{\lambda}}[\mathbf{E}], \mathbf{p}_{\vec{\mu}}[\mathbf{X}] \rangle$$

$$\chi_{G_{\lambda}}^{\vec{\lambda}}(d_{\vec{\mu}}) = \langle \mathbf{s}_{\vec{\lambda}}[\mathbf{X}], \mathbf{p}_{\vec{\mu}}[\mathbf{X}] \rangle$$

Characters, symmetric functions, and plethysm

Theorem (OSSZ 2022)

$$\chi_{\mathcal{U}_k}^{\vec{\lambda}}(d_{\vec{\mu}}) = \langle \mathbf{s}_{\vec{\lambda}}[\mathbf{E}], \mathbf{p}_{\vec{\mu}}[\mathbf{X}] \rangle$$

$$\chi_{G_{\lambda}}^{\vec{\lambda}}(d_{\vec{\mu}}) = \langle \mathbf{s}_{\vec{\lambda}}[\mathbf{X}], \mathbf{p}_{\vec{\mu}}[\mathbf{X}] \rangle$$


Corollary

Multiplicity of $V_{S_k}^{\mu}$ in $\text{Res}_{S_k}^{\mathcal{U}_k} V_{\mathcal{U}_k}^{\vec{\lambda}}$ is $\langle \mathbf{s}_{\lambda(1)}[s_1] \mathbf{s}_{\lambda(2)}[s_2] \cdots \mathbf{s}_{\lambda(k)}[s_k], \mathbf{s}_{\mu} \rangle$



Outline

- 1 The plethysm problem
- 2 Diagram algebras
- 3 Uniform block permutation algebra
- 4 Symmetric chain decompositions

Young lattice for partitions in box



Young lattice for partitions in box





- Partitions in box of size $w \times h$
- Crystal $B(w)$ of type A_h
- Plethysm

$$s_w[s_h[x + y]] = \sum_{\nu} a_{wh}^{\nu} s_{\nu}$$

ν at most two parts

Young lattice for partitions in box



- Partitions in box of size $w \times h$

- Crystal $B(w)$ of type A_h

- Plethysm

$$s_w[s_h[x + y]] = \sum_{\nu} a_{wh}^{\nu} s_{\nu}$$

ν at most two parts

- Example:

$$s_2[s_2[x + y]] = s_4 + s_{22}$$

Thank you !

Thank you !

Remark (Take away)

Plethysm is hard!

Thank you !

Remark (Take away)

Plethysm is hard!

Remark (Take away)

The random walk exploring plethysm leads to interesting mathematics!

Thank you !

Remark (Take away)

Plethysm is hard!

Remark (Take away)

The random walk exploring plethysm leads to interesting mathematics!

